

1 **Eutrophication mitigation in rivers: 30 years of trends in**  
2 **spatial and seasonal patterns of biogeochemistry of the**  
3 **Loire River (1980-2012)**

4 **C. Minaudo<sup>1</sup>, M. Meybeck<sup>2</sup>, F. Moatar<sup>1</sup>, N. Gassama<sup>1</sup>, F. Curie<sup>1</sup>**

5 [1]{University of Tours, E.A. 6293 GéHCO, Tours, France }

6 [2]{Paris VI University, UMR METIS, Paris, France }

7 Correspondence to: C. Minaudo (camille.minaudo@etu.univ-tours.fr)

8

9 **Abstract**

10 Trends and seasonality analysis since 1980 and longitudinal distribution from headwaters to  
11 estuary of chlorophyll *a*, nitrate and phosphate were investigated in the eutrophic Loire River.  
12 The continuous decline of phosphate concentrations recorded since 1991 both in the main  
13 river and in the tributaries led to the conclusion that it was responsible for the significant  
14 reduction in phytoplanktonic biomass across the whole river system, although *Corbicula*  
15 clams species invaded the river during the same period and probably played a significant role  
16 in the phytoplankton decline. While eutrophication remained lower in the main tributaries  
17 than in the Loire itself, they were found to contribute up to  $\approx 35\%$  to the total nutrient load of  
18 the main river. The seasonality analysis revealed significant seasonal variations for the  
19 different eutrophication metrics and called into question the classical monthly survey  
20 recommended by national or international authorities. Reducing P-inputs impacted these  
21 seasonal variations: the decline of seasonal amplitudes of chlorophyll *a* reduced the seasonal  
22 amplitude of orthophosphate and of daily variations of dissolved oxygen and pH but did not  
23 significantly affect the seasonal amplitude of nitrate. Thus, the influence of phytoplankton on  
24 seasonal variations of nitrate was minor throughout the period of study.

25

## 1 **1 Introduction**

2 For several decades, eutrophication has become a major issue affecting most surface waters  
3 (Smith et al., 1999; Hilton et al., 2006; Smith and Schindler, 2009; Grizzetti et al., 2012;  
4 Romero et al., 2012). The regulation of nutrient inputs in waters by the elimination of N and P  
5 during waste-water treatment, better agricultural practices and restriction of the use of  
6 phosphorus products (EEC 1991a and b) led to a decrease in phosphate and/or nitrate content  
7 which is recorded in several European rivers presenting temperate and continental regimes  
8 since the mid-1990s, including the Elbe (Lehmann and Rode, 2001), the Seine (Billen et al.,  
9 2007), the Thames (Howden et al., 2010), the Danube (Istvánovics and Honti, 2012), the  
10 Rhine (Hartmann et al., 2007) as well as some Mediterranean rivers (Ludwig et al., 2009) and  
11 Scandinavian rivers (Grimvall et al., 2014).

12 Surface water quality is also affected by variations in hydro-climatic conditions (Durance and  
13 Ormerod, 2010) and nutrients availability is not the only limiting factor of phytoplanktonic  
14 growth in rivers: successful phytoplankton species in rivers are selected on their ability to  
15 survive high frequency irradiance fluctuations and the important determinants are turbidity (or  
16 its impact upon underwater light) and the water residence time (Istvánovics and Honti, 2012;  
17 Krogstad and Lovstad, 1989; Reynolds and Descy, 1996; Reynolds et al., 1994). In Europe,  
18 both climatic models and observations show a general rise in air and water temperature since  
19 the 1970s (Moatar and Gailhard, 2006; Whitehead et al., 2009; Bustillo et al., 2013) and  
20 models predict lower water discharge and rising temperatures during summer, potentially  
21 intensifying the risk of eutrophication (Arheimer et al., 2005; Barlocher et al., 2008; Lecerf et  
22 al., 2007; Whitehead et al., 2009) as shallow rivers are particularly susceptible to  
23 eutrophication (Istvánovics et al., 2014). Besides, phytoplanktonic biomass remains at a high  
24 level in many water bodies, evidencing that leaching of long last stored nutrient in soils is still  
25 significant: if nutrient mobility should increase with global warming because of the  
26 acceleration of organic matter mineralization and of higher soil leaching (Bouraoui et al.,  
27 2002; Arheimer et al., 2005), the river system response time to a nitrogen input reduction is  
28 limited by the time required for nitrate to be released from soil to receiving waters (Jackson et  
29 al., 2008; Bouraoui and Grizzetti, 2011). Therefore we should expect that changes in current  
30 agricultural practices may improve water quality only after several decades (Behrendt et al.,  
31 2002; Howden et al., 2010).

1 The first regulatory studies of the largest French river eutrophication, i.e. the Loire River,  
2 were made in the 1980s in the Middle and Lower segments (Crouzet, 1983; Meybeck et al.,  
3 1988; Lair and Reyes-Marchant, 1997; Etcheber et al., 2007). The Middle reaches (Fig. 1)  
4 were recognized as being the most eutrophic sector (Lair and Reyes-Marchant, 1997)  
5 resulting from high P levels (Floury et al., 2012), low river velocity and shallow waters, its  
6 multiple channels morphology with numerous vegetated islands slowing down flow velocity  
7 (Latapie et al., 2014). In recent years, Loire eutrophication indicators and their trends  
8 recorded several variations: (i) decline of chlorophyll *a* in the Middle segment since the late  
9 1990s (Floury et al., 2012), (ii) decline of phosphorus as well in the Middle Loire (Gosse et  
10 al., 1990; Moatar and Meybeck, 2005; Oudin et al., 2009), (iii) development of *Corbicula*  
11 *fluminea* as an invasive species since the 1990s (Brancotte and Vincent, 2002) and (iv)  
12 dominance of small centric diatoms and green algae in phytoplankton population, for most of  
13 the year in the Middle and Lower river sectors (Abonyi et al., 2012, 2014; Descy et al., 2011).

14 Most previous studies focused on the Middle Loire, which represents only 25% of the total  
15 drainage basin and excluded the main tributaries and their possible influences on the main  
16 river course. Besides, most studies on river eutrophication stayed at the inter-annual variations  
17 and did not investigate how long term trends might affect the river biogeochemistry at the  
18 seasonal or the daily scale, while seasonal and daily cycles are especially amplified in  
19 eutrophic rivers (Moatar et al., 2001). This paper examines longitudinal distributions and  
20 long-time trends of chlorophyll *a* and nutrients over three decades (1980-2012) and for the  
21 whole Loire basin. Thus, it includes the study of the main tributaries variations and their  
22 potential influences on the Loire main stem. It also focuses on how the noticeable long term  
23 changes affected the biogeochemical functioning of the river at the seasonal scale, exploring  
24 the seasonal variations of chlorophyll *a* and nutrients since 1980 and examining both seasonal  
25 and daily fluctuations of dissolved oxygen and pH since 1990.

26

## 27 **2 Study area and data compilation**

### 28 **2.1 Geographical and physical characteristics**

29 The Loire River basin (110,000 km<sup>2</sup>) covers 20% of the French territory. Its hydrological  
30 regime is pluvial with some snow-melt influences because of high headwater elevation (6% of  
31 the basin area is over 800 m above sea level). The main stem can be divided into three parts

1 (Fig.1, Table 1): (i) the Upper Loire (18% of basin area; stations 1 to 9) extending from the  
2 headwaters to the confluence with the River Allier; (ii) the Middle Loire (24%; stations 10 to  
3 18) from the Loire-Allier confluence to the Loire-Cher confluence which receives only minor  
4 inputs from small tributaries; (iii) the Lower Loire (65%; stations 19 to 21) which receives  
5 major tributaries (Cher, Indre, Vienne and Maine Rivers) doubling the river basin area and the  
6 average river water discharge.

7 As summer low flows can reach critically low levels in the Middle reaches where four nuclear  
8 power plants are located (Fig. 1), two dams were constructed on the Allier and Upper Loire  
9 (Naussac 1981 and Villerest, 1984) to maintain low flows over a minimum of  $60 \text{ m}^3 \text{ s}^{-1}$ .  
10 Grangent dam was constructed in 1957 for electricity production purposes. The median  
11 annual discharge over the last 30 years is  $850 \text{ m}^3 \text{ s}^{-1}$  at the basin outlet (station 21) and the  
12 median in the driest period from July to September is only  $250 \text{ m}^3 \text{ s}^{-1}$ , corresponding to only 2  
13  $\text{L s}^{-1} \text{ km}^{-2}$ . The driest years were 1990, 1991, 2003 and 2011 with a daily discharge average at  
14 station 21 reaching sometimes  $100 \text{ m}^3 \text{ s}^{-1}$ .

15 The headwater catchment is a mountainous area and the Loire itself runs through narrow  
16 gorges and valleys (Latapie, 2011). After the confluence with the Allier, the geomorphology  
17 of the Middle Loire favors phytoplankton development, its multiple channels with numerous  
18 vegetated islands slowing down flow velocity and the valleys becoming wider (Latapie et al.,  
19 2014). As a consequence, average water depth can be low in the summer ( $\approx 1 \text{ m}$ ), contributing  
20 to warming and lighting up the water column.

21 The temperature is always at least  $2^\circ\text{C}$  lower in the Upper part than in the lower reaches  
22 (annual medians are around  $15^\circ\text{C}$  in the Upper Loire during April-October versus  $19^\circ\text{C}$  in the  
23 Middle and Lower segments) and is affected by global warming. Hence, Moatar and Gailhard  
24 (2006) showed that mean water temperature has increased by 2.4 to  $3^\circ\text{C}$  in spring and  
25 summer since 1975 due to rising air temperature (Gosse et al., 2008) without a significant  
26 impact on phytoplanktonic development (Floury et al., 2012). Approximately 60% of this  
27 general rise in water temperature during the warm period was explained by rising air  
28 temperature and 40% by a decrease in the May/June river discharge since 1977 (Moatar and  
29 Gailhard 2006, Floury et al., 2012). The water returning to the Loire from the nuclear power  
30 plants only raises the temperature by a few tenths of a degree thanks to an atmospheric  
31 cooling system (Vicaud, 2008) and does not influence the thermal regime of the river studied  
32 here.

1 Urban pressure is significant with 8 million people living in the Loire Basin (2008 population  
2 census by the French National Institute of Statistics and Economic Studies, INSEE), mainly  
3 concentrated near the main river course. It corresponds to an overall population density of 73  
4 inhabitant km<sup>-2</sup>. The density is greater in the Upper Loire (144 inhab. km<sup>-2</sup>, Table 1) due to the  
5 city of Saint Etienne (180,000 inhabitants). The Middle and Lower catchments contain some  
6 major riparian cities (Fig. 1) with a stable population density around 76 inhab. km<sup>-2</sup>.

7 Agricultural pressure is defined here with two indicators: the percentage of the basin occupied  
8 by arable land and the Agricultural Pressure Indicator (API) represented as the quotient of  
9 (pasture + forest) over (pasture + forest + arable land). According to the Corine Land Cover  
10 database (2006), the headwater areas are mostly forested or pastureland (Table 1). Arable land  
11 increases from headwaters going downstream to reach 30% of the total basin area at station  
12 21. Land use distribution in the major tributaries differs widely (Table 2): the Allier  
13 (catchment at station A) is mostly composed of pasture; the Cher at station B has similar  
14 amounts of pasture and arable land, most of the rest being forested; half of the Indre basin at  
15 station C is arable land, but this tributary drains only 3% of the total basin; the Vienne and the  
16 Maine contribute very significantly to the total area of arable land in the Loire basin. Urban  
17 pressure is also significant in the Maine catchment due to the cities of Le Mans and Angers  
18 (Fig.1).

19

## 20 **2.2 River monitoring datasets**

21 Water quality databases from regulatory surveys (Loire Brittany river basin agency, AELB)  
22 used here (chlorophyll a, pheopigments, nitrate (NO<sub>3</sub><sup>-</sup>), nitrite (NO<sub>2</sub><sup>-</sup>), Kjeldahl nitrogen  
23 (NKj), orthophosphate (PO<sub>4</sub><sup>3-</sup>) and total phosphorus (Ptot)) are available online  
24 (<http://osur.eau-loire-bretagne.fr/exportosur/Accueil>). Sixty-nine monitoring stations were set  
25 up along an 895 km stretch. Stations sampled at least monthly between 1980 and 2012 (bi-  
26 monthly or weekly for some variables) were selected for analysis in this paper (17 stations,  
27 Fig. 1). To take into account the influence of major tributaries, five sampling sites at each of  
28 the major tributary outlets were also included (stations A to E).

29 The water quality of the Loire River has also been assessed during several other surveys,  
30 generally with high sampling frequency, but these data have seldom been used and/or  
31 compared in previous studies. They included:

- 1 i) Water quality surveys upstream and downstream of nuclear power plants carried out  
2 since the early 1980s by the French Electricity Company (EDF) (Moatar and  
3 Gailhard, 2006; Moatar et al., 2013); see stations 12, 14, 16 and 19 on Fig. 1.  
4 These datasets were used to improve the spatial analysis. These surveys included  
5 temperature, dissolved oxygen and pH recorded hourly at station 19 enabling us to  
6 analyze possible changes in day/night variations (variables hereafter named *delta*  
7 *O<sub>2</sub>* and *delta pH* corresponding to the daily range of *O<sub>2</sub>* and pH).
- 8 ii) The Orléans city experimental survey carried out by the Loire basin authority at  
9 station 15 from 1981 to 1985, measuring nutrients and chlorophyll a every three  
10 days (Crouzet, 1983; Moatar and Meybeck, 2005).

11 River flow datasets on a daily basis were taken from the national “Banque Hydro” database  
12 (<http://www.hydro.eaufrance.fr/>). The local population census (INSEE, 2008) and the Corine  
13 Land Cover (2006) were also used to estimate the general characteristics at different water  
14 quality stations (Tables 1 and 2).

15

## 16 **3 Methods**

### 17 **3.1 Data pre-processing**

18 To validate the AELB datasets and eliminate remaining outliers, log-log relationships  
19 between concentration and discharge were analyzed and compared with previous research  
20 studies carried out during targeted periods (Grosbois et al., 2001; Moatar and Meybeck,  
21 2005). The separation of living phytoplankton biomass (characterized by chlorophyll a) and  
22 algal detritus (characterized by pheopigments) depends on the protocol used and since this  
23 protocol may have changed over the last 30 years, we worked with the sum of chlorophyll *a*  
24 and pheopigments, which increased the robustness of the data and corresponded better to  
25 phytoplanktonic biomass as an active biomass and organic detritus (Dessery et al., 1984;  
26 Meybeck et al., 1988). Thus, for clarity further in the text, “Chl. *a*” corresponds to the sum  
27 chlorophyll *a* + pheopigments.

28  $\text{PO}_4^{3-}$  time series included periods reaching the limit of quantification. When evidenced, such  
29 data were not taken into account to avoid mis-interpretation of such constant values. The  
30 datasets also included periods with missing values. In all cases, no infilling were realized.

1 Sampling frequencies were most of the time monthly (only 10% of datasets were sampled on  
2 average every two weeks or more often), but in order to homogenize the time series, the  
3 following analysis was conducted on monthly medians.

4 To assess longitudinal distribution of nutrients and phytoplanktonic biomass, each year was  
5 divided into two seasons: “*summer*”, here considered as the phytoplankton growth period  
6 from April to October, when more than 90% of the phytoplankton bloom is observed (Leitão  
7 and Lepretre, 1998) and “*winter*”, here November to March when Chl. *a* concentrations are  
8 usually under  $20 \mu\text{g L}^{-1}$  (average *winter* Chl. *a* in the Middle Loire  $\approx 20 \mu\text{g L}^{-1}$  for the  
9 considered period).

10 Uncertainties on estimates of concentration averages were assessed using Monte Carlo  
11 random draws (Moatar and Meybeck, 2005) on experimental high frequency data at Orléans  
12 city (station 15). Uncertainties on seasonal means varied between 10% ( $\text{NO}_3^-$ ) and 30% ( $\text{PO}_4^{3-}$ )  
13 ) in *summer* and between 6% ( $\text{NO}_3^-$ ) and 10% ( $\text{PO}_4^{3-}$ ) in *winter*.

14 When both river discharge and nutrient concentration datasets were available during the  
15 period considered, average annual fluxes were calculated to assess the contribution of each  
16 major tributary to the Loire. This calculation was possible during 1980-86 and 1994-2006 for  
17 the Allier input, 1985-90 and 1999-2009 for the Cher, 2006-2011 for the Vienne and 1981-  
18 2012 for the Maine.

19 In order to assess potential changes in the nitrogen to phosphorus molar ratio (N:P further in  
20 the text) and make the link with possible nutrient limitation of phytoplankton, this ratio was  
21 calculated using  $\text{N}_{\text{tot}}$  (sum of  $\text{NO}_3^-$ ,  $\text{NO}_2^-$  and  $\text{NKj}$ ) and  $\text{P}_{\text{tot}}$ .

## 22 **3.2 Building up spatio-temporal diagrams**

23 Time series were represented with a 2D spatial x-axis and seasonal y-axis. This allowed the  
24 observation of both longitudinal and seasonal distribution during a certain period, between the  
25 river headwaters to the estuary and from January to December. When needed and possible,  
26 missing data were interpolated both spatially and temporally to represent a smoother diagram.  
27 Three periods were defined and separated the last three decades in three sub-periods on the  
28 basis of Chl. *a* concentrations: 1980-1989, 1990-2001 and 2002-2012.

### 1 3.3 Time series decomposition

2 Long-term trends and seasonal variations analysis were carried out using Dynamic Harmonic  
3 Regression (DHR) technique, extensively described in Taylor et al., (2007) (a brief outline of  
4 it is also explained in Halliday et al., 2012 and 2013). It decomposes an observed time series  
5 into its component parts:

$$6 \quad f(t) = T(t) + S(t) + Irr(t) \quad (1)$$

7 where  $f$  is the observed time series,  $T$  is the identified trend,  $S$  the seasonal component and  $Irr$   
8 the “irregular” component defined as white noise, representing the residuals.

9 The trend was defined using an Integrated Random Walk model. It is a special case of the  
10 Generalized Random Walk model (GRW) and has been shown to be useful for extracting  
11 smoothed trends (Pedregal et al., 2007). This provided the identified trend and the slope of the  
12 trend.

13 The seasonal components were defined as follow:

$$14 \quad S(t) = \sum_i^{N/2} [a_{i,t} \cos(\omega_i t) + b_{i,t} \sin(\omega_i t)] \quad \omega_i = \frac{2\pi \cdot i}{N} \quad i = 1, 2, \dots, \left[ \frac{N}{2} \right] \quad (2)$$

15 where  $\omega_i$  are the fundamental and harmonic frequencies associated with the periodicity in the  
16 observed time series chosen by reference to the spectral properties. For instance, the period 12  
17 was corresponding to a monthly sampling in an annual cycle.

18 The phase and amplitude parameters were modeled as GRW processes and estimated  
19 recursively using the Kalman Filter and the Fixed Interval Smoother. These parameters were  
20 defined as non-stationary stochastic variables to allow variation with time i.e. allow non-  
21 stationary seasonality and represent better the dynamic of the observed parameters.

22 Significance of the seasonality was based on the squared correlation coefficient between  
23 calculated seasonal component and detrended data. Similarly, significance of the trend was  
24 determined based on the squared correlation coefficient between calculated trend and  
25 deseasonalized data.

26 Stations 4 (Upper Loire), 18 (Middle) and 21(Lower) presented a large amount of data and  
27 were selected here to present and discuss the DHR analysis. Similarly, water discharge data at  
28 station 15 was daily and continuous since 1980 and was selected for the DHR analysis  
29 presented in the Results section.



## 1 4 Results

### 2 4.1 Long term trends and longitudinal distributions of Chl. *a* and nutrients

3 Chl. *a* *summer* medians (used as the prime indicator of eutrophication) showed a very clear  
4 longitudinal increase from headwaters to river mouth (Fig. 2a). At the headwaters, Chl. *a*  
5 concentrations remained below  $30 \mu\text{g L}^{-1}$  between 1981 and 2012. It has been shown in other  
6 studies that in the Upper Loire reservoirs which have always been eutrophic since the 1980s  
7 (Aleya et al., 1994; Jugnia et al., 2004), the phytoplankton assemblage is lake-like and these  
8 species do not survive very long in the turbulent and quite turbid river downstream (Abonyi et  
9 al., 2011, 2014), explaining why Chl. *a* remains at low levels. In the lowest reaches of the  
10 Upper Loire (station 9), Chl. *a* was higher but showed a descending trend for the whole  
11 period. In the Middle segments, Chl. *a* levels increased between 1981 and 1990 by a factor of  
12 two (Table 3). The maximum ever measured occurred at station 18 in early October 1990  
13 ( $365 \mu\text{g L}^{-1}$ ). The next decade, the situation already started to decrease in the Middle Loire ( $-5$   
14  $\mu\text{g L}^{-1} \text{ year}^{-1}$ ) and even more in the Lower ( $-9 \mu\text{g L}^{-1} \text{ year}^{-1}$ ). Finally, since 2002, the decline  
15 generalized to the whole river and trends slopes were  $\approx -5 \mu\text{g L}^{-1} \text{ year}^{-1}$  in the Middle Loire  
16 and  $-4 \mu\text{g L}^{-1} \text{ year}^{-1}$  in the Lower reach.

17 *Winter* medians of phosphate concentrations increased downstream of station 2 (Fig. 2b) and  
18 the maximum for the Upper segment was reached at station 4, where population density is  
19  $143 \text{ inhab. km}^{-2}$ , a maximum for the whole basin. Population density decreased to  $75 \text{ inhab.}$   
20  $\text{km}^{-2}$  between stations 4 and 9, with a corresponding reduction in the phosphate levels.  $\text{PO}_4^{3-}$   
21 levels were stabilized in the Middle Loire (stations 10 to 18).

22 The general phosphorus decline during the last decade can be observed along the whole  
23 longitudinal profile. Phosphate was at its maximum in the 1980s (above  $100 \mu\text{g P L}^{-1}$ ) for  
24 almost the whole main stem. It then decreased gradually to reach lower levels  $<70 \mu\text{g P L}^{-1}$ . In  
25 the urbanized Upper part (stations 3 and 4), from a *winter* median of  $190 \mu\text{g P L}^{-1}$  during  
26 1980-1989, phosphate decreased to its current level ( $60 \mu\text{g P L}^{-1}$ ). Average phosphate in the  
27 Middle and Lower reaches has reduced at least two-fold since 1980. At the Lower Loire outlet  
28 (station 21), phosphate contents increased during 1980-1989 and then decreased at the rate of  
29  $\approx -4 \mu\text{g P L}^{-1} \text{ y}^{-1}$ . Downstream the main reservoirs (Upper Loire), a noticeable decrease in  
30 phosphorus concentration was observed. This was probably partly due to P retention between  
31 stations 4 and 5 (Fig. 1) as a large part of the particulate matter is stored in the reservoir.

1 The *winter* nitrate longitudinal profile showed a regular increase from 1 mg N L<sup>-1</sup> in the  
2 headwaters to 3.5 mg N L<sup>-1</sup> at the river mouth (Fig. 2c). This longitudinal rise could be  
3 observed throughout the period of study. The upstream reservoirs did not seem to impact the  
4 nitrogen concentration as nitrate represented most of the total nitrogen and the  
5 phytoplanktonic uptake within these reservoirs is not questioned here: Fig. 2c present *winter*  
6 nitrate concentration. Annual median nitrate concentration remained stable in the Upper  
7 Loire, with no significant trends since 1980. In the Middle segment, it only presented an  
8 increasing trend during the 1990s (+0.1 mg N L<sup>-1</sup> y<sup>-1</sup>) but the more significant variations were  
9 observed in the Lower reaches at station 21 where nitrate increased on average at +0.3 mg N  
10 L<sup>-1</sup> y<sup>-1</sup> during the 1980s, a bit less the next decade (+0.1 mg N L<sup>-1</sup> y<sup>-1</sup>) and finally slightly  
11 decreased since 2002.

12 These trends provided by the DHR model were always significant and explained at least 50%  
13 of the variations in the deseasonalized time series (Table 3). The most significant trends were  
14 observed in Chl. *a* and PO<sub>4</sub><sup>3-</sup>. The long term variations in NO<sub>3</sub><sup>-</sup> were less pronounced  
15 justifying a lower corresponding strength.

#### 16 **4.2 Seasonal shifts across the longitudinal distribution of Chl. *a* and nutrients**

17 Throughout the period of study, Chl. *a* concentrations reached their maximum in July or  
18 August for the whole Loire River. During the 1980s and 90s, phytoplankton production  
19 usually started in early April, reached a peak in early May with a second peak in late August  
20 (Fig. 3a) suggesting different phytoplankton communities growth (Abonyi et al., 2012, 2014).  
21 After mid-November, Chl. *a* concentrations were very low. A slight change is nevertheless  
22 evidenced: between 1980 and 2000 in the Middle and Lower Loire, Chl. *a* concentrations  
23 reached occasionally their maximum in October (it is the case of the years 1985, 1988, 1989,  
24 1990, 1995); since 1996, it never happened again.

25 Phosphate spatio-temporal variations showed inverted seasonal patterns between the Upper  
26 and Middle-Lower Loire (Fig. 3b). Maximum phosphorus levels were observed in the middle  
27 part of the Upper section (stations 3 to 5) as a result of urban pressure, previously mentioned  
28 in the longitudinal profile description. In this upstream reach where phytoplankton  
29 development is limited, the seasonal maximum level was observed in *summer* when low flows  
30 cannot dilute urban phosphorus inputs; during the period 2002-2012, PO<sub>4</sub><sup>3-</sup> medians reached  
31 140 µg P L<sup>-1</sup> at station 4 in June. In the lower reaches of the Upper Loire, the Middle and the

1 Lower reaches (stations 8 to 21), the seasonality of phosphate was inverted compared to the  
2 Upper Loire and clearly controlled by eutrophication with a minimum ( $<30 \mu\text{g P L}^{-1}$ )  
3 occurring during *summer* due to phytoplankton uptake.

4 Nitrate concentrations had a very clear seasonality (Fig. 3c) with maximum levels during  
5 *winter* (leaching) along the whole Loire River. In *summer*, nitrate was very low with  
6 concentrations around 1 to 2.5 mg N L<sup>-1</sup> along the whole river profile and the lowest  
7 concentrations were recorded in August in the Middle Loire. The *summer* nitrate minima have  
8 increased since 1980: around 0.4 mg N L<sup>-1</sup> in the Middle Loire between 1980 and 1999, the  
9 average *summer* 10% percentile increased to 1 mg N L<sup>-1</sup> this last decade. A seasonal Kendall  
10 test analysis (station 15, 1980-2012) revealed that water discharge explained 26% of the  
11 nitrate variance.

12 The Dynamic Harmonic Regression model represented well the time series, depending on the  
13 river reach and the type of variable (Table 4). Seasonal components were stronger in Middle  
14 and Lower Loire than in Upper, with better correlations between detrended time series and  
15 calculated seasonal component (45 to 85% variance explained by the seasonal component in  
16 the Middle and Lower against 15-45% in the Upper). Chl. *a* series were well represented by  
17 the seasonal component, whereas PO<sub>4</sub><sup>3-</sup> was sometime poorly explained, illustrating the high  
18 variability of this parameter. Nitrate time series presented the best fits, with around 80% of  
19 the variance explained by the seasonal component in the Middle and Lower reaches.

#### 20 **4.3 Analysis of the main tributaries variations and their impacts on the Loire** 21 **long-term trends**

22 Trends in the main tributaries of the Loire River (stations A to E) mimicked the Loire River  
23 variations with high signs of eutrophication during the 1980s and 1990s followed by a general  
24 decline (Table 5).

25 Chl. *a* in the tributaries remained under the Loire main stem levels in each of the major  
26 tributaries except for the Cher River (station B): its highest Chl. *a* concentrations during the  
27 1990s were very close to the extreme values reached at the same time in the Middle Loire  
28 (average seasonal variation  $\approx 190 \mu\text{g L}^{-1}$  during the 1990s). Nonetheless, trends in Chl. *a*  
29 concentrations were everywhere following the same pattern, with high seasonal variations and  
30 high annual medians between 1980 and 2001, and then clearly declined the last decade.

1 Phosphate concentrations decreased everywhere continuously from high values in the 1980s  
2 ( $\approx 200 \mu\text{g P L}^{-1}$ ) down to  $\approx 50 \mu\text{g P L}^{-1}$  except at station E (Maine River) where  $\text{PO}_4^{3-}$  first  
3 increased during the 1980s from  $200 \mu\text{g P L}^{-1}$  to peak in 1992 at  $300 \mu\text{g P L}^{-1}$  and finally  
4 declined towards  $50 \mu\text{g P L}^{-1}$ .

5 Like in the Loire River, nitrate concentrations in the main tributaries increased slightly since  
6 1980, but levels and seasonal amplitudes progressed differently: quite low in the Upper  
7 tributary (station A, annual medians  $\approx 1.5 \text{ mg N L}^{-1}$ ),  $\text{NO}_3^-$  reached higher concentrations in  
8 the other tributaries and extreme values in the Maine River with *winter* maximums over  $10$   
9  $\text{mg N L}^{-1}$  during the 1990s. At each station but station A,  $\text{NO}_3^-$  seasonal amplitudes slightly  
10 started to decrease since 2002 i.e. the *summer* minimum slightly increased.

11 At each major tributary confluence, the tributaries inputs could contribute on average to 35%  
12 of the main river nutrient fluxes. The more significant inputs were coming from the Allier  
13 River (station A) discharging almost the same amount of  $\text{NO}_3^-$  and  $\text{PO}_4^{3-}$  as the Upper Loire  
14 River. Because of the lack of data allowing nutrient fluxes calculations on a fine temporal  
15 scale, these results are to be considered with caution. But they are certainly giving good  
16 approximations of how much these tributaries can influence the Loire main stem  
17 eutrophication trajectory.

#### 18 **4.4 Seasonal amplitudes of Chl. *a*, nutrients, $\text{O}_2$ and pH in the Middle Loire**

19 As described above, Chl. *a*, nitrate and phosphate concentrations presented different patterns  
20 of seasonality depending on the location. This paragraph focuses on seasonality of nutrients  
21 and Chl. *a* at station 18 and on dissolved oxygen, pH and temperature at station 19. Both of  
22 these stations are representative of the Middle Loire reach where the highest signs of  
23 eutrophication occurred in the early 1990s.

24 Chl. *a* seasonal variation at station 18 increased during the 1980s (Fig. 4a) from  $150$  to  $240 \mu\text{g}$   
25  $\text{L}^{-1}$  (1990) and then presented a spectacular decline in two steps: first, it went down to  $150 \mu\text{g}$   
26  $\text{L}^{-1}$  in 1992 and remained at the same level the next 8 years; then, it kept on decreasing since  
27 2000 to finally reach levels of amplitude around  $50 \mu\text{g L}^{-1}$ . Phosphate seasonal variations  
28 decreased continuously from  $150 \mu\text{g P L}^{-1}$  in 1980 to  $30 \mu\text{g L}^{-1}$  in 2012 (Fig. 4b), at the rate of  
29  $-6 \mu\text{g P L}^{-1} \text{ year}^{-1}$  in the 1980s,  $-4 \mu\text{g P L}^{-1} \text{ year}^{-1}$  in the 1990s and finally reached a stable  
30 variation since 2008 (Table 4). The seasonal variations of  $\text{NO}_3^-$  presented another pattern  
31 through the last 30 years (Fig. 4c): it increased from  $2.2 \text{ mg N L}^{-1}$  in 1980 to  $2.8 \text{ mg N L}^{-1}$  in

1 1991, then remained stable around  $2.9 \text{ mg N L}^{-1}$  the next 7 years to finally decrease slightly  
2 down to  $2 \text{ mg N L}^{-1}$ .

3 Interannual dissolved oxygen concentration and pH at station 19 did not present any  
4 significant trend (Fig. 4d and 4e): since 1990, annual average  $\text{O}_2 = 10.8 \text{ mg L}^{-1}$  and  $\text{pH} = 8.3$ .  
5 At the daily scale, the variations of  $\text{O}_2$  were synchronous with water temperature: the typical  
6  $\text{O}_2$  daily cycle corresponded to a minimum at sunrise, followed by a rapid increase and a  
7 maximum observed two hours after solar mid-day; the daily range could reach  $10 \text{ mg L}^{-1}$ ,  
8 with oxygen saturation ranging from 60% to 200%. These daily variations greatly challenge  
9 the validity of  $\text{O}_2$  measurements as a water quality indicator within the regulatory monthly  
10 survey of such eutrophic river. Alongside daily oxygen cycles, significant daily pH cycles  
11 were observed (see also Moatar et al., 2001). Dissolved  $\text{CO}_2$  and/or bicarbonate uptake by  
12 primary producers during the solar day led to increasing pH. By contrast, night-time  
13 respiration was reducing pH. In the Loire, daily pH cycles were pronounced with the same  
14 phase as the  $\text{O}_2$  cycle. The common daily pH range in *summer* was 0.8 unit and could reach 1  
15 pH unit. Because these variations are linked to the in-stream biological activities, daily  $\text{O}_2$  and  
16 daily pH amplitudes presented a well-defined seasonality, with maximum reached in *summer*.

17 DHR model applied on water temperature ( $^{\circ}\text{C}$ ) successfully represented the observations  
18 with squared correlation coefficients  $R^2$  of 0.96. Performances were lower for discharge (Q)  
19 with  $R^2 = 0.57$ . Both  $^{\circ}\text{C}$  and Q trends were weak (only 20% of the variances of  
20 deseasonalized datasets were explained); however,  $^{\circ}\text{C}$  was increasing, and Q slightly  
21 decreasing.

22

## 23 **5 Discussion**

### 24 **5.1 Role of agricultural and urban pressures on the Loire long-term variations**

25 The population density profile (Fig. 2) illustrates well the fact that phosphate concentrations  
26 are linked with urban P-inputs. Thus, most changes in phosphate levels are connected to more  
27 efficient sewage treatment plants (de-phosphatation steps were set up) and the use of  
28 phosphate-free detergents. De-phosphatation technologies were not implemented at the same  
29 time across the basin, explaining different trends for different catchments. These observations  
30 support previous studies highlighting the need for phosphorus control (Gosse et al., 1990;  
31 Oudin, 1990). This control has considerably reduced phosphate concentration in the surface

1 waters of the Loire basin (Bouraoui and Grizzetti, 2011). Nevertheless, Descy et al. (2011)  
2 assessed the biogeochemical processes using numerical models of the Middle reaches during  
3 the year 2005 and the phosphorus reduction could not totally explain the phytoplankton  
4 diminution: it was necessary to introduce the effect of grazing by a benthic lamellibranch,  
5 *Corbicula fluminea*. The role played by this invasive clam definitely needs to be assessed, as  
6 it has propagated dramatically in the Loire Basin since 1990 (Brancotte and Vincent, 2002)  
7 like it did in some other European rivers with significant impacts on the phytoplankton  
8 biomass (Hardenbicker et al., 2014; Pigneur et al., 2014). Orthophosphate series were  
9 sometimes poorly explained by the DHR model, giving an indication of its time variability. In  
10 summertime, very little augmentations in water discharge could refill the system with more  
11 available phosphorus, allowing more phytoplankton developments, but this would only be  
12 seen at a fine temporal scale. We are here using monthly datasets, which is obviously not  
13 detailed enough to discuss variations we shall observe at the daily scale:  $\text{PO}_4^{3-}$  concentration  
14 is very much sensitive to TSS concentration and consequently to water discharge variations.

15 The relationship between the *winter* nitrate levels and the percentage of the catchment under  
16 arable land is strong (Fig. 2), illustrating the fact that nitrate levels originate mainly from  
17 diffuse agricultural sources. The slightly increasing trend in nitrate could partly be explained  
18 by the delayed response of the environment to external changes (Behrendt et al., 2002;  
19 Howden et al., 2010), or, according to Bouraoui and Grizzetti (2008), this could be showing a  
20 lack of appropriate agro-environmental methods, or a delay in implementing the 1991  
21 European Nitrates Directive. It has been shown that mitigation measures in agriculture did  
22 decrease nitrogen loads in several Swedish rivers (Grimvall et al., 2014) and in the Rhine and  
23 Danube Rivers (Hartmann et al., 2007) making a great contrast with many other temperate  
24 lowland rivers where nitrate increasing trends are still recorded: the Mississippi (Sprague et  
25 al., 2011), Ebro, Po and Rhone Rivers (Ludwig et al., 2009) and also the Thames (Howden et  
26 al., 2010). Another potential reason for this increase could be climate change: higher  
27 mineralization of organic matter in the arable soils is expected and caused by an increased  
28 temperature over time (Arheimer et al., 2005) together with higher soil mineralization  
29 (Bouraoui et al., 2002). This thesis would seem reasonably concomitant with the rising water  
30 temperature which was recorded in the Loire River (Fig 4f), but it seems too early to fully  
31 determine the link between climate change and nitrate trends.

1 Such diffuse N sources are seasonal and this depends on leaching of bare soils by rainfall in  
2 winter and retention by vegetation in the growing season. Thus, it is possible that the decrease  
3 of nitrate seasonal amplitude recorded since 2005 was linked to lower discharge variations,  
4 however the increasing nitrate trend contradicts with a slightly descending discharge trend.  
5 Figure 3 clearly indicates the antivaration of phytoplankton and nitrate in their seasonal  
6 cycle: nitrate minima were reached when Chl. *a* concentrations were maximum, i.e. in  
7 *summer* in the Middle and Lower sectors. In addition, increasing of nitrate concentration have  
8 been seen in *summer* in the Middle and Lower sectors (see section 4.2), which is concomitant  
9 with a reduced phytoplankton biomass. However, seasonal amplitudes of nitrate did not  
10 decrease significantly in the Middle Loire while the decline of phytoplanktonic biomass  
11 started since the 1990s and was generalized to the whole basin since 2002 (section 4.3).  
12 Hence, it is likely that N uptake by phytoplankton had only a minor influence on nitrate  
13 seasonal variations. Denitrification could play a significant role on nitrate seasonal variations,  
14 like in the neighboring Seine basin (Curie et al., 2011), but further investigations would be  
15 needed to fully assess the processes involved. A complete N budget in the watershed plus the  
16 development of a N-surplus model would better explain why nitrate levels remain this high in  
17 the Loire Basin.

## 18 **5.2 Nutrient limitation variation since 1980**

19 The N:P molar ratio allows to determine whether the system studied is potentially under  
20 nitrate or phosphate limitation (Koerselman and Meuleman, 1996; Ludwig et al., 2009) and  
21 may constitute the basis of some indicators to assess the risk of eutrophication in freshwaters  
22 (Dupas et al., 2015). Given other controlling factors as non-limiting factors of phytoplankton  
23 growth, if N:P is under 14, the system is limited by N; over 16, it is considered P-limited. In-  
24 between, N and P availabilities might be sufficient or the ecosystem might be co-limited by N  
25 and P (Koerselman and Meuleman, 1996).

26 In the Loire River, a slight increase in annual concentrations of nitrogen during the last 30  
27 years while phosphorus inputs decreased greatly resulted in the modification of the N:P molar  
28 ratio (Fig. 5). In the Middle Loire, the annual average ratio kept on increasing since 1980. In  
29 *summer* during the 1980s, the lowest values observed were occasionally under the Redfield  
30 limitation but most of the time over the limitation. Since 1992, the system never reached  
31 again the Redfield limit and remained in the P-limitation domain as a result of reducing  
32 significantly phosphorus direct inputs. Similar variations were observed in other river systems

1 (e.g. the Ebro, Rhone, Po, Danube, Ludwig et al., 2009; the Seine, Billen and Garnier, 2007;  
2 the Mississippi, Turner et al., 2003) where similar trends in N and P were recorded. The N:P  
3 ratio was subjected to a significant seasonality. Its pattern and strength has changed from low  
4 seasonal variations during the 1980s and a minimum reached in *summer* to a well-defined  
5 seasonality since 2002 in the Middle and Lower reaches with a maximum reached in *summer*,  
6 reinforcing the P-limitation characteristic of the Loire River during the phytoplanktonic  
7 growth period. These results indicate that P-limitation of phytoplankton growth has become a  
8 significant factor. When the river hydrology remains stable in the summer, phytoplankton is  
9 potentially under P-limitation. This is suggesting a potential explanation for the apparent  
10 shift in seasonal phases of Chl. *a* concentrations (late summer blooms no longer occur,  
11 described in section 3.2): in those cases, the P-limitation is reached before any other  
12 limitation. This shift could also correspond to a significant grazing by invasive *Corbicula*  
13 clams species which would abate significantly the phytoplankton biomass (Pigneur et al.,  
14 2014).

### 15 **5.3 Daily O<sub>2</sub> and pH amplitudes as indicators of eutrophication mitigation**

16 The *delta O<sub>2</sub>* and *delta pH* seasonal amplitudes decreased greatly since 1990: around 3.5 mg  
17 L<sup>-1</sup> in 1990-95, *delta O<sub>2</sub>* amplitude declined down to 1.25 mg L<sup>-1</sup>. Similarly, from a seasonal  
18 amplitude at 0.25 pH unit, *delta pH* seasonal amplitude was maximum in 1998 (0.35) and  
19 went down to 0.3 since 2007. These descending trends are linked to the apparent decrease of  
20 phytoplanktonic biomass: the seasonal amplitude of Chl. *a* concentrations explained 80% of  
21 the seasonal variations of *delta O<sub>2</sub>* and only 59% for *delta pH* amplitudes. Continuous records  
22 of O<sub>2</sub> and pH take into account the whole in-stream primary activity, that is to say not only  
23 the phytoplankton respiration but also macrophytes and periphyton activities. While Chl. *a*  
24 concentrations kept on declining since 1991, *delta O<sub>2</sub>* and *delta pH* stopped decreasing  
25 suggesting that a non-phytoplanktonic activity was rising. Besides, one would expect that  
26 since phytoplankton biomass declined, water column irradiance increased and macrophyte  
27 abundance would have risen. We unfortunately lack data about macrophyte and periphyton  
28 developments in the Loire River, but the biological reserve Saint-Mesmin located near  
29 Orléans City (station 15) studied the development of macrophytes species since 1998 on 24  
30 river sections (60 m long by 5m width) and showed the increasing abundance and biodiversity  
31 of such aquatic plants since 2002. Two species were dominant, *Myriophyllum spicatum* and  
32 *Ranunculus fluitans*. The role played by fixed aquatic vegetation on the river biogeochemistry



1 is probably very significant as macrophytes are known to get nutrients contained in the water  
2 compartment as well as in the sediments (Carignan and Kalff, 1980; Hood, 2012). Hence,  
3 during low  $\text{PO}_4^{3-}$  concentration in *summer*, macrophyte growth is not limited by the in-stream  
4 nutrient limitation.

5 A major change occurred in the seasonal patterns of daily maximum of dissolved  $\text{O}_2$ . From a  
6 maximum reached in June or July at least between 1990 and 2001, the seasonal pattern of  
7 daily maximum shifted dramatically to a maximum reached in *winter*. On the contrary, daily  
8  $\text{O}_2$  and pH minimum always reached their maximum in *winter* and their minimum in *summer*  
9 (due to biomass respiration). Such a spectacular change in daily  $\text{O}_2$  maximum because of a  
10 declining eutrophication has never been shown in other major European rivers.

11 When unusual late floods occurred, higher flow velocity, increased turbidity and a reduced  
12 water column irradiance probably disrupted the well-established dominance of  
13 production/respiration cycles. Therefore both dissolved oxygen and pH levels dropped for a  
14 few days. Such episodes happened in 1992 (event described in Moatar et al., 2001), 1998 and  
15 2008. In those cases, phytoplankton growth is under hydrologic limitation.

## 16 **6 Conclusions**

17 The Loire River is a relevant case of a river recovering from severe eutrophication by  
18 controlling phosphorus direct inputs. However, other recent changes should also be  
19 considered. For example, it would be interesting to investigate the impact of the development  
20 of *Corbicula* clams (Brancotte and Vincent, 2002) on the biogeochemistry of the Loire basin  
21 surface waters. A potential numerical model of the Loire basin eutrophication should not only  
22 take into account climate and land-use changes, but also recent ecological changes (Descy et  
23 al., 2011; Pigneur et al., 2014) and this model would probably be able to answer many  
24 questions about the occurrence of invasive grazers in the Loire River.

25 This study highlighted how contrasted can be the different long term trajectories of Chl. *a* and  
26 nutrient concentrations in the different reaches of a eutrophic river and contributed to better  
27 understand the current biogeochemical functioning. Although the Upper Loire received the  
28 highest concentrations of phosphorus, the signs of eutrophication were expressed only in the  
29 lowest part of the Upper River because of its morphology. The Middle Loire is very favorable  
30 to eutrophication and the Lower reach functioning and trends remained close to the Middle  
31 Loire trajectory although it receives most of the tributaries inputs. Signs of eutrophication

1 remained lower in the major tributaries than the main river stem, but it has been shown that  
2 their influence on the Loire River nutrient fluxes (and consequently on the phytoplanktonic  
3 biomass) at the confluences can reach up to 35%.

4 This study also support the previous works on the Loire eutrophication, but the analysis of the  
5 long term changes in seasonality in this paper could bring more elements:

6 i) Controlling P-inputs also impacted the river biogeochemistry at the seasonal scale:  
7 seasonal amplitudes of Chl. *a* and orthophosphate greatly decreased; and this  
8 impacted O<sub>2</sub> and pH both daily and seasonally. However, nitrate amplitudes  
9 remained quite stable, evidencing the fact that phytoplankton growth had a minor  
10 influence on nitrate seasonal variations questioning the exact role played by fixed  
11 aquatic vegetation and denitrification on the nitrogen cycle.

12 ii) When hydrologic conditions remain favorable for phytoplankton growth in *summer*,  
13 orthophosphate concentration becomes the limiting factor.

14 iii) Combined to Chl. *a* concentration time series, *delta O<sub>2</sub>* and *delta pH* are relevant  
15 metrics for studying eutrophication variations. High frequency records of Chl. *a*,  
16 O<sub>2</sub> and pH could potentially enable the separation between phytoplankton and  
17 macrophytes impacts on the river biogeochemistry.

18 In addition, this study highlights the temporal variability of the different eutrophication  
19 metrics: in *summer*, the river biogeochemistry is essentially controlled by  
20 production/respiration processes. Thus, daily and seasonal variations are very significant and  
21 call into question the classical monthly survey recommended by national or international  
22 authorities.

23

## 24 **Acknowledgements**

25 This study started within the VARIQUAL national program financed by ANR (2008-2010)  
26 and continued with the help of the Plan Loire Grandeur Nature, European funds FEDER and  
27 AELB (2013-2014, Eutrophication Trends Project). The authors are grateful to A. Poirel  
28 (EdF) for providing access to water quality records at Nuclear Power Plant stations and to M.  
29 Chanterau (Réserve St Mesmin) for communicating macrophytes details and data. Comments  
30 and suggestions by O. Coulon (AELB), Ph. Gosse (EdF), G. Pinay (Rennes 1 Univ.) and by  
31 the Biogeosciences Referees and Editor significantly improved the manuscript.

## 1 **References**

- 2 Abonyi, A., Leitão, M., Lançon, A. M. and Padisák, J.: Phytoplankton functional groups as  
3 indicators of human impacts along the River Loire (France), *Hydrobiologia*, 689, 233–249,  
4 doi:10.1007/s10750-012-1130-0, 2012.
- 5 Abonyi, A., Leitão, M., Stanković, I., Borics, G., Várbíró, G. and Padisák, J.: A large river  
6 (River Loire, France) survey to compare phytoplankton functional approaches: Do they  
7 display river zones in similar ways?, *Ecol. Indic.*, 46, 11–22,  
8 doi:10.1016/j.ecolind.2014.05.038, 2014.
- 9 Aleya, L., Desmolles, F., Michard, M., Bonnet, M.-P. and Devaux, J.: The deterministic  
10 factors of the *Microcystis aeruginosa* blooms over a biyearly survey in the hypereutrophic  
11 reservoir of Villerest (Roanne, France)., *Arch. Hydrobiol.*, 120, 489–515, 1994.
- 12 Arheimer, B., Andréasson, J., Fogelberg, S., Johnsson, H., Pers, C. B. and Persson, K.:  
13 Climate change impact on water quality: model results from southern Sweden., *Ambio*, 34,  
14 559–566, 2005.
- 15 Barlocher, F., Seena, S., Wilson, K. P. and Williams, D. D.: Raised water temperature lowers  
16 diversity of hyporheic aquatic hyphomycetes, *Freshw. Biol.*, 53, 368–379,  
17 doi:10.1111/j.1365-2427.2007.01899.x, 2008.
- 18 Behrendt, H., Kornmilch, M., Opitz, D., Schmoll, O. and Scholz, G.: Estimation of the  
19 nutrient inputs into river systems - experiences from German rivers, *Reg. Environ. Chang.*, 3,  
20 107–117, doi:10.1007/s10113-002-0042-3, 2002.
- 21 Billen, G. and Garnier, J.: River basin nutrient delivery to the coastal sea: Assessing its  
22 potential to sustain new production of non-siliceous algae, *Mar. Chem.*, 106, 148–160,  
23 doi:10.1016/j.marchem.2006.12.017, 2007.
- 24 Billen, G., Garnier, J., Némery, J., Sebilo, M., Sferratore, A., Barles, S., Benoit, P. and  
25 Benoît, M.: A long-term view of nutrient transfers through the Seine river continuum., *Sci.*  
26 *Total Environ.*, 375, 80–97, doi:10.1016/j.scitotenv.2006.12.005, 2007.
- 27 Bouraoui, F., Galbiati, L. and Bidoglio, G.: Climate change impacts on nutrient loads in the  
28 Yorkshire Ouse catchment (UK), *Hydrol. Earth Syst. Sci.*, 6, 197–209, 2002.
- 29 Bouraoui, F. and Grizzetti, B.: An integrated modelling framework to estimate the fate of  
30 nutrients: Application to the Loire (France), *Ecol. Modell.*, 212, 450–459,  
31 doi:10.1016/j.ecolmodel.2007.10.037, 2008.
- 32 Bouraoui, F. and Grizzetti, B.: Long term change of nutrient concentrations of rivers  
33 discharging in European seas., *Sci. Total Environ.*, 409, 4899–916,  
34 doi:10.1016/j.scitotenv.2011.08.015, 2011.
- 35 Brancotte, V. and Vincent, T.: L'invasion du réseau hydrographique français par les  
36 mollusques *Corbicula* Spp. Modalité de colonisation et rôle prépondérant des canaux de  
37 navigation., *B. Fr. Peche Piscic.*, 325–337, 2002 (in French).

- 1 Bustillo, V., Moatar, F., Ducharne, A., Thiéry, D. and Poirel, A.: A multimodel comparison  
2 for assessing water temperatures under changing climate conditions via the equilibrium  
3 temperatures concept: case study of the Middle Loire River, France, *Hydrol. Process.*, 28,  
4 1507-1524, doi:10.1002/hyp.9683, 2013.
- 5 Carignan, R. and Kalff, J.: Phosphorus sources for aquatic weeds: water or sediments?,  
6 *Science*, 207, 987–989, doi:10.1126/science.207.4434.987, 1980.
- 7 CLC2006: Corine Land Cover, available at: <http://clc.developpement-durable.gouv.fr/>, last  
8 access: 20 September 2014, 2006.
- 9 Crouzet, P.: L'eutrophisation de la Loire, *Water Supply*, 1, 131–144, 1983 (in French).
- 10 Curie, F., Ducharne, A., Bendjoudi, H. and Billen, G.: Spatialization of denitrification by  
11 river corridors in regional-scale watersheds: Case study of the Seine river basin, *Phys. Chem.  
12 Earth, Parts A/B/C*, 36, 530–538, doi:10.1016/j.pce.2009.02.004, 2011.
- 13 Descy, J.-P., Leitão, M., Everbecq, E., Smitz, J. S. and Deliege, J.-F.: Phytoplankton of the  
14 River Loire, France: a biodiversity and modelling study, *J. Plankton Res.*, 34, 120–135,  
15 doi:10.1093/plankt/fbr085, 2011.
- 16 Dessery, S., Dulac, C., Laurenceau, J. M. and Meybeck, M.: Evolution du carbone organique  
17 «algal» et «détritique» dans trois rivières du bassin parisien., *Arch. Hydrobiol.*, 100, 235–260,  
18 1984 (in French).
- 19 Dupas, R., Delmas, M., Dorioz, J., Garnier, J., Moatar F. and Gascuel-Oudou C.: Assessing  
20 the impact of agricultural pressures on N and P loads and eutrophication risk, *Ecol. Indic.*, 48,  
21 396–407, doi:10.1016/j.ecolind.2014.08.007, 2015.
- 22 Durance, I. and Ormerod, S. J.: Evidence for the role of climate in the local extinction of a  
23 cool-water triclade, *J. North Am. Benthol. Soc.*, 29, 1367–1378, doi:10.1899/09-159.1, 2010.
- 24 EEC: Council Directive 91/676/EEC Concerning the Protection of Waters Against Pollution  
25 Caused by Nitrates from Agricultural Sources, L 375, Official Journal of the European  
26 Community, Brussels, 1991a.
- 27 EEC: Council Directive 91/271/EEC Concerning Urban Waste Water Treatment, L 135,  
28 Official Journal of the European Community, Brussels, 1991b.
- 29 Etcheber, H., Taillez, A., Abril, G., Garnier, J., Servais, P., Moatar, F. and Commarieu, M.-  
30 V.: Particulate organic carbon in the estuarine turbidity maxima of the Gironde, Loire and  
31 Seine estuaries: origin and lability, *Hydrobiologia*, 588, 245–259, doi:10.1007/s10750-007-  
32 0667-9, 2007.
- 33 Flourey, M., Delattre, C., Ormerod, S. J. and Souchon, Y.: Global versus local change effects  
34 on a large European river., *Sci. Total Environ.*, 441, 220–229,  
35 doi:10.1016/j.scitotenv.2012.09.051, 2012.

- 1 Gosse, P., Gailhard, J. and Hendrickx, F.: Analyse de la température de la Loire moyenne en  
2 été sur la période 1949 à 2003, *Hydroécologie Appliquée*, 16, 233–274, 2008 (in French).
- 3 Gosse, P., Lascombe, C. and Salleron, J. L.: Modélisation de la qualité de grands cours d'eau  
4 Eutrophisation : cas de la Moselle et du Doubs, *La Houille Blanche*, 3-4, 187–194,  
5 doi:10.1051/lhb/1990010, 1990 (in French).
- 6 Grimvall, A., Von Brömssen, C. and Lindström, G.: Using process-based models to filter out  
7 natural variability in observed concentrations of nitrogen and phosphorus in river water,  
8 *Environ. Monit. Assess.*, 186, 5135–5152, doi:10.1007/s10661-014-3765-y, 2014.
- 9 Grizzetti, B., Bouraoui, F. and Aloe, A.: Changes of nitrogen and phosphorus loads to  
10 European seas, *Glob. Chang. Biol.*, 18, 769–782, doi:10.1111/j.1365-2486.2011.02576.x,  
11 2012.
- 12 Grosbois, C., Négrel, P., Grimaud, D. and Fouillac, C.: An Overview of Dissolved and  
13 Suspended Matter Fluxes in the Loire River Basin : Natural and Anthropogenic Inputs, *Aquat.  
14 Geochemistry*, 7, 81–105, 2001.
- 15 Halliday, S. J., Skeffington, R. a., Wade, a. J., Neal, C., Reynolds, B., Norris, D. and  
16 Kirchner, J. W.: Upland streamwater nitrate dynamics across decadal to sub-daily timescales:  
17 a case study of Plynlimon, Wales, *Biogeosciences*, 10, 8013–8038, doi:10.5194/bg-10-8013-  
18 2013, 2013.
- 19 Halliday, S. J., Wade, A. J., Skeffington, R. a, Neal, C., Reynolds, B., Rowland, P., Neal, M.  
20 and Norris, D.: An analysis of long-term trends, seasonality and short-term dynamics in water  
21 quality data from Plynlimon, Wales., *Sci. Total Environ.*, 434, 186–200,  
22 doi:10.1016/j.scitotenv.2011.10.052, 2012.
- 23 Hardenbicker, P., Rolinski, S., Weitere, M. and Fischer, H.: Contrasting long-term trends and  
24 shifts in phytoplankton dynamics in two large rivers, *Int. Rev. Hydrobiol.*, 99, 326–334,  
25 doi:10.1002/iroh.201301680, 2014.
- 26 Hartmann, J., Jansen, N., Kempe, S. and Dürr, H. H.: Geochemistry of the river Rhine and the  
27 Upper Danube : recent trends and lithological influence on baselines, *J. Environ. Sci. Sustain.  
28 Soc.*, 1, 39–46, 2007.
- 29 Hilton, J., O'Hare, M., Bowes, M. J. and Jones, J. I.: How green is my river? A new paradigm  
30 of eutrophication in rivers., *Sci. Total Environ.*, 365, 66–83,  
31 doi:10.1016/j.scitotenv.2006.02.055, 2006.
- 32 Hood, J.: The role of submersed macrophytes in river eutrophication and biogeochemical  
33 nutrient cycling, Ph.D. thesis, University of Waterloo, Canada 2012.
- 34 Howden, N. J. K., Burt, T. P., Worrall, F., Whelan, M. J. and Bieroza, M.: Nitrate  
35 concentrations and fluxes in the River Thames over 140 years (1868-2008): are increases  
36 irreversible?, *Hydrol. Process.*, 24, 2657–2662, doi:10.1002/hyp.7835, 2010.

- 1 INSEE: Annuaire Statistique de la France. Institut National de Statistiques et d' Etudes  
2 Economiques, Paris, available at: <http://www.insee.fr/fr/bases-de-donnees/>, last access: 20  
3 September 2014, 2008.
- 4 Istvánovics, V. and Honti, M.: Efficiency of nutrient management in controlling  
5 eutrophication of running waters in the Middle Danube Basin, *Hydrobiologia*, 686, 55–71,  
6 2012.
- 7 Istvánovics, V., Honti, M., Kovács, Á., Kocsis, G. and Stier, I.: Phytoplankton growth in  
8 relation to network topology: time-averaged catchment-scale modelling in a large lowland  
9 river, *Freshw. Biol.*, 59, 1856–1871, doi:10.1111/fwb.12388, 2014.
- 10 Jackson, B. M., Browne, C. a., Butler, A. P., Peach, D., Wade, A. J. and Wheater, H. S.:  
11 Nitrate transport in Chalk catchments: monitoring, modelling and policy implications,  
12 *Environ. Sci. Policy*, 11, 125–135, doi:10.1016/j.envsci.2007.10.006, 2008.
- 13 Jugnia, L., Debroas, D., Romagoux, J. and Dévaux, J.: Initial results of remediation activities  
14 to restore hypereutrophic Villerest Reservoir ( Roanne , France ), *Lakes Reserv. Res. Manag.*,  
15 9, 109–117, 2004.
- 16 Koerselman, W. and Meuleman, A. F. M.: The vegetation N : P ratio : a new tool to detect the  
17 nature of nutrient limitation, *J. Appl. Ecol.*, 33, 1441–1450, 1996.
- 18 Krogstad, T. and Lovstad, O.: Erosion, phosphorus and phytoplankton response in rivers of  
19 South-Eastern Norway, *Hydrobiologia*, 183, 33–41, 1989.
- 20 Lair, N. and Reyes-Marchant, P.: The potamoplankton of the Middle Loire and the role of the  
21 “moving littoral ” in downstream transfer of algae and rotifers, *Hydrobiologia*, 356, 33–52,  
22 1997.
- 23 Latapie, A.: Modelisation de l'évolution morphologique d'un lit alluvial : application à la  
24 Loire moyenne, Ph.D. thesis, Université François Rabelais de Tours, 2011 (in French).
- 25 Latapie, A., Camenen, B., Rodrigues, S., Paquier, A., Bouchard, J. P. and Moatar, F.:  
26 Assessing channel response of a long river influenced by human disturbance, *CATENA*, 121,  
27 1–12, doi:10.1016/j.catena.2014.04.017, 2014.
- 28 Lecerf, A., Risnoveanu, G., Popescu, C., Gessner, M. and Chauvet, E.: Decomposition of  
29 diverse litter mixtures in streams., *Ecology*, 88, 219–227, 2007.
- 30 Lehmann, A. and Rode, M.: Long-term behaviour and cross-correlation water quality analysis  
31 of the River Elbe, Germany., *Water Res.*, 35, 2153–60, 2001.
- 32 Leitão, M. and Lepretre, A.: The phytoplankton of the River Loire, France: a typological  
33 approach., *Verhandlungen der Internationale Vereinigung für theoretische und angewandte*  
34 *Limnologie*, 26, 1050–1056, 1998.
- 35 Ludwig, W., Dumont, E., Meybeck, M. and Heussner, S.: River discharges of water and  
36 nutrients to the Mediterranean and Black Sea: Major drivers for ecosystem changes during

- 1 past and future decades?, *Prog. Oceanogr.*, 80, 199–217, doi:10.1016/j.pocean.2009.02.001,  
2 2009.
- 3 Meybeck, M., Cauwet, G., Dessery, S., Somville, M., Gouleau, D. and Billen, G.: Nutrients  
4 (organic C, P, N, Si) in the eutrophic River Loire (France) and its estuary, *Estuar. Coast. Shelf*  
5 *S.*, 27, 595–624, 1988.
- 6 Moatar, F. and Gailhard, J.: Water temperature behaviour in the River Loire since 1976 and  
7 1881, *Comptes Rendus Geosci.*, 338, 319–328, doi:10.1016/j.crte.2006.02.011, 2006.
- 8 Moatar, F. and Meybeck, M.: Compared performances of different algorithms for estimating  
9 annual nutrient loads discharged by the eutrophic River Loire, *Hydrol. Process.*, 19, 429–444,  
10 doi:10.1002/hyp.5541, 2005.
- 11 Moatar, F., Meybeck, M., Raymond, S., Birgand, F. and Curie, F.: River flux uncertainties  
12 predicted by hydrological variability and riverine material behaviour, *Hydrol. Process.*, 27,  
13 3535–3546, doi:10.1002/hyp.9464, 2013.
- 14 Moatar, F., Miquel, J. and Poirel, A.: A quality-control method for physical and chemical  
15 monitoring data. Application to dissolved oxygen levels in the river Loire (France), *J.*  
16 *Hydrol.*, 252, 25–36, doi:10.1016/S0022-1694(01)00439-5, 2001.
- 17 Oudin, L. C.: Modélisation de l'eutrophisation en Loire, *Houille Blanche*, 3, 195–206, 1990.
- 18 Oudin, L. C., Lair, N., Leitão, M., Reyes-Marchant, P., Mignot, J.-F., Steinbach, P., Vigneron,  
19 T., Berton, J.-P., Bacchi, M., Roché, J. E. and Descy, J.-P.: The Loire Basin, in *Rivers of*  
20 *Europe* (Eds Tockner K. & C.T. Robinson). British Library, London., 2009.
- 21 Pedregal, D. J. and Trapero, J. R.: Electricity prices forecasting by automatic dynamic  
22 harmonic regression models, *Energ. Convers. Manage.*, 48, 1710–1719,  
23 doi:10.1016/j.enconman.2006.11.004, 2007.
- 24 Pigneur, L.-M., Falisse, E., Roland, K., Everbecq, E., Deliège, J.-F., Smitz, J. S., Van  
25 Doninck, K. and Descy, J.-P.: Impact of invasive Asian clams, *Corbicula* spp., on a large river  
26 ecosystem, *Freshw. Biol.*, 59, 573–583, doi:10.1111/fwb.12286, 2014.
- 27 Reynolds, C. S. and Descy, J.-P.: The production, biomass and structure of phytoplankton in  
28 large rivers, *Arch. Hydrobiol. Suppl.*, 113, 161–187, 1996.
- 29 Reynolds, C. S., Descy, J.-P. and Padisdak, J.: Are phytoplankton dynamics in rivers so  
30 different from those in shallow lakes ?, *Hydrobiologia*, 289, 1–7, 1994.
- 31 Romero, E., Garnier, J., Lassaletta, L., Billen, G., Gendre, R., Riou, P. and Cugier, P.: Large-  
32 scale patterns of river inputs in southwestern Europe: seasonal and interannual variations and  
33 potential eutrophication effects at the coastal zone, *Biogeochemistry*, 113, 481–505,  
34 doi:10.1007/s10533-012-9778-0, 2012.

- 1 Sprague, L. A., Hirsch, R. M. and Aulenbach, B. T.: Nitrate in the Mississippi River and Its  
2 Tributaries , 1980 to 2008 : Are We Making Progress ?, *Environ. Sci. Technol.*, 45, 7209–  
3 7216, 2011.
- 4 Smith, V. H. and Schindler, D. W.: Eutrophication science: where do we go from here?,  
5 *Trends Ecol. Evol.*, 24, 201–7, doi:10.1016/j.tree.2008.11.009, 2009.
- 6 Smith, V. H., Tilman, G. D. and Nekola, J. C.: Eutrophication: impacts of excess nutrient  
7 inputs on freshwater, marine, and terrestrial ecosystems., *Environ. Pollut.*, 100, 179–96, 1999.
- 8 Taylor, C., Pedregal, D., Young, P. and Tych, W.: Environmental time series analysis and  
9 forecasting with the Captain toolbox, *Environ. Model. Softw.*, 22, 797–814,  
10 doi:10.1016/j.envsoft.2006.03.002, 2007.
- 11 Turner, R. E., Rabalais, N. N., Justic, D. and Dortch, Q.: Global patterns of dissolved N , P  
12 and Si in large rivers, *Biogeochemistry*, 64, 297–317, 2003.
- 13 Vicaud, A.: Les besoins en eau de refroidissement des centrales thermiques de production  
14 d'électricité, *La Houille Blanche*, 6, 34–40, doi:10.1051/lhb:2008069, 2008 (in French).
- 15 Whitehead, P. G., Wilby, R., Battarbee, R., Kernan, M. and Wade, A.: A review of the  
16 potential impacts of climate change on surface water quality, *Hydrol. Sci. J.*, 54, 101–123,  
17 doi:10.1623/hysj.54.1.101, 2009.
- 18



1 Table 1. Loire main stem stations characteristics. Kilometric point (KP): distance from  
 2 headwaters; Drained area; Q: average annual discharge; population density in 2008; arable  
 3 land as percentage of the drained catchment; API: agricultural pressure indicator = (pasture +  
 4 forest) / (pasture + forest + arable land) expressed in percentage. See paragraph 1.2. for source  
 5 information.

	Upper Loire								Middle Loire							Lower	
Station	1	2	3	4	5	6	7	8	9	10	11	13	15	17	18	20	21
KP (km)	44	92	200	224	273	292	344	417	451	465	500	564	633	712	772	822	895
Drained area (10 <sup>3</sup> km <sup>2</sup> )	0.5	1	4	5	7	8	13	15	18	33	34	36	37	41	43	82	109
Q (m <sup>3</sup> s <sup>-1</sup> )	6	10	-	47	67	-	89	-	180	300	320	327	-	360	366	680	850
Population density (inhab.km <sup>-2</sup> )	13	50	144	143	122	128	101	91	80	75	74	74	73	80	83	-	73
Arable land (%)	0.6	3	1	3	4	4	3	4	6	9	11	13	13	15	17	24	30
API (%)	99	97	99	96	96	96	96	96	93	90	89	87	86	84	82	75	69

6

1 Table 2. Major tributaries station characteristics.

Station	A	B	C	D	E
Drained area ( $10^3 \text{ km}^2$ )	14	13	33	21	22
Average discharge ( $\text{m}^3 \text{ s}^{-1}$ )	143	81	37	-	135
Population density (inhab. $\text{km}^{-2}$ )	67	52	76	55	82
Arable land (%)	13	36	52	25	49
API (%)	87	63	46	74	50

2

1 Table 3. Long term trends at three stations representative of the Upper, Middle and Lower  
 2 Loire.

	years	Annual median			Trend			Significance of trend 1980-2012 (%)		
		Chl. <i>a</i> µg L <sup>-1</sup>	PO <sub>4</sub> <sup>3-</sup> µg P L <sup>-1</sup>	NO <sub>3</sub> <sup>-</sup> mg N L <sup>-1</sup>	Chl. <i>a</i> µg L <sup>-1</sup> y <sup>-1</sup>	PO <sub>4</sub> <sup>3-</sup> µg P L <sup>-1</sup> y <sup>-1</sup>	NO <sub>3</sub> <sup>-</sup> mg N L <sup>-1</sup> y <sup>-1</sup>	Chl. <i>a</i>	PO <sub>4</sub> <sup>3-</sup>	NO <sub>3</sub> <sup>-</sup>
Upper Loire	80-89	9	183	1.4	+2	+16	0.0			
Station 4	90-01	12	169	1.8	0	-16	0.0	74	87	77
	02-12	11	88	1.4	-1	-3	0.0			
Middle Loire	80-89	47	121	1.8	+3	-6	0.0			
Station 18	90-01	83	58	1.9	-5	-3	+0.1	82	91	53
	02-12	17	26	2.2	-5	-2	0.0			
Lower Loire	80-89	50	79	2.5	+5	+12	+0.3			
Station 21	90-01	58	89	3.3	-9	-3	+0.1	83	76	71
	02-12	14	37	2.6	-4	-5	-0.1			

3

4

1 Table 4. Seasonality analysis and changes since 1980 at three stations representative of the  
 2 Upper, Middle and Lower Loire.


	years	Seasonal amplitude			Significance (%)			Amplitude trend		
		Chl. <i>a</i> $\mu\text{g L}^{-1}$	$\text{PO}_4^{3-}$ $\mu\text{g P L}^{-1}$	$\text{NO}_3^-$ $\text{mg N L}^{-1}$	Chl. <i>a</i>	$\text{PO}_4^{3-}$	$\text{NO}_3^-$	Chl. <i>a</i> $\mu\text{g L}^{-1} \text{y}^{-1}$	$\text{PO}_4^{3-}$ $\mu\text{g P L}^{-1} \text{y}^{-1}$	$\text{NO}_3^-$ $\text{mg N L}^{-1} \text{y}^{-1}$
Upper Loire Station 4	80-89	61	101	0.7	41	16	25	0.0	-0.2	+0.1
	90-01	114	107	0.9	31	33	38	-0.2	+0.7	0.0
	02-12	17	26	2.2	24	41	42	-1.2	+0.5	+0.1
Middle Loire Station 18	80-89	182	123	2.2	61	44	80	+7.8	-5.6	+0.1
	90-01	152	71	2.8	64	43	85	-9.8	-3.7	0.0
	02-12	57	38	2.1	55	47	84	-8.1	-2.1	0.0
Lower Loire Station 21	80-89	184	125	3.2	68	46	78	-2.7	+2.0	+0.4
	90-01	82	120	5.5	52	62	81	-9.6	-1.6	-0.2
	02-12	53	65	3.2	62	51	85	-1.1	-10.4	+0.1

3

1 Table 5. Annual medians, DHR-model seasonal amplitudes and nutrients flux contributions of  
 2 the main tributaries.


	Annual median			Seasonal amplitude			Nutrient flux contribution		
	Chl. <i>a</i> μg L <sup>-1</sup>	PO <sub>4</sub> <sup>3-</sup> μg P L <sup>-1</sup>	NO <sub>3</sub> <sup>-</sup> mg N L <sup>-1</sup>	Chl. <i>a</i> μg L <sup>-1</sup>	PO <sub>4</sub> <sup>3-</sup> μg P L <sup>-1</sup>	NO <sub>3</sub> <sup>-</sup> mg N L <sup>-1</sup>	PO <sub>4</sub> <sup>3-</sup>	NO <sub>3</sub> <sup>-</sup>	
A	1980-89	20	124	1.4	85	180	1.7	54%	47%
	1990-01	23	83	1.5	134	112	2.1	44%	43%
	2002-12	17	51	1.7	83	74	2.3	42%	36%
B	1980-89	44	108	3.6	147	190	4.3	17%	32%
	1990-01	61	79	4.1	197	181	5.9	31%	37%
	2002-12	13	45	4.7	57	57	3.9	33%	33%
C	1980-89	28	166	4.0	104	234	3.7	-	-
	1990-01	44	90	4.2	109	144	5.0	-	-
	2002-12	16	59	4.6	37	79	4.2	-	-
D	1980-89	43	126	3.0	102	137	2.2	-	-
	1990-01	50	68	2.7	107	87	2.8	-	-
	2002-12	6	30	2.8	18	36	2.5	27%	35%
E	1980-89	50	191	4.0	142	326	4.2	38%	24%
	1990-01	62	181	4.4	132	236	8.1	33%	23%
	2002-12	21	73	4.1	51	102	5.6	35%	27%

3




1  
2  
3  
4  
5  
6  
7

Figure 1. Loire River Basin. Dark circles: sites of regulatory surveys. White circles: Nuclear Power Plants sampling sites. A to E: regulatory survey stations at tributaries outlets. G, V, N: three major dams, respectively Grangent, Villerest and Naussac. The estuarine influence begins downstream of station 21.



1  
2  
3  
4  
5  
6  
7  
8


Figure 2. Longitudinal profiles of *summer* median Chl. *a* (a), *winter* median  $\text{PO}_4^{3-}$  (b) and  $\text{NO}_3^-$  (c). Averages for three periods, in relation to % arable land (2006) and population density (2008) tested as eutrophication control variables. Uncertainty bars are due to sampling frequency. Arrows and capital letters (A to E) represent confluences with major tributaries (Fig. 1).



1  
2  
3  
4  
5  
6


Figure 3. Spatio-temporal diagrams of monthly median levels of Chl. *a* (a),  $\text{PO}_4^{3-}$  (b) and  $\text{NO}_3^-$  (c) during three periods along a longitudinal profile. Dotted vertical lines correspond to the monitoring stations (Fig.1).





1  
2  
3  
4  
5  
6  
7

Figure 4. Trends and seasonal components at station 18 of Chl. *a* (a), phosphate (b) and nitrate (c). Corresponding time series of monthly medians of both daily min and max of  $\text{O}_2$  (d) and pH (e) and their amplitude dynamics at station 19 (i.e.  $\text{delta O}_2$  and  $\text{delta pH}$ ). Water temperature trend at station 19 (f) and discharge trend at station 15 since 1980.



1

2

3 Figure 5. Variations of total nitrogen over total phosphorus molar ratios ranges during  
 4 *summer* and *winter* in the Middle Loire (station 18) since 1980 and compared to the Redfield  
 5 limit (dotted line). Each patch is composed at the bottom by the percentile 10% of the  
 6 recorded data and percentile 90% at the top, and y-axis is logarithmic.

7