

This discussion paper is/has been under review for the journal Biogeosciences (BG).
Please refer to the corresponding final paper in BG if available.

Identifying vital effects in *Halimeda* algae with Ca isotopes

C. L. Blättler^{1,*}, S. M. Stanley², G. M. Henderson¹, and H. C. Jenkyns¹

¹University of Oxford, Department of Earth Sciences, Oxford, UK

²University of Hawaii, Department of Geology & Geophysics, Honolulu, HI, USA

* now at: Princeton University, Department of Geosciences, Princeton, NJ, USA

Received: 21 February 2014 – Accepted: 21 February 2014 – Published: 5 March 2014

Correspondence to: C. L. Blättler (blattler@princeton.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

BGD

11, 3559–3580, 2014

Ca isotopes of
Halimeda algae

C. L. Blättler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Geochemical records of biogenic carbonates provide some of the most valuable records of the geological past, but are often difficult to interpret without a mechanistic understanding of growth processes. In this experimental study, *Halimeda* algae are used as a test organism to untangle some of the specific factors that influence their skeletal composition, in particular their Ca-isotope composition. Algae were stimulated to precipitate both calcite and aragonite by growth in artificial Cretaceous seawater. The Ca-isotope fractionation of the algal calcite is much smaller than that for the algal aragonite, similar to the behaviour observed in inorganic precipitates. However, the carbonate from *Halimeda* is isotopically heavier than inorganic forms, likely due to Rayleigh distillation within the algal intercellular space. In identifying specific vital effects and the magnitude of their influence on Ca-isotope ratios, this study suggests that mineralogy has a first-order control on the Ca-isotope budget of the carbonate sink and the Ca-isotope composition of seawater.

1 Introduction

This work investigates the mechanism of Ca-isotope fractionation in one of the major producers of aragonite sediment, *Halimeda* algae, to test models of biocalcification and the Ca-isotope cycle. Biogenic carbonate is the source of many important sedimentary records, which can be useful for interpreting ancient environments if the mechanisms of skeletal formation are understood. The elemental and isotopic compositions of biogenically precipitated minerals deviate from those of inorganic precipitates – differences sometimes referred to as “vital effects” – which complicate the application of geochemical proxies in skeletal material. Ca-isotope ratios, defined as $\delta^{44/40}\text{Ca} = \left[\left(\frac{^{44}\text{Ca}/^{40}\text{Ca}}{\text{sample}} \right) / \left(\frac{^{44}\text{Ca}/^{40}\text{Ca}}{\text{standard}} \right) - 1 \right] \cdot 1000$, may provide a tool for understanding the environmental factors affecting carbonate geochemistry and are

BGD

11, 3559–3580, 2014

Ca isotopes of *Halimeda* algae

C. L. Blättler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



themselves a tool to assess past environmental conditions. Ca is the dominant cation in biogenic calcite and aragonite and should be subject to similar vital effects as other cations. The experiments presented here use skeletal material from *Halimeda* to isolate particular aspects of the biocalcification process and quantify their effects on the Ca-isotope system.

Skeletal growth can be divided broadly into two categories, “biologically induced” and “biologically controlled”, reflecting the location of biomineralization, the transport of chemical species, and the influence of environmental conditions (Weiner and Dove, 2003). The green algae *Halimeda* spp. are recognized as an example of a biologically induced system, based on the crystallography and organization of their aragonite needles (Borowitzka and Larkum, 1987) and confirmed by observation of changing skeletal mineralogy under experimental conditions (Stanley et al., 2010). Their simplicity provides an ideal starting point for addressing the mechanisms of cation fractionation in biogenic aragonite.

Samples for this study were generated in a controlled *Halimeda* growth experiment, where the Mg/Ca ratio of seawater was changed to reflect hypothetical conditions during Earth history. The current molar Mg/Ca ratio of the oceans is 5.2, which generally favours the precipitation of aragonite and high-Mg calcite (cation composition > 4 mol% Mg) over low-Mg calcite (cation composition < 4 mol% Mg), subject to favourable temperature conditions, $p\text{CO}_2$, phosphate, or sulfate concentrations (e.g. Burton and Walter, 1987, 1990; Morse et al., 1997; Bots et al., 2011). Several lines of evidence, such as halite fluid inclusions, nonskeletal carbonate mineralogy, and fossil echinoderm Mg/Ca, suggest that seawater Mg/Ca may have been as low as 1.0–1.5 during periods such as the Cretaceous, which would have favoured the precipitation of low-Mg calcite (Sandberg, 1983; Hardie, 1996; Stanley and Hardie, 1998; Lowenstein et al., 2001; Horita et al., 2002; Dickson, 2004). Growth experiments on a variety of biocalcifiers under such low Mg/Ca conditions have produced changes in calcification rate and skeletal geochemistry (e.g. Stanley et al., 2002, 2005; Ries et al., 2006) and, in the case of *Halimeda incrassata*, a change in carbonate mineralogy from aragonite to cal-

BGD

11, 3559–3580, 2014

Ca isotopes of *Halimeda* algae

C. L. Blättler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



cite (Stanley et al., 2010). For this study, the *Halimeda* experiment was repeated with a close relative, *Halimeda discoidea*, to reproduce the unusual case of aragonite and calcite grown simultaneously under the same conditions. The growth of two different carbonate minerals from the same fluid presents a test case for the isotopic effects of carbonate mineral precipitation, modified by simple algal biology, with implications for the balance of the Ca-isotope budget.

2 Background to Ca-isotope fractionation

2.1 Inorganic Ca-isotope fractionation

The behaviour of Ca isotopes in inorganic carbonates has been explored in a variety of precipitation experiments, with all calcium carbonate minerals expressing fractionation in favour of the lighter isotopes of Ca. In isotope notation, this is expressed as $\alpha < 1$ (where α is the ratio of precipitate $^{44}\text{Ca}/^{40}\text{Ca}$ to solution $^{44}\text{Ca}/^{40}\text{Ca}$) or $\Delta^{44/40}\text{Ca} < 0$ (where $\Delta^{44/40}\text{Ca}$ is equal to $\delta^{44/40}\text{Ca}_{\text{carbonate}} - \delta^{44/40}\text{Ca}_{\text{solution}}$). Fractionation of laboratory-grown aragonite ($\Delta^{44/40}\text{Ca}$) has an average value of -1.7‰ at 15°C , while showing a small positive temperature dependence of 0.015‰ (Gussone et al., 2003). Smaller fractionations are generally observed during calcite growth, with samples precipitated in a beaker averaging -0.8‰ at 15°C , with a similar positive temperature dependence as for aragonite (Marriott et al., 2004). Much greater fractionations have been observed in laboratory speleothem-like calcite growth (Reynard et al., 2011), and relationships between $\Delta^{44/40}\text{Ca}$ and other factors such as growth rate and $[\text{CO}_3^{2-}]$ have also been identified (Lemarchand et al., 2004; Tang et al., 2008). Gussone et al. (2011) showed that the metastable CaCO_3 polymorph ikaite expresses smaller fractionations of -0.55‰ and -0.62‰ for experimental and natural samples, respectively, while laboratory-grown vaterite has an even smaller average fractionation of -0.36‰ . Measured fractionations of less than -0.2‰ for synthetic amorphous calcium carbonate (ACC) suggest that Ca-isotope ratios are not strongly affected in the

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



absence of a mineral structure (Gagnon et al., 2011), although the cause of isotopic offsets for the different carbonate minerals is not understood.

Models of Ca-isotope fractionation suggest that the preference for light isotopes in solid mineral phases is due to a kinetic effect associated with ion dehydration and attachment onto a surface, moderated by the kinetic fractionations of a backwards detachment reaction and ion transport effects (Gussone et al., 2003; DePaolo, 2011). The isotopic fractionation between fluid and carbonate in equilibrium appears to be close to zero, based on evidence from deep-sea carbonate sediment and pore fluids (Fantle and DePaolo, 2007) and from a long-lived carbonate aquifer (Jacobson and Holmden, 2008). These observations suggest that maximum fractionation occurs at intermediate growth rates when isotopically light surface-layer ions are incorporated into the crystal, and ion transport in the fluid is rapid enough not to limit the expression of the attachment fractionation (DePaolo, 2011). This model can explain the relationships between $\delta^{44/40}\text{Ca}$ and several growth variables as observed during inorganic precipitation experiments, but it may not translate directly to biological settings.

2.2 Biogenic Ca-isotope fractionation

In biological calcification systems, carbonate skeletons are also isotopically lighter than the fluid (usually seawater) from which they precipitate, but the degree of fractionation and responses to variable growth conditions differ from the inorganic data presented above, confirming the presence of vital effects. For example, temperature changes induce a range of responses in various organisms. Many forms of biogenic carbonate exhibit a similar temperature dependence to that of inorganic precipitates of 0.01–0.03‰ per °C (Gussone et al., 2003, 2005; Böhm et al., 2006; Gussone et al., 2006), although there are significant exceptions to this trend. The planktic foraminifera *G. sacculifer* expresses a much greater sensitivity of 0.22–0.24‰ per °C (Nägler et al., 2000; Hippler et al., 2006), although subsequent studies of this and other planktic foraminifera show a negligible temperature dependence (Chang et al., 2004; Sime et al., 2005; Griffith et al., 2008; Kasemann et al., 2008). A study of benthic foraminifera revealed a tem-

Ca isotopes of *Halimeda* algae

C. L. Blättler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



perature sensitivity of $\sim 0.02\text{‰}$ per $^{\circ}\text{C}$ for shells formed at temperatures $> 5^{\circ}\text{C}$, but significant deviations from this relationship at temperatures $< 5^{\circ}\text{C}$ (Gussone and Filipsson, 2010). These different responses in biocalcifiers, in contrast to the relatively consistent temperature sensitivity of inorganic carbonates, demonstrate that metabolic or physiological processes can contribute significantly to skeletal $\delta^{44/40}\text{Ca}$ and indicate the potential for Ca isotopes to provide information about the processes of biomineralization.

Isotopic vital effects are also observed in response to carbonate-ion concentration and growth rate. A strong correlation with both $[\text{CO}_3^{2-}]$ and growth rate was observed for inorganic calcite (Lemarchand et al., 2004), but no correlation was apparent with $[\text{CO}_3^{2-}]$ for *O. universa* (Gussone et al., 2005) nor several coccolithophorid species (Gussone et al., 2006, 2007; Langer et al., 2007). Growth rate was shown to have an effect on $\delta^{44/40}\text{Ca}$ in *G. siphonifera* (Kısakürek et al., 2011), but not for *E. huxleyi* (Langer et al., 2007). Observed variations in skeletal $\delta^{44/40}\text{Ca}$ in response to salinity may also be related to changes in growth rate, given that assemblages are usually adapted to local salinity conditions and are stressed by environmental changes (Gussone et al., 2009; Kısakürek et al., 2011). These studies suggest that Ca-isotope responses to $[\text{CO}_3^{2-}]$ and growth rate are determined by biological control over internal pH and carbonate chemistry, and are likely to vary according to specific biocalcification mechanisms.

Estimating the precise magnitude of vital effects in biocalcifiers is complicated by the additional isotopic fractionations associated with ion reservoirs and fractional utilization of Ca. The blue mussel, *Mytilus edulis*, demonstrates the significance of these effects by precipitating its skeleton from an extra-pallial fluid that has higher $\delta^{44/40}\text{Ca}$ than the surrounding seawater (Heinemann et al., 2008), producing a difference in $\delta^{44/40}\text{Ca}$ between its calcitic and aragonitic skeletal components of only 0.25‰ , rather than the expected 0.9‰ for an inorganic process. Transport effects and Rayleigh distillation have also been implicated in coccolith calcification (Gussone et al., 2006) and

BGD

11, 3559–3580, 2014

Ca isotopes of *Halimeda* algae

C. L. Blättler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



coral skeletal growth (Böhm et al., 2006). For foraminifera, conflicting estimates of the magnitude of these internal isotopic fractionations have been made, balancing their utilization of an internal pool of Ca against its compositional difference from seawater (Erez, 2003; Gussone et al., 2009). Griffith et al. (2008) developed a Ca-isotope model ($\alpha = 0.9985$, 85% utilization) suggesting that this internal reservoir could be offset by -0.8 to -0.9‰ from seawater, whereas Kısakürek et al. (2011) proposed that utilization of only 10–40% of the reservoir indicates a smaller offset of -0.2 to -0.4‰ , with additional fractionation originating from active Ca-pumping. Before proxy $\delta^{44/40}\text{Ca}$ measurements can be linked to environmental variables, the magnitudes and relative contributions of these competing biological and physico-chemical effects must be resolved.

3 Samples and methods

3.1 *Halimeda* experiment

This case study investigates the balance of isotopic effects linked to specific biocalcification mechanisms in the sample organism, *Halimeda discoidea*. Experiments were performed at the University of Hawai'i to induce the growth of both calcite and aragonite from the normally aragonitic algae. Individual specimens of *Halimeda discoidea* were collected from offshore Hawai'i and transplanted into an aquarium, where growth conditions were modified to simulate Cretaceous–Eocene seawater (Stanley et al., 2010). For the aquarium water, molar Mg/Ca was maintained at 1.5 and $[\text{Ca}^{2+}]$ at 25.3 mM, while the sum of $([\text{Mg}^{2+}] + [\text{Ca}^{2+}])$ remained equal to that of modern seawater to match total salinity.

After approximately six weeks of growth, eight individuals were harvested and sent to the University of Oxford. Four to five terminal segments from each individual were sampled by hand, bleached in $\sim 15\%$ NaClO to remove organic matter, and crushed to a powder by mortar and pestle. A number of wild specimens collected in the vicin-

BGD

11, 3559–3580, 2014

Ca isotopes of *Halimeda* algae

C. L. Blättler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Ca isotopes of
Halimeda algae

C. L. Blättler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ity of Lee Stocking Island in the Bahamas, including several *Halimeda* species, were also treated in the same fashion. X-ray diffractometry (XRD) was used to quantify the percentage of calcite and aragonite in each sample using a PANalytical X'Pert Pro diffractometer in the Department of Chemistry, University of Oxford. Approximately 10 mg of sample powder was scattered onto a glass slide with a frosted surface to randomize particle orientation. Aragonite produces prominent peaks at $2\theta = 26.2^\circ$ (3.40 Å, [111]), 27.2° (3.27 Å, [021]), and 33.0° (2.70 Å, [012]), whereas calcite produces a much stronger relative signal at $2\theta = 29.4^\circ$ (3.04 Å, [104]). The dominant calcite peak has a greater magnitude than that of any aragonite peak for a mass contribution of only 10% calcite. For the purpose of assessing the mass fraction of carbonate minerals, a calibration curve was generated using mixtures of aragonite (a fire coral, *Millepora*, from the Bahamas) and calcite (mineral spar) standards, using areas under the aragonite peak at $2\theta = 26.2^\circ$ and the calcite peak at $2\theta = 29.4^\circ$ (Fig. 1).

Ca-isotope analysis by multi-collector inductively coupled plasma mass spectrometry (MC-ICP-MS) was performed on the same powders used for XRD analysis, using between 0.5 and 2.0 mg of carbonate, and on the growth solutions. Methods are derived from those in Blättler et al. (2011). Sample Ca was isolated using ion-exchange chromatographic columns and prepared as 10 ppm Ca solutions in 2% HNO₃, with blanks contributing < 5 ng Ca to the total sample. Ratios of $^{44}\text{Ca}/^{42}\text{Ca}$ were measured on a Nu Instruments MC-ICP-MS, with the international carbonate standard NIST SRM915a used as a reference. Mass-dependent fractionation was confirmed by measuring ^{44}Ca , ^{43}Ca , and ^{42}Ca beams, correcting for the interference of double-charged Sr by monitoring the beam at mass 43.5. Multiple internal standard measurements and replicate analyses of samples generate standard errors of $\sim 0.05\%$. Values of $\delta^{44/42}\text{Ca}$, equal to $\left[\left(^{44}\text{Ca}/^{42}\text{Ca} \right)_{\text{sample}} / \left(^{44}\text{Ca}/^{42}\text{Ca} \right)_{\text{standard}} - 1 \right] \cdot 1000$, are converted to $\delta^{44/40}\text{Ca}$ relative to modern seawater and $\Delta^{44/40}\text{Ca}$ (Table 1).

4 Results

Combined XRD and Ca-isotope analyses allow a relationship between mineral composition and Ca-isotope ratios to be defined for *Halimeda* algae (Fig. 3). The carbonate skeleton of natural, wild samples contained 0–5% calcite. Seven of the experimental samples had slightly increased levels of 2.8–8.8% calcite, while the eighth experimental individual had a skeleton of > 85% calcite (Table 1 and Fig. 3). These data are supported by the morphologies of the experimental *Halimeda*, with the lowest amounts of calcite (within the range of natural *Halimeda*) present in thick, well-calcified segments, whereas higher levels (up to 8.8%) were present in flimsier, malformed samples (Fig. 2). The most calcite-rich sample was particularly shriveled and yielded a skeletal mass fraction < 10% of total sample mass, in comparison to other experimental and natural algae where the skeletons represented 45–75% of the total mass. The low mass of carbonate retrieved from this sample generated a very poor XRD spectrum with a significant background, a small calcite peak, and no identifiable aragonite peaks, so the calculated value of 88% calcite within the skeletal fraction should be treated as a rough minimum.

Ca-isotope ratios of various experimental materials were measured to establish $\Delta^{44/40}\text{Ca}$, the difference between fluid and precipitate, for all samples (Table 1). Analyses include the calcium chloride salt added to the aquarium water, the aquarium water from both before and after the growth period, terminal algal samples which grew entirely in artificial Cretaceous–Eocene seawater (eight individual *Halimeda*), and basal algal samples whose calcification commenced in natural seawater and may have continued in experimental seawater after transplanting (three individuals). The Ca in the artificial seawater was a mixture of modern seawater Ca and added calcium chloride salt in an approximate proportion of 1 : 3, based on the $\delta^{44/40}\text{Ca}$ of seawater (0‰, as the reference value), the salt (–1.19‰), and the resulting solution (–0.87‰). The Ca-isotope ratio in the aquarium rose slightly (by +0.1‰) during the course of the experiment, likely due to the preferential uptake of light Ca isotopes by algal calcification.

BGD

11, 3559–3580, 2014

Ca isotopes of *Halimeda* algae

C. L. Blättler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The seven experimental algae with < 10% calcite had an average $\delta^{44/40}\text{Ca}$ of -2.31‰ relative to natural seawater, or $\Delta^{44/40}\text{Ca} = -1.44\text{‰}$. This fractionation is in good agreement with six natural *Halimeda* samples which express $\Delta^{44/40}\text{Ca} = -1.42\text{‰}$. Three samples taken from the base of the experimental algae, rather than the terminal segments which grew entirely in the aquarium, had an average $\delta^{44/40}\text{Ca}$ of -1.65‰ relative to natural seawater, reflecting a mixture of carbonate precipitated before the transplant and continued calcification from the isotopically lighter aquarium water. The anomalous, calcite-rich sample had a much smaller fractionation than other *Halimeda*, with $\delta^{44/40}\text{Ca}$ of -1.47‰ and $\Delta^{44/40}\text{Ca} = -0.61\text{‰}$. This single sample allows the relationship between skeletal mineralogy and Ca-isotope fractionation of *Halimeda* to be established.

5 Discussion

The identification of specific fractionation mechanisms in the *Halimeda* Ca-isotope data is possible due to the relatively simple biology of the algae. *Halimeda* exert minimal control on the calcification process and the environment in which it takes place. Calcium carbonate oversaturation is achieved due to photosynthetic CO_2 consumption and subsequent local CO_3^{2-} elevation, without additional control over the carbonate system (Borowitzka and Larkum, 1987). Cell walls provide surfaces for precipitation of aragonite needles which subsequently fill the intercellular space, and their morphology suggests that the algae do not further control the crystallography of the precipitate with organic templates (Borowitzka and Larkum, 1987; Macintyre and Reid, 1995). Insensitivity to Ca-channel blocking drugs indicates that algal tissue has no influence on the supply of Ca, leaving diffusion of seawater as the only uptake mechanism (de Beer and Larkum, 2001). This lack of biological cation transport is consistent with the observation of similar Mg/Ca for inorganic precipitates and *Halimeda* during experiments in solutions with a variety of Mg/Ca ratios (Stanley et al., 2010). Without the complicating

BGD

11, 3559–3580, 2014

Ca isotopes of *Halimeda* algae

C. L. Blättler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



effects of organic matrices and active transport observed in some biological systems, the simplicity of *Halimeda* skeletal formation makes it possible to focus on other specific factors affecting Ca-isotope ratios.

The influence of mineralogy on the chemical composition of skeletal carbonate is reflected in the Ca-isotope composition for *Halimeda*. The calcite–aragonite mixing line that is developed for experimental *Halimeda* (Fig. 3) can be used to define endmember Ca-isotope values for the pure calcite and pure aragonite phases. The offset in $\delta^{44/40}\text{Ca}$ between aragonite and calcite endmembers is nearly identical, at ca. 0.9‰, for both the experimental *Halimeda* and inorganic precipitates, suggesting that the intrinsic fractionation factor for these carbonate polymorphs is expressed even when these crystals are formed in biogenic settings. Although a range of $\Delta^{44/40}\text{Ca}$ values have been observed for calcite and aragonite, the consistent offset between the phases suggests that effects from growth rate or CO_3^{2-} content are secondary to the first-order relationship between mineralogy and $\Delta^{44/40}\text{Ca}$. Given this behaviour for Ca-isotope ratios, it is expected that other metal/Ca ratios experience a similarly strong influence from carbonate mineralogy in *Halimeda* skeletons.

These *Halimeda* data also allow the effect of substrate on Ca-isotope fractionation to be assessed. Beaker experiments usually probe carbonate-nucleated growth of calcite or aragonite, but understanding the effect of an organic substrate on carbonate geochemistry could be important for interpreting ooid or microbial mat carbonate. The *Halimeda* experiment is able to isolate this factor to some degree, due to the space and manner in which aragonite needles grow (Borowitzka and Larkum, 1987). The results suggest that the organic substrate may be important for determining carbonate mineralogy, but not for further affecting cation geochemistry. The prevalence of aragonite in most of the experimental *Halimeda*, despite a fluid composition that mimics Cretaceous seawater (molar $\text{Mg}/\text{Ca} = 1.5$) and strongly favours calcite, suggests that the cell walls exerted influence over carbonate mineralogy, even without a three-dimensional organic matrix. By influencing the mineralogy of the first nucleated crystals, the organic surface may affect the structure and therefore geochemistry of further growth, even when the

Ca isotopes of *Halimeda* algae

C. L. Blättler et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

ambient fluid promotes an alternative mineralogy. *Halimeda* appears to behave in this manner, and other carbonate associated with, but not entirely controlled by, organic material may show similar mineralogical and geochemical behaviour.

Although many aspects of *Halimeda* calcification resemble inorganic calcification, $\Delta^{44/40}\text{Ca}$ for both the aragonitic and calcitic endmembers is $\sim 0.25\%$ smaller than for inorganic precipitates in the experiments of Gussone et al. (2003) and Marriott et al. (2004). Among the several factors which could introduce a vital effect of this magnitude, the calcification process of *Halimeda* suggests two primary candidates for this difference. One possibility is increased carbonate saturation and calcification rate, which has been shown to decrease fractionation on both an experimental and theoretical basis (Lemarchand et al., 2004; DePaolo, 2011). Within the intercellular spaces, consumption of CO_2 by photosynthesis drives carbonate oversaturation and promotes calcification. However, differences in calcification rate were visibly apparent in the samples, which ranged from thick and fleshy to stunted and shriveled, and did not always accompany changes in $\delta^{44/40}\text{Ca}$. This relationship implies that growth rate had a minor impact on Ca-isotope ratios. The second possibility is the occurrence of Rayleigh distillation, or a reservoir effect. As light isotopes are preferentially consumed within the intercellular space, the $\delta^{44/40}\text{Ca}$ of that reservoir will rise and the apparent fractionation between the external fluid and the skeleton will decrease. The rate of calcification relative to the recharge of Ca ions into the calcifying space by diffusion would determine the magnitude of this effect. This mechanism could be responsible for the difference in $\Delta^{44/40}\text{Ca}$ of inorganic aragonite and aragonitic sclerosponges, as the sponges also have a simple system of induced biocalcification (Gussone et al., 2005). Although the balance of calcification to diffusion rate could be specific to individual organisms, some inherent link between the two could arise as DIC (dissolved inorganic carbon) supply eventually limits calcification. If such a link occurs in other settings, then the magnitude of the Rayleigh effect in *Halimeda*, $\sim 0.25\%$, may reflect a characteristic vital effect among macroorganisms with a similar calcification mechanism.

Ca isotopes of *Halimeda* algae

C. L. Blättler et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

The Ca-isotope fractionation that is expressed by *Halimeda* has implications for the marine Ca-isotope budget. Due to the large difference in $\Delta^{44/40}\text{Ca}$ between aragonite and calcite, it is possible that the overall proportion of these two minerals in the global carbonate sink can affect the steady-state Ca-isotope composition of seawater (Farkaš et al., 2007; Blättler et al., 2012). This mechanism can help explain variation in reconstructed seawater $\delta^{44/40}\text{Ca}$ over the Phanerozoic only if the difference between sedimentary aragonite and calcite $\delta^{44/40}\text{Ca}$ is maintained over time. Although few major calcifying species remained quantitatively important contributors to the carbonate sink over long timescales (10^7 – 10^8 years), and *Halimeda* only became a major sediment producer in the late Cenozoic aragonite sea (Stanley and Hardie, 1998), the first-order control of mineralogy on biogenic $\delta^{44/40}\text{Ca}$ in this experimental setup appears to support this hypothesis.

6 Conclusions

This work presents a growth experiment with the sediment-producing algae *Halimeda* that investigates mechanisms of biocalcification and their geochemical expression. Developing a mechanistic understanding of biogenic carbonate formation and associated vital effects is necessary for linking skeletal proxy measurements to environmental interpretations and mass-balance calculations. While not applicable to every calcifying organism, the principles derived here show that it is possible to understand a combination of factors with superimposed effects on skeletal geochemistry. The aragonite and calcite endmember phases derived from the *Halimeda* experiment demonstrate how mineralogy strongly influences Ca-isotope ratios in biominerals. Biological substrates can exert first-order control on Ca-isotope fractionation by promoting a particular mineral form. The most substantial vital effect in this example is likely to be Rayleigh distillation of Ca within the algal intercellular space, which may have analogues in other simple macroorganisms. Stronger vital effects in other biocalcifiers are likely due to

BDG

11, 3559–3580, 2014

Ca isotopes of *Halimeda* algae

C. L. Blättler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



more sophisticated ion-transport mechanisms and internal reservoir dynamics, but the expression of an intrinsic mineralogical signal within *Halimeda* suggests that the calcite and aragonite content of marine carbonate can significantly affect the Ca-isotope balance of seawater. Further work on this and other biocalcifiers may continue to unravel the factors that control carbonate composition and increase the value of these important geological archives.

Acknowledgements. Thanks are due to Jennifer Engels for harvesting and shipping experimental samples to Oxford, Phil Holdship for analyzing cation concentrations in experimental seawater, and Phil Wiseman for generously accommodating XRD analyses in the Department of Chemistry, University of Oxford.

References

- Blättler, C. L., Jenkyns, H. C., Reynard, L. M., and Henderson, G. M.: Significant increases in global weathering during Oceanic Anoxic Events 1a and 2 indicated by calcium isotopes, *Earth Planet. Sc. Lett.*, 309, 77–88, doi:10.1016/j.epsl.2011.06.029, 2011. 3566
- Blättler, C. L., Henderson, G. M., and Jenkyns, H. C.: Explaining the Phanerozoic Ca-isotope history of seawater, *Geology*, 40, 843–846, doi:10.1130/G33191.1, 2012. 3571
- Böhm, F., Gussone, N., Eisenhauer, A., Dullo, W.-C., Reynaud, S., and Paytan, A.: Calcium isotope fractionation in modern scleractinian corals, *Geochim. Cosmochim. Ac.*, 70, 4452–4462, doi:10.1016/j.gca.2006.06.1546, 2006. 3563, 3565
- Borowitzka, M. A. and Larkum, A. W. D.: Calcification in algae: mechanisms and the role of metabolism, *Crit. Rev. Plant Sci.*, 6, 1–45, doi:10.1080/07352688709382246, 1987. 3561, 3568, 3569
- Bots, P., Benning, L. G., Rickaby, R. E. M., and Shaw, S.: The role of SO₄ in the switch from calcite to aragonite seas, *Geology*, 39, 331–334, 2011. 3561
- Burton, E. A. and Walter, L. M.: Relative precipitation rates of aragonite and Mg calcite from seawater: temperature or carbonate ion control?, *Geology*, 15, 111–114, 1987. 3561
- Burton, E. A. and Walter, L. M.: The role of pH in phosphate inhibition of calcite and aragonite precipitation rates in seawater, *Geochim. Cosmochim. Ac.*, 54, 797–808, 1990. 3561

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Ca isotopes of
Halimeda algae**

C. L. Blättler et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Chang, V. T.-C., Williams, R. J. P., Makishima, A., Belshaw, N. S., and O'Nions, R. K.: Mg and Ca isotope fractionation during CaCO_3 biomineralisation, *Biochem. Bioph. Res. Co.*, 323, 79–85, doi:10.1016/j.bbrc.2004.08.053, 2004. 3563
- de Beer, D. and Larkum, A. W. D.: Photosynthesis and calcification in the calcifying algae *Halimeda discoidea* studied with microsensors, *Plant Cell Environ.*, 24, 1209–1217, doi:10.1046/j.1365-3040.2001.00772.x, 2001. 3568
- DePaolo, D. J.: Surface kinetic model for isotopic and trace element fractionation during precipitation of calcite from aqueous solutions, *Geochim. Cosmochim. Ac.*, 75, 1039–1056, doi:10.1016/j.gca.2010.11.020, 2011. 3563, 3570
- Dickson, J. A. D.: Echinoderm skeletal preservation: calcite–aragonite seas and the Mg/Ca ratio of Phanerozoic oceans, *J. Sediment. Res.*, 74, 355–365, doi:10.1306/112203740355, 2004. 3561
- Erez, J.: The source of ions for biomineralization in foraminifera and their implications for paleoceanographic proxies, *Rev. Mineral. Geochem.*, 54, 115–149, doi:10.2113/0540115, 2003. 3565
- Fantle, M. S. and DePaolo, D. J.: Ca isotopes in carbonate sediment and pore fluid from ODP Site 807A: the $\text{Ca}^{2+}(\text{aq})$ –calcite equilibrium fractionation factor and calcite recrystallization rates in Pleistocene sediments, *Geochim. Cosmochim. Ac.*, 71, 2524–2546, doi:10.1016/j.gca.2007.03.006, 2007. 3563
- Farkaš, J., Böhm, F., Wallmann, K., Blenkinsop, J., Eisenhauer, A., van Geldern, R., Munnecke, A., Voigt, S., and Veizer, J.: Calcium isotope record of Phanerozoic oceans: implications for chemical evolution of seawater and its causative mechanisms, *Geochim. Cosmochim. Ac.*, 71, 5117–5134, doi:10.1016/j.gca.2007.09.004, 2007. 3571
- Gagnon, A. C., DePaolo, D. J., Adkins, J. F., and De Yoreo, J. J.: Calcium isotopes during coral biomineralization, in: *Goldschmidt Conference Abstracts*, vol. 75, p. 880, *Mineralogical Magazine*, 2011. 3563
- Griffith, E. M., Paytan, A., Kozdon, R., Eisenhauer, A., and Ravelo, A. C.: Influences on the fractionation of calcium isotopes in planktonic foraminifera, *Earth Planet. Sc. Lett.*, 268, 124–136, doi:10.1016/j.epsl.2008.01.006, 2008. 3563, 3565
- Gussone, N. and Filipsson, H. L.: Calcium isotope ratios in calcitic tests of benthic foraminifers, *Earth Planet. Sc. Lett.*, 290, 108–117, doi:10.1016/j.epsl.2009.12.010, 2010. 3564
- Gussone, N., Eisenhauer, A., Heuser, A., Dietzel, M., Bock, B., Böhm, F., Spero, H. J., Lea, D. W., Bijma, J., and Nägler, T. F.: Model for kinetic effects on calcium isotope frac-

Ca isotopes of
Halimeda algae

C. L. Blättler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- tionation ($\delta^{44}\text{Ca}$) in inorganic aragonite and cultured planktonic foraminifera, *Geochim. Cosmochim. Ac.*, 67, 1375–1382, doi:10.1016/S0016-7037(02)01296-6, 2003. 3562, 3563, 3570
- Gussone, N., Böhm, F., Eisenhauer, A., Dietzel, M., Heuser, A., Teichert, B. M. A., Reiter, J., Wörheide, G., and Dullo, W.-C.: Calcium isotope fractionation in calcite and aragonite, *Geochim. Cosmochim. Ac.*, 69, 4485–4494, doi:10.1016/j.gca.2005.06.003, 2005. 3563, 3564, 3570, 3580
- Gussone, N., Langer, G., Thoms, S., Nehrke, G., Eisenhauer, A., Riebesell, U., and Wefer, G.: Cellular calcium pathways and isotope fractionation in *Emiliania huxleyi*, *Geology*, 34, 625–628, doi:10.1130/G22733.1, 2006. 3563, 3564
- Gussone, N., Langer, G., Geisen, M., Steel, B. A., and Riebesell, U.: Calcium isotope fractionation in coccoliths of cultured *Calcidiscus leptoporus*, *Helicosphaera carteri*, *Syracosphaera pulchra* and *Umbilicosphaera foliosa*, *Earth Planet. Sc. Lett.*, 260, 505–515, doi:10.1016/j.epsl.2007.06.001, 2007. 3564
- Gussone, N., Hönisch, B., Heuser, A., Eisenhauer, A., Spindler, M., and Hemleben, C.: A critical evaluation of calcium isotope ratios in tests of planktonic foraminifers, *Geochim. Cosmochim. Ac.*, 73, 7241–7255, doi:10.1016/j.gca.2009.08.035, 2009. 3564, 3565
- Gussone, N., Nehrke, G., and Teichert, B. M. A.: Calcium isotope fractionation in ikaite and vaterite, *Chem. Geol.*, 285, 194–202, doi:10.1016/j.chemgeo.2011.04.002, 2011. 3562
- Hardie, L. A.: Secular variation in seawater chemistry: an explanation for the coupled secular variation in the mineralogies of marine limestones and potash evaporites over the past 600 m.y., *Geology*, 24, 279–283, doi:10.1130/0091-7613(1996)024<0279:SVISCA>2.3.CO;2, 1996. 3561
- Heinemann, A., Fietzke, J., Eisenhauer, A., and Zumholz, K.: Modification of Ca isotope and trace metal composition of the major matrices involved in shell formation of *Mytilus edulis*, *Geochem. Geophys. Geosy.*, 9, Q01006, doi:10.1029/2007GC001777, 2008. 3564
- Hippler, D., Eisenhauer, A., and Nägler, T. F.: Tropical Atlantic SST history inferred from Ca isotope thermometry over the last 140ka, *Geochim. Cosmochim. Ac.*, 70, 90–100, doi:10.1016/j.gca.2005.07.022, 2006. 3563
- Horita, J., Zimmermann, H., and Holland, H. D.: Chemical evolution of seawater during the Phanerozoic: implications from the record of marine evaporites, *Geochim. Cosmochim. Ac.*, 66, 3733–3756, doi:10.1016/S0016-7037(01)00884-5, 2002. 3561

Ca isotopes of
Halimeda algae

C. L. Blättler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Jacobson, A. D. and Holmden, C.: $\delta^{44}\text{Ca}$ evolution in a carbonate aquifer and its bearing on the equilibrium isotope fractionation factor for calcite, *Earth Planet. Sc. Lett.*, 270, 349–353, doi:10.1016/j.epsl.2008.03.039, 2008. 3563
- 5 Kasemann, S. A., Schmidt, D. N., Pearson, P. N., and Hawkesworth, C. J.: Biological and ecological insights into Ca isotopes in planktic foraminifers as a palaeotemperature proxy, *Earth Planet. Sc. Lett.*, 271, 292–302, doi:10.1016/j.epsl.2008.04.007, 2008. 3563
- Kisakürek, B., Eisenhauer, A., Böhm, F., Hathorne, E. C., and Erez, J.: Controls on calcium isotope fractionation in cultured planktic foraminifera, *Globigerinoides ruber* and *Globigerinella siphonifera*, *Geochim. Cosmochim. Ac.*, 75, 427–443, doi:10.1016/j.gca.2010.10.015, 2011. 3564, 3565
- 10 Langer, G., Gussone, N., Nehrke, G., Riebesell, U., Eisenhauer, A., and Thoms, S.: Calcium isotope fractionation during coccolith formation in *Emiliania huxleyi*: independence of growth and calcification rate, *Geochem. Geophys. Geosy.*, 8, Q05007, doi:10.1029/2006GC001422, 2007. 3564
- 15 Lemarchand, D., Wasserburg, G. J., and Papanastassiou, D. A.: Rate-controlled calcium isotope fractionation in synthetic calcite, *Geochim. Cosmochim. Ac.*, 68, 4665–4678, doi:10.1016/j.gca.2004.05.029, 2004. 3562, 3564, 3570
- Lowenstein, T. K., Timofeeff, M. N., Brennan, S. T., Hardie, L. A., and Demicco, R. V.: Oscillations in Phanerozoic seawater chemistry: evidence from fluid inclusions, *Science*, 294, 1086–1088, doi:10.1126/science.1064280, 2001. 3561
- 20 Macintyre, I. G. and Reid, R. P.: Crystal alteration in a living calcareous alga (*Halimeda*): implications for studies in skeletal diagenesis, *J. Sediment. Res.*, 65, 143–153, doi:10.1306/D4268054-2B26-11D7-8648000102C1865D, 1995. 3568
- Marriott, C. S., Henderson, G. M., Belshaw, N. S., and Tudhope, A. W.: Temperature dependence of $\delta^7\text{Li}$, $\delta^{44}\text{Ca}$ and Li/Ca during growth of calcium carbonate, *Earth Planet. Sc. Lett.*, 222, 615–624, doi:10.1016/j.epsl.2004.02.031, 2004. 3562, 3570, 3580
- 25 Morse, J. W., Wang, Q., and Tsio, M. Y.: Influences of temperature and Mg: Ca ratio on CaCO_3 precipitates from seawater, *Geology*, 25, 85–87, 1997. 3561
- Nägler, T. F., Eisenhauer, A., Müller, A., Hemleben, C., and Kramers, J.: The $\delta^{44}\text{Ca}$ -temperature calibration on fossil and cultured *Globigerinoides sacculifer*: new tool for reconstruction of past sea surface temperatures, *Geochem. Geophys. Geosy.*, 1, 1052, doi:10.1029/2000GC000091, 2000. 3563
- 30

**Ca isotopes of
Halimeda algae**

C. L. Blättler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Reynard, L. M., Day, C. C., and Henderson, G. M.: Large fractionation of calcium isotopes during cave-analogue calcium carbonate growth, *Geochim. Cosmochim. Ac.*, 75, 3726–3740, doi:10.1016/j.gca.2011.04.010, 2011. 3562

Ries, J. B., Stanley, S. M., and Hardie, L. A.: Scleractinian corals produce calcite, and grow more slowly, in artificial Cretaceous seawater, *Geology*, 34, 525–528, doi:10.1130/G22600.1, 2006. 3561

Sandberg, P. A.: An oscillating trend in Phanerozoic non-skeletal carbonate mineralogy, *Nature*, 305, 19–22, doi:10.1038/305019a0, 1983. 3561

Sime, N. G., De La Rocha, C. L., and Galy, A.: Negligible temperature dependence of calcium isotope fractionation in 12 species of planktonic foraminifera, *Earth Planet. Sc. Lett.*, 232, 51–66, doi:10.1016/j.epsl.2005.01.011, 2005. 3563

Stanley, S. M. and Hardie, L. A.: Secular oscillations in the carbonate mineralogy of reef-building and sediment-producing organisms driven by tectonically forced shifts in seawater chemistry, *Palaeogeogr. Palaeoclimatol.*, 144, 3–19, doi:10.1016/S0031-0182(98)00109-6, 1998. 3561, 3571

Stanley, S. M., Ries, J. B., and Hardie, L. A.: Low-magnesium calcite produced by coralline algae in seawater of Late Cretaceous composition, *P. Natl. Acad. Sci. USA*, 99, 15323–15326, doi:10.1073/pnas.232569499, 2002. 3561

Stanley, S. M., Ries, J. B., and Hardie, L. A.: Seawater chemistry, coccolithophore population growth, and the origin of Cretaceous chalk, *Geology*, 33, 593–596, doi:10.1130/G21405.1, 2005. 3561

Stanley, S. M., Ries, J. B., and Hardie, L. A.: Increased production of calcite and slower growth for the major sediment-producing alga *Halimeda* as the Mg/Ca ratio of seawater is lowered to a “calcite sea” level, *J. Sediment. Res.*, 80, 6–16, doi:10.2110/jsr.2010.011, 2010. 3561, 3562, 3565, 3568

Tang, J., Dietzel, M., Böhm, F., Köhler, S. J., and Eisenhauer, A.: Sr²⁺/Ca²⁺ and ⁴⁴Ca/⁴⁰Ca fractionation during inorganic calcite formation: II. Ca isotopes, *Geochim. Cosmochim. Ac.*, 72, 3733–3745, doi:10.1016/j.gca.2008.05.033, 2008. 3562

Weiner, S. and Dove, P. M.: An overview of biomineralization processes and the problem of the vital effect, *Rev. Mineral. Geochem.*, 54, 1–29, doi:10.2113/0540001, 2003. 3561

Ca isotopes of
Halimeda algae

C. L. Blättler et al.

Table 1. Ca-isotope data for natural and experimental *Halimeda* algae, along with other materials from the experiment. Percent calcite (% cc) determined by quantifying calcite (cc) and aragonite (arag) by X-ray diffractometry and calculating % cc = cc/(cc + arag), with errors of $\pm 5\%$ (except for sample HAL-6 for which they are $\pm 15\%$). Ca-isotope ratios are reported as $\delta^{44/42}\text{Ca} = [({}^{44}\text{Ca}/{}^{42}\text{Ca})_{\text{sample}}/({}^{44}\text{Ca}/{}^{42}\text{Ca})_{\text{SRM915a}} - 1] \cdot 1000$, $\delta^{44/40}\text{Ca} = [({}^{44}\text{Ca}/{}^{40}\text{Ca})_{\text{sample}}/({}^{44}\text{Ca}/{}^{40}\text{Ca})_{\text{seawater}} - 1] \cdot 1000$, and $\Delta^{44/40}\text{Ca} = \delta^{44/40}\text{Ca}_{\text{carbonate}} - \delta^{44/40}\text{Ca}_{\text{solution}}$.

Sample name	$\delta^{44/42}\text{Ca}$ (‰) rel. to SRM915a	<i>n</i>	$\delta^{44/40}\text{Ca}$ (‰) rel. to seawater	2 SE	avg. $\delta^{44/40}\text{Ca}$	$\Delta^{44/40}\text{Ca}$ (difference)	% cc	wt.% carbonate
Ca salt used for Cretaceous "seawater"								
CaCl ₂ -A	0.40	4	-1.08	0.07	-1.19			
CaCl ₂ -B	0.38	4	-1.13	0.07				
CaCl ₂ -C	0.33	4	-1.23	0.10				
CaCl ₂ -D	0.27	3	-1.34	0.11				
Pre-experimental Cretaceous "seawater"								
PRE-A	0.54	4	-0.80	0.07	-0.87	0.00		
PRE-B	0.51	4	-0.86	0.07				
PRE-C	0.49	4	-0.91	0.10				
PRE-D	0.49	3	-0.91	0.11				
Post-experimental Cretaceous "seawater"								
EXP-A	0.58	4	-0.72	0.07	-0.77	0.10		
EXP-B	0.55	4	-0.77	0.07				
EXP-C	0.54	3	-0.81	0.10				
EXP-D	0.54	4	-0.79	0.11				
Experimental <i>Halimeda</i> samples, post-XRD analysis (terminal segments only)								
HAL-1	-0.30	5	-2.49	0.10	-2.20	-1.62	2.8	73
HAL-2	-0.19	4	-2.25	0.11		-1.38	4.9	70
HAL-3	-0.21	4	-2.31	0.11		-1.44	7.6	63
HAL-4	-0.21	4	-2.31	0.11		-1.44	8.8	49
HAL-5	-0.15	4	-2.17	0.11		-1.30	3.7	57
HAL-6 ^a	0.20	4	-1.47	0.09		-0.61	88.3	4
HAL-7	-0.15	4	-2.19	0.11		-1.32	8.5	45
HAL-8 ^b	-0.27	5	-2.42	0.10		-1.56	4.0	50
Transplanted <i>Halimeda</i> samples (basal samples)								
HAL-2X	0.14	5	-1.59	0.09	-1.65	-1.59		
HAL-4X	0.01	5	-1.85	0.09		-1.85		
HAL-6X	0.19	5	-1.50	0.09		-1.50		
Wild <i>Halimeda</i> samples (from the Bahamas)								
<i>H. monile</i>	0.30	4	-1.28	0.13	-1.42	-1.28		
<i>H. tuna</i>	0.18	4	-1.52	0.13		-1.52		
<i>H. discoidea</i>	0.26	4	-1.36	0.13		-1.36		
<i>H. discoidea</i>	0.22	4	-1.44	0.13		-1.44		
<i>H. incrassata</i>	0.19	4	-1.50	0.13		-1.50		
<i>H. incrassata</i>	0.22	4	-1.44	0.13		-1.44		

^a poorly calcified sample

^b offshot of another individual, entirely grown in aquarium






Fig. 1. Standard calibration curve for quantifying the proportions of calcite and aragonite using XRD. Analyses used a $\text{Cu-K}\alpha$ X-ray source (40 mA and 40 kV) with the sample loaded on a fixed stage while 2θ varied through $25\text{--}40^\circ$ with a 0.004° step size (0.05°s^{-1}). Areas under the calcite (cc) peak at 29.4° and the aragonite (arag) peak at 26.2° were used to calculate the proportional intensity of calcite as $\text{cc}/(\text{cc} + \text{arag})$. Insets show standard intensity vs. 2θ -angle XRD spectra for 0% (pure aragonite), 49.7%, and 100% calcite (note the relative change in the vertical axis for the 0% calcite/pure aragonite standard).



wild *Halimeda discoidea*



Sample #1, 2.8% cc



Sample #2, 4.9% cc



Sample #3, 7.6% cc



Sample #4, 8.8% cc



Sample #5, 3.7% cc



Sample #6, >88% cc



Sample #7, 8.5% cc



Sample #8, 4.0% cc*

*offshoot, grown entirely in aquarium

Fig. 2. Photographs of the eight individual *Halimeda* samples from the experimental Cretaceous seawater. Note that Sample #6, which contained mostly calcite rather than aragonite, appears particularly malformed.

Ca isotopes of *Halimeda* algae

C. L. Blättler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 3. Ca-isotope ratios for *Halimeda* grown in experimental Cretaceous seawater, plotted against the percentage of calcite determined by XRD analysis. The open circle represents the average value for natural, ocean-grown *Halimeda*. The black line represents a mixing relationship between the inorganic calcite and aragonite end-members from Marriott et al. (2004) and Gussone et al. (2005), respectively, at 15 °C. The *Halimeda* data define a best-fit line with similar slope and approximately 0.25‰ heavier values than inorganic carbonate.