

This discussion paper is/has been under review for the journal Biogeosciences (BG).
Please refer to the corresponding final paper in BG if available.

Soil organic matter dynamics under different land-use in grasslands in Inner Mongolia (northern China)

L. Zhao, W. Wu, X. Xu, and Y. Xu

Moe Key Laboratory for Earth Surface Processes, College of Urban and Environmental Sciences, Peking University, Beijing 100871, China

Received: 20 January 2014 – Accepted: 26 March 2014 – Published: 11 April 2014

Correspondence to: Y. Xu (yunpingxu@pku.edu.cn)

Published by Copernicus Publications on behalf of the European Geosciences Union.

BGD

11, 5613–5637, 2014

Soil organic matter dynamics

L. Zhao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

We examined bulk soil properties and molecular biomarker distributions in surface soils from Inner Mongolia grasslands in order to understand the responses of soil organic matter to different land-use. The total of sixteen soils were collected from severely degraded grassland by overgrazing (DG), native grassland without apparent anthropogenic disturbance (NG), groundwater-sustaining grassland (GG) and restored grassland from previous potato cropland (RG). Compared to NG, soil organic carbon content was lower by 50 % in DG, but higher by six-fold in GG and one-fold in RG. The $\delta^{13}\text{C}$ values of soil organic carbon were -24.2 ± 0.6 ‰ in DG, -24.9 ± 0.6 ‰ in NG, -25.1 ± 0.1 ‰ in RG and -26.2 ± 0.6 ‰ in GG, reflecting different degradation degrees of soil organic matter or different water use efficiencies. The soils in DG contained the lowest abundance of aliphatic lipids (*n*-alkanes, *n*-alkanols, *n*-alkanoic acids, ω -hydroxylalkanoic acids and α -hydroxylalkanoic acids) and lignin-phenols, suggesting selective removal of these biochemically recalcitrant biomarkers with grassland degradation by microbial respiration or wind erosion. Compared to NG, the soils in GG and RG increased ω -hydroxylalkanoic acids by 60–70 %, a biomarker for suberin from roots, and increased α -hydroxylalkanoic acids by 10–20 %, a biomarker for both cutin and suberin. Our results demonstrate that the groundwater supply and cultivation-restoration practices in Inner Mongolia grasslands not only enhance soil organic carbon sequestration, but also change the proportions of shoot vs. root-derived carbon in soils. This finding has important implications for global carbon cycle since root derived aliphatic carbon has a longer residence time than the aboveground tissue-derived carbon in soils.

1 Introduction

Soil organic carbon, representing 80 % of terrestrial active carbon pool, plays a key role in the global carbon cycle (Janzen, 2004; Schlesinger and Andrews, 2000; Watson, 2000). Recently increasing human activities such as stock grazing, cultivation, defor-

BGD

11, 5613–5637, 2014

Soil organic matter dynamics

L. Zhao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Soil organic matter
dynamics**

L. Zhao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I◀

▶I

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



estation and plantation substantially change the carbon balance between soil and atmosphere and thereby mitigate or accelerate global climate changes. Guo et al. (2007) reported that soil carbon stock to 100 cm decreased by 20 % after 16 year plantation in a native pasture of Australia, while Islam et al. (2000) found 44 % decline in soil quality in a tropical forest ecosystem of Bangladesh under cultivation. The change in quality and quantity of soil organic matter inevitably affects various terrestrial ecosystem functions such as the soil fertility, biological diversity and biomass productivity (Kaiser et al., 2010).

Grasslands are important ecological landscapes in China and account for 40 % of the national land area (ca. 4 000 000 km²) (National Statistics Bureau of China, 2002). These grasslands are mainly distributed in the semiarid and semi-humid northern China (e.g., Inner Mongolia) and Tibet Plateau and are highly susceptible ecosystems for changes in climate and land-use (He et al., 2008; Xiao et al., 1995; Zhou et al., 2007). Previous studies have demonstrated that improper land-use can cause grassland degradation and even desertification in Inner Mongolia (e.g. Cui et al., 2005; Su et al., 2005; Wang et al., 2013; Zhao et al., 2007). However, most of these studies focused on bulk soil properties such as total organic carbon and nitrogen contents. Since SOM consists of heterogeneous mixtures of plant, microbe and animal-derived residues with different environmental stabilities (Baldock and Skjemstad, 2000; Kögel-Knabner, 2002), the investigation for different fractions and even molecular compositions of SOM is indispensable for understanding carbon dynamics under different land-use practices (Leifeld and Kögel-Knabner, 2005).

Organic matter biomarkers are compounds specific for certain organisms (e.g., higher plants, fungi and bacteria) or tissues (e.g., leaves and root) (Kögel-Knabner, 2000; Otto et al., 2005). Free lipids from plant waxes and biopolymers such as suberin and cutin are the major sources of the aliphatic constituents of SOM, while lignin is an important contributor to aromatic components of SOM (Kögel-Knabner, 2002). Over the past decades, biomarker approaches have been successfully used to estimate SOM responses to different environmental changes such as soil warming (e.g. Feng et al.,

2008), vegetation shifts (e.g. Pisani et al., 2013) and elevated atmospheric CO₂ (e.g. Wiesenberg et al., 2008).

In this study, we analyzed the composition and distribution of bulk organic matter and molecular biomarkers in grassland soils from Inner Mongolia, northern China (Fig. S1).

5 These grasslands have developed in the same climate region and have same dominant grass species, but are subject to different levels of human disturbances. The main objectives of our study are to understand the chemical characteristics of SOM at the bulk and molecular levels in temperate grasslands in Inner Mongolia and to evaluate influences of land-use practices on SOM. We also attempt to answer whether certain components of SOM such as aliphatic lipids, cutin/suberin and lignin, usually thought to be refractory components, are preferentially preserved with changes in grassland uses.

2 Materials and methods

2.1 Study area and sampling

15 Our study was conducted on the Keshiketengqi Grassland in the Inner Mongolia Autonomous Region (northern China), being 400 km away from Beijing (Fig. S1). The semi-arid climate has a long, cold winter (November to March) and a short summer (June to August). The mean annual temperature is -1.5 °C and the mean annual precipitation is 400 mm (Feng and Zhao, 2011; Tan et al., 2013). The soil is classified as a chestnut soil.

20 Four experimental fields with the size of 100m × 100m were chosen in our study (Fig. 1), representing severely degraded grassland (DG; 42°37' N, 117°05' E), native grassland (NG; 42°35' N, 117°05' E), groundwater-sustaining grassland (GG; 42°32' N, 117°08' E) and restored grassland from previous cropland (RG; 42°33' N, 117°13' E).

25 The DG has been subject to intense sheep and cattle grazing and trampling for over 15 years and its vegetation is sparse. At NG, grazing activity is seldom and there was no

BGD

11, 5613–5637, 2014

Soil organic matter dynamics

L. Zhao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Soil organic matter
dynamics

L. Zhao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



apparent anthropogenic disturbance. Pristine grassland at the site RG was converted into a potato cropland in 1990, and the cultivation activities lasted until 2010. Since then, this cropland was restored to the grassland by planting indigenous grasses and banning stock grazing. The GG is in a lowland area and supplied by groundwater in summer (June to September) when rainfall is relatively high. Similar to NG, the grazing activity is seldom at GG. Among four sampling sites, vegetation in GG is the best developed. Since the NG is subject to the least anthropogenic impact and resembles the major native grasslands in the Keshiketengqi Grassland, it is regarded as the reference site. In May 2012, four soil samples (0–10 cm) were collected from each site by a soil auger. After sampling, the soils were freeze dried at -40°C and passed through a 2 mm sieve to remove large roots and coarse particles. Soil samples were stored at -20°C until further analyses.

2.2 Grain size, elemental and stable isotope analyses

The detailed methods for grain size measurement has been described in Sun et al. (2011). About 1.0 g dried soils were reacted with excess 1 M hydrochloric acid and then with hydrogen peroxide to remove carbonates and oxidize organic matter, respectively. After that, sodium hexametaphosphate was added, and the solutions were allowed to settle for 24 h. After ultra-sonication for 1 min, the grain size distributions were measured by a Mastersizer 2000 Laser Particle Size Analyzer. The scan range was from 0.02 to 2000 μm , and categorized into three fractions: sand (20 ~ 2000 μm), silt (2 ~ 20 μm) and clay ($< 2 \mu\text{m}$).

The procedures for elemental and stable isotopic analyses were modified from Wu et al. (2014). The dried soil samples were treated with 1 M hydrochloric acid to remove inorganic carbon. The residues were analyzed by a CHNOS Element Analyzer (Elementar Germany) for total organic carbon (TOC) and total nitrogen (TN), while the $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ were determined on a Flash EA1112HT coupled with MAT 253 (Thermo Fisher Scientific, Inc). The standard deviations based on the replicate analyses were 0.02 % for TOC, 0.005 % for TN, 0.15 ‰ for $\delta^{13}\text{C}$ and 0.10 ‰ for $\delta^{15}\text{N}$.

2.3 Extractions of free lipids, bound lipids and lignin-phenols

Figure 2 shows the procedures for biomarkers analyses. About 6 g soil samples were mixed with 2 μg squalane and extracted with ultrasonication three times, each with 15 mL dichloromethane:methanol (1:1; v/v) for 15 min. The combined solvent extracts were filtered through glass fiber columns, concentrated by rotary evaporation, and dried under N_2 gas in 2 mL glass vials.

Soil residues after the solvent extraction were subjected to base hydrolysis to cleave bound lipids (Otto et al., 2005). After addition of 1 M methanolic KOH (15 mL) and 20 μg 5 α -cholestane (internal standard), the air-dried soil residues (~ 3 g) were sealed and heated at 100 $^\circ\text{C}$ for 3 h. After cooling, the suspension was acidified to pH 1 by addition of 6 M HCl. The bound lipids were recovered from the water phase by liquid-liquid extraction with 20 mL ethyl acetate three times. Anhydrous Na_2SO_4 was added to remove any remaining water. The ethyl acetate extracts were concentrated by rotary evaporation, transferred to 2 mL glass vials and dried under nitrogen gas.

Subsamples after the base hydrolysis were oxidized with CuO to release lignin-derived phenols. Teflon-lined bombs were loaded with 2 g of dry soil, 1 g CuO, 100 mg ammonium iron (II) sulfate hexahydrate and 15 mL of 2 M NaOH. The teflon vessels were purged with nitrogen gas for oxygen exhaustion, sealed and heated for 2.5 h at 170 $^\circ\text{C}$. After reaction, the liquid was decanted into a 50 mL centrifuge tube, and the residue was washed twice, each with 10 mL deionized water. The combined washings were centrifuged for 10 min at 3000 rpm. The supernatant was transferred into another centrifuge tube, acidified to pH 1 by addition of 6 M HCl and kept for 1 h at room temperature in the dark to prevent reactions of cinnamic acids. After centrifugation for 10 min at 3000 rpm again, the supernatant was transferred to a separation funnel and liquid-liquid extracted three times, each with 20 mL ethyl acetate. Anhydrous Na_2SO_4 was added to remove water. The ethyl acetate extracts were concentrated by rotary evaporation, transferred to 2 mL glass vials and dried under nitrogen gas.

BGD

11, 5613–5637, 2014

Soil organic matter dynamics

L. Zhao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.4 Derivatization of extracts and instrumental analyses

The base hydrolysis extracts were first methylated by reaction of 14 % BF_3 : MeOH for 90 min at 70 °C. The bound lipids were obtained by liquid-liquid extraction with 2 mL hexane (3×), and dried under N_2 stream. After that, all biomarkers from solvent extracts, base hydrolysis and CuO oxidation were trimethylsilyl (TMS) derivatized by reaction with 50 μL *N,O*-bis-(trimethylsilyl) trifluoroacetamide (BSTFA) and 10 μL pyridine for 2 h at 60 °C. After cooling, DCM was added to dilute to 1 mL.

The biomarkers were identified by a gas chromatography-mass spectrometry (GC-MS). This instrument was composed of an Agilent 7890A GC coupled to an Agilent 5973N quadruple mass selective detector. The biomarkers were separated by a DB-5 capillary column (30 m; 0.25 mm i.d.; 0.25 μm film thickness). A 1 μL of samples was injected in the splitless mode under a constant flow (1.0 mL min^{-1}). The injector temperature was 300 °C. High pure helium (> 99.999 %) was used as carrier gas. The mass spectrometer was operated in the electron impact mode (70 eV). The mass scan range was 50 to 550 Da.

An Agilent 7890A GC coupled to a flame ionization detector (FID) was used for quantification of biomarkers. The separation was achieved on an HP 5 capillary column (30 m; 0.32 mm i.d.; 0.25 μm film thickness). The injector temperature was 300 °C. A 1 μL volume of sample was injected in splitless mode under a constant flow (1.5 mL min^{-1}). Helium (purity > 99.999 %) was the carrier gas.

Oven programs for GC-MS and GC-FID were set as follows: temperature held at 60 °C for 1 min, increased to 300 °C at a rate of 6 °C min^{-1} with a final isothermal hold at 300 °C for 20 min. For solvent extracts and base hydrolysis, the biomarkers were quantified by comprising the peak areas of biomarkers to an internal standard (squalane for free lipids and 5 α -cholestane for bound lipids), while the amounts of lignin phenols were determined by the calibration curves of eight lignin-phenol standards.

BGD

11, 5613–5637, 2014

Soil organic matter dynamics

L. Zhao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.5 Statistical analysis

The program package SPSS 18.0 (Illinois, USA) was used for statistical analyses. One-way ANOVA analysis was conducted to examine the differences in bulk properties and molecular biomarkers among soils from different grasslands. All analyses were performed with a significance level of $P < 0.05$.

3 Results

3.1 Elemental and isotopic compositions

Table 1 shows the grain size distributions, elemental and isotopic compositions of grassland soils. All soils were dominated by sand (64.4 to 87.3%), while clay fractions accounted for only 1.7 to 6.4% in an increasing order of DG, NG, GG and RG. The TOC content ranged from $3.2 \pm 0.2 \text{ g kg}^{-1}$ (DG) to $46.6 \pm 16.7 \text{ g kg}^{-1}$ (GG), while the TN content gradually increased from $0.32 \pm 0.15 \text{ g kg}^{-1}$ (DG) to $3.9 \pm 1.5 \text{ g kg}^{-1}$ (GG). The values of $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ varied from $-24.2 \pm 0.6\text{‰}$ (DG) to $-26.2 \pm 0.6\text{‰}$ (GG) and $0.36 \pm 0.3\text{‰}$ (DG) to $4.8 \pm 1.6\text{‰}$ (GG), respectively. The C/N ratio increased from 9.9 to 12.5 in an order of DG, NG, GG and RG. A strong linear correlation was observed between the $\delta^{13}\text{C}$ and C/N ratio ($r = -0.667$, $n = 16$, $p < 0.01$), suggesting that both parameters are controlled by same factors such as the source and degradation stage of SOM.

3.2 Compositions and distributions of solvent extracts

The major components of the solvent extracts were *n*-alkanoic acids, *n*-alkanols, ω -hydroxyalkanoic acids, *n*-alkanes, steroids and sugars (Fig. S2). The concentrations of each type of biomarkers were summarized in Fig. 3. Long-chain ($> \text{C}_{20}$) aliphatic lipids including *n*-alkanes, *n*-alkanols and *n*-alkanoic acid are derived from epicuticular waxes of vascular plants, while short-chain ($< \text{C}_{20}$) aliphatic lipids are of a microbial

BGD

11, 5613–5637, 2014

Soil organic matter dynamics

L. Zhao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Soil organic matter
dynamics

L. Zhao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



origin (Otto et al., 2005; Simoneit, 2005). The soils from NG contained the highest concentrations of free aliphatic lipids ($3.53 \pm 1.18 \text{ mg g}^{-1} \text{ OC}$), followed by those from DG ($2.23 \pm 1.03 \text{ mg g}^{-1} \text{ OC}$), RG ($1.36 \pm 0.73 \text{ mg g}^{-1} \text{ OC}$) and GG ($1.20 \pm 0.65 \text{ mg g}^{-1} \text{ OC}$). Since the biomarker abundances have been normalized to organic carbon, the lowest concentrations of free aliphatic lipids at GG and RG were attributed to a dilution effect of their relatively high TOC contents. The homologues of *n*-alkanoic acids (C_{14} to C_{32}) were detected at the total concentrations of $1.20 \pm 0.39 \text{ mg g}^{-1} \text{ OC}$ in NG, $0.54 \pm 0.24 \text{ mg g}^{-1} \text{ OC}$ in DG, $0.40 \pm 0.37 \text{ mg g}^{-1} \text{ OC}$ in RG and $0.38 \pm 0.23 \text{ mg g}^{-1} \text{ OC}$ in GG. These *n*-alkanoic acids were characterized by the strong even-over-odd carbon numbered predominance and the C_{max} at C_{16} and C_{24} , suggesting a mixed input of soil microbes and vascular plants. Besides saturated acids, mono- and di-unsaturated C_{16} and C_{18} *n*-alkanoic acids were also detected at the concentrations lower than $0.25 \text{ mg g}^{-1} \text{ OC}$. These short-chain unsaturated alkanolic acids are mainly biosynthesized by bacteria and fungi.

The homologous *n*-alkanes (C_{22} to C_{33}) displayed a strong odd-over-even predominance and the maximum abundance (C_{max}) at C_{29} . The soils in NG were composed of the highest abundant *n*-alkanes ($0.33 \pm 0.08 \text{ mg g}^{-1} \text{ OC}$), followed by soils in DG ($0.32 \pm 0.12 \text{ mg g}^{-1} \text{ OC}$), RG ($0.31 \pm 0.20 \text{ mg g}^{-1} \text{ OC}$) and GG ($0.06 \pm 0.02 \text{ mg g}^{-1} \text{ OC}$). Series of *n*-alkanols (C_{18} – C_{32}) presented a strong even-over-odd predominance with the concentration of $0.66 \pm 0.18 \text{ mg g}^{-1} \text{ OC}$ (NG), $0.50 \pm 0.13 \text{ mg g}^{-1} \text{ OC}$ (DG), $0.32 \pm 0.19 \text{ mg g}^{-1} \text{ OC}$ (RG) and $0.19 \pm 0.08 \text{ mg g}^{-1} \text{ OC}$ (GG). Five ω -hydroxyalkanoic acids (C_{22} to C_{28}) were detected with the concentrations of $0.63 \pm 0.24 \text{ mg g}^{-1} \text{ OC}$ in NG, $0.26 \pm 0.16 \text{ mg g}^{-1} \text{ OC}$ in GG, $0.23 \pm 0.13 \text{ mg g}^{-1} \text{ OC}$ in DG and $0.16 \pm 0.19 \text{ mg g}^{-1} \text{ OC}$ in RG. These compounds were abundant in suberin of roots and bark of higher plants (Otto and Simpson, 2006b; Pisani et al., 2013). Steroids including β -sitosterol, cholesterol, campesterol and stigmasterol were only minor components ($< 0.21 \pm 0.14 \text{ mg g}^{-1} \text{ OC}$). Cholesterol is of a mixture origin from plants, fungi and animals, while other steroids are predominantly derived from higher plants (Otto et al., 2005 and ref-

erences therein). Several free carbohydrates including glucose, mannose and sucrose were detected and their abundance was lower than $0.30 \pm 0.46 \text{ mgg}^{-1} \text{ OC}$.

3.3 Compositions and distributions of bound lipids

Homologous aliphatic lipids (alkanoic acids, alkanols, alkanedioic acids, hydroxyalkanoic acids) and steroids were major constituents of the base hydrolysis products (Figs. 3 and S2). These bound aliphatic lipids are mainly derived from the biomacromolecular suberin and cutin of vascular plants (Kögel-Knabner, 2002; Nierop et al., 2003; Otto et al., 2005). Among them, *n*-alkanoic acids ranging from C_{14} to C_{32} and with C_{max} at C_{16} , mono- and di-unsaturated *n*- C_{16} and *n*- C_{18} acids are dominant with the total concentrations of $7.1 \pm 0.9 \text{ mgg}^{-1} \text{ OC}$ (RG) to $13.4 \pm 2.1 \text{ mgg}^{-1} \text{ OC}$ (NG). Branched alkanoic acids including *iso*- C_{14} , *iso*- C_{15} , *iso*- C_{16} , *iso*- C_{17} and *iso*- C_{18} were also detected and varied in concentrations from $1.05 \pm 0.23 \text{ mgg}^{-1} \text{ OC}$ (GG) to $2.28 \pm 0.37 \text{ mgg}^{-1} \text{ OC}$ (NG). Branched alkanoic acids, along with short-chain *n*-alkanoic acids ($< C_{20}$) reflect inputs from soil microbes since these compounds are usually not biosynthesized by higher plants (e.g., Otto et al., 2005). After *n*-alkanoic acids, ω -hydroxyalkanoic acids from C_{16} to C_{26} and with C_{max} at C_{20} or C_{22} were the secondarily most abundant type of biomarkers with the concentration of $1.85 \pm 1.80 \text{ mgg}^{-1} \text{ OC}$ (DG) to $5.05 \pm 2.07 \text{ mgg}^{-1} \text{ OC}$ (GG). The α, ω -alkanedioic acids including C_8 , C_9 , C_{10} , C_{11} and long-chain even numbered acids from C_{16} to C_{28} varied from 1.68 ± 0.51 to $2.38 \pm 0.38 \text{ mgg}^{-1} \text{ OC}$. The homologues of C_{16} – C_{28} α -hydroxyalkanoic acids and *n*-alkanols (C_{16} – C_{30}) were minor components of bound lipids ($< 2.0 \text{ mgg}^{-1} \text{ OC}$).

3.4 Lignin degradation products after CuO oxidation

Eight lignin derived phenols were identified in the CuO oxidation extracts of the soil residues after base hydrolysis, including three vanillyl (V; vanillin, acetovanillone, vanillic acid), three syringyl (S; syringaldehyde, acetosyringone, syringic acid) and two cinnamyl (C; *p*-coumaric acid and ferulic acid) groups (Fig. S2). The concentrations of

BGD

11, 5613–5637, 2014

Soil organic matter dynamics

L. Zhao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



total lignin phenols (V + S + C) were $9.5 \pm 2.1 \text{ mgg}^{-1} \text{ OC}$ in GG, $12.9 \pm 2.7 \text{ mgg}^{-1} \text{ OC}$ in RG, $13.3 \pm 5.1 \text{ mgg}^{-1} \text{ OC}$ in NG, and $22.2 \pm 7.7 \text{ mgg}^{-1} \text{ OC}$ in DG (Fig. 3). The ratios of C : V and S : V were calculated to estimate the source of lignins since S-unit is absent in gymnosperms, while C-unit is specific for non-woody tissues (Hedges and Mann, 1979). In addition, the ratios of acids over corresponding aldehydes (Ad/AI) are indicative of degradation stage of SOM since aldehydes are oxidized into corresponding acids with lignin degradation by white-rot and brown-rot fungi (e.g. Hedges et al., 1988). Non-woody tissues such as grasses and leaves have been reported to have higher Ad/AI values (0.2–1.6) than angiosperm and conifer wood (0.1–0.5) (Hedges et al., 1988; Otto and Simpson, 2006a). In our study, all soils showed similar S/V values ranging from 0.67 to 0.79 (Table 2), in agreement with the same vegetation cover in all grasslands. Significant differences of $(\text{Ad/AI})_V$ (0.35 ~ 1.47 in an increasing order of NG < RG < DG < GG) and $(\text{Ad/AI})_S$ (0.58 ~ 2.26 in an increasing order of NG < DG < RG < GG) were observed (Table 2), reflecting different degradation degrees of lignin.

4 Discussion

4.1 Changes in bulk soil organic matter properties under different land-use

Over recent decades, the land use such as cultivation and grazing have caused considerable reduction of vegetation cover, destruction of topsoil structure and compaction of soil in the Inner Mongolia grasslands in northern China (Cui et al., 2005; Wang et al., 2013). Consequently, degradation and even desertification have become a common phenomenon in the Inner Mongolia grasslands (Su et al., 2005). A series of strategies including grazing exclusion, planting indigenous plants, and returning farmlands to grasslands have been implemented to protect the regional environments (Jiang et al., 2006). However, the effects of these strategies on the quality and quantity of SOM are not fully understood.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Soil organic matter
dynamics

L. Zhao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



In this study, a strong correlation was observed between TOC and TN ($r = 0.999$, $n = 16$, $p < 0.01$), suggesting that nitrogen is primarily associated with SOM. There are significant differences in soil organic carbon and nitrogen contents among four sampling sites. A six-fold increase of soil organic carbon content in GG compared to NG suggests that enhanced water supply can be an effective strategy for the sequestration of organic carbon in soils. This is not surprising because the Inner Mongolia grasslands are mainly distributed in semiarid and arid regions where water is one of limiting factors for vegetation growth. In addition, increasing water content in soils can reduce oxygen diffusion rate and thereby decreases the decomposition of SOM. Relative to NG, soil organic carbon and nitrogen contents in RG is about one-fold higher, likely owing to two factors: (1) potato cultivation and fertilization for two decades (1990–2010) resulted in the accumulation of organic carbon and nitrogen in soils; and (2) banning grazing for 2 years (2010–2012) increased organic carbon inputs to soil. Among four field sites, RG is characterized by the highest $\delta^{15}\text{N}$ values ($4.5 \pm 0.4\%$; Table 1), suggesting an important effect of previous cultivation activities. In the Inner Mongolia, stock manures were commonly used as fertilizers, which has remarkably higher $\delta^{15}\text{N}$ values (10 to 20‰) than from plants (close to 0‰) (Bateman et al., 2005; Kohl et al., 1973).

In contrast to GG and RG, about 50% reduction of soil organic carbon and nitrogen contents was observed for DG, suggesting that overgrazing caused sparse vegetation cover and thereby decreased organic matter inputs to soils. In addition, with the lack of vegetation, the strong northwestern wind during Asian winter monsoon can accelerate soil erosion particularly for fine particles (e.g., Cui et al., 2005; Su et al., 2005; Zhao et al., 2007). This wind erosion effect is reflected by significantly lower clay fraction in DG ($1.7 \pm 0.9\%$) than NG ($4.6 \pm 1.6\%$; Table 1). Since SOM is enriched in fine particles (Anderson et al., 1981; Nichols, 1984), the loss of clay fractions further reduced SOM levels in DG.

The $\delta^{13}\text{C}$ value is a useful tracer for organic matter sources (e.g., C3 and C4 plants) degradation stage (Farquhar et al., 1989; O’Leary, 1981) and water use efficiency (Wang et al., 2008). In our study, the dominant grass species, *Leymus chinensis* and

Stipa grandis, are C3 plants, and thus the effect of C3/C4 plant shift on the $\delta^{13}\text{C}$ value was not considered. The degradation of soil organic matter can cause $\delta^{13}\text{C}$ values to positively shift by several per mil due to the preferential utilization of light carbon isotope by soil microbes (Feng, 2002; Natelhoffer and Fry, 1988), while the $\delta^{13}\text{C}$ values of C3 plant leaves decreases with increasing rainfall by a coefficient of ca. $-0.4\text{‰}/100\text{mm}$ (Wang et al., 2008). Among four sampling sites, GG presented significant difference in $\delta^{13}\text{C}$ values of SOM ($-26.2 \pm 0.6\text{‰}$) from other three sites (-24.2 to -25.1‰ ; Table 1), reflecting the largest inputs of fresh, undegraded organic matter due to high primary productivity or the change of water use efficiency with groundwater supply.

4.2 Molecular compositions of SOM in different grasslands

Compared to NG, DG contained significantly lower concentrations of total free lipids, cutin/suberin-monomers and lignin-phenols (Fig. 3). Such difference can be explained by two mechanisms. First, aliphatic and lignin biomarkers are preferentially degraded constituents of bulk SOM. This explanation is consistent with previous results that the turnover of lignin is faster than bulk SOM (e.g., Dignac et al., 2005; Gleixner et al., 2002; Thevenot et al., 2010) as well as soil fungi and bacteria synthesized compounds (e.g., Amelung et al., 2008). In contrast, carbohydrates such as glucose, mannose and sucrose, usually thought to be biochemically labile compounds (Lorenz et al., 2007; Schmidt et al., 2011), were significantly higher abundance in DG ($304.0 \pm 460.5 \mu\text{g g}^{-1} \text{OC}$) than NG ($56.1 \pm 21.9 \mu\text{g g}^{-1} \text{OC}$; Fig. 3). The enrichment of free carbohydrates in DG is indicative of higher proportions of microbial carbon in degraded soils, supported by their heaviest $\delta^{13}\text{C}$ values of SOM (Table 1). Our results support the hypothesis that molecular structure does not control long-term decomposition of organic matter in mineral soils (Schmidt et al., 2011 and references therein). The persistence of SOM is actually dependent on complex interactions between organic matter and its environment such as compound chemistry, climate, water availability, soil pH and soil microbial community (Schmidt et al., 2011; Thevenot et al., 2010).

BGD

11, 5613–5637, 2014

Soil organic matter dynamics

L. Zhao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



about 50% reduction of soil organic carbon and nitrogen, whereas the groundwater supply and potato cultivation-restoration activities are beneficial for the accumulation of soil organic carbon and nitrogen. With land-use changes, the root-derived aliphatic carbon (e.g., suberin) are preferentially preserved in soils compared to aboveground-derived carbon. This finding is important for prediction of global carbon cycling under climate and land-use changes since root-derived carbon (particularly aliphatic suberin) has longer resident time than aboveground carbon (Rasse et al., 2005) and other type organic carbon (e.g., lignin) in soils (Feng and Simpson, 2007; Lorenz et al., 2007). Future studies will be investigation for other types of inert carbon such as charcoal to understand the response of different soil organic matter to land-use changes.

Supplementary material related to this article is available online at <http://www.biogeosciences-discuss.net/11/5613/2014/bgd-11-5613-2014-supplement.pdf>.

Acknowledgements. This study was financially supported by the National Basic Research Program of China (2014CB954001) and the National Science Foundation of China (41006042, 41176164). We thank Chengyang Zheng and Hongyan Liu for assistance in sampling.

References

- Amelung, W., Brodowski, S., Sandhage-Hofmann, A., and Bol, R.: Combining biomarker with stable isotope analyses for assessing the transformation and turnover of soil organic matter, *Adv. Agron.*, 100, 155–250, 2008.
- Anderson, D., Sagar, S., Bettany, J., and Stewart, J.: Particle size fractions and their use in studies of soil organic matter: I. The nature and distribution of forms of carbon, nitrogen, and sulfur, *Soil Sci. Soc. Am. J.*, 45, 767–772, 1981.
- Baldock, J. A. and Skjemstad, J. O.: Role of the soil matrix and minerals in protecting natural organic materials against biological attack, *Org. Geochem.*, 31, 697–710, 2000.

BGD

11, 5613–5637, 2014

Soil organic matter dynamics

L. Zhao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Soil organic matter dynamics

L. Zhao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Bateman, A. S., Kelly, S. D., and Jickells, T. D.: Nitrogen isotope relationships between crops and fertilizer: implications for using nitrogen isotope analysis as an indicator of agricultural regime, *J. Agr. Food Chem.*, 53, 5760–5765, 2005.

Cui, X., Wang, Y., Niu, H., Wu, J., Wang, S., Schnug, E., Rogasik, J., Fleckenstein, J., and Tang, Y.: Effect of long-term grazing on soil organic carbon content in semiarid steppes in Inner Mongolia, *Ecol. Res.*, 20, 519–527, 2005.

Deleeuw, J. W., Rijpstra, W. I. C., and Nienhuis, P. H.: Free and bound fatty-acids and hydroxy fatty-acids in the living and decomposing eelgrass *zostera-marina* L., *Org. Geochem.*, 23, 721–728, 1995.

Dignac, M. F., Bahri, H., Rumpel, C., Rasse, D. P., Bardoux, G., Balesdent, J., Girardin, C., Chenu, C., and Mariotti, A.: Carbon-13 natural abundance as a tool to study the dynamics of lignin monomers in soil: an appraisal at the Closeaux experimental field (France), *Geoderma*, 128, 3–17, 2005.

Farquhar, G. D., Ehleringer, J. R., and Hubick, K. T.: Carbon isotope discrimination and photosynthesis, *Annu. Rev. Plant Biol.*, 40, 503–537, 1989.

Feng, X.: A theoretical analysis of carbon isotope evolution of decomposing plant litters and soil organic matter, *Global Biogeochem. Cy.*, 16, 1119, doi:10.1029/2002GB001867, 2002.

Feng, X., Simpson, A. J., Wilson, K. P., Dudley Williams, D., and Simpson, M. J.: Increased cuticular carbon sequestration and lignin oxidation in response to soil warming, *Nat. Geosci.*, 1, 836–839, 2008.

Feng, X. J. and Simpson, M. J.: The distribution and degradation of biomarkers in Alberta grassland soil profiles, *Org. Geochem.*, 38, 1558–1570, 2007.

Feng, X. M. and Zhao, Y. S.: Grazing intensity monitoring in Northern China steppe: integrating CENTURY model and MODIS data, *Ecol. Indic.*, 11, 175–182, 2011.

Gleixner, G., Poirier, N., Bol, R., and Balesdent, J.: Molecular dynamics of organic matter in a cultivated soil, *Org. Geochem.*, 33, 357–366, 2002.

Guo, L., Wang, M., and Gifford, R.: The change of soil carbon stocks and fine root dynamics after land use change from a native pasture to a pine plantation, *Plant Soil*, 299, 251–262, 2007.

He, N., Yu, Q., Wu, L., Wang, Y., and Han, X.: Carbon and nitrogen store and storage potential as affected by land-use in a *Leymus chinensis* grassland of northern China, *Soil Biol. Biochem.*, 40, 2952–2959, 2008.

Soil organic matter
dynamics

L. Zhao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Hedges, J. I. and Mann, D. C.: The characterization of plant tissues by their lignin oxidation products, *Geochim. Cosmochim. Ac.*, 43, 1803–1807, 1979.
- Hedges, J. I., Blanchette, R. A., Weliky, K., and Devol, A. H.: Effects of fungal degradation on the CuO oxidation products of lignin: a controlled laboratory study, *Geochim. Cosmochim. Ac.*, 52, 2717–2726, 1988.
- Islam, K. R. and Weil, R. R.: Land use effects on soil quality in a tropical forest ecosystem of Bangladesh, *Agr. Ecosyst. Environ.*, 79, 9–16, 2000.
- Janzen, H. H.: Carbon cycling in earth systems – a soil science perspective, *Agr. Ecosyst. Environ.*, 104, 399–417, 2004.
- Jiang, G., Han, X., and Wu, J.: Restoration and management of the Inner Mongolia grassland require a sustainable strategy, *Ambio*, 35, 269–270, 2006.
- Kögel-Knabner, I.: Analytical approaches for characterizing soil organic matter, *Org. Geochem.*, 31, 609–625, 2000.
- Kögel-Knabner, I.: The macromolecular organic composition of plant and microbial residues as inputs to soil organic matter, *Soil Biol. Biochem.*, 34, 139–162, 2002.
- Kaiser, M., Wirth, S., Ellerbrock, R. H., and Sommer, M.: Microbial respiration activities related to sequentially separated, particulate and water-soluble organic matter fractions from arable and forest topsoils, *Soil Biol. Biochem.*, 42, 418–428, 2010.
- Kohl, D. H., Shearer, G. B., and Commoner, B.: Variation of ^{15}N in corn and soil following application of fertilizer nitrogen, *Soil Sci. Soc. Am. J.*, 37, 888–892, 1973.
- Kolattukudy, P. E. and Espelie, K. E.: Chemistry, biochemistry, and function of suberin and associated waxes, in: *Natural Products of Woody Plants*, edited by: Rowe, J., Springer Series in Wood Science, Springer, Berlin, Heidelberg, 304–367, 1989.
- Leifeld, J. and Kögel-Knabner, I.: Soil organic matter fractions as early indicators for carbon stock changes under different land-use?, *Geoderma*, 124, 143–155, 2005.
- Lorenz, K., Lal, R., Preston, C. M., and Nierop, K. G.: Strengthening the soil organic carbon pool by increasing contributions from recalcitrant aliphatic bio (macro) molecules, *Geoderma*, 142, 1–10, 2007.
- Natelhofer, K. J. and Fry, B.: Controls on natural nitrogen-15 and carbon-13 abundances in forest soil organic matter, *Soil Sci. Soc. Am. J.*, 52, 1633–1640, 1988.
- National Statistics Bureau of China: *China Statistics Yearbook*, Beijing, 6 pp., 2002.
- Nichols, J. D.: Relation of organic carbon to soil properties and climate in the southern Great Plains, *Soil Sci. Soc. Am. J.*, 48, 1382–1384, 1984.

Soil organic matter
dynamics

L. Zhao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Nierop, K. G. J., Naafs, D. F. W., and Verstraten, J. M.: Occurrence and distribution of ester-bound lipids in Dutch coastal dune soils along a pH gradient, *Org. Geochem.*, 34, 719–729, 2003.

O'Leary, M. H.: Carbon isotope fractionation in plants, *Phytochemistry*, 20, 553–567, 1981.

5 Otto, A. and Simpson, M. J.: Evaluation of CuO oxidation parameters for determining the source and stage of lignin degradation in soil, *Biogeochemistry*, 80, 121–142, 2006a.

Otto, A. and Simpson, M. J.: Sources and composition of hydrolysable aliphatic lipids and phenols in soils from western Canada, *Org. Geochem.*, 37, 385–407, 2006b.

10 Otto, A., Shunthirasingham, C., and Simpson, M. J.: A comparison of plant and microbial biomarkers in grassland soils from the Prairie Ecozone of Canada, *Org. Geochem.*, 36, 425–448, 2005.

Pisani, O., Hills, K. M., Courtier-Murias, D., Simpson, A. J., Mellor, N. J., Paul, E. A., Morris, S. J., and Simpson, M. J.: Molecular level analysis of long term vegetative shifts and relationships to soil organic matter composition, *Org. Geochem.*, 62, 7–16, 2013.

15 Quenea, K., Derenne, S., Largeau, C., Rumpel, C., and Mariotti, A.: Variation in lipid relative abundance and composition among different particle size fractions of a forest soil, *Org. Geochem.*, 35, 1355–1370, 2004.

Rasse, D., Rumpel, C., and Dignac, M.-F.: Is soil carbon mostly root carbon? Mechanisms for a specific stabilisation, *Plant Soil*, 269, 341–356, 2005.

20 Schlesinger, W. H. and Andrews, J. A.: Soil respiration and the global carbon cycle, *Biogeochemistry*, 48, 7–20, 2000.

Schmidt, M. W. I., Torn, M. S., Abiven, S., Dittmar, T., Guggenberger, G., Janssens, I. A., Kleber, M., Kogel-Knabner, I., Lehmann, J., Manning, D. A. C., Nannipieri, P., Rasse, D. P., Weiner, S., and Trumbore, S. E.: Persistence of soil organic matter as an ecosystem property, *Nature*, 478, 49–56, 2011.

25 Simoneit, B. R.: A review of current applications of mass spectrometry for biomarker/molecular tracer elucidations, *Mass Spectrom. Rev.*, 24, 719–765, 2005.

Su, Y.-Z., Li, Y.-L., Cui, J.-Y., and Zhao, W.-Z.: Influences of continuous grazing and livestock exclusion on soil properties in a degraded sandy grassland, Inner Mongolia, northern China, *Catena*, 59, 267–278, 2005.

30 Sun, D. Y., Tan, W. B., Pei, Y. D., Zhou, L. P., Wang, H., Yang, H., and Xu, Y. P.: Late Quaternary environmental change of Yellow River Basin: an organic geochemical record in Bohai Sea (North China), *Org. Geochem.*, 42, 575–585, 2011.

Soil organic matter
dynamics

L. Zhao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Tan, W. B., Zhou, L. P., and Liu, K. X.: Soil aggregate fraction-based C-14 analysis and its application in the study of soil organic carbon turnover under forests of different ages, *Chinese Sci. Bull.*, 58, 1936–1947, 2013.

Thevenot, M., Dignac, M.-F., and Rumpel, C.: Fate of lignins in soils: a review, *Soil Biol. Biochem.*, 42, 1200–1211, 2010.

Wang, G., Feng, X., Han, J., Zhou, L., Tan, W., and Su, F.: Paleovegetation reconstruction using $\delta^{13}\text{C}$ of Soil Organic Matter, *Biogeosciences*, 5, 1325–1337, doi:10.5194/bg-5-1325-2008, 2008.

Wang, Z.-P., Han, X.-G., Chang, S. X., Wang, B., Yu, Q., Hou, L.-Y., and Li, L.-H.: Soil organic and inorganic carbon contents under various land uses across a transect of continental steppes in Inner Mongolia, *Catena*, 109, 110–117, 2013.

Watson, R. T.: *Land Use, Land-Use Change, and Forestry: a Special Report of the Intergovernmental Panel on Climate Change*, Cambridge University Press, 2000.

Wiesenberg, G. L. B., Schmidt, M. W. I., and Schwark, L.: Plant and soil lipid modifications under elevated atmospheric CO_2 conditions: I. Lipid distribution patterns, *Org. Geochem.*, 39, 91–102, 2008.

Wu, W., Ruan, J., Ding, S., Zhao, L., Xu, Y., Yang, H., Ding, W., and Pei, Y.: Source and distribution of glycerol dialkyl glycerol tetraethers along lower Yellow River-estuary-coast transect, *Mar. Chem.*, 158, 17–26, 2014.

Xiao, X., Ojima, D., Parton, W., Chen, Z., and Chen, D.: Sensitivity of Inner Mongolia grasslands to climate change, *J. Biogeogr.*, 22, 643–648, 1995.

Zhao, Y., Peth, S., Krümmelbein, J., Horn, R., Wang, Z., Steffens, M., Hoffmann, C., and Peng, X.: Spatial variability of soil properties affected by grazing intensity in Inner Mongolia grassland, *Ecol. Model.*, 205, 241–254, 2007.

Zhou, Z., Sun, O., Huang, J., Li, L., Liu, P., and Han, X.: Soil carbon and nitrogen stores and storage potential as affected by land-use in an agro-pastoral ecotone of northern China, *Biogeochemistry*, 82, 127–138, 2007.

Soil organic matter dynamics

L. Zhao et al.

Table 1. Grain size distribution, total organic carbon (TOC), total nitrogen (TN), $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values of soils from DG (degraded grassland), NG (native grassland), GG (groundwater-sustaining grassland) and RG (restored grassland) in Inner Mongolia.

	Clay ¹ (%)	Silt (%)	Sand (%)	TOC (g kg ⁻¹)	TN (g kg ⁻¹)	$\delta^{13}\text{C}$ (‰)	$\delta^{15}\text{N}$ (‰)
DG	1.7 ± 0.9 ² c	11.0 ± 0.8b	87.3 ± 5.6a	3.2 ± 0.2d	0.32 ± 0.15d	-24.2 ± 0.6a	4.14 ± 0.50a
NG	4.6 ± 1.6b	21.6 ± 7.8a	73.8 ± 9.4b	6.6 ± 0.25c	0.65 ± 0.25c	-24.9 ± 0.6a	3.31 ± 0.45b
GG	5.3 ± 0.7a, b	26.0 ± 4.5a	68.8 ± 5.1b	46.6 ± 16.7a	3.9 ± 1.5a	-26.2 ± 0.6b	2.77 ± 0.27b
RG	6.4 ± 1.0a	29.2 ± 5.6a	64.4 ± 6.5b	15.9 ± 3.4b	1.3 ± 0.03b	-25.1 ± 0.1a	4.51 ± 0.37a

¹ Clay (< 2 μm), silt (2 ~ 200 μm) and sand (200 ~ 2000 μm).

² All values are expressed as means of four replicates ± standard errors.

* Different letters (a, b, c, d) in column indicate significant difference ($P < 0.05$).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Soil organic matter dynamics

L. Zhao et al.

Table 2. Source and degradation parameters of lignin phenols in grassland soils in Inner Mongolia. DG: degraded grassland by overgrazing; NG: native grassland without apparent anthropogenic disturbance; GG: groundwater-sustaining grassland; RG: restored grassland from previous cultivated land.

Fields	C/V	S/V	(Ad/Al) _V	(Ad/Al) _S
DG	0.26 ± 0.04b	0.75 ± 0.29a	1.43 ± 0.55a	0.59 ± 0.49c
NG	0.24 ± 0.02b	0.67 ± 0.05a	0.58 ± 0.07b	0.35 ± 0.03c
GG	0.47 ± 0.16a	0.79 ± 0.4a	2.26 ± 0.99a	1.47 ± 0.32a
RG	0.28 ± 0.04b	0.71 ± 0.09a	0.98 ± 0.13a	0.76 ± 0.09b

* Values are expressed as mean ± SD of four determinations. Different letters (a, b, c, d) indicate significant difference ($P < 0.05$).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I ◀

▶ I

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Soil organic matter dynamics

L. Zhao et al.

Table 3. Enrichment factors (Ec) of biomarkers in the grassland soils (Inner Mongolia) under different land use practices. DG: degraded grassland by overgrazing; NG: native grassland without apparent anthropogenic disturbance; GG: groundwater-sustaining grassland; RG: restored grassland from previous cultivated land. FA = fatty acids.

Field	Free lipids	ω -FA	α -FA	Lignin
DG	0.63 ± 0.29b	0.62 ± 0.60c	0.31 ± 0.18b	0.90 ± 0.03b
NG	1.00 ± 0.33a	1.00 ± 1.52b	1.00 ± 0.53a	1.00 ± 0.06a
GG	0.34 ± 0.18c	1.69 ± 0.69a	1.24 ± 0.43a	0.64 ± 0.14c
RG	0.39 ± 0.21c	1.66 ± 0.58a	1.18 ± 0.37a	0.87 ± 0.18b

* Values are expressed as mean ± SD of four determinations. Different letters (a, b, c, d) indicate significant difference ($P < 0.05$).

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[I ◀](#)
[▶ I](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)





Fig. 1. Pictures for Inner Mongolia grasslands, DG: degraded grassland by overgrazing; NG: native grassland without apparent anthropogenic disturbance; GG: groundwater-sustaining grassland; RG: restored grassland from potato cropland.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Soil organic matter dynamics

L. Zhao et al.




Fig. 2. Flow chart of bulk parameters and molecular biomarker analyses for grassland soils, Inner Mongolia. FA = fatty acids.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)





Fig. 3. Concentrations ($\text{mg g}^{-1} \text{OC}$) of free lipids, bound lipids and lignin-phenols in the grassland soils in Inner Mongolia. DG: degraded grassland by overgrazing; NG: native grassland without apparent anthropogenic disturbance; GG: groundwater-sustaining grassland; RG: restored grassland from previous cultivated land. The values presented are means of four replicates \pm standard errors. Different letters (a, b, c, d) above each column bar indicate significant difference ($P < 0.05$).

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)
