



This discussion paper is/has been under review for the journal Biogeosciences (BG).  
Please refer to the corresponding final paper in BG if available.

# Factors controlling shell carbon isotopic composition of land snail *Acusta despecta sieboldiana* estimated from lab culturing experiment

N. Zhang<sup>1</sup>, K. Yamada<sup>1</sup>, N. Suzuki<sup>2</sup>, and N. Yoshida<sup>1,3</sup>

<sup>1</sup>Department of Environmental Chemistry and Engineering, Tokyo Institute of Technology, Yokohama, Japan

<sup>2</sup>Department of Geology and Paleontology, National Museum of Nature and Science, Japan

<sup>3</sup>Earth-Life Science Institute, Tokyo Institute of Technology, Tokyo, Japan

Received: 8 April 2014 – Accepted: 20 April 2014 – Published: 6 May 2014

Correspondence to: N. Zhang (zhang.n.aa@m.titech.ac.jp)

Published by Copernicus Publications on behalf of the European Geosciences Union.

BGD

11, 6555–6590, 2014

Factors controlling  
shell carbon isotopic  
composition of land  
snail

N. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Abstract

The carbon isotopic composition ( $\delta^{13}\text{C}$ ) of land snail shell carbonate derives from three potential sources: diet, atmospheric  $\text{CO}_2$ , and ingested carbonate (limestone). However, their relative contributions remain unclear. Under various environmental conditions, we cultured one land snail species, *Acusta despecta sieboldiana* collected from Yokohama, Japan, and confirmed that all of these sources affect shell carbonate  $\delta^{13}\text{C}$  values. Herein, we consider the influences of metabolic rates and temperature on the carbon isotopic composition of the shell carbonate. Based on previous works and on results obtained in this study, a simple but credible framework is presented for discussion of how each source and environmental parameter can affect shell carbonate  $\delta^{13}\text{C}$  values. According to this framework and some reasonable assumptions, we have estimated the contributions of different carbon sources for each snail individual: for cabbage ( $\text{C}_3$  plant) fed groups, the contributions of diet, atmospheric  $\text{CO}_2$  and ingested limestone respectively vary as 66–80 %, 16–24 %, and 0–13 %. For corn ( $\text{C}_4$  plant) fed groups, because of the possible food stress (lower consumption ability of  $\text{C}_4$  plant), the values vary respectively as 56–64 %, 18–20 %, and 16–26 %. Moreover, we present new evidence that snails have discrimination to choose  $\text{C}_3$  and  $\text{C}_4$  plants as food. Therefore, we suggest that food preferences must be considered adequately when applying  $\delta^{13}\text{C}$  in paleo-environment studies. Finally, we inferred that, during egg laying and hatching of our cultured snails, carbon isotope fractionation is controlled only by the isotopic exchange of the calcite– $\text{HCO}_3^-$ –aragonite equilibrium.

## 1 Introduction

Land snail shells are widely applied for studying paleo and present environment characteristics because they can be well preserved in Quaternary fossils and because most species still exist today (e.g., Yapp, 1979; Lecolle, 1985; Goodfriend et al., 1989, 1992; Colonese et al., 2007, 2010, 2011; Yanes et al., 2008a, b, 2009, 2011, 2012a, b, 2013;

BGD

11, 6555–6590, 2014

### Factors controlling shell carbon isotopic composition of land snail

N. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Kehrwald et al., 2010; Zaarur et al., 2011). Specifically, the carbon isotopic composition ( $\delta^{13}\text{C}$  value) of land snail shell carbonate is regarded as useful to reconstruct the distribution of terrestrial  $\text{C}_3/\text{C}_4$  vegetation, which itself is related to some environmental parameters such as rainfall amounts and temperature variations (Goodfriend and Magaritz, 1987; Goodfriend, 1992; Stott, 2002; Metref et al., 2003).

Land snail shell carbon has three sources: food, atmospheric  $\text{CO}_2$ , and ingested carbonate (limestone) (Goodfriend and Hood, 1983). Food sources such as plants, fungi, and other organic matter can be transformed into metabolic  $\text{CO}_2$  in the body of a land snail by two pathways: direct digestion and breakdown of urea (Stott, 2002). The resulting  $\text{CO}_2$  will dissolve into the bicarbonate pool in the hemolymph and then precipitate as shell carbonate. Atmospheric  $\text{CO}_2$  can be introduced into the bicarbonate pool via respiration. Similarly, ingested carbonate can first dissolve into stomach acid to produce gaseous  $\text{CO}_2$ . Then it can be introduced into the bicarbonate pool, too. Some previous works have been undertaken to clarify their relative contributions both from observations and model calculations. Nevertheless, this topic remains poorly understood (Goodfriend and Hood, 1983; Goodfriend and Stipp, 1983; Stott, 2002; Metref et al., 2003; Balakrishnan and Yapp, 2004; Romaniello et al., 2008). Following are estimations of each carbon source based on published works: food source ratios vary as 25–40 % (Goodfriend and Hood, 1983), 36–73 % (Romainello et al., 2008), and 100 % (Stott, 2002; Metref et al., 2003); atmospheric  $\text{CO}_2$  has been estimated as negligible (Stott, 2002; Metref et al., 2003), 16–48 % (Romaiello et al., 2008), and 30–60 % (Goodfriend and Hood, 1983); ingested carbonate has been inferred as negligible (Stott, 2002), up to 30 % (Goodfriend and Stipp, 1983), often negligible for small terrestrial gastropods of less than 10 mm, and as always much less than 20–30 % for larger species (Pigati et al., 2004, 2010), ~ 20 % up to ~ 40 % (Yanes et al., 2008a).

For solving problems of this kind, more studies including lab culturing experiments must be done. Stott (2002) and Meterf et al. (2003) reported two independent works related to land snail culture experiments. Both show a marked but discrepant correlation between land snail shell carbonate  $\delta^{13}\text{C}$  and diet  $\delta^{13}\text{C}$  with slopes less than one.

**BGD**

11, 6555–6590, 2014

## Factors controlling shell carbon isotopic composition of land snail

N. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Factors controlling shell carbon isotopic composition of land snail

N. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Secondly, Stott pointed out that ingested carbonate does not contribute to shell carbonate  $\delta^{13}\text{C}$  values based on results of snails fed with and without an added  $\text{CaCO}_3$  source. Finally, according to their calculations and discussions, both papers reported that atmospheric  $\text{CO}_2$  does not contribute to shell carbonate  $\delta^{13}\text{C}$  values. Those reports notwithstanding, inconsistent observations and discussions emerged soon thereafter. Yanes et al. (2012a) reported that higher  $\delta^{13}\text{C}$  values were observed during the younger growth stages of both living and fossil snails, which showed signs of a higher contribution of ingested limestone. Results of this research suggest strongly that environmental carbonate is incorporated as an important source for precipitating land snail shells (at least some snail species) and suggest that it can affect their shell  $\delta^{13}\text{C}$  values. To the atmospheric  $\text{CO}_2$ , after summarizing the previous studies of relation between land snail shell carbonate  $\delta^{13}\text{C}$  and shell organics/body tissue organics/diet (Stott, 2002; Goodfriend and Ellis, 2002; Balakrishnan and Yapp, 2004), McConnaughey and Gillikin (2008) pointed out that the offsets between  $\delta^{13}\text{C}$  of shell carbonate and their diet (or body organics) are greater when the  $\delta^{13}\text{C}$  of the latter diverge more from atmospheric  $\text{CO}_2$ , suggesting that the atmospheric  $\text{CO}_2$  does contribute to shell carbonate.

A better understanding of the contribution of each carbon source can promote this isotopic tool in the field of paleo environment reconstruction. Therefore, we cultured one land snail species (*Acusta despecta sieboldiana*) under different controlled conditions. A simple but credible framework was raised to discuss the mechanism of how each possible source and environmental parameter can affect shell carbonate  $\delta^{13}\text{C}$  values based on previous works and results of this study. According to this frame and some reasonable assumptions, we estimated the contribution of different carbon sources for each snail individual. This report is the first describing an attempt to estimate the contributing proportion of limestone ingested by snail individuals using stable isotope values compared with a previous method using calculations including radiocarbon dating (Goodfriend and Stipp, 1983; Pigati et al., 2010).



## 2 Materials and methods

### 2.1 Culturing of land snails

Eight adult snails of *Acusta despecta sieboldiana* were collected in Suzukakedai, Yokohama, Japan and were cultured at room temperature (ca. 25 °C) from January 2012. These snails began to lay eggs in March 2012. Then the eggs were transferred into a stainless steel container and were covered with moist cloth in a 25 °C incubator. Most eggs hatched at around 3–4 weeks.

Larvae were distributed randomly into around 30 small transparent plastic boxes that had been perforated to allow air and vapor exchange: each box contained 3–5 snails. Some later hatched larvae were added to May and June 2012. Then four small boxes were put into semi-sealed big plastic boxes: two parallel groups were fed cabbage ( $C_3$  plant,  $\delta^{13}C = -28.4 \pm 1.2\text{‰}$ ,  $n = 12$ ) that had been sprinkled with fine calcium carbonate powder ( $CaCO_3$ ,  $\delta^{13}C = 4.0\text{‰}$ ). Another two were fed cabbage sprinkled with calcium phosphate ( $Ca_3(PO_4)_2$ ). A NaCl saturated solution was used to produce high humidity conditions in each large box. The snails grew under natural light/dark cycles. The air and food were changed every two days. Then the big plastic boxes were put into three incubators with respective temperatures of 20 °C, 25 °C, and 30 °C, until January 2013; some snails from 20 °C group were cultured until May 2013. The growth phases of snails are presented in Fig. 1.

We also cultured some snails using corn ( $C_4$  plant,  $\delta^{13}C = -12.0 \pm 0.7\text{‰}$ ,  $n = 4$ ; first we used *Miscanthus sinensis*, but the snails did not eat it; then we changed the food to corn) sprinkled with fine calcium carbonate powder in a 20 °C incubator.

Ultimately, snails were collected into labeled sampling bottles and were preserved at -4 °C for additional treatment.

BGD

11, 6555–6590, 2014

### Factors controlling shell carbon isotopic composition of land snail

N. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 2.2 Isotopic samples preparation and analysis

Frozen snails were dried using a cryogenic vacuum line. They were then washed with distilled water. The body tissues of respective snails were separated from shells using a nipper. Then they were immersed into 3N HCl for 2–4 h, rinsed with distilled water and then lyophilized. All dry tissues were ground to powder and wrapped in tinfoil (Sn) capsules. Each sample was introduced into a combustion tube from the auto-sampler and converted into gaseous CO<sub>2</sub> at 980 °C one by one. Then the resultant CO<sub>2</sub> was injected into a Cavity Ring-Down Spectroscopy (CRDS, L1121-I; Picarro, Inc., Santa Clara, CA, USA) for  $\delta^{13}\text{C}$  measurement. More details describing the CRDS method and system are presented by Wahl et al. (2006) and Balslev-Clausen et al. (2013). All of the measured  $\delta^{13}\text{C}$  values were normalized against two simultaneously measured standards, acetanilide ( $\delta^{13}\text{C} = -33.62\text{‰}$ ; Costech Analytical Technologies, Inc., Valencia, CA, USA) and sucrose ( $\delta^{13}\text{C} = -13.55\text{‰}$ ; Kanto Chemical Co., Inc., Tokyo, Japan) by two-point calibration method (Coplen et al., 2006). The analytical precision was better than 0.4 ‰ ( $n = 3$ ).

Shells were washed successively with distilled water, then with acetone under an ultrasonic bath for 60 min to eliminate the organic residues. After washing again using distilled water, they were put into labeled sampling bottles containing diluted hydrogen peroxide (10 %) to remove the remaining organic matter. These bottles were kept overnight in a shaking machine with speed around 100 r/min. Finally, all the shells were dried using lyophilization and were crushed into powder using an agate mortar for isotopic analyses. To the collected adults and cultured snails, their shell powder (7–10 mg) was reacted with 103 % phosphoric acid for more than 2 h in a 25 °C water bath. Then the purified CO<sub>2</sub> was analyzed using isotope ratio mass spectrometry (MAT 253; Thermo Fisher Scientific, Inc., Waltham, MA, USA) and calibrated against an synthetic calcium carbonate standard, Wako ( $\delta^{13}\text{C} = -9.13\text{‰}$ ; Wako Pure Chemical Industries, Ltd., Osaka, Japan). The precision was better than 0.1 ‰ ( $n = 3$ ).

BGD

11, 6555–6590, 2014

### Factors controlling shell carbon isotopic composition of land snail

N. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

To the larvae and eggshell, the powder ( $400 \pm 50 \mu\text{g}$ ) was reacted with 103 % phosphoric acid for more than 1 h at  $72^\circ\text{C}$ . The resultant  $\text{CO}_2$  was analyzed using an isotope ratio mass spectrometer (DELTA XL; ThermoFisher Scientific, Inc., Waltham, MA, USA) coupled with a PAL autosampler (GCPAL; CTC Analytics, Zwingen, Switzerland), and a GasBench II preparation device (Thermo Fisher Scientific, Inc., Waltham, MA, USA). The  $\delta^{13}\text{C}$  values were normalized against two simultaneously measured synthetic calcium carbonate standards, Wako ( $\delta^{13}\text{C} = -9.13\text{‰}$ ; Wako Pure Chemical Industries, Ltd., Osaka, Japan) and Kanto ( $\delta^{13}\text{C} = -20.62\text{‰}$ ; Kanto Chemical Co., Inc., Tokyo, Japan) by two-point calibration method. The analytical precision was around 0.1 ‰ ( $n = 3$ ).

All of our in-house working standards mentioned here are calibrated with respect to the IAEA standards such as NBS-19 and the  $\delta^{13}\text{C}$  values are reported relative to VPDB (Vienna Pee Dee belemnite).

In addition, the crystal structure of shell carbonate powder was checked using X-ray diffractometry (XRD, MXP3TA; Mac Science Ltd.). The results revealed that our snail shell carbonate is aragonite, although the eggshell carbonate is calcite.

## 3 Results

### 3.1 Summary of culturing experiment

Figure 2a shows that snails that grew at low temperature ( $20^\circ\text{C}$ ) have a higher survival rate (72 %) than those at  $30^\circ\text{C}$  (16 %). The reason probably is that, in high-temperature groups, the humidity is slightly higher so that the leaves are more easily putrified. Because the cultured environment is a sealed system with an air exchange per two days, the perishable leaves consumed oxygen and perhaps produced noxious gases which might have caused all the snails in one small box to have died. Therefore, these snails show higher mortality.

BGD

11, 6555–6590, 2014

## Factors controlling shell carbon isotopic composition of land snail

N. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Factors controlling shell carbon isotopic composition of land snail

N. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Egonmwan (2008) reported a positive correlation between calcium provision, snail body weight and shell length. High mortality was observed for snails that were deprived of a calcium source. Similarly, in our study, snails fed  $\text{Ca}_3(\text{PO}_4)_2$  exhibited lower weight and shorter length. They were more translucent, more fragile, and had thinner shells, than snails fed  $\text{CaCO}_3$  during our cultivation. Moreover, Fig. 2b shows a lower survival rate (30 %) of snails fed  $\text{Ca}_3(\text{PO}_4)_2$  than those fed  $\text{CaCO}_3$  (5 %), which is also consistent with results of the previous study, suggesting that calcium from the powder of  $\text{Ca}_3(\text{PO}_4)_2$  was ingested only to a slight degree by land snails.

For the cabbage groups, shell weight proportions of the snails, which is equal to the mass of dry shell divide by their total body mass, cultured with  $\text{CaCO}_3$  at different temperature are:  $27.5 \pm 2.7\%$  ( $20^\circ\text{C}$ ),  $31.1 \pm 6.1\%$  ( $25^\circ\text{C}$ ),  $17.9 \pm 5.5\%$  ( $30^\circ\text{C}$ ); those without  $\text{CaCO}_3$  are:  $7.8 \pm 2.9\%$  ( $20^\circ\text{C}$ ),  $5.7 \pm 1.8\%$  ( $25^\circ\text{C}$ ), respectively. The discrepancy also reveals worse growth conditions at  $30^\circ\text{C}$  among the temperatures discussed above. Those without  $\text{CaCO}_3$  can get only a very small amount of calcium from either  $\text{Ca}_3(\text{PO}_4)_2$ , drinking water or diet. To the corn groups, preliminary results demonstrate that the snails probably have a higher shell weight proportion than those fed with cabbage (*t* test,  $p < 0.01$ ):  $34.7 \pm 0.3\%$  (corn,  $\text{CaCO}_3$ ,  $20^\circ\text{C}$ ) vs.  $27.5 \pm 2.7\%$  (cabbage,  $\text{CaCO}_3$ ,  $20^\circ\text{C}$ ).

### 3.2 Carbon isotope results

Body tissues of cultured snails yield a similar  $\delta^{13}\text{C}$  value with same food source (Table 1), which are  $-27.9 \pm 0.6\text{‰}$  (fed cabbage,  $n = 38$ ) and  $-10.8 \pm 0.7\text{‰}$  (fed corn,  $n = 3$ ). No marked differences were found among temperatures or between those fed with and without  $\text{CaCO}_3$ .

Shell carbonate of cultured snails fed cabbage and  $\text{CaCO}_3$  under  $20^\circ\text{C}$ ,  $25^\circ\text{C}$ , and  $30^\circ\text{C}$  respectively yielded  $\delta^{13}\text{C}$  values of  $-9.7 \pm 0.5\text{‰}$ ,  $-10.4 \pm 0.8\text{‰}$  and  $-13.2 \pm 0.8\text{‰}$  (Table 1., Fig. 6a). Those fed cabbage and  $\text{Ca}_3(\text{PO}_4)_2$  respectively yielded values of  $-12.5 \pm 0.4\text{‰}$ ,  $-13.7 \pm 0.7\text{‰}$ ,  $-14.2\text{‰}$ . Snails fed corn and  $\text{CaCO}_3$  at  $20^\circ\text{C}$  yielded a value of  $3.9 \pm 0.5\text{‰}$ .

We also measured the  $\delta^{13}\text{C}$  values of collected adult snails, newly hatched snails, and eggshells, the results are, respectively,  $-11.5 \pm 1.2\text{‰}$  ( $n = 6$ ),  $-11.0 \pm 0.1\text{‰}$  (mixed samples,  $n > 10$ ),  $-13.1 \pm 0.9\text{‰}$  (two groups of eggs, for each group,  $n > 10$ ).

## 4 Discussion

### 4.1 Land snail shell carbonate precipitation and carbon source estimation

The solid line of Fig. 3 shows a model of diet-controlled snail shell carbonate  $\delta^{13}\text{C}$  (Francey, 1983; Stott, 2002), whereas the dashed line shows a flux balance model considering both metabolic  $\text{CO}_2$  and atmospheric  $\text{CO}_2$  based on  $\text{CO}_2$  diffusion (Balakrishnan and Yapp, 2004). However, additional enrichments were observed in both the published results and this study (e.g. snails cultured at  $20^\circ\text{C}$  and  $25^\circ\text{C}$  with carbonate, this study), suggesting a contribution from ingested carbonate. In addition, because a common model seems unsuitable for all snails because of the different metabolic rates among snail individuals growing up at different conditions (Barnhart and McMahon, 1987), the estimation of shell carbon sources for each snail is expected to be useful and necessary.

The main sources of shell carbon are from diet, atmospheric  $\text{CO}_2$  and ingested carbonate (Goodfriend and Hood, 1983). A simple framework of the possible mechanisms to precipitate shell carbonate is presented in Fig. 4 based on results of a previous study as well as results from the present study. Almost all of these three carbon sources are expected to transform into gaseous  $\text{CO}_2$  and then dissolve into the hemolymph of snail (so-called bicarbonate pool) for reaching isotopic equilibrium with bicarbonate ( $\text{HCO}_3^-$ ). The fractionation between gaseous  $\text{CO}_2$  and  $\text{HCO}_3^-$  under equilibrium is controlled by temperature, shown as  $\Delta^{13}\text{C}_{\text{HCO}_3^-(\text{g})} = -(0.114 \pm 0.003)T(^{\circ}\text{C}) + (10.78 \pm 0.05)\text{‰}$  (Mook et al., 1974; Zhang et al., 1995; Szaran, 1997). Therefore, the carbon isotope fractionations under our culturing temperatures might be  $8.5\text{‰}$ ,  $7.9\text{‰}$ , and  $7.4\text{‰}$ , respectively,

BGD

11, 6555–6590, 2014

## Factors controlling shell carbon isotopic composition of land snail

N. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Factors controlling shell carbon isotopic composition of land snail

N. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



at 20 °C, 25 °C, and 30 °C. Finally, the shell aragonite will precipitate from bicarbonate pool in isotopic equilibrium (also designated as carbon isotope steady state, Balakrishnan and Yapp, 2004) and the carbon isotope ratio might be enriched to around 2.7 ‰ (Rubinson and Clayton, 1969; Romanek et al., 1991). Therefore, the total fractionation between each carbon source (diet, atmospheric CO<sub>2</sub> and ingested carbonate) and shell aragonite is expected to be an enrichment of 11.2 ‰, 10.6 ‰, and 10.1 ‰, respectively, at 20 °C, 25 °C, and 30 °C. In addition, trace amounts of DIC dissolved in leaf water and drinking water might also introduce isotopic fractionations of shell carbonate. For instance, Pigati et al. (2004), based on radiocarbon dating technique, reported that aqueous carbon sources account for approximately 10 % of the shell carbon for one species of *Catinella*, which is a semi-aquatic gastropod. However, no report to date describes a study of this source related to terrestrial gastropods.

For simplification of the model and calculations, we made two assumptions. First, no carbon isotopic fractionation takes place when the ingested carbonate dissolves into the water and reacts with stomach acid. Second, because of the variable sources and probably low contribution of DIC from directly ingested water, which are similar to all earlier published discussions, we will not involve such sources in our calculation. These assumptions can be evaluated for their propriety in future research. Results of those studies can be expected to improve the accuracy of this estimation method.

According to this framework, we will explain briefly how to calculate the contribution of each source to shell carbon based on the snail data we measured. First, from the mass balance model, we obtain the following.

$$\delta^{13}\text{C}_s = x\delta^{13}\text{C}_t + y\delta^{13}\text{C}_a + z\delta^{13}\text{C}_c + b \quad (1)$$

$$x + y + z = 1 \quad (2)$$

Therein,  $\delta^{13}\text{C}_s$  = the isotopic composition of shell aragonite.

$\delta^{13}\text{C}_t$  = the isotopic composition of snail tissue. Based on our observations and discussions, see Sect. 4.2.2, we use the measurements for each snail, not an average value, because of the individual differences among snails.

# Factors controlling shell carbon isotopic composition of land snail

N. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



$\delta^{13}\text{C}_a$  = the isotopic composition of atmospheric air ( $\delta^{13}\text{C}$  of atmospheric  $\text{CO}_2$  was observed from June 2010 to July 2011 in Suzukakedai, Yokohama, Japan. The average value is  $-9.5\text{‰}$  (Zhang et al., 2014,  $n = 42$ ); the year decrease rate reported by Keeling et al. (2010) is approximately  $0.02\text{‰}\cdot\text{year}^{-1}$ . Therefore, we use  $-9.5\text{‰}$  as the  $\delta^{13}\text{C}_a$  value).

$\delta^{13}\text{C}_c$  = the isotopic composition of ingested carbonate ( $4.0\text{‰}$ ).

$x$  = proportion of metabolic  $\text{CO}_2$ ;  $y$  = proportion of atmospheric  $\text{CO}_2$ ;  $z$  = proportion of  $\text{CO}_2$  from ingested carbonate.

$b$  = isotope fractionation value from gaseous  $\text{CO}_2$  to aragonite at different temperatures ( $11.2\text{‰}$ ,  $10.6\text{‰}$ , and  $10.1\text{‰}$ , at  $20^\circ\text{C}$ ,  $25^\circ\text{C}$ , and  $30^\circ\text{C}$ , respectively).

Here we chose  $20^\circ\text{C}$  group snails to demonstrate how we calculated the contribution of each source. When there is no  $\text{CaCO}_3$  added,  $z = 0$ .

$$\delta^{13}\text{C}_s = x\delta^{13}\text{C}_t + (1 - x)\delta^{13}\text{C}_a + 11.2\text{‰} \quad (3)$$

$$x = (\delta^{13}\text{C}_s - \delta^{13}\text{C}_a - 11.2)/(\delta^{13}\text{C}_t - \delta^{13}\text{C}_a) \cdot 100\% \quad (4)$$

$$x + y = 1 \quad (5)$$

From inputting the measured data, we obtain the  $x$  and  $y$  values shown in Table 1. For simplifying the calculation, to the snails fed  $\text{CaCO}_3$ , we assumed a similar  $x/y$  ratio to those of all the snails, although individual differences might happen among snails attributable to different growth rates. Therefore we calculated the average  $x/y$  ratio in non-carbonate groups, and obtained

$$x/y = 3.2 \quad (6)$$

at  $20^\circ\text{C}$ ,  $b = 11.2\text{‰}$ . Therefore,

$$y = (\delta^{13}\text{C}_s - \delta^{13}\text{C}_c - 11.2)/(3.2\delta^{13}\text{C}_t + \delta^{13}\text{C}_a - 4.2\delta^{13}\text{C}_c) \cdot 100\% \quad (7)$$

By combining Eqs. (2), (6), and (7), we calculated  $x$ ,  $y$ ,  $z$  values separately. Similar calculations were done at  $25^\circ\text{C}$  and  $30^\circ\text{C}$ . Finally, for the  $\text{C}_4$  plant groups, we assumed



the same  $x/y$  ratio to the  $C_3$  plant groups, which might not be accurate because the food consumption preference is different and  $C_3$  plant groups have a higher growth rate than  $C_4$  plant groups have. Then we calculated  $x$ ,  $y$ , and  $z$  using the same method. All calculated results are presented in Table 1.

The calculated  $x$ ,  $y$ ,  $z$  values for snails fed  $C_3$  plants reveal that the contributions of diet, atmospheric  $CO_2$ , and ingested limestone varied respectively as 66–80 %, 16–24 %, and 0–13 %. Furthermore, for those fed  $C_4$  plants, because of the potential food stress (lower consumption ability of  $C_4$  plant), they vary respectively as 56–64 %, 18–20 %, and 16–26 %. We observed a higher shell weight proportion of snails that had been fed corn compared with those that had been fed cabbage, suggesting a higher ingested limestone contribution, which is coincident with our calculations.

Figure 5a presents a positive correlation between the calculated contribution of ingested carbonate and the shell weight proportion. Calcium carbonate can be transformed into  $Ca^{2+}$  in the stomach of land snails and can then be ingested into the hemolymph, and can finally be precipitated into the shell carbonate. Therefore, high calcium carbonate consumption is always correlated with a high shell weight proportion. The  $\delta^{13}C$  of fed carbonate is more positive than  $\delta^{13}C$  of food and atmospheric  $CO_2$ . For that reason, if snails consume more carbonate, then their shell  $\delta^{13}C$  values are expected to be more positive, which is consistent to the relation presented in Fig. 5b.

As described earlier, the  $x/y$  ratio can vary among snail individuals. Therefore, the robustness of our estimation method was tested using the maximum and minimum  $x/y$  values observed under a no- $CaCO_3$  condition at different temperatures. The results are presented in Table 2. For snails fed with cabbage and  $CaCO_3$  at 20 °C, the  $x/y$  values vary as 2.9–3.3. The maximum discrepancies of estimations by the maximum and minimum  $x/y$  value could be 1.3 %, –2.9 %, and 1.6 % for  $x$ ,  $y$ , and  $z$  values, respectively. For those fed with cabbage and  $CaCO_3$  at 25 °C,  $x/y$  values vary as 3.4–5.2. The maximum discrepancies could be 3.0 %, –6.9 %, and 4.1 % for  $x$ ,  $y$ , and  $z$  values, respectively. Considering individual differences among snails (calculated stan-

# Factors controlling shell carbon isotopic composition of land snail

N. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



dard deviations of  $x$ ,  $y$ , and  $z$  to the snails fed with cabbage and  $\text{CaCO}_3$  are 3.9%, 1.6%, and 3.9%, respectively), these discrepancies are acceptable, especially for the estimated contribution of diet and ingested carbonate. Moreover, almost all of these re-estimated  $x$ ,  $y$ , and  $z$  values are within in the scopes we have calculated based on an average  $x/y$  value, showing a satisfactory robustness of this estimation method.

From the information presented above, we are confident that our calculation method and results are suitable and reasonable.

## 4.2 Contribution of each carbon source to shell $\delta^{13}\text{C}$

### 4.2.1 Ingested carbonate (limestone in nature)

Stott (2002) reported no apparent differences in shell  $\delta^{13}\text{C}$  of land snail *Helix aspersa* fed with and without  $\text{CaCO}_3$ . However, the deviation of  $^{14}\text{C}$  ages estimation for some snail species in nature shows an incorporation of limestone (Goodfriend and Stipp, 1983; Pigati et al., 2010). Consequently, ingested limestone is expected to play an important role in controlling shell carbonate  $\delta^{13}\text{C}$ , at least for some species. Heavier  $\delta^{13}\text{C}$  values were observed in our cultivated land snail *Acusta despecta sieboldiana* fed  $\text{CaCO}_3$  compared with those fed  $\text{Ca}_3(\text{PO}_4)_2$ , especially at 20°C and 25°C (Fig. 6a). These values are even heavier than the values predicted from a flux balance model (Balakrishnan and Yapp), which considers both metabolic  $\text{CO}_2$  and atmospheric  $\text{CO}_2$  (Fig. 3), strongly suggesting the involvement of ingested  $\text{CaCO}_3$  powder (Fig. 6b). The estimated contribution for  $\text{C}_3$  plant group is 0–13%; and for  $\text{C}_4$  plant, it varies: 16–26%. The different estimated proportions reveal different growth rates (metabolic rates) among snails. Apparently, snails fed  $\text{C}_4$  plants might prefer involving higher proportions of limestone, which is probably caused by diet stress.

Figure 7a (see Table 3) shows a decreasing trend of shell  $\delta^{13}\text{C}$  along with the snail growth direction, which is consistent to the measurements of living land snail individuals reported by Yanes et al. (2012), who observed higher  $\delta^{13}\text{C}$  values during the younger growth stages of land snails. This fact is explainable by the larger proportion of lime-

BGD

11, 6555–6590, 2014

## Factors controlling shell carbon isotopic composition of land snail

N. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Factors controlling shell carbon isotopic composition of land snail

N. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



stone ingested during the early period of land snails to enhance their growth rates. Egonmwan (2008) reported that the amount of food ingestion increases gradually, although calcium ingestion first increases (1–3 months, but increase rate is lower than diet consumption) and then decreases (4–6 months) during the first 6 months, suggesting that the contribution of ingested carbonate can be expected to decrease along with the snail growth. Our estimated contributions of ingested carbonate (Fig. 7b) support this supposition. To elucidate this phenomenon, similar studies should be done to observe snails fed with  $C_4$  plants and those fed without  $CaCO_3$ .

### 4.2.2 Diet and food selectivity

DeNiro and Epstein (1978) reported a slight enrichment (approximately 1‰) of the snail body tissue  $\delta^{13}C$  relative to their diet: romaine lettuce. Stott (2002) shown no significant isotopic offset between snail body tissue  $\delta^{13}C$  fed lettuce and their diet, whereas about 2–3‰ depletion was observed for snails fed corn (Fig. 8a), which is regarded as attributable to the contribution of preculture carbon because they cultured snails from juveniles but not eggs. Our observations show slight enrichment among all snails relative to their diet: approximately 0.5‰ for cabbage and approximately 1.2‰ for corn. However, this enrichment is negligible when considering measurement precision among both snail individuals and vegetable samples. Therefore, we infer that the  $\delta^{13}C$  values of snail body tissue are similar and that they should directly reflect the  $\delta^{13}C$  values of their food.

Figure 8b shows the observed relations between shell carbonate  $\delta^{13}C$  of snails and their diet (This study; Stott, 2002; Metref et al., 2003). Evidently, all of these slopes are less than the expected slope 1, which is considered diet as a single source controlling shell carbon (Francey, 1993; Stott, 2002). Offsets suggest the influence of other sources such as limestone and atmospheric  $CO_2$  causing these more positive values. The relative contributions of  $C_3$  and  $C_4$  type food might be different. Our estimation confirms this inference: the proportions of shell carbon from cabbage are 66–80 % and from corn are 56–64 %, which might reflect food stress among different plants. Metref

## Factors controlling shell carbon isotopic composition of land snail

N. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

et al. (2003) reported a  $C_3$  vs.  $C_4$  mixed experiment. Most snails show a preferential use of  $C_3$  food. This study also found a preference of this kind in consuming  $C_3$  but not  $C_4$  plants, i.e., snails were growing faster when eating  $C_3$  plants such as lettuce and cabbage than those fed a  $C_4$  plant such as corn, which agrees with the observations reported by Metref et al. (2003). Moreover, almost all the snails fed *Miscanthus sinensis* ( $C_4$  plant) died after 20–30 days, except one or two large individuals and their shell  $\delta^{13}C$  values ( $-11.1 \pm 3.4\%$ ) have no marked differences from newly hatched snail larvae ( $-11.0 \pm 0.1\%$ ), suggesting that the snails were unable to consume *Miscanthus sinensis* at all. However, Stott (2002) reported a lower growth rate of snails cultured with dried sour orange tree leaves ( $C_3$  plant) compared to those cultured by lettuce and corn, suggesting food quality such as water content or physical structure as the reason for food preference. This food selectivity of land snails increases the difficulty of their application in the paleo-environment reconstruction, especially for the accurate study of  $C_3/C_4$  vegetation distribution. Therefore, future studies of food quality's influence on snail shell  $\delta^{13}C$  values might be helpful.

### 4.2.3 Atmospheric $CO_2$

Although Stott (2002) and Metref et al. (2003) reported that atmospheric  $CO_2$  does not contribute to shell carbonate  $\delta^{13}C$  values based on culturing land snail *Helix aspersa*, some works clarify it as an important source of shell carbon. For instance, Balakrishnan and Yapp (2004) pointed out that  $pCO_2$  ratio between the snail and its environment affects shell incorporation of respired  $CO_2$ . McConnaughey and Gillikin (2008) summarized the previous studies of relation between land snail shell carbonate  $\delta^{13}C$  and shell organics (or body tissue organics/diet) and pointed out that the offsets are greater when the  $\delta^{13}C$  of organics diverge more from atmospheric  $CO_2$ , i.e., when the slope of the shell vs. organic regression line is less than unity, suggesting that the latter does contribute to shell carbonate.

In this study, our calculation results revealed that atmospheric CO<sub>2</sub> can affect shell carbonate isotopic values, and our estimated contributions are, respectively, 16–24 % (fed cabbage), 18–20 % (fed corn). More clear evidence might be that, for snails growing up without CaCO<sub>3</sub> (Fig. 3), shell  $\delta^{13}\text{C}$  values became heavier than the expected values controlled by one end member (diet).

However, the limited accuracy of our assumption of similar diet/atmospheric CO<sub>2</sub> ratio at the same temperature must be acknowledged because the snails have individual differences of metabolic rates during their growth (Barnhart and McMahon, 1987). Therefore, to learn more accurate contributions of atmospheric CO<sub>2</sub>, further incubation experiment is necessary, which is expected to include several parallel groups with labeled  $\delta^{14}\text{C}$  or  $\delta^{13}\text{C}$ -different CO<sub>2</sub> compositions.

### 4.3 Metabolic rate

Snail individuals have different metabolic rates in different environment conditions, or even in different growth phases of a single individual. Different metabolic rates reflect different partial pressures of CO<sub>2</sub> produced by diet, which consequently causes different flux ratios among CO<sub>2</sub> produced from metabolism, digestion of limestone, and atmosphere to approach a carbon isotope steady state in snail body fluids, and eventually produce different  $\delta^{13}\text{C}$  values. Many discussions we presented above show influences of this kind, such as (1) snails cultured at 30 °C groups with CaCO<sub>3</sub> show more depleted  $\delta^{13}\text{C}$  values than those at 20 °C and 25 °C groups (Fig. 6a). Correspondingly, they have a lower survival rate, smaller size and lighter mass, suggesting stress from poor growth conditions (e.g., unsuitable temperature, perishable leaves, lack of oxygen, etc.); (2) for one snail individual, the trend of decrease in both  $\delta^{13}\text{C}$  values of shell carbonate (Fig. 7a) and estimated that the contribution of ingested carbonate (Fig. 7b) reflects a decreasing metabolic rate along with snail growing up; (3) the different slopes presented in Fig. 8b reveal different metabolic rates among different snail species or different cultivation conditions.

**BGD**

11, 6555–6590, 2014

## Factors controlling shell carbon isotopic composition of land snail

N. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Factors controlling shell carbon isotopic composition of land snail

N. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Therefore, although we usually infer that a difference of  $\delta^{13}\text{C}$  values (e.g. approx. 3‰; Yanes et al., 2011) in snail shell fossils reflects a shift of  $\text{C}_3/\text{C}_4$  plant proportion among ages, it might be caused by variation in metabolic rates attributable to shifting environmental conditions, which might produce a discrepancy of shell carbonate  $\delta^{13}\text{C}$  values as large as the 3.5‰ found in this study.

Moreover, these results suggest a lack of a common model for all land snail species. We must calculate the relative contribution of each source to each snail individual. Therefore, the metabolic rates among individuals might prevent the reconstruction of the paleo-environment parameters (e.g. radiocarbon dating) precisely in quantitative analysis from shell carbonate  $\delta^{13}\text{C}$  values, but sometimes it might help us understand the environmental conditions better where and when a snail grew up, which should be addressed in a future study.

### 4.4 Temperature

Theoretically, temperature can affect the fractionation factor between gaseous  $\text{CO}_2$  and  $\text{HCO}_3^-$  in land snail body fluid from a relation presented as  $\Delta^{13}\text{C}_{\text{HCO}_3^-(\text{g})} = -(0.114 \pm 0.003)T(^{\circ}\text{C}) + (10.78 \pm 0.05)\text{‰}$  (Mook et al., 1974; Zhang et al., 1995; Szaran, 1997). Therefore, for shells precipitated at two different temperatures, the discrepancy between  $\delta^{13}\text{C}$  values ( $\Delta^{13}\text{C}$ , ‰) is  $-0.11/^{\circ}\text{C} \times T$  (Zhang et al., 1995; see also in Romanek et al., 1991,  $-0.13/^{\circ}\text{C}$  and Szaran, 1997,  $-0.10/^{\circ}\text{C}$ ). We observed a relation of  $-0.17 \pm 0.04/^{\circ}\text{C}$  among snails fed without  $\text{CaCO}_3$  (Fig. 9), which is not significantly different from the theoretically expected relation. The small sharper trend might somehow reflect metabolic differences among the snails at different temperature conditions. The  $\delta^{13}\text{C}$  discrepancy of 0.17‰ per degree is small compared with the contributions of different carbon sources. For that reason, no special consideration of environmental parameters of this kind is necessary on most occasions.

## 4.5 Carbon isotopic fractionations during snail laying and hatching

Metref et al. (2003) found that hatched and 1 day old snails showed depletion of 2.5‰ compared to their parents. They inferred that shells of the hatching juvenile were built not only from eggshell calcite but also from isotopic depleted albumen. We have observed these  $\delta^{13}\text{C}$  results for *Acusta despecta sieboldiana*: adult snails (parents),  $-11.5 \pm 1.2\text{‰}$  ( $n = 6$ ); hatched snails:  $-11.0 \pm 0.1\text{‰}$  ( $n > 10$ , mixed samples); eggs:  $-13.1 \pm 0.9\text{‰}$  (two groups of eggs, for each group,  $n > 10$ ). No obvious depletion was found between hatched snails and adults.

Romanek et al. (1992) pointed out that fractionation factors among calcite,  $\text{HCO}_3^-$  and aragonite are temperature independent when they are in equilibrium conditions. Their values were reported as  $\Delta^{13}\text{C}_{\text{Calcite-HCO}_3^-} = 0.9 \pm 0.2\text{‰}$ ,  $\Delta^{13}\text{C}_{\text{Aragonite-HCO}_3^-} = 2.7 \pm 0.2\text{‰}$ ,  $\Delta^{13}\text{C}_{\text{Aragonite-Calcite}} = 1.8 \pm 0.2\text{‰}$  (Rubinson and Clayton, 1969; see also in Romanek et al., 1992,  $\Delta^{13}\text{C}_{\text{Calcite-HCO}_3^-} = 1.0 \pm 0.2\text{‰}$ ;  $\Delta^{13}\text{C}_{\text{Aragonite-HCO}_3^-} = 2.7 \pm 0.6\text{‰}$ ;  $\Delta^{13}\text{C}_{\text{Aragonite-Calcite}} = 1.7 \pm 0.4\text{‰}$ ). According to the XRD results, all of our cultured snail shells are made of aragonite, whereas eggshells are made of calcite. We are striving to ascertain the isotopic fractionations based on calcite- $\text{HCO}_3^-$ -aragonite equilibrium. At this isotopic equilibrium condition, (a) for the laying process: when the snail precipitate shell (aragonite) from bicarbonate pool,  $\delta^{13}\text{C}$  will enrich 2.7‰; and when precipitate eggshell (calcite),  $\delta^{13}\text{C}$  will enrich 0.9–1.0‰. Therefore, the difference between shell aragonite and eggshell calcite is expected to be 1.7–1.8‰, we obtain a similar value of 1.6‰, suggesting that snail eggshells are also precipitated from the bicarbonate pool and that they follow the equilibrium fractionation. (b) For the hatching process, when the larvae are hatching, we assume that the shell calcite will dissolve gradually into the egg water to form a bicarbonate pool to precipitate egg aragonite. Then the fractionation between egg calcite and hatched snail shell aragonite is expected to be around  $-1.8\text{‰}$  to  $-1.7\text{‰}$ . Here we obtained a value of  $-2.1\text{‰}$ , which is also consistent with

BGD

11, 6555–6590, 2014

### Factors controlling shell carbon isotopic composition of land snail

N. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

the assumed value if we consider the measurement error, suggesting that the shell aragonite is transferred from eggshell calcite at an isotopic equilibrium condition.

## 5 Conclusions

From culturing the land snail *Acusta despecta sieboldiana* under a controlled environment, we confirmed that diet, atmospheric CO<sub>2</sub>, and ingested limestone are important sources controlling shell carbon isotopic composition. We have also discussed the influences of metabolic rates. Furthermore, temperature can affect shell carbonate  $\delta^{13}\text{C}$  values by controlling the carbon isotopic fractionation of gaseous CO<sub>2</sub>–bicarbonate equilibrium. We presented a simple but credible framework to discuss the mechanisms of how each source and environmental parameter might affect shell carbonate  $\delta^{13}\text{C}$  values based on this and earlier studies. According to this framework and some reasonable assumptions, we estimated the contribution of different carbon sources for each snail individual: to cabbage (C<sub>3</sub> plant) fed snails, the contributions of diet, atmospheric CO<sub>2</sub> and ingested limestone vary respectively as 66–80 %, 16–24 %, and 0–13 %. Furthermore, to corn (C<sub>4</sub> plant) fed snails, because of the potential food stress (lower consumption ability of C<sub>4</sub> plant), they vary respectively in 56–64 %, 18–20 %, and 16–26 %.

Secondly, we found that snails discriminate in their choices of C<sub>3</sub>/C<sub>4</sub> plants as food. For instance, they can grow faster when eating C<sub>3</sub> plants such as cabbage compared with C<sub>4</sub> plant such as corn than they can when they eat C<sub>4</sub> plants such as *Miscanthus sinensis*. This kind of food selectivity of land snails increases the difficulty of their application in the paleo-environment reconstruction, especially for the accurate study of C<sub>3</sub>/C<sub>4</sub> vegetation distribution. Finally, we inferred that during egg laying and hatching of our cultured snails, carbon isotope fractionation is controlled only by the isotopic exchange of calcite–HCO<sub>3</sub><sup>−</sup>–aragonite equilibrium.

To prompt the application of carbon isotope in paleo-environment reconstruction, additional work should be undertaken in future culture experiments, especially in the

BGD

11, 6555–6590, 2014

### Factors controlling shell carbon isotopic composition of land snail

N. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Factors controlling shell carbon isotopic composition of land snail

N. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

following several aspects: (1) intra-shell measurements of snails fed with  $C_4$  plants and those fed without  $CaCO_3$  should be taken in future studies to verify the phenomenon of decreasing trend of shell  $\delta^{13}C$  along with the snail growth direction; (2) influences of food quality such as water contents or physical structure on snail shell  $\delta^{13}C$  values should be investigated; (3) snails can be cultured in air with labeled  $\Delta^{14}C$  or  $\delta^{13}C$ -different  $CO_2$  composition to ascertain the contribution of atmospheric  $CO_2$  more accurately; (4) snails fed with  $C_4$  plants under a no-carbonate condition should be evaluated; (5) more land snail species preferring varied environmental conditions should be studied to ascertain which inferences are common to all species and which are suitable for a given species.

**Acknowledgements.** This study was supported by a Grant-in-Aid for Scientific Research (#23224013) and a grant for the Global COE Program “From the Earth to Earths” to Tokyo Institute of Technology. We thank Kazunori Hasegawa for helping us ascertain the species of cultured land snails. We thank Yingying Cheng for helping us run XRD measurements. Furthermore, our appreciation is extended to Xueqin Chi for her kind help during our cultivation of land snails. Finally, we are very grateful to Alexis Gilbert for his helpful comments.

## References

- Balakrishnan, M. and Yapp, C. J.: Flux balance models for the oxygen and carbon isotope compositions of land snail shells, *Geochim. Cosmochim. Ac.*, 68, 2007–2024, 2004.
- Balslev-Clausen, D., Dahl, T. W., Saad, N., and Rosing, M. T.: Precise and accurate  $\delta^{13}C$  analysis of rock samples using Flash Combustion–Cavity Ring Down Laser Spectroscopy, *J. Anal. Atom. Spectrom.*, 28, 516–523, 2013.
- Barnhart, M. C. and McMahon, B. R.: Discontinuous carbon dioxide release and metabolic depression in dormant land snails, *J. Exp. Biol.*, 128, 123–138, 1987.
- Colonese, A. C., Zanchetta, G., Fallick, A. E., Martini, F., Manganelli, G., and Lo-Vetro, D.: Stable isotope composition of Late Glacial land snail shells from Grotta del Romito (Southern Italy): palaeoclimatic implications, *Palaeogeogr. Palaeoclimatol.*, 254, 550–560, 2007.



# Factors controlling shell carbon isotopic composition of land snail

N. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Colonese, A. C., Zanchetta, G., Fallick, A. E., Martini, F., Manganelli, G., and Drysdale, R. N.: Stable isotope composition of *Helix ligata* (Müller, 1774) from Late Pleistocene–Holocene archaeological record from Grotta della Serratura (Southern Italy): palaeoclimatic implications, *Global Planet. Change*, 71, 249–257, 2010.
- 5 Colonese, A. C., Zanchetta, G., Drysdale, R. N., Fallick, A. E., Manganelli, G., Lo-Vetro, D., Martini, F., and Di Giuseppe, Z.: Stable isotope composition of Late Pleistocene–Holocene *Eobania vermiculata* (Müller, 1774) (Pulmonata, Stylommatophora) shells from the Central Mediterranean Basin: data from Grotta d'Oriente (Favignana, Sicily), *Quatern. Int.*, 244, 76–87, 2011.
- 10 Coplen, T. B., Brand, W. A., Gehre, M., Gröning, M., Meijer, H. A., Toman, B., and Verhoeven, R. M.: New guidelines for  $\delta^{13}\text{C}$  measurements, *Anal. Chem.*, 78, 2439–2441, 2006.
- DeNiro, M. J. and Epstein, S.: Influence of diet on the distribution of carbon isotopes in animals, *Geochim. Cosmochim. Ac.*, 42, 495–506, 1978.
- Egonmwan, R. I.: Effects of dietary calcium on growth and oviposition of the African land snail *Limicolaria flammea* (Pulmonata: Achatinidae), *Rev. Biol. Trop.*, 56, 333–343, 2008.
- 15 Francey, R. J.: A comment on  $^{13}\text{C}/^{12}\text{C}$  in land snail shells, *Earth Planet. Sc. Lett.*, 63, 142–143, 1983.
- Goodfriend, G. A.: The use of land snail shells in paleoenvironmental reconstruction, *Quaternary Sci. Rev.*, 11, 665–685, 1992.
- 20 Goodfriend, G. A. and Hood, D. G.: Carbon isotope analysis of land snail shells; implications for carbon sources and radiocarbon dating, *Radiocarbon*, 25, 810–830, 1983.
- Goodfriend, G. A. and Magaritz, M.: Carbon and oxygen isotope composition of shell carbonate of desert land snails, *Earth Planet. Sc. Lett.*, 86, 377–388, 1987.
- Goodfriend, G. A. and Stipp, J. J.: Limestone and the problem of radiocarbon dating of land-snail shell carbonate, *Geology*, 11, 575–577, 1983.
- 25 Goodfriend, G. A., Magaritz, M., and Gat, J. R.: Stable isotope composition of land snail body water and its relation to environmental waters and shell carbonate, *Geochim. Cosmochim. Ac.*, 53, 3215–3221, 1989.
- Keeling, R. F., Piper, S. C., Bollenbacher, A. F., and Walker, S. J.: Monthly atmospheric  $^{13}\text{C}/^{12}\text{C}$  isotopic ratios for 11 SIO stations, in: *Trends: a Compendium of Data on Global Change*, Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, US Department of Energy, Oak Ridge, Tenn., USA, 2010.
- 30

- Kehrwald, N. M., McCoy, W. D., Thibeault, J., Burns, S. J., and Oches, E. A.: Paleoclimatic implications of the spatial patterns of modern and LGM European land-snail shell  $\delta^{18}\text{O}$ , *Quaternary Res.*, 74, 166–176, 2010.
- Lécolle, P.: The oxygen isotope composition of landsnail shells as a climatic indicator: applications to hydrogeology and paleoclimatology, *Chem. Geol.*, 58, 157–181, 1985.
- McConnaughey, T. A. and Gillikin, D. P.: Carbon isotopes in mollusk shell carbonates, *Geo-Mar. Lett.*, 28, 287–299, 2008.
- Metref, S., Rousseau, D. D., Bentaleb, I., Labonne, M., and Vianey-Liaud, M.: Study of the diet effect on  $\delta^{13}\text{C}$  of shell carbonate of the land snail *Helix aspersa* in experimental conditions, *Earth Planet. Sc. Lett.*, 211, 381–393, 2003.
- Mook, W. G., Bommerson, J. C., and Staverman, W. H.: Carbon isotope fractionation between dissolved bicarbonate and gaseous carbon dioxide, *Earth Planet. Sc. Lett.*, 22, 169–176, 1974.
- Pigati, J. S., Quade, J., Shahanan, T. M., and Haynes Jr., C. V.: Radiocarbon dating of minute gastropods and new constraints on the timing of late Quaternary spring–discharge deposits in southern Arizona, USA, *Palaeogeogr. Palaeoclimatol.*, 204, 33–45, 2004.
- Pigati, J. S., Rech, J. A., and Nekola, J. C.: Radiocarbon dating of small terrestrial gastropod shells in North America, *Quat. Geochronol.*, 5, 519–532, 2010.
- Romanek, C. S., Grossman, E. L., and Morse, J. W.: Carbon isotopic fractionation in synthetic aragonite and calcite: effects of temperature and precipitation rate, *Geochim. Cosmochim. Ac.*, 56, 419–430, 1992.
- Romaniello, L., Quarta, G., Mastronuzzi, G., D’Elia, M., and Calcagnile, L.:  $^{14}\text{C}$  age anomalies in modern land snails shell carbonate from Southern Italy, *Quat. Geochronol.*, 3, 68–75, 2008.
- Rubinson, M. and Clayton, R. N.: Carbon-13 fractionation between aragonite and calcite, *Geochim. Cosmochim. Ac.*, 33, 997–1002, 1969.
- Stott, L. D.: The influence of diet on the  $\delta^{13}\text{C}$  of shell carbon in the pulmonate snail *Helix aspersa*, *Earth Planet. Sc. Lett.*, 195, 249–259, 2002.
- Szaran, J.: Carbon isotope fractionation between dissolved and gaseous carbon dioxide, *Chem. Geol.*, 150, 331–337, 1998.
- Wahl, E. H., Fidric, B., Rella, C. W., Koulikov, S., Kharlamov, B., Tan, S., Kachanov, A. A., Richman, B. A., Crosson, E. R., Paldus, B. A., Kalaskar, S., and Bowling, D. R.: Applications

**BGD**

11, 6555–6590, 2014

## Factors controlling shell carbon isotopic composition of land snail

N. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

of cavity ring-down spectroscopy to high precision isotope ratio measurement of  $^{13}\text{C}/^{12}\text{C}$  in carbon dioxide, *Isot. Environ. Health. S.*, 42, 21–35, 2006.

Yanes, Y., Delgado, A., Castillo, C., Alonso, M. R., Ibáñez, M., De-la-Nuez, J., and Kowalewski, M.: Stable isotope ( $\delta^{18}\text{O}$ ,  $\delta^{13}\text{C}$ , and  $\delta\text{D}$ ) signatures of recent terrestrial communities from a low-latitude, oceanic setting: endemic land snails, plants, rain, and carbonate sediments from the eastern Canary Islands, *Chem. Geol.*, 249, 377–392, 2008a.

Yanes, Y., Tomašových, A., Kowalewski, M., Castillo, C., Aguirre, J., Alonso, M. R., and Ibáñez, M.: Taphonomy and compositional fidelity of Quaternary fossil assemblages of terrestrial gastropods from carbonate-rich environments of the Canary Islands, *Lethaia*, 41, 235–256, 2008b.

Yanes, Y., Romanek, C. S., Delgado, A., Brant, H. A., Noakes, J. E., Alonso, M. R., and Ibáñez, M.: Oxygen and carbon stable isotopes of modern land snail shells as environmental indicators from a low-latitude oceanic island, *Geochim. Cosmochim. Ac.*, 73, 4077–4099, 2009.

Yanes, Y., Yapp, C. J., Ibáñez, M., Alonso, M. R., De-la-Nuez, J., Quesada, M. L., Castillo, C., and Delgado, A.: Pleistocene–Holocene environmental change in the Canary Archipelago as inferred from the stable isotope composition of land snail shells, *Quaternary Res.*, 75, 658–669, 2011.

Yanes, Y., Gutiérrez-Zugasti, I., and Delgado, A.: Late-glacial to Holocene transition in northern Spain deduced from land-snail shelly accumulations, *Quaternary Res.*, 78, 373–385, 2012a.

Yanes, Y., Riquelme, J. A., Cámara, J. A., and Delgado, A.: Stable isotope composition of middle to late Holocene land snail shells from the Marroquíes archeological site (Jaén, Southern Spain): paleoenvironmental implications, *Quatern. Int.*, 302, 77–87, 2012b.

Yanes, Y., García-Alix, A., Asta, M. P., Ibáñez, M., Alonso, M. R., and Delgado, A.: Late Pleistocene–Holocene environmental conditions in Lanzarote (Canary Islands) inferred from calcitic and aragonitic land snail shells and bird bones, *Palaeogeogr. Palaeoclimatol.*, 378, 91–102, 2013.

Yapp, C. J.: Oxygen and carbon isotope measurements of land snail shell carbonate, *Geochim. Cosmochim. Ac.*, 43, 629–635, 1979.

Zaarur, S., Olack, G., and Affek, H. P.: Paleo-environmental implication of clumped isotopes in land snail shells, *Geochim. Cosmochim. Ac.*, 75, 6859–6869, 2011.

Zhang, J., Quay, P. D., and Wilbur, D. O.: Carbon isotope fractionation during gas–water exchange and dissolution of  $\text{CO}_2$ , *Geochim. Cosmochim. Ac.*, 59, 107–114, 1995.

**BGD**

11, 6555–6590, 2014

## Factors controlling shell carbon isotopic composition of land snail

N. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Zhang, N. Z., Yamada, K., and Yoshida, N.: Variations of  $^{13}\text{C}^{16}\text{O}^{18}\text{O}$  and its controlling factors in the atmosphere of Yokohama, Japan based on the observation during 2007–2011, in preparation, 2014.

**BGD**

11, 6555–6590, 2014

**Factors controlling  
shell carbon isotopic  
composition of land  
snail**

N. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Table 1.** Stable isotope results of snails cultured under different conditions and the estimated contributions of their shell carbon sources.

Snail No.	Temp. (°C)	CaCO <sub>3</sub> <sup>a</sup>	Diet	Diet $\delta^{13}\text{C}$ (‰)	Shell $\delta^{13}\text{C}$ (‰)	Tissue $\delta^{13}\text{C}$ (‰)	x <sup>e</sup> (%)	y <sup>e</sup> (%)	z <sup>e</sup> (%)	Shell weight proportion (%)
S1	20	+	Cabbage	-28.4	-9.6	-28.4	67.7	21.2	11.1	30.1
S2	20	+	Cabbage	-28.4	-9.1	-28.6	66.0	20.6	13.3	28.3
S9	20	+	Cabbage	-28.4	-9.1	-28.5	66.2	20.7	13.1	29.7
S10	20	+	Cabbage	-28.4	-9.7	-28.8	67.3	21.0	11.7	27.0
S11	20	+	Cabbage	-28.4	-9.7	-27.6	69.5	21.7	8.7	32.7
S12	20	+	Cabbage	-28.4	-9.5	-27.9	68.4	21.4	10.3	25.4
S5	20	+	Cabbage	-28.4	-9.5	-27.7	68.7	21.5	9.8	25.9
S6	20	+	Cabbage	-28.4	-9.8	-27.4	70.2	21.9	7.9	26.4
S13	20	+	Cabbage	-28.4	-9.4	-28.1	67.9	21.2	10.9	28.2
S14	20	+	Cabbage	-28.4	-10.2	-27.7	70.7	22.1	7.2	23.6
S15	20	+	Cabbage	-28.4	-11.0	-26.9	74.5	23.3	2.2	24.8
S3	20	-	Cabbage	-28.4	-13.1	-29.4	74.3	25.7	0.0	4.6
S7	20	-	Cabbage	-28.4	-12.3	-27.7	76.9	23.1	0.0	10.1
S8	20	-	Cabbage	-28.4	-12.3	-27.8	76.6	23.4	0.0	8.8
S16	20	+	Corn	-12.0	4.2	-10.3	59.1	18.5	22.4	34.4
S17	20	+	Corn	-12.0	4.0	-11.6	56.3	17.6	26.1	34.9
S18	20	+	Corn	-12.0	3.3	-10.5	63.7	19.9	16.4	-
S19	25	+	Cabbage	-28.4	-9.7	-27.5	69.6	17.4	12.9	33.1
S21	25	+	Cabbage	-28.4	-11.0	-27.7	72.8	18.2	9.0	31.0
S25	25	+	Cabbage	-28.4	-12.1	-28.7	74.1	18.5	7.4	19.9
S26	25	+	Cabbage	-28.4	-10.1	-27.7	70.5	17.6	11.9	34.5
S29	25	+	Cabbage	-28.4	-10.2	-27.3	71.6	17.9	10.5	37.8
S31	25	+	Cabbage	-28.4	-9.8	-27.0	71.0	17.8	11.2	30.2
S33 <sup>b</sup>	25	+	Cabbage	-28.4	-9.9	-26.5	72.3	18.1	9.6	9.7
S22	25	-	Cabbage	-28.4	-14.9	-28.5	83.9	16.1	0.0	4.3
S23	25	-	Cabbage	-28.4	-13.7	-27.5	82.1	17.9	0.0	5.9
S27	25	-	Cabbage	-28.4	-13.6	-28.5	77.3	22.7	0.0	5.1
S28	25	-	Cabbage	-28.4	-13.4	-27.9	78.8	21.2	0.0	4.2
S32	25	-	Cabbage	-28.4	-13.0	-27.7	77.8	22.2	0.0	8.7
S34	30	+	Cabbage	-28.4	-13.7	-28.0	78.5	20.1	1.4	13.8
S35	30	+	Cabbage	-28.4	-13.5	-27.2	79.6	20.4	0.0	18.2
S36	30	+	Cabbage	-28.4	-13.1	-28.7	75.2	19.3	5.6	18.3
S37	30	+	Cabbage	-28.4	-13.3	-28.3	76.5	19.6	3.9	16.2
S38	30	+	Cabbage	-28.4	-13.4	-28.5	76.4	19.6	3.9	14.5
S39	30	+	Cabbage	-28.4	-11.4	-27.3	73.3	18.8	7.9	31.0
S40	30	+	Cabbage	-28.4	-13.8	-29.3	75.9	19.5	4.6	14.4
S41	30	+	Cabbage	-28.4	-13.5	-28.0	77.9	20.0	2.2	17.1
S42 <sup>c</sup>	30	-	Cabbage	-28.4	-14.2	-28.2 <sup>d</sup>	79.5	20.5	0.0	-

<sup>a</sup> "-" means those fed Ca<sub>3</sub>(PO<sub>4</sub>)<sub>2</sub>, which is thought to be hard and rarely ingested by snails;

<sup>b</sup> snails regarded as a control group, cultured under an air-free-exchange system with a small amount of soil,

<sup>c</sup> mixture of four individuals cultured in the same condition (sizes are similar);

<sup>d</sup> average value of snails from S34 to S41.

<sup>e</sup> x, calculated proportion of metabolic CO<sub>2</sub>; y, proportion of atmospheric CO<sub>2</sub>; z, proportion of CO<sub>2</sub> produced from ingested carbonate.

BGD

11, 6555–6590, 2014

## Factors controlling shell carbon isotopic composition of land snail

N. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Table 2.** Estimations of  $x$ ,  $y$ , and  $z$  under observed maximum and minimum  $x/y$  values at 20 °C and 25 °C.

Snail No.	$x_{\min}$ (%)	$y_{\min}$ (%)	$z_{\min}$ (%)	$x_{\max}$ (%)	$y_{\max}$ (%)	$z_{\max}$ (%)	$\Delta x$ (%)	$\Delta y$ (%)	$\Delta z$ (%)
20 °C, Cabbage	min: $x/y = 2.9$			max: $x/y = 3.3$					
S1	66.9	23.1	10.0	68.0	20.4	11.5	1.1	-2.6	1.5
S2	65.3	22.5	12.2	66.3	19.9	13.8	1.1	-2.6	1.5
S9	65.4	22.5	12.1	66.5	20.0	13.6	1.1	-2.6	1.5
S10	66.5	22.9	10.6	67.6	20.3	12.1	1.1	-2.6	1.5
S11	68.7	23.7	7.6	69.9	21.0	9.2	1.2	-2.7	1.6
S12	67.5	23.3	9.2	68.7	20.6	10.7	1.1	-2.7	1.5
S5	67.9	23.4	8.7	69.0	20.7	10.2	1.1	-2.7	1.5
S6	69.3	23.9	6.7	70.5	21.2	8.3	1.2	-2.7	1.6
S13	67.0	23.1	9.8	68.2	20.5	11.4	1.1	-2.7	1.5
S14	69.8	24.1	6.1	71.0	21.3	7.7	1.2	-2.8	1.6
S15	73.6	25.4	1.0	74.9	22.5	2.6	1.3	-2.9	1.6
20 °C, Corn	min: $x/y = 2.9$			max: $x/y = 3.3$					
S16	57.7	19.9	22.3	59.6	17.9	22.5	1.9	-2.0	0.1
S17	55.1	19.0	25.9	56.8	17.0	26.2	1.7	-1.9	0.3
S18	62.2	21.5	16.3	64.2	19.3	16.5	2.0	-2.2	0.1
25 °C, Cabbage	min: $x/y = 3.4$			max: $x/y = 5.2$					
S19	68.5	20.1	11.4	71.3	13.7	15.1	2.8	-6.5	3.7
S21	71.6	21.1	7.4	74.5	14.3	11.3	2.9	-6.8	3.9
S25	72.9	21.4	5.7	75.7	14.5	9.8	2.9	-6.9	4.1
S26	69.3	20.4	10.3	72.1	13.8	14.1	2.8	-6.6	3.8
S29	70.4	20.7	8.9	73.2	14.0	12.7	2.9	-6.7	3.8
S31	69.8	20.5	9.7	72.7	13.9	13.4	2.9	-6.6	3.7
S33	71.1	20.9	8.0	74.0	14.2	11.8	3.0	-6.7	3.7

" $x_{\min}$ ", " $y_{\min}$ ", and " $z_{\min}$ " are calculated by the minimum  $x/y$  values and " $x_{\max}$ ", " $y_{\max}$ ", and " $z_{\max}$ " are calculated by the maximum  $x/y$  values; " $\Delta x$ ", " $\Delta y$ ", and " $\Delta z$ " show the discrepancies of estimations by maximum and minimum  $x/y$  values for each snail individual.

**BGD**

11, 6555–6590, 2014

## Factors controlling shell carbon isotopic composition of land snail

N. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Table 3.** Stable isotope results of snail individual fractions and the estimated contributions of their shell carbon sources.

Sample	Spirals	Shell $\delta^{13}\text{C}$ (‰)	x (%)	y (%)	z (%)
S13_1	1–3	–8.2	64.6	20.2	15.2
S13_2	3–4	–9.2	67.2	21.0	11.9
S13_3	4–4.5	–9.3	67.5	21.1	11.4
S13_4	4.5–5	–10.0	69.5	21.7	8.8
S19_1	1–3	–8.1	65.1	16.3	18.6
S19_2	3–4	–9.0	67.8	16.9	15.3
S19_3	4–4.5	–9.1	67.9	17.0	15.1
S19_4	4.5–5	–9.2	68.1	17.0	14.9
S19_5	5–5.5	–12.6	77.9	19.5	2.6
S25_1	1–3	–9.2	66.0	16.5	17.5
S25_2	3–4	–10.8	70.6	17.6	11.8
S25_3	4–4.5	–12.0	73.9	18.5	7.6
S25_4	4.5–5	–12.4	75.0	18.8	6.2
S25_5	5–5.5	–13.1	76.9	19.2	3.9
S31_1	1–3	–8.1	66.1	16.5	17.4
S31_2	3–4	–9.2	69.4	17.3	13.3
S31_3	4–4.5	–8.8	68.3	17.1	14.7
S31_4	4.5–5	–10.8	74.0	18.5	7.5
S39_1	1–3	–9.1	66.7	17.1	16.2
S39_2	3–4	–10.4	70.5	18.1	11.4
S39_3	4–4.5	–11.4	73.5	18.9	7.6
S39_4	4.5–5	–12.8	77.3	19.8	2.8

Relative culturing conditions are shown in Table 1. Meanings of symbols “x”, “y”, and “z” are the same as those shown in Table 1.

**BGD**

11, 6555–6590, 2014

## Factors controlling shell carbon isotopic composition of land snail

N. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶


Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



**Fig. 1.** Growth phases of *Acusta despecta sieboldiana*: **(A)** adult snails collected from Suzukakedai, Yokohama, Japan (length: ca. 10–17 mm); **(B)** eggs (ca. 2 mm); **(C)** larva (1–3 days, ca. 2 mm); **(D)** juvenile (3–4 months, ca. 5–8 mm); **(E)** juvenile (> 6 months, > 10 mm); **(F)** shells for isotopic measurement.





**Fig. 2.** Snail survival rate: **(A)** at different temperatures; **(B)** with different calcium sources.

**BGD**

11, 6555–6590, 2014

## Factors controlling shell carbon isotopic composition of land snail

N. Zhang et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)


[Printer-friendly Version](#)

[Interactive Discussion](#)



# Factors controlling shell carbon isotopic composition of land snail

N. Zhang et al.



**Fig. 3.** Measured  $\delta^{13}\text{C}$  of snail shell aragonite against  $\delta^{13}\text{C}$  of associated organic matter (this study; Stott, 2002; Metref et al., 2003; Balakrishnan and Yapp, 2004).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

# Factors controlling shell carbon isotopic composition of land snail

N. Zhang et al.



**Fig. 4.** Simple framework of shell carbon sources and their relative isotope fractionation processes based on this and earlier studies.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶


Back

Close

Full Screen / Esc

Printer-friendly Version


Interactive Discussion



**Fig. 5.** Relation between **(A)** estimated contribution of ingested carbonate and shell weight percentage and **(B)**  $\delta^{13}\text{C}$  of snail shell aragonite and shell weight percentage.

# Factors controlling shell carbon isotopic composition of land snail

N. Zhang et al.




**Fig. 6.** Bulk  $\delta^{13}\text{C}$  of snail shell aragonite and estimated contribution of ingested carbonate at different culturing conditions.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


# Factors controlling shell carbon isotopic composition of land snail

N. Zhang et al.




**Fig. 7.**  $\delta^{13}\text{C}$  of snail shell sections and their estimated contribution of ingested carbonate at different growth phases.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)



**Fig. 8. (A)**  $\delta^{13}\text{C}$  of snail body tissue against their diet. **(B)**  $\delta^{13}\text{C}$  of snail shell aragonite against their diet.



**Fig. 9.** Relation between  $\delta^{13}\text{C}$  of snail shell aragonite (fed without carbonate) and different culture temperatures.