

1 **Increased winter soil temperature variability enhances nitrogen cycling and soil biotic**
2 **activity in temperate heathland and grassland mesocosms**

3

4 J. Schuerings^{1,2}, A. Jentsch¹, V. Hammer³, K. Lenz², H. A. L. Henry⁴, A. V. Malyshev² and J.
5 Kreyling²

6

7 ¹ Disturbance Ecology, BayCEER, University of Bayreuth, D-95440 Bayreuth, Germany

8 ² Biogeography, BayCEER, University of Bayreuth, D-95440 Bayreuth, Germany

9 ³ Research Unit Environmental Genomics, Helmholtz Zentrum München, D-85764
10 Neuherberg, Germany

11 ⁴ Biology, University of Western Ontario, London, Ontario, Canada

12

13 Corresponding author: Jan Schuerings, email: Jan.Schuerings@uni-bayreuth.de, phone:
14 +49(0)921-552345, fax: +49(0)921-2315

15 **ABSTRACT**

16 Winter air temperatures are projected to increase in the temperate zone, whereas snow
17 cover is projected to decrease, leading to increased soil temperature variability, and
18 potentially to changes in nutrient cycling. Here, we experimentally evaluated the effects of
19 increased winter soil temperature variability on selected aspects of the N-cycle in
20 mesocosms containing different plant community compositions. The experiment was
21 replicated at two sites, a colder mountainous upland site with high snow accumulation and a
22 warmer and dryer lowland site.

23 Increased soil temperature variability enhanced soil biotic activity for both sites during winter,
24 as indicated by 35% higher nitrogen (N) availability in the soil solution, 40% higher
25 belowground decomposition and a 25% increase in the potential activity of the enzyme
26 cellobiohydrolase. The mobilization of N differed between sites, and the ¹⁵N signal in leaves
27 was reduced by 31% in response to winter warming pulses, but only at the cold site, with
28 significant reductions occurring for three of four tested plant species at this site. Furthermore,
29 there was a trend of increased N leaching in response to the recurrent winter warming
30 pulses.

31 Overall, projected winter climate change in the temperate zone, with less snow and more
32 variable soil temperatures, appears important for shifts in ecosystem functioning (i.e. nutrient
33 cycling). While the effects of warming pulses on plant N mobilization did not differ among
34 sites, reduced plant ¹⁵N incorporation at the colder temperate site suggests that frost damage
35 may reduce plant N uptake in a warmer world, with important implications for nitrogen cycling
36 and nitrogen losses from ecosystems.

37 1 Introduction

38 Winter soil temperature is an important driver for many ecological and biogeochemical
39 processes in the cold-temperate and boreal zone, and it can influence the activity of plants
40 and soil biota (Matzner and Borken, 2008; Kreyling, 2010). While microbial activity and
41 nitrogen (N) cycling continue below freezing (Clein and Schimel, 1995; Mikan et al., 2002),
42 higher mean soil temperatures are generally expected to cause exponentially higher soil
43 biotic activity (Rustad et al., 2001; Melillo et al., 2002). Consequently, winter warming can
44 result in increased N mineralization and N availability in the soil solution in the following
45 growing season (Turner and Henry, 2010). Warmer soils over winter increase soil biotic
46 activity, e.g. soil respiration, decomposition by soil fauna and microbes, higher enzymatic
47 activity, higher N mineralization, etc. This holds true especially towards the end of winter, and
48 can accelerate plant productivity (Schuerings et al., 2013). Since plants are capable of winter
49 N uptake (Grogan et al., 2004; Andresen and Michelsen, 2005), their activity could
50 counteract N leaching (Patil et al., 2010). The general effectiveness of plants in taking up N
51 over winter, however, is not fully clear until now. Comparable N uptake rates over winter and
52 summer have been reported for some species (Nasholm et al., 2000; Bardgett et al., 2003),
53 but there is also evidence that cold acclimation reduces the potential for N uptake (Malyshev
54 and Henry, 2012a).

55 Due to increased winter air temperatures, snow cover will decrease in many regions of the
56 temperate zone (Christensen et al., 2007; Kreyling and Henry, 2011). However, air frost
57 events will still occur with unchanged magnitude and duration as nowadays in many
58 temperate regions (Kodra et al., 2011), and with less insulating snow cover, winter soil
59 temperatures can become more variable, particularly in upland and cold temperate regions
60 (Henry, 2008; Brown and DeGaetano, 2011). The resulting more variable soil temperature
61 conditions with frequent soil frost and freeze-thaw cycles (FTC) can affect N cycling. Soil
62 frost and FTC can physically damage plant roots (Tierney et al., 2001) and therefore reduce
63 the plants ability to take up N (Campbell et al., 2014), break up soil aggregates (Oztas and
64 Fayetorbay, 2003), and lyse microbial cells what enlarges the easily available N pool
65 (Skogland et al., 1988), thereby affecting N cycling and leading to N losses in dissolved
66 (Boutin and Robitaille, 1995; Brooks et al., 1998; Joseph and Henry, 2008) or gaseous forms
67 (Matzner and Borken, 2008). For warmer, lowland temperate regions, however, although soil
68 temperature variability might still increase (Kreyling, 2010), an increase in winter air
69 temperatures could lead to fewer soil FTC due to less frost (e.g. lowland Germany, Kreyling
70 and Henry, 2011). Contrasting effects of winter climate change can therefore be expected for
71 colder (stronger effects due to greater increase in soil temperature variability) versus warmer
72 (naturally higher soil temperature variability) temperate regions, and studies of
73 biogeochemical responses to increased soil temperature variability should be designed to
74 account for these differences.

75 Finally, plant species and vegetation types are known to influence N cycling (Hooper and
76 Vitousek, 1998; Knops et al., 2002). Different plant species and communities further show
77 different reactions to increased winter temperature variability in the temperate zone, with
78 grasses appearing more responsive than dwarf shrubs (Kreyling et al., 2010; Schuerings et
79 al., 2014) regarding their productivity, probably due to their faster life-cycle. However, this
80 increased responsiveness in productivity of grasses can either be beneficial (Kreyling et al.,
81 2008), or detrimental (Schuerings et al., 2014), probably depending on whether the minimum

82 temperatures experienced after warm phases induce frost damage. Altered plant productivity
83 can therefore indirectly affect N cycling. Generally, stress resistance is linked to nitrogen or
84 nutrient stress tolerance (Macgillivray et al., 1995). Moreover, increased N availability over
85 winter can increase the risk of frost damage to plants (Malyshev and Henry, 2012b).

86 In this experiment we tested the effects of more variable winter temperature conditions, i.e.
87 recurrent, short winter warming pulses, on soil biotic and potential extracellular enzyme
88 activity, N availability in the soil solution, and N uptake by plants in different plant
89 communities (grassland, heathland; same communities as in Schuerings et al., 2014) at two
90 sites with contrasting winter climate (a warm, snow-poor lowland and a cold, snow-rich
91 upland site). We hypothesised that (1) recurrent winter warming pulses would enhance N-
92 cycling (i.e. increased N availability, soil biotic activity and N uptake into plants). (2) We
93 further expected different responsiveness to the recurrent warming pulses at the two sites,
94 with more variable soil temperatures and stronger frost, therefore frost damage negatively
95 affecting plant N uptake at the colder upland site. (3) Finally, we expected differences among
96 the plant communities in the response of N cycling to the recurrent warming pulses, with a
97 higher ability for winter N uptake in grassland than in heathland plants.

98

99 **2 METHODS**

100 2.1 Experimental design and site description

101 This research is part of the EVENT IV experiment, testing the effects of increased winter
102 temperature variability on temperate heath and grassland communities. The effects of the
103 recurrent warming pulses on plant growth (above- and below-ground) are summarized in
104 Schuerings et al. (2014), whereas here we concentrate on nitrogen cycling. The experiment
105 was replicated at two sites: the warm site was located in the Ecological-Botanical Garden of
106 the University of Bayreuth (49° 55' 36.32" N, 11° 34' 57.28" E, 358 m a.s.l.) and the cold site
107 was located at the Waldstein mountain in the Fichtelgebirge (50° 8' 35.81" N, 11° 51' 50.92"
108 E, 781 m a.s.l.). The cold site generally experiences more precipitation and harsher winter
109 conditions (Table 1).

110 The experiment consisted of three fully crossed factors: (1) increased winter temperature
111 variability by application of winter warming pulses versus ambient reference conditions, (2)
112 two experimental sites with naturally different winter climate, (3) six different plant
113 communities and an additional bare ground control. The plant communities consisted of
114 three grassland communities (monocultures of the grass *Holcus lanatus* (L.) and the herb
115 *Plantago lanceolata* (L.), and a community with a mix of both species) and three heathland
116 communities (monocultures of the dwarf shrub *Calluna vulgaris* (L.) and the grass
117 *Deschampsia flexuosa* (L.) and a community with a mix of both species). All species present
118 in this experiment are very common perennial species in Central Europe. In addition, there
119 was a bare ground control in every block. Plant communities were blocked and randomly
120 assigned to the winter warming pulses manipulation and ambient reference. Temperature
121 manipulation blocks, and therefore each factorial combination, were replicated five times.
122 This setup was fully replicated at both experimental sites. For the 140 plots, plastic barrels
123 with 0.2 m² surface (50 cm diameter) and 80 cm depth were used as mesocosms. Each of
124 the six mesocosms per treatment was placed in a corner of a hexagon, with 30 cm distance

125 between mesocosms and at least 50 cm separation from the hexagon edge. The bare
126 ground control was placed in the middle of the hexagons. All space between the mesocosms
127 was filled with the same substrate as used within the mesocosms. The soil substrate was
128 homogenized loamy sand (77% sand, 16% silt, 7% clay) from a nearby sand quarry (where
129 all used plant species naturally occur), with a pH=7.35 (measured in 1 M KCl) and a total
130 carbon content of 2.37%. The barrels were attached with outlet hoses at the bottom of each
131 mesocosm, so that the mesocosms functioned as zero tension lysimeters. Sixteen plants per
132 mesocosm were planted in a systematic grid in May 2010. All plants were grown from seed
133 in January 2010, except for the dwarf-shrub *C. vulgaris*, which was obtained as 2-year old
134 individuals in February 2010.

135 2.2 Manipulation of winter temperature variability

136 Winter warming pulses were applied with six IR-heating lamps (250 W) located in between
137 the mesocosms at a height of 60 cm and surface heating wires (distance 20 cm, 400 W per
138 block), which resulted in 1900 W per block (7 mesocosms). The ambient reference
139 mesocosms were equipped with dummy lamps. Six warming pulses were administered
140 simultaneously for both sites between 15 December 2010 and 28 February 2011 (see Fig. 1).
141 Warming pulses were administered when there was soil frost at both sites and weather
142 forecast predicted further air frost for at least the next 48 h.

143 Soil temperature (-2 cm; once in every treatment and reference block; 10 measurements per
144 site and 20 in total) and air temperature (+5 cm; one treatment and reference block per site;
145 2 measurements per site and 4 in total) were measured hourly by thermistors (B57863-S302-
146 F40, EPCOS AG, Germany) connected to a datalogger (dl2, Delta-T Devices Ltd, UK). To
147 quantify the effect of the warming pulses treatment on soil temperature variability, we
148 calculated the coefficient of variation ($CV = \text{standard deviation} \times \text{hourly mean}^{-1} \times 100$;
149 temperatures were converted to K for this). Snow height was measured each morning via a
150 webcam picture of a measuring stick.

151 2.3 Response parameters

152 Plant available N was measured via the resin stick method (Plant-root-simulator (PRSTM)-
153 probes; Western Ag Innovations Inc., Canada). Two cation and two anion PRSTM-probes
154 were installed vertically with a distance of 20 cm to each other (0 -15 cm depth) per
155 mesocosm prior to the warming pulse manipulation on 18 December 2010 and collected on
156 17 March 2011 after the winter warming pulses treatment. PRSTM-probes were cleaned and
157 kept in a fridge until being sent to Western Ag Innovations Inc. (Canada) in a cool box for
158 analysis. For the statistical analysis, nitrate and ammonium were pooled due to low
159 ammonium concentrations. The maximum ion capacity of the probes for nitrate is 2088 $\mu\text{g } 10$
160 cm^{-2} . The values in our study are far lower, showing that the system was not saturated. For
161 better comparability to other studies we give mean plant available N per cm^{-2} and day. But it
162 is important to note that N uptake by resin sticks is not a linear process.

163 Soil biotic activity, i.e. decomposition by microorganisms and feeding by soil fauna, was
164 measured via bait-lamina sticks (terra protecta GmbH, Germany) (Kratz, 1998). One bait-
165 lamina stick containing 16 baits was inserted vertically in the top soil layer of every
166 mesocosm prior to the warming pulses treatment on 18 December. The baits consisted of a
167 mixture of powdered cellulose, bran flakes and active coal. These baits are potentially eaten

168 by earthworms, macro- to micro arthropods and additionally are decomposed by soil
169 microorganisms. The sticks were collected after the winter warming pulses treatment on 17
170 March, cleaned, and the number of eaten baits was counted. For the latter, sticks were
171 placed on a light bench and when light shined through the baits they were counted as eaten.
172 This analysis was done by a single person who was blind to the factors.

173 For the potential extracellular enzymatic activity (PEEA), which we used as another proxy for
174 soil biotic activity and decomposition, three soil samples (2 cm diameter, 10 cm depth) per
175 mesocosm were collected and mixed for assays of potential extracellular enzyme activity in
176 soil on 21 February 2011. Soil samples were stored in airtight plastic zip-bags at 4°C and
177 were analysed within 3 days. PEEA assays were carried out with Methylumbelliferone
178 substrates (MUF) (Pritsch et al., 2004; Pritsch et al., 2005). The following PEEAs were
179 measured: MU-β-D-glucopyranoside (MU-G), for β-glucosidase, MU-β-cellobioside (MU-C)
180 for cellobiohydrolase, MU-β-D-xylopyranoside (MU-X) for xylosidase, MU-phosphate (MU-P)
181 for acid phosphatase. Substrates and calibration saturation and incubation times were
182 determined in pre-experiments (data not shown) as follows: MU-G and MU-X each 500 μM
183 incubating for 60 min, MU-C 500 μM incubating for 120 min, MU-P 800 μM incubating for 40
184 min. Fluorescence was detected at an excitation wavelength of 360 nm and an emission
185 wavelength of 450 nm with a Gemini EM Fluorescence Microplate Reader from Molecular
186 Device, California.

187 Prior to the warming pulses treatment (18 December 2010), plots were labelled with 0.02 g
188 Potassium Nitrate-¹⁵N (min. 99.19 atom % ¹⁵N; Campro Scientific GmbH, Germany),
189 dissolved in 250 ml deionized water, resulting in 0.1 g ¹⁵N m⁻². Leaf (2-3 medium aged leaves
190 per plot and species, randomly chosen), root (fine roots from a soil sample taken directly next
191 to a randomly chosen plant per mesocosm and species) and soil samples (3 soil samples per
192 plot were mixed; 2 cm diameter, 10 cm depth) were taken on 17 March 2011, after the winter
193 warming pulses treatment. The samples were kept frozen until they were cleaned, dried (48
194 h at 50° C) and ball milled. Mass spectroscopy analysis was done at the laboratory of Isotope
195 Biogeochemistry, BayCEER, University of Bayreuth, with a combination of an elemental
196 analyzer (Carlo Erba NC 2500, CE Instruments, Italy) and an isotope mass spectrometer
197 (delta plus, Thermo Fisher Scientific, Germany). Atom % increase values for plant and soil
198 material collected after the winter warming pulses treatment were calculated by comparing to
199 values obtained from unlabelled reference plants (n = 5 per species) and soil material taken
200 prior to the winter warming pulses treatment (n = 3 per experimental site). Due to missing
201 volume readings, the isotopic signature of leachate could only be determined and related to
202 volume of leachate for four mesocosms (*Holcus lanatus* and *Plantago lanceolata* mixed
203 mesocosms at both sites for both winter warming pulses treatments), which were
204 permanently equipped by tipping buckets (7041.3000X, Theodor Friedrichs & Co., Germany).
205 Therefore, no mass balancing of the label was possible, and we report ¹⁵N-atom% here. For
206 interpretation of the data it is important to note that overall above-ground biomass
207 significantly decreased by 9.2 % due to the warming pulses treatment (Schuerings et al.,
208 2014). For single species, only *H. lanatus* showed a strong decrease by 29.2 % whereas the
209 other species showed no significant treatment effects (Schuerings et al., 2014).

210 2.4 Data analyses

211 Linear mixed-effect models combined with analysis of variance (ANOVA) were applied to test
212 for significant winter warming pulses treatment, site and plant community effects. All possible
213 interactions of community or species and site with the warming pulses treatment were
214 included as fixed effects (s. Table 2 & 3 for all tested interactions). For the analysis of ¹⁵N
215 content in plants, species identity was included as a fixed factor instead of community
216 composition, whereas community was included as a random effect. Block identity was set as
217 a random effect in all models, thereby accounting for the blocked design. Before statistical
218 analysis, we tested for normality and homogeneity of variance by examining the residuals
219 versus fitted plots and the normal qq-plots of the linear models (Faraway, 2005). If conditions
220 were not satisfactorily met, we applied log(x)- (plant available N; ¹⁵N atom% increase of
221 leaves and roots; PEEA of beta-glucosidase, cellobiohydrolase, xylosidase), log(x+1)- (¹⁵N
222 atom% increase in soil), or sqrt(x)- (PEEA acid phosphatase) transformation. Significance
223 level was set to p < 0.05. All statistical analyses were performed using R 2.12.2 (R
224 Development Core Team 2011) and additional packages nlme (Version 3.1-98, 2011) and
225 sciplot (Version 1.0-9, 2011) for graphical illustrations.

226

227 3 RESULTS

228 The winter warming pulses manipulation successfully decreased snow cover and resulted in
229 increased soil temperature variability (Fig. 1). At the warm site, variation in soil temperature
230 during the manipulation period (15 December 2010 to 28 February 2011) was increased to
231 CV = 0.99 in comparison to CV = 0.66 in the reference mesocosms. Mean soil temperature
232 increased to 1.8°C in the manipulation as compared to 0.1°C in the ambient reference.
233 Minimum temperature reached -4.2 °C and -4.0 °C, respectively. For the cold site, variation
234 in soil temperature during the manipulation period increased to CV= 0.68 in comparison to
235 CV= 0.43 in the reference mesocosms. Mean soil temperature was almost unchanged with -
236 0.1°C in the warming pulses manipulation and -0.3°C under ambient reference conditions.
237 However, minimum temperature was considerably lower in the warming pulses mesocosms,
238 reaching -4.7 °C, as compared to -2.6 °C in the reference mesocosms. The number of soil
239 freeze thaw cycles was not altered noticeably at any site (warm site: 7 vs. 8, cold site: 6 vs.
240 5).

241 Plant available nitrate and ammonium significantly increased by 34.5% in response to the
242 winter warming pulses treatment (F=13.5, p<0.001; Table 2, Fig. 2). The cold site overall had
243 a 48.4% higher amount of N available than the warm site (F=20.0, p<0.001; Table 2, Fig. 2).
244 Plant community composition also influenced plant available N (F=18.4, p<0.001; Table 2,
245 Fig. 2). Bare ground control mesocosms had the highest N values, followed by the heathland
246 communities and then the grassland communities, with only monocultures of *H. lanatus*
247 reaching levels of the heathland communities. Winter warming pulse effects were not
248 influenced by site or plant community (no significant interactions, Table 2).

249 Soil biotic activity, i.e. the number of eaten baits, increased by 40% (F=17.5, p<0.001; Table
250 2, Fig. 2) due to the winter warming pulses treatment in comparison to reference conditions.
251 Soil biotic activity did not significantly differ between sites or plant communities. The warming
252 pulses effect, however, was influenced by the plant communities (F=2.3, p=0.037), with
253 slightly decreasing activities in monocultures of *P. lanceolata* and mixed communities of *C.*

254 *vulgaris* & *D. flexuosa* due to the warming pulses (Fig. 2). All other communities showed an
255 increase in soil biotic activity due to the warming pulses. No other interaction with the
256 warming pulses treatment yielded significance for soil biotic activity (Table 2).

257 Regarding PEEA there was a general trend towards higher values under the winter warming
258 pulses treatment, yet only for cellobiohydrolase was this effect statistically significant ($F=5.3$,
259 $p=0.035$). For the other three tested enzymes no significant effect of the winter warming
260 pulses treatment was observed. Generally, there were significantly higher PEEAs at the cold
261 site than at the warm site (Table 2, Fig. 3) and plant community composition effects differed
262 such that, except for acid phosphatase, grassland communities showed higher PEEA than
263 heathland communities (Table 2, Fig. 3). No significant interactions between the warming
264 pulses treatment and site or plant community were observed (Table 2).

265 The AT% ^{15}N values in leaves were significantly reduced by 21.7% (relative difference) under
266 the winter warming pulses treatment in comparison to reference conditions ($F=5.9$, $p=0.016$),
267 whereas for root and soil material no significant winter warming pulse effect was observed
268 (Table 3, Fig. 4). For leachate, no statistical analysis was performed due to the low
269 replication, but for the existing samples ($n=2$ per winter warming pulses treatment), a clear
270 trend towards increased leaching of the ^{15}N -tracer was observed (Fig. 4). Generally, the cold
271 site showed significantly higher plant AT% ^{15}N values than the warm site (Table 3, Fig. 4). *D.*
272 *flexuosa* exhibited the highest AT% ^{15}N values, followed by *P. lanceolata*, with the same
273 pattern observed for leaves and roots. Significant decreases in the ^{15}N signal in plant leaves
274 (-30.7%) in response to warming pulses only occurred at the cold site (winter warming pulses
275 treatment x site interaction: $F=8.6$, $p=0.004$; Table 3, Fig. 4). The significant three-way
276 interaction between warming pulses treatment, site, and species identity ($F=3.4$, $p=0.004$)
277 indicated that the decrease in ^{15}N values only happened at the cold site and only for three of
278 the four species (*C. vulgaris*, *D. flexuosa* and *H. lanatus*; Fig. 4).

279

280 4 DISCUSSION

281 Recurrent winter warming pulses led to increased soil temperature variability and influenced
282 N cycling in our experiment. As expected, N availability was increased (+35%) in the
283 mesocosms which received the winter warming pulses treatment. Increased N availability
284 during winter/early spring is often explained by freeze-thaw events resulting in increased
285 biological and physical decomposition of soil organic matter (SOM) (Matzner and Borken,
286 2008) and increased N mineralization (Rustad et al., 2001; Melillo et al., 2002). Yet, in our
287 study FTC frequency was merely changed between winter warming pulses and references
288 plots (± 1), implying that the warming pulses treatment affected N availability either through
289 increased temperature variability or the increase in mean temperature. Due to the winter
290 warming pulses soil biotic activity increased by 40%. This increase in soil biotic activity is in
291 line with results from other winter warming experiments which measured soil respiration as
292 an index of soil biotic activity (Davidson and Janssens, 2006; Allison and Treseder, 2011).
293 The soil enzymes we examined play a major role in the decomposition of biological material
294 (Marx et al., 2001). We observed significantly increased PEEA for cellobiohydrolase,
295 whereas for the other three tested enzymes the observed increases were not significant.
296 Therefore in our experiment, increased soil temperature variability led to increased biotic

297 decomposition as indicated by increased soil biotic activity and increased PEEA of
298 cellobiohydrolase. In winter warming experiments, increased N cycling is often attributed to
299 changes in the frequency of soil FTC (Mikan et al., 2002). Despite only small changes in FTC
300 frequency in our mesocosms, however, we observed increased N availability, increased soil
301 biotic and soil potential enzymatic activity. However, for the cold site, where it is important to
302 note that mean soil temperature only increased by 0.2 °C, mean minimum temperature was
303 considerably lower in the warming pulses mesocosms, reaching -4.7 °C, as compared to -2.6
304 °C. Since we found lowered N incorporation into plants (see discussion further down) and
305 stable or lower plant biomass (Schuerings et al., 2014) at the cold site, this could have
306 lowered N immobilization by plants. The temporal dynamics of soil temperature, in particular
307 the intensity of freezing right after warming pulses, is therefore another important
308 determinant of N cycling responses, possibly leading to frost damaging of dehardend plants.
309 While changed FTCs (Joseph and Henry, 2008), warmer mean soil temperatures (Rustad et
310 al., 2001; Melillo et al., 2002) and single extreme frost events (Elliott and Henry, 2009) are
311 known to be important drivers of N cycling, our results imply that soil temperature variability,
312 i.e. temperature dynamics, can also affect N availability and soil biotic activity.

313 We found significantly higher N availability and potential activity of all four tested potential
314 soil enzymes for the cold site despite lower mean temperatures at the site. Groffman et al.
315 (2009) found the same pattern along an altitudinal gradient in a northern hardwood forest.
316 This suggests that the local climate may have an important influence on the magnitude of N
317 mobilization processes. However, since we found no significant interaction between winter
318 warming pulses treatment and site, the effects of winter warming pulses on N availability, soil
319 biotic activity and potential soil enzymatic activity therefore appear independent of the local
320 climate.

321 The mobilization of N was influenced by the plant community composition, with the bare
322 ground control showing highest levels of available N. Since there were no roots in the bare
323 ground plots competing with the PRSTM-probes for N, this result is not surprising. Regarding
324 plant communities, there was no clear pattern in N availability, although the heathland
325 communities showed higher values than grassland communities with the exception of
326 monocultures of *H. lanatus*, which showed similar values as the heathland communities. The
327 interaction between the warming pulses treatment and plant community indicated that plant
328 species composition influenced soil biotic activity differently under winter warming pulses.
329 However, there was no clear pattern, since all communities showed increased soil biotic
330 activity in response to the winter warming pulses, except for monocultures of *P. lanceolata*
331 and mixed cultures of *C. vulgaris* and *D. flexuosa*. Potential soil enzymatic activity was
332 generally higher in grassland mesocosms in comparison to heathland mesocosms, with the
333 exception of acid phosphatase.

334 The ¹⁵N signal in plants leaves was, contrary to our expectations, decreased by the winter
335 warming pulses treatment. Plants can lose their cold hardiness within hours in response to
336 elevated temperatures (Kalberer et al., 2006), and subsequent frost events after a winter
337 warm spell can thus damage plants substantially (Bokhorst et al., 2009). Freezing intensity is
338 also an important determinant of plant frost damage, and while most temperate species can
339 tolerate temperatures at or below freezing, there is often a threshold subfreezing temperature
340 where damage intensifies (Malyshev and Henry, 2012a). Notably, the minimum temperatures
341 reached in the reference mesocosms at the cold site were the least severe, and the highest

342 AT% ¹⁵N values were observed in these plots, whereas minimum soil temperatures of at
343 least -4 °C were reached in the treatment plots at the cold site and in all of the warm site
344 mesocosms, all of which featured relatively low ¹⁵N values. Similarly, in other systems, grass
345 ecotypes located at northern sites that are protected from cold air by thick snow cover have
346 developed lower frost tolerance than conspecific ecotypes located in warmer locations that
347 feature less snow cover, because the latter ecotypes experience more intense frost (Dionne
348 et al., 2010).

349 We also observed significant differences among the tested species in the increase of AT%
350 ¹⁵N values, which is not surprising, given that species exhibit wide variation in their nutrient
351 uptake capacities (Hooper and Vitousek, 1998; Knops et al., 2002). The interesting point is
352 that the reduction in ¹⁵N values only happened at the cold site and only for *C. vulgaris*, *D.*
353 *flexuosa* and *H. lanatus* (interaction: winter warming pulses treatment x site x species). Total
354 above-ground biomass of all tested species decreased by 9.2 % in response to the winter
355 warming pulses treatment (Schuerings et al., 2014), thus dilution effects on N-tracer uptake
356 can be excluded. Lower or stable above-ground biomass and lower AT% ¹⁵N values
357 combined are a clear hint for reduced N uptake by the affected plant species. Such
358 differences among species in frost susceptibility could have important consequences for
359 competitive balances and shifts in community composition over the long term (Joseph and
360 Henry, 2008; Cornelissen and Makoto, 2014).

361 Chronic winter warming can increase above-ground biomass (Hutchison and Henry, 2010;
362 Natali et al., 2012; Schuerings et al., 2013). This additional growth may be fuelled by
363 increased N mobilization in early spring. Pulsed winter warming increasing the risk of frost
364 damage, however, complicates this simple expectation of increased plant growth under
365 winter climate change. The inability of frost-damaged plants to take up the available N in the
366 soil solution might trigger N losses from ecosystems by N leaching or gaseous losses
367 (Ineson et al., 1998; Campbell et al., 2014). In this experiment we also found species-specific
368 responses in above-ground biomass production due to the winter warming pulses
369 (Schuerings et al., 2014); only *H. lanatus* showed a decrease in above-ground biomass,
370 whereas the other tested species remained unaffected by the winter warming pulses
371 treatment in their above-ground productivity. Taken together, species- or vegetation type-
372 specific responses have to be taken into account when forecasting effects of climate change
373 on N-cycling (Makoto et al., 2014). Furthermore, regarding winter climate change, pulsed
374 warming events can result in opposing effects on N cycling and biomass accumulation than
375 chronic warming.

376

377 **5 CONCLUSIONS**

378 Future winters in the temperate zone are expected to be characterized by more variable soil
379 temperatures due to increasing air temperature variability and due to missing insulation by
380 snow. Our experiment implies that more variable soil temperatures enhance nitrogen
381 mobilization in the soil independent from vegetation types and the local climate. Plant
382 performance, however, depended on local climate, with plant ¹⁵N immobilization during
383 winter and early spring after exposure to winter warming pulses being reduced at colder
384 sites, probably due to frost damage after the warming pulses. This pattern implies increased

385 risk for nitrogen leaching at colder temperate sites in response to increased winter
386 temperature variability. Taken together, our findings emphasize the importance of
387 temperature variability, plant performance, and frost damage in a warmer world for nitrogen
388 cycling and nitrogen losses from ecosystems.

389

390 **ACKNOWLEDGEMENTS**

391 This study was funded by the German Science Foundation (DFG JE 282/5-1). The isotope
392 analysis was done by the BayCEER Laboratory of Isotope Biogeochemistry led by Prof. Dr.
393 Gebauer. We thank Elke and Stefan König for installing the field experiment.

REFERENCES

- 394
395
396 Allison, S. D. and Treseder, K. K.: Climate change feedbacks to microbial decomposition in
397 boreal soils, *Fungal Ecol.* 4, 362–374, 2011.
- 398 Andresen, L. C. and Michelsen, A.: Off-season uptake of nitrogen in temperate heath
399 vegetation, *Oecologia*, 144, 585–597, 2005.
- 400 Bardgett, R. D., Streeter, T. C., and Bol, R.: Soil microbes compete effectively with plants for
401 organic-nitrogen inputs to temperate grasslands, *Ecology*, 84, 1277–1287, 2003.
- 402 Bokhorst, S. F., Bjerke, J. W., Tommervik, H., Callaghan, T. V., and Phoenix, G. K.: Winter
403 warming events damage sub-Arctic vegetation: consistent evidence from an experimental
404 manipulation and a natural event, *J. Ecol.*, 97, 1408–1415, 2009.
- 405 Boutin, R. and Robitaille, G.: Increased soil nitrate losses under mature sugar maple trees
406 affected by experimentally induced deep frost, *Can. J. Forest Res.*, 25, 588–602, 1995.
- 407 Brooks, P. D., Williams, M. W., and Schmidt, S. K.: Inorganic nitrogen and microbial biomass
408 dynamics before and during spring snowmelt, *Biogeochemistry*, 43, 1–15, 1998.
- 409 Brown, P. J. and DeGaetano, A. T.: A paradox of cooling winter soil surface temperatures in
410 a warming northeastern United States, *Agr. Forest Meteorol.*, 151, 947–956, 2011.
- 411 Christensen, J. H., Hewitson, B., Busuioc, A., Chen, A., Gao, X., Jones, R., et al.: Regional
412 climate projections. In: Solomon, S., Qin, D., Manning, M., Chen, Z., and Marquis, M.
413 (eds) *Climate change 2007: The physical science basis*, Contribution of Working Group I
414 to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change,
415 Cambridge Univ. Press, Cambridge, pp 847–940, 2007.
- 416 Clein, J. S. and Schimel, J. P.: Microbial activity of tundra and taiga soils at sub-zero
417 temperatures, *Soil Biol. Biochem.*, 27, 1231–1234, 1995.
- 418 Cornelissen, J. H. C. and Makoto, K.: Winter climate change, plant traits and nutrient and
419 carbon cycling in cold biomes, *Ecol. Res.*, 29, 517–527, 2014.
- 420 Davidson, E. A. and Janssens, I. A.: Temperature sensitivity of soil carbon decomposition
421 and feedbacks to climate change, *Nature*, 440, 165–173, 2006.
- 422 Dionne, J., Rochefort, S., Huff, D. R., Desjardins, Y., Bertrand, A., and Castonguay, Y.:
423 Variability for Freezing Tolerance among 42 Ecotypes of Green-Type Annual Bluegrass,
424 *Crop Sci.*, 50, 321–336, 2010.
- 425 Elliott, A. C. and Henry, H. A. L.: Freeze–thaw cycle amplitude and freezing rate effects on
426 extractable nitrogen in a temperate old field soil, *Biol. Fertil. Soils*, 45, 469–476, 2009.
- 427 Faraway J. J.: *Linear models with R*, Chapman & Hall/CRC, Boca Raton, 2005.
- 428 Groffman, P. M., Hardy, J. P., Fisk, M. C., Fahey, T. J., and Driscoll, C. T.: Climate Variation
429 and Soil Carbon and Nitrogen Cycling Processes in a Northern Hardwood Forest,
430 *Ecosystems*, 12, 927–943, 2009.
- 431 Grogan, P., Michelsen, A., Ambus, P., and Jonasson, S.: Freeze-thaw regime effects on
432 carbon and nitrogen dynamics in sub-arctic heath tundra mesocosms, *Soil Biol. Biochem.*,
433 36, 641–654, 2004.
- 434 Henry, H. A. L.: Climate change and soil freezing dynamics: historical trends and projected
435 changes, *Climatic Change*, 87, 421–434, 2008.
- 436 Hooper, D. U. and Vitousek, P. M.: Effects of plant composition and diversity on nutrient
437 cycling. *Ecol. Monogr.*, 68, 121–149, 1998.

- 438 Hutchison, J. S. and Henry, H. A. L.: Additive effects of warming and increased nitrogen
439 deposition in a temperate old field: Plant productivity and the importance of winter,
440 *Ecosystems*, 13, 661–672, 2010.
- 441 Ineson, P., Taylor, K., Harrison, A. F., Poskitt, J., Benham, D. G., Tipping, E., and Woof, C.:
442 Effects of climate change on nitrogen dynamics in upland soils. 1. A transplant approach,
443 *Global Change Biol.*, 4, 143–152, 1998.
- 444 Joseph, G. and Henry, H. A. L.: Soil nitrogen leaching losses in response to freeze-thaw
445 cycles and pulsed warming in a temperate old field, *Soil Biol. Biochem.*, 40, 1947–1953,
446 2008.
- 447 Kalberer, S. R., Wisniewski, M., and Arora, R.: Deacclimation and reacclimation of cold-
448 hardy plants: Current understanding and emerging concepts, *Plant Sci.*, 171, 3–16, 2006.
- 449 Knops, J. M. H., Bradley, K. L., and Wedin, D. A.: Mechanisms of plant species impacts on
450 ecosystem nitrogen cycling, *Ecol. Lett.*, 5, 454–466, 2002.
- 451 Kodra, E., Steinhäuser, K., and Ganguly, A. R.: Persisting cold extremes under 21st-century
452 warming scenarios, *Geophys. Res. Lett.*, 38, 2011.
- 453 Kratz, W.: The bait-lamina test – General aspects, applications and perspectives, *Environ.*
454 *Sci. Pollut. Res.*, 5, 94–96, 1998.
- 455 Kreyling, J.: Winter climate change: A critical factor for temperate vegetation performance,
456 *Ecology*, 91, 1939–1948, 2010.
- 457 Kreyling, J., Beierkuhnlein, C., and Jentsch, A.: Effects of soil freeze-thaw cycles differ
458 between experimental plant communities, *Basic Appl. Ecol.*, 11, 65–75, 2010.
- 459 Kreyling, J., Beierkuhnlein, C., Pritsch, K., Schloter, M., and Jentsch, A.: Recurrent soil
460 freeze–thaw cycles enhance grassland productivity, *New Phytol.*, 177, 938–945, 2008.
- 461 Kreyling, J. and Henry, H. A. L.: Vanishing winters in Germany: soil frost dynamics and snow
462 cover trends, and ecological implications, *Climate Res.*, 46, 269–276, 2011.
- 463 Makoto, K., Kajimoto, T., Koyama, L., Kudo, G., Shibata, H., Yanai, Y., and Cornelissen, J.
464 H. C.: Winter climate change in plant–soil systems: summary of recent findings and future
465 perspectives, *Ecol. Res.* 29, 593–606, 2014.
- 466 Malyshev, A. V. and Henry, H. A. L.: N uptake and growth responses to sub-lethal freezing in
467 the grass *Poa pratensis* L., *Plant Soil*, 360, 175–185, 2012a.
- 468 Malyshev, A. V. and Henry, H. A. L.: Frost damage and winter nitrogen uptake by the grass
469 *Poa pratensis* L.: Consequences for vegetative versus reproductive growth, *Plant Ecol.*,
470 213, 1739–1747, 2012b.
- 471 Marx, M. C., Wood, M., and Jarvis, S. C.: A microplate fluorimetric assay for the study of
472 enzyme diversity in soils, *Soil Biol. Biochem.*, 33, 1633–1640, 2001.
- 473 Matzner, E. and Borken, W.: Do freeze-thaw events enhance C and N losses from soils of
474 different ecosystems? A review, *Eur. J. Soil Sci.*, 59, 274–284, 2008.
- 475 Macgillivray, C. W., Grime, J. P., Band, S. R., Booth, R. E., Campbell, B., Hendry, G. A. F., et
476 al.: Testing predictions of the resistance and resilience of vegetation subjected to extreme
477 events, *Funct. Ecol.*, 9, 640–9, 1995.
- 478 Melillo, J. M., Steudler, P. A., Aber, J. D., Newkirk, K., Lux, H., Bowles, F. P., Catricala, C.,
479 Magill, A., Ahrens, T., and Morrisseau, S.: Soil warming and carbon-cycle feedbacks to
480 the climate system, *Science*, 298, 2173–2176, 2002.
- 481 Mikan, C. J., Schimel, J. P., and Doyle, A. P.: Temperature controls of microbial respiration in
482 arctic tundra soils above and below freezing, *Soil Biol. Biochem.*, 34, 1785–1795, 2002.

- 483 Nasholm, T., Huss-Danell, K., and Hogberg, P.: Uptake of organic nitrogen in the field by four
484 agriculturally important plant species, *Ecology*, 81, 1155–1161, 2000.
- 485 Natali, S. M., Edward, A. G., Schuur, G., and Rubin, R. L.: Increased plant productivity in
486 Alaskan tundra as a result of experimental warming of soil and permafrost, *J. Ecol.*, 100,
487 488–498, 2012.
- 488 Oztas, T. and Fayetorbay, F.: Effect of freezing and thawing processes on soil aggregate
489 stability, *Catena*, 52, 1–8, 2003.
- 490 Patil, R. H., Laegdsmand, M., Olesen, J. E., and Porter, J. R.: Effect of soil warming and
491 rainfall patterns on soil N cycling in Northern Europe, *Agr. Ecosyst. Environ.*, 139, 195–
492 205, 2010.
- 493 Pritsch, K., Luedemann, G., Matyssek, R., Hartmann, A., Schloter, M., Scherb, H., and
494 Grams, T. E. E.: Mycorrhizosphere Responsiveness to Atmospheric Ozone and
495 Inoculation with *Phytophthora citricola* in a Phytotron Experiment with Spruce/Beech
496 Mixed Cultures, *Plant Biol.*, 7, 718–727, 2005.
- 497 Pritsch, K., Raidl, S., Marksteiner, E., Blaschke, H., Agerer, R., Schloter, M., and Hartmann,
498 A.: A rapid and highly sensitive method for measuring enzyme activities in single
499 mycorrhizal tips using 4-methylumbelliferone-labelled fluorogenic substrates in a
500 microplate system, *J. Microbiol. Meth.*, 58, 233–241, 2004.
- 501 Rustad, L. E., Campbell, J. L., Marion, G. M., Norby, R. J., Mitchell, M. J., Hartley, A. E.,
502 Cornelissen, J. H. C., and Gurevitch, J.: A meta-analysis of the response of soil
503 respiration, net nitrogen mineralization, and aboveground plant growth to experimental
504 ecosystem warming, *Oecologia*, 126, 543–562, 2001.
- 505 Schuerings, J., Beierkuhnlein, C., Grant, K., Jentsch, A., Malyshev, A., Penuelas, J.,
506 Sardans, J., and Kreyling, J.: Absence of soil frost affects plant-soil interactions in
507 temperate grasslands, *Plant Soil*, 371, 559–572, 2013.
- 508 Schuerings, J., Jentsch, A., Walter, J., and Kreyling, J.: Winter warming pulses differently
509 affect plant performance in temperate heathland and grassland communities, *Ecol. Res.*,
510 29, 561–570, 2014.
- 511 Skogland, T., Lomeland, S., and Goksoyr, J.: Respiratory burst after freezing and thawing of
512 soil – experiments with soil bacteria, *Soil Biol. Biochem.*, 20, 851–856, 1988.
- 513 Sturm, M., Holmgren, J., König, M., and Morris, K.: The thermal conductivity of seasonal
514 snow, *J. Glaciol.*, 43, 26–41, 1997.
- 515 Tierney, G. L., Fahey, T. J., Groffman, P. M., Hardy, J. P., Fitzhugh, R. D., and Driscoll, C.
516 T.: Soil freezing alters fine root dynamics in a northern hardwood forest, *Biogeochemistry*,
517 56, 175–190, 2001.
- 518 Turner, M. M. and Henry, H. A. L.: Interactive effects of warming and nitrogen deposition on
519 net nitrogen mineralization and leaching losses in a temperate old field: the importance of
520 winter, *Oecologia*, 162, 227–236, 2010.

521

TABLES

522

523 Table 1: Climate characteristics of the two experimental sites, measured on site by the
524 department of Micrometeorology until 2008; University of Bayreuth, Prof. T. Foken
525 (Schuerings et al., 2014)

526

Parameter (Unit; start of measurements warm site / cold site)	Warm site	Cold site
Mean annual temperature (°C; 1998 / 1994)	8.8	5.0
Mean winter temperature (DJF; °C; 1998 / 1994)	0.6	-2.0
Mean annual precipitation (mm; 1998 / 1994)	717	1002
Mean winter precipitation (DJF; mm; 1998 / 1994)	158	237
Mean # of days with soil frost (-5 cm) (2003 / 1999)	19	31


527 Table 2: ANOVA-results of all tested main and interaction effects for N mobilization, i.e. N availability in the soil solution (NH₄⁺ and NO₃⁻), soil biotic
 528 activity (bait-lamina test), and the four tested potential soil enzyme activities. Warming pulses: Winter warming pulses treatment.
 529

Factor	N availability in soil solution		Soil biotic activity		Beta-glucosidase activity		Cellobiohydrolase activity		Acid phosphatase activity		Xylosidase activity	
	F	P	F	P	F	P	F	P	F	P	F	P
Warming pulses	13.5	<0.001	17.5	<0.001	1.8	0.199	5.3	0.035	2.6	0.127	2.0	0.173
Site	20.0	<0.001	0.6	0.441	67.2	<0.001	69.2	<0.001	12.6	0.003	33.6	<0.001
Community	18.4	<0.001	0.3	0.912	23.5	<0.001	16.2	<0.001	32.5	<0.001	44.5	<0.001
Warming pulses x Site	0.6	0.425	0.9	0.358	3.2	0.094	1.3	0.266	0.9	0.359	3.8	0.068
Warming pulses x Community	0.2	0.961	2.3	0.037	1.4	0.213	0.7	0.663	1.1	0.388	0.6	0.694
Warming pulses x Site x Community	0.6	0.715	1.1	0.370	0.7	0.685	1.0	0.400	0.9	0.500	1.4	0.212

530


531 Table 3: ANOVA-results of all tested main and interaction effects for the fate of a ¹⁵N label
 532 (increase in atom % ¹⁵N in the compartments leaves, fine roots, and bulk soil). Warming
 533 pulses: Winter warming pulses treatment.
 534

Factor	¹⁵ N atom % increase					
	Leaves		Roots		Bulk soil	
	F	P	F	P	F	P
Warming pulses	5.9	0.016	1.5	0.228	0.9	0.331
Site	144.5	<0.001	19.3	<0.001	29.9	<0.001
Species/Community (Soil)	7.4	<0.001	9.6	<0.001	1.7	0.134
Warming pulses x Site	8.6	0.004	2.1	0.153	2.0	0.162
Warming pulses x Species	1.2	0.313	0.5	0.695	0.7	0.647
Warming pulses x Site x Species	3.4	0.004	1.0	0.422	1.2	0.292




536

537 Fig. 1 Mean daily air temperature at +5 cm (a), snow depth (b) and mean daily soil
 538 temperature at -2 cm (c) at the two experimental sites for the winter warming pulses
 539 treatment (black line) and reference conditions (grey line). Warming pulses (grey boxes)
 540 were applied between 15th December 2010 and 28th February 2011 (Schuerings et al.,
 541 2014).



542


543 Fig. 2 (a) Plant available nitrogen (nitrate and ammonium; PRSTM-probes) and (b) soil biotic
 544 activity (bait-lamina test) during the manipulation period (18 December 2010 - 17 March
 545 2011). Main winter warming pulses treatment, site and community effects and all significant
 546 interactions between the winter warming pulses treatment with site and community are
 547 shown. Mean (\pm S.E.) values are shown (n=140).



548

549 Fig. 3 Mean potential soil enzymatic activity for the four tested enzymes (a) β -glucosidase,
 550 (b) cellobiohydrolase, (c) acid phosphatase and (d) xylosidase (all \pm S.E.) during the
 551 manipulation period (18 December 2010 - 17 March 2011). Main winter warming pulses

552 treatment, site and community effects are shown. No significant interactions between the
553 winter warming pulses treatment with site and community were detected.



554

555 Fig. 4 Mean increase in atom% values (\pm S.E.) for leaves (n=80), roots (n=80), bulk soil
556 (n=70) and leachate (n=2). Before the warming pulses treatment all plots were watered with
557 0.25 l of water with 0.02 g Potassium Nitrate-¹⁵N (min. 99.19 atom % ¹⁵N). Main winter
558 warming pulses treatment, site and community effects and all significant interactions
559 between the winter warming pulses treatment with site and community are shown. It is

560 important to note that total above-ground biomass declined by 9.2 % in the growing season
561 after manipulations, so that tracer dilution effects due to increasing biomass can be excluded
562 (Schuerings et al., 2014).