

BGD

11, 9103–9134, 2014

Maize yield development in Lower Saxony

J. F. Degener and
M. Kappas

Biomass yield development of early, medium and late Maize varieties under a future climate in Lower Saxony, Germany

J. F. Degener and M. Kappas

University of Göttingen, Institute of Geography, Göttingen, Germany

Received: 30 April 2014 – Accepted: 2 June 2014 – Published: 16 June 2014

Correspondence to: J. F. Degener (jdegene@gwdg.de)

Published by Copernicus Publications on behalf of the European Geosciences Union.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Abstract

Lower Saxony, with a total land area of about 46 500 km², constitutes one of the most important agricultural areas in Germany and thus within Europe. Roughly one third of its agricultural area is used for maize cultivation and as of today only few information exist on how a future changing climate will affect its local growing conditions. Thus the newly developed carbon-based crop model BioSTAR and a high-resolution regional climate data-set (WETTREG) were used to evaluate the change in biomass yields of an early, medium and late maize variety. The climate input data is based on the SRES A1B scenario, with a potential fertilization effect or better still, an increased water use efficiency due to rising CO₂ levels, taken into account. The biomass yield for all varieties was calculated for each year from 2001 until 2099 on a total of 91 014 sites. The results suggest clearly differentiated development paths of all varieties. All three show a significant positive trend until the end of the century. However the medium variety shows a statistical significant decline of 5 % during the first 30 years and only a slight recovery towards +5 % around the century's end. The late variety has the clearest and strongest positive trend, with partially more than 30 % increase of biomass yields around the end of the century or +25 % mean increase in the last three decades. The early variety can be seen as in-between, with no negative but also not an as strong positive development path. All varieties have their strongest increase in yields after the mid of the 21st century. Statistical evaluation of these results suggests that the shift from a summer rain to a winter rain climate in Germany will be the main limiting factor for all varieties. In addition summer temperatures will become less optimal for all maize crops. Only if the plants can supply themselves sufficiently with water outside of the increasingly dry summer months, when also temperatures are much more favorable, an increase in biomass yields is feasible. As the data suggests the increasing atmospheric CO₂ concentrations will play a critical role in reducing the crops water uptake, thus enabling yield increases in the first place.

BGD

11, 9103–9134, 2014

Maize yield development in Lower Saxony

J. F. Degener and
M. Kappas

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



1 Introduction

With a production of around 875 Mt, maize was the second most grown crop on earth in 2012, only surpassed by sugarcane and surpassing rice (3rd 718 Mt) and wheat (4th 675 Mt). However in terms of nutrition, rice and wheat provided around 3.8 times more calories to the world's average human (FAO, 2014). This spread in the data is a clear indicator for the variety of usage that maize allows for, from its first and foremost use as feed for livestock to a raw material for energy purposes.

As of 2012, like most years before, Lower Saxony (LS) constituted Germany's largest maize producer, accounting for more than a quarter of the 94.56 Mt total German production, while extending over merely 13 % of Germany's overall territory. This is due to an over-average yield of 50.6 t ha^{-1} (avg. Germany 46.4 t ha^{-1}) combined with a relative large cropping area of 27 % (avg. Germany 17 %) of the total utilized agricultural area (DeStatis, 2013).

Around the early 1980's the cropping area of LS for silage Maize leveled out at around 220 000 ha for several years. Around 2004 this began to change rapidly. Within five years the area nearly doubled, after less than a decade the area already amounted to 514 000 ha in 2012 (LWK, 2014a). An early look into the matter (Hoeher, 2007) did not show any increase in the local livestock nor a dramatic change in livestock diet or related im- or exports. Even more, the maize cropping area for feed receded by 30 000 ha between 2004 and 2007. Energy Maize however, in LS used predominantly as a regenerative power source, showed an increase in cropping area by 38 000 ha in only one year. Therefore it can be safely assumed that this increase in cropping area was due to reasons other than livestock farming. While there are some propositions for alternatives to this extensive maize cultivation (NMELVL, 2012; LWK, 2014b) its known production strategies and biomass yields will make it hard for any competing crop to replace maize. Thus it can be assumed that maize will be around for some time, raising the question how changing regional or local conditions will affect its yield potential.

BGD

11, 9103–9134, 2014

Maize yield development in Lower Saxony

J. F. Degener and
M. Kappas

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Maize yield development in Lower Saxony

J. F. Degener and
M. Kappas

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Wolf and van Diepen (1995) did an early estimation of the European grain maize yield potential, basically coming to the conclusion that no large changes are to be expected for the central part of the European Community and thus for LS. This outlook has not changed dramatically in present day studies (Supit et al., 2012), generally suggesting no trend or seldom a positive trend in rainfed maize yields for most parts of Germany. Spatial surveys covering only Germany in its entirety are relatively rare. However, it is often pointed out that maize already grows near optimal conditions in Germany and is as of today already limited through drought stress in its main growing period of July and August (Meyer et al., 2009; Taube and Herrmann, 2009). The expected further decline in summer precipitation of around 30 % for some areas in Germany (Jacob et al., 2008), however uncertain this change might be, would thus strongly limit the growing conditions of maize in these regions.

In part probably owed to the administrative structure in Germany, the assessment of climate change impacts on crops yields was mostly done on a federal state level. A wide variety of approaches (differences in climate model and dataset, crop model, reference period etc.) make a direct comparison often difficult at least. However, results from regions close to LS are still of special interest for comparison.

The federal state of Hesse, directly to the south of LS, shows a regional differentiated pattern with a positive maize yield trend (up to +15 %) in the southern part and a neutral to negative (mostly around –10 %) northern part in middle of the 21st century under the SRES B2 scenario (USF, 2005). To the south-west of LS lies North Rhine-Westphalia. Fröhlich (2010) did show that most of the state will profit from a changing climate from an silage maize yield increase of around 2–4 % (B1 scenario) or 3–7 % (A1B scenario) until 2050. Saxony and Thuringia, both to the south-east of LS, have a generally negative development until 2050 (A1B) with a decline in maize yields of roughly –10 % (Mirschel et al., 2008, 2012). Especially the study for Thuringia does show how wide these results may spread, even if climate model, scenario and crop model are kept the same. Four alternative approaches, including more or less progress in cultivation and

breeding, further differentiated by dry or moist conditions, resulted here in an average yield changes from -8.2% to $+38.6\%$.

Furthermore Buttlar et al. (2013) took a closer look at a part of LS, the region connecting the cities of Hanover, Brunswick, Göttingen and Hildesheim. This study was however rather site specific, with biomass yield changes of maize between -3% and $+7\%$ (until mid-century) and -4% to $+13\%$ (end of century).

While the current study does not expect to diverge largely from these findings, a regional or even local approach was necessary as a probable basis for action of regional decision makers. An important difference to the mentioned studies lies however in the selection of different maize varieties. For simplification many studies omit the use of different varieties that are differentiated only by their required temperature sums to reach their respective development stages. As Southworth et al. (2000) could show in a study in the Midwestern United States this differentiation can indeed make a difference, as heat-resistant late (or long-term) varieties did show a considerably better yield development in a future climate than varieties with less temperature requirements. However rare, if studies do evaluate distinct varieties, the findings are similar as Liu et al. (2013) could show for Northeast China. Most studies however only hint in a more general way towards the influence of variety choice (Wolf and van Diepen, 1994; Kwabiah, 2004; Meza et al., 2008).

2 Materials and methods

The basic approach used in this study was to use high-resolution climate data in combination with detailed soil information as the input for a crop model. All components involved are introduced in the following sections.

BGD

11, 9103–9134, 2014

Maize yield development in Lower Saxony

J. F. Degener and
M. Kappas

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.1 Research area

25 Lower Saxony (LS), with roughly 46 500 km² of land area, is the second largest of the 16 federal states of Germany, providing around 15 % of the nations agricultural land (DeStatis, 2013). Located to the North-West of Germany (Fig. 1) the state lies in a transition zone between a more maritime (NW) towards a more continental climate (SE) (Seedorf and Meyer, 1992) with an average annual temperature of around 9 °C
5 and a mean precipitation of 749 mm in the period of 1971–2000 (DWD, 2014).

Principally LS consists of three distinguishable landscape structures: the coast, including the East Frisian Islands, the German North-Western Lowland (amounting for three quarters of LS' total land area) as well as a low mountain range to its south, with the Harz as its most prominent representative (Drachenfels, 2010). The broad
10 loess valleys to the south and especially the fertile “Börde” that fronts the low mountain range to the north are the main cultivation areas for high-demand crops like winter wheat. The Lowland mainly consists of “Geest”-land, Quaternary sediments that are particularly sandy to the North-East, with precipitation as low as 500 mm, making irrigation already today necessary on several sites. The west of LS is dominated by
15 livestock farming with the coastal area predominantly used for grassland farming as high ground water levels prevent intensive use (Heunisch et al., 2007).

The regional differences manifest themselves in the average regional yields. In the period of 2003–2008 the average winter wheat yield south of Hanover was always above 8 t ha⁻¹, above 7 t ha⁻¹ south of Oldenburg and generally below 7 t ha⁻¹ in the
20 North-East. Maize yields behave rather similar, with dry maize silage (33 % dry matter content) having the best yields to the south. The margin between the different parts of LS is however smaller for maize than for wheat and varies generally around 15 t ha⁻¹. As can be seen in Fig. 1 the areas with the largest maize production coincide with areas where only little wheat is grown and where feed for livestock is in high demand.

Maize yield development in Lower Saxony

J. F. Degener and
M. Kappas

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.2 Crop model

The crop model used in this study is a relatively new model called *BioSTAR*, developed at the Georg-August-University in Göttingen (Bauböck, 2013). The model uses a CO₂ based crop development engine, thus taking a potential CO₂ fertilization effect into account.

The basic working principle uses temperature to determine the plants development stages and a combination of temperature, solar radiation and CO₂ concentration the maximum photosynthesis rate. Both incrementally build the plants maximum possible biomass that is then limited by precipitation respectively the soil water content. The model is suitable for large-scale as well as parcel size yield assessments. The philosophy behind it is an easy to use model with a robust output and a manageable amount of required input parameters.

The model was validated on sample sites in Lower Saxony with a general disagreement between actual and modeled yield of around 10%. The required climatic input variables are precipitation, temperature, atmospheric CO₂ concentration, solar radiation, relative air humidity and wind-speed at 2 m altitude. In addition, information on the soil type is required. As the model was initially conceived as a tool for the estimation of bio-energy potentials, the maize crops only contain silage maize (no food maize). Furthermore the three varieties do not consist of single breeds but represent an average of several early, medium or late breeds respectively.

As a rather robust approach the model leaves out some aspects that might well be of great importance for a future yield development. Results in this study should thus be read as what would happen if nothing but the climatic input variables would change. These neglected aspects include any technological advances, including any changes in farm management. Irrigation was not included in the modeling, whether for current nor future yields, even if there do exist some areas today that are under irrigation. The sowing date was always the 115th day of the year and was not changed throughout the century. No extra fertilization was included and soil water content expected to be at

BGD

11, 9103–9134, 2014

Maize yield development in Lower Saxony

J. F. Degener and
M. Kappas

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Maize yield development in Lower Saxony

J. F. Degener and
M. Kappas

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



25 100% at the beginning of the growing season. No effects of a prior crop on a specific site are taken into account. The model either stopped on day 300 of a given year or when full maturity was attained, depending on what happened first.

At the time of the actual modeling the BioSTAR model could be understood as being in beta stage development. All crops had been validated, however some minor changes in the model engine and more heavy changes in the programs layout have happened since. No version number was in use when the model was applied.

5 2.3 Soil data

The soil data used in this study is part of the official digital soil survey map of LS in a resolution of 1 : 50000, called BÜK50 (Boess et al., 2004). This map was intersected with data from the CORINE land-use classification of 2005 for Lower Saxony to extract sites that are used for agricultural purposes only. The result between the intersected
10 soil and land-use map was a data-set of 91 014 sites with each used as a unique modeling area. The soil map contained codified information on the soil type and its thickness that were translated into the format required by BioSTAR. Fifteen 10 cm soil levels had to be identified, each containing the information on prevalent soil type with a 16th level representing everything below the initial 1.5 m. The crop model uses these
15 information solely for the calculation of soil water content and flows.

2.4 Climate model and data

The climate data was derived from the regional climate model WETTREG, a German portmanteau word translating into “weather condition based regional model”. The model uses a statistical downscaling method where large scale atmospheric patterns are brought into a statistical relationship with local climate station data (Enke and Spekat, 1997). The initial link is created by using known measured data at these stations and globally gridded reanalysis data, with both ERA40 and NCEP/NCAR data using a k-means cluster approach. This link is then reestablished through GCM de-
20

speed in a height of 2 m above grass. Typical wind-speed ranges from 5–6 m s⁻¹ at the coast to around 1–2 m s⁻¹ in the south of LS. To present knowledge no significant change in the wind speed pattern is anticipated for the future (NMUEK, 2012), hence the data was applied without further changes.

Relative Air Humidity was calculated backwards from the WETTREG2010 data on evapotranspiration, as this was derived through the Penman/Monteith approach.

All data was then intersected with the soil sites using the respective variables mean value.

2.5 Statistics

To account for extreme or unrealistic outliers, a two-way approach was devised for the original resulting data-set. At first all sites with a biomass yield of 0 g m⁻² were excluded. This typically amounted to 706 sites that contain only bedrock in their soil levels. In a second step all data below the 0.1 and above the 99.9 percentile were excluded, as values close to zero or unreasonably large yields were present. This proved to well eliminate outliers while preserving as much data as possible.

Basic statistics in this study include standard deviation, coefficient of variability (cov), linear regression models and the coefficient of determination as described in Schönwiese (2006). The time series could well be described using linear regression models, however tests with exponential, logarithmic, second and third order polynomial and potential models did show about equal results.

The data was further explicitly tested for trends using a robust trend/noise ratio (t/n), where the difference in yield from the years 2099 and 2001 was divided through the time series' standard deviation. A significant trend is assumed at a ratio of 1.96 or above, representing the $\alpha = 0.05$ level. As this test is often considered to be relatively weak, the non-parametric Mann–Kendall-Test (MK) was applied as well. A further advantage of MK is its ability to detect non-linear trends. Most statistics were applied for the time series of 2001–2030, 2001–2050 and 2001–2099.

BGD

11, 9103–9134, 2014

Maize yield development in Lower Saxony

J. F. Degener and
M. Kappas

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



a near (2011–2040), middle (2041–2070) and long-term (2071–2100) climatic development.

25 There are no areas at any time that do show a stagnant or even decreasing temperature development. However, warming in spring is always below the annual average while the winter months are always above. Fall temperatures are slightly below annual average and summer months above, though both deviate less from the mean than spring and winter seasons do. The mean temperature increase is 0.95 °C for near, 2.30 °C for middle and 3.40 °C for long-term scenarios. The development is relatively
5 uniform throughout LS with a slightly stronger (but still less than 0.5 °C difference) development to the south-east.

The precipitation development is different in terms of being positive or negative depending on time and space. If only annual means are considered, almost no change in precipitation can be detected, although a moderate decline is visible. It however
10 becomes increasingly obvious that the winter and summer seasons are drifting into opposite directions. While in the near future all seasonal differences remain in a window of more or less $\pm 10\%$, these changes drastically amplify towards the end. The mean decline in precipitation is around -25% in the long-term perspective with some areas at a nearly -50% decrease. Winter increases are also substantial but at around
15 15% towards the end of the century they cannot fully counterbalance the summery losses.

In summary, all deviations from today's values will increase with passing time, fostering a local development towards a more winter rain climate that features increasingly hot and dry summers and mild wet winters.

20 4 Results

The results in this study will describe the *change* in biomass yields during the 21st century. Changes are relative to the mean yields of the decade 2001–2010 as a representation of the present time.

Maize yield development in Lower Saxony

J. F. Degener and
M. Kappas

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



4.1 Mean yield development

25 The results in this section give the average yield development of all modeled sites. As can be seen in Fig. 3 all three varieties visually show a positive yield development throughout the century. This is further underpinned by the actual biomass yields after 2060, where the average yield per decade is always higher than for the reference period. Not as evident is the shared pattern of the decadic coefficient of variability. All
5 three varieties have their lowest value in the present (3–4%) with an increase (except for the comparably low variability between 2031–2040) towards mid-century (above 8%) and a slight decline towards 6–7% at the end of the century. Actual yields will therefore vary more widely around the decadic mean at mid-century.

Apart from these shared aspects there are also obvious differences in the overall development. The following description will thus cover each variety on its own.

10 The *early* variety shows an $R^2 = 0.24$ and an average increase in yields of 0.12% p.a. throughout the century. This trend could be slightly better explained through a polynomial model of second or third order with $R^2 = 0.27$, however no big advantage would be expected from such an approach. The trend/noise (t/n) ratio 1.68 shows no significant trend for $\alpha = 0.95$ but would for $\alpha = 0.9$. Mann–Kendall delivers a more unambiguous result with $p < 0.001$ over the century. It is therefore assumed that a significant
15 trend exists throughout the entire time-period.

This trend can basically be split into two parts: the period 2001–2050 has an $R^2 = 0.003$ in a linear regression model with an average yield development of $\pm 0\%$ p.a. The period 2051–2099 has an $R^2 = 0.11$ with an average yield increase of 0.2% p.a. The lack of a trend in the first half of the century is confirmed by its t/n ratio of 0.5 and $p = 0.39$ for MK. For the period 2001–2030 there even seems to appear a slight
20 negative development, with a t/n ratio of -0.63 and a $p = 0.08$ for MK that is however not recognized as being significant.

25 All in all it seems clear that a change in biomass yields is expected to happen, however only after the mid of the century and especially after 2070 when there is only

BGD

11, 9103–9134, 2014

Maize yield development in Lower Saxony

J. F. Degener and
M. Kappas

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



This is evident by just looking at the graph as well as in the numbers of the linear regression model with an $R^2 = 0.65$ and an average yield increase of 0.31 % p.a. A change to another regression model does not show any different results. Also the t/n ratio (2.74) and Mann–Kendall ($p < 0.001$) are more explicit in determining significance than for the other varieties.

However clear, this trend is not entirely constant. While the t/n ratio (2.31) and MK ($p < 0.001$) already show a highly significant trend towards 2050 the data seems to have a break around the year 2030. There appears to be no trend for the period 2001–2030 as a t/n ratio of 0.65 and $p = 0.35$ for MK suggest. A linear regression for 2001–2030 shows an $R^2 = 0.09$ and a mean yield increase of 0.11 % p.a., a non-significant but positive trend, whereas 2031–2099 has a $R^2 = 0.37$ and an increase of 0.26 % p.a.

The late variety has thus the most positive development throughout the century. The mean yields in the last three decades increase about at least 10 % compared to today, with a mean of 25 % and a maximum of 36 %. The same holds true for a production comparison of both half-centuries, as the first 50 years would only contribute 47.1 % to a potential overall production.

4.2 Regional yield development

As Fig. 4 indicates there are also certain differences in the regional distribution of potential yields increases or decreases. The late variety does clearly have the most uniform development as it is positive for almost all times and sites. The share of sites with a positive development lies around 53 % for the early variety in the period and increases to 87 %, 96 % and 95 % towards the end of the century. In a similar fashion the medium variety starts out at very low share of 16 % positive sites, increasing to 49 %, 88 % and 83 %.

If the coefficient of variation is calculated regarding all sites and years from the respective periods, all three varieties show an increasing *cov* with progressing time. With 9.3 % (2021–2040) to 10.7 % (2081–2099) the late variety does have the least variabil-

BGD

11, 9103–9134, 2014

Maize yield development in Lower Saxony

J. F. Degener and
M. Kappas

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



with higher p values increased. Even the late variety now did have around 1500 sites exceeding this threshold, with early and medium variety on about 7000 sites. Roughly a third of these exceptional sites have p values > 0.05 . In conclusion, the models are slightly better for the description of the long-term development than for the first half of the century.

For 2001–2099 two main influence variables are detected. Summer precipitation shows a strong positive correlation for all three varieties. As the amount of rain is expected to drastically decline throughout the century, this seems to be the main factor to limit future maize yields. On the other hand, atmospheric CO_2 concentrations have a comparable positive correlation and are thus possibly the main agent for a positive yield development. The amount of spring precipitation seems to be of higher importance for medium ($p = 0.09$) and especially early ($p = 0.04$) variety. Both do also show a negative connection with the rising summer temperatures (early $p = 0.09$, medium $p = 0.08$), at least to some degree. The late variety shows basically similar dependencies, however weaker. Instead spring temperatures ($p < 0.01$) seem to be of much higher importance than for the other two varieties.

For 2001–2050 these indicators change only slightly. Still summer precipitation and CO_2 concentrations remain the determining variables ($p < 0.001$). The late variety still shows some dependency towards spring temperatures ($p = 0.09$). For all three varieties fall temperatures seem to be of higher importance in the first 50 years ($0.1 > p > 0.05$), whereas summer temperatures and spring precipitation have no apparent influence.

That these multivariate models are not entirely perfect becomes evident when for example winter precipitation and late maize are considered for 2001–2099. While not being *highly* significant a certain connection between both variables is suggested. However, as winter months include December, January and February, when no maize is grown, this also seems to be highly improbable. While the statistical model was believed to be reasonably good in determining the relative influence of each variable, there was a need to exclude variables that are not necessarily important.

BGD

11, 9103–9134, 2014

Maize yield development in Lower Saxony

J. F. Degener and
M. Kappas

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Maize yield development in Lower Saxony

J. F. Degener and
M. Kappas

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Therefore the next step was to identify the one linear multivariate regression model for each site that best describes its yield development, as shown on the right side in Fig. 5. These results were largely in accordance with the results from the models with eleven input variables. Models in the following approach are numbered from R1 to R2047. R is here short for *Run* while the number indicates the variable combination. Higher numbers relate to more input variables. The relevant numbers are explained further in Fig. 5.

For 2001–2099 the model best describing the yield development of all varieties was one containing only summer precipitation and CO₂ concentrations (R37). 92 % of the early variety, 80 % of the medium variety and 85 % of the late variety sites had this as the optimal model. The remaining sites of the early variety were best described by a model *only* containing CO₂ concentrations (R10). The same is partially true for the late variety, as 3 % of the models show their best results when only including CO₂ (R10), however models that only used spring temperatures (R6) accounted for the remaining 12 %. This connection to spring temperatures was also identified in the models featuring all variables. The two runs R6 and R10 have a combined share of about 10 % of the medium variety's remaining sites, while the other remaining 10 % are a combination of summer precipitation, summer temperatures and CO₂ concentration (R171).

For the period 2001–2050 the varieties did show a more differentiated picture. The late variety did still have R37 as the dominant model on 90 % of its sites. 5 % were made up of R6 and another 5 % of other not further distinguished models. The early variety had R37 on just 58 % of its sites, 5 % showing R171 and almost the entire rest of 34 % from R2 with summer precipitation only. The medium variety had only have 24 % comprising of R37, 6 % of R171 and a dominating 66 % of R2.

5 Discussion

The results from this study basically agree with the findings of e.g. Southworth et al. (2000) in that the choice of variety will have a critical effect on how maize yields will develop under a future climate. It even agrees to the point that late varieties will show the most positive development which can be quite substantial with > 25% in Lower Saxony towards the end of the century.

One reason for this beneficial development of late maize is clearly the fact that today's temperature sums in LS are not suitable for a full completion of its growing cycle. Temperature sums from 20 April to 15 October (minus 6°C temperature basis) vary today around 1500°C in LS and are therefore perfect for medium varieties but below optimum for late varieties. It seems that around 2030, when temperature sums have increased by about 100°C, the late variety can fully benefit from these temperatures. That the late variety disproportionately benefits from the generally rising temperatures is further supported by statistical analysis, as the late variety is the only one to show a substantial positive correlation to rising spring temperatures.

The future climatic conditions are however not entirely beneficial for the growth of late maize varieties. The main limiting factor, for all varieties, is the decline in summer precipitation. However, the time spent within these dry months in relation to the total growing time is shorter for the late variety as e.g. the medium one. Thus the late variety can use the moister spring or fall conditions for a successful growth. Similarly the early variety profits from the spring conditions while the medium variety would need more water during the summer months of which August will be the driest.

An adaptation of sowing date could mitigate these negative effects to some degree. Some testing on single sites however suggests that the general yield development series of *late* > *early* > *medium* variety is not changed, though the absolute difference might change. The influence of the sowing date on the results of this work are currently under evaluation.

BGD

11, 9103–9134, 2014

Maize yield development in Lower Saxony

J. F. Degener and
M. Kappas

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Maize yield development in Lower Saxony

J. F. Degener and
M. Kappas

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



In a related matter, critical development stages of maize, e.g. during flowering, where water shortage disproportionately restrains plant growth (Ehlers, 1996), were not sufficiently implemented in the used version of the crop model. This has changed in the current version and is under evaluation as well. While a shift in sowing dates is expected to be generally beneficial for maize yields, this increase in sensitivity is believed to have a rather detrimental effect.

While increasing temperatures are generally good for the late variety, temperatures during the growing season should not exceed an optimum growing temperatures of 25–30°C over longer time periods, as this would inhibit photosynthesis rates (Endlicher, 2007). However, heat days will increase in LS throughout the century. The effects of this are visible in the multivariate regression models with an increasing negative impact of summer temperatures on yields.

In reality this effect is expected to be even worse. The main reason is the use of a relatively smooth time-series. This is firstly caused by the statistical nature of the climate model. The usage of mean values from 10 climate model runs tends to eliminate extreme values. Secondly, the results of this downscaled climate data-set were 10-day-values that were further combined into monthly averages. Temperature peaks were thus eliminated within the monthly means. The same is true for the monthly values of precipitation, as BioSTAR simply assumes that the monthly value is distributed evenly over each day of the month. This is clearly not the case in nature where a steady flow of water would be optimal for the plants water supply. It will make some difference if 20 mm precipitates in one day followed by nine dry days or if 2 mm for each of the ten days is assumed. While a shortage in precipitation might be worse than elevated temperatures, it can be mitigated relatively easy by irrigation while the latter can hardly be opposed. As some areas in LS are already today under irrigation it would be interesting to estimate probable changes in irrigation practices, meaning an estimation of the amount of water needed for optimal growth and taking the actually available amount of water into account.

Maize yield development in Lower Saxony

J. F. Degener and
M. Kappas

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



That even relatively smooth precipitation series lead to varying yields becomes evident when the observed decadal variability of the yields is compared with the variability of precipitation. More precisely the yields have their greatest variability in the decade 2051–2060, when summer precipitation has the greatest variability too. The same link can be found in the decade 2021–2030 but not for 2031–2050, as both decades have either high yield or high precipitation variability but not both. The underlying cause for the change in yield variability therefore seems to be more complex than a single dependency on summer precipitation.

There further exist some climatic circumstances that might negatively affect yield development that were not accounted for in this study. One is tropospheric ozone, as 30 ppb are sufficient to induce ozone intoxication in plants (Long et al., 2006). Since 1950 the concentration of tropospheric ozone has nearly doubled. Studies suggest that maize yields might be 2–5.5 % higher today if this rise would not have happened (Avnery et al., 2011). It is however debatable if tropospheric ozone concentrations will further increase, at least in Europe, due to anthropogenic emission as CMIP 5 runs suggest (Fiore et al., 2012).

A greater potential risk arises through common or invasive pests. Complicated interactions and feedbacks between climate, crop and pests make concise predictions difficult (Schaller and Weigel, 2007). However, as Fröhlich (2010) points out, there is no expectation at all that the climatic change will lead to a reduction in infestation of any pest. In how far new cropping techniques or breeds will be able to counteract such problems is beyond the scope of this study.

The general outlook considering the climatic changes could thus be interpreted as quite severe. However, the results from this study suggest the contrary with rising yields towards the century's end. The only variable contributing significantly towards rising yields is atmospheric CO₂.

Maize as a C4 crop is not expected to profit from rising CO₂ through an elevated photosynthesis rate (Lambers et al., 2008). However an increased water use efficiency is expected in C3 as well as C4 plants. This effect is accounted for by the crop model,

Maize yield development in Lower Saxony

J. F. Degener and
M. Kappas

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



resulting in a relatively linear decrease in the amount of water that is needed to produce the same amount of plant matter. The reduction is comparable for all three varieties and ranges between 25–30 %.

In an environment where water is getting increasingly scarce this is a desirable development. It appears that the negative impacts of summer temperature and precipitation are stronger until mid-century, especially for medium or early variety. The positive influence of CO₂ steadily increases to a point where the positive effects prevail and yields are rising.

That water saving through increased CO₂ concentrations can have such a strong effect is also pointed out by Taube and Herrmann (2009), where Grasslands profit from a rise even under increasing drought stress during summer months. This would be in line with Morgan et al. (2004) who are emphasizing the importance of water saving through increased CO₂ in contrast to a direct fertilization effect.

CO₂ might still not be solely responsible for the rising yields. It undoubtedly plays a major role in doing so, however other factors that have not been included in the process evaluation might contribute as well. Mera et al. (2006) included the effect of solar radiation in their research and found a non-linear contribution to yield development, however not as prominent as changes in precipitation or water availability.

6 Conclusions

As could be shown, the changing climate will have a predominantly positive effect on the yield development of maize and its varieties in Lower Saxony. A real positive development is however not expected to set in before the second half of the 21st century.

The first half will be stagnant in yields for the early variety. In the last decades of the century the yields will on average increase about 9 %. The medium variety even shows a negative development in the first half that is later reversed. Towards the century's end the yields then increase about 5 % in comparison to today's yields. The late variety has the all-out best yield development, with an average increase of 25 % for 2071–

Maize yield development in Lower Saxony

J. F. Degener and
M. Kappas

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2099 and a strong positive trend beginning already around 2030. In addition to this above-average rise the yields itself are higher so that a transition of local agricultural practices towards the late variety is conceivable. The yield development of all yields is accompanied by an increase in yield variability during mid-century that seems to partially follow precipitation patterns.

Thus the development will generally be positive in the long run, though the path for each variety diverges. As the varieties react in different ways to the changing annual pattern of temperature and precipitation, the results do indicate that the consideration of different varieties might also change the outcome of studies at different study sites.

At any rate the few other existing studies are hinting towards the same result. Varieties with longer or shorter growing periods will have an advantage in areas where medium varieties are predominantly grown today.

Besides, for Lower Saxony or Germany in general, a decline in summer precipitation is not seen as an insurmountable obstacle for local agriculture, as there is no necessity for irrigation on most sites today and present water reserves would allow an expansion of irrigated areas at least to some degree. Intensive groundwater management will a basis for this, as increasing winter precipitation could cover the water extraction during the summer months. New breeds and cropping techniques will also aid to counteract the negative effects of climate change, including the expansion of pests or hitherto unknown effects that might arise.

In conclusion the maize yields in Lower Saxony will not suffer from long lasting declines but will have a generally positive outlook over the course of the 21st century.

Acknowledgements. We acknowledge support by the Open Access Publication Funds of the Göttingen University. We would like to thank the LBEG for the provision of the climate data-set. This Open Access Publication is funded by the University of Göttingen.

Maize yield development in Lower Saxony

J. F. Degener and
M. Kappas

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- 25 Fiore, A. M., Naik, V., Spracklen, D. V., Steiner, A., Unger, N., Prather, M., Bergmann, D., Cameron-Smith, P. J., Cionni, I., Collins, W. J., Dalsøren, S., Eyring, V., Folberth, G. A., Ginoux, P., Horowitz, L. W., Josse, B., Lamarque, J.-F., MacKenzie, I. A., Nagashima, T., O'Connor, F. M., Righi, M., Rumbold, S. T., Shindell, D. T., Skeie, R. B., Sudo, K., Szopa, S., Takemura, T., and Zeng, G.: Global air quality and climate, *Chem. Soc. Rev.*, 41, 6663–6683, 2012.
- 30 Fröhlich, M.: Klimawandel und Landwirtschaft: Auswirkungen der globalen Erwärmung auf die Entwicklung der Pflanzenproduktion in Nordrhein-Westfalen, Ministerium für Klimaschutz, Umwelt, Landwirtschaft, Natur- und Verbraucherschutz des Landes Nordrhein-Westfalen, 2010.
- Heunisch, C., Caspers, G., Elbracht, J., Langer, A., Röhling, H.-G., Schwarz, C., and Streif, H.: Erdgeschichte von Niedersachsen: Geologie und Landschaftsentwicklung, Landesamt für Bergbau, Energie und Geologie (LBEG), Hannover, Geoberichte 6, 2007.
- 5 Hoehner, G.: Entwicklung Energiepflanzenanbau und Biogas in Niedersachsen, available at: www.ifeu.org/landwirtschaft/pdf/6_Hoehner_Maisanbau_Nieders.pdf, (11 March 2014), 2007.
- Hunter, J. D.: Matplotlib: a 2D graphics environment, *Comput. Sci. Eng.*, 9, 90–95, 2007.
- Jacob, D., Göttel, H., Kotlarski, S., Lorenz, P., and Sieck, K.: Klimaauswirkungen und Anpassung in Deutschland: Phase 1: Erstellung regionaler Klimaszenarien für Deutschland, Im Auftrag des Umweltbundesamtes, Dessau-Roßlau, Climate Change (11/2008), 2008.
- 10 Jones, E., Oliphant, T., Peterson, P., et al.: SciPy: open source scientific tools for Python, (17.07.2013), 2001.
- Kreienkamp, F., Spekat, A., and Enke, W.: Weiterentwicklung von WETTREG bezüglich neuer Wetterlagen, 2010.
- 15 Kwabiah, A.: Growth and yield of sweet corn (*Zea mays* L.) cultivars in response to planting date and plastic mulch in a short-season environment, *Sci. Hortic.*, 102, 147–166, 2004.
- Lambers, H., Pons, T. L., and Chapin, F. S.: *Plant Physiological Ecology*, 2. edn., Springer, New York, 2008.
- Liu, Z., Hubbard, K. G., Lin, X., and Yang, X.: Negative effects of climate warming on maize yield are reversed by the changing of sowing date and cultivar selection in Northeast China, *Glob. Change Biol.*, 19, 3481–3492, 2013.
- 20

Maize yield development in Lower Saxony

J. F. Degener and
M. Kappas

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Long, S. P., Ainsworth, E. A., Leakey, A. D. B., Nösberger, J., and Ort, D. R.: Food for thought: lower-than-expected crop yield stimulation with rising CO₂ concentrations, *Science*, 312, 1918–1921, 2006.

25 LSN: Statistical Data of the Statistical Office of Lower Saxony, available at: <http://www.nls.niedersachsen.de/> (last access: 14 March 2014), 2014.

LWK: Homepage of the Chamber of Agriculture of Lower Saxony, available at: <http://www.lwk-niedersachsen.de> (last access: 10 March 2014), 2014a.

30 LWK: Energiepflanzenfeldtage: Es muss nicht immer Mais sein, available at: <http://www.lwk-niedersachsen.de/index.cfm/portal/2/nav/74/article/15172.html> (last access: 10 March 2014), 2014b.

Mera, R., Niyogi, D., Buol, G., Wilkerson, G., and Semazzi, F.: Potential individual versus simultaneous climate change effects on soybean (C₃) and maize (C₄) crops: an agrotechnology model based study, *Global Planet. Change*, 54, 163–182, 2006.

5 Meyer, U., Hüther, L., Manderscheid, R., Weigel, H.-J., Lohölter, M., and Schenderlein, A.: Nutritional value of maize 2050, in: *Optimierung des Futterwertes von Mais und Maisprodukten*, vol. 331 of *Landbauforschung/Sonderheft*, edited by: Schwarz, F. J., VTI, Braunschweig, 107–114, 2009.

10 Meza, F. J., Silva, D., and Vigil, H.: Climate change impacts on irrigated maize in Mediterranean climates: Evaluation of double cropping as an emerging adaptation alternative, *Agr. Syst.*, 98, 21–30, 2008.

Mirschel, W., Wenkel, K.-O., Wieland, R., Köstner, B., Albert, E., and Luzi, K.: Auswirkungen des Klimawandels auf die Ertragsleistung ausgewählter landwirtschaftlicher Fruchtarten im Freistaat Sachsen, ZALF, Müncheberg, 2008.

15 Mirschel, W., Wieland, R., Wenkel, K.-O., Guddat, C., Michel, H., Luzi, K., and Groth, K.: Regionaldifferenzierte Abschätzung der Auswirkungen des Klimawandels auf die Erträge von wichtigen Fruchtarten im Freistaat Thüringen mittels Ertragssimulation mit YIELDSTAT, ZALF, Müncheberg, 2012.

20 Morgan, J. A., Pataki, D. E., Körner, C., Clark, H., Del Grosso, S. J., Grünzweig, J. M., Knapp, A. K., Mosier, A. R., Newton, P. C. D., Niklaus, P. A., Nippert, J. B., Nowak, R. S., Parton, W. J., Polley, H. W., and Shaw, M. R.: Water relations in grassland and desert ecosystems exposed to elevated atmospheric CO₂, *Oecologia*, 140, 11–25, 2004.

Maize yield development in Lower Saxony

J. F. Degener and
M. Kappas

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Müller, U., Engel, N., Heidt, L., Schäfer, W., Kunkel, R., Wendland, F., Röhm, H., and Elbracht, J.: Klimawandel und Bodenwasserhaushalt, landesamt für Bergbau, Energie und Geologie (LBEG), Hannover, Geoberichte 20, 2012.

25 NMELVL: Maisanbau: Mehr Vielfalt durch Alternativen und Blühstreifen, available at: <http://www.ml.niedersachsen.de/download/78012>, 2012.

NMUEK: Empfehlung für eine niedersächsische Klimaanpassungsstrategie, 2012.

Pandas: pandas: Python Data Analysis Library, (17.07.2013), 2012.

30 Roeckner, E.: ensembles stream2 echam5c-mpi-om sra1b run1: World Data Center for Climate, 2009.

Schaller, M. and Weigel, H. J.: Analyse des Sachstands zu Auswirkungen von Klimaveränderungen auf die deutsche Landwirtschaft und Maßnahmen zur Anpassung, 2007.

Schönwiese, C.-D.: Praktische Statistik für Meteorologen und Geowissenschaftler, 4. edn., Borntraeger, Berlin (u.a.), 2006.

Seedorf, H. H. and Meyer, H.-H.: Landeskunde Niedersachsen: Historische Grundlagen und naturräumliche Ausstattung, Wachholtz, Neumünster, 1992.

5 Southworth, J., Randolph, J., Habeck, M., Doering, O., Pfeifer, R., Rao, D., and Johnston, J.: Consequences of future climate change and changing climate variability on maize yields in the midwestern United States, *Agr. Ecosyst. Environ.*, 82, 139–158, 2000.

Supit, I., van Diepen, C., Wit, A. d., Wolf, J., Kabat, P., Baruth, B., and Ludwig, F.: Assessing climate change effects on European crop yields using the Crop Growth Monitoring System and a weather generator, *Agr. Forest Meteorol.*, 164, 96–111, 2012.

10 Taube, F. and Herrmann, A.: Relative benefit of maize and grass under conditions of climatic change, in: Optimierung des Futterwertes von Mais und Maisprodukten, vol. 331 of *Landbauforschung/Sonderheft*, edited by: Schwarz, F. J., VTI, Braunschweig, 115–126, 2009.

USF: Klimawandel und Landwirtschaft in Hessen: Mögliche Auswirkungen des Klimawandels auf landwirtschaftliche Erträge, *iNKLIM Baustein 2*, Universität Kassel, 2005.

715 WMO: Guide to Climatological Practices: 2011 edition, 2011.

Wolf, J. and van Diepen, C.: Effects of climate change on silage maize production potential in the European community, *Agr. Forest Meteorol.*, 71, 33–60, 1994.

720 Wolf, J. and van Diepen, C.: Effects of climate change on grain maize yield potential in the European community, *Climatic Change*, 29, 299–331, 1995.




Figure 1. Average Maize production in 2010 as fresh-matter in 1000 metric tonnes by district (LSN, 2014).

BGD

11, 9103–9134, 2014

Maize yield development in Lower Saxony

J. F. Degener and
M. Kappas

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Maize yield development in Lower Saxony

J. F. Degener and M. Kappas




Figure 2. Indicators of a regional climatic change. Box-Whisker-Plots of temperature (left) and precipitation (right) changes for three different periods in relation to 1971–2000 by season. The data is taken from the WETTREG2010 data-set and represents the mean over Lower Saxony.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



BGD

11, 9103–9134, 2014

Maize yield development in Lower Saxony


J. F. Degener and
M. Kappas

Figure 3. Maize yield development in relation to the mean of 2001–2010 for the three varieties. The black lines and data indicate the linear trend over the century. Dotted lines represent two linear trends during the century with a breaking point at 2050 (early) or 2030 (medium and late variety).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Maize yield development in Lower Saxony

J. F. Degener and
M. Kappas




Figure 4. Local percentage variation of biomass yields of early, medium and late maize varieties compared to 2001–2010.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏪](#)
[⏩](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Maize yield development in Lower Saxony

J. F. Degener and
M. Kappas




Figure 5. Left side: output of 9 variables from the linear multivariate regression analysis for 2001–2099 (top) and 2001–2050 (bottom) – Right side: relative share of different linear (multivariate) models to the total number of models R_x represents the number of the 2047 possible runs through variable combination – P_x is precipitation by respective season, T_x for temperature.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

