

1

2 **Methane related changes in prokaryotes along geochemical profiles in sediments**
3 **of Lake Kinneret (Israel)**

4 **I. Bar Or¹, E. Ben-Dov^{2,3}, A. Kushmaro^{*2,5,6}, W. Eckert⁴ and O. Sivan¹**

5

6 ¹Department of Geological and Environmental Sciences, Ben-Gurion University of the Negev,
7 Be'er- Sheva, P.O. Box 653, 8410501, Israel.

8 ²Avram and Stella Goldstein-Goren Department of Biotechnology Engineering, Ben-Gurion
9 University of the Negev, Be'er- Sheva, P.O. Box 653, 8410501, Israel.

10 ³Department of Life Sciences, Achva Academic College, Achva, M.P. Shikmim 79800, Israel.

11 ⁴ Israel Oceanographic and Limnological Research, The Yigal Allon Kinneret Limnological
12 Laboratory, PO Box 447, 14950 Migdal, Israel.

13 ⁵National Institute for Biotechnology in the Negev, Ben-Gurion University of the Negev, P.O. Box
14 653, Be'er-Sheva 8410501, Israel

15 ⁶School of Materials Science and Engineering, Nanyang Technological University, Singapore

16

17 * Corresponding author: Mailing address: Department of Biotechnology Engineering, Ben-Gurion
18 University of the Negev, P.O. Box 653, Be'er-Sheva 8410501, Israel. Phone: 972-8-6479.024. Fax:
19 972-8-6472.983. E-mail: arielkus@bgu.ac.il.

20

21 **Keywords:** anaerobic methane oxidation, iron reduction, archaeal classification, bacterial
22 classification, Lake Kinneret.

23

24 **Abstract.** Microbial methane oxidation is the primary control on the emission of the greenhouse
25 gas methane into the atmosphere. In terrestrial environments, aerobic methanotrophic bacteria are
26 largely responsible for this process. In marine sediments a coupling of anaerobic oxidation of
27 methane (AOM) with sulfate reduction, often carried out by a consortium of anaerobic
28 methanotrophic archaea (ANME) and sulfate reducing bacteria, consumes almost all methane
29 produced within marine sediments. Motivated by recent evidence for AOM with iron(III) in Lake
30 Kinneret sediments, the goal of the present study was to link the geochemical gradients in the lake
31 porewater to the microbial community structure. Screening of archaeal 16S rRNA gene sequences
32 revealed a shift from hydrogenotrophic to acetoclastic methanogens with depth. The observed
33 changes in microbial community structure suggest possible direct and indirect mechanisms for the

34 AOM coupled to iron reduction in deep sediments. Members of the *Burkholderiales* and
35 *Nitrospirales* orders increased with depth, suggesting their involvement in iron reduction together
36 with *Geobacter* genus and "reverse methanogenesis". An indirect mechanism through sulfate and
37 ANMEs seems less probable to explain the data due to the absence of ANME sequences. This is
38 despite the abundant sequences related to sulfate reducing bacteria (*Deltaproteobacteria*) together
39 with the occurrence of *dsrAB* in the deep sediment that could indicate the production of sulfate
40 (disproportionation) from S^0 for sulfate-driven AOM. The presence of the functional gene *pmoA* in
41 the deep anoxic sediment together with sequences related to *Methylococcales* suggests the existence
42 of a second unexpected indirect pathway - aerobic methane oxidation pathway in an anaerobic
43 environment.

44

45

46 1. Introduction

47 Chemical profiles in the porewater of aquatic sediments reflect the sequence of microbially
48 mediated redox reactions that are driven by the availability of both, electron donors and of suitable
49 electron acceptors. The latter are depleted in the order of decreasing chemical potential, beginning
50 with oxygen and proceeding through nitrate, manganese and iron oxides, and then sulfate. Below
51 the main zone of sulfate reduction, the fermentation of organic carbon leads to the formation of
52 methane (CH₄) by the process of methanogenesis (Froelich et al., 1979).

53 The produced methane is isotopically depleted in ¹³C, with values of ~-50 to -110‰ (Schoell,
54 1988), and the residual dissolved inorganic carbon (DIC) pool is enriched by an isotopic
55 fractionation factor (ε) of 50 to 70‰ (e.g., (Borowski et al., 2000; Whiticar, 1999)). When the
56 produced methane diffuses into a zone with a suitable electron acceptor, it can be consumed by
57 microbial oxidation (methanotrophy), the main process by which the important greenhouse methane
58 is prevented from escaping into the atmosphere. While in the terrestrial environment, aerobic
59 methanotrophy is the dominant process (Chistoserdova et al., 2005), in anaerobic marine sediments
60 archaea are found to consume the majority of upward diffusing methane coupled to sulfate
61 reduction (Knittel and Boetius, 2009; Thauer, 2010; Valentine, 2002).

62 Evidence from lipids and from fluorescence *in situ* hybridization (FISH) showed that a
63 consortium of archaea and sulfate reducing bacteria are involved in this anaerobic methane
64 oxidation (AOM) (Boetius et al., 2000; Hinrichs et al., 1999; Orphan et al., 2001). To date, three
65 groups of anaerobic methanotrophic archaea (ANME), named ANME-1, ANME-2, and ANME-3,
66 are known to perform sulfate driven AOM (Niemann et al., 2006; Orphan et al., 2002). However
67 Milucka et al. (2012) demonstrated AOM mediated solely by archaea, where the archaea was shown
68 to oxidize the methane and reduce the sulfate to elemental sulfur. Disproportionating bacteria, also
69 involved in this mechanism, oxidize and reduce this elemental sulfur to sulfate and sulfide,
70 respectively. The carbon isotopic fractionation factor (ε) for this methanotrophic process was shown
71 to be in the range of 4-30‰ (Kinnaman et al., 2007; Whiticar, 1999). Enrichment cultures of
72 ANME from different environments showed a carbon isotopic fractionation of 12-39‰ (Holler et
73 al., 2009).

74 Other electron acceptors were recently shown to drive AOM. Nitrite driven AOM by oxygenic
75 bacteria was observed in two different freshwater ecosystems in Netherlands (Ettwig et al., 2009;
76 Raghoebarsing et al., 2006) and also in peatlands (Zhu et al., 2012). Beal et al. (Beal et al., 2009)
77 showed the potential of manganese and iron-driven AOM in marine sediments, and Egger et al.,
78 (2014) showed it in brackish coastal sediments. In our recent study (Sivan et al., 2011), we provided
79 *in situ* geochemical evidence for AOM coupled to microbial iron reduction below the main

80 methanogenesis zone in Lake Kinneret (LK) sediments, where dissolved sulfate and nitrate are
81 absent. However, the mechanism that is responsible for this process was not investigated. The goal
82 of the present study was to explore the possible microbial pathways that may be involved in the iron
83 driven AOM in LK sediments. This is by combining chemical and isotope analyses of porewater
84 samples along a depth profile from LK sediments with molecular biological techniques. By using
85 specific functional genes related to dissimilatory sulfate reductase (*dsr*) for sulfate reducers (Klein
86 et al., 2001), methyl-coenzyme M reductase (*mcr*) for methanogens and anaerobic methanotrophs
87 (Hallam et al., 2003) and particulate methane monooxygenase (*pmmo*) for aerobic methanotrophs
88 (McDonald et al., 2008) we pinpoint the most likely candidate for this relatively unknown process.

89 **1.1. Study site**

90 Located in northern Israel, Lake Kinneret (LK, Fig.1) is a warm monomictic subtropical lake.
91 Typical concentrations of major electron acceptors in the water column during the mixed period are
92 35–50 μM nitrate and 600 μM sulfate (Adler et al., 2011; Nishri et al., 2000; Serruya et al., 1974).
93 In the spring, the newly formed epilimnion is characterized by increasing temperatures and
94 enhanced phytoplankton development, while in the hypolimnion heterotrophic microorganisms
95 gradually deplete oxygen and then nitrate. Organic matter degradation by bacterial iron and
96 manganese reduction takes place below the thermocline in the summer, and in the end of the
97 stratification period, sulfate reduction starts in the bottom water. In the upper part of the sediment,
98 sulfate reduction is the dominant microbial process year round, and below depth of 5 cm this
99 process is mostly replaced by methanogenesis (Adler et al., 2011; Eckert and Conrad, 2007). Total
100 iron content ($\text{Fe}_{(\text{tot})}$) in the sediment increases with depth till 18 cm and then remains uniform
101 around 550 $\mu\text{mol/g}$ Dry Weight (Eckert, 2000). Mn(II) concentration is about 13 $\mu\text{mol/g}$ Dry
102 Weight in the sediment (Serruya, 1971). The total carbon (C_{tot}) in the solid phase shows a decrease
103 from 14 $\mu\text{mol/g}$ Dry Weight in the top part of the sediment to 8 $\mu\text{mol/g}$ Dry Weight in the deep part
104 of the sediment (~30 cm) (Eckert, 2000).

105

106 **2. Material and Methods**

107 **2.1. Sampling**

108 Several sediment cores were collected from the center of the lake (Station A, Fig. 1) at a water
109 depth of ~42 m (maximum lake water column depth) at different times using Perspex tubes,
110 measuring 55 cm long by 5 cm in diameter, with a gravity corer. The cores were stored in the dark
111 at 4°C until they were sliced (on the same day or the day after). Core sampling for the microbial
112 community study took place in December 2009. Over a dozen of porewater chemical and isotope

113 profiles were conducted seasonally (every 3-4 months) from 2007 to 2013. The slight seasonal
114 changes allowed using typical geochemical profiles (Adler et al., 2011) in order to sample for the
115 microbiology communities in the different electron acceptors zones and to correlate between the
116 microbial sampling and the geochemical profiles.

117 **2.2. Geochemical analyses**

118 Cores were divided into 2-cm slices from top to bottom under a constant flow of N₂ using a slicing
119 device. About 1.5 ml of each sediment slice was transferred into N₂-flushed crimp bottles
120 containing 5 ml of 1.5 N NaOH for the headspace measurements of CH₄ and δ¹³C_{CH4} (after Sivan et
121 al., 2011). CH₄ concentrations were measured on a SHIMADZU gas chromatograph (GC 8IF)
122 equipped with a FID detector at a precision of 2 μmol·L⁻¹.

123 Porewater was extracted immediately from each slice by centrifugation at 27,000 g at 4°C in a
124 N₂ atmosphere, and the supernatant was filtered through 0.45-μm filters. A 1-ml subsample was
125 fixed with Ferrozine and analyzed for dissolved Fe(II) (Stookey, 1970). Four millimeters of
126 subsample were poured into an acidified vial (1 ml of 0.5N nitric acid) to measure dissolved Mn(II).
127 The sample was analyzed via an Inductively Coupled Plasma Mass Spectrometer (ICP-MS, Elan
128 DRC II, Perkin Elmer) at a precision of ±10%. For sulfide profiles, 1 ml of subsample was added to
129 zinc acetate and hydrogen sulfide concentrations were determined by the methylene blue method
130 with MDL of the method is 1 μM (Cline, 1969). For sulfate measurements, 5 ml of the subsamples
131 were analyzed with a Dionex DX500 high-pressure liquid chromatograph with a precision of ±3%.
132 Iron isotope analysis was done by acidifying the subsamples with 10% HCl for one week (to
133 dissolve any precipitated iron) and then purifying them by anion exchange chromatography (Borrok
134 et al., 2007). δ⁵⁶Fe was measured on a Neptune multi-collector ICP-MS in high resolution mode
135 according to standard methods and standardized against isotopic reference material (IRMM-014)
136 with a precision of ±0.1‰ (John and Adkins, 2010). Total lipids were extracted using the Bligh–
137 Dyer procedure (Bligh and Dyer, 1959) with solvent mixture of 2:1:0.8
138 (methanol:dichloromethane:buffer). δ¹³C of the total lipid extraction (TLE) was measured on an
139 Elemental Analyzer Isotopic Ratio Mass Spectrometer (EA-IRMS) with a precision of 0.1‰.

140 **2.3. DNA extraction and quantitative PCR (qPCR) amplification from sediment samples**

141 Part of the sediment slices were kept frozen at -20°C for the microbial work. Samples from three
142 different depths (0–3 cm representing the sulfate reduction zone, 6–9 cm representing the
143 methanogenesis zone, and 29–32 cm representing the deep AOM zone) were defrosted. Those depth
144 zones were chosen based on the geochemical profiles sampled at different times, showing quasi
145 steady state and stabilization of the sediments (Adler et al., 2011; Sivan et al., 2011). Therefore,

146 microbial community sampling of the three depth zones at a single time point should represent the
147 different microbial habitats as they pertain to the respective geochemical zones. Total genomic
148 DNA was extracted from the sediment samples using the MoBio Power Soil DNA isolation kit
149 (MoBio Laboratories, Solana Beach, CA). Genomic DNA was eluted using 60 µl of elution buffer
150 and stored at -20°C. Concentrations of DNA were determined via UV-Vis spectrophotometry (ND-
151 1000 NanoDrop Technologies, Wilmington, DE) (sulfate reduction zone 22ng/µl, methanogenesis
152 zone 35.8ng/µl and AOM zone 14ng/µl).

153 Quantification of functional genes were performed using the *mcrA* primers (Luton et al., 2002),
154 F 'GGTGGTGTMGGATTCACACARTAYGCWACAG' and R
155 'TTCATTGCRTAGTTWGGRTAGTT', *dsrA* primers, DSR1F 'ACSCACTGGAAGCACG'
156 (Wagner et al., 1998) and RH3-dsr-R 'GGTGGAGCCGTGCATGTT' (Ben-Dov et al., 2007) and
157 *pmoA* primers, a189F 'GGNGACTGGGACTTCTGG' and mb661R
158 'CCGGMGCAACGTCYTTACC' (Yan et al., 2006) with a ABI prism 7000 Sequence Detection
159 System (Applied Biosystems). The qPCR reaction consisted of 10 µL of Absolute Blue SYBR
160 Green ROX, 150 nM each of forward and reverse primers and 5 µL of each DNA template.
161 Thermal cycling conditions were as follows: 15 minutes at 95°C for enzyme activation, followed by
162 40 rounds of 15 seconds at 95°C for denaturation and 1 minute at 60°C for annealing/extension. To
163 verify that the used primer pair produced only a single specific product, a dissociation protocol was
164 added after thermo cycling, to determine dissociation of the PCR products from 60°C to 95°C.
165 Standards for the calibration curves for quantification were made using pGEM-T Easy plasmid
166 cloned with 1.9 kb *dsrAB* and amplicons of the *mcrA* and *pmoA* encoding for functional genes at a
167 known concentration with six serial dilution points (in steps of tenfold). All runs included a no-
168 template control. Plasmid standards and environmental samples were simultaneously assayed in
169 triplicates. The ABI prism 7000 Sequence Detection System and SDS Software were used for data
170 analysis. qC values were exported into a Microsoft Excel Work sheet for further statistical analysis.

171 **2.4. Sequence analysis**

172 454 tag-encoded FLX amplicon pyrosequencing was performed by the Research and Testing
173 Laboratory (Lubbock, Texas, USA) as previously described (Dowd et al., 2008). The bacterial and
174 archaeal 16S rRNA gene primers that were used are 28F 'GAGTTTGATCNTGGCTCAG' and
175 519R 'GTNTTACNGCGGCKGCTG' and Arch349F 'GYGCASCAGKCGMGA AW' and Arch806R
176 'GGACTACVSGGGTATCTAAT' respectively. Data analysis was made using two different
177 methodologies (MOTHUR and SILVA ngs). The initial trimming of the sequences was made by
178 MOTHUR v1.33 (Schloss et al., 2009) and generated around 300 bp length of sequences. The

179 trimmed sequences were taken to further analysis by MOTHUR (Schloss et al., 2011) using
180 SILVA.nr_v119 database. MOTHUR 454 pipeline filtration and denoising remove from the
181 analysis sequences that were <150 bp, when they contained homopolymers longer than 8 bp,
182 ambiguous bases, more than one mismatch to barcode sequences or more than two mismatches to
183 the forward primer sequence. We further removed sequences that did not align in the same
184 nucleotide position on the reference database. Most of the bacterial sequences were in the same
185 region, however the archaeal sequences were spared between two different regions. We used one
186 position window that was dominant in the top sample and implied it on the other archaeal analysis.
187 Identical sequences were grouped and then were aligned against SILVA.nr_v119. Chimers were
188 removed using MOTHUR CHIMERA UCHIME (Edgar et al., 2011). The lengths of the remaining
189 sequences were around 200 bp. A further screening step (pre-cluster) was applied to reduce
190 sequencing noise by clustering reads differing by only one base every 100 bases (Huse et al., 2010).
191 In addition, the rest of the sequences were classified in order to remove eukaryote, mitochondria
192 and chloroplast classified sequences. The remaining sequences were used to generate a distance
193 matrix and clustering into operational taxonomic units (OTUs) defined at 97% cutoff using the
194 average neighbor algorithm. The OTUs were classified using SILVA.nr_v119 database with a
195 confidence threshold of 80%. Alpha diversity was calculated by MOTHUR using the remaining
196 sequences at 97% similarity. The beta diversity was calculated using comparable data, the number
197 of sequences per sample was made equal through subsampling (Table 1 and 2). The second program
198 which we used for analysis was SILVA ngs pipeline (Quast et al., 2013). The trimmed sequences
199 from mother were Aligner against the SILVA SSU rRNA seed. Sequences shorter than 50 aligned
200 nucleotides and with more than 2% of ambiguities or 2% of homopolymers, respectively, were
201 removed. The sequences that were not aligned as being putative contaminations/artifacts were
202 removed. Then sequences were clustered to OTUs with 98% similarity and classified by local
203 nucleotide BLAST search against SILVA database v119. To filter out low identity and artificial
204 BLAST hits, hits for which the function $(\%sequence\ identity + \%alignment\ coverage)/2$ did not
205 exceed the value of 93% were discarded. Sequences with weak low score were classified as 'No
206 Relatives' and will be disregarded from now on. Standard deviation between the percentages of the
207 two classification showed that at the phylum level there are small differences between the two
208 pipelines (supplementary Table 1). In the order classification, the standard deviations increased but
209 the sequences still showed close similarity. Estimates of phylotype richness, diversity coverage and
210 similarity were calculated according to the abundance-based coverage estimate (ACE), Chao's
211 estimator (Chao, 1984; Chao and Ma, 1993), the Shannon diversity index, Good's coverage (Good,
212 1953) were calculated in MOTHUR (Table 1 and 2). The distribution and abundance matrix of the
213 OTUs was normalized to the sample with the smallest number of reads by randomly resampling of

214 the MOTHUR data set for statistical comparisons. Raw sequencing data was deposited in the MG-
215 RAST (metagenomics.anl.gov) archive.

216

217 **2.5. Microbial community structure related to the environment conditions**

218 To estimate community similarity among samples, we used PC-ORD 6 software. Subsample OTU
219 data from MOTHUR was used to create a distance matrix based on the Sørensen (Bray–Curtis)
220 dissimilarities of the OTU composition of the samples. The data was normalized to percentages
221 before the analysis. Community relationships were visualized using principal coordinate analysis
222 (PCoA) based on this distance matrix. Different environmental variables were added as well (CH₄,
223 $\delta^{13}\text{C}_{\text{CH}_4}$, Fe(II), $\delta^{56}\text{Fe(II)}$, Mn(II), H₂S, SO₄²⁻, dissolved organic carbon (DOC), $\delta^{13}\text{C}_{\text{TLE}}$, NH₄⁺ and
224 PO₄³⁻) in order to identify potential explanatory variables. The environmental vector were applied
225 and projected to the PCoA ordination. In addition, Venn diagrams for graphical descriptions of
226 unshared and shared OTUs between the three samples were constructed using MOTHUR

227 **3. Results**

228 **3.1. Porewater profiles**

229 This study focused on microbial community shifts along the porewater profiles of electron acceptor
230 gradients related to the methane cycle. Over dozen of porewater profiles were performed seasonally
231 from station A (Fig. 1) to characterize shifts in main electron acceptors with depth. The shown
232 profiles (Fig. 2) are representatives of the slight seasonal trends. Sulfate and sulfide profiles show
233 depletion from ~100 μM at the top part of the sediment, to below detection limits within the upper
234 15 cm (Fig. 2A). This typical concave curvature profile of sulfate in the porewater indicates
235 intensive sulfate reduction in the upper few centimeters of the sediment throughout the year. A
236 typical LK methane concentration profile (Fig. 2B) shows increase from 250 μM at the water-
237 sediment interface to a maximum of about 1.25 mM in the depth range of 7 to 15 cm, and then a
238 decrease below 15 cm depth. The profile of $\delta^{13}\text{C}_{\text{CH}_4}$ (Fig. 2B) shows a decrease from -60‰ at a
239 depth of 1 cm to about -65‰ at a depth of 7 cm and then an increase in the deeper sediments to a
240 maximum value of -53.5‰ at a depth of 25 cm. Also, the profile of $\delta^{13}\text{C}_{\text{TLE}}$ (Fig. 2C) shows a
241 decrease in this deepest part of the sediment.

242 Manganese and iron oxides are the most probable electron accepters in the deep part of the sediment
243 where methane is decreased. Dissolved Mn(II) concentration (Fig. 2D) increased from 5 μM at the
244 top of the sediment to a plateau of about 23 μM from depths of 23 to 36 cm. The Fe(II)
245 concentration profile also showed an increase with depth (Fig. 2D), however in a different pattern.

246 In the upper 15 cm, dissolved Fe(II) concentrations were below the detection limit, and gradually
247 increased below 15 cm to about 90 μM at a depth of 36 cm. The $\delta^{56}\text{Fe}$ profile (Fig. 2C) showed a
248 decrease with depth from 0.5‰ in the upper part to -1.7‰ to -2.3‰ in the deep part of the
249 sediment.

250 3.2. Sediment microbial communities

251 To study the sediment depth-dependent microbial community shifts, 16S rRNA gene sequences of
252 the prokaryotic community from three different depth zones (0–3, 6–9 and 29–32 cm, see arrows in
253 Fig. 2a) were extracted and analyzed. As mentioned above, these three zones represent the main
254 sulfate reduction zone, the methanogenesis zone, and the deep iron driven AOM zone, respectively.
255 The diverse bacterial and archaeal communities at these depth zones varied in composition and
256 richness throughout the sediment. A total of 13227 and 4881 bacterial and archaeal sequences
257 representing 3852 and 705 OTUs (cut-off value of 98%), respectively, were identified by SILVA
258 database project. The normalized Shannon and Chao index were used as proxies for diversity and
259 richness, respectively (Table 1 and 2). The upper layer showed lower diversity (Shannon index,
260 3.91 and 1.8 for bacteria and archaea) and richness (Chao index, 925.7 and 83.4 for bacteria and
261 archaea) than the other depths. Moreover, the bacterial coverage of the upper layer was the highest
262 (91%), while that of the deeper layers was about 80%. The archaeal coverage was almost the same
263 (~88%) in all the sampled layers. Most of our sequences were affiliated with uncultured
264 microorganisms mainly from various sediment environments.

265 Although the bacterial OTUs were distributed over 43 phyla, we represent here only the phyla
266 that were over 1% sequences of the 454 library. The most abundant phyla were: *Proteobacteria*,
267 *Chloroflexi*, *Nitrospirae*, *Bacteroidetes*, *Firmicutes* and *Chlorobi* observed in at least one of the
268 libraries (Fig. 3). Dominant phyla had different trends; *Chloroflexi* and *Nitrospirae* showed increase
269 with depth while *Proteobacteria* showed decrease with depth. *Firmicutes* were highly abundant
270 only in the top layer. *Bacteroidetes* and *Chlorobi* were more abundant in the middle layer of the
271 sediment.

272 The communities of microorganisms in the top layer (0-3cm) of the sediment had less sequence
273 overlap with those of the deeper layers (70 and 33 shared bacterial sequences with the middle and
274 bottom layers respectively) and of the deeper layers (Supplementary Fig 1 and 2). The most
275 dominant (~48%) phylum in this bacterial community was *Proteobacteria*; class
276 *Gammaproteobacteria* (91%) and the genera *Acinetobacter* (28%) and *Pseudomonas* (14%) (Fig.
277 3). The second most abundant (38%) phylum was *Firmicutes*; class *Clostridia* (21%) and *Bacilli*
278 (17%), family *Clostridiaceae* (15%) and genus *Bacillus* (14%). Further sequences were related to
279 other phyla and order (Fig 3 and supplementary Table 1).

280 In the archaeal community the sequences number were much lower, however the coverage
281 calculation shows about 90% coverage. The dominant phylum was *Euryarchaeota* (98%), and the
282 rest of the 2% was *Thaumarchaeota*. *Methanomicrobia* (92%) was the dominant class of
283 *Euryarchaeota* divided between genera of *Methanoregula* (58%), *Methanosaeta* (22%) and
284 *Methanolinea* (11%). The other abundant classes of *Euryarchaeota* were *Halobacteria* (2%) and
285 *Thermoplasmata* (3%). *Thaumarchaeota* was divided between two classes: Marine Benthic Group
286 B (MBGB 1%) and Miscellaneous Crenarchaeotic Group (MCG 1%) (Fig. 4).

287 The middle layer (6-9cm) was the richest and most diverse in its bacterial and archaeal
288 populations comparing to the other layers, but with the lowest community coverage (bacterial 77%
289 and archaeal 86%). The dominant phyla in this depth were divided to *Proteobacteria* (40%),
290 *Chloroflexi* (14%), *Bacteroidetes* (7%), *Chlorobi* (5%) and *Nitrospirae* (4%). The dominant classes
291 in *Proteobacteria* were *Deltaproteobacteria* (68%), *Betaproteobacteria* (14%) and
292 *Gammaproteobacteria* (14%). Sva0485 (8%), *Syntrophobacterales* (7%), *Desulfarculales* (3%) and
293 *Desulfuromonadales* (2%) were the dominant affiliated order in *Deltaproteobacteria*.
294 *Methylococcaceae* family of *Gammaproteobacteria* had 1% affiliated sequences. The dominant
295 family in *Chloroflexi* was *Anaerolineaceae* (6%). *Ignavibacteriales* (4%) order was the dominant in
296 *Chlorobi*. *Nitrospirae* increased from the upper layer and was mainly represented by the
297 *Nitrospiraceae* (4%) family. Further sequences were related to other phyla and order (Fig. 3,
298 supplementary Table 1).

299 The archaeal community was the richest and most diverse in the middle zone. The dominant
300 phyla were *Euryarchaeota* (96%) while *Thaumarchaeota* has the rest of the 4%. *Euryarchaeota* was
301 divided between 3 dominant classes: *Methanomicrobia* (78%), *Thermoplasmata* (11%) and
302 *Halobacteria* (6%). *Methanomicrobia* was divided between *Methanoregula* (19%),
303 *Methanosaeta*(50%) and *Methanolinea* (10%) genus's. *Thermoplasmata* was divided between
304 Marine Benthic Group D and DHVEG-1 (5%) and Terrestrial Miscellaneous Gp(TMEG) (3%).
305 *Halobacteria* dominant family was Deep Sea Hydrothermal Vent Gp 6(DHVEG-6) (6%)
306 *Thaumarchaeota* was divided mainly between two classes: Marine Benthic Group B (MBGB 2%)
307 and Miscellaneous Crenarchaeotic Group (MCG 1%) (Fig. 4).

308 The bottom layer (29-32 cm) bacterial community dominant phyla included: *Proteobacteria*
309 (39%), *Chloroflexi* (19%) and *Nitrospirae* (8%). *Proteobacteria* were divided to three main classes:
310 *Deltaproteobacteria* (20%), *Gammaproteobacteria* (10%) and *Betaproteobacteria* (6%). Sva0485
311 (11%) and *Syntrophobacterales* (4%) were the dominant affiliated order in *Deltaproteobacteria*.
312 The dominant family in *Chloroflexi* was *Anaerolineaceae* (7%) and *Ignavibacteriales* (4%) order is
313 the dominant *Chlorobi*. *Nitrospirae* increased from the upper layer and was mainly represented by

314 *Nitrospiraceae* (8%) family. Further sequences were related to other phyla and order (Fig. 3,
315 supplementary Table 1)

316 At this depth, the dominant archaeal phyla were *Euryarchaeota* (98%) while *Thaumarchaeota*
317 has the rest of the 2%. *Euryarchaeota* was divided between 3 dominant classes: *Methanomicrobia*
318 (86%), *Thermoplasmata* (6%) and *Halobacteria* (5%). *Methanomicrobia* was divided between
319 *Methanoregula* (16%), *Methanosaeta*(58%) and *Methanolinea* (11%) genus's. *Thermoplasmata* was
320 divided between Marine Benthic Group D and DHVEG-1 (5%) and Terrestrial Miscellaneous
321 Gp(TMEG) (3%) family. *Halobacteria* dominant family was Deep Sea Hydrothermal Vent Gp
322 6(DHVEG-6) (1%) *Thaumarchaeota* was divided mainly between two classes: Marine Benthic
323 Group B (MBGB 1%) and Miscellaneous Crenarchaeotic Group (MCG 1%) (Fig. 4). The detailed
324 description of the major communities in the sediment is described at the discussion and the
325 appendix. No ANME sequences were detected even though specific primers (ANME2C-AR468F,
326 ANME3-1249, ANME1-395F, ANME1-1417, ANME3-140F, ANME3-1249, ANME2a-426 and
327 ANME2a-1242R, see Miyashita et al., 2009) were used.

328

329 **3.3. Depth comparison of microbial communities**

330 An ordination plot from the 454 pyrosequencing by MOTHUR subsample OTUs dataset was
331 derived from Principal coordinate's analysis (PCoA) of the bacterial and archaeal. The PCoA
332 display the similarities and differences between the bacterial and archaeal communities varied with
333 depth (Fig. 5a and 5b respectively). The PCoA of bacteria and archaea show very similar separation
334 between the communities of different layers related to the different environment conditions of each
335 layer. The two deeper layers are on the same distant on the X axis showing relatively more
336 similarity than the upper layer. The vectors of the sulfate and sulfide are correlative with
337 communities of the top layer, while methane, $\delta^{13}\text{C}_{\text{TLE}}$ and $\delta^{56}\text{Fe(II)}$ were more associated to the
338 communities of the middle layer. The communities in the bottom layer were more correlated to
339 Fe(II), Mn(II) and NH_4^+ . The Venn diagram shows also that more OTUs from the deep layers were
340 shared than between the upper layer and the deeper layers (Supplementary Fig 1 and 2).

341 **3.4. Functional gene profile**

342 In order to better understand the abundance of microbial functionality in the different depth we used
343 qPCR for different functional genes. The functional genes *mcrA* and *dsrA* had very similar pattern -
344 low concentrations in the upper layer ($9 \times 10^5 \pm 6.4 \times 10^4$ and $1 \times 10^6 \pm 1.7 \times 10^5$ respectively) and highest
345 concentrations ($6.9 \times 10^6 \pm 6.7 \times 10^5$ and $6.9 \times 10^6 \pm 9.9 \times 10^4$) in the middle layer. *pmoA* gene showed also
346 the same pattern, however with lower concentration ($2.3 \times 10^5 \pm 9.7 \times 10^3$ to $1.6 \times 10^6 \pm 1.7 \times 10^5$) (Fig. 6).

348 4. Discussion

349 In this study of LK sediments we investigated changes in the microbial diversity associated with
350 porewater geochemistry and the transition of the dominant electron acceptors with depth (0-40 cm).
351 The geochemical porewater profiles in LK (Fig. 2) suggest that the sediment can be broadly divided
352 into three different regions of microbial processes: the upper 5 cm which is dominated by sulfate
353 reduction, the methanogenesis zone between 5 to 17 cm, and the deep sediments, dominated by
354 anaerobic oxidation of methane coupled to iron reduction. This division of the sediment and the
355 deep iron-driven AOM processes were confirmed by *in situ* profiles of methane, $\delta^{13}\text{C}_{\text{CH}_4}$, sulfate,
356 and ferrous iron and subsequent use of a numerical mass conservation model (based on the
357 geochemical profiles of DIC and $\delta^{13}\text{C}_{\text{DIC}}$ (Adler et al., 2011) and a set of geochemical incubation
358 experiments conducted in our previous work (Sivan et al., 2011).

359 The mechanisms that enable the process of AOM via iron reduction can be characterized by
360 the existence of specific prokaryotic populations at each depth and their resemblance to similar
361 environments with a distinct characterization. There are only few studies that analyzed microbial
362 communities relative to geochemical zones in fresh water sediments (Deutzmann and Schink, 2011;
363 Koizumi et al., 2004; Ye et al., 2009). Therefore, this study can deepen the information on
364 microbial community shifts under different electron acceptors conditions, especially those related to
365 the methane cycle. It is also the first study in LK that divides the sediment to different layers and
366 attribute significantly to the diversity of the microbial ecology in the different zones.

367 Pipeline of SILVA ngs was used for the alpha-diversity to get more classified sequences and
368 better coverage of the classification and population. However, for the beta-diversity the data from
369 the MOTHUR pipeline applied to get more statistical results without the bias of different sample
370 size. The changes between the pipeline results for the alpha diversity were not large in the phyla
371 level (standard deviation of ~0.5%) but increased with higher taxonomy levels.

372 4.1. Sediment geochemical conditions

373 In the upper layer of sediment (0-3 cm), sulfate concentrations were the highest and decreased with
374 depth as particulate organic matter from the water column accumulates, making organic carbon
375 degradation available (Hadas and Pinkas, 1992). Adler et al. (2011) calculated that the bulk of
376 bacterial sulfate reduction occurs in the top 1 cm of sediment, a finding that was supported by the
377 microbial work of Hadas and Pinkas, (1992). Using sulfide and particulate organic carbon
378 measurements, Eckert and Conrad. (2007) showed that sulfate reduction accounts for most of the
379 mineralization of organic matter in the upper 15 cm of sediment. Methane concentrations in the

380 upper layer were relatively low and probably reached from upward diffusion as methanogens are
381 outcompeted by sulfate reducers there (Lovley and Klug, 1983). The depletion in methane
382 concentrations and the decline $\delta^{13}\text{C}_{\text{TLE}}$ values and the heavier isotopic values of $\delta^{13}\text{C}_{\text{CH}_4}$ in the
383 upper layer (Fig 2) compared to the methanogenic zone may be explained by AOM via sulfate
384 reduction, although no ANME sequences were found using specific primers or by phylogenetic
385 alignment at any depth in the sediment.

386 Below the sulfate reduction zone, in the middle layer of sediment (6-9 cm), methane reached
387 its maximum concentrations (Fig. 2B). The low values of $\delta^{13}\text{C}_{\text{CH}_4}$ in the methanogenesis zone are
388 typical due to the large carbon isotope fractionation that occurs during methane production, thereby
389 leaving the produced methane isotopically light and the DIC isotopically heavy. Below the
390 maximum methane production zone (around 20 cm), methane concentrations began to decrease
391 while $\delta^{13}\text{C}_{\text{CH}_4}$ values increased (Fig. 2B), an outcome that may be due to the AOM process, which
392 leaves residual methane isotopically heavier. Depleted $\delta^{13}\text{C}_{\text{TLE}}$ (Fig. 2C) in the deep part of the
393 sediment also supports the presence of AOM with the production of light biomass from methane
394 oxidation. In addition, although all the suitable electron acceptors at this depth were depleted, both
395 Fe(II) and Mn(II) increased (Fig. 2D). Also, $\delta^{56}\text{Fe}$ values (Fig. 2C) of dissolved iron in the deep
396 sediment were isotopically negative, similar to sediments with active dissimilatory bacterial iron
397 reduction (Severmann et al., 2006) and are an indication of active iron reduction in the deep
398 sediments and not just diffusion. The increase in Fe(II) concentrations below a depth of 15 cm was
399 probably supported by the absence of sulfide. The potential of highly reactive Fe(III) oxides to drive
400 AOM in LK was also supported by our set of mesocosm incubation studies (Sivan et al., 2011).
401 Since manganese oxide concentrations were very low (~0.04%) throughout the LK sediment
402 column (Serruya et al., 1974), iron oxide seemed to play a bigger role in this AOM process.

403 **4.2. Microbial population that could be involved in methane cycle**

404 *4.2.1. Deltaproteobacteria*

405 *Deltaproteobacteria* were the most abundant in the middle and deep sediment samples.
406 Many members of the families of *Syntrophobacterales* are sulfate reducers or as fermentative
407 organism, depending on the environmental conditions (McInerney et al., 2007). *Syntrophaceae*, the
408 dominant family of *Syntrophobacterales*, are acetate degrading sulfate reducing bacterium (Jiang et
409 al., 2009). *Desulfarculales* are strictly anaerobic mesophilic sulfate-reducing bacterium with the
410 capability to oxidize acetate and fatty acids (Sun et al., 2010). *Desulfobacterales* are sulfate
411 reducer, capable of oxidizing hydrogen in soils and sediments (Burow et al., 2014).
412 *Desulfuromonadales* was shown as S^0 respiring (Pjevac et al., 2014) and may also reduce Fe(III)
413 and Mn(IV) in marine surface sediments (Lovley, 2006). About 1% of the sequences in the middle

414 and bottom layers or the cores were affiliated to the *Geobacter* genus members which are involved
415 in iron reduction. In the deep layer the most abundant class was Sva0485. Sequences of the SVa485
416 order were similar to those found in different aquatic environments, but the metabolic functions of
417 members of this order are not clear. *Pelobacter carbinolious* (Lovley et al., 1995), a member of
418 SVa485 is capable of Fe(III) and sulfur reduction. Schwarz et al. (2007a) showed that in the upper
419 part of LK sediment *Deltaproteobacteria* was one of the dominant classes, and that most of them
420 were affiliated with acetate oxidizing sulfate reducing bacteria, that outcompete the acetoclastic
421 methanogens. Even though *Deltaproteobacteria* are best known for their sulfate reduction
422 metabolism, they can shift their metabolism in response to depleted sulfate concentrations (Plugge
423 et al., 2011).

424 4.2.2. *Betaproteobacteria*

425 *Betaproteobacteria* percentages increased slightly with depth. These percentages are comprised of
426 chemoheterotrophs and chemoautotrophs that derive nutrients from decomposition of organic
427 material. *Burkholderiales* was the dominant order in *Betaproteobacteria* and increased with depth
428 (1.5% to 4%). (Schwarz et al., 2007b) showed that *Burkholderiales* is active in the upper sediment
429 of LK. Members of this group are able to carry out anaerobic oxidation of acetate with (per)chlorate
430 as electron acceptor (Yoshida et al., 2005) or oxidize hydrogen in deep subsurface (Orcutt et al.,
431 2011).

432 4.2.3. *Gammaproteobacteria*

433 Our *Gammaproteobacteria* affiliated sequences have different phylogenetic classification
434 distribution in each layer. In the middle and bottom layers members of the *Methylococcales* order
435 were observed (~1%). *Methylococcales* are aerobic methanotrophs which some members were
436 present within the sediments and overlying water column from dysoxic, methane-rich vent and seep
437 systems (Tavormina et al., 2008). The most abundant order in the deep layer was HOC36 (5%)
438 which is an uncultured *Gammaproteobacteria*. however when compared to NCBI data base it was
439 found to be closely related to uncultured LK clones (99%) and to cultured *Methylocaldum* sp. (94%)
440 (Bodrossy et al., 1997) which is a Thermophilic methanotroph isolated from landfill cover soil.

441 4.2.4. *Nitrospirae*

442 Members of the *Nitrospirae* phylum, increased with depth. Sequences from our samples were
443 classified to *Nitrospirales* order in *Nitrospirea* and were most abundant in the bottom layer of the
444 sediment (7%). Part of our sequences were similar to those found in a previous study (Schwarz et
445 al., 2007b). . Using SIP-RNA, (Schwarz et al., 2007b) showed t not only that *Nitrospirae* were
446 present, but also that they were functionally bioactive. Although *Nitrospirae* is a known nitrate
447 oxidizer (Ehrich et al., 1995), the conditions of this environment suggest that it utilized another
448 metabolic pathway. *Nitrospirae* also include the iron reducing candidates such as

449 *Magnetobacterium bavaricum* (Spring et al., 1993) and sulfur reducers (Sonne-Hansen and Ahring,
450 1999). Part of our sequences were aligned to uncultured clone (98%) from freshwater sediment
451 which enhanced degradation of phenanthrene and pyrene by amorphous ferric hydroxide (Yan et
452 al., 2012) .

453 4.2.5. *Methanomicrobia*

454 The majority of our sequences were classified within the *Euryarchaeota*, in the
455 *Methanomicrobia* class. Our 16S rRNA sequences were divided between the *Methanomicrobiales*
456 and *Methanosarcinales* order. *Methanosaeta* genus of *Methanosarcinales* increases with depth.
457 *Methanosaeta* are acetoclastic methanogens which only able to grow on acetate even at low
458 concentrations (Jetten et al., 1990). *Methanosaeta* was also shown in previous studies of LK
459 (Schwarz et al., 2007a, 2007b) and in other meso to eutrophic freshwater lakes (Glissman et al.,
460 2004; Koizumi et al., 2003). Yamada et al. (2014) showed that *Methanosaeta* also have the
461 ability to reduce ferrihydrite with H₂ as the electron source. The *Methanosaeta concilii* was the
462 most similar cultured acetoclastic methanogen (96%) to our sequences which was observed also in
463 the previous study. The *Methanolinea* genus of *Methanomicrobiales* a hydrogenotrophic
464 methanogen was observed in constant percentages for all depths. Members of the *Methanoregula*
465 genus, a genus that is also hydrogenotrophic methanogen, decreased with depth in the core.
466 *Methanoregula*, *Methanosaeta* and *Methanolinea* represent 60% of sequences of methanogens
467 retrieved from freshwater lakes (Borrel et al., 2011).

468 4.2.6. *Thermoplasmata*

469 The dominant family in *Thermoplasmata* is The Marine Benthic Group D and DHVEG-1
470 (MBG-D). Members of the MBG-D have been shown to exist in a variety of freshwater and marine
471 environments (Beal et al., 2009; Borrel et al., 2012), and it is the most widely encountered,
472 uncultured lineage in freshwater lake sediments. Even though their metabolism is unknown,
473 hypotheses about their functionalities are based on the environments in which they were found.
474 Methanogenesis was suggested, as they were found in deep lake sediments with high methane
475 concentrations (Borrel et al., 2012), and they were also hypothesized to be involved in AOM, as
476 they were found in AOM zones (Schubert et al., 2011) and in marine seep sediment (Beal et al.,
477 2009). However, in other environments, in which methane concentrations were low, the utilization
478 of waste products, intermediates, or dead cells by MBG-D was also suggested (Smith et al., 1975).
479 Recently, a single cell genomics study showed that members of MBG-D were capable of exogenous
480 protein degradation in cold anoxic environments (Lloyd et al., 2013).

481 4.2.7. *Thaumarchaeota*

482 *Thaumarchaeota* are mostly known as archaeal ammonia oxidizers but in the sediments they may
483 contribute significantly to the reservoir of nitrogen oxides in ocean waters and thus to productivity,

484 including the assimilation of carbon (Pester et al., 2011). The enzyme ammonia monooxygenase,
485 which belongs to the enzyme family of copper containing membrane bound monooxygenases that
486 possess wide substrate ranges. Closely related enzymes, which can often perform the same function
487 but with different substrates, include ammonia oxidizing bacteria with ammonia (in
488 *Gammaproteobacteria*) and methane as a substitute (Lontoh et al., 2000). However, a
489 microorganism's metabolic energy is mainly defined by its downstream enzyme machinery
490 (Tavormina et al., 2011). In our sediment *Thaumarchaeota* observed in low percentages with 2
491 main classes: Marine Benthic Group B and Miscellaneous Crenarchaeotic Group.

492 4.2.8. Miscellaneous Crenarchaeotic Group

493 The Miscellaneous Crenarchaeotic Group (MCG) is a cosmopolitan class assigned to the
494 *Thaumarchaeota* phylum whose members are found mostly in anoxic habitats and have the
495 capability to take up organic carbon (Biddle et al., 2006). Because of their huge environmental
496 range and their complex phylogeny, the MCG collectively possess great metabolic diversity (Jiang
497 et al., 2008). Current evidence for members of the MCG lineage suggests that they may obtain
498 energy from the anaerobic oxidation of methane, but they do so via a dissimilatory methane-
499 oxidizing process and they do not assimilate its carbon (Biddle et al., 2006)

500 4.2.9. Marine Benthic Group B

501 Members of the MBGB were detected in a growing number of benthic marine environments,
502 including Atlantic deep-sea sediments, marine mud volcanoes, marine carbonate crusts from the
503 Black Sea, organic-poor deep subsurface sediments of the central oceanic basins, and organic-rich
504 methane or methane hydrate containing sediments near continental margins (Sørensen and Teske,
505 2006). Recent results from Ocean Drilling Program showed that no known methanotroph was
506 detected however methane oxidation appeared to be mediated by Marine Benthic Group B and the
507 Miscellaneous Crenarchaeotal Group (Biddle et al., 2006).

508 4.3. Concept for methane cycle mechanisms

509 A few possible mechanisms for the novel process of iron driven AOM in the deep part of the
510 sediment can be suggested based on the geochemical profiles and the microbial communities. A
511 possible direct process could be through new, currently unknown bacteria/archaea that reduce iron
512 and utilize methane. Possible indirect processes include: 1) Reduction of Fe(III) oxides by oxidation
513 of sulfide (in pyrite or FeS minerals) to sulfur intermediates, and then disproportionation to sulfide
514 and sulfate (such as in (Holmkvist et al., 2011)), and sulfate driven AOM but without ANME. 2)
515 Reduction of Fe(III) by utilizing H₂ (Lovley, 1991), which creates a low concentration of H₂ and
516 drives "reverse methanogenesis" (Hallam et al., 2004; Hoehler et al., 1994) by the archaea in the
517 deep sediment. 3) An oxygenic methane oxidation pathway in an anaerobic environment as

518 described by Ettwig et al. (2010), when methane is oxidized by oxygen that is released from iron
519 oxides.

520 The microbial population changes can be used to support the deep iron driven AOM
521 mechanism (presented by the sample from 29-32cm). The bacterial diversity in this depth was lower
522 than in the methanogenesis zone (6-9 cm) but higher than in the upper layer (0-3 cm) of the
523 sediment. The changes in bacterial diversity with depth could be related to the availability of
524 different electron acceptors and of organic matter (Nam et al., 2008). However, there is an overlap
525 between bacterial and archaeal sequences in the middle and the bottom layers (PCoA and Venn
526 diagram), indicating that the environmental conditions affecting the processes. Alpha diversity
527 indexes (Table 1 and 2) were compared to previous studies conducted only in the upper layer of the
528 sediment of LK using different methods (Schwarz et al., 2007a; Wobus et al., 2003), and showed a
529 similarity in the upper parts of the sediment. However, the deeper layers in this study showed higher
530 diversity and richness than the previous studies (Nusslein et al., 2001; Schwarz et al., 2007a, 2007b)
531 and was close to other studies using 454 sequencing (Hollister et al., 2010; Röske et al., 2012).

532 The direct mechanism of iron reduction coupled with methane oxidation can occur by one
533 microorganism, as methanogens have the ability to reduce iron (Van Bodegom et al., 2004; Bond et
534 al., 2002) however not with methane as the electron donor. According to our results, this kind of
535 unique methanotrophy could be from MBGD (*Thermoplasmata* class) or MCG/MBGB
536 (*Thaumarchaeota* phylum). Although their metabolisms are unknown, these groups were found in
537 methane rich environments. Another option is a consortium of microbes including those belonging
538 *Geobacter* genus, which are known for iron reduction, or *Nitrospirales*, known to reduce iron,
539 together with methanotrophs or with methanogens capable of "reverse methanogenesis". The
540 "reverse methanogenesis" pathway could be carried out by the dominant deep layer methanogens
541 *Methanosaeta*, which have been shown in a previous study as the dominant active methanogens in
542 the upper layer (Schwarz et al., 2007b). *Methanosaeta* could perform a different metabolic process
543 or may have been inhibited. *mcrA* functional gene of methanogens/anaerobic methanotroph shows
544 that in the middle layer methanogens were more abundant than in the upper and bottom layers (Fig
545 6). ANME were not found in the sediment using specific primers and Fluorescent *in situ*
546 hybridization methods. An additional reason for not finding any ANME sequences in our samples is
547 likely because that the *mcrA* observed in the deep layer belongs to the methanogens which maybe
548 cause the reverse methanogenesis.

549 The results suggest that *Desulfuromonadales* of *Deltaproteobacteria* class could be involved in
550 indirect mechanisms of disproportionation of sulfur together with other *Deltaproteobacteria* sulfate
551 reducer. The *dsrAB* functional gene shows that sulfate reducers are present at the same level of the
552 core where *mcrA* functional gene (Fig. 6) of methanogens or methanotrophs are found in the deep

553 part of the sediment. Additionally, the presence of the *Deltaproteobacteria* at the deepest sediment
554 depth and the observed accumulation of acetate with depth (data not shown) could indicate their
555 exploitation of a different metabolic path in the deep sediment than in the upper part of the
556 sediment.

557 The third indirect mechanism of anaerobic methane oxidation via an oxygenic pathway was
558 shown clearly by the *pmoA* functional gene. The *pmoA* concentration in the deep part of the
559 sediment was higher than in the upper part, indicating an oxygenic pathway (Fig. 6). However,
560 pMMO is a homologue enzyme of ammonia monooxygenase and might be sequenced together with
561 pMMO (Tavormina et al., 2011). The source of pMMO could be *Methylococcales*
562 (*Gammaproteobacteria*), which is an aerobic methanotroph found in the deep part of our samples.
563 However, bias of *pmoA* with ammonia monooxygenase could be caused by *Thaumarchaeota*,
564 Betaproteobacteria or Nitrospirae, which were found in all depth.

565 *Thaumarchaeota* comprise not only all known archaeal ammonia oxidizers, but also several
566 clusters of environmental sequences representing microorganisms with unknown energy
567 metabolisms (Pester et al., 2011). Members of *Thaumarchaeota* phylum could have monooxygenases
568 like enzymes that are able to capture methane due to the enzymes' phylogenetic proximities to
569 methane monooxygenases, but that activity requires the necessary downstream metabolic pathway. If
570 they could have that ability, they would have been good candidates for group of methanotrophic
571 archaea. This indirect pathway can be similar to that found in the NC10 phylum (Zhu et al., 2012),
572 which produces oxygen via the reduction of nitrite and the oxidation of methane, but with iron
573 oxides. Also many bacterial ammonia oxidizer sequences were found in our environment.
574 Ammonium profiles show increase with depth mainly due to decomposition of the organic matter,
575 and theoretically the large amount of ammonium could be oxidized by ferric iron minerals and
576 produce nitrite (Clement et al., 2005). Maybe ammonia monooxygenase can function for
577 ammonium up take for iron reduction and for methane up take for oxidation, but that is only a
578 speculation. However, no nitrate or nitrite was detected in the deep sediment, moreover no NC10
579 phylum (Ettwig et al., 2010) was observed in our sequences.

580 To summarize, this study attempted to correlate between the performed geochemical and
581 microbial profiles in lake sediments. The geochemical data suggest three main depth related zones
582 of electron acceptor activities in the sediment: sulfate reduction, methanogenesis and a novel, deep
583 iron-driven AOM. The prokaryotic analysis provided clues regarding the microorganisms that may
584 be involved in this novel process and the metabolic paths that occur throughout the microbial
585 assemblage. For AOM via iron reduction to occur, a number of potential pathways and their
586 combinations have been suggested. Orders that become enriched (Sva0485, *Methanosarcinales* and
587 *Nitrospirales*) with depth can be assumed to participate in the AOM process either directly or

588 indirectly. A possible direct process could be through new, currently unknown bacteria/archaea that
589 reduce iron and utilize methane, which may be carried out by a MCG as a methanotroph in a
590 consortium with an iron reducer like *Geobacter*. Possible indirect processes could be Fe(III)
591 reduction by sulfide, oxidation of the sulfide to elemental sulfur and other sulfur intermediates and
592 then disproportionation to sulfide and sulfate and sulfate driven AOM. However, this process is less
593 likely because ANMEs were not found. It could also be via reduction of Fe(III) by utilizing H₂,
594 creating a low concentration of H₂ and driveing “reverse methanogenesis”. Fe(III) reduction
595 processes could be carried out by *Nitrospirae* and/or *Deltaproteobacteria*. Members of both groups
596 can reduce iron while in a consortium with methanogenic *Methanosarcinales*, or can reduce the iron
597 with sulfur minerals, creating sulfate (see above) that *Deltaproteobacteria* can utilize while in a
598 consortium with a MCG as a methanotrophic archaea. An oxidation of methane coupled to iron
599 reduction pathway as described by Ettwig et al., (2010) could occur by *Thaumarchaeota* or
600 *Nitrospirae* with a monooxygenase enzyme that can utilize methane while using iron oxides to
601 generate the oxygen needed to oxidize the methane. Further research involving even larger sample
602 of the microbial community and characterization of more diverse functional genes will provide
603 better indication for the composition of the microbial communities at the different depths. Also,
604 microbial and geochemical experiments have the potential to give more clues on the mechanism of
605 this novel iron driven AOM.

606

607 **Acknowledgments**

608 We thank M. Adler for her assistance in the field and in the laboratory. Thanks to the members
609 of Orit's and Ariel's laboratories for all the help. Many thanks to V. Orphan, S. Connon and K.
610 Dawson from Caltech for their help and for their fruitful discussions. This research was funded by
611 the Water Authority of Israel (O.S. and W.E.).

612

613 **References**

- 614 Adler, M., Eckert, W. and Sivan, O.: Quantifying rates of methanogenesis and methanotrophy in
615 Lake Kinneret sediments (Israel) using pore-water profiles, *Limnol. Oceanogr.*, 56(4), 1525–1535,
616 2011.
- 617 Beal, E. J., House, C. H. and Orphan, V. J.: Manganese- and Iron-Dependent Marine Methane
618 Oxidation, *Science*, 325(5937), 184–187, 2009.
- 619 Biddle, J. F., Lipp, J. S., Lever, M. a, Lloyd, K. G., Sørensen, K. B., Anderson, R., Fredricks, H. F.,
620 Elvert, M., Kelly, T. J., Schrag, D. P., Sogin, M. L., Brenchley, J. E., Teske, A., House, C. H. and

- 621 Hinrichs, K.-U.: Heterotrophic Archaea dominate sedimentary subsurface ecosystems off Peru.,
622 Proc. Natl. Acad. Sci. U. S. A., 103(10), 3846–3851, 2006.
- 623 Bligh, E. G. and Dyer, W. J.: A rapid method of total lipid extraction and purification, Can. J.
624 Biochem. Physiol., 37(8), 911–917, 1959.
- 625 Van Bodegom, P. M., Scholten, J. C. M. and Stams, A. J. M.: Direct inhibition of methanogenesis
626 by ferric iron., FEMS Microbiol. Ecol., 49(2), 261–8, 2004.
- 627 Bodrossy, L., Holmes, E. M., Holmes, A. J., Kovács, K. L. and Murrell, J. C.: Analysis of 16S
628 rRNA and methane monooxygenase gene sequences reveals a novel group of thermotolerant, Arch
629 microbiol, 168, 493–503, 1997.
- 630 Boetius, A., Ravensschlag, K., Schubert, C. J., Rickert, D., Widdel, F., Gieseke, A., Amann, R.,
631 Jørgensen, B. B., Witte, U. and Pfannkuche, O.: A marine microbial consortium apparently
632 mediating anaerobic oxidation of methane, Nature, 407(6804), 623–626, 2000.
- 633 Bond, D. R., Lovley, D. R. and Methanococcus, A.: Reduction of Fe (III) oxide by methanogens in
634 the presence and absence of extracellular quinones, Environ. Microbiol., 4(2), 115–124, 2002.
- 635 Borowski, W. S., Cagatay, N., Ternois, Y. and Paull, C. K.: Data report: carbon isotopic
636 composition of dissolved CO₂, CO₂ gas, and methane, Blake-Bahama Ridge and northeast
637 Bermuda Rise, ODP Leg 172, Proc. Ocean Drill. Program, Sci. Results, 172, 1–16, 2000.
- 638 Borrel, G., Jézéquel, D., Biderre-Petit, C., Morel-Desrosiers, N., Morel, J.-P., Peyret, P., Fonty, G.
639 and Lehours, A.-C.: Production and consumption of methane in freshwater lake ecosystems., Res.
640 Microbiol., 162(9), 832–847, 2011.
- 641 Borrel, G., Lehours, A.-C., Crouzet, O., Jézéquel, D., Rockne, K., Kulczak, A., Duffaud, E., Joblin,
642 K. and Fonty, G.: Stratification of Archaea in the deep sediments of a freshwater meromictic lake:
643 vertical shift from methanogenic to uncultured archaeal lineages., PLoS One, 7(8), e43346, 1–14,
644 2012.
- 645 Borrok, D. M., Wanty, R. B., Ridley, W. I., Wolf, R., Lamothe, P. J. and Adams, M.: Separation of
646 copper, iron, and zinc from complex aqueous solutions for isotopic measurement, Chem. Geol.,
647 242(3-4), 400–414, 2007.
- 648 Burow, L. C., Woebken, D., Marshall, I. P. G., Singer, S. W., Pett-Ridge, J., Prufert-Bebout, L.,
649 Spormann, a M., Bebout, B. M., Weber, P. K. and Hoehler, T. M.: Identification of
650 Desulfobacterales as primary hydrogenotrophs in a complex microbial mat community.,
651 Geobiology, 12(3), 221–230, 2014.
- 652 Chao, A.: Nonparametric Estimation of the Number of Classes in a Population, Scand. J. Stat.,
653 11(4), 265–270, 1984.
- 654 Chao, B. Y. A. and Ma, M.: Stopping rules and estimation for recapture debugging with unequal
655 failure rates, Biometrika, 80(1), 193–201, 1993.
- 656 Chistoserdova, L., Vorholt, J. A. and Lidstrom, M. E.: A genomic view of methane oxidation by
657 aerobic bacteria and anaerobic archaea, Genome Biol., 6(2), 208.1–208.6, 2005.

- 658 Clement, J., Shrestha, J., Ehrenfeld, J. and Jaffe, P.: Ammonium oxidation coupled to dissimilatory
659 reduction of iron under anaerobic conditions in wetland soils, *Soil Biol. Biochem.*, 37(12), 2323–
660 2328, 2005.
- 661 Cline, J. D.: Spectrophotometric determination of hydrogen sulfide In natural waters, *Limnol.*
662 *Oceanogr.*, 14, 454–458, 1969.
- 663 Deutzmann, J. S. and Schink, B.: Anaerobic oxidation of methane in sediments of Lake Constance,
664 an oligotrophic freshwater lake., *Appl. Environ. Microbiol.*, 77(13), 4429–36,.
- 665 Dowd, S. E., Callaway, T. R., Wolcott, R. D., Sun, Y., McKeehan, T., Hagevoort, R. G. and
666 Edrington, T. S.: Evaluation of the bacterial diversity in the feces of cattle using 16S rDNA
667 bacterial tag-encoded FLX amplicon pyrosequencing (bTEFAP)., *BMC Microbiol.*, 8(125), 1–8,
668 2008.
- 669 Eckert, T.: The Influence of Chemical Stratification in the Water Column on Sulfur and Iron
670 Dynamics in Pore Waters and Sediments of Lake Kinneret, Israel, M.Sc. thesis, University of
671 Bayreuth, Germany., 2000.
- 672 Eckert, W. and Conrad, R.: Sulfide and methane evolution in the hypolimnion of a subtropical lake:
673 A three-year study, *Biogeochemistry*, 82(1), 67–76, 2007.
- 674 Edgar, R. C., Haas, B. J., Clemente, J. C., Quince, C. and Knight, R.: UCHIME improves
675 sensitivity and speed of chimera detection., *Bioinformatics*, 27(16), 2194–2200, 2011.
- 676 Egger, M., Rasigraf, O., Sapart, C. J., Jilbert, T., Jetten, M. S. M., Röckmann, T., van der Veen, C.,
677 Banda, N., Kartal, B., Ettwig, K. F. and Slomp, C. P.: Iron-mediated anaerobic oxidation of
678 methane in brackish coastal sediments., *Environ. Sci. Technol.*, doi:10.1021/es503663z, 2014.
- 679 Ehrlich, S., Behrens, D., Lebedeva, E., Ludwig, W. and Bock, E.: A new obligately
680 chemolithoautotrophic, nitrite-oxidizing bacterium, *Nitrospira moscoviensis* sp. nov. and its
681 phylogenetic relationship., *Arch. Microbiol.*, 164(1), 16–23, 1995.
- 682 Ettwig, K. F., Van Alen, T., van de Pas-Schoonen, K. T., Jetten, M. S. M. and Strous, M.:
683 Enrichment and molecular detection of denitrifying methanotrophic bacteria of the NC10 phylum,
684 *Appl. Environ. Microbiol.*, 75(11), 3656–3662, 2009.
- 685 Ettwig, K. F., Butler, M. K., Le Paslier, D., Pelletier, E., Mangenot, S., Kuypers, M. M. M.,
686 Schreiber, F., Dutilh, B. E., Zedelius, J. and de Beer, D.: Nitrite-driven anaerobic methane oxidation
687 by oxygenic bacteria, *Nature*, 464(7288), 543–548, 2010.
- 688 Froelich, P. N., Klinkhammer, G. P., Bender, M. L., Luedtke, N. A., Heath, G. R., Cullen, D.,
689 Dauphin, P., Hammond, D., Hartman, B. and Maynard, V.: Early oxidation of organic matter in
690 pelagic sediments of the eastern equatorial Atlantic: suboxic diagenesis, *Geochim. Cosmochim.*
691 *Acta*, 43(7), 1075–1090, 1979.
- 692 Glissman, K., Chin, K.-J., Casper, P. and Conrad, R.: Methanogenic pathway and archaeal
693 community structure in the sediment of eutrophic Lake Dagow: effect of temperature., *Microb.*
694 *Ecol.*, 48(3), 389–399, 2004.
- 695 Good, I. J.: The population frequencies of species and the estimation of population parameters,
696 *Biometrika*, 40(3-4), 237–264, 1953.

- 697 Hadas, O. and Pinkas, R.: Sulfate-reduction process in sediments of Lake Kinneret, Israel,
698 *Hydrobiologia*, 235(1), 295–301, 1992.
- 699 Hallam, S. J., Girguis, P. R., Preston, C. M., Richardson, P. M. and DeLong, E. F.: Identification of
700 methyl coenzyme M reductase A (*mcrA*) genes associated with methane-oxidizing archaea, *Appl.*
701 *Environ. Microbiol.*, 69(9), 5483–5491, 2003.
- 702 Hallam, S. J., Putnam, N., Preston, C. M., Detter, J. C., Rokhsar, D., Richardson, P. M. and
703 DeLong, E. F.: Reverse methanogenesis: testing the hypothesis with environmental genomics,
704 *Science*, 305(5689), 1457–1462, 2004.
- 705 Hinrichs, K. U., Hayes, J. M., Sylva, S. P., Brewer, P. G. and DeLong, E. F.: Methane-consuming
706 archaeobacteria in marine sediments, *Nature*, 398(6730), 802–805, 1999.
- 707 Hoehler, T. M., Alperin, M. J., Albert, D. B. and Martens, C. S.: Field and laboratory studies of
708 methane oxidation in an anoxic marine sediment: evidence for a methanogen-sulfate reducer
709 consortium, *Global Biogeochem. Cycles*, 8(4), 451–463, 1994.
- 710 Holler, T., Wegener, G., Knittel, K., Boetius, A., Brunner, B., Kuypers, M. M. M. and Widdel, F.:
711 Substantial $^{13}\text{C}/^{12}\text{C}$ and D/H fractionation during anaerobic oxidation of methane by marine
712 consortia enriched in vitro, *Environ. Microbiol. Rep.*, 1(5), 370–376, 2009.
- 713 Hollister, E. B., Engledow, A. S., Hammett, A. J. M., Provin, T. L., Wilkinson, H. H. and Gentry,
714 T. J.: Shifts in microbial community structure along an ecological gradient of hypersaline soils and
715 sediments., *ISME J.*, 4(6), 829–838, 2010.
- 716 Holmkvist, L., Ferdelman, T. G. and Jørgensen, B. B.: A cryptic sulfur cycle driven by iron in the
717 methane zone of marine sediment (Aarhus Bay, Denmark), *Geochim. Cosmochim. Acta*, 75(12),
718 3581–3599, 2011.
- 719 Huse, S. M., Welch, D. M., Morrison, H. G. and Sogin, M. L.: Ironing out the wrinkles in the rare
720 biosphere through improved OTU clustering., *Environ. Microbiol.*, 12(7), 1889–1898, 2010.
- 721 Jetten, M. S. M., Stams, A. J. M. and Zehnder, A. J. B.: Acetate threshold values and acetate
722 activating enzymes in methanogenic bacteria, *FEMS Microbiol. Lett.*, 73(4), 339–344, 1990.
- 723 Jiang, H., Dong, H., Yu, B., Ye, Q., Shen, J., Rowe, H. and Zhang, C.: Dominance of putative
724 marine benthic Archaea in Qinghai Lake, north-western China., *Environ. Microbiol.*, 10(9), 2355–
725 2367, 2008.
- 726 Jiang, L., Zheng, Y., Peng, X., Zhou, H., Zhang, C., Xiao, X. and Wang, F.: Vertical distribution
727 and diversity of sulfate-reducing prokaryotes in the Pearl River estuarine sediments, Southern
728 China., *FEMS Microbiol. Ecol.*, 70(2), 93–106, 2009.
- 729 John, S. G. and Adkins, J. F.: Analysis of dissolved iron isotopes in seawater, *Mar. Chem.*, 119(1-
730 4), 65–76, 2010.
- 731 Kinnaman, F. S., Valentine, D. L. and Tyler, S. C.: Carbon and hydrogen isotope fractionation
732 associated with the aerobic microbial oxidation of methane, ethane, propane and butane, *Geochim.*
733 *Cosmochim. Acta*, 71(2), 271–283, 2007.

- 734 Klein, M., Friedrich, M., Roger, a J., Hugenholtz, P., Fishbain, S., Abicht, H., Blackall, L. L., Stahl,
735 D. a and Wagner, M.: Multiple lateral transfers of dissimilatory sulfite reductase genes between
736 major lineages of sulfate-reducing prokaryotes., *J. Bacteriol.*, 183(20), 6028–6035, 2001.
- 737 Knittel, K. and Boetius, A.: Anaerobic oxidation of methane: progress with an unknown process.,
738 *Annu. Rev. Microbiol.*, 63, 311–334, 2009.
- 739 Koizumi, Y., Takii, S. and Fukui, M.: Depth-related change in archaeal community structure in a
740 freshwater lake sediment as determined with denaturing gradient gel electrophoresis of amplified
741 16S rRNA genes and reversely transcribed rRNA fragments., *FEMS Microbiol. Ecol.*, 48(2), 285–
742 92, doi:10.1016/j.femsec.2004.02.013, 2004.
- 743 Koizumi, Y., Takii, S., Nishino, M. and Nakajima, T.: Vertical distributions of sulfate-reducing
744 bacteria and methane-producing archaea quantified by oligonucleotide probe hybridization in the
745 profundal sediment of a mesotrophic lake., *FEMS Microbiol. Ecol.*, 44(1), 101–108, 2003.
- 746 Lloyd, K. G., Schreiber, L., Petersen, D. G., Kjeldsen, K. U., Lever, M. a., Steen, A. D.,
747 Stepanauskas, R., Richter, M., Kleindienst, S., Lenk, S., Schramm, A. and Jørgensen, B. B.:
748 Predominant archaea in marine sediments degrade detrital proteins, *Nature*, 496(7444), 215–218,
749 2013.
- 750 Lontoh, S., DiSpirito, a a, Krema, C. L., Whittaker, M. R., Hooper, a B. and Semrau, J. D.:
751 Differential inhibition in vivo of ammonia monooxygenase, soluble methane monooxygenase and
752 membrane-associated methane monooxygenase by phenylacetylene., *Environ. Microbiol.*, 2(5), 485–
753 494, 2000.
- 754 Lovley, D.: Dissimilatory Fe (III) - and Mn (IV) -Reducing Prokaryotes Microorganisms, in
755 *Prokaryotes*, pp. 635–658, Springer New York, New York., 2006.
- 756 Lovley, D. R.: Dissimilatory Fe(III) and Mn(IV) reduction., *Microbiol. Rev.*, 55(2), 259–287, 1991.
- 757 Lovley, D. R. and Klug, M. J.: Sulfate Reducers Can Outcompete Methanogens at Concentrations
758 Sulfate Reducers Can Outcompete Methanogens Sulfate Concentrationst at Freshwater, *Appl.*
759 *Environ. Microbiol.*, 45(1), 187–192, 1983.
- 760 Lovley, D. R., Phillips, E. J. P., Lonergan, D. J. and Widman, P. K.: Fe (III) and S₀ reduction by
761 *Pelobacter carbinolicus*, *Appl. Environ. Microbiol.*, 61(6), 2132–2138, 1995.
- 762 McDonald, I. R., Bodrossy, L., Chen, Y. and Murrell, J. C.: Molecular ecology techniques for the
763 study of aerobic methanotrophs., *Appl. Environ. Microbiol.*, 74(5), 1305–15,
764 doi:10.1128/AEM.02233-07, 2008.
- 765 McInerney, M. J., Rohlin, L., Mouttaki, H., Kim, U., Krupp, R. S., Rios-Hernandez, L., Sieber, J.,
766 Struchtemeyer, C. G., Bhattacharyya, A., Campbell, J. W. and Gunsalus, R. P.: The genome of
767 *Syntrophus aciditrophicus*: life at the thermodynamic limit of microbial growth., *Proc. Natl. Acad.*
768 *Sci. U. S. A.*, 104(18), 7600–7605, 2007.
- 769 Milucka, J., Ferdelman, T. G., Polerecky, L., Franzke, D., Wegener, G., Schmid, M., Lieberwirth,
770 I., Wagner, M., Widdel, F. and Kuypers, M. M. M.: Zero-valent sulphur is a key intermediate in
771 marine methane oxidation., *Nature*, 491(7425), 541–546, 2012.

- 772 Miyashita, A., Mochimaru, H., Kazama, H., Ohashi, A., Yamaguchi, T., Nunoura, T., Horikoshi,
773 K., Takai, K. and Imachi, H.: Development of 16S rRNA gene-targeted primers for detection of
774 archaeal anaerobic methanotrophs (ANMEs), *FEMS Microbiol. Lett.*, 297(1), 31–37, 2009.
- 775 Nam, Y.-D., Sung, Y., Chang, H.-W., Roh, S. W., Kim, K.-H., Rhee, S.-K., Kim, J.-C., Kim, J.-Y.,
776 Yoon, J.-H. and Bae, J.-W.: Characterization of the depth-related changes in the microbial
777 communities in Lake Hovsgol sediment by 16S rRNA gene-based approaches., *J. Microbiol.*, 46(2),
778 125–136, 2008.
- 779 Niemann, H., Lösekann, T., de Beer, D., Elvert, M., Nadalig, T., Knittel, K., Amann, R., Sauter, E.
780 J., Schlüter, M. and Klages, M.: Novel microbial communities of the Haakon Mosby mud volcano
781 and their role as a methane sink, *Nature*, 443(7113), 854–858, 2006.
- 782 Nishri, a., Imberger, J., Eckert, W., Ostrovsky, I. and Geifman, Y.: The physical regime and the
783 respective biogeochemical processes in the lower water mass of Lake Kinneret, *Limnol. Oceanogr.*,
784 45(4), 972–981, 2000.
- 785 Nusslein, B., Chin, K. J., Eckert, W. and Conrad, R.: Evidence for anaerobic syntrophic acetate
786 oxidation during methane production in the profundal sediment of subtropical Lake Kinneret
787 (Israel), *Environ. Microbiol.*, 3(7), 460–470, 2001.
- 788 Orcutt, B. N., Sylvan, J. B., Knab, N. J. and Edwards, K. J.: Microbial ecology of the dark ocean
789 above, at, and below the seafloor., *Microbiol. Mol. Biol. Rev.*, 75(2), 361–422, 2011.
- 790 Orphan, V. J., House, C. H., Hinrichs, K. U., McKeegan, K. D. and DeLong, E. F.: Methane-
791 consuming archaea revealed by directly coupled isotopic and phylogenetic analysis., *Science*,
792 293(5529), 484–487, 2001.
- 793 Orphan, V. J., House, C. H., Hinrichs, K. U., McKeegan, K. D. and DeLong, E. F.: Multiple
794 archaeal groups mediate methane oxidation in anoxic cold seep sediments, *Proc. Natl. Acad. Sci.*,
795 99(11), 7663–7668, 2002.
- 796 Pester, M., Schleper, C. and Wagner, M.: The Thaumarchaeota: an emerging view of their
797 phylogeny and ecophysiology., *Curr. Opin. Microbiol.*, 14(3), 300–306, 2011.
- 798 Pjevac, P., Kamyshny, A., Dyksma, S. and Mußmann, M.: Microbial consumption of zero-valence
799 sulfur in marine benthic habitats., *Environ. Microbiol.*, 16(11), 3416–3430, 2014.
- 800 Plugge, C. M., Zhang, W., Scholten, J. C. M. and Stams, A. J. M.: Metabolic flexibility of sulfate-
801 reducing bacteria., *Front. Microbiol.*, 2(May), 1–8, 2011.
- 802 Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J. and Glöckner, F.
803 O.: The SILVA ribosomal RNA gene database project: improved data processing and web-based
804 tools., *Nucleic Acids Res.*, 41(Database issue), D590–D596, 2013.
- 805 Raghoebarsing, A. A., Pol, A., Van de Pas-Schoonen, K. T., Smolders, A. J. P., Ettwig, K. F.,
806 Rijpstra, W. I. C., Schouten, S., Damsté, J. S. S., den Camp, H. J. M. O. and Jetten, M. S. M.: A
807 microbial consortium couples anaerobic methane oxidation to denitrification, *Nature*, 440(7086),
808 918–921, 2006.

- 809 Röske, K., Sachse, R., Scheerer, C. and Röske, I.: Microbial diversity and composition of the
810 sediment in the drinking water reservoir Saidenbach (Saxonia, Germany)., *Syst. Appl. Microbiol.*,
811 35(1), 35–44, 2012.
- 812 Schloss, P. D., Gevers, D. and Westcott, S. L.: Reducing the effects of PCR amplification and
813 sequencing artifacts on 16S rRNA-based studies., *PLoS One*, 6(12), e27310, 2011.
- 814 Schloss, P. D., Westcott, S. L., Ryabin, T., Hall, J. R., Hartmann, M., Hollister, E. B., Lesniewski,
815 R. a, Oakley, B. B., Parks, D. H., Robinson, C. J., Sahl, J. W., Stres, B., Thallinger, G. G., Van
816 Horn, D. J. and Weber, C. F.: Introducing mothur: open-source, platform-independent, community-
817 supported software for describing and comparing microbial communities., *Appl. Environ.*
818 *Microbiol.*, 75(23), 7537–7541, 2009.
- 819 Schoell, M.: Multiple origins of methane in the earth, *Chem. Geol.*, 71(1-3), 1–10, 1988.
- 820 Schubert, C. J., Vazquez, F., Lösekann-Behrens, T., Knittel, K., Tonolla, M. and Boetius, A.:
821 Evidence for anaerobic oxidation of methane in sediments of a freshwater system (Lago di
822 Cadagno)., *FEMS Microbiol. Ecol.*, 76(1), 26–38, 2011.
- 823 Schwarz, J. I. K., Eckert, W. and Conrad, R.: Community structure of Archaea and Bacteria in a
824 profundal lake sediment Lake Kinneret (Israel), *Syst. Appl. Microbiol.*, 30(3), 239–254, 2007a.
- 825 Schwarz, J. I. K., Lueders, T., Eckert, W. and Conrad, R.: Identification of acetate-utilizing Bacteria
826 and Archaea in methanogenic profundal sediments of Lake Kinneret (Israel) by stable isotope
827 probing of rRNA., *Environ. Microbiol.*, 9(1), 223–237, 2007b.
- 828 Serruya, C.: Lake Kinneret: the nutrient chemistry of the sediments, *Limnol. Oceanogr.*, 16(May),
829 510–521, 1971.
- 830 Serruya, C., Edelstein, M., Pollinger, U. and Serruya, S.: Lake Kinneret sediments: nutrient
831 composition of the pore water and mud water exchanges, *Limnol. Oceanogr.*, 19(3), 489–508, 1974.
- 832 Severmann, S., Johnson, C. M., Beard, B. L. and McManus, J.: The effect of early diagenesis on the
833 Fe isotope compositions of porewaters and authigenic minerals in continental margin sediments,
834 *Geochim. Cosmochim. Acta*, 70(8), 2006–2022, 2006.
- 835 Sivan, O., Adler, M., Pearson, A., Gelman, F., Bar-Or, I., John, S. G. and Eckert, W.: Geochemical
836 evidence for iron-mediated anaerobic oxidation of methane, *Limnol. Oceanogr.*, 56(4), 1536–1544,
837 2011.
- 838 Smith, P. F., Langworthy, T. A. and Smith, M. R.: Polypeptide nature of growth requirement in
839 yeast extract for *Thermoplasma* Polypeptide Nature of Growth Requirement in Yeast Extract for
840 *Thermoplasma acidophilum*, *J. Bacteriol.*, 124(2), 884–892, 1975.
- 841 Sonne-Hansen, J. and Ahring, B. K.: *Thermodesulfobacterium hveragerdense* sp. nov., and
842 *Thermodesulfobacterium islandicus* sp. nov., two thermophilic sulfate reducing bacteria isolated from a
843 Icelandic hot spring., *Syst. Appl. Microbiol.*, 22(4), 559–564, 1999.
- 844 Sørensen, K. B. and Teske, A.: Stratified communities of active Archaea in deep marine subsurface
845 sediments., *Appl. Environ. Microbiol.*, 72(7), 4596–4603, 2006.

- 846 Spring, S., Amann, R., Ludwig, W., Schleifer, K. H., van Gemerden, H. and Petersen, N.:
847 Dominating role of an unusual magnetotactic bacterium in the microaerobic zone of a freshwater
848 sediment., *Appl. Environ. Microbiol.*, 59(8), 2397–2403, 1993.
- 849 Stookey, L. L.: Ferrozine-a new spectrophotometric reagent for iron, *Anal. Chem.*, 42(7), 779–781,
850 1970.
- 851 Sun, H., Spring, S., Lapidus, A., Davenport, K., Del Rio, T. G., Tice, H., Nolan, M., Copeland, A.,
852 Cheng, J.-F., Lucas, S., Tapia, R., Goodwin, L., Pitluck, S., Ivanova, N., Pagani, I., Mavromatis, K.,
853 Ovchinnikova, G., Pati, A., Chen, A., Palaniappan, K., Hauser, L., Chang, Y.-J., Jeffries, C. D.,
854 Detter, J. C., Han, C., Rohde, M., Brambilla, E., Göker, M., Woyke, T., Bristow, J., Eisen, J. a,
855 Markowitz, V., Hugenholtz, P., Kyrpides, N. C., Klenk, H.-P. and Land, M.: Complete genome
856 sequence of *Desulfarculus baarsii* type strain (2st14)., *Stand. Genomic Sci.*, 3(3), 276–284, 2010.
- 857 Tavormina, P. L., Orphan, V. J., Kalyuzhnaya, M. G., Jetten, M. S. M. and Klotz, M. G.: A novel
858 family of functional operons encoding methane/ammonia monooxygenase-related proteins in
859 gammaproteobacterial methanotrophs, *Environ. Microbiol. Rep.*, 3(1), 91–100, 2011.
- 860 Tavormina, P. L., Ussler, W. and Orphan, V. J.: Planktonic and sediment-associated aerobic
861 methanotrophs in two seep systems along the North American margin., *Appl. Environ. Microbiol.*,
862 74(13), 3985–2995, 2008.
- 863 Thauer, R. K.: Functionalization of methane in anaerobic microorganisms., *Angew. Chem. Int. Ed.*
864 *Engl.*, 49(38), 6712–6713, 2010.
- 865 Valentine, D. L.: Biogeochemistry and microbial ecology of methane oxidation in anoxic
866 environments: a review, *Antonie Van Leeuwenhoek*, 81(1-4), 271–282, 2002.
- 867 Whiticar, M. J.: Carbon and hydrogen isotope systematics of bacterial formation and oxidation of
868 methane, *Chem. Geol.*, 161(1-3), 291–314, 1999.
- 869 Wobus, A., Bleul, C., Maassen, S., Scheerer, C., Schuppler, M., Jacobs, E. and Röske, I.: Microbial
870 diversity and functional characterization of sediments from reservoirs of different trophic state.,
871 *FEMS Microbiol. Ecol.*, 46(3), 331–347, 2003.
- 872 Yamada, C., Kato, S., Kimura, S., Ishii, M. and Igarashi, Y.: Reduction of Fe(III) oxides by
873 phylogenetically and physiologically diverse thermophilic methanogens., *FEMS Microbiol. Ecol.*,
874 89(3), 637–645, 2014.
- 875 Yan, Z., Song, N., Cai, H., Tay, J.-H. and Jiang, H.: Enhanced degradation of phenanthrene and
876 pyrene in freshwater sediments by combined employment of sediment microbial fuel cell and
877 amorphous ferric hydroxide., *J. Hazard. Mater.*, 199-200, 217–225, 2012.
- 878 Ye, W., Liu, X., Lin, S., Tan, J., Pan, J., Li, D. and Yang, H.: The vertical distribution of bacterial
879 and archaeal communities in the water and sediment of Lake Taihu., *FEMS Microbiol. Ecol.*, 70(2),
880 107–20, 2009.
- 881 Yoshida, N., Takahashi, N. and Hiraishi, A.: Phylogenetic Characterization of a Polychlorinated-
882 Dioxin- Dechlorinating Microbial Community by Use of Microcosm Studies, , 71(8), 4325–4334,
883 2005.

884 Zhu, B., van Dijk, G., Fritz, C., Smolders, A. J. P., Pol, A., Jetten, M. S. M. and Ettwig, K. F.:
 885 Anaerobic oxidization of methane in a minerotrophic peatland: enrichment of nitrite-dependent
 886 methane-oxidizing bacteria., *Appl. Environ. Microbiol.*, 78(24), 8657–8665, 2012.

887

888 Tables

889 **Table 1.** Bacterial sequences used for classification in SILVA ngs and for subsample in MOTHUR for the
 890 alpha diversity.

Bacteria	# seq SILVA ngs	OTUs SILVA ngs	# seq MOTHUR	OTUs MOTHUR	coverage	invsimpson	chao	ace	shannon
0-3 cm	3631	599	2588	388	0.92	10/10.8/11.7	805/925.7/1092.8	1246.5/1384.0/1546.7	3.83/3.91/3.98
6-9 cm	4641	1735	3365	1337	0.77	286.6/315.9/351.8	2437.9/2673.0/2958.4	3876.8/4135.2/4420.0	6.48/6.53/6.57
29-32 cm	5038	1516	3615	1214	0.81	127.8/142.4/160.6	2104.4/2318.7/2583.1	3047.7/3259.5/3495.4	6.13/6.19/6.24

892 **Table 2.** Archaeal sequences used for classification in SILVA ngs and for subsample in MOTHUR for the
 893 alpha diversity.

Archaea	# seq SILVA ngs	OTUs SILVA ngs	# seq MOTHUR	OTUs MOTHUR	coverage	invsimpson	chao	ace	shannon
0-3 cm	288	48	268	32	0.90	2.5/2.9/3.5	53.9/83.4/164.5	232.8/344.6/520.4	1.6/1.8/2.0
6-9 cm	1382	184	1408	200	0.86	8.4/10.2/13.2	39.0/131.4/220.4	144.7/195.2/277.4	2.87/3.05/3.22
29-32 cm	3110	383	2829	245	0.88	7.9/9.2/10.9	73.5/104.8/181.7	133.2/190.9/289.6	2.58/2.74/2.91

895 Figure Captions:

896 **Figure 1:** Schematic location of Lake Kinneret. Numbers correspond to altitude measured in meters
 897 from the level of Lake Kinneret. Cores were taken from station A (taken from Hambright et al.,
 898 2004).

899 **Figure 2:** Geochemical porewater profiles in LK sediment. **A.** profile of SO_4^{-2} (black circles) and
 900 sulfide (gray diamonds) in the porewater **B.** Headspace measurements of methane (gray circles) and
 901 $\delta^{13}\text{C}_{\text{CH}_4}$ (black squares) in the sediments **C.** $\delta^{13}\text{C}$ of total lipids extraction (TLE) (black diamonds)
 902 from the sediment, and $\delta^{56}\text{Fe}$ (gray circles) of the dissolved iron in the porewater. **D.** Electron
 903 acceptor profiles of dissolved Fe(II) (gray triangles) and Mn (II) (black squares) in the porewater.,
 904 Black arrows indicate the sampled sections for 16S rRNA gene analysis. The presented methane
 905 profile was taken two weeks prior to the sampling for the microbial communities on December
 906 2009. Sampling for Fe(II), Mn(II) and SO_4^{-2} profiles was carried out between 2007 and 2011 and
 907 for sulfide from 2007 to 2013. The $\delta^{13}\text{C}_{\text{CH}_4}$ and $\delta^{56}\text{Fe}$ profiles were performed four months prior to
 908 the microbial sampling on August 2009.


909 **Figure 3:** Classification of bacterial sequences using SILVA ngs pipeline. Phyla and order
 910 distribution of sequences of the 454 sequencing at the different depths.

911 **Figure 4:** Classification of archaeal sequences using SILVA ngs pipeline. Order distributions of
912 sequences of the 454 sequencing at the different depths.

913 **Figure 5:** Principal coordinate analysis (PCoA) of the **A.** bacterial and **B.** archaeal community
914 matrix (based on operational taxonomic units) and vector fitting of the environmental geochemical
915 conditions in the different depth. The bacterial community from the upper layer is labeled with red
916 triangle, the middle layer is labeled with green triangle and bottom layer is labeled with blue
917 triangle.

918 **Figure 6:** profiles of functional genes from the 3 different depths. Green bar represent *pmoA*
919 functional gene of aerobic methanotroph. Red bar represent *dsrAB* functional gene of sulfate
920 reducers and blue bar represent *mcrA* functional gene from methanogens and anaerobic
921 methanotrophs.

922



923

924


925

926

927

928

Figure 1.




929
930

931

932

Figure 2.



933


934

935

936

Figure 3.

937




938


939

Figure 4.

940



941




942

943

Figure 5.

944

945



946

947

948

949


950

Figure 6

951

952

953



955


956

957

Figure 1

958 **Supplementary Fig. 1.** Venn diagram showing archaeal sequences overlap between the different
959 depths. The green circle represents the top sample (0-3 cm), the blue circle represents the middle
960 sample (6-9 cm) and the red circle represents the bottom sample (29-32 cm).

961



962

963

Figure 2

964 **Supplementary Fig. 2.** Venn diagram showing bacterial sequences overlap between the different
965 depths. The green circle represents the top sample (0-3 cm), the blue circle represents the middle
966 sample (6-9 cm) and the red circle represents the bottom sample (29-32 cm).

967

968 **Supplementary Table 1.** The classification percentage and number of sequences of archaea A.
 969 phylum, B. class and C. order by SILVA ngs.

Archaea	0-3 cm		6-9 cm		29-32 cm	
Phylum	# seq	%	# seq	%	# seq	%
Euryarchaeota	283	98.3	1430	96.4	3048	98.0
Thaumarchaeota	5	1.7	53	3.6	62	2.0

Archaea	0-3 cm		6-9 cm		29-32 cm	
Class	# seq	%	# seq	%	# seq	%
AK59					1	0.0
AK8			1	0.1	4	0.1
Group C3			2	0.1	2	0.1
Halobacteria	7	2.4	96	6.5	165	5.3
Marine Benthic Group B	2	0.7	24	1.6	30	1.0
Marine Group I					2	0.1
Methanobacteria	1	0.3				
Methanomicrobia	266	92.4	1164	78.5	2679	86.1
Miscellaneous Crenarchaeotic Group	3	1.0	21	1.4	19	0.6
pSL12			2	0.1	3	0.1
South African Gold Mine Gp 1(SAGMCG-1)			3	0.2	1	0.0
Thermoplasmata	9	3.1	170	11.5	204	6.6

Archaea	0-3 cm		6-9 cm		29-32 cm	
Order	# seq	%	# seq	%	# seq	%
20a-9			5	0.3	3	0.1
ANME-1					1	0.0
Cenarchaeales					2	0.1
Halobacteriales	7	2.4	96	6.5	165	5.3
Methanobacteriales	1	0.3				
Methanomicrobiales	202	70.1	422	28.5	878	28.2
Methanosarcinales	64	22.2	742	50.0	1800	57.9
Thermoplasmatales	8	2.8	165	11.1	201	6.5
WCHA1-57	1	0.3				
Uncultured	5	1.7	53	3.6	60	1.9

973 **Supplementary Table 2.** The classification percentage and number of sequences of bacterial A.
 974 phylum, B. class and C. order by SILVA ngs.

Bacteria	0-3 cm		6-9 cm		29-32 cm	
Phylum	# seq	%	# seq	%	# seq	%
Acidobacteria	33	0.9	112	2.4	123	2.4
Actinobacteria	56	1.5	61	1.3	90	1.8
Armatimonadetes	21	0.6	12	0.3	14	0.3
Bacteroidetes	80	2.2	349	7.5	126	2.5
BD1-5	2	0.1	2	0.0	2	0.0
Caldiserica					2	0.0
Candidate division BRC1	2	0.1	14	0.3	8	0.2
Candidate division JS1	1	0.0	2	0.0		
Candidate division KB1					3	0.1
Candidate division OD1			1	0.0	18	0.4
Candidate division OP11	2	0.1	17	0.4	1	0.0
Candidate division OP3	1	0.0	108	2.3	84	1.7
Candidate division OP8	9	0.2	81	1.7	106	2.1
Candidate division SR1			4	0.1		
Candidate division TM7	6	0.2	5	0.1		
Candidate division WS3	15	0.4	70	1.5	103	2.0
Chlorobi	38	1.0	235	5.1	206	4.1
Chloroflexi	65	1.8	672	14.5	969	19.2
CKC4	1	0.0				
Cyanobacteria	18	0.5	69	1.5	21	0.4
Deferribacteres	1	0.0	122	2.6	126	2.5
Deinococcus-Thermus	1	0.0	8	0.2	9	0.2
Elusimicrobia			7	0.2	13	0.3
Fibrobacteres			20	0.4	9	0.2
Firmicutes	1376	37.9	140	3.0	152	3.0
Gemmatimonadetes	2	0.1	15	0.3	15	0.3
Hyd24-12	4	0.1	8	0.2		
GOUTA4					4	0.1
JL-ETNP-Z39			1	0.0	3	0.1
Lentisphaerae	12	0.3	38	0.8	10	0.2
Nitrospirae	31	0.9	195	4.2	409	8.1
NPL-UPA2	1	0.0	33	0.7	5	0.1
OC31			1	0.0		
Planctomycetes	22	0.6	140	3.0	172	3.4
Proteobacteria	1762	48.5	1837	39.6	1954	38.8
SHA-109	6	0.2	6	0.1	1	0.0
Spirochaetae	1	0.0	102	2.2	134	2.7
Synergistetes					1	0.0
TA06	5	0.1	66	1.4	102	2.0
Tenericutes	13	0.4	1	0.0	1	0.0
Thermotogae			8	0.2	1	0.0
TM6	19	0.5	27	0.6	21	0.4
Verrucomicrobia	3	0.1	39	0.8	20	0.4
WCHB1-60	22	0.6	13	0.3		

Bacteria	0-3 cm		6-9 cm		29-32 cm	
	# seq	%	# seq	%	# seq	%
Acidimicrobiia	5	0.1	4	0.1	2	0.0
Acidobacteria	26	0.7	80	1.7	83	1.6
Actinobacteria	7	0.2	5	0.1	5	0.1
Alphaproteobacteria	23	0.6	25	0.5	50	1.0
Anaerolineae	31	0.9	296	6.4	375	7.4
Ardenticatenia			1	0.0		
ARKDMS-49			2	0.0		
ARKICE-90			11	0.2		
Bacilli	611	16.8	23	0.5	49	1.0
Bacteroidia			5	0.1	1	0.0
BD2-2	2	0.1	16	0.3	8	0.2
Betaproteobacteria	65	1.8	268	5.8	344	6.8
BS5	3	0.1	7	0.2		
BSV13			9	0.2	1	0.0
Caldilineae	1	0.0	8	0.2	3	0.1
Caldisericia					2	0.0
Chlorobia	6	0.2	42	0.9	18	0.4
Chloroplast	3	0.1	42	0.9	4	0.1
Clostridia	761	21.0	116	2.5	99	2.0
Coriobacteriia	3	0.1	4	0.1	7	0.1
Cyanobacteria	13	0.4	22	0.5	7	0.1
Cytophagia	1	0.0	4	0.1	8	0.2
Deferribacteres	1	0.0	122	2.6	126	2.5
Dehalococcoidia	2	0.1	276	5.9	453	9.0
Deinococci	1	0.0	8	0.2	9	0.2
Deltaproteobacteria	57	1.6	1257	27.1	1022	20.3
DEV055	6	0.2	9	0.2		
Elusimicrobia			7	0.2	13	0.3
Epsilonproteobacteria			12	0.3	1	0.0
Erysipelotrichia					3	0.1
FFCH16263	1	0.0			2	0.0
Fibrobacteria			20	0.4	9	0.2
Flavobacteriia	9	0.2	34	0.7	3	0.1
Gammaproteobacteria	1614	44.5	257	5.5	536	10.6
Gemmatimonadetes	2	0.1	15	0.3	15	0.3
Holophagae	5	0.1	24	0.5	29	0.6
Ignavibacteria	32	0.9	193	4.2	188	3.7
JG30-KF-CM66			12	0.3	6	0.1
KD4-96	9	0.2	5	0.1	4	0.1
Ktedonobacteria	1	0.0				
LD1-PB3			1	0.0	1	0.0
Lentisphaeria	1	0.0	3	0.1		
MB-A2-108	6	0.2	4	0.1	14	0.3
MBMPE71			1	0.0	1	0.0
Melainabacteria	2	0.1	1	0.0	4	0.1
ML635J-21			4	0.1	6	0.1
Mollicutes	13	0.4	1	0.0	1	0.0
MSB-5B2					4	0.1
MSBL3			6	0.1		
Negativicutes	4	0.1	1	0.0	1	0.0
Nitrospira	31	0.9	195	4.2	409	8.1
OM190	1	0.0	4	0.1	2	0.0
Oligosphaeria			7	0.2	5	0.1
OPB35 soil group	3	0.1	25	0.5	11	0.2
OPB41	20	0.6	21	0.5	35	0.7
Opiritae			12	0.3	7	0.1
PBS-III-20	2	0.1	4	0.1		
Phycisphaerae	11	0.3	101	2.2	130	2.6
Pla3 lineage			10	0.2	7	0.1
Pla4 lineage			9	0.2	8	0.2
Planctomycetacia	10	0.3	15	0.3	19	0.4
Proteobacteria Incertae Sedis			1	0.0	1	0.0
S085					1	0.0
SB-1			7	0.2	10	0.2
SB-5	19	0.5	33	0.7	12	0.2
S-BQ2-57 soil group					1	0.0
SHA-26			7	0.2	14	0.3
Spartobacteria			1	0.0	1	0.0
Sphingobacteriia	30	0.8	95	2.0	30	0.6
Spirochaetes	1	0.0	102	2.2	134	2.7
Subgroup 22	2	0.1	8	0.2	11	0.2
Synergistia					1	0.0
TA18	3	0.1	4	0.1		
Thermoleophilia	14	0.4	23	0.5	25	0.5
Thermotogae			8	0.2	1	0.0
TK10					2	0.0
vadinHA17	18	0.5	145	3.1	51	1.0
vadinHA49					5	0.1
VC2.1 Bac22			1	0.0		
Verrucomicrobiae			1	0.0		
WCHB1-41			1	0.0	2	0.0
WCHB1-32	1	0.0			1	0.0
uncultured	138	3.8	538	11.6	585	11.6

Bacteria	0-3 cm		6-9 cm		29-32 cm	
Order	# seq	%	# seq	%	# seq	%
10bav-F6					2	0.0
43F-1404R			2	0.0	3	0.1
Acidimicrobiales	5	0.1	4	0.1	2	0.0
Alteromonadales			9	0.2	14	0.3
Amsterdam-1B-07			1	0.0	4	0.1
Anaerolineales	31	0.9	296	6.4	375	7.4
B1-7BS			1	0.0		
B276-D12			1	0.0		
Bacillales	602	16.6	23	0.5	47	0.9
Bacteroidales			5	0.1	1	0.0
BD2-11 terrestrial group			2	0.0	5	0.1
Bdellovibrionales			34	0.7	13	0.3
BP-U1C-1g10			1	0.0		
Burkholderiales	56	1.5	86	1.9	208	4.1
C86					2	0.0
Caldilineales	1	0.0	8	0.2	3	0.1
Caldisericales					2	0.0
Campylobacterales			12	0.3	1	0.0
Caulobacterales	12	0.3	1	0.0	1	0.0
CCM11a			9	0.2	6	0.1
Chlorobiales	6	0.2	42	0.9	18	0.4
Chromatiales	3	0.1	11	0.2	21	0.4
Chthoniobacterales			1	0.0	1	0.0
Clostridiales	760	20.9	107	2.3	99	2.0
Coriobacteriales	3	0.1	4	0.1	7	0.1
Corynebacteriales	1	0.0	1	0.0	3	0.1
Cytophagales	1	0.0	3	0.1		
D8A-2	1	0.0				
Deferribacterales	1	0.0	122	2.6	126	2.5
Dehalococcoidales			6	0.1	2	0.0
Deltaproteobacteria Incertae Sedis	2	0.1	55	1.2	2	0.0
Desulfarculales	12	0.3	132	2.8	62	1.2
Desulfobacterales	13	0.4	118	2.5	88	1.7
Desulfuromonadales			80	1.7	41	0.8
Erysipelotrichales					3	0.1
Fibrobacterales			18	0.4	6	0.1
Flavobacteriales	9	0.2	34	0.7	3	0.1
Frankiales	1	0.0	2	0.0	3	0.1
FS117-23B-02			2	0.0	4	0.1
FW22	1	0.0	2	0.0	2	0.0
Gaiellales	11	0.3	19	0.4	20	0.4
Gammaproteobacteria Incertae Sedis			4	0.1	4	0.1
Gastranaerophilales			1	0.0	4	0.1
Gemmatimonadales	2	0.1	13	0.3	10	0.2
GIF3			2	0.0	7	0.1
GIF9			117	2.5	251	5.0

GR-WP33-30	1	0.0	14	0.3	34	0.7
Haloplasmales	13	0.4				
HOC36	2	0.1	72	1.6	267	5.3
Holophagae Incertae Sedis			5	0.1	12	0.2
Holophagales			2	0.0	2	0.0
Hydrogenophilales			48	1.0	44	0.9
Ignavibacteriales	32	0.9	193	4.2	188	3.7
KCLunmb-38-53	1	0.0	9	0.2		
KD3-62	1	0.0	8	0.2	9	0.2
KI89A clade			4	0.1	5	0.1
Lactobacillales	2	0.1			2	0.0
Legionellales	5	0.1	17	0.4	15	0.3
Lineage IIb					4	0.1
Lineage IIc			5	0.1	4	0.1
Lineage IV			1	0.0	1	0.0
MD2894-B20			1	0.0	3	0.1
Methylococcales	9	0.2	70	1.5	69	1.4
Methylophilales			27	0.6	10	0.2
Micrococcales	1	0.0	2	0.0	1	0.0
mle1-8	6	0.2	5	0.1	8	0.2
MSB-3A7 sediment group			3	0.1	3	0.1
MSBL8			1	0.0		
MSBL5	1	0.0	139	3.0	169	3.4
MSBL9	1	0.0	22	0.5	59	1.2
MVP-21	1	0.0				
MVP-88					1	0.0
Myxococcales	1	0.0	82	1.8	45	0.9
NB1-n			1	0.0	1	0.0
Nitrosomonadales	2	0.1	36	0.8	22	0.4
Nitrospirales	31	0.9	195	4.2	409	8.1
ODP1230B30.09					1	0.0
Oceanospirillales			1	0.0		
Oligosphaerales			7	0.2	5	0.1
Opitutales			12	0.3	7	0.1
Order II (Bacteroidetes)			1	0.0	5	0.1
Order Incertae Sedis			2	0.0		
PBS-18			3	0.1	1	0.0
PeM15	1	0.0				
Phycisphaerales	2	0.1	10	0.2	10	0.2
Pla1 lineage			18	0.4	17	0.3
Planctomycetales	10	0.3	15	0.3	19	0.4
Propionibacteriales	3	0.1				
possible order 07			2	0.0	3	0.1
Pseudomonadales	1523	41.9	1	0.0		
Rhizobiales	6	0.2	16	0.3	45	0.9
Rhodobacterales	1	0.0	1	0.0		
Rhodocyclales	4	0.1	43	0.9	29	0.6
Rhodospirillales			3	0.1	1	0.0
Rickettsiales	3	0.1	4	0.1		

S15A-MN16	1	0.0	18	0.4	14	0.3
S-70			3	0.1		
SAR324 clade(Marine group B)			2	0.0		
SC-I-84	2	0.1	27	0.6	31	0.6
Sh765B-AG-111					2	0.0
Sh765B-TzT-29			1	0.0	3	0.1
SHA-43			2	0.0	6	0.1
Selenomonadales	4	0.1	1	0.0	1	0.0
Solirubrobacterales	3	0.1	4	0.1	5	0.1
Sphingobacteriales	30	0.8	95	2.0	30	0.6
Sphingomonadales	1	0.0			3	0.1
Spirochaetales	1	0.0	98	2.1	133	2.6
Streptomycetales					1	0.0
Subgroup 17	1	0.0	3	0.1		
Subgroup 18	3	0.1	27	0.6	21	0.4
Subgroup 23	2	0.1	6	0.1	7	0.1
Subgroup 4	2	0.1				
Subgroup 6	20	0.6	31	0.7	25	0.5
Subgroup 7	2	0.1	3	0.1		
Subgroup 13			5	0.1	5	0.1
Subgroup 19			1	0.0	13	0.3
Subgroup 2			1	0.0	6	0.1
Subgroup 21			9	0.2		
Subgroup 25			1	0.0	6	0.1
Subgroup 3			1	0.0	1	0.0
Subgroup 11					1	0.0
Subgroup 9					1	0.0
SubsectionII (Cyanobacteria)					1	0.0
SubsectionI (Cyanobacteria)	6	0.2	8	0.2	3	0.1
SubsectionIV (Cyanobacteria)	7	0.2	14	0.3	3	0.1
Sva0485	20	0.6	390	8.4	546	10.8
Synergistales					1	0.0
Syntrophobacterales	8	0.2	347	7.5	183	3.6
SZB30	4	0.1	9	0.2	7	0.1
Thermoanaerobacterales			9	0.2		
Thermotogales			8	0.2	1	0.0
Thiotrichales	3	0.1	5	0.1	3	0.1
TPD-58	1	0.0	6	0.1	8	0.2
TRA3-20	1	0.0				
vadinBA26			8	0.2	16	0.3
Verrucomicrobiales			1	0.0		
Vampirovibrionales	2	0.1				
VAN12	7	0.2				
Victivallales	1	0.0	3	0.1		
WD2101 soil group			1	0.0		
Xanthomonadales	65	1.8	50	1.1	129	2.6
uncultured	237	6.5	951	20.5	816	16.2

978 **Appendix**

979 The major bacterial populations in the sediment:

980 *Proteobacteria*

981 *Proteobacteria* was the most dominant bacterial phylum in all the depths. Their percentage was
982 slowly decreases with depth however the composition within its classes were changing with depth.
983 *Proteobacteria* is also among the most abundant phyla in LK (Schwarz et al., 2007a) and in other
984 freshwater sediments (Tamaki et al., 2005; Wobus et al., 2003). *Proteobacteria* is a very diverse
985 phylum, divided in to number of classes which most of them have been observed in LK sediments.

986 *Betaproteobacteria*

987 *Betaproteobacteria* percentages were slowly increasing with depth. They are comprised of
988 chemoheterotrophs and chemoautotrophs which derive nutrients from decomposition of organic
989 material. *Burkholderiales* was the dominant order in *Betaproteobacteria* and increased with depth
990 (1.5% to 4%). Schwarz et al. (2007a) showed that *Burkholderiales* is active in the upper sediment of
991 LK. Members of this group are able of anaerobic oxidation of acetate with (per)chlorate as electron
992 acceptor (Yoshida et al., 2005) or oxidize hydrogen in deep subsurface(Orcutt et al., 2011).
993 *Hydrogenophilales* and *Nitrosomonadales* orders were observed in lower abundant but in the
994 middle and deep layer of the sediment. *Hydrogenophilales* can utilize hydrogen (Ontiveros-
995 Valencia et al., 2013) and *Hydrogenophilaceae* family are obligatory chemolithoautotrophic,
996 aerobic or facultative anaerobic sulfur oxidizers and reduction of nitrate (Kelly and Wood, 2000)
997 which are often found also in wastewater treatment systems(Luo et al., 2011). *Nitrosomonadales*
998 are freshwater bacteria which presumed to have a role in ammonia oxidation and/or denitrification
999 processes (Edlund et al., 2008). McBeth et al. (2013) also showed that *Nitrosomonadales* are
1000 commonly found as freshwater Fe oxidizers. It should be noted that the upper sediment may
1001 contain denitrifying bacteria that arrived from the water column where nitrification conditions exists
1002 throughout the year.

1003 *Gammaproteobacteria*

1004 Our sequences which were affiliated to the *Gammaproteobacteria* have different phylogenetic
1005 classification distribution in each layer. The upper layer was dominated by *Pseudomonadales* (42%)
1006 which were divided to 2 main genera: *Acinetobacter* (28%) and *Pseudomonas* (14%). *Acinetobacter*
1007 was shown to have the ability to degrade organic matter. Our sequences which similar to
1008 *Acinetobacter* genus were closest to *Acinetobacter lwoffii* (99%) found in sewage and growndwater
1009 (Nazina et al., 2000). our sequences classified as *Pseudomonas* were similar to *Pseudomonas*
1010 *stutzeri* (99%) which has the ability to oxidize a large variety of organic compounds, and some

1011 members also involved in nitrification and denitrification, degradation of aromatic compounds, and
1012 nitrogen fixation (Chen et al., 2011). In the middle and bottom layers *Methylococcales* order was
1013 observed (~1%). *Methylococcales* are aerobic methanotrophs which some members were present
1014 within the sediments and overlying water column from dysoxic, methane-rich vent and seep
1015 systems (Tavormina et al., 2008). The most abundant order in the deep layer was HOC36 (5%)
1016 which is an uncultured *Gammaproteobacteria*. however when compared to NCBI data base it was
1017 closely related to uncultured LK clones (99%) and to cultured *Methylocaldum* sp. (94%) (Bodrossy
1018 et al., 1997) which was a Thermophilic methanotroph isolated from landfill cover soil.

1019 *Deltaproteobacteria*

1020 *Deltaproteobacteria* were the most abundant in the middle and deep sediment samples. In the
1021 middle layer the abundant classes were *Syntrophobacterales* (8%), *Desulfarculales* (3%),
1022 *Desulfobacterales* (2%) and *Desulfuromonadales* (2%). Many members of the families of
1023 *Syntrophobacterales* are sulfate reducers or as fermentative organism, depending on the
1024 environmental conditions (McInerney et al., 2007). *Syntrophaceae* the dominant family of
1025 *Syntrophobacterales* are acetate degrading sulfate reducing bacterium (Jiang et al., 2009).
1026 *Desulfarculales* are strictly anaerobic are mesophilic sulfate-reducing bacterium with the capability
1027 to oxidize acetate and fatty acids (Sun et al., 2010) . *Desulfobacterales* are sulfate reducers capable
1028 to oxidize hydrogen in soils and sediments (Burow et al., 2014). *Desulfuromonadales* was shown as
1029 S⁰ respiring (Pjevac et al., 2014) and may also reduce Fe(III) and Mn(IV) in marine surface
1030 sediments (Lovley, 2006). About 1% of the sequences in the middle and bottom layers were
1031 affiliated to *Geobacter* genus which involve in iron reduction. In the deep layer the most abundant
1032 class was Sva0485. Sequences of the SVa485 order were similar to those found in different aquatic
1033 environments, but the metabolic functions of members of this order are not clear. *Pelobacter*
1034 *carbinolious* (Lovley et al., 1995), a member of SVa485 capable of Fe(III) and sulfur reduction.
1035 Schwarz et al. (2007a) showed in the upper part of LK sediment that *Deltaproteobacteria* was one
1036 of the dominant classes, most of which were affiliated with acetate oxidizing sulfate reducing
1037 bacteria, which outcompete the acetoclastic methanogens. Even though *Deltaproteobacteria* are
1038 best known for their sulfate reduction metabolism, they can shift their metabolism in response to
1039 depleted sulfate concentrations (Plugge et al., 2011).

1040 *Chloroflexi*

1041 *Chloroflexi* was found to be one of the most abundant bacterial phyla in LK and represented diverse
1042 functional groups that increased with depth. The abundant classes that increase with depth were
1043 *Anaerolineae* and *Dehalococcoidia*. Although they are found in a variety of anaerobic

1044 environments, only a few representatives have been cultivated so far (Yamada and Sekiguchi,
1045 2009). The metabolisms of the cultured representatives showed that they utilized organic matter, but
1046 no specific electron acceptor has as yet been reported (Yamada et al., 2006). *Chloroflexi* was one of
1047 the most dominant phyla in contaminated soil environment which had a lot of polycyclic aromatic
1048 hydrocarbons (Winderl et al., 2008). Members of *Chloroflexi* were found throughout the entire core,
1049 indicating that *Anaerolineae* representatives could be the main aromatic organic matter
1050 decomposers in LK. *Dehalococcoidia* are widely distributed in the marine subsurface, yet metabolic
1051 properties of the many uncultivated lineages are completely unknown. *Dehalococcoidia* could be
1052 capable of oxidizing various fatty acids and/or structurally related substrates (Wasmund et al.,
1053 2013) or reductive dechlorination of chlorinated hydrocarbon (Maymo et al., 1995).

1054 *Nitrospirae*

1055 *Nitrospirae* phylum, were increased with depth. Sequences from our samples were classified to
1056 *Nitrospirales* order in *Nitrospirea* which were most abundant in the bottom layer of the sediment
1057 (7%). Part of our sequences were similar to previous study (Schwarz et al., 2007b). (Schwarz et al.,
1058 2007b) showed by SIP-RNA not only that *Nitrospirae* was present, but also that it was functionally
1059 bioactive. Although *Nitrospirae* is a known nitrate oxidizer (Ehrich et al., 1995), the conditions of
1060 this environment suggest that it utilized another metabolic pathway. *Nitrospirae* also include the
1061 iron reducing candidates *Magnetobacterium bavaricum* (Spring et al., 1993) and sulfur reducers
1062 (Sonne-Hansen and Ahring, 1999). Part of our sequences were aligned to uncultured clone (98%)
1063 from freshwater sediment which enhanced degradation of phenanthrene and pyrene by amorphous
1064 ferric hydroxide (Yan et al., 2012).

1065 *Chlorobi*

1066 *The Chlorobi*, green sulfur bacteria, are a group of anoxygenic photosynthetic bacteria observed
1067 frequently in anoxic microbial mats and meromictic lakes. *Chlorobi* in found in deeper layers of the
1068 sediment since they support higher H₂S concentrations and require less light intensity. They
1069 photosynthesize using H₂S as electron donor, although they can also grow photoheterotrophically,
1070 and some of them form symbiotic consortia with heterotrophic partners that are widely distributed
1071 in chemoclines of meromictic lakes throughout the world (Bri e et al., 2007). *Chlorobi* metabolic
1072 features include fixing CO₂, fixing N₂ and oxidize sulfide or other reduced sulfur compounds (Liu
1073 et al., 2012). Most our sequences were classified to *Ignavibacteria* order which capable of
1074 organoheterotrophy under both oxic and anoxic conditions.

1075 *Firmicutes*

1076 *Firmicutes* are mostly abundant in the upper part of the sediment. Our sequences were classified to
1077 2 main orders: *Bacillales* and *Clostridia*. *Bacillales* have been observed in diverse environments
1078 and with diverse metabolic functions. Our sequences were closely related to a few species: *Bacillus*
1079 *thioparans* (99%) which grew chemolithoautotrophically by oxidation of thiosulfate to sulfate and
1080 was found in deep sea sediment (Pérez-Ibarra et al., 2007). *Bacillus subterraneus* (99%) utilize
1081 amorphous iron(III), Mn(II)(IV), nitrate, nitrite and fumarate as electron acceptors (Kanso et al.,
1082 2002). *Clostridia* are anaerobic bacteria which play an important role in the biogeochemical cycling
1083 of carbon, sulfur and iron. *Clostridia* are key microbial drivers in iron reduction in estuarine
1084 sediments and acid mine drainage environments (Shah et al., 2014). Part of our sequences were
1085 closely related to *Clostridium tunisiense* (96%) which uses elemental sulfur but not sulfate,
1086 thiosulfate, nor sulfite as terminal electron acceptors (Thabet et al., 2004).

1087 *Bacteroidetes*

1088 Sequences of *Bacteroidetes* were mostly abundant in the middle layer of the LK sediment core.
1089 *Bacteroidetes* are known as hydrolytic fermentative bacteria, they are major utilizers of high-
1090 molecular-mass dissolved organic matter in marine ecosystems (Cottrell and Kirchman, 2000).
1091 Therefore, fresh organic matter (high in high molecular mass dissolved organic matter) that
1092 descends from the water column could be utilized by the *Bacteroidetes* present in the upper part of
1093 the sediment.

1094 The major archaeal populations in the sediment:

1095 The distribution of the population was described above. However the environmental conditions and
1096 metabolic functionality of our sequences were not shown. The uncultured clones similar to our
1097 sequences might help indicate on microbial process of the dominant sequences in our samples.

1098 *Euryarchaeota*

1099 *Methanomicrobia*

1100 The majority of our sequences were classified within *Euryarchaeota*, in *Methanomicrobia* class.
1101 Our 16S rRNA sequences were divided between *Methanomicrobiales* and *Methanosarcinales*
1102 order. *Methanosaeta* genus of *Methanosarcinales* increases with depth. *Methanosaeta* are
1103 acetoclastic methanogens which only able to grow on acetate even at low concentrations (Jetten et
1104 al., 1990). *Methanosaeta* was also shown in previous studies of LK (Schwarz et al., 2007a, 2007b)
1105 and in other meso to eutrophic freshwater lakes (Glissman et al., 2004; Koizumi et al., 2003).
1106 Yamada et al. (2014) showed that *Methanosaeta* also have the ability to reduce ferrihydrite with
1107 H₂ as the electron source. The closest relative cultured acetoclastic methanogen to our sequences
1108 was *Methanosaeta concilii* (96%) similar results to the previous studies. *Methanolinea* genus of

1109 *Methanomicrobiales* which is hydrogenotrophic methanogen was observed in constant percentages
1110 for all depths. *Methanoregula* genus which is also hydrogenotrophic methanogen is decreasing
1111 with depth. *Methanoregula*, *Methanosaeta* and *Methanolinea* represent 60% of sequences of
1112 methanogens retrieved from freshwater lakes (Borrel et al., 2011).

1113 *Thermoplasmata*

1114 The dominant family in *Thermoplasmata* is The Marine Benthic Group D and DHVEG-1 (MBG-
1115 D). Members of the MBG-D have been shown to exist in a variety of freshwater and marine
1116 environments (Beal et al., 2009; Borrel et al., 2012), and it is the most widely encountered,
1117 uncultured lineage in freshwater lake sediments. Even though their metabolism is unknown,
1118 hypotheses about their functionalities are based on the environments in which they were found.
1119 Methanogenesis was suggested, as they were found in deep lake sediments with high methane
1120 concentrations (Borrel et al., 2012), and they were also hypothesized to be involved in AOM, as
1121 they were found in AOM zones (Schubert et al., 2011) and in marine seep sediment (Beal et al.,
1122 2009). However, in other environments, in which methane concentrations were low, the utilization
1123 of waste products, intermediates, or dead cells by MBG-D was also suggested (Smith et al., 1975).
1124 Recently, a single cell genomics study showed that members of MBG-D were capable of exogenous
1125 protein degradation in cold anoxic environments (Lloyd et al., 2013).

1126 In addition Terrestrial Miscellaneous Gp(TMEG) family was observed only in the middle and
1127 bottom layers of the sediment. This lineage was includes clones from the terrestrial subsurface and
1128 from soils, marine sediments and freshwater lakes (Teske and Sørensen, 2008). Sequences closely
1129 related to this group were found in sediment influenced by sulfur-rich, hypoxic groundwater and
1130 aquatic sediment samples.

1131 *Thaumarchaeota*

1132 *Thaumarchaeota* are mostly known as archaeal ammonia oxidizers but in the sediments they may
1133 contribute significantly to the reservoir of nitrogen oxides in ocean waters and thus to productivity,
1134 including the assimilation of carbon (Pester et al., 2011). The enzyme ammonia monooxygenase,
1135 which belongs to the enzyme family of copper containing membrane bound monooxygenases that
1136 possess wide substrate ranges. Closely related enzymes, which can often perform the same function
1137 but with different substrates, include ammonia oxidizing bacteria with ammonia (in
1138 *Gammaproteobacteria*) and methane as a substitute (Lontoh et al., 2000). However, a
1139 microorganism's metabolic energy is mainly defined by its downstream enzyme machinery
1140 (Tavormina et al., 2011). In our sediment *Thaumarchaeota* observed in low percentages with 2
1141 main classes: Marine Benthic Group B and Miscellaneous Crenarchaeotic Group.

1142 Miscellaneous Crenarchaeotic Group

1143 The Miscellaneous Crenarchaeotic Group (MCG) is a cosmopolitan class assigned to the
1144 *Thaumarchaeota* phylum whose members are found mostly in anoxic habitats and have the
1145 capability to take up organic carbon (Biddle et al., 2006). Because of their huge environmental
1146 range and their complex phylogeny, the MCG collectively possess great metabolic diversity(Jiang
1147 et al., 2008). Current evidence for members of the MCG lineage suggests that they may obtain
1148 energy from the anaerobic oxidation of methane, but they do so via a dissimilatory methane-
1149 oxidizing process and they do not assimilate its carbon (Biddle et al., 2006)

1150 Marine Benthic Group B

1151 Members of the MBGB were detected in a growing number of benthic marine environments,
1152 including Atlantic deep-sea sediments, marine mud volcanoes, marine carbonate crusts from the
1153 Black Sea, organic-poor deep subsurface sediments of the central oceanic basins, and organic-rich
1154 methane or methane hydrate containing sediments near continental margins(Sørensen and Teske,
1155 2006). Recent results from Ocean Drilling Program showed that no known methantroph was
1156 detected however methane oxidation appeared to be mediated by Marine Benthic Group B and the
1157 Miscellaneous Crenarchaeotal Group (Biddle et al., 2006)

1158 **References**

1159 Beal, E. J., House, C. H. and Orphan, V. J.: Manganese-and Iron-Dependent Marine Methane
1160 Oxidation, *Science*, 325(5937), 184–187, 2009.

1161 Biddle, J. F., Lipp, J. S., Lever, M. a, Lloyd, K. G., Sørensen, K. B., Anderson, R., Fredricks, H. F.,
1162 Elvert, M., Kelly, T. J., Schrag, D. P., Sogin, M. L., Brenchley, J. E., Teske, A., House, C. H. and
1163 Hinrichs, K.-U.: Heterotrophic Archaea dominate sedimentary subsurface ecosystems off Peru.,
1164 *Proc. Natl. Acad. Sci. U. S. A.*, 103(10), 3846–3851, 2006.

1165 Bodrossy, L., Holmes, E. M., Holmes, A. J., Kovács, K. L. and Murrell, J. C.: Analysis of 16S
1166 rRNA and methane monooxygenase gene sequences reveals a novel group of thermotolerant, *Arch*
1167 *microbiol*, 168, 493–503, 1997.

1168 Borrel, G., Jézéquel, D., Biderre-Petit, C., Morel-Desrosiers, N., Morel, J.-P., Peyret, P., Fonty, G.
1169 and Lehours, A.-C.: Production and consumption of methane in freshwater lake ecosystems., *Res.*
1170 *Microbiol.*, 162(9), 832–847, 2011.

1171 Borrel, G., Lehours, A.-C., Crouzet, O., Jézéquel, D., Rockne, K., Kulczak, A., Duffaud, E., Joblin,
1172 K. and Fonty, G.: Stratification of Archaea in the deep sediments of a freshwater meromictic lake:
1173 vertical shift from methanogenic to uncultured archaeal lineages., *PLoS One*, 7(8), e43346, 1–14,
1174 2012.

- 1175 Briée, C., Moreira, D. and López-García, P.: Archaeal and bacterial community composition of
1176 sediment and plankton from a suboxic freshwater pond., *Res. Microbiol.*, 158(3), 213–227, 2007.
- 1177 Burow, L. C., Woebken, D., Marshall, I. P. G., Singer, S. W., Pett-Ridge, J., Prufert-Bebout, L.,
1178 Spormann, a M., Bebout, B. M., Weber, P. K. and Hoehler, T. M.: Identification of
1179 Desulfobacterales as primary hydrogenotrophs in a complex microbial mat community.,
1180 *Geobiology*, 12(3), 221–230, 2014.
- 1181 Chen, M., Yan, Y., Zhang, W., Lu, W., Wang, J., Ping, S. and Lin, M.: Complete genome sequence
1182 of the type strain *Pseudomonas stutzeri* CGMCC 1.1803., *J. Bacteriol.*, 193(21), 6095, 2011.
- 1183 Cottrell, M. T. and Kirchman, D. L.: Natural Assemblages of Marine Proteobacteria and Members
1184 of the Cytophaga-Flavobacter Cluster Consuming Low- and High-Molecular-Weight Dissolved
1185 Organic Matter, *Appl. Environ. Microbiol.*, 66(4), 1692–1697, 2000.
- 1186 Edlund, A., Hårdeman, F., Jansson, J. K. and Sjöling, S.: Active bacterial community structure
1187 along vertical redox gradients in Baltic Sea sediment., *Environ. Microbiol.*, 10(8), 2051–2063,
1188 2008.
- 1189 Ehrich, S., Behrens, D., Lebedeva, E., Ludwig, W. and Bock, E.: A new obligately
1190 chemolithoautotrophic, nitrite-oxidizing bacterium, *Nitrospira moscoviensis* sp. nov. and its
1191 phylogenetic relationship., *Arch. Microbiol.*, 164(1), 16–23, 1995.
- 1192 Glissman, K., Chin, K.-J., Casper, P. and Conrad, R.: Methanogenic pathway and archaeal
1193 community structure in the sediment of eutrophic Lake Dagow: effect of temperature., *Microb.*
1194 *Ecol.*, 48(3), 389–399, 2004.
- 1195 Jetten, M. S. M., Stams, A. J. M. and Zehnder, A. J. B.: Acetate threshold values and acetate
1196 activating enzymes in methanogenic bacteria, *FEMS Microbiol. Lett.*, 73(4), 339–344, 1990.
- 1197 Jiang, H., Dong, H., Yu, B., Ye, Q., Shen, J., Rowe, H. and Zhang, C.: Dominance of putative
1198 marine benthic Archaea in Qinghai Lake, north-western China., *Environ. Microbiol.*, 10(9), 2355–
1199 2367, 2008.
- 1200 Jiang, L., Zheng, Y., Peng, X., Zhou, H., Zhang, C., Xiao, X. and Wang, F.: Vertical distribution
1201 and diversity of sulfate-reducing prokaryotes in the Pearl River estuarine sediments, Southern
1202 China., *FEMS Microbiol. Ecol.*, 70(2), 93–106, 2009.
- 1203 Kanso, S., Greene, A. C. and Patel, B. K. C.: *Bacillus subterraneus* sp. nov., an iron- and
1204 manganese-reducing bacterium from a deep subsurface Australian thermal aquifer, , 869–874, 2002.
- 1205 Kelly, D. P. and Wood, A. P.: Confirmation of *Thiobacillus denitrificans* as a species of the genus
1206 *Thiobacillus*, in the β -subclass of the Proteobacteria, with strain NCIMB 9548 as the type strain, ,
1207 547–550, 2000.
- 1208 Koizumi, Y., Takii, S., Nishino, M. and Nakajima, T.: Vertical distributions of sulfate-reducing
1209 bacteria and methane-producing archaea quantified by oligonucleotide probe hybridization in the
1210 profundal sediment of a mesotrophic lake., *FEMS Microbiol. Ecol.*, 44(1), 101–108, 2003.
- 1211 Liu, Z., Frigaard, N.-U., Vogl, K., Iino, T., Ohkuma, M., Overmann, J. and Bryant, D. a: Complete
1212 Genome of *Ignavibacterium album*, a Metabolically Versatile, Flagellated, Facultative Anaerobe
1213 from the Phylum Chlorobi., *Front. Microbiol.*, 3(May), 185, 2012.

- 1214 Lloyd, K. G., Schreiber, L., Petersen, D. G., Kjeldsen, K. U., Lever, M. a., Steen, A. D.,
 1215 Stepanauskas, R., Richter, M., Kleindienst, S., Lenk, S., Schramm, A. and Jørgensen, B. B.:
 1216 Predominant archaea in marine sediments degrade detrital proteins, *Nature*, 496(7444), 215–218,
 1217 2013.
- 1218 Lontoh, S., DiSpirito, a a, Krema, C. L., Whittaker, M. R., Hooper, a B. and Semrau, J. D.:
 1219 Differential inhibition in vivo of ammonia monooxygenase, soluble methane monooxygenase and
 1220 membrane-associated methane monooxygenase by phenylacetylene., *Environ. Microbiol.*, 2(5), 485–
 1221 494, 2000.
- 1222 Lovley, D.: Dissimilatory Fe (III) - and Mn (IV) -Reducing Prokaryotes Microorganisms, in
 1223 Prokaryotes, pp. 635–658, Springer New York, New York., 2006.
- 1224 Lovley, D. R., Phillips, E. J. P., Lonergan, D. J. and Widman, P. K.: Fe (III) and S₀ reduction by
 1225 *Pelobacter carbinolicus*, *Appl. Environ. Microbiol.*, 61(6), 2132–2138, 1995.
- 1226 Luo, J.-F., Lin, W.-T. and Guo, Y.: Functional genes based analysis of sulfur-oxidizing bacteria
 1227 community in sulfide removing bioreactor., *Appl. Microbiol. Biotechnol.*, 90(2), 769–778, 2011.
- 1228 Maymo, X., Chien, Y., Gossett, J. M. and Zinder, S. H.: Isolation of a Bacterium That Reductively
 1229 Dechlorinates Tetrachloroethene to Ethene, 1995.
- 1230 McBeth, J. M., Fleming, E. J. and Emerson, D.: The transition from freshwater to marine iron-
 1231 oxidizing bacterial lineages along a salinity gradient on the Sheepscot River, Maine, USA., *Environ.*
 1232 *Microbiol. Rep.*, 5(3), 453–463, 2013.
- 1233 McInerney, M. J., Rohlin, L., Mouttaki, H., Kim, U., Krupp, R. S., Rios-Hernandez, L., Sieber, J.,
 1234 Struchtemeyer, C. G., Bhattacharyya, A., Campbell, J. W. and Gunsalus, R. P.: The genome of
 1235 *Syntrophus aciditrophicus*: life at the thermodynamic limit of microbial growth., *Proc. Natl. Acad.*
 1236 *Sci. U. S. A.*, 104(18), 7600–7605, 2007.
- 1237 Nazina, T. N., Kosareva, I. M., Davidov, a. S., Tourova, T. P., Novikova, E. V., Khafizov, R. R.
 1238 and Poltarau, a. B.: Physicochemical and microbiological characteristics of groundwater from
 1239 observation wells of a deep radioactive liquid waste repository, *Microbiology*, 69(1), 89–95, 2000.
- 1240 Ontiveros-Valencia, A., Ilhan, Z. E., Kang, D.-W., Rittmann, B. and Krajmalnik-Brown, R.:
 1241 Phylogenetic analysis of nitrate- and sulfate-reducing bacteria in a hydrogen-fed biofilm., *FEMS*
 1242 *Microbiol. Ecol.*, 85(1), 158–167, 2013.
- 1243 Orcutt, B. N., Sylvan, J. B., Knab, N. J. and Edwards, K. J.: Microbial ecology of the dark ocean
 1244 above, at, and below the seafloor., *Microbiol. Mol. Biol. Rev.*, 75(2), 361–422, 2011.
- 1245 Pérez-Ibarra, B. M., Flores, M. E. and García-Varela, M.: Isolation and characterization of *Bacillus*
 1246 *thioparus* sp. nov., chemolithoautotrophic, thiosulfate-oxidizing bacterium., *FEMS Microbiol. Lett.*,
 1247 271(2), 289–296, 2007.
- 1248 Pester, M., Schleper, C. and Wagner, M.: The Thaumarchaeota: an emerging view of their
 1249 phylogeny and ecophysiology., *Curr. Opin. Microbiol.*, 14(3), 300–306, 2011.
- 1250 Pjevac, P., Kamyshny, A., Dyksma, S. and Mußmann, M.: Microbial consumption of zero-valence
 1251 sulfur in marine benthic habitats., *Environ. Microbiol.*, 16(11), 3416–3430, 2014.

- 1252 Plugge, C. M., Zhang, W., Scholten, J. C. M. and Stams, A. J. M.: Metabolic flexibility of sulfate-
1253 reducing bacteria., *Front. Microbiol.*, 2(May), 1–8, 2011.
- 1254 Schubert, C. J., Vazquez, F., Lösekann-Behrens, T., Knittel, K., Tonolla, M. and Boetius, A.:
1255 Evidence for anaerobic oxidation of methane in sediments of a freshwater system (Lago di
1256 Cadagno)., *FEMS Microbiol. Ecol.*, 76(1), 26–38, 2011.
- 1257 Schwarz, J. I. K., Eckert, W. and Conrad, R.: Community structure of Archaea and Bacteria in a
1258 profundal lake sediment Lake Kinneret (Israel), *Syst. Appl. Microbiol.*, 30(3), 239–254, 2007a.
- 1259 Schwarz, J. I. K., Lueders, T., Eckert, W. and Conrad, R.: Identification of acetate-utilizing Bacteria
1260 and Archaea in methanogenic profundal sediments of Lake Kinneret (Israel) by stable isotope
1261 probing of rRNA., *Environ. Microbiol.*, 9(1), 223–237, 2007b.
- 1262 Shah, M., Lin, C.-C., Kukkadapu, R., Engelhard, M. H., Zhao, X., Wang, Y., Barkay, T. and Yee,
1263 N.: Syntrophic Effects in a Subsurface Clostridial Consortium on Fe(III)-(Oxyhydr)oxide Reduction
1264 and Secondary Mineralization, *Geomicrobiol. J.*, 31(2), 101–115, 2014.
- 1265 Smith, P. F., Langworthy, T. A. and Smith, M. R.: Polypeptide nature of growth requirement in
1266 yeast extract for *Thermoplasma* Polypeptide Nature of Growth Requirement in Yeast Extract for
1267 *Thermoplasma acidophilum*, *J. Bacteriol.*, 124(2), 884–892, 1975.
- 1268 Sonne-Hansen, J. and Ahring, B. K.: *Thermodesulfobacterium hveragerdense* sp. nov., and
1269 *Thermodesulfobacterium islandicus* sp. nov., two thermophilic sulfate reducing bacteria isolated from a
1270 Icelandic hot spring., *Syst. Appl. Microbiol.*, 22(4), 559–564, 1999.
- 1271 Sørensen, K. B. and Teske, A.: Stratified communities of active Archaea in deep marine subsurface
1272 sediments., *Appl. Environ. Microbiol.*, 72(7), 4596–4603, 2006.
- 1273 Spring, S., Amann, R., Ludwig, W., Schleifer, K. H., van Gemerden, H. and Petersen, N.:
1274 Dominating role of an unusual magnetotactic bacterium in the microaerobic zone of a freshwater
1275 sediment., *Appl. Environ. Microbiol.*, 59(8), 2397–2403, 1993.
- 1276 Sun, H., Spring, S., Lapidus, A., Davenport, K., Del Rio, T. G., Tice, H., Nolan, M., Copeland, A.,
1277 Cheng, J.-F., Lucas, S., Tapia, R., Goodwin, L., Pitluck, S., Ivanova, N., Pagani, I., Mavromatis, K.,
1278 Ovchinnikova, G., Pati, A., Chen, A., Palaniappan, K., Hauser, L., Chang, Y.-J., Jeffries, C. D.,
1279 Detter, J. C., Han, C., Rohde, M., Brambilla, E., Göker, M., Woyke, T., Bristow, J., Eisen, J. a,
1280 Markowitz, V., Hugenholtz, P., Kyrpides, N. C., Klenk, H.-P. and Land, M.: Complete genome
1281 sequence of *Desulfarculus baarsii* type strain (2st14)., *Stand. Genomic Sci.*, 3(3), 276–284, 2010.
- 1282 Tamaki, H., Sekiguchi, Y., Hanada, S., Nomura, N., Matsumura, M., Nakamura, K. and Kamagata,
1283 Y.: Comparative Analysis of Bacterial Diversity in Freshwater Sediment of a Shallow Eutrophic
1284 Lake by Molecular and Improved Cultivation-Based Techniques Comparative Analysis of Bacterial
1285 Diversity in Freshwater Sediment of a Shallow Eutrophic Lake by Molecul, *Appl. Environ.*
1286 *Microbiol.*, 71(4), 2162–2169, 2005.
- 1287 Tavormina, P. L., Orphan, V. J., Kalyuzhnaya, M. G., Jetten, M. S. M. and Klotz, M. G.: A novel
1288 family of functional operons encoding methane/ammonia monooxygenase-related proteins in
1289 gammaproteobacterial methanotrophs, *Environ. Microbiol. Rep.*, 3(1), 91–100, 2011.

- 1290 Tavormina, P. L., Ussler, W. and Orphan, V. J.: Planktonic and sediment-associated aerobic
1291 methanotrophs in two seep systems along the North American margin., *Appl. Environ. Microbiol.*,
1292 74(13), 3985–2995, 2008.
- 1293 Teske, A. and Sørensen, K. B.: Uncultured archaea in deep marine subsurface sediments: have we
1294 caught them all?, *ISME J.*, 2(1), 3–18, 2007.90, 2008.
- 1295 Thabet, O. B. D., Fardeau, M.-L., Joulian, C., Thomas, P., Hamdi, M., Garcia, J.-L. and Ollivier, B.:
1296 *Clostridium tunisiense* sp. nov., a new proteolytic, sulfur-reducing bacterium isolated from an olive
1297 mill wastewater contaminated by phosphogypse., *Anaerobe*, 10(3), 185–190, 2004.
- 1298 Wasmund, K., Lars, S., Lloyd, K. G., Petersen, D. G., Schramm, A., Stepanauskas, R., Jørgensen,
1299 B. B. and Adrian, L.: Genome sequencing of a single cell of the widely distributed marine
1300 subsurface Dehalococcoidia, phylum Chloroflexi, *ISME J.*, 8, 383–397, 2013.
- 1301 Wobus, A., Bleul, C., Maassen, S., Scheerer, C., Schuppler, M., Jacobs, E. and Röske, I.: Microbial
1302 diversity and functional characterization of sediments from reservoirs of different trophic state.,
1303 *FEMS Microbiol. Ecol.*, 46(3), 331–347, 2003.
- 1304 Yamada, C., Kato, S., Kimura, S., Ishii, M. and Igarashi, Y.: Reduction of Fe(III) oxides by
1305 phylogenetically and physiologically diverse thermophilic methanogens., *FEMS Microbiol. Ecol.*,
1306 89(3), 637–645, 2014.
- 1307 Yamada, T. and Sekiguchi, Y.: Cultivation of Uncultured Chloroflexi Subphyla: Significance and
1308 Ecophysiology of Formerly Uncultured Chloroflexi “Subphylum I” with Natural and
1309 Biotechnological Relevance, *Microbes Environ.*, 24(3), 205–216, 2009.
- 1310 Yamada, T., Sekiguchi, Y., Hanada, S., Imachi, H., Ohashi, A., Harada, H. and Kamagata, Y.:
1311 *Anaerolinea thermolimosa* sp. nov., *Levilinea saccharolytica* gen. nov., sp. nov. and *Leptolinea*
1312 *tardivitalis* gen. nov., sp. nov., novel filamentous anaerobes, and description of the new classes
1313 *Anaerolineae* classis nov. and *Caldilineae* classis nov. in the, *Int. J. Syst. Evol. Microbiol.*, 56(Pt 6),
1314 1331–1340, 2006.
- 1315 Yan, Z., Song, N., Cai, H., Tay, J.-H. and Jiang, H.: Enhanced degradation of phenanthrene and
1316 pyrene in freshwater sediments by combined employment of sediment microbial fuel cell and
1317 amorphous ferric hydroxide., *J. Hazard. Mater.*, 199-200, 217–225, 2012.
- 1318 Yoshida, N., Takahashi, N. and Hiraishi, A.: Phylogenetic Characterization of a Polychlorinated-
1319 Dioxin- Dechlorinating Microbial Community by Use of Microcosm Studies, , 71(8), 4325–4334,
1320 2005.
- 1321
- 1322