

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35

Dear Dr. Kirsten Kuesel,

Co-Editor of Biogeosciences

We would like to thank you and the reviewers for the positive and constructive reviews that have improved the manuscript significantly. We have accepted all comments and addressed them in detail below (our response is in bold).

Main comments:

1) The archaeal sequences belonging to uncultured lineages (MCG, MBG-B, MBG-D) represent less than 5% of the sequences. It is impossible in this case to infer on their potential metabolisms based solely on the geochemical characteristics of the environment in which they were detected. I would suggest deleting all paragraphs pertaining to these groups in the discussion and discuss the dominant archaeal groups which are also known described cultured organisms.

**It is correct and the sequences of archaeal classes that were below 5% of the community (MCG, MBG-B) were not discussed, as it impossible to infer their potential metabolisms based only on the geochemical characteristics of the environment in which they were detected. However MBG-D (as important group in methane cycle mechanism) was represented by about 5% of the sequences in the deep sediment and thus discussed in the manuscript.**

2) Part 4.2 is long and descriptive, whereas part 4.3 contains very interesting information. Please delete part 4.2 and integrate the most relevant data from this part in part 4.3.

**Part 4.2 was deleted as suggested and the relevant information was integrated to part 4.3.**

3) Most of the discussion in part 4.3 needs to be checked for English and the authors need to clearly write what data type they are discussing. Here are some examples:

**The revised version was edited as requested and additional round of language editing was done.**

L. 543 states ‘mcrA functional gene of methanogens/anaerobic methanotrophs shows’, which should be written ‘xxx analysis of mcrA functional genes ... shows.’

**The sentence was changed according to the advice.**

L. 546-548 ‘is likely because that the mcrA observed ...’, what are the authors referring to here? qPCR data or mcrA gene sequences?

**The sentence was clarified in the text.**

L. 547 ‘methanogens which maybe cause the reverse methanogenesis’, methanogens possibly perform reverse methanogenesis but they don’t cause it. Etc ...

**This sentence was corrected in the text.**

36

37 Specific comments

38 L. 82 : Please change ‘microbial pathways’ to ‘microbial communities’

39 **The change was made.**

40 L. 187: ‘Chimeras?’

41 **Using the position window in the denoising pipeline of MOTHUR helps us to look at the same**  
42 **region in the 16S rRNA gene in our sequenced dataset. In this way we can better compare the**  
43 **data set between other samples of the same position window. After using the position window**  
44 **we also use chimera check and discard all the chimeras. This was explained better in the text.**

45 L. 266-271: This paragraph has information that is repeated after in more detail, please delete. Same  
46 comment for paragraph L. 280-286.

47 **The paragraph was deleted as suggested.**

48 L. 285-286: Recent data suggests that the MCG and MBG-B are separate phyla and are not  
49 considered to belong to the Thaumarchaeota anymore. Hence some of the further discussions need  
50 to be amended.

51 **The MCG and MBG-B were almost not mentioned in the revised manuscript, however the**  
52 **recent change in phylogenetics of MCG and MBG-B was added to the revised version.**

53 L. 304: MBG-D and DHVEG-1 belong to the same cluster, they can all be labelled MBG-D.

54 **This was corrected.**

55 L. 384-385: please explain this sentence, what was analysed exactly?

56 **The sentence was clarified.**

57 L. 403: ‘microbial populations’, ‘methane cycling’

58 **This paragraph was deleted.**

59 L. 515-517: ‘which creates ... in deep sediments’, please rewrite this sentence

60 **The sentence was rewritten and clarified.**

61 L. 526: there is a verb missing in this sentence

62 **The sentence was fixed.**

63 L. 533: ‘can occur by one microorganism?’ Please rephrase.

64 **The sentence was rephrased.**

65 L. 534-536: 16S rRNA gene data is not enough to infer on metabolism. Please delete this sentence.

66 **The possible direct mechanism that could be performed by unique methanotrophy like MBG-**  
67 **D (Thermoplasmata class) or MCG/MBG-B, which some of our sequences were similar was**  
68 **suggested.**

69

70

Revised version to Biogeosciences

71 **Methane related changes in prokaryotes along geochemical profiles in sediments**  
72 **of Lake Kinneret (Israel)**

73 **I. Bar Or<sup>1</sup>, E. Ben-Dov<sup>2,3</sup>, A. Kushmaro<sup>\*2,5,6</sup>, W. Eckert<sup>4</sup> and O. Sivan<sup>1</sup>**

74

75 <sup>1</sup>Department of Geological and Environmental Sciences, Ben-Gurion University of the Negev,  
76 Be'er- Sheva, P.O. Box 653, 8410501, Israel.

77 <sup>2</sup>Avram and Stella Goldstein-Goren Department of Biotechnology Engineering, Ben-Gurion  
78 University of the Negev, Be'er- Sheva, P.O. Box 653, 8410501, Israel.

79 <sup>3</sup>Department of Life Sciences, Achva Academic College, Achva, M.P. Shikmim 79800, Israel.

80 <sup>4</sup> Israel Oceanographic and Limnological Research, The Yigal Allon Kinneret Limnological  
81 Laboratory, PO Box 447, 14950 Migdal, Israel.

82 <sup>5</sup>National Institute for Biotechnology in the Negev, Ben-Gurion University of the Negev, P.O. Box  
83 653, Be'er-Sheva 8410501, Israel

84 <sup>6</sup>School of Materials Science and Engineering, Nanyang Technological University, Singapore

85

86 \* Corresponding author: Mailing address: Department of Biotechnology Engineering, Ben-Gurion  
87 University of the Negev, P.O. Box 653, Be'er-Sheva 8410501, Israel. Phone: 972-8-6479.024. Fax:  
88 972-8-6472.983. E-mail: arielkus@bgu.ac.il.

89

90 **Keywords:** anaerobic methane oxidation, iron reduction, archaeal classification, bacterial  
91 classification, Lake Kinneret.

92

93 **Abstract.** Microbial methane oxidation is the primary control on the emission of the greenhouse  
94 gas methane into the atmosphere. In terrestrial environments, aerobic methanotrophic bacteria are  
95 largely responsible for this process. In marine sediments a coupling of anaerobic oxidation of  
96 methane (AOM) with sulfate reduction, often carried out by a consortium of anaerobic  
97 methanotrophic archaea (ANME) and sulfate reducing bacteria, consumes almost all methane  
98 produced within marine sediments. Motivated by recent evidence for AOM with iron(III) in Lake

99 Kinneret sediments, the goal of the present study was to link the geochemical gradients in the lake  
100 porewater to the microbial community structure. Screening of archaeal 16S rRNA gene sequences  
101 revealed a shift from hydrogenotrophic to acetoclastic methanogens with depth. The observed  
102 changes in microbial community structure suggest possible direct and indirect mechanisms for the  
103 AOM coupled to iron reduction in deep sediments. Members of the *Burkholderiales* and  
104 *Nitrospirales* orders increased with depth, suggesting their involvement in iron reduction together  
105 with *Geobacter* genus and "reverse methanogenesis". An indirect mechanism through sulfate and  
106 ANMEs seems less probable ~~to explain the data~~ due to the absence of ANME sequences. This is  
107 despite the abundant sequences related to sulfate reducing bacteria (*Deltaproteobacteria*) together  
108 with the occurrence of *dsrAB* in the deep sediment that could indicate the production of sulfate  
109 (disproportionation) from S<sup>0</sup> for sulfate-driven AOM. The presence of the functional gene *pmoA* in  
110 the deep anoxic sediment together with sequences related to *Methylococcales* suggests the existence  
111 of a second unexpected indirect pathway - aerobic methane oxidation pathway in an anaerobic  
112 environment.

113

114

## 115 1. Introduction

116 Chemical profiles in the porewater of aquatic sediments reflect the sequence of microbially  
117 mediated redox reactions that are driven by the availability of both, electron donors and of suitable  
118 electron acceptors. The latter are depleted in the order of decreasing chemical potential, beginning  
119 with oxygen and proceeding through nitrate, manganese and iron oxides, and then sulfate. Below  
120 the main zone of sulfate reduction, the fermentation of organic carbon leads to the formation of  
121 methane (CH<sub>4</sub>) by the process of methanogenesis (Froelich et al., 1979).

122 The produced methane is isotopically depleted in <sup>13</sup>C, with values of ~-50 to -110‰ (Schoell,  
123 1988), and the residual dissolved inorganic carbon (DIC) pool is enriched by an isotopic  
124 fractionation factor (ε) of 50 to 70‰ (e.g., (Borowski et al., 2000; Whiticar, 1999)). When the  
125 produced methane diffuses into a zone with a suitable electron acceptor, it can be consumed by  
126 microbial oxidation (methanotrophy), the main process by which the important greenhouse methane  
127 is prevented from escaping into the atmosphere. While in the terrestrial environment, aerobic  
128 methanotrophy is the dominant process (Chistoserdova et al., 2005), in anaerobic marine sediments  
129 archaea are found to consume the majority of upward diffusing methane coupled to sulfate  
130 reduction (Knittel and Boetius, 2009; Thauer, 2010; Valentine, 2002).

131 Evidence from lipids and from fluorescence *in situ* hybridization (FISH) showed that a  
132 consortium of archaea and sulfate reducing bacteria are involved in this anaerobic methane  
133 oxidation (AOM) (Boetius et al., 2000; Hinrichs et al., 1999; Orphan et al., 2001). To date, three  
134 groups of anaerobic methanotrophic archaea (ANME), named ANME-1, ANME-2, and ANME-3,  
135 are known to perform sulfate driven AOM (Niemann et al., 2006; Orphan et al., 2002). **However**  
136 **Recently** Milucka et al. (2012) demonstrated AOM mediated solely by archaea, where the archaea  
137 was shown to oxidize the methane and reduce the sulfate to elemental sulfur. Disproportionating  
138 bacteria, also involved in this mechanism, oxidize and reduce this elemental sulfur to sulfate and  
139 sulfide, respectively. The carbon isotopic fractionation factor (ε) for this methanotrophic process  
140 was shown to be in the range of 4-30‰ (Kinnaman et al., 2007; Whiticar, 1999). Enrichment  
141 cultures of ANME from different environments showed a carbon isotopic fractionation of 12-39‰  
142 (Holler et al., 2009).

143 Other electron acceptors were recently shown to drive AOM. Nitrite driven AOM by oxygenic  
144 bacteria was observed in two different freshwater ecosystems in Netherlands (Ettwig et al., 2009;  
145 Raghoebarsing et al., 2006) and also in peatlands (Zhu et al., 2012). Beal et al. (Beal et al., 2009)  
146 showed the potential of manganese and iron-driven AOM in marine sediments, and Egger et al.,  
147 (2014) showed it in brackish coastal sediments. In our recent study (Sivan et al., 2011), we provided  
148 *in situ* geochemical evidence for AOM coupled to microbial iron reduction below the main

149 methanogenesis zone in Lake Kinneret (LK) sediments, where dissolved sulfate and nitrate are  
150 absent. However, the mechanism that is responsible for this process was not investigated. The goal  
151 of the present study ~~was-is~~ to explore the possible microbial ~~pathways-communities~~ that may be  
152 involved in the iron driven AOM in LK sediments. This is accomplished by combining chemical  
153 and isotope analyses of porewater samples along a depth profile from LK sediments with molecular  
154 biological techniques. By using specific functional genes related to dissimilatory sulfate reductase  
155 (*dsr*) for sulfate reducers (Klein et al., 2001), methyl-coenzyme M reductase (*mcr*) for methanogens  
156 and anaerobic methanotrophs (Hallam et al., 2003) and particulate methane monooxygenase  
157 (*pmmo*) for aerobic methanotrophs (McDonald et al., 2008) we pinpoint the most likely candidate  
158 for this relatively unknown process.

## 159 1.1. Study site

160 Located in northern Israel, Lake Kinneret (LK, Fig.1) is a warm monomictic subtropical lake.  
161 Typical concentrations of major electron acceptors in the water column during the mixed period are  
162 35–50  $\mu\text{M}$  nitrate and 600  $\mu\text{M}$  sulfate (Adler et al., 2011; Nishri et al., 2000; Serruya et al., 1974).  
163 In the spring, the newly formed epilimnion is characterized by increasing temperatures and  
164 enhanced phytoplankton development, while in the hypolimnion heterotrophic microorganisms  
165 gradually deplete oxygen and then nitrate. Organic matter degradation by bacterial iron and  
166 manganese reduction takes place below the thermocline in the summer, and in the end of the  
167 stratification period, sulfate reduction starts in the bottom water. In the upper part of the sediment,  
168 sulfate reduction is the dominant microbial process year round, and below depth of 5 cm this  
169 process is mostly replaced by methanogenesis (Adler et al., 2011; Eckert and Conrad, 2007). Total  
170 iron content ( $\text{Fe}_{(\text{tot})}$ ) in the sediment increases with depth till 18 cm and then remains uniform  
171 around 550  $\mu\text{mol/g}$  Dry Weight (Eckert, 2000).  $\text{Mn(II)}$  concentration is about 13  $\mu\text{mol/g}$  Dry  
172 Weight in the sediment (Serruya, 1971). The total carbon ( $\text{C}_{(\text{tot})}$ ) in the solid phase shows a decrease  
173 from 14  $\mu\text{mol/g}$  Dry Weight in the top part of the sediment to 8  $\mu\text{mol/g}$  Dry Weight in the deep part  
174 of the sediment ( $\sim 30$  cm) (Eckert, 2000).

175

## 176 2. Material and Methods

### 177 2.1. Sampling

178 Several sediment cores were collected from the center of the lake (Station A, Fig. 1) at a water  
179 depth of  $\sim 42$  m (maximum lake water column depth) at different times using Perspex tubes,  
180 measuring 55 cm long by 5 cm in diameter, with a gravity corer. The cores were stored in the dark  
181 at  $4^\circ\text{C}$  until they were sliced (on the same day or the day after). Core sampling for the microbial

182 community study took place in December 2009. Over a dozen of porewater chemical and isotope  
183 profiles were conducted seasonally (every 3-4 months) from 2007 to 2013. The slight seasonal  
184 changes allowed using typical geochemical profiles (Adler et al., 2011) in order to sample for the  
185 microbiology communities in the different electron acceptors zones and to correlate between the  
186 microbial sampling and the geochemical profiles.

## 187 **2.2. Geochemical analyses**

188 Cores were divided into 2-cm slices from top to bottom under a constant flow of N<sub>2</sub> using a  
189 slicing device. About 1.5 ml of each sediment slice was transferred into N<sub>2</sub>-flushed crimp bottles  
190 containing 5 ml of 1.5 N NaOH for the headspace measurements of CH<sub>4</sub> and δ<sup>13</sup>C<sub>CH<sub>4</sub></sub> (after Sivan et  
191 al., 2011). CH<sub>4</sub> concentrations were measured on a SHIMADZU gas chromatograph (GC 8IF)  
192 equipped with a FID detector at a precision of 2 μmol·L<sup>-1</sup>.

193 Porewater was extracted immediately from each slice by centrifugation at 27,000 g at 4°C in a  
194 N<sub>2</sub> atmosphere, and the supernatant was filtered through 0.45-μm filters. A 1-ml subsample was  
195 fixed with Ferrozine and analyzed for dissolved Fe(II) (Stookey, 1970). Four millimeters of  
196 subsample were poured into an acidified vial (1 ml of 0.5N nitric acid) to measure dissolved Mn(II).  
197 The sample was analyzed via an Inductively Coupled Plasma Mass Spectrometer (ICP-MS, Elan  
198 DRC II, Perkin Elmer) at a precision of ±10%. For sulfide profiles, 1 ml of subsample was added to  
199 zinc acetate and hydrogen sulfide concentrations were determined by the methylene blue method  
200 with MDL of the method is 1 μM (Cline, 1969). For sulfate measurements, 5 ml of the subsamples  
201 were analyzed with a Dionex DX500 high-pressure liquid chromatograph with a precision of ±3%.  
202 Iron isotope analysis was done by acidifying the subsamples with 10% HCl for one week (to  
203 dissolve any precipitated iron) and then purifying them by anion exchange chromatography (Borrok  
204 et al., 2007). δ<sup>56</sup>Fe was measured on a Neptune multi-collector ICP-MS in high resolution mode  
205 according to standard methods and standardized against isotopic reference material (IRMM-014)  
206 with a precision of ±0.1‰ (John and Adkins, 2010). Total lipids were extracted using the Bligh–  
207 Dyer procedure (Bligh and Dyer, 1959) with solvent mixture of 2:1:0.8  
208 (methanol:dichloromethane:buffer). δ<sup>13</sup>C of the total lipid extraction (TLE) was measured on an  
209 Elemental Analyzer Isotopic Ratio Mass Spectrometer (EA-IRMS) with a precision of 0.1‰.

## 210 **2.3. DNA extraction and quantitative PCR (qPCR) amplification from sediment samples**

211 Part of the sediment slices were kept frozen at -20°C for the microbial work. Samples from  
212 three different depths (0–3 cm representing the sulfate reduction zone, 6–9 cm representing the  
213 methanogenesis zone, and 29–32 cm representing the deep AOM zone) were defrosted. Those depth  
214 zones were chosen based on the geochemical profiles sampled at different times, showing quasi

215 steady state and stabilization of the sediments (Adler et al., 2011; Sivan et al., 2011). Therefore,  
216 microbial community sampling of the three depth zones at a single time point should represent the  
217 different microbial habitats as they pertain to the respective geochemical zones. Total genomic  
218 DNA was extracted from the sediment samples using the MoBio Power Soil DNA isolation kit  
219 (MoBio Laboratories, Solana Beach, CA). Genomic DNA was eluted using 60 µl of elution buffer  
220 and stored at -20°C. Concentrations of DNA were determined via UV-Vis spectrophotometry (ND-  
221 1000 NanoDrop Technologies, Wilmington, DE) (sulfate reduction zone 22ng/µl, methanogenesis  
222 zone 35.8ng/µl and AOM zone 14ng/µl).

223 Quantification of functional genes were performed using the *mcrA* primers (Luton et al., 2002),  
224 F 'GGTGGTGTMGGATTCACACARTAYGCWACAG' and R  
225 'TTCATTGCRTAGTTWGGRTAGTT', *dsrA* primers, DSR1F 'ACSCACTGGAAGCACG'  
226 (Wagner et al., 1998) and RH3-dsr-R 'GGTGGAGCCGTGCATGTT' (Ben-Dov et al., 2007) and  
227 *pmoA* primers, a189F 'GGNGACTGGGACTTCTGG' and mb661R  
228 'CCGGMGCAACGTCYTTACC' (Yan et al., 2006) with a ABI prism 7000 Sequence Detection  
229 System (Applied Biosystems). The qPCR reaction consisted of 10 µL of Absolute Blue SYBR  
230 Green ROX, 150 nM each of forward and reverse primers and 5 µL of each DNA template.  
231 Thermal cycling conditions were as follows: 15 minutes at 95°C for enzyme activation, followed by  
232 40 rounds of 15 seconds at 95°C for denaturation and 1 minute at 60°C for annealing/extension. To  
233 verify that the used primer pair produced only a single specific product, a dissociation protocol was  
234 added after thermo cycling, to determine dissociation of the PCR products from 60°C to 95°C.  
235 Standards for the calibration curves for quantification were made using pGEM-T Easy plasmid  
236 cloned with 1.9 kb *dsrAB* and amplicons of the *mcrA* and *pmoA* encoding for functional genes at a  
237 known concentration with six serial dilution points (in steps of tenfold). All runs included a no-  
238 template control. Plasmid standards and environmental samples were simultaneously assayed in  
239 triplicates. The ABI prism 7000 Sequence Detection System and SDS Software were used for data  
240 analysis. qC values were exported into a Microsoft Excel Work sheet for further statistical analysis.

#### 241 **2.4. Sequence analysis**

242 454 tag-encoded FLX amplicon pyrosequencing was performed by the Research and Testing  
243 Laboratory (Lubbock, Texas, USA) as previously described (Dowd et al., 2008). The bacterial and  
244 archaeal 16S rRNA gene primers that were used are 28F 'GAGTTTGATCNTGGCTCAG' and  
245 519R 'GTNTTACNGCGGCKGCTG' and Arch349F 'GYGCASCAGKCGMGA AW' and Arch806R  
246 'GGACTACVSGGGTATCTAAT' respectively.



247 Data analysis was made using two different methodologies (MOTHUR and SILVA ngs). The initial  
248 trimming of the sequences was made by MOTHUR v1.33 (Schloss et al., 2009) and generated  
249 around 300 bp length of sequences. The trimmed sequences were taken to further analysis by  
250 MOTHUR (Schloss et al., 2011) using SILVA.nr\_v119 database. MOTHUR 454 pipeline filtration  
251 and denoising remove from the analysis sequences that were <150 bp, when they contained  
252 homopolymers longer than 8 bp, ambiguous bases, more than one mismatch to barcode sequences  
253 or more than two mismatches to the forward primer sequence. We further removed sequences that  
254 did not align in the same nucleotide position on the reference database.

255 Most of the bacterial sequences were in the same region, however the archaeal sequences were  
256 ~~spared-spaced~~ between two different regions. We used one position window that was dominant in  
257 the top sample and implied it on the other archaeal analysis [in order to better compare the dataset](#)  
258 [between other samples of the same position window](#). Identical sequences were grouped and then  
259 were aligned against SILVA.nr\_v119. Chimeras were removed using MOTHUR CHIMERA  
260 UCHIME (Edgar et al., 2011). The lengths of the remaining sequences were around 200 bp. A  
261 further screening step (pre-cluster) was applied to reduce sequencing noise by clustering reads  
262 differing by only one base every 100 bases (Huse et al., 2010). In addition, the rest of the sequences  
263 were classified in order to remove eukaryote, mitochondria and chloroplast classified sequences.  
264 The remaining sequences were used to generate a distance matrix and clustering into operational  
265 taxonomic units (OTUs) defined at 97% cutoff using the average neighbor algorithm. The OTUs  
266 were classified using SILVA.nr\_v119 database with a confidence threshold of 80%. Alpha diversity  
267 was calculated by MOTHUR using the remaining sequences at 97% similarity. The beta diversity  
268 was calculated using comparable data, the number of sequences per sample was made equal through  
269 subsampling (Table 1 and 2).

270 The second program which we used for analysis was SILVA ngs pipeline (Quast et al., 2013).  
271 The trimmed sequences from mother were Aligner against the SILVA SSU rRNA seed. Sequences  
272 shorter than 50 aligned nucleotides and with more than 2% of ambiguities or 2% of homopolymers,  
273 respectively, were removed. The sequences that were not aligned as being putative  
274 contaminations/artifacts were removed. Then sequences were clustered to OTUs with 98%  
275 similarity and classified by local nucleotide BLAST search against SILVA database v119. To filter  
276 out low identity and artificial BLAST hits, hits for which the function ‘(%sequence identity  
277 +%alignment coverage)/2’ did not exceed the value of 93% were discarded. Sequences with weak  
278 low score were classified as ‘No Relatives’ and will be disregarded from now on. Standard  
279 deviation between the percentages of the two classification showed that at the phylum level there  
280 are small differences between the two pipelines (supplementary Table 1). In the order classification,

281 the standard deviations increased but the sequences still showed close similarity. Estimates of  
282 phylotype richness, diversity coverage and similarity were calculated according to the abundance-  
283 based coverage estimate (ACE), Chao's estimator (Chao, 1984; Chao and Ma, 1993), the Shannon  
284 diversity index, Good's coverage (Good, 1953) were calculated in MOTHUR (Table 1 and 2). The  
285 distribution and abundance matrix of the OTUs was normalized to the sample with the smallest  
286 number of reads by randomly resampling of the MOTHUR data set for statistical comparisons. Raw  
287 sequencing data was deposited in the MG-RAST (metagenomics.anl.gov) archive.

## 288 2.5. Microbial community structure related to the environment conditions

289 To estimate community similarity among samples, we used PC-ORD 6 software. Subsample  
290 OTU data from MOTHUR was used to create a distance matrix based on the Sørensen (Bray-  
291 Curtis) dissimilarities of the OTU composition of the samples. The data was normalized to  
292 percentages before the analysis. Community relationships were visualized using principal  
293 coordinate analysis (PCoA) based on this distance matrix. Different environmental variables were  
294 added as well ( $\text{CH}_4$ ,  $\delta^{13}\text{C}_{\text{CH}_4}$ , Fe(II),  $\delta^{56}\text{Fe}(\text{II})$ , Mn(II),  $\text{H}_2\text{S}$ ,  $\text{SO}_4^{2-}$ , dissolved organic carbon (DOC),  
295  $\delta^{13}\text{C}_{\text{TLE}}$ ,  $\text{NH}_4^+$  and  $\text{PO}_4^{3-}$ ) in order to identify potential explanatory variables. The environmental  
296 vector were applied and projected to the PCoA ordination. In addition, Venn diagrams for graphical  
297 descriptions of unshared and shared OTUs between the three samples were constructed using  
298 MOTHUR.

299

## 300 3. Results

### 301 3.1. Porewater profiles

302 This study focused on microbial community shifts along the porewater profiles of electron  
303 acceptor gradients related to the methane cycle. To characterize shifts in main electron acceptors  
304 with depth, Over a dozen of porewater profiles were performed seasonally from station A (Fig. 1)  
305 ~~to characterize shifts in main electron acceptors with depth.~~ The shown profiles (Fig. 2) are  
306 representatives of the slight seasonal trends. Sulfate and sulfide profiles show depletion from ~100  
307  $\mu\text{M}$  at the top part of the sediment, to below detection limits within the upper 15 cm (Fig. 2A). This  
308 typical concave curvature profile of sulfate profile in the porewater indicates intensive sulfate  
309 reduction in the upper few centimeters of the sediment throughout the year. A typical LK methane  
310 concentration profile (Fig. 2B) shows increase from 250  $\mu\text{M}$  at the water-sediment interface to a  
311 maximum of about 1.25 mM in the depth range of 7 to 15 cm, and then a decrease below 15 cm  
312 depth. The profile of  $\delta^{13}\text{C}_{\text{CH}_4}$  (Fig. 2B) shows a decrease from -60‰ at a depth of 1 cm to about

313 -65‰ at a depth of 7 cm and then an increase in the deeper sediments to a maximum value of  
314 -53.5‰ at a depth of 25 cm. Also, the profile of  $\delta^{13}\text{CTLE}$  (Fig. 2C) shows a decrease in this  
315 deepest part of the sediment.

316 Manganese and iron oxides are the most probable electron ~~accepters~~ acceptors in the deep part  
317 of the sediment where methane is decreased. Dissolved Mn(II) concentration (Fig. 2D) increased  
318 from 5  $\mu\text{M}$  at the top of the sediment to a plateau of about 23  $\mu\text{M}$  from depths of 23 to 36 cm. The  
319 dissolved Fe(II) concentration profile also showed an increase with depth (Fig. 2D), however  
320 however infollowing a different pattern. In the upper 15 cm, dissolved Fe(II) concentrations were  
321 below the detection limit, and gradually increased below 15 cm to about 90  $\mu\text{M}$  at a depth of 36 cm.  
322 The  $\delta^{56}\text{Fe}$  profile (Fig. 2C) showed a decrease with depth from 0.5‰ in the upper part to -1.7‰ to  
323 -2.3‰ in the deep part of the sediment.

### 324 3.2. Sediment microbial communities

325 To study the sediment depth-dependent microbial community shifts, 16S rRNA gene sequences  
326 of the prokaryotic community from three different depth zones (0–3, 6–9 and 29–32 cm, see arrows  
327 in Fig. 2a) were extracted and analyzed. As mentioned above, these three zones represent the main  
328 sulfate reduction zone, the methanogenesis zone, and the deep iron driven AOM zone, respectively.  
329 The diverse bacterial and archaeal communities at these depth zones varied in composition and  
330 richness throughout the sediment. A total of 13227 and 4881 bacterial and archaeal sequences  
331 representing 3852 and 705 OTUs (cut-off value of 98%), respectively, were identified by SILVA  
332 database project. The normalized Shannon and Chao index were used as proxies for diversity and  
333 richness, respectively (Table 1 and 2). The upper layer showed lower diversity (Shannon index,  
334 3.91 and 1.8 for bacteria and archaea) and richness (Chao index, 925.7 and 83.4 for bacteria and  
335 archaea) than the other depths. Moreover, the bacterial coverage of the upper layer was the highest  
336 (91%), while that of the deeper layers was about 80%. The archaeal coverage was almost the same  
337 (~88%) in all the sampled layers. Most of our sequences were affiliated with uncultured  
338 microorganisms mainly from various sediment environments.

339 Although the bacterial OTUs were distributed over 43 phyla, we ~~represent~~ here only the phyla  
340 that were over 1% sequences of the 454 library. The most abundant phyla were: *Proteobacteria*,  
341 *Chloroflexi*, *Nitrospirae*, *Bacteroidetes*, *Firmicutes* and *Chlorobi* observed in at least one of the  
342 libraries (Fig. 3). ~~Dominant phyla had different trends; Chloroflexi and Nitrospirae showed increase~~  
343 ~~with depth while Proteobacteria showed decrease with depth. Firmicutes were highly abundant~~  
344 ~~only in the top layer. Bacteroidetes and Chlorobi were more abundant in the middle layer of the~~  
345 ~~sediment.~~

346 The communities of microorganisms in the top layer (0-3cm) of the sediment had less sequence  
347 overlap with those of the deeper layers (70 and 33 shared bacterial sequences with the middle and  
348 bottom layers respectively) and of the deeper layers (Supplementary Fig 1 and 2). The most  
349 dominant (~48%) phylum in this bacterial community was *Proteobacteria*; class  
350 *Gammaproteobacteria* (91%) and the genera *Acinetobacter* (28%) and *Pseudomonas* (14%) (Fig.  
351 3). The second most abundant (38%) phylum was *Firmicutes*; class *Clostridia* (21%) and *Bacilli*  
352 (17%), family *Clostridiaceae* (15%) and genus *Bacillus* (14%). Further sequences were related to  
353 other phyla and order (Fig 3 and supplementary Table 1).

354 In the archaeal community the sequences number were much lower, however the coverage  
355 calculation shows about 90% coverage. The dominant phylum in the upper layer was  
356 *Euryarchaeota* (98%), and the rest of the 2% was *Thaumarchaeota*. *Methanomicrobia* (92%) was  
357 the dominant class of *Euryarchaeota* divided between genera of *Methanoregula* (58%),  
358 *Methanosaeta* (22%) and *Methanolinea* (11%). The other abundant classes of *Euryarchaeota* were  
359 *Halobacteria* (2%) and *Thermoplasmata* (3%). ~~*Thaumarchaeota* was divided between two classes:~~  
360 Marine Benthic Group B (MBG-B 0.7%) and Miscellaneous Crenarchaeotic Group (MCG 1%)  
361 (Fig. 4 and supplementary table 1).

362 The middle layer (6-9cm) was the richest and most diverse in its bacterial and archaeal  
363 populations comparing to the other layers, but with the lowest community coverage (bacterial 77%  
364 and archaeal 86%). The dominant phyla in this depth were divided to *Proteobacteria* (40%),  
365 *Chloroflexi* (14%), *Bacteroidetes* (7%), *Chlorobi* (5%) and *Nitrospirae* (4%). The dominant classes  
366 in *Proteobacteria* were *Deltaproteobacteria* (68%), *Betaproteobacteria* (14%) and  
367 *Gammaproteobacteria* (14%). Sva0485 (8%), *Syntrophobacterales* (7%), *Desulfarculales* (3%) and  
368 *Desulfuromonadales* (2%) were the dominant affiliated order in *Deltaproteobacteria*.  
369 *Methylococcaceae* family of *Gammaproteobacteria* had 1% affiliated sequences. The dominant  
370 family in *Chloroflexi* was *Anaerolineaceae* (6%). *Ignavibacteriales* (4%) order was the dominant in  
371 *Chlorobi*. *Nitrospirae* increased from the upper layer and was mainly represented by the  
372 *Nitrospiraceae* (4%) family. Further sequences were related to other phyla and order (Fig. 3,  
373 supplementary Table 1).

374 The archaeal community was the richest and most diverse in the middle zone. The dominant  
375 phyla were *Euryarchaeota* (96%) while *Thaumarchaeota* has the rest of the 4%. *Euryarchaeota* was  
376 divided between 3 dominant classes: *Methanomicrobia* (78%), *Thermoplasmata* (11%) and  
377 *Halobacteria* (6%). *Methanomicrobia* was divided between *Methanoregula* (19%),  
378 *Methanosaeta*(50%) and *Methanolinea* (10%) genus's. *Thermoplasmata* was divided between  
379 Marine Benthic Group D ~~and DHVEG-1~~ (5%) and Terrestrial Miscellaneous Gp(TMEG) (3%).  
380 *Halobacteria* dominant family was Deep Sea Hydrothermal Vent Gp 6(DHVEG-6) (6%).

381 ~~Thaumarchaeota was divided mainly between two classes:~~ Marine Benthic Group B (MBGB 1.5%)  
382 and Miscellaneous Crenarchaeotic Group (MCG 1.5%) (Fig. 4 [and supplementary table 1](#)).

383 The bottom layer (29-32 cm) bacterial community dominant phyla included: *Proteobacteria*  
384 (39%), *Chloroflexi* (19%) and *Nitrospirae* (8%). *Proteobacteria* were divided to three main classes:  
385 *Deltaproteobacteria* (20%), *Gammaproteobacteria* (10%) and *Betaproteobacteria* (6%). Sva0485  
386 (11%) and *Syntrophobacterales* (4%) were the dominant affiliated order in *Deltaproteobacteria*.  
387 The dominant family in *Chloroflexi* was *Anaerolineaceae* (7%) and *Ignavibacteriales* (4%) order is  
388 the dominant *Chlorobi*. *Nitrospirae* increased from the upper layer and was mainly represented by  
389 *Nitrospiraceae* (8%) family. Further sequences were related to other phyla and order (Fig. 3,  
390 supplementary Table 1)

391 At this depth, the dominant archaeal phyla were *Euryarchaeota* (98%) while *Thaumarchaeota*  
392 has the rest of the 2%. *Euryarchaeota* was divided between 3 dominant classes: *Methanomicrobia*  
393 (86%), *Thermoplasmata* (6%) and *Halobacteria* (5%). *Methanomicrobia* was divided between  
394 *Methanoregula* (16%), *Methanosaeta* (58%) and *Methanolinea* (11%) genus's. *Thermoplasmata*  
395 was divided between Marine Benthic Group D (~~MBG-D~~) and ~~DHVEG-1~~ (5%) and Terrestrial  
396 Miscellaneous Gp(TMEG) (3%) family. *Halobacteria* dominant family was Deep Sea  
397 Hydrothermal Vent Gp 6(DHVEG-6) (1%). ~~Thaumarchaeota was divided mainly between two~~  
398 ~~classes:~~ Marine Benthic Group B (MBG-B 1%) and Miscellaneous Crenarchaeotic Group (MCG  
399 0.6%) (Fig. 4 [and supplementary table 1](#)). The detailed description of the major communities in the  
400 sediment is described at the discussion and the appendix. No ANME sequences were detected even  
401 though specific primers (ANME2C-AR468F, ANME3-1249, ANME1-395F, ANME1-1417,  
402 ANME3-140F, ANME3-1249, ANME2a-426 and ANME2a-1242R, see Miyashita et al., 2009)  
403 were used.

404

### 405 3.3. Depth comparison of microbial communities

406 An ordination plot from the 454 pyrosequencing by MOTHUR subsample OTUs dataset was  
407 derived from Principal coordinate's analysis (PCoA) of the bacterial and archaeal. The PCoA  
408 display the similarities and differences between the bacterial and archaeal communities varied with  
409 depth (Fig. 5a and 5b respectively). The PCoA of bacteria and archaea show very similar separation  
410 between the communities of different layers related to the different environment conditions of each  
411 layer. The two deeper layers are on the same distant on the X axis showing relatively more  
412 similarity than the upper layer. The vectors of the sulfate and sulfide are correlative with  
413 communities of the top layer, while methane,  $\delta^{13}\text{C}_{\text{TLE}}$  and  $\delta^{56}\text{Fe}(\text{II})$  were more associated to the  
414 communities of the middle layer. The communities in the bottom layer were more correlated to

415 Fe(II), Mn(II) and  $\text{NH}_4^+$ . The Venn diagram shows also that more OTUs from the deep layers were  
416 shared than between the upper layer and the deeper layers (Supplementary Fig 1 and 2).

### 417 **3.4. Functional gene profile**

418 In order to better understand the abundance of microbial functionality in the different depth we  
419 used qPCR for different functional genes. The functional genes *mcrA* and *dsrA* had very similar  
420 pattern - low concentrations in the upper layer ( $9 \times 10^5 \pm 6.4 \times 10^4$  and  $1 \times 10^6 \pm 1.7 \times 10^5$  respectively) and  
421 highest concentrations ( $6.9 \times 10^6 \pm 6.7 \times 10^5$  and  $6.9 \times 10^6 \pm 9.9 \times 10^4$ ) in the middle layer. *pmoA* gene  
422 showed also the same pattern, however with lower concentration ( $2.3 \times 10^5 \pm 9.7 \times 10^3$  to  
423  $1.6 \times 10^6 \pm 1.7 \times 10^5$ ) (Fig. 6).

424

## 425 **4. Discussion**

426 In this study of LK sediments we investigated changes in the microbial diversity associated  
427 with porewater geochemistry and the transition of the dominant electron acceptors with depth (0-  
428 40 cm). The geochemical porewater profiles in LK (Fig. 2) suggest that the sediment can be broadly  
429 divided into three different regions of microbial processes: the upper 5 cm which is dominated by  
430 sulfate reduction, the methanogenesis zone between 5 to 17 cm, and the deep sediments, dominated  
431 by anaerobic oxidation of methane coupled to iron reduction. This division of the sediment and the  
432 deep iron-driven AOM processes were confirmed by *in situ* profiles of methane,  $\delta^{13}\text{C}_{\text{CH}_4}$ , sulfate,  
433 and ferrous iron and subsequent use of a numerical mass conservation model (based on the  
434 geochemical profiles of DIC and  $\delta^{13}\text{C}_{\text{DIC}}$  (Adler et al., 2011) and a set of geochemical incubation  
435 experiments conducted in our previous work (Sivan et al., 2011).

436 The mechanisms that enable the process of AOM via iron reduction can be characterized by the  
437 existence of specific prokaryotic populations at each depth and their resemblance to similar  
438 environments with a distinct characterization. There are only few studies that analyzed microbial  
439 communities relative to geochemical zones in fresh water sediments (Deutzmann and Schink, 2011;  
440 Koizumi et al., 2004; Ye et al., 2009). Therefore, this study can deepen the information on  
441 microbial community shifts under different electron acceptors conditions, especially those related to  
442 the methane cycle. It is also the first study in LK that divides the sediment to different layers and  
443 attribute significantly to the diversity of the microbial ecology in the different zones.

444 Pipeline of SILVA ngs was used for the alpha-diversity to get more classified sequences and  
445 better coverage of the classification and population. However, for the beta-diversity the data from  
446 the MOTHUR pipeline applied to get more statistical results without the bias of different sample

447 size. The changes between the pipeline results for the alpha diversity were not large in the phyla  
448 level (standard deviation of ~0.5%) but increased with higher taxonomy levels.

#### 449 **4.1. Sediment geochemical conditions**

450 In the upper layer of sediment (0-3 cm), sulfate concentrations were the highest and decreased  
451 with depth as particulate organic matter from the water column accumulates, making organic carbon  
452 degradation available (Hadas and Pinkas, 1992). Adler et al. (2011) calculated that the bulk of  
453 bacterial sulfate reduction occurs in the top 1 cm of sediment, a finding that was supported by the  
454 microbial work of Hadas and Pinkas, (1992). Using sulfide and particulate organic carbon  
455 measurements, Eckert and Conrad. (2007) showed that sulfate reduction accounts for most of the  
456 mineralization of organic matter in the upper 15 cm of sediment. Methane concentrations in the  
457 upper layer were relatively low and probably reached from upward diffusion as methanogens are  
458 outcompeted by sulfate reducers there (Lovley and Klug, 1983). The depletion in methane  
459 concentrations and the decline  $\delta^{13}\text{C}_{\text{TLE}}$  values and the heavier isotopic values of  $\delta^{13}\text{C}_{\text{CH}_4}$  in the  
460 upper layer (Fig 2) compared to the methanogenic zone may be explained by AOM via sulfate  
461 reduction, although no ANME sequences were found using specific primers [and Fluorescent \*in situ\*](#)  
462 [hybridization methods](#) or [by our phylogenetic alignment of 16S rRNA gene libraries at any depth](#)  
463 in the sediment.

464 Below the sulfate reduction zone, in the middle layer of sediment (6-9 cm), methane reached its  
465 maximum concentrations (Fig. 2B). The low values of  $\delta^{13}\text{C}_{\text{CH}_4}$  in the methanogenesis zone are  
466 typical due to the large carbon isotope fractionation that occurs during methane production, thereby  
467 leaving the produced methane isotopically light and the DIC isotopically heavy. Below the  
468 maximum methane production zone (around 20 cm), methane concentrations began to decrease  
469 while  $\delta^{13}\text{C}_{\text{CH}_4}$  values increased (Fig. 2B), an outcome that may be due to the AOM process, which  
470 leaves residual methane isotopically heavier. Depleted  $\delta^{13}\text{C}_{\text{TLE}}$  (Fig. 2C) in the deep part of the  
471 sediment also supports the presence of AOM with the production of light biomass from methane  
472 oxidation. In addition, although all the suitable electron acceptors at this depth were depleted, both  
473 Fe(II) and Mn(II) increased (Fig. 2D). Also,  $\delta^{56}\text{Fe}$  values (Fig. 2C) of dissolved iron in the deep  
474 sediment were isotopically negative, similar to sediments with active dissimilatory bacterial iron  
475 reduction (Severmann et al., 2006) and are an indication of active iron reduction in the deep  
476 sediments and not just diffusion. The increase in Fe(II) concentrations below a depth of 15 cm was  
477 probably supported by the absence of sulfide. The potential of highly reactive Fe(III) oxides to drive  
478 AOM in LK was also supported by our set of mesocosm incubation studies (Sivan et al., 2011).  
479 Since manganese oxide concentrations were very low (~0.04%) throughout the LK sediment  
480 column (Serruya et al., 1974), iron oxide seemed to play a bigger role in this AOM process.

## 481 **4.2. Microbial population that could be involved in methane cycle**

### 482 *4.2.1. Deltaproteobacteria*

483 *Deltaproteobacteria* were the most abundant in the middle and deep sediment samples.  
484 Many members of the families of *Syntrophobacterales* are sulfate reducers or as fermentative  
485 organism, depending on the environmental conditions (McInerney et al., 2007). *Syntrophaceae*, the  
486 dominant family of *Syntrophobacterales*, are acetate-degrading sulfate-reducing bacterium (Jiang et  
487 al., 2009). *Desulfarculales* are strictly anaerobic mesophilic sulfate-reducing bacterium with the  
488 capability to oxidize acetate and fatty acids (Sun et al., 2010). *Desulfobacterales* are sulfate  
489 reducer, capable of oxidizing hydrogen in soils and sediments (Bürow et al., 2014).  
490 *Desulfuromonadales* was shown as  $S^0$ -respiring (Pjevac et al., 2014) and may also reduce Fe(III)  
491 and Mn(IV) in marine surface sediments (Lovley, 2006). About 1% of the sequences in the middle  
492 and bottom layers or the cores were affiliated to the *Geobacter* genus members which are involved  
493 in iron reduction. In the deep layer the most abundant class was Sva0485. Sequences of the Sva485  
494 order were similar to those found in different aquatic environments, but the metabolic functions of  
495 members of this order are not clear. *Pelobacter carbinolius* (Lovley et al., 1995), a member of  
496 Sva485 is capable of Fe(III) and sulfur reduction. Schwarz et al. (2007a) showed that in the upper  
497 part of LK sediment *Deltaproteobacteria* was one of the dominant classes, and that most of them  
498 were affiliated with acetate-oxidizing sulfate-reducing bacteria, that outcompete the acetoclastic  
499 methanogens. Even though *Deltaproteobacteria* are best known for their sulfate reduction  
500 metabolism, they can shift their metabolism in response to depleted sulfate concentrations (Plugge  
501 et al., 2011).

### 502 *4.2.2. Betaproteobacteria*

503 *Betaproteobacteria* percentages increased slightly with depth. These percentages are comprised of  
504 chemoheterotrophs and chemoautotrophs that derive nutrients from decomposition of organic  
505 material. *Burkholderiales* was the dominant order in *Betaproteobacteria* and increased with depth  
506 (1.5% to 4%). (Schwarz et al., 2007b) showed that *Burkholderiales* is active in the upper sediment  
507 of LK. Members of this group are able to carry out anaerobic oxidation of acetate with (per)chlorate  
508 as electron acceptor (Yoshida et al., 2005) or oxidize hydrogen in deep subsurface (Orcutt et al.,  
509 2011).

### 510 *4.2.3. Gammaproteobacteria*

511 Our *Gammaproteobacteria* affiliated sequences have different phylogenetic classification  
512 distribution in each layer. In the middle and bottom layers members of the *Methylococcales* order  
513 were observed (~1%). *Methylococcales* are aerobic methanotrophs which some members were  
514 present within the sediments and overlying water column from dysoxic, methane-rich vent and seep  
515 systems (Tavormina et al., 2008). The most abundant order in the deep layer was HOC36 (5%)



516 which is an uncultured *Gammaproteobacteria*. however when compared to NCBI data base it was  
517 found to be closely related to uncultured LK clones (99%) and to cultured *Methylocaldum* sp. (94%)  
518 (Bodrossy et al., 1997) which is a Thermophilic methanotroph isolated from landfill cover soil.

#### 519 4.2.4. *Nitrospirae*

520 Members of the *Nitrospirae* phylum, increased with depth. Sequences from our samples were  
521 classified to *Nitrospirales* order in *Nitrospirea* and were most abundant in the bottom layer of the  
522 sediment (7%). Part of our sequences were similar to those found in a previous study (Schwarz et  
523 al., 2007b). . Using SIP-RNA, (Schwarz et al., 2007b) showed not only that *Nitrospirae* were  
524 present, but also that they were functionally bioactive. Although *Nitrospirae* is a known nitrate  
525 oxidizer (Ehrich et al., 1995), the conditions of this environment suggest that it utilized another  
526 metabolic pathway. *Nitrospirae* also include the iron reducing candidates such as  
527 *Magnetobacterium bavaricum* (Spring et al., 1993) and sulfur reducers (Sonne Hansen and Ahring,  
528 1999). Part of our sequences were aligned to uncultured clone (98%) from freshwater sediment  
529 which enhanced degradation of phenanthrene and pyrene by amorphous ferric hydroxide (Yan et  
530 al., 2012).

#### 531 4.2.5. *Methanomicrobia*

532 The majority of our sequences were classified within the *Euryarchaeota*, in the  
533 *Methanomicrobia* class. Our 16S rRNA sequences were divided between the *Methanomicrobiales*  
534 and *Methanosarcinales* order. *Methanosaeta* genus of *Methanosarcinales* increases with depth.  
535 *Methanosaeta* are acetoclastic methanogens which only able to grow on acetate even at low  
536 concentrations (Jetten et al., 1990). *Methanosaeta* was also shown in previous studies of LK  
537 (Schwarz et al., 2007a, 2007b) and in other meso to eutrophic freshwater lakes (Glissman et al.,  
538 2004; Koizumi et al., 2003). Yamada et al. (2014) showed that *Methanosaeta* can also have the  
539 ability to reduce ferrihydrite with H<sub>2</sub> as the electron source. The *Methanosaeta concilii* was the  
540 most similar cultured acetoclastic methanogen (96%) to our sequences which was observed also in  
541 the previous study. The *Methanolinea* genus of *Methanomicrobiales* a hydrogenotrophic  
542 methanogen was observed in constant percentages for all depths. Members of the *Methanoregula*  
543 genus, a genus that is also hydrogenotrophic methanogen, decreased with depth in the core.  
544 *Methanoregula*, *Methanosaeta* and *Methanolinea* represent 60% of sequences of methanogens  
545 retrieved from freshwater lakes (Borrel et al., 2011).

#### 546 4.2.6. *Thermoplasmata*

547 The dominant family in *Thermoplasmata* is The Marine Benthic Group D and DHVEG-1  
548 (MBG-D). Members of the MBG-D have been shown to exist in a variety of freshwater and marine  
549 environments (Beal et al., 2009; Borrel et al., 2012), and it is the most widely encountered,  
550 uncultured lineage in freshwater lake sediments. Even though their metabolism is unknown,

551 hypotheses about their functionalities are based on the environments in which they were found.  
552 Methanogenesis was suggested, as they were found in deep lake sediments with high methane  
553 concentrations (Borrel et al., 2012), and they were also hypothesized to be involved in AOM, as  
554 they were found in AOM zones (Schubert et al., 2011) and in marine seep sediment (Beal et al.,  
555 2009). However, in other environments, in which methane concentrations were low, the utilization  
556 of waste products, intermediates, or dead cells by MBG-D was also suggested (Smith et al., 1975).  
557 Recently, a single cell genomics study showed that members of MBG-D were capable of exogenous  
558 protein degradation in cold anoxic environments (Lloyd et al., 2013).

#### 559 4.2.7. *Thaumarchaeota*

560 *Thaumarchaeota* are mostly known as archaeal ammonia oxidizers but in the sediments they may  
561 contribute significantly to the reservoir of nitrogen oxides in ocean waters and thus to productivity,  
562 including the assimilation of carbon (Pester et al., 2011). The enzyme ammonia monooxygenase,  
563 which belongs to the enzyme family of copper containing membrane bound monooxygenases that  
564 possess wide substrate ranges. Closely related enzymes, which can often perform the same function  
565 but with different substrates, include ammonia oxidizing bacteria with ammonia (in  
566 *Gammaproteobacteria*) and methane as a substitute (Lontoh et al., 2000). However, a  
567 microorganism's metabolic energy is mainly defined by its downstream enzyme machinery  
568 (Tavormina et al., 2011). In our sediment *Thaumarchaeota* observed in low percentages with 2  
569 main classes: Marine Benthic Group B and Miscellaneous Crenarchaeotic Group.

#### 570 4.2.8. Miscellaneous Crenarchaeotic Group

571 The Miscellaneous Crenarchaeotic Group (MCG) is a cosmopolitan class assigned to the  
572 *Thaumarchaeota* phylum whose members are found mostly in anoxic habitats and have the  
573 capability to take up organic carbon (Biddle et al., 2006). Because of their huge environmental  
574 range and their complex phylogeny, the MCG collectively possess great metabolic diversity (Jiang  
575 et al., 2008). Current evidence for members of the MCG lineage suggests that they may obtain  
576 energy from the anaerobic oxidation of methane, but they do so via a dissimilatory methane-  
577 oxidizing process and they do not assimilate its carbon (Biddle et al., 2006)

#### 578 4.2.9. Marine Benthic Group B

579 Members of the MBGB were detected in a growing number of benthic marine environments,  
580 including Atlantic deep sea sediments, marine mud volcanoes, marine carbonate crusts from the  
581 Black Sea, organic poor deep subsurface sediments of the central oceanic basins, and organic rich  
582 methane or methane hydrate containing sediments near continental margins (Sørensen and Teske,  
583 2006). Recent results from Ocean Drilling Program showed that no known methantroph was  
584 detected however methane oxidation appeared to be mediated by Marine Benthic Group B and the  
585 Miscellaneous Crenarchaeotal Group (Biddle et al., 2006).

586

## 4.2 Concept for methane cycle mechanisms

587 A few possible mechanisms for the novel process of iron driven AOM in the deep part of the  
588 sediment can be suggested based on the geochemical profiles and the microbial communities. A  
589 possible direct process could be through new, currently unknown bacteria/archaea that reduce iron  
590 and utilize methane. ~~Possible indirect processes~~ could include: 1) Reduction of Fe(III) oxides by  
591 oxidation of sulfide (in pyrite or FeS minerals) to sulfur intermediates, and then disproportionation  
592 to sulfide and sulfate (such as in (Holmkvist et al., 2011)), and/or sulfate driven AOM but without  
593 ANME. 2) Reduction of Fe(III) by utilizing H<sub>2</sub> (Lovley, 1991), could consume most of produced  
594 H<sub>2</sub> in the deep sediment. Thus which ~~creates~~ ing a low concentration of H<sub>2</sub> ~~and which~~ drives  
595 methanogenesis backwards ("reverse methanogenesis") (Hallam et al., 2004; Hoehler et al., 1994)  
596 by the methanogens ~~archaea~~ in the deep sediment. 3) An oxygenic methane oxidation pathway in an  
597 anaerobic environment as described by Ettwig et al. (2010), when methane is oxidized by oxygen  
598 that is released from iron oxides.

599 Changes in ~~t~~The microbial population shown in the present study ~~changes can be used~~ may ~~to~~  
600 support the deep iron driven AOM mechanism (presented by the sample from 29-32cm). The  
601 bacterial diversity in this depth was lower than in the methanogenesis zone (6-9 cm) but higher than  
602 in the upper layer (0-3 cm) of the sediment. The changes in bacterial diversity with depth could be  
603 related to the availability of different electron acceptors and of organic matter (Nam et al., 2008).  
604 However, there is an overlap between bacterial and archaeal sequences in the middle and the bottom  
605 layers (PCoA and Venn diagram), indicating that the environmental conditions are affecting the  
606 processes. Alpha diversity indexes (Table 1 and 2) were compared to previous studies conducted  
607 only in the upper layer of the sediment of LK using different methods (Schwarz et al., 2007a;  
608 Wobus et al., 2003), and showed a similarity in the upper parts of the sediment. However, the  
609 deeper layers in this study showed higher diversity and richness than the previous studies (Nusslein  
610 et al., 2001; Schwarz et al., 2007a, 2007b) and was close to other studies using 454 sequencing  
611 (Hollister et al., 2010; Röske et al., 2012).

612 The direct mechanism of iron reduction coupled with methane oxidation could be performed by  
613 a single microorganism ~~can occur by one microorganism~~, as methanogens have the ability to reduce  
614 iron (Van Bodegom et al., 2004; Bond et al., 2002) however not with methane as the electron donor.  
615 ~~According to our results, this kind of~~ This direct mechanism could be performed by unique  
616 methanotrophy ~~could be from~~ like MBG-D (*Thermoplasmata* class) or ~~by~~ by MCG/MBG-B  
617 (*Thaumarchaeota* phylum ~~which that~~ MCG may represent as new archaeal phylum (Lloyd et al.,  
618 2013) or as a sister lineage with *Thaumarchaeota* (Meng et al., 2014). However MBG-B is still  
619 classified as *Thaumarchaeota* (Marlow et al., 2014), to which some of our sequences were similar

620 ~~to. The representation of sequences similar to~~About 5% of the archaeal sequences in the middle and  
621 ~~deep layers were affiliated to MBG-D, were close to 5% of the archaeal community in the middle~~  
622 ~~and deep layers, however~~ MCG and MBG-B showed much lower representation and ~~will not be thus~~  
623 ~~are not~~ discussed. ~~the~~ Members of the MBG-D have been shown to exist in a variety of freshwater  
624 and marine environments (Beal et al., 2009; Borrel et al., 2012), and it is the most widely  
625 encountered, uncultured lineage in freshwater lake sediments. Even though their metabolism is  
626 unknown, hypotheses about their functionalities are based on the environments in which they were  
627 found. Methanogenesis was suggested, as they were found in deep lake sediments with high  
628 methane concentrations (Borrel et al., 2012), and they were also hypothesized to be involved in  
629 AOM, as they were found in AOM zones (Schubert et al., 2011) and in marine seep sediment (Beal  
630 et al., 2009). ~~However, in~~In other environments, in which methane concentrations were low, the  
631 utilization of waste products, intermediates, or dead cells by MBG-D was also suggested (Smith et  
632 al., 1975). Recently, a single cell genomics study showed that members of MBG-D were capable of  
633 exogenous protein degradation in cold anoxic environments (Lloyd et al., 2013). ~~Although their~~  
634 ~~metabolisms are unknown, these groups were found in methane rich environments.~~

635 Another option is a consortium of ~~iron reducing microbes~~ ~~microbes including those belonging~~  
636 ~~Geobacter~~ genus, which are known for iron reduction, or ~~Nitrospirales~~, known to reduce iron,  
637 together with methanotrophs or with methanogens capable of "reverse methanogenesis".

638 The well-known iron reducers are *Geobacter* genus, however only about 1% of the sequences  
639 in the middle and bottom layers or the cores were affiliated to the *Geobacter* genus members. In the  
640 deep layer the most abundant class was Sva0485. Sequences of the Sva485 order were similar to  
641 those found in different aquatic environments, but the metabolic functions of members of this order  
642 are not clear. *Pelobacter carbinolious* (Lovley et al., 1995), a member of Sva485 is capable of  
643 Fe(III) and sulfur reduction. *Desulfuromonadales*, which was found in our sequences in the deep  
644 parts of the sediment, was shown as  $S^0$  respiring (Pjevac et al., 2014) and may also reduce Fe(III)  
645 and Mn(IV) in marine surface sediments (Lovley, 2006). Schwarz et al. (2007a) showed that in the  
646 upper part of LK sediment *Deltaproteobacteria* was one of the dominant classes, and that most of  
647 them were affiliated with acetate oxidizing sulfate reducing bacteria, that outcompete the  
648 acetoclastic methanogens. Even though *Deltaproteobacteria* are best known for their sulfate  
649 reduction metabolism, they can shift their metabolism in response to depleted sulfate concentrations  
650 (Plugge et al., 2011).

651 In addition, our ~~The~~ results suggest that *Desulfuromonadales* of *Deltaproteobacteria* class  
652 could be also involved in indirect mechanisms of disproportionation of sulfur together with other  
653 *Deltaproteobacteria* sulfate reducer. The *dsrAB* functional gene shows that sulfate reducers are  
654 present at the same level of the core where *mcrA* functional gene (Fig. 6) of methanogens or

655 methanotrophs are found in the deep part of the sediment. Additionally, the presence of the  
656 *Deltaproteobacteria* at the deepest sediment depth and the observed accumulation of acetate with  
657 depth (data not shown) could indicate their exploitation of a different metabolic path in the deep  
658 sediment than in the upper part of the sediment.

659 Another options for functioning iron reducers are include members of the *Nitrospirae* phylum  
660 which—that increased with depth. Indeed, Ssequences from our samples were classified to  
661 *Nitrospirales* order in *Nitrospirea* and were most abundant in the bottom layer of the sediment  
662 (7%). Part of our sequences were similar to those found in a previous study (Schwarz et al., 2007b).  
663 Using SIP-RNA, (Schwarz et al., (2007e2007b) showed not only that *Nitrospirae* were present, but  
664 also that they were functionally bioactive. Although *Nitrospirae* is a known nitrate oxidizer (Ehrich  
665 et al., 1995), the conditions of this environment suggest that it utilized another metabolic pathway.  
666 *Nitrospirae* also include the iron reducing candidates such as *Magnetobacterium bavaricum* (Spring  
667 et al., 1993) and sulfur reducers (Sonne-Hansen and Ahring, 1999). Part of our sequences were  
668 aligned to uncultured clone (98%) from freshwater sediment which enhanced degradation of  
669 phenanthrene and pyrene by amorphous ferric hydroxide (Yan et al., 2012).

670 The "reverse methanogenesis" pathway could be carried out by the dominant deep layer  
671 methanogens *Methanosaeta*-. *Methanosaeta* are acetoclastic methanogens, which are ~~only~~ able to  
672 grow only on acetate; even at low concentrations (Jetten et al., 1990). *Methanosaeta* was also  
673 shown ~~asto be~~ the dominant active methanogens in the upper layer in previous studies of LK  
674 (Schwarz et al., 2007a, 2007b) and in other meso to eutrophic freshwater lakes (Glissman et al.,  
675 2004; Koizumi et al., 2003). Yamada et al. (2014) showed that *Methanosaetacan* also have the  
676 ability to reduce ferrihydrite with H<sub>2</sub> as the electron source. The *Methanosaeta concilii* was the  
677 most similar cultured acetoclastic methanogen (96%) to our sequences which was observed also in  
678 the previous study. ~~which have been shown in a previous study as the dominant active methanogens~~  
679 in the upper layer (Schwarz et al., 2007b). *Methanosaeta* could perform a different metabolic  
680 process or may have been inhibited. The other methanogens that could perform the "revers  
681 methanogenesis" are from the *Methanomicrobiales* order. The *Methanolinea* genus of  
682 *Methanomicrobiales* a hydrogenotrophic methanogen was observed in constant percentages for all  
683 depths. Members of the *Methanoregula* genus, a genus that is also hydrogenotrophic methanogen,  
684 decreased with depth in the core. qPCR analysis of *mcrA* functional gene of methanogens/anaerobic  
685 methanotroph shows that in the middle layer methanogens were more abundant than in the upper  
686 and bottom layers (Fig 6). ANME were not found in the sediment using specific primers and  
687 Fluorescent *in situ* hybridization methods. An additional reason for not finding any ANME  
688 sequences in our samples is likely because 454 sequences related to methanogens were found in

689 high percentages and they are probably that dominant in the qPCR analysis of *mcrA* gene observed  
690 in the deep layer ~~belongs to the methanogens which maybe cause the reverse methanogenesis.~~

691 ~~The results suggest that *Desulfuromonadales* of *Deltaproteobacteria* class could be involved in~~  
692 ~~indirect mechanisms of disproportionation of sulfur together with other *Deltaproteobacteria* sulfate~~  
693 ~~reducer. The *dsrAB* functional gene shows that sulfate reducers are present at the same level of the~~  
694 ~~core where *mcrA* functional gene (Fig. 6) of methanogens or methanotrophs are found in the deep~~  
695 ~~part of the sediment. Additionally, the presence of the *Deltaproteobacteria* at the deepest sediment~~  
696 ~~depth and the observed accumulation of acetate with depth (data not shown) could indicate their~~  
697 ~~exploitation of a different metabolic path in the deep sediment than in the upper part of the~~  
698 ~~sediment.~~

699 The ~~third~~ additional indirect mechanism of anaerobic methane oxidation via an oxygenic  
700 pathway was shown clearly by the *pmoA* functional gene. The *pmoA* concentration in the deep part  
701 of the sediment was higher than in the upper part, indicating an oxygenic pathway (Fig. 6).  
702 However, pMMO is a homologue enzyme of ammonia monooxygenase and might be sequenced  
703 together with pMMO (Tavormina et al., 2011). The source of pMMO could be *Methylococcales*  
704 (~1%), of *Gammaproteobacteria*), which were observed in the middle and bottom sediment layers.  
705 *Methylococcales* are aerobic methanotrophs, which some members were present within the  
706 sediments and overlying water column from dysoxic, methane-rich vent and seep systems  
707 (Tavormina et al., 2008). The most abundant order in the deep layer was HOC36 (5%), which is an  
708 uncultured *Gammaproteobacteria*. However, when compared to NCBI data base, it was found to  
709 be closely related to uncultured LK clones (99%) and to cultured *Methylocaldum* sp. (94%)  
710 (Bodrossy et al., 1997), which is a Thermophilic methanotroph isolated from landfill cover soil.  
711 ~~(*Gammaproteobacteria*), which is an aerobic methanotroph found in the deep part of our samples.~~  
712 However, bias of *pmoA* with ammonia monooxygenase could be ~~eased~~ caused by *Thaumarchaeota*,  
713 Betaproteobacteria or Nitrospirae, which were found in all depth.

714 *Thaumarchaeota* comprise not only all known archaeal ammonia oxidizers, but also several  
715 clusters of environmental sequences representing microorganisms with unknown energy  
716 metabolisms (Pester et al., 2011). Members of *Thaumarchaeota* phylum could have monooxygenases  
717 like enzymes that are able to capture methane due to the enzymes' phylogenetic proximities to  
718 methane monooxygenases, but that activity requires the necessary downstream metabolic pathway. If  
719 they could have that ability, they would have been good candidates for group of methanotrophic  
720 archaea. This indirect pathway can be similar to that found in the NC10 phylum (Zhu et al., 2012),  
721 which produces oxygen via the reduction of nitrite and the oxidation of methane, but with iron  
722 oxides. Also many bacterial ammonia oxidizer sequences were found in our environment.  
723 Ammonium profiles show increase with depth mainly due to decomposition of the organic matter,

724 and theoretically the large amount of ammonium could be oxidized by ferric iron minerals and  
725 produce nitrite (Clement et al., 2005). Maybe ammonia monooxygenase can function for  
726 ammonium up take for iron reduction and for methane up take for oxidation, but that is only a  
727 speculation. However, no nitrate or nitrite was detected in the deep sediment, moreover no NC10  
728 phylum (Ettwig et al., 2010) was observed in our sequences.

729 To summarize, this study attempted to correlate between the performed geochemical and  
730 microbial profiles in lake sediments. The geochemical data suggest three main depth related zones  
731 of electron acceptor activities in the sediment: sulfate reduction, methanogenesis and a novel, deep  
732 iron-driven AOM. The prokaryotic analysis provided clues regarding the microorganisms that may  
733 be involved in this novel process and the metabolic paths that occur throughout the microbial  
734 assemblage. For AOM via iron reduction to occur, a number of potential pathways and their  
735 combinations have been suggested. Orders that become enriched (*Sva0485*, *Methanosarcinales* and  
736 *Nitrospirales*) with depth can be assumed to participate in the AOM process either directly or  
737 indirectly. A possible direct process could be through new, currently unknown bacteria/archaea that  
738 reduce iron and utilize methane, which may be carried out by a [MBG-D MCG](#) as a methanotroph in  
739 a consortium with an iron reducer like *Geobacter*. Possible indirect processes could be Fe(III)  
740 reduction by sulfide, oxidation of the sulfide to elemental sulfur and other sulfur intermediates and  
741 then disproportionation to sulfide and sulfate and sulfate driven AOM. However, this process is less  
742 likely because ANMEs were not found. It could also be via reduction of Fe(III) by utilizing H<sub>2</sub>,  
743 creating a low concentration of H<sub>2</sub> and ~~driveing~~driving “reverse methanogenesis”. Fe(III) reduction  
744 processes could be carried out by *Nitrospirae* and/or *Deltaproteobacteria*. Members of both groups  
745 can reduce iron while in a consortium with methanogenic *Methanosarcinales*/[Methanomicrobiales](#),  
746 or can reduce the iron with sulfur minerals, creating sulfate (see above) that *Deltaproteobacteria*  
747 can utilize while in a consortium with a [MBG-D MCG](#) as a methanotrophic archaea ([Schubert et al.,](#)  
748 [2011](#)). An oxidation of methane coupled to iron reduction pathway as described by Ettwig et al.,  
749 (2010) could occur by *Thaumarchaeota* or *Nitrospirae* with a monooxygenase enzyme that can  
750 utilize methane while using iron oxides to generate the oxygen needed to oxidize the methane.  
751 Further research involving even larger sample of the microbial community and characterization of  
752 more diverse functional genes will provide better indication for the composition of the microbial  
753 communities at the different depths. Also, microbial and geochemical experiments have the  
754 potential to give more clues on the mechanism of this novel iron driven AOM.

755

## 756 **Acknowledgments**

757 We [would like to](#) thank [the editor and the reviewers for the helpful and constructive comments.](#) We  
758 [thank](#) M. Adler for her assistance in the field and in the laboratory. Thanks to the members of Orit's  
759 and Ariel's laboratories for all the help. Many thanks to V. Orphan, S. Connon and K. Dawson from  
760 Caltech for their help and ~~for their~~ fruitful discussions. This research was funded by the Water  
761 Authority of Israel (O.S. and W.E.). [E. Kramarsky-Winter is gratefully acknowledged for the](#)  
762 [thorough scrutiny of the manuscript and for heris critical and helpful comments.](#)

763

764

## 765 **References**

- 766 Adler, M., Eckert, W. and Sivan, O.: Quantifying rates of methanogenesis and methanotrophy in  
767 Lake Kinneret sediments (Israel) using pore-water profiles, *Limnol. Oceanogr.*, 56(4), 1525–1535,  
768 2011.
- 769 Beal, E. J., House, C. H. and Orphan, V. J.: Manganese- and Iron-Dependent Marine Methane  
770 Oxidation, *Science*, 325(5937), 184–187, 2009.
- 771 Bligh, E. G. and Dyer, W. J.: A rapid method of total lipid extraction and purification, *Can. J.*  
772 *Biochem. Physiol.*, 37(8), 911–917, 1959.
- 773 Van Bodegom, P. M., Scholten, J. C. M. and Stams, A. J. M.: Direct inhibition of methanogenesis  
774 by ferric iron., *FEMS Microbiol. Ecol.*, 49(2), 261–8, 2004.
- 775 Bodrossy, L., Holmes, E. M., Holmes, A. J., Kovács, K. L. and Murrell, J. C.: Analysis of 16S  
776 rRNA and methane monooxygenase gene sequences reveals a novel group of thermotolerant, *Arch*  
777 *microbiol.*, 168, 493–503, 1997.
- 778 Boetius, A., Ravensschlag, K., Schubert, C. J., Rickert, D., Widdel, F., Gieseke, A., Amann, R.,  
779 Jørgensen, B. B., Witte, U. and Pfannkuche, O.: A marine microbial consortium apparently  
780 mediating anaerobic oxidation of methane, *Nature*, 407(6804), 623–626, 2000.
- 781 Bond, D. R., Lovley, D. R. and Methanococcus, A.: Reduction of Fe ( III ) oxide by methanogens in  
782 the presence and absence of extracellular quinones, *Environ. Microbiol.*, 4(2), 115–124, 2002.
- 783 Borowski, W. S., Cagatay, N., Ternois, Y. and Paull, C. K.: Data report: carbon isotopic  
784 composition of dissolved CO<sub>2</sub>, CO<sub>2</sub> gas, and methane, Blake-Bahama Ridge and northeast  
785 Bermuda Rise, ODP Leg 172, Proc. Ocean Drill. Program, *Sci. Results*, 172, 1–16, 2000.
- 786 Borrel, G., Lehours, A.-C., Crouzet, O., Jézéquel, D., Rockne, K., Kulczak, A., Duffaud, E., Joblin,  
787 K. and Fonty, G.: Stratification of Archaea in the deep sediments of a freshwater meromictic lake:  
788 vertical shift from methanogenic to uncultured archaeal lineages., *PLoS One*, 7(8), e43346, 1–14,  
789 2012.



- 790 Borrok, D. M., Wanty, R. B., Ridley, W. I., Wolf, R., Lamothe, P. J. and Adams, M.: Separation of  
791 copper, iron, and zinc from complex aqueous solutions for isotopic measurement, *Chem. Geol.*,  
792 242(3-4), 400–414, 2007.
- 793 Chao, A.: Nonparametric Estimation of the Number of Classes in a Population, *Scand. J. Stat.*,  
794 11(4), 265–270, 1984.
- 795 Chao, B. Y. A. and Ma, M.: Stopping rules and estimation for recapture debugging with unequal  
796 failure rates, *Biometrika*, 80(1), 193–201, 1993.
- 797 Chistoserdova, L., Vorholt, J. A. and Lidstrom, M. E.: A genomic view of methane oxidation by  
798 aerobic bacteria and anaerobic archaea, *Genome Biol.*, 6(2), 208.1–208.6, 2005.
- 799 Clement, J., Shrestha, J., Ehrenfeld, J. and Jaffe, P.: Ammonium oxidation coupled to dissimilatory  
800 reduction of iron under anaerobic conditions in wetland soils, *Soil Biol. Biochem.*, 37(12), 2323–  
801 2328, doi:10.1016/j.soilbio.2005.03.027, 2005.
- 802 Cline, J. D.: Spectrophotometric determination of hydrogen sulfide In natural waters, *Limnol.*  
803 *Oceanogr.*, 14, 454–458, 1969.
- 804 Deutzmann, J. S. and Schink, B.: Anaerobic oxidation of methane in sediments of Lake Constance,  
805 an oligotrophic freshwater lake., *Appl. Environ. Microbiol.*, 77(13), 4429–36,  
806 doi:10.1128/AEM.00340-11, 2011.
- 807 Dowd, S. E., Callaway, T. R., Wolcott, R. D., Sun, Y., McKeehan, T., Hagevoort, R. G. and  
808 Edrington, T. S.: Evaluation of the bacterial diversity in the feces of cattle using 16S rDNA  
809 bacterial tag-encoded FLX amplicon pyrosequencing (bTEFAP)., *BMC Microbiol.*, 8(125), 1–8,  
810 2008.
- 811 Eckert, T.: The Influence of Chemical Stratification in the Water Column on Sulfur and Iron  
812 Dynamics in Pore Waters and Sediments of Lake Kinneret, Israel, M.Sc. thesis, University of  
813 Bayreuth, Germany., 2000.
- 814 Eckert, W. and Conrad, R.: Sulfide and methane evolution in the hypolimnion of a subtropical lake:  
815 A three-year study, *Biogeochemistry*, 82(1), 67–76, 2007.
- 816 Edgar, R. C., Haas, B. J., Clemente, J. C., Quince, C. and Knight, R.: UCHIME improves  
817 sensitivity and speed of chimera detection., *Bioinformatics*, 27(16), 2194–2200, 2011.
- 818 Egger, M., Rasigraf, O., Sapart, C. J., Jilbert, T., Jetten, M. S. M., Röckmann, T., van der Veen, C.,  
819 Banda, N., Kartal, B., Ettwig, K. F. and Slomp, C. P.: Iron-mediated anaerobic oxidation of  
820 methane in brackish coastal sediments., *Environ. Sci. Technol.*, 49, 277–283, 2014.
- 821 Ehrich, S., Behrens, D., Lebedeva, E., Ludwig, W. and Bock, E.: A new obligately  
822 chemolithoautotrophic, nitrite-oxidizing bacterium, *Nitrospira moscoviensis* sp. nov. and its  
823 phylogenetic relationship., *Arch. Microbiol.*, 164(1), 16–23, 1995.
- 824 Ettwig, K. F., Van Alen, T., van de Pas-Schoonen, K. T., Jetten, M. S. M. and Strous, M.:  
825 Enrichment and molecular detection of denitrifying methanotrophic bacteria of the NC10 phylum,  
826 *Appl. Environ. Microbiol.*, 75(11), 3656–3662, 2009.

- 827 Ettwig, K. F., Butler, M. K., Le Paslier, D., Pelletier, E., Mangenot, S., Kuypers, M. M. M.,  
828 Schreiber, F., Dutilh, B. E., Zedelius, J. and de Beer, D.: Nitrite-driven anaerobic methane oxidation  
829 by oxygenic bacteria, *Nature*, 464(7288), 543–548, 2010.
- 830 Froelich, P. N., Klinkhammer, G. P., Bender, M. L., Luedtke, N. A., Heath, G. R., Cullen, D.,  
831 Dauphin, P., Hammond, D., Hartman, B. and Maynard, V.: Early oxidation of organic matter in  
832 pelagic sediments of the eastern equatorial Atlantic: suboxic diagenesis, *Geochim. Cosmochim.*  
833 *Acta*, 43(7), 1075–1090, 1979.
- 834 Glissman, K., Chin, K.-J., Casper, P. and Conrad, R.: Methanogenic pathway and archaeal  
835 community structure in the sediment of eutrophic Lake Dagow: effect of temperature., *Microb.*  
836 *Ecol.*, 48(3), 389–399, 2004.
- 837 Good, I. J.: The population frequencies of species and the estimation of population parameters,  
838 *Biometrika*, 40(3-4), 237–264, 1953.
- 839 Hadas, O. and Pinkas, R.: Sulfate-reduction process in sediments of Lake Kinneret, Israel,  
840 *Hydrobiologia*, 235(1), 295–301, 1992.
- 841 Hallam, S. J., Girguis, P. R., Preston, C. M., Richardson, P. M. and DeLong, E. F.: Identification of  
842 methyl coenzyme M reductase A (*mcrA*) genes associated with methane-oxidizing archaea, *Appl.*  
843 *Environ. Microbiol.*, 69(9), 5483–5491, 2003.
- 844 Hallam, S. J., Putnam, N., Preston, C. M., Detter, J. C., Rokhsar, D., Richardson, P. M. and  
845 DeLong, E. F.: Reverse methanogenesis: testing the hypothesis with environmental genomics,  
846 *Science*, 305(5689), 1457–1462, 2004.
- 847 Hinrichs, K. U., Hayes, J. M., Sylva, S. P., Brewer, P. G. and DeLong, E. F.: Methane-consuming  
848 archaeobacteria in marine sediments, *Nature*, 398(6730), 802–805, 1999.
- 849 Hoehler, T. M., Alperin, M. J., Albert, D. B. and Martens, C. S.: Field and laboratory studies of  
850 methane oxidation in an anoxic marine sediment: evidence for a methanogen-sulfate reducer  
851 consortium, *Global Biogeochem. Cycles*, 8(4), 451–463, 1994.
- 852 Holler, T., Wegener, G., Knittel, K., Boetius, A., Brunner, B., Kuypers, M. M. M. and Widdel, F.:  
853 Substantial  $^{13}\text{C}/^{12}\text{C}$  and D/H fractionation during anaerobic oxidation of methane by marine  
854 consortia enriched in vitro, *Environ. Microbiol. Rep.*, 1(5), 370–376, 2009.
- 855 Hollister, E. B., Engledow, A. S., Hammett, A. J. M., Provin, T. L., Wilkinson, H. H. and Gentry,  
856 T. J.: Shifts in microbial community structure along an ecological gradient of hypersaline soils and  
857 sediments., *ISME J.*, 4(6), 829–838, 2010.
- 858 Holmkvist, L., Ferdelman, T. G. and Jørgensen, B. B.: A cryptic sulfur cycle driven by iron in the  
859 methane zone of marine sediment (Aarhus Bay, Denmark), *Geochim. Cosmochim. Acta*, 75(12),  
860 3581–3599, 2011.
- 861 Huse, S. M., Welch, D. M., Morrison, H. G. and Sogin, M. L.: Ironing out the wrinkles in the rare  
862 biosphere through improved OTU clustering., *Environ. Microbiol.*, 12(7), 1889–1898, 2010.
- 863 Jetten, M. S. M., Stams, A. J. M. and Zehnder, A. J. B.: Acetate threshold values and acetate  
864 activating enzymes in methanogenic bacteria, *FEMS Microbiol. Lett.*, 73(4), 339–344, 1990.

- 865 John, S. G. and Adkins, J. F.: Analysis of dissolved iron isotopes in seawater, *Mar. Chem.*, 119(1-  
866 4), 65–76, 2010.
- 867 Kinnaman, F. S., Valentine, D. L. and Tyler, S. C.: Carbon and hydrogen isotope fractionation  
868 associated with the aerobic microbial oxidation of methane, ethane, propane and butane, *Geochim.*  
869 *Cosmochim. Acta*, 71(2), 271–283, 2007.
- 870 Klein, M., Friedrich, M., Roger, a J., Hugenholtz, P., Fishbain, S., Abicht, H., Blackall, L. L., Stahl,  
871 D. a and Wagner, M.: Multiple lateral transfers of dissimilatory sulfite reductase genes between  
872 major lineages of sulfate-reducing prokaryotes., *J. Bacteriol.*, 183(20), 6028–6035, 2001.
- 873 Knittel, K. and Boetius, A.: Anaerobic oxidation of methane: progress with an unknown process.,  
874 *Annu. Rev. Microbiol.*, 63, 311–334, 2009.
- 875 Koizumi, Y., Takii, S. and Fukui, M.: Depth-related change in archaeal community structure in a  
876 freshwater lake sediment as determined with denaturing gradient gel electrophoresis of amplified  
877 16S rRNA genes and reversely transcribed rRNA fragments., *FEMS Microbiol. Ecol.*, 48(2), 285–  
878 292, 2004.
- 879 Koizumi, Y., Takii, S., Nishino, M. and Nakajima, T.: Vertical distributions of sulfate-reducing  
880 bacteria and methane-producing archaea quantified by oligonucleotide probe hybridization in the  
881 profundal sediment of a mesotrophic lake., *FEMS Microbiol. Ecol.*, 44(1), 101–108, 2003.
- 882 Lloyd, K. G., Schreiber, L., Petersen, D. G., Kjeldsen, K. U., Lever, M. a., Steen, A. D.,  
883 Stepanauskas, R., Richter, M., Kleindienst, S., Lenk, S., Schramm, A. and Jørgensen, B. B.:  
884 Predominant archaea in marine sediments degrade detrital proteins, *Nature*, 496(7444), 215–218,  
885 2013.
- 886 Lovley, D.: Dissimilatory Fe ( III ) - and Mn ( IV ) -Reducing Prokaryotes Microorganisms, in  
887 *Prokaryotes*, pp. 635–658, Springer New York, New York., 2006.
- 888 Lovley, D. R.: Dissimilatory Fe(III) and Mn(IV) reduction., *Microbiol. Rev.*, 55(2), 259–287, 1991.
- 889 Lovley, D. R. and Klug, M. J.: Sulfate Reducers Can Outcompete Methanogens at Concentrations  
890 Sulfate Reducers Can Outcompete Methanogens Sulfate Concentrationst at Freshwater, *Appl.*  
891 *Environ. Microbiol.*, 45(1), 187–192, 1983.
- 892 Lovley, D. R., Phillips, E. J. P., Lonergan, D. J. and Widman, P. K.: Fe ( III ) and S<sub>0</sub> reduction by  
893 *Pelobacter carbinolicus*, *Appl. Environ. Microbiol.*, 61(6), 2132–2138, 1995.
- 894 Marlow, J. J., Steele, J. a., Case, D. H., Connon, S. a., Levin, L. a. and Orphan, V. J.: Microbial  
895 abundance and diversity patterns associated with sediments and carbonates from the methane seep  
896 environments of Hydrate Ridge, OR, *Front. Mar. Sci.*, 1(October), 1–16, 2014.
- 897 McDonald, I. R., Bodrossy, L., Chen, Y. and Murrell, J. C.: Molecular ecology techniques for the  
898 study of aerobic methanotrophs., *Appl. Environ. Microbiol.*, 74(5), 1305–1315, 2008.
- 899 Meng, J., Xu, J., Qin, D., He, Y., Xiao, X. and Wang, F.: Genetic and functional properties of  
900 uncultivated MCG archaea assessed by metagenome and gene expression analyses., *ISME J.*, 8(3),  
901 650–659, 2014.

- 902 Milucka, J., Ferdelman, T. G., Polerecky, L., Franzke, D., Wegener, G., Schmid, M., Lieberwirth,  
903 I., Wagner, M., Widdel, F. and Kuypers, M. M. M.: Zero-valent sulphur is a key intermediate in  
904 marine methane oxidation., *Nature*, 491(7425), 541–546, 2012.
- 905 Miyashita, A., Mochimaru, H., Kazama, H., Ohashi, A., Yamaguchi, T., Nunoura, T., Horikoshi,  
906 K., Takai, K. and Imachi, H.: Development of 16S rRNA gene-targeted primers for detection of  
907 archaeal anaerobic methanotrophs (ANMEs), *FEMS Microbiol. Lett.*, 297(1), 31–37, 2009.
- 908 Nam, Y.-D., Sung, Y., Chang, H.-W., Roh, S. W., Kim, K.-H., Rhee, S.-K., Kim, J.-C., Kim, J.-Y.,  
909 Yoon, J.-H. and Bae, J.-W.: Characterization of the depth-related changes in the microbial  
910 communities in Lake Hovsgol sediment by 16S rRNA gene-based approaches., *J. Microbiol.*, 46(2),  
911 125–136, 2008.
- 912 Niemann, H., Lösekann, T., de Beer, D., Elvert, M., Nadalig, T., Knittel, K., Amann, R., Sauter, E.  
913 J., Schlüter, M. and Klages, M.: Novel microbial communities of the Haakon Mosby mud volcano  
914 and their role as a methane sink, *Nature*, 443(7113), 854–858, 2006.
- 915 Nishri, a., Imberger, J., Eckert, W., Ostrovsky, I. and Geifman, Y.: The physical regime and the  
916 respective biogeochemical processes in the lower water mass of Lake Kinneret, *Limnol. Oceanogr.*,  
917 45(4), 972–981, 2000.
- 918 Nusslein, B., Chin, K. J., Eckert, W. and Conrad, R.: Evidence for anaerobic syntrophic acetate  
919 oxidation during methane production in the profundal sediment of subtropical Lake Kinneret  
920 (Israel), *Environ. Microbiol.*, 3(7), 460–470, 2001.
- 921 Orphan, V. J., House, C. H., Hinrichs, K. U., McKeegan, K. D. and DeLong, E. F.: Methane-  
922 consuming archaea revealed by directly coupled isotopic and phylogenetic analysis., *Science*,  
923 293(5529), 484–487, 2001.
- 924 Orphan, V. J., House, C. H., Hinrichs, K. U., McKeegan, K. D. and DeLong, E. F.: Multiple  
925 archaeal groups mediate methane oxidation in anoxic cold seep sediments, *Proc. Natl. Acad. Sci.*,  
926 99(11), 7663–7668, 2002.
- 927 Pester, M., Schleper, C. and Wagner, M.: The Thaumarchaeota: an emerging view of their  
928 phylogeny and ecophysiology., *Curr. Opin. Microbiol.*, 14(3), 300–306, 2011.
- 929 Pjevac, P., Kamyshny, A., Dykstra, S. and Mußmann, M.: Microbial consumption of zero-valence  
930 sulfur in marine benthic habitats., *Environ. Microbiol.*, 16(11), 3416–3430, 2014.
- 931 Plugge, C. M., Zhang, W., Scholten, J. C. M. and Stams, A. J. M.: Metabolic flexibility of sulfate-  
932 reducing bacteria., *Front. Microbiol.*, 2(May), 1–8, 2011.
- 933 Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J. and Glöckner, F.  
934 O.: The SILVA ribosomal RNA gene database project: improved data processing and web-based  
935 tools., *Nucleic Acids Res.*, 41(Database issue), D590–D596, 2013.
- 936 Raghoebarsing, A. A., Pol, A., Van de Pas-Schoonen, K. T., Smolders, A. J. P., Ettwig, K. F.,  
937 Rijpstra, W. I. C., Schouten, S., Damsté, J. S. S., den Camp, H. J. M. O. and Jetten, M. S. M.: A  
938 microbial consortium couples anaerobic methane oxidation to denitrification, *Nature*, 440(7086),  
939 918–921, 2006.

- 940 Röske, K., Sachse, R., Scheerer, C. and Röske, I.: Microbial diversity and composition of the  
941 sediment in the drinking water reservoir Saidenbach (Saxonia, Germany)., *Syst. Appl. Microbiol.*,  
942 35(1), 35–44, 2012.
- 943 Schloss, P. D., Gevers, D. and Westcott, S. L.: Reducing the effects of PCR amplification and  
944 sequencing artifacts on 16S rRNA-based studies., *PLoS One*, 6(12), e27310, 2011.
- 945 Schloss, P. D., Westcott, S. L., Ryabin, T., Hall, J. R., Hartmann, M., Hollister, E. B., Lesniewski,  
946 R. a, Oakley, B. B., Parks, D. H., Robinson, C. J., Sahl, J. W., Stres, B., Thallinger, G. G., Van  
947 Horn, D. J. and Weber, C. F.: Introducing mothur: open-source, platform-independent, community-  
948 supported software for describing and comparing microbial communities., *Appl. Environ.*  
949 *Microbiol.*, 75(23), 7537–7541, 2009.
- 950 Schoell, M.: Multiple origins of methane in the earth, *Chem. Geol.*, 71(1-3), 1–10, 1988.
- 951 Schubert, C. J., Vazquez, F., Lösekann-Behrens, T., Knittel, K., Tonolla, M. and Boetius, A.:  
952 Evidence for anaerobic oxidation of methane in sediments of a freshwater system (Lago di  
953 Cadagno)., *FEMS Microbiol. Ecol.*, 76(1), 26–38, 2011.
- 954 Schwarz, J. I. K., Eckert, W. and Conrad, R.: Community structure of Archaea and Bacteria in a  
955 profundal lake sediment Lake Kinneret (Israel), *Syst. Appl. Microbiol.*, 30(3), 239–254, 2007a.
- 956 Schwarz, J. I. K., Lueders, T., Eckert, W. and Conrad, R.: Identification of acetate-utilizing Bacteria  
957 and Archaea in methanogenic profundal sediments of Lake Kinneret (Israel) by stable isotope  
958 probing of rRNA., *Environ. Microbiol.*, 9(1), 223–237, 2007b.
- 959 Serruya, C.: Lake Kinneret: the nutrient chemistry of the sediments, *Limnol. Oceanogr.*, 16(May),  
960 510–521, 1971.
- 961 Serruya, C., Edelstein, M., Pollinger, U. and Serruya, S.: Lake Kinneret sediments: nutrient  
962 composition of the pore water and mud water exchanges, *Limnol. Oceanogr.*, 19(3), 489–508, 1974.
- 963 Severmann, S., Johnson, C. M., Beard, B. L. and McManus, J.: The effect of early diagenesis on the  
964 Fe isotope compositions of porewaters and authigenic minerals in continental margin sediments,  
965 *Geochim. Cosmochim. Acta*, 70(8), 2006–2022, 2006.
- 966 Sivan, O., Adler, M., Pearson, A., Gelman, F., Bar-Or, I., John, S. G. and Eckert, W.: Geochemical  
967 evidence for iron-mediated anaerobic oxidation of methane, *Limnol. Oceanogr.*, 56(4), 1536–1544,  
968 2011.
- 969 Smith, P. F., Langworthy, T. A. and Smith, M. R.: Polypeptide nature of growth requirement in  
970 yeast extract for *Thermoplasma* Polypeptide Nature of Growth Requirement in Yeast Extract for  
971 *Thermoplasma acidophilum*, *J. Bacteriol.*, 124(2), 884–892, 1975.
- 972 Sonne-Hansen, J. and Ahring, B. K.: *Thermodesulfobacterium hveragerdense* sp. nov., and  
973 *Thermodesulfobacterium islandicus* sp. nov., two thermophilic sulfate reducing bacteria isolated from a  
974 Icelandic hot spring., *Syst. Appl. Microbiol.*, 22(4), 559–564, 1999.
- 975 Spring, S., Amann, R., Ludwig, W., Schleifer, K. H., van Gemerden, H. and Petersen, N.:  
976 Dominating role of an unusual magnetotactic bacterium in the microaerobic zone of a freshwater  
977 sediment., *Appl. Environ. Microbiol.*, 59(8), 2397–2403, 1993.

- 978 Stookey, L. L.: Ferrozine-a new spectrophotometric reagent for iron, *Anal. Chem.*, 42(7), 779–781,  
979 1970.
- 980 Tavormina, P. L., Orphan, V. J., Kalyuzhnaya, M. G., Jetten, M. S. M. and Klotz, M. G.: A novel  
981 family of functional operons encoding methane/ammonia monooxygenase-related proteins in  
982 gammaproteobacterial methanotrophs, *Environ. Microbiol. Rep.*, 3(1), 91–100, 2011.
- 983 Tavormina, P. L., Ussler, W. and Orphan, V. J.: Planktonic and sediment-associated aerobic  
984 methanotrophs in two seep systems along the North American margin., *Appl. Environ. Microbiol.*,  
985 74(13), 3985–2995, 2008.
- 986 Thauer, R. K.: Functionalization of methane in anaerobic microorganisms., *Angew. Chem. Int. Ed.*  
987 *Engl.*, 49(38), 6712–6713, 2010.
- 988 Valentine, D. L.: Biogeochemistry and microbial ecology of methane oxidation in anoxic  
989 environments: a review, *Antonie Van Leeuwenhoek*, 81(1-4), 271–282, 2002.
- 990 Whitticar, M. J.: Carbon and hydrogen isotope systematics of bacterial formation and oxidation of  
991 methane, *Chem. Geol.*, 161(1-3), 291–314, 1999.
- 992 Wobus, A., Bleul, C., Maassen, S., Scheerer, C., Schuppler, M., Jacobs, E. and Röske, I.: Microbial  
993 diversity and functional characterization of sediments from reservoirs of different trophic state.,  
994 *FEMS Microbiol. Ecol.*, 46(3), 331–347, 2003.
- 995 Yamada, C., Kato, S., Kimura, S., Ishii, M. and Igarashi, Y.: Reduction of Fe(III) oxides by  
996 phylogenetically and physiologically diverse thermophilic methanogens., *FEMS Microbiol. Ecol.*,  
997 89(3), 637–645, 2014.
- 998 Yan, Z., Song, N., Cai, H., Tay, J.-H. and Jiang, H.: Enhanced degradation of phenanthrene and  
999 pyrene in freshwater sediments by combined employment of sediment microbial fuel cell and  
1000 amorphous ferric hydroxide., *J. Hazard. Mater.*, 199-200, 217–225, 2012.

1001 Ye, W., Liu, X., Lin, S., Tan, J., Pan, J., Li, D. and Yang, H.: The vertical distribution of bacterial  
1002 and archaeal communities in the water and sediment of Lake Taihu., *FEMS Microbiol. Ecol.*, 70(2),  
1003 107–120, 2009.

1004 Zhu, B., van Dijk, G., Fritz, C., Smolders, A. J. P., Pol, A., Jetten, M. S. M. and Ettwig, K. F.:  
1005 Anaerobic oxidization of methane in a minerotrophic peatland: enrichment of nitrite-dependent  
1006 methane-oxidizing bacteria., *Appl. Environ. Microbiol.*, 78(24), 8657–8665, 2012.

1007

## 1008 **Tables**

1009 **Table 1.** Bacterial sequences used for classification in SILVA ngs and for subsample in MOTUR for the  
1010 alpha diversity.

Bacteria	# seq SILVA ngs	OTUs SILVA ngs	# seq MOTUR	OTUs MOTUR	coverage	invsimpson	chao	ace	shannon
0-3 cm	3631	599	2588	388	0.92	10/10.8/11.7	805/925.7/1092.8	1246.5/1384.0/1546.7	3.83/3.91/3.98
6-9 cm	4641	1735	3365	1337	0.77	286.6/315.9/351.8	2437.9/2673.0/2958.4	3876.8/4135.2/4420.0	6.48/6.53/6.57
29-32 cm	5038	1516	3615	1214	0.81	127.8/142.4/160.6	2104.4/2318.7/2583.1	3047.7/3259.5/3495.4	6.13/6.19/6.24

1011

1012 **Table 2.** Archaeal sequences used for classification in SILVA ngs and for subsample in MOTHR for the  
 1013 alpha diversity.

Archaea	# seq SILVA ngs	OTUs SILVA ngs	# seq MOTHR	OTUs MOTHR	coverage	invsimpson	chao	ace	shannon
0-3 cm	288	48	268	32	0.90	2.5/2.9/3.5	53.9/83.4/164.5	232.8/344.6/520.4	1.6/1.8/2.0
6-9 cm	1382	184	1408	200	0.86	8.4/10.2/13.2	39.0/131.4/220.4	144.7/195.2/277.4	2.87/3.05/3.22
29-32 cm	3110	383	2829	245	0.88	7.9/9.2/10.9	73.5/104.8/181.7	133.2/190.9/289.6	2.58/2.74/2.91

1014

1015 **Figure Captions:**

1016 **Figure 1:** Schematic location of Lake Kinneret. Numbers correspond to altitude measured in meters  
 1017 from the level of Lake Kinneret. Cores were taken from station A (taken from Hambright et al.,  
 1018 2004).

1019 **Figure 2:** Geochemical porewater profiles in LK sediment. **A.** profile of  $\text{SO}_4^{-2}$  (black circles) and  
 1020 sulfide (gray diamonds) in the porewater **B.** Headspace measurements of methane (gray circles) and  
 1021  $\delta^{13}\text{C}_{\text{CH}_4}$  (black squares) in the sediments **C.**  $\delta^{13}\text{C}$  of total lipids extraction (TLE) (black diamonds)  
 1022 from the sediment, and  $\delta^{56}\text{Fe}$  (gray circles) of the dissolved iron in the porewater. **D.** Electron  
 1023 acceptor profiles of dissolved Fe(II) (gray triangles) and Mn (II) (black squares) in the porewater.,  
 1024 Black arrows indicate the sampled sections for 16S rRNA gene analysis. The presented methane  
 1025 profile was taken two weeks prior to the sampling for the microbial communities on December  
 1026 2009. Sampling for Fe(II), Mn(II) and  $\text{SO}_4^{-2}$  profiles was carried out between 2007 and 2011 and  
 1027 for sulfide from 2007 to 2013. The  $\delta^{13}\text{C}_{\text{CH}_4}$  and  $\delta^{56}\text{Fe}$  profiles were performed four months prior to  
 1028 the microbial sampling on August 2009.


1029 **Figure 3:** Classification of bacterial sequences using SILVA ngs pipeline. Phyla and classes  
 1030 distribution of sequences of the 454 sequencing at the different depths above 1% in the  
 1031 communities.

1032 **Figure 4:** Classification of archaeal sequences using SILVA ngs pipeline. Phyla and orders  
 1033 distributions of sequences of the 454 sequencing at the different depths above 1% in the  
 1034 communities.

1035 **Figure 5:** Principal coordinate analysis (PCoA) of the **A.** bacterial and **B.** archaeal community  
 1036 matrix (based on operational taxonomic units) and vector fitting of the environmental geochemical  
 1037 conditions in the different depth. The bacterial community from the upper layer is labeled with red  
 1038 triangle, the middle layer is labeled with green triangle and bottom layer is labeled with blue  
 1039 triangle.

1040 **Figure 6:** profiles of functional genes from the 3 different depths. Green bar represent *pmoA*  
1041 functional gene of aerobic methanotroph. Red bar represent *dsrAB* functional gene of sulfate  
1042 reducers and blue bar represent *mcrA* functional gene from methanogens and anaerobic  
1043 methanotrophs.

1044



1045

1046

1047


1048

1049

1050

**Figure 1.**






1051  
1052

1053

1054

Figure 2.



1055


1056

1057

1058

Figure 3.

1059




1060


1061

1062

Figure 4.



1063




1064

1065

1066

1067

Figure 5.



1068

1069

1070

1071


1072

**Figure 6**

1073

1074

1075



1077


1078

1079

**Figure 1**

1080 **Supplementary Fig. 1.** Venn diagram showing archaeal sequences overlap between the different  
1081 depths. The green circle represents the top sample (0-3 cm), the blue circle represents the middle  
1082 sample (6-9 cm) and the red circle represents the bottom sample (29-32 cm).

1083



1084

1085

**Figure 2**

1086 **Supplementary Fig. 2.** Venn diagram showing bacterial sequences overlap between the different  
1087 depths. The green circle represents the top sample (0-3 cm), the blue circle represents the middle  
1088 sample (6-9 cm) and the red circle represents the bottom sample (29-32 cm).

1089

1090 **Supplementary Table 1.** The classification percentage and number of sequences of archaea A.  
 1091 phylum, B. class and C. order by SILVA ngs.

1092

Archaea	0-3 cm		6-9 cm		29-32 cm	
<b>Phylum</b>	# seq	%	# seq	%	# seq	%
Euryarchaeota	283	98.3	1430	96.4	3048	98.0
Thaumarchaeota	5	1.7	53	3.6	62	2.0

1093

Archaea	0-3 cm		6-9 cm		29-32 cm	
<b>Class</b>	# seq	%	# seq	%	# seq	%
AK59					1	0.0
AK8			1	0.1	4	0.1
Group C3			2	0.1	2	0.1
Halobacteria	7	2.4	96	6.5	165	5.3
Marine Benthic Group B	2	0.7	24	1.6	30	1.0
Marine Group I					2	0.1
Methanobacteria	1	0.3				
Methanomicrobia	266	92.4	1164	78.5	2679	86.1
Miscellaneous Crenarchaeotic Group	3	1.0	21	1.4	19	0.6
pSL12			2	0.1	3	0.1
South African Gold Mine Gp 1(SAGMCG-1)			3	0.2	1	0.0
Thermoplasmata	9	3.1	170	11.5	204	6.6

1094

Archaea	0-3 cm		6-9 cm		29-32 cm	
<b>Order</b>	# seq	%	# seq	%	# seq	%
20a-9			5	0.3	3	0.1
ANME-1					1	0.0
Cenarchaeales					2	0.1
Halobacteriales	7	2.4	96	6.5	165	5.3
Methanobacteriales	1	0.3				
Methanomicrobiales	202	70.1	422	28.5	878	28.2
Methanosarcinales	64	22.2	742	50.0	1800	57.9
Thermoplasmatales	8	2.8	165	11.1	201	6.5
WCHA1-57	1	0.3				
Uncultured	5	1.7	53	3.6	60	1.9

1095 **Supplementary Table 2.** The classification percentage and number of sequences of bacterial A.  
 1096 phylum, B. class and C. order by SILVA ngs.

Bacteria	0-3 cm		6-9 cm		29-32 cm	
Phylum	# seq	%	# seq	%	# seq	%
Acidobacteria	33	0.9	112	2.4	123	2.4
Actinobacteria	56	1.5	61	1.3	90	1.8
Armatimonadetes	21	0.6	12	0.3	14	0.3
Bacteroidetes	80	2.2	349	7.5	126	2.5
BD1-5	2	0.1	2	0.0	2	0.0
Caldiserica					2	0.0
Candidate division BRC1	2	0.1	14	0.3	8	0.2
Candidate division JS1	1	0.0	2	0.0		
Candidate division KB1					3	0.1
Candidate division OD1			1	0.0	18	0.4
Candidate division OP11	2	0.1	17	0.4	1	0.0
Candidate division OP3	1	0.0	108	2.3	84	1.7
Candidate division OP8	9	0.2	81	1.7	106	2.1
Candidate division SR1			4	0.1		
Candidate division TM7	6	0.2	5	0.1		
Candidate division WS3	15	0.4	70	1.5	103	2.0
Chlorobi	38	1.0	235	5.1	206	4.1
Chloroflexi	65	1.8	672	14.5	969	19.2
CKC4	1	0.0				
Cyanobacteria	18	0.5	69	1.5	21	0.4
Deferribacteres	1	0.0	122	2.6	126	2.5
Deinococcus-Thermus	1	0.0	8	0.2	9	0.2
Elusimicrobia			7	0.2	13	0.3
Fibrobacteres			20	0.4	9	0.2
Firmicutes	1376	37.9	140	3.0	152	3.0
Gemmatimonadetes	2	0.1	15	0.3	15	0.3
Hyd24-12	4	0.1	8	0.2		
GOUTA4					4	0.1
JL-ETNP-Z39			1	0.0	3	0.1
Lentisphaerae	12	0.3	38	0.8	10	0.2
Nitrospirae	31	0.9	195	4.2	409	8.1
NPL-UPA2	1	0.0	33	0.7	5	0.1
OC31			1	0.0		
Planctomycetes	22	0.6	140	3.0	172	3.4
Proteobacteria	1762	48.5	1837	39.6	1954	38.8
SHA-109	6	0.2	6	0.1	1	0.0
Spirochaetae	1	0.0	102	2.2	134	2.7
Synergistetes					1	0.0
TA06	5	0.1	66	1.4	102	2.0
Tenericutes	13	0.4	1	0.0	1	0.0
Thermotogae			8	0.2	1	0.0
TM6	19	0.5	27	0.6	21	0.4
Verrucomicrobia	3	0.1	39	0.8	20	0.4
WCHB1-60	22	0.6	13	0.3		



Bacteria	0-3 cm		6-9 cm		29-32 cm	
	# seq	%	# seq	%	# seq	%
Acidimicrobiia	5	0.1	4	0.1	2	0.0
Acidobacteria	26	0.7	80	1.7	83	1.6
Actinobacteria	7	0.2	5	0.1	5	0.1
Alphaproteobacteria	23	0.6	25	0.5	50	1.0
Anaerolineae	31	0.9	296	6.4	375	7.4
Ardenticatenia			1	0.0		
ARKDMS-49			2	0.0		
ARKICE-90			11	0.2		
Bacilli	611	16.8	23	0.5	49	1.0
Bacteroidia			5	0.1	1	0.0
BD2-2	2	0.1	16	0.3	8	0.2
Betaproteobacteria	65	1.8	268	5.8	344	6.8
BS5	3	0.1	7	0.2		
BSV13			9	0.2	1	0.0
Caldilineae	1	0.0	8	0.2	3	0.1
Caldisericia					2	0.0
Chlorobia	6	0.2	42	0.9	18	0.4
Chloroplast	3	0.1	42	0.9	4	0.1
Clostridia	761	21.0	116	2.5	99	2.0
Coriobacteriia	3	0.1	4	0.1	7	0.1
Cyanobacteria	13	0.4	22	0.5	7	0.1
Cytophagia	1	0.0	4	0.1	8	0.2
Deferribacteres	1	0.0	122	2.6	126	2.5
Dehalococcoidia	2	0.1	276	5.9	453	9.0
Deinococci	1	0.0	8	0.2	9	0.2
Deltaproteobacteria	57	1.6	1257	27.1	1022	20.3
DEV055	6	0.2	9	0.2		
Elusimicrobia			7	0.2	13	0.3
Epsilonproteobacteria			12	0.3	1	0.0
Erysipelotrichia					3	0.1
FFCH16263	1	0.0			2	0.0
Fibrobacteria			20	0.4	9	0.2
Flavobacteriia	9	0.2	34	0.7	3	0.1
Gammaproteobacteria	1614	44.5	257	5.5	536	10.6
Gemmatimonadetes	2	0.1	15	0.3	15	0.3
Holophagae	5	0.1	24	0.5	29	0.6
Ignavibacteria	32	0.9	193	4.2	188	3.7
JG30-KF-CM66			12	0.3	6	0.1
KD4-96	9	0.2	5	0.1	4	0.1
Ktedonobacteria	1	0.0				
LD1-PB3			1	0.0	1	0.0
Lentisphaeria	1	0.0	3	0.1		
MB-A2-108	6	0.2	4	0.1	14	0.3
MBMPE71			1	0.0	1	0.0
Melainabacteria	2	0.1	1	0.0	4	0.1
ML635J-21			4	0.1	6	0.1
Mollicutes	13	0.4	1	0.0	1	0.0
MSB-5B2					4	0.1
MSBL3			6	0.1		
Negativicutes	4	0.1	1	0.0	1	0.0
Nitrospira	31	0.9	195	4.2	409	8.1
OM190	1	0.0	4	0.1	2	0.0
Oligosphaeria			7	0.2	5	0.1
OPB35 soil group	3	0.1	25	0.5	11	0.2
OPB41	20	0.6	21	0.5	35	0.7
Opitutae			12	0.3	7	0.1
PBS-III-20	2	0.1	4	0.1		
Phycisphaerae	11	0.3	101	2.2	130	2.6
Pla3 lineage			10	0.2	7	0.1
Pla4 lineage			9	0.2	8	0.2
Planctomycetacia	10	0.3	15	0.3	19	0.4
Proteobacteria Incertae Sedis			1	0.0	1	0.0
S085					1	0.0
SB-1			7	0.2	10	0.2
SB-5	19	0.5	33	0.7	12	0.2
S-BQ2-57 soil group					1	0.0
SHA-26			7	0.2	14	0.3
Spartobacteria			1	0.0	1	0.0
Sphingobacteriia	30	0.8	95	2.0	30	0.6
Spirochaetes	1	0.0	102	2.2	134	2.7
Subgroup 22	2	0.1	8	0.2	11	0.2
Synergistia					1	0.0
TA18	3	0.1	4	0.1		
Thermoleophilia	14	0.4	23	0.5	25	0.5
Thermotogae			8	0.2	1	0.0
TK10					2	0.0
vadinHA17	18	0.5	145	3.1	51	1.0
vadinHA49					5	0.1
VC2.1 Bac22			1	0.0		
Verrucomicrobiae			1	0.0		
WCHB1-41			1	0.0	2	0.0
WCHB1-32	1	0.0			1	0.0
uncultured	138	3.8	538	11.6	585	11.6

Bacteria	0-3 cm		6-9 cm		29-32 cm	
Order	# seq	%	# seq	%	# seq	%
10bav-F6					2	0.0
43F-1404R			2	0.0	3	0.1
Acidimicrobiales	5	0.1	4	0.1	2	0.0
Alteromonadales			9	0.2	14	0.3
Amsterdam-1B-07			1	0.0	4	0.1
Anaerolineales	31	0.9	296	6.4	375	7.4
B1-7BS			1	0.0		
B276-D12			1	0.0		
Bacillales	602	16.6	23	0.5	47	0.9
Bacteroidales			5	0.1	1	0.0
BD2-11 terrestrial group			2	0.0	5	0.1
Bdellovibrionales			34	0.7	13	0.3
BP-U1C-1g10			1	0.0		
Burkholderiales	56	1.5	86	1.9	208	4.1
C86					2	0.0
Caldilineales	1	0.0	8	0.2	3	0.1
Caldisericales					2	0.0
Campylobacterales			12	0.3	1	0.0
Caulobacterales	12	0.3	1	0.0	1	0.0
CCM11a			9	0.2	6	0.1
Chlorobiales	6	0.2	42	0.9	18	0.4
Chromatiales	3	0.1	11	0.2	21	0.4
Chthoniobacterales			1	0.0	1	0.0
Clostridiales	760	20.9	107	2.3	99	2.0
Coriobacteriales	3	0.1	4	0.1	7	0.1
Corynebacteriales	1	0.0	1	0.0	3	0.1
Cytophagales	1	0.0	3	0.1		
D8A-2	1	0.0				
Deferribacterales	1	0.0	122	2.6	126	2.5
Dehalococcoidales			6	0.1	2	0.0
Deltaproteobacteria Incertae Sedis	2	0.1	55	1.2	2	0.0
Desulfarculales	12	0.3	132	2.8	62	1.2
Desulfobacterales	13	0.4	118	2.5	88	1.7
Desulfuromonadales			80	1.7	41	0.8
Erysipelotrichales					3	0.1
Fibrobacterales			18	0.4	6	0.1
Flavobacteriales	9	0.2	34	0.7	3	0.1
Frankiales	1	0.0	2	0.0	3	0.1
FS117-23B-02			2	0.0	4	0.1
FW22	1	0.0	2	0.0	2	0.0
Gaiellales	11	0.3	19	0.4	20	0.4
Gammaproteobacteria Incertae Sedis			4	0.1	4	0.1
Gastranaerophilales			1	0.0	4	0.1
Gemmatimonadales	2	0.1	13	0.3	10	0.2
GIF3			2	0.0	7	0.1
GIF9			117	2.5	251	5.0

GR-WP33-30	1	0.0	14	0.3	34	0.7
Haloplasmales	13	0.4				
HOC36	2	0.1	72	1.6	267	5.3
Holophagae Incertae Sedis			5	0.1	12	0.2
Holophagales			2	0.0	2	0.0
Hydrogenophilales			48	1.0	44	0.9
Ignavibacteriales	32	0.9	193	4.2	188	3.7
KCLunmb-38-53	1	0.0	9	0.2		
KD3-62	1	0.0	8	0.2	9	0.2
KI89A clade			4	0.1	5	0.1
Lactobacillales	2	0.1			2	0.0
Legionellales	5	0.1	17	0.4	15	0.3
Lineage IIb					4	0.1
Lineage IIc			5	0.1	4	0.1
Lineage IV			1	0.0	1	0.0
MD2894-B20			1	0.0	3	0.1
Methylococcales	9	0.2	70	1.5	69	1.4
Methylophilales			27	0.6	10	0.2
Micrococcales	1	0.0	2	0.0	1	0.0
mle1-8	6	0.2	5	0.1	8	0.2
MSB-3A7 sediment group			3	0.1	3	0.1
MSBL8			1	0.0		
MSBL5	1	0.0	139	3.0	169	3.4
MSBL9	1	0.0	22	0.5	59	1.2
MVP-21	1	0.0				
MVP-88					1	0.0
Myxococcales	1	0.0	82	1.8	45	0.9
NB1-n			1	0.0	1	0.0
Nitrosomonadales	2	0.1	36	0.8	22	0.4
Nitrospirales	31	0.9	195	4.2	409	8.1
ODP1230B30.09					1	0.0
Oceanospirillales			1	0.0		
Oligosphaerales			7	0.2	5	0.1
Opitutales			12	0.3	7	0.1
Order II (Bacteroidetes)			1	0.0	5	0.1
Order Incertae Sedis			2	0.0		
PBS-18			3	0.1	1	0.0
PeM15	1	0.0				
Phycisphaerales	2	0.1	10	0.2	10	0.2
Pla1 lineage			18	0.4	17	0.3
Planctomycetales	10	0.3	15	0.3	19	0.4
Propionibacteriales	3	0.1				
possible order 07			2	0.0	3	0.1
Pseudomonadales	1523	41.9	1	0.0		
Rhizobiales	6	0.2	16	0.3	45	0.9
Rhodobacterales	1	0.0	1	0.0		
Rhodocyclales	4	0.1	43	0.9	29	0.6
Rhodospirillales			3	0.1	1	0.0
Rickettsiales	3	0.1	4	0.1		

S15A-MN16	1	0.0	18	0.4	14	0.3
S-70			3	0.1		
SAR324 clade(Marine group B)			2	0.0		
SC-I-84	2	0.1	27	0.6	31	0.6
Sh765B-AG-111					2	0.0
Sh765B-TzT-29			1	0.0	3	0.1
SHA-43			2	0.0	6	0.1
Selenomonadales	4	0.1	1	0.0	1	0.0
Solirubrobacterales	3	0.1	4	0.1	5	0.1
Sphingobacteriales	30	0.8	95	2.0	30	0.6
Sphingomonadales	1	0.0			3	0.1
Spirochaetales	1	0.0	98	2.1	133	2.6
Streptomycesetales					1	0.0
Subgroup 17	1	0.0	3	0.1		
Subgroup 18	3	0.1	27	0.6	21	0.4
Subgroup 23	2	0.1	6	0.1	7	0.1
Subgroup 4	2	0.1				
Subgroup 6	20	0.6	31	0.7	25	0.5
Subgroup 7	2	0.1	3	0.1		
Subgroup 13			5	0.1	5	0.1
Subgroup 19			1	0.0	13	0.3
Subgroup 2			1	0.0	6	0.1
Subgroup 21			9	0.2		
Subgroup 25			1	0.0	6	0.1
Subgroup 3			1	0.0	1	0.0
Subgroup 11					1	0.0
Subgroup 9					1	0.0
SubsectionII (Cyanobacteria)					1	0.0
SubsectionI (Cyanobacteria)	6	0.2	8	0.2	3	0.1
SubsectionIV (Cyanobacteria)	7	0.2	14	0.3	3	0.1
Sva0485	20	0.6	390	8.4	546	10.8
Synergistales					1	0.0
Syntrophobacterales	8	0.2	347	7.5	183	3.6
SZB30	4	0.1	9	0.2	7	0.1
Thermoanaerobacterales			9	0.2		
Thermotogales			8	0.2	1	0.0
Thiotrichales	3	0.1	5	0.1	3	0.1
TPD-58	1	0.0	6	0.1	8	0.2
TRA3-20	1	0.0				
vadinBA26			8	0.2	16	0.3
Verrucomicrobiales			1	0.0		
Vampirovibrionales	2	0.1				
VAN12	7	0.2				
Victivallales	1	0.0	3	0.1		
WD2101 soil group			1	0.0		
Xanthomonadales	65	1.8	50	1.1	129	2.6
uncultured	237	6.5	951	20.5	816	16.2

## 1100 **Appendix**

1101 The major bacterial populations in the sediment:

### 1102 *Proteobacteria*

1103 *Proteobacteria* was the most dominant bacterial phylum in all the depths. Their percentage was  
1104 slowly decreases with depth however the composition within its classes were changing with depth.  
1105 *Proteobacteria* is also among the most abundant phyla in LK (Schwarz et al., 2007a) and in other  
1106 freshwater sediments (Tamaki et al., 2005; Wobus et al., 2003). *Proteobacteria* is a very diverse  
1107 phylum, divided in to number of classes which most of them have been observed in LK sediments.

### 1108 *Betaproteobacteria*

1109 *Betaproteobacteria* percentages were slowly increasing with depth. They are comprised of  
1110 chemoheterotrophs and chemoautotrophs which derive nutrients from decomposition of organic  
1111 material. *Burkholderiales* was the dominant order in *Betaproteobacteria* and increased with depth  
1112 (1.5% to 4%). Schwarz et al. (2007a) showed that *Burkholderiales* is active in the upper sediment of  
1113 LK. Members of this group are able of anaerobic oxidation of acetate with (per)chlorate as electron  
1114 acceptor (Yoshida et al., 2005) or oxidize hydrogen in deep subsurface (Orcutt et al., 2011).  
1115 *Hydrogenophilales* and *Nitrosomonadales* orders were observed in lower abundant but in the  
1116 middle and deep layer of the sediment. *Hydrogenophilales* can utilize hydrogen (Ontiveros-  
1117 Valencia et al., 2013) and *Hydrogenophilaceae* family are obligatory chemolithoautotrophic,  
1118 aerobic or facultative anaerobic sulfur oxidizers and reduction of nitrate (Kelly and Wood, 2000)  
1119 which are often found also in wastewater treatment systems (Luo et al., 2011). *Nitrosomonadales*  
1120 are freshwater bacteria which presumed to have a role in ammonia oxidation and/or denitrification  
1121 processes (Edlund et al., 2008). McBeth et al. (2013) also showed that *Nitrosomonadales* are  
1122 commonly found as freshwater Fe oxidizers. It should be noted that the upper sediment may  
1123 contain denitrifying bacteria that arrived from the water column where nitrification conditions exists  
1124 throughout the year.

### 1125 *Gammaproteobacteria*

1126 Our sequences which were affiliated to the *Gammaproteobacteria* have different phylogenetic  
1127 classification distribution in each layer. The upper layer was dominated by *Pseudomonadales* (42%)  
1128 which were divided to 2 main genera: *Acinetobacter* (28%) and *Pseudomonas* (14%). *Acinetobacter*  
1129 was shown to have the ability to degrade organic matter. Our sequences which similar to  
1130 *Acinetobacter* genus were closest to *Acinetobacter lwoffii* (99%) found in sewage and groundwater  
1131 (Nazina et al., 2000). our sequences classified as *Pseudomonas* were similar to *Pseudomonas*  
1132 *stutzeri* (99%) which has the ability to oxidize a large variety of organic compounds, and some

1133 members also involved in nitrification and denitrification, degradation of aromatic compounds, and  
1134 nitrogen fixation (Chen et al., 2011). In the middle and bottom layers *Methylococcales* order was  
1135 observed (~1%). *Methylococcales* are aerobic methanotrophs which some members were present  
1136 within the sediments and overlying water column from dysoxic, methane-rich vent and seep  
1137 systems (Tavormina et al., 2008). The most abundant order in the deep layer was HOC36 (5%)  
1138 which is an uncultured *Gammaproteobacteria*. however when compared to NCBI data base it was  
1139 closely related to uncultured LK clones (99%) and to cultured *Methylocaldum* sp. (94%) (Bodrossy  
1140 et al., 1997) which was a Thermophilic methanotroph isolated from landfill cover soil.

#### 1141 *Deltaproteobacteria*

1142 ~~*Deltaproteobacteria* were the most abundant in the middle and deep sediment samples. In the~~  
1143 ~~middle layer the abundant classes were *Syntrophobacterales* (8%), *Desulfarculales* (3%),~~  
1144 ~~*Desulfobacterales* (2%) and *Desulfuromonadales* (2%). Many members of the families of~~  
1145 ~~*Syntrophobacterales* are sulfate reducers or as fermentative organism, depending on the~~  
1146 ~~environmental conditions (McInerney et al., 2007). *Syntrophaceae* the dominant family of~~  
1147 ~~*Syntrophobacterales* are acetate degrading sulfate reducing bacterium (Jiang et al., 2009).~~  
1148 ~~*Desulfarculales* are strictly anaerobic are mesophilic sulfate reducing bacterium with the capability~~  
1149 ~~to oxidize acetate and fatty acids (Sun et al., 2010). *Desulfobacterales* are sulfate reducers capable~~  
1150 ~~to oxidize hydrogen in soils and sediments (Burow et al., 2014). *Desulfuromonadales* was shown as~~  
1151 ~~S<sup>0</sup> respiring (Pjevac et al., 2014) and may also reduce Fe(III) and Mn(IV) in marine surface~~  
1152 ~~sediments (Lovley, 2006). About 1% of the sequences in the middle and bottom layers were~~  
1153 ~~affiliated to *Geobacter* genus which involve in iron reduction. In the deep layer the most abundant~~  
1154 ~~class was Sva0485. Sequences of the Sva485 order were similar to those found in different aquatic~~  
1155 ~~environments, but the metabolic functions of members of this order are not clear. *Pelobacter*~~  
1156 ~~*earbinolious* (Lovley et al., 1995), a member of Sva485 capable of Fe(III) and sulfur reduction.~~  
1157 ~~Schwarz et al. (2007a) showed in the upper part of LK sediment that *Deltaproteobacteria* was one~~  
1158 ~~of the dominant classes, most of which were affiliated with acetate oxidizing sulfate reducing~~  
1159 ~~bacteria, which outcompete the acetoclastic methanogens. Even though *Deltaproteobacteria* are~~  
1160 ~~best known for their sulfate reduction metabolism, they can shift their metabolism in response to~~  
1161 ~~depleted sulfate concentrations (Plugge et al., 2011).~~

#### 1162 *Chloroflexi*

1163 ~~*Chloroflexi* was found to be one of the most abundant bacterial phyla in LK and represented diverse~~  
1164 ~~functional groups that increased with depth. The abundant classes that increase with depth were~~  
1165 ~~*Anaerolineae* and *Dehalococcoidia*. Although they are found in a variety of anaerobic~~

1166 environments, only a few representatives have been cultivated so far (Yamada and Sekiguchi,  
1167 2009). The metabolisms of the cultured representatives showed that they utilized organic matter, but  
1168 no specific electron acceptor has as yet been reported (Yamada et al., 2006). *Chloroflexi* was one of  
1169 the most dominant phyla in contaminated soil environment which had a lot of polycyclic aromatic  
1170 hydrocarbons (Winderl et al., 2008). Members of *Chloroflexi* were found throughout the entire core,  
1171 indicating that *Anaerolineae* representatives could be the main aromatic organic matter  
1172 decomposers in LK. *Dehalococcoidia* are widely distributed in the marine subsurface, yet metabolic  
1173 properties of the many uncultivated lineages are completely unknown. *Dehalococcoidia* could be  
1174 capable of oxidizing various fatty acids and/or structurally related substrates (Wasmund et al.,  
1175 2013) or reductive dechlorination of chlorinated hydrocarbon (Maymo et al., 1995).

#### 1176 *Nitrospirae*

1177 *Nitrospirae* phylum, were increased with depth. Sequences from our samples were classified to  
1178 *Nitrospirales* order in *Nitrospirea* which were most abundant in the bottom layer of the sediment  
1179 (7%). Part of our sequences were similar to previous study (Schwarz et al., 2007b). (Schwarz et al.,  
1180 2007b) showed by SIP RNA not only that *Nitrospirae* was present, but also that it was functionally  
1181 bioactive. Although *Nitrospirae* is a known nitrate oxidizer (Ehrich et al., 1995), the conditions of  
1182 this environment suggest that it utilized another metabolic pathway. *Nitrospirae* also include the  
1183 iron reducing candidates *Magnetobacterium bavaricum* (Spring et al., 1993) and sulfur reducers  
1184 (Sonne-Hansen and Ahring, 1999). Part of our sequences were aligned to uncultured clone (98%)  
1185 from freshwater sediment which enhanced degradation of phenanthrene and pyrene by amorphous  
1186 ferric hydroxide (Yan et al., 2012).

#### 1187 *Chlorobi*

1188 *The Chlorobi*, green sulfur bacteria, are a group of anoxygenic photosynthetic bacteria observed  
1189 frequently in anoxic microbial mats and meromictic lakes. *Chlorobi* in found in deeper layers of the  
1190 sediment since they support higher H<sub>2</sub>S concentrations and require less light intensity. They  
1191 photosynthesize using H<sub>2</sub>S as electron donor, although they can also grow photoheterotrophically,  
1192 and some of them form symbiotic consortia with heterotrophic partners that are widely distributed  
1193 in chemoclines of meromictic lakes throughout the world (Bri e et al., 2007). *Chlorobi* metabolic  
1194 features include fixing CO<sub>2</sub>, fixing N<sub>2</sub> and oxidize sulfide or other reduced sulfur compounds (Liu  
1195 et al., 2012). Most our sequences were classified to *Ignavibacteria* order which capable of  
1196 organoheterotrophy under both oxic and anoxic conditions.

#### 1197 *Firmicutes*

1198 *Firmicutes* are mostly abundant in the upper part of the sediment. Our sequences were classified to  
1199 2 main orders: *Bacillales* and *Clostridia*. *Bacillales* have been observed in diverse environments  
1200 and with diverse metabolic functions. Our sequences were closely related to a few species: *Bacillus*  
1201 *thioparans* (99%) which grew chemolithoautotrophically by oxidation of thiosulfate to sulfate and  
1202 was found in deep sea sediment (Pérez-Ibarra et al., 2007). *Bacillus subterraneus* (99%) utilize  
1203 amorphous iron(III), Mn(II)(IV), nitrate, nitrite and fumarate as electron acceptors (Kanso et al.,  
1204 2002). *Clostridia* are anaerobic bacteria which play an important role in the biogeochemical cycling  
1205 of carbon, sulfur and iron. *Clostridia* are key microbial drivers in iron reduction in estuarine  
1206 sediments and acid mine drainage environments (Shah et al., 2014). Part of our sequences were  
1207 closely related to *Clostridium tunisiense* (96%) which uses elemental sulfur but not sulfate,  
1208 thiosulfate, nor sulfite as terminal electron acceptors (Thabet et al., 2004).

#### 1209 *Bacteroidetes*

1210 Sequences of *Bacteroidetes* were mostly abundant in the middle layer of the LK sediment core.  
1211 *Bacteroidetes* are known as hydrolytic fermentative bacteria, they are major utilizers of high-  
1212 molecular-mass dissolved organic matter in marine ecosystems (Cottrell and Kirchman, 2000).  
1213 Therefore, fresh organic matter (high in high molecular mass dissolved organic matter) that  
1214 descends from the water column could be utilized by the *Bacteroidetes* present in the upper part of  
1215 the sediment.

1216 The major archaeal populations in the sediment:

1217 The distribution of the population was described above. However the environmental conditions and  
1218 metabolic functionality of our sequences were not shown. The uncultured clones [similar](#) to  
1219 our sequences might help indicate on microbial process of the dominant sequences in our samples.

#### 1220 *Euryarchaeota*

#### 1221 *Methanomicrobia*

1222 The majority of our sequences were classified within *Euryarchaeota*, in *Methanomicrobia* class.  
1223 Our 16S rRNA sequences were divided between *Methanomicrobiales* and *Methanosarcinales*  
1224 order. *Methanosaeta* genus of *Methanosarcinales* increases with depth. *Methanosaeta* are  
1225 acetoclastic methanogens which only able to grow on acetate even at low concentrations (Jetten et  
1226 al., 1990). *Methanosaeta* was also shown in previous studies of LK (Schwarz et al., 2007a, 2007b)  
1227 and in other meso to eutrophic freshwater lakes (Glissman et al., 2004; Koizumi et al., 2003).  
1228 Yamada et al. (2014) showed that *Methanosaeta* can also have the ability to reduce ferrihydrite with  
1229 H<sub>2</sub> as the electron source. The closest relative cultured acetoclastic methanogen to our sequences  
1230 was *Methanosaeta concilii* (96%) similar results to the previous [studies](#). *Methanolinea* genus



1231 of *Methanomicrobiales* which is hydrogenotrophic methanogen was observed in constant  
1232 percentages for all depths. *Methanoregula* genus which is also hydrogenotrophic methanogen is  
1233 decreasing with depth. *Methanoregula*, *Methanosaeta* and *Methanolinea* represent 60% of  
1234 sequences of methanogens retrieved from freshwater lakes (Borrel et al., 2011).

#### 1235 *Thermoplasmata*

1236 The dominant family in *Thermoplasmata* is The Marine Benthic Group D and DHVEG-1 (MBG-  
1237 D). Members of the MBG-D have been shown to exist in a variety of freshwater and marine  
1238 environments (Beal et al., 2009; Borrel et al., 2012), and it is the most widely encountered,  
1239 uncultured lineage in freshwater lake sediments. Even though their metabolism is unknown,  
1240 hypotheses about their functionalities are based on the environments in which they were found.  
1241 Methanogenesis was suggested, as they were found in deep lake sediments with high methane  
1242 concentrations (Borrel et al., 2012), and they were also hypothesized to be involved in AOM, as  
1243 they were found in AOM zones (Schubert et al., 2011) and in marine seep sediment (Beal et al.,  
1244 2009). However, in other environments, in which methane concentrations were low, the utilization  
1245 of waste products, intermediates, or dead cells by MBG-D was also suggested (Smith et al., 1975).  
1246 Recently, a single cell genomics study showed that members of MBG-D were capable of exogenous  
1247 protein degradation in cold anoxic environments (Lloyd et al., 2013).

1248 In addition Terrestrial Miscellaneous Gp(TMEG) family was observed only in the middle and  
1249 bottom layers of the sediment. This lineage was includes clones from the terrestrial subsurface and  
1250 from soils, marine sediments and freshwater lakes (Teske and Sørensen, 2008). Sequences closely  
1251 related to this group were found in sediment influenced by sulfur-rich, hypoxic groundwater and  
1252 aquatic sediment samples.

#### 1253 *Thaumarchaeota*

1254 ~~*Thaumarchaeota* are mostly known as archaeal ammonia oxidizers but in the sediments they may~~  
1255 ~~contribute significantly to the reservoir of nitrogen oxides in ocean waters and thus to productivity,~~  
1256 ~~including the assimilation of carbon(Pester et al., 2011). The enzyme ammonia monooxygenase,~~  
1257 ~~which belongs to the enzyme family of copper containing membrane bound monooxygenases that~~  
1258 ~~possess wide substrate ranges. Closely related enzymes, which can often perform the same function~~  
1259 ~~but with different substrates, include ammonia oxidizing bacteria with ammonia (in~~  
1260 ~~*Gammaproteobacteria*) and methane as a substitute (Lontoh et al., 2000). However, a~~  
1261 ~~microorganism's metabolic energy is mainly defined by its downstream enzyme machinery~~  
1262 ~~(Tavormina et al., 2011). In our sediment *Thaumarchaeota* observed in low percentages with 2~~  
1263 ~~main classes: Marine Benthic Group B and Miscellaneous Crenarchaeotic Group.~~

1264 ~~Miscellaneous Crenarchaeotic Group~~

1265 ~~The Miscellaneous Crenarchaeotic Group (MCG) is a cosmopolitan class assigned to the~~  
1266 ~~*Thaumarchaeota* phylum whose members are found mostly in anoxic habitats and have the~~  
1267 ~~capability to take up organic carbon (Biddle et al., 2006). Because of their huge environmental~~  
1268 ~~range and their complex phylogeny, the MCG collectively possess great metabolic diversity (Jiang~~  
1269 ~~et al., 2008). Current evidence for members of the MCG lineage suggests that they may obtain~~  
1270 ~~energy from the anaerobic oxidation of methane, but they do so via a dissimilatory methane-~~  
1271 ~~oxidizing process and they do not assimilate its carbon (Biddle et al., 2006)~~

1272 ~~Marine Benthic Group B~~

1273 ~~Members of the MBGB were detected in a growing number of benthic marine environments,~~  
1274 ~~including Atlantic deep-sea sediments, marine mud volcanoes, marine carbonate crusts from the~~  
1275 ~~Black Sea, organic-poor deep subsurface sediments of the central oceanic basins, and organic-rich~~  
1276 ~~methane or methane hydrate-containing sediments near continental margins (Sørensen and Teske,~~  
1277 ~~2006). Recent results from Ocean Drilling Program showed that no known methanotroph was~~  
1278 ~~detected however methane oxidation appeared to be mediated by Marine Benthic Group B and the~~  
1279 ~~Miscellaneous Crenarchaeotal Group (Biddle et al., 2006)~~

1280 **References**

1281 Adler, M., Eckert, W. and Sivan, O.: Quantifying rates of methanogenesis and methanotrophy in  
1282 Lake Kinneret sediments (Israel) using pore-water profiles, *Limnol. Oceanogr.*, 56(4), 1525–1535,  
1283 2011.

1284 Beal, E. J., House, C. H. and Orphan, V. J.: Manganese- and Iron-Dependent Marine Methane  
1285 Oxidation, *Science*, 325(5937), 184–187, 2009.

1286 Biddle, J. F., Lipp, J. S., Lever, M. a, Lloyd, K. G., Sørensen, K. B., Anderson, R., Fredricks, H. F.,  
1287 Elvert, M., Kelly, T. J., Schrag, D. P., Sogin, M. L., Brenchley, J. E., Teske, A., House, C. H. and  
1288 Hinrichs, K.-U.: Heterotrophic Archaea dominate sedimentary subsurface ecosystems off Peru.,  
1289 *Proc. Natl. Acad. Sci. U. S. A.*, 103(10), 3846–3851, 2006.

1290 Bligh, E. G. and Dyer, W. J.: A rapid method of total lipid extraction and purification, *Can. J.*  
1291 *Biochem. Physiol.*, 37(8), 911–917, 1959.

1292 Van Bodegom, P. M., Scholten, J. C. M. and Stams, A. J. M.: Direct inhibition of methanogenesis  
1293 by ferric iron., *FEMS Microbiol. Ecol.*, 49(2), 261–8, 2004.

1294 Bodrossy, L., Holmes, E. M., Holmes, A. J., Kovács, K. L. and Murrell, J. C.: Analysis of 16S  
1295 rRNA and methane monooxygenase gene sequences reveals a novel group of thermotolerant, *Arch*  
1296 *microbiol.*, 168, 493–503, 1997.

- 1297 Boetius, A., Ravenschlag, K., Schubert, C. J., Rickert, D., Widdel, F., Gieseke, A., Amann, R.,  
1298 Jørgensen, B. B., Witte, U. and Pfannkuche, O.: A marine microbial consortium apparently  
1299 mediating anaerobic oxidation of methane, *Nature*, 407(6804), 623–626, 2000.
- 1300 Bond, D. R., Lovley, D. R. and Methanococcus, A.: Reduction of Fe ( III ) oxide by methanogens in  
1301 the presence and absence of extracellular quinones, *Environ. Microbiol.*, 4(2), 115–124, 2002.
- 1302 Borowski, W. S., Cagatay, N., Ternois, Y. and Paull, C. K.: Data report: carbon isotopic  
1303 composition of dissolved CO<sub>2</sub>, CO<sub>2</sub> gas, and methane, Blake-Bahama Ridge and northeast  
1304 Bermuda Rise, ODP Leg 172, Proc. Ocean Drill. Program, Sci. Results, 172, 1–16, 2000.
- 1305 Borrel, G., Jézéquel, D., Biderre-Petit, C., Morel-Desrosiers, N., Morel, J.-P., Peyret, P., Fonty, G.  
1306 and Lehours, A.-C.: Production and consumption of methane in freshwater lake ecosystems., *Res.*  
1307 *Microbiol.*, 162(9), 832–847, 2011.
- 1308 Borrel, G., Lehours, A.-C., Crouzet, O., Jézéquel, D., Rockne, K., Kulczak, A., Duffaud, E., Joblin,  
1309 K. and Fonty, G.: Stratification of Archaea in the deep sediments of a freshwater meromictic lake:  
1310 vertical shift from methanogenic to uncultured archaeal lineages., *PLoS One*, 7(8), e43346, 1–14,  
1311 2012.
- 1312 Borrok, D. M., Wanty, R. B., Ridley, W. I., Wolf, R., Lamothe, P. J. and Adams, M.: Separation of  
1313 copper, iron, and zinc from complex aqueous solutions for isotopic measurement, *Chem. Geol.*,  
1314 242(3-4), 400–414, 2007.
- 1315 Briée, C., Moreira, D. and López-García, P.: Archaeal and bacterial community composition of  
1316 sediment and plankton from a suboxic freshwater pond., *Res. Microbiol.*, 158(3), 213–227, 2007.
- 1317 Burow, L. C., Woebken, D., Marshall, I. P. G., Singer, S. W., Pett-Ridge, J., Prufert-Bebout, L.,  
1318 Spormann, a M., Bebout, B. M., Weber, P. K. and Hoehler, T. M.: Identification of  
1319 Desulfobacterales as primary hydrogenotrophs in a complex microbial mat community.,  
1320 *Geobiology*, 12(3), 221–230, 2014.
- 1321 Chao, A.: Nonparametric Estimation of the Number of Classes in a Population, *Scand. J. Stat.*,  
1322 11(4), 265–270, 1984.
- 1323 Chao, B. Y. A. and Ma, M.: Stopping rules and estimation for recapture debugging with unequal  
1324 failure rates, *Biometrika*, 80(1), 193–201, 1993.
- 1325 Chen, M., Yan, Y., Zhang, W., Lu, W., Wang, J., Ping, S. and Lin, M.: Complete genome sequence  
1326 of the type strain *Pseudomonas stutzeri* CGMCC 1.1803., *J. Bacteriol.*, 193(21), 6095, 2011.
- 1327 Chistoserdova, L., Vorholt, J. A. and Lidstrom, M. E.: A genomic view of methane oxidation by  
1328 aerobic bacteria and anaerobic archaea, *Genome Biol.*, 6(2), 208.1–208.6, 2005.
- 1329 Clement, J., Shrestha, J., Ehrenfeld, J. and Jaffe, P.: Ammonium oxidation coupled to dissimilatory  
1330 reduction of iron under anaerobic conditions in wetland soils, *Soil Biol. Biochem.*, 37(12), 2323–  
1331 2328, doi:10.1016/j.soilbio.2005.03.027, 2005.
- 1332 Cline, J. D.: Spectrophotometric determination of hydrogen sulfide In natural waters, *Limnol.*  
1333 *Oceanogr.*, 14, 454–458, 1969.

- 1334 Cottrell, M. T. and Kirchman, D. L.: Natural Assemblages of Marine Proteobacteria and Members  
1335 of the Cytophaga-Flavobacter Cluster Consuming Low- and High-Molecular-Weight Dissolved  
1336 Organic Matter, *Appl. Environ. Microbiol.*, 66(4), 1692–1697, 2000.
- 1337 Deutzmann, J. S. and Schink, B.: Anaerobic oxidation of methane in sediments of Lake Constance,  
1338 an oligotrophic freshwater lake., *Appl. Environ. Microbiol.*, 77(13), 4429–36,  
1339 doi:10.1128/AEM.00340-11, 2011.
- 1340 Dowd, S. E., Callaway, T. R., Wolcott, R. D., Sun, Y., McKeehan, T., Hagevoort, R. G. and  
1341 Edrington, T. S.: Evaluation of the bacterial diversity in the feces of cattle using 16S rDNA  
1342 bacterial tag-encoded FLX amplicon pyrosequencing (bTEFAP)., *BMC Microbiol.*, 8(125), 1–8,  
1343 2008.
- 1344 Eckert, T.: The Influence of Chemical Stratification in the Water Column on Sulfur and Iron  
1345 Dynamics in Pore Waters and Sediments of Lake Kinneret, Israel, M.Sc. thesis, University of  
1346 Bayreuth, Germany., 2000.
- 1347 Eckert, W. and Conrad, R.: Sulfide and methane evolution in the hypolimnion of a subtropical lake:  
1348 A three-year study, *Biogeochemistry*, 82(1), 67–76, 2007.
- 1349 Edgar, R. C., Haas, B. J., Clemente, J. C., Quince, C. and Knight, R.: UCHIME improves  
1350 sensitivity and speed of chimera detection., *Bioinformatics*, 27(16), 2194–2200, 2011.
- 1351 Edlund, A., Hårdeman, F., Jansson, J. K. and Sjöling, S.: Active bacterial community structure  
1352 along vertical redox gradients in Baltic Sea sediment., *Environ. Microbiol.*, 10(8), 2051–2063,  
1353 2008.
- 1354 Egger, M., Rasigraf, O., Sapart, C. J., Jilbert, T., Jetten, M. S. M., Röckmann, T., van der Veen, C.,  
1355 Banda, N., Kartal, B., Ettwig, K. F. and Slomp, C. P.: Iron-mediated anaerobic oxidation of  
1356 methane in brackish coastal sediments., *Environ. Sci. Technol.*, 49, 277–283, 2014.
- 1357 Ehrich, S., Behrens, D., Lebedeva, E., Ludwig, W. and Bock, E.: A new obligately  
1358 chemolithoautotrophic, nitrite-oxidizing bacterium, *Nitrospira moscoviensis* sp. nov. and its  
1359 phylogenetic relationship., *Arch. Microbiol.*, 164(1), 16–23, 1995.
- 1360 Ettwig, K. F., Van Alen, T., van de Pas-Schoonen, K. T., Jetten, M. S. M. and Strous, M.:  
1361 Enrichment and molecular detection of denitrifying methanotrophic bacteria of the NC10 phylum,  
1362 *Appl. Environ. Microbiol.*, 75(11), 3656–3662, 2009.
- 1363 Ettwig, K. F., Butler, M. K., Le Paslier, D., Pelletier, E., Mangenot, S., Kuypers, M. M. M.,  
1364 Schreiber, F., Dutilh, B. E., Zedelius, J. and de Beer, D.: Nitrite-driven anaerobic methane oxidation  
1365 by oxygenic bacteria, *Nature*, 464(7288), 543–548, 2010.
- 1366 Froelich, P. N., Klinkhammer, G. P., Bender, M. L., Luedtke, N. A., Heath, G. R., Cullen, D.,  
1367 Dauphin, P., Hammond, D., Hartman, B. and Maynard, V.: Early oxidation of organic matter in  
1368 pelagic sediments of the eastern equatorial Atlantic: suboxic diagenesis, *Geochim. Cosmochim.  
1369 Acta*, 43(7), 1075–1090, 1979.
- 1370 Glissman, K., Chin, K.-J., Casper, P. and Conrad, R.: Methanogenic pathway and archaeal  
1371 community structure in the sediment of eutrophic Lake Dagow: effect of temperature., *Microb.  
1372 Ecol.*, 48(3), 389–399, 2004.

- 1373 Good, I. J.: The population frequencies of species and the estimation of population parameters,  
1374 *Biometrika*, 40(3-4), 237–264, 1953.
- 1375 Hadas, O. and Pinkas, R.: Sulfate-reduction process in sediments of Lake Kinneret, Israel,  
1376 *Hydrobiologia*, 235(1), 295–301, 1992.
- 1377 Hallam, S. J., Girguis, P. R., Preston, C. M., Richardson, P. M. and DeLong, E. F.: Identification of  
1378 methyl coenzyme M reductase A (*mcrA*) genes associated with methane-oxidizing archaea, *Appl.*  
1379 *Environ. Microbiol.*, 69(9), 5483–5491, 2003.
- 1380 Hallam, S. J., Putnam, N., Preston, C. M., Detter, J. C., Rokhsar, D., Richardson, P. M. and  
1381 DeLong, E. F.: Reverse methanogenesis: testing the hypothesis with environmental genomics,  
1382 *Science*, 305(5689), 1457–1462, 2004.
- 1383 Hinrichs, K. U., Hayes, J. M., Sylva, S. P., Brewer, P. G. and DeLong, E. F.: Methane-consuming  
1384 archaeobacteria in marine sediments, *Nature*, 398(6730), 802–805, 1999.
- 1385 Hoehler, T. M., Alperin, M. J., Albert, D. B. and Martens, C. S.: Field and laboratory studies of  
1386 methane oxidation in an anoxic marine sediment: evidence for a methanogen-sulfate reducer  
1387 consortium, *Global Biogeochem. Cycles*, 8(4), 451–463, 1994.
- 1388 Holler, T., Wegener, G., Knittel, K., Boetius, A., Brunner, B., Kuypers, M. M. M. and Widdel, F.:  
1389 Substantial  $^{13}\text{C}/^{12}\text{C}$  and D/H fractionation during anaerobic oxidation of methane by marine  
1390 consortia enriched in vitro, *Environ. Microbiol. Rep.*, 1(5), 370–376, 2009.
- 1391 Hollister, E. B., Engledow, A. S., Hammett, A. J. M., Provin, T. L., Wilkinson, H. H. and Gentry,  
1392 T. J.: Shifts in microbial community structure along an ecological gradient of hypersaline soils and  
1393 sediments., *ISME J.*, 4(6), 829–838, 2010.
- 1394 Holmkvist, L., Ferdelman, T. G. and Jørgensen, B. B.: A cryptic sulfur cycle driven by iron in the  
1395 methane zone of marine sediment (Aarhus Bay, Denmark), *Geochim. Cosmochim. Acta*, 75(12),  
1396 3581–3599, 2011.
- 1397 Huse, S. M., Welch, D. M., Morrison, H. G. and Sogin, M. L.: Ironing out the wrinkles in the rare  
1398 biosphere through improved OTU clustering., *Environ. Microbiol.*, 12(7), 1889–1898, 2010.
- 1399 Jetten, M. S. M., Stams, A. J. M. and Zehnder, A. J. B.: Acetate threshold values and acetate  
1400 activating enzymes in methanogenic bacteria, *FEMS Microbiol. Lett.*, 73(4), 339–344, 1990.
- 1401 Jiang, H., Dong, H., Yu, B., Ye, Q., Shen, J., Rowe, H. and Zhang, C.: Dominance of putative  
1402 marine benthic Archaea in Qinghai Lake, north-western China., *Environ. Microbiol.*, 10(9), 2355–  
1403 2367, 2008.
- 1404 Jiang, L., Zheng, Y., Peng, X., Zhou, H., Zhang, C., Xiao, X. and Wang, F.: Vertical distribution  
1405 and diversity of sulfate-reducing prokaryotes in the Pearl River estuarine sediments, Southern  
1406 China., *FEMS Microbiol. Ecol.*, 70(2), 93–106, 2009.
- 1407 John, S. G. and Adkins, J. F.: Analysis of dissolved iron isotopes in seawater, *Mar. Chem.*, 119(1-  
1408 4), 65–76, 2010.
- 1409 Kanso, S., Greene, A. C. and Patel, B. K. C.: *Bacillus subterraneus* sp. nov., an iron- and  
1410 manganese-reducing bacterium from a deep subsurface Australian thermal aquifer, , 869–874, 2002.

- 1411 Kelly, D. P. and Wood, A. P.: Confirmation of *Thiobacillus denitrificans* as a species of the genus  
1412 *Thiobacillus*, in the  $\beta$  - subclass of the Proteobacteria, with strain NCIMB 9548 as the type strain, ,  
1413 547–550, 2000.
- 1414 Kinnaman, F. S., Valentine, D. L. and Tyler, S. C.: Carbon and hydrogen isotope fractionation  
1415 associated with the aerobic microbial oxidation of methane, ethane, propane and butane, *Geochim.*  
1416 *Cosmochim. Acta*, 71(2), 271–283, 2007.
- 1417 Klein, M., Friedrich, M., Roger, a J., Hugenholtz, P., Fishbain, S., Abicht, H., Blackall, L. L., Stahl,  
1418 D. a and Wagner, M.: Multiple lateral transfers of dissimilatory sulfite reductase genes between  
1419 major lineages of sulfate-reducing prokaryotes., *J. Bacteriol.*, 183(20), 6028–6035, 2001.
- 1420 Knittel, K. and Boetius, A.: Anaerobic oxidation of methane: progress with an unknown process.,  
1421 *Annu. Rev. Microbiol.*, 63, 311–334, 2009.
- 1422 Koizumi, Y., Takii, S. and Fukui, M.: Depth-related change in archaeal community structure in a  
1423 freshwater lake sediment as determined with denaturing gradient gel electrophoresis of amplified  
1424 16S rRNA genes and reversely transcribed rRNA fragments., *FEMS Microbiol. Ecol.*, 48(2), 285–  
1425 292, 2004.
- 1426 Koizumi, Y., Takii, S., Nishino, M. and Nakajima, T.: Vertical distributions of sulfate-reducing  
1427 bacteria and methane-producing archaea quantified by oligonucleotide probe hybridization in the  
1428 profundal sediment of a mesotrophic lake., *FEMS Microbiol. Ecol.*, 44(1), 101–108, 2003.
- 1429 Liu, Z., Frigaard, N.-U., Vogl, K., Iino, T., Ohkuma, M., Overmann, J. and Bryant, D. a: Complete  
1430 Genome of *Ignavibacterium album*, a Metabolically Versatile, Flagellated, Facultative Anaerobe  
1431 from the Phylum Chlorobi., *Front. Microbiol.*, 3(May), 185, 2012.
- 1432 Lloyd, K. G., Schreiber, L., Petersen, D. G., Kjeldsen, K. U., Lever, M. a., Steen, A. D.,  
1433 Stepanauskas, R., Richter, M., Kleindienst, S., Lenk, S., Schramm, A. and Jørgensen, B. B.:  
1434 Predominant archaea in marine sediments degrade detrital proteins, *Nature*, 496(7444), 215–218,  
1435 2013.
- 1436 Lontoh, S., DiSpirito, a a, Krema, C. L., Whittaker, M. R., Hooper, a B. and Semrau, J. D.:  
1437 Differential inhibition in vivo of ammonia monooxygenase, soluble methane monooxygenase and  
1438 membrane-associated methane monooxygenase by phenylacetylene., *Environ. Microbiol.*, 2(5), 485–  
1439 494, 2000.
- 1440 Lovley, D.: Dissimilatory Fe ( III ) - and Mn ( IV ) -Reducing Prokaryotes Microorganisms, in  
1441 *Prokaryotes*, pp. 635–658, Springer New York, New York., 2006.
- 1442 Lovley, D. R.: Dissimilatory Fe(III) and Mn(IV) reduction., *Microbiol. Rev.*, 55(2), 259–287, 1991.
- 1443 Lovley, D. R. and Klug, M. J.: Sulfate Reducers Can Outcompete Methanogens at Concentrations  
1444 Sulfate Reducers Can Outcompete Methanogens Sulfate Concentrationst at Freshwater, *Appl.*  
1445 *Environ. Microbiol.*, 45(1), 187–192, 1983.
- 1446 Lovley, D. R., Phillips, E. J. P., Lonergan, D. J. and Widman, P. K.: Fe ( III ) and S<sub>0</sub> reduction by  
1447 *Pelobacter carbinolicus*, *Appl. Environ. Microbiol.*, 61(6), 2132–2138, 1995.
- 1448 Luo, J.-F., Lin, W.-T. and Guo, Y.: Functional genes based analysis of sulfur-oxidizing bacteria  
1449 community in sulfide removing bioreactor., *Appl. Microbiol. Biotechnol.*, 90(2), 769–778, 2011.

- 1450 Maymo, X., Chien, Y., Gossett, J. M. and Zinder, S. H.: Isolation of a Bacterium That Reductively  
1451 Dechlorinates Tetrachloroethene to Ethene, 1995.
- 1452 McBeth, J. M., Fleming, E. J. and Emerson, D.: The transition from freshwater to marine iron-  
1453 oxidizing bacterial lineages along a salinity gradient on the Sheepscot River, Maine, USA., *Environ.*  
1454 *Microbiol. Rep.*, 5(3), 453–463, 2013.
- 1455 McDonald, I. R., Bodrossy, L., Chen, Y. and Murrell, J. C.: Molecular ecology techniques for the  
1456 study of aerobic methanotrophs., *Appl. Environ. Microbiol.*, 74(5), 1305–1315, 2008.
- 1457 McInerney, M. J., Rohlin, L., Mouttaki, H., Kim, U., Krupp, R. S., Rios-Hernandez, L., Sieber, J.,  
1458 Struchtemeyer, C. G., Bhattacharyya, A., Campbell, J. W. and Gunsalus, R. P.: The genome of  
1459 *Syntrophus aciditrophicus*: life at the thermodynamic limit of microbial growth., *Proc. Natl. Acad.*  
1460 *Sci. U. S. A.*, 104(18), 7600–7605, 2007.
- 1461 Meng, J., Xu, J., Qin, D., He, Y., Xiao, X. and Wang, F.: Genetic and functional properties of  
1462 uncultivated MCG archaea assessed by metagenome and gene expression analyses., *ISME J.*, 8(3),  
1463 650–659, 2014.
- 1464 Milucka, J., Ferdelman, T. G., Polerecky, L., Franzke, D., Wegener, G., Schmid, M., Lieberwirth,  
1465 I., Wagner, M., Widdel, F. and Kuypers, M. M. M.: Zero-valent sulphur is a key intermediate in  
1466 marine methane oxidation., *Nature*, 491(7425), 541–546, 2012.
- 1467 Miyashita, A., Mochimaru, H., Kazama, H., Ohashi, A., Yamaguchi, T., Nunoura, T., Horikoshi,  
1468 K., Takai, K. and Imachi, H.: Development of 16S rRNA gene-targeted primers for detection of  
1469 archaeal anaerobic methanotrophs (ANMEs), *FEMS Microbiol. Lett.*, 297(1), 31–37, 2009.
- 1470 Nam, Y.-D., Sung, Y., Chang, H.-W., Roh, S. W., Kim, K.-H., Rhee, S.-K., Kim, J.-C., Kim, J.-Y.,  
1471 Yoon, J.-H. and Bae, J.-W.: Characterization of the depth-related changes in the microbial  
1472 communities in Lake Hovsgol sediment by 16S rRNA gene-based approaches., *J. Microbiol.*, 46(2),  
1473 125–136, 2008.
- 1474 Nazina, T. N., Kosareva, I. M., Davidov, a. S., Tourova, T. P., Novikova, E. V., Khafizov, R. R.  
1475 and Poltarau, a. B.: Physicochemical and microbiological characteristics of groundwater from  
1476 observation wells of a deep radioactive liquid waste repository, *Microbiology*, 69(1), 89–95, 2000.
- 1477 Niemann, H., Lösekann, T., de Beer, D., Elvert, M., Nadalig, T., Knittel, K., Amann, R., Sauter, E.  
1478 J., Schlüter, M. and Klages, M.: Novel microbial communities of the Haakon Mosby mud volcano  
1479 and their role as a methane sink, *Nature*, 443(7113), 854–858, 2006.
- 1480 Nishri, a., Imberger, J., Eckert, W., Ostrovsky, I. and Geifman, Y.: The physical regime and the  
1481 respective biogeochemical processes in the lower water mass of Lake Kinneret, *Limnol. Oceanogr.*,  
1482 45(4), 972–981, 2000.
- 1483 Nusslein, B., Chin, K. J., Eckert, W. and Conrad, R.: Evidence for anaerobic syntrophic acetate  
1484 oxidation during methane production in the profundal sediment of subtropical Lake Kinneret  
1485 (Israel), *Environ. Microbiol.*, 3(7), 460–470, 2001.
- 1486 Ontiveros-Valencia, A., Ilhan, Z. E., Kang, D.-W., Rittmann, B. and Krajmalnik-Brown, R.:  
1487 Phylogenetic analysis of nitrate- and sulfate-reducing bacteria in a hydrogen-fed biofilm., *FEMS*  
1488 *Microbiol. Ecol.*, 85(1), 158–167, 2013.

- 1489 Orcutt, B. N., Sylvan, J. B., Knab, N. J. and Edwards, K. J.: Microbial ecology of the dark ocean  
1490 above, at, and below the seafloor., *Microbiol. Mol. Biol. Rev.*, 75(2), 361–422, 2011.
- 1491 Orphan, V. J., House, C. H., Hinrichs, K. U., McKeegan, K. D. and DeLong, E. F.: Methane-  
1492 consuming archaea revealed by directly coupled isotopic and phylogenetic analysis., *Science*,  
1493 293(5529), 484–487, 2001.
- 1494 Orphan, V. J., House, C. H., Hinrichs, K. U., McKeegan, K. D. and DeLong, E. F.: Multiple  
1495 archaeal groups mediate methane oxidation in anoxic cold seep sediments, *Proc. Natl. Acad. Sci.*,  
1496 99(11), 7663–7668, 2002.
- 1497 Pérez-Ibarra, B. M., Flores, M. E. and García-Varela, M.: Isolation and characterization of *Bacillus*  
1498 *thioparus* sp. nov., chemolithoautotrophic, thiosulfate-oxidizing bacterium., *FEMS Microbiol. Lett.*,  
1499 271(2), 289–296, 2007.
- 1500 Pester, M., Schleper, C. and Wagner, M.: The Thaumarchaeota: an emerging view of their  
1501 phylogeny and ecophysiology., *Curr. Opin. Microbiol.*, 14(3), 300–306, 2011.
- 1502 Pjevac, P., Kamyshny, A., Dykma, S. and Mußmann, M.: Microbial consumption of zero-valence  
1503 sulfur in marine benthic habitats., *Environ. Microbiol.*, 16(11), 3416–3430, 2014.
- 1504 Plugge, C. M., Zhang, W., Scholten, J. C. M. and Stams, A. J. M.: Metabolic flexibility of sulfate-  
1505 reducing bacteria., *Front. Microbiol.*, 2(May), 1–8, 2011.
- 1506 Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J. and Glöckner, F.  
1507 O.: The SILVA ribosomal RNA gene database project: improved data processing and web-based  
1508 tools., *Nucleic Acids Res.*, 41(Database issue), D590–D596, 2013.
- 1509 Raghoebarsing, A. A., Pol, A., Van de Pas-Schoonen, K. T., Smolders, A. J. P., Ettwig, K. F.,  
1510 Rijpstra, W. I. C., Schouten, S., Damsté, J. S. S., den Camp, H. J. M. O. and Jetten, M. S. M.: A  
1511 microbial consortium couples anaerobic methane oxidation to denitrification, *Nature*, 440(7086),  
1512 918–921, 2006.
- 1513 Röske, K., Sachse, R., Scheerer, C. and Röske, I.: Microbial diversity and composition of the  
1514 sediment in the drinking water reservoir Saidenbach (Saxonia, Germany)., *Syst. Appl. Microbiol.*,  
1515 35(1), 35–44, 2012.
- 1516 Schloss, P. D., Gevers, D. and Westcott, S. L.: Reducing the effects of PCR amplification and  
1517 sequencing artifacts on 16S rRNA-based studies., *PLoS One*, 6(12), e27310, 2011.
- 1518 Schloss, P. D., Westcott, S. L., Ryabin, T., Hall, J. R., Hartmann, M., Hollister, E. B., Lesniewski,  
1519 R. a, Oakley, B. B., Parks, D. H., Robinson, C. J., Sahl, J. W., Stres, B., Thallinger, G. G., Van  
1520 Horn, D. J. and Weber, C. F.: Introducing mothur: open-source, platform-independent, community-  
1521 supported software for describing and comparing microbial communities., *Appl. Environ.*  
1522 *Microbiol.*, 75(23), 7537–7541, 2009.
- 1523 Schoell, M.: Multiple origins of methane in the earth, *Chem. Geol.*, 71(1-3), 1–10, 1988.
- 1524 Schubert, C. J., Vazquez, F., Lösekann-Behrens, T., Knittel, K., Tonolla, M. and Boetius, A.:  
1525 Evidence for anaerobic oxidation of methane in sediments of a freshwater system (Lago di  
1526 Cadagno)., *FEMS Microbiol. Ecol.*, 76(1), 26–38, 2011.



- 1527 Schwarz, J. I. K., Eckert, W. and Conrad, R.: Community structure of Archaea and Bacteria in a  
1528 profundal lake sediment Lake Kinneret (Israel), *Syst. Appl. Microbiol.*, 30(3), 239–254, 2007a.
- 1529 Schwarz, J. I. K., Eckert, W. and Conrad, R.: Community structure of Archaea and Bacteria in a  
1530 profundal lake sediment Lake Kinneret (Israel)., *Syst. Appl. Microbiol.*, 30(3), 239–254 [online]  
1531 Available from: <http://www.ncbi.nlm.nih.gov/pubmed/16857336> (Accessed 15 March 2013b),  
1532 2007.
- 1533 Schwarz, J. I. K., Lueders, T., Eckert, W. and Conrad, R.: Identification of acetate-utilizing Bacteria  
1534 and Archaea in methanogenic profundal sediments of Lake Kinneret (Israel) by stable isotope  
1535 probing of rRNA., *Environ. Microbiol.*, 9(1), 223–237, 2007c.
- 1536 Serruya, C.: Lake Kinneret: the nutrient chemistry of the sediments, *Limnol. Oceanogr.*, 16(May),  
1537 510–521, 1971.
- 1538 Serruya, C., Edelstein, M., Pollinger, U. and Serruya, S.: Lake Kinneret sediments: nutrient  
1539 composition of the pore water and mud water exchanges, *Limnol. Oceanogr.*, 19(3), 489–508, 1974.
- 1540 Severmann, S., Johnson, C. M., Beard, B. L. and McManus, J.: The effect of early diagenesis on the  
1541 Fe isotope compositions of porewaters and authigenic minerals in continental margin sediments,  
1542 *Geochim. Cosmochim. Acta*, 70(8), 2006–2022, 2006.
- 1543 Shah, M., Lin, C.-C., Kukkadapu, R., Engelhard, M. H., Zhao, X., Wang, Y., Barkay, T. and Yee,  
1544 N.: Syntrophic Effects in a Subsurface Clostridial Consortium on Fe(III)-(Oxyhydr)oxide Reduction  
1545 and Secondary Mineralization, *Geomicrobiol. J.*, 31(2), 101–115, 2014.
- 1546 Sivan, O., Adler, M., Pearson, A., Gelman, F., Bar-Or, I., John, S. G. and Eckert, W.: Geochemical  
1547 evidence for iron-mediated anaerobic oxidation of methane, *Limnol. Oceanogr.*, 56(4), 1536–1544,  
1548 2011.
- 1549 Smith, P. F., Langworthy, T. A. and Smith, M. R.: Polypeptide nature of growth requirement in  
1550 yeast extract for *Thermoplasma* Polypeptide Nature of Growth Requirement in Yeast Extract for  
1551 *Thermoplasma acidophilum*, *J. Bacteriol.*, 124(2), 884–892, 1975.
- 1552 Sonne-Hansen, J. and Ahring, B. K.: *Thermodesulfobacterium hveragerdense* sp. nov., and  
1553 *Thermodesulfobacterium islandicus* sp. nov., two thermophilic sulfate reducing bacteria isolated from a  
1554 Icelandic hot spring., *Syst. Appl. Microbiol.*, 22(4), 559–564, 1999.
- 1555 Sørensen, K. B. and Teske, A.: Stratified communities of active Archaea in deep marine subsurface  
1556 sediments., *Appl. Environ. Microbiol.*, 72(7), 4596–4603, 2006.
- 1557 Spring, S., Amann, R., Ludwig, W., Schleifer, K. H., van Gemerden, H. and Petersen, N.:  
1558 Dominating role of an unusual magnetotactic bacterium in the microaerobic zone of a freshwater  
1559 sediment., *Appl. Environ. Microbiol.*, 59(8), 2397–2403, 1993.
- 1560 Stookey, L. L.: Ferrozine-a new spectrophotometric reagent for iron, *Anal. Chem.*, 42(7), 779–781,  
1561 1970.
- 1562 Sun, H., Spring, S., Lapidus, A., Davenport, K., Del Rio, T. G., Tice, H., Nolan, M., Copeland, A.,  
1563 Cheng, J.-F., Lucas, S., Tapia, R., Goodwin, L., Pitluck, S., Ivanova, N., Pagani, I., Mavromatis, K.,  
1564 Ovchinnikova, G., Pati, A., Chen, A., Palaniappan, K., Hauser, L., Chang, Y.-J., Jeffries, C. D.,  
1565 Detter, J. C., Han, C., Rohde, M., Brambilla, E., Göker, M., Woyke, T., Bristow, J., Eisen, J. a,

- 1566 Markowitz, V., Hugenholtz, P., Kyrpides, N. C., Klenk, H.-P. and Land, M.: Complete genome  
1567 sequence of *Desulfarculus baarsii* type strain (2st14)., *Stand. Genomic Sci.*, 3(3), 276–284, 2010.
- 1568 Tamaki, H., Sekiguchi, Y., Hanada, S., Nomura, N., Matsumura, M., Nakamura, K. and Kamagata,  
1569 Y.: Comparative Analysis of Bacterial Diversity in Freshwater Sediment of a Shallow Eutrophic  
1570 Lake by Molecular and Improved Cultivation-Based Techniques Comparative Analysis of Bacterial  
1571 Diversity in Freshwater Sediment of a Shallow Eutrophic Lake by Molecul, *Appl. Environ.*  
1572 *Microbiol.*, 71(4), 2162–2169, 2005.
- 1573 Tavormina, P. L., Orphan, V. J., Kalyuzhnaya, M. G., Jetten, M. S. M. and Klotz, M. G.: A novel  
1574 family of functional operons encoding methane/ammonia monooxygenase-related proteins in  
1575 gammaproteobacterial methanotrophs, *Environ. Microbiol. Rep.*, 3(1), 91–100, 2011.
- 1576 Tavormina, P. L., Ussler, W. and Orphan, V. J.: Planktonic and sediment-associated aerobic  
1577 methanotrophs in two seep systems along the North American margin., *Appl. Environ. Microbiol.*,  
1578 74(13), 3985–2995, 2008.
- 1579 Teske, A. and Sørensen, K. B.: Uncultured archaea in deep marine subsurface sediments: have we  
1580 caught them all?, *ISME J.*, 2(1), 3–18, 2008.
- 1581 Thabet, O. B. D., Fardeau, M.-L., Joulain, C., Thomas, P., Hamdi, M., Garcia, J.-L. and Ollivier, B.:  
1582 *Clostridium tunisiense* sp. nov., a new proteolytic, sulfur-reducing bacterium isolated from an olive  
1583 mill wastewater contaminated by phosphogypse., *Anaerobe*, 10(3), 185–190, 2004.
- 1584 Thauer, R. K.: Functionalization of methane in anaerobic microorganisms., *Angew. Chem. Int. Ed.*  
1585 *Engl.*, 49(38), 6712–6713, 2010.
- 1586 Valentine, D. L.: Biogeochemistry and microbial ecology of methane oxidation in anoxic  
1587 environments: a review, *Antonie Van Leeuwenhoek*, 81(1-4), 271–282, 2002.
- 1588 Wasmund, K., Lars, S., Lloyd, K. G., Petersen, D. G., Schramm, A., Stepanauskas, R., Jørgensen,  
1589 B. B. and Adrian, L.: Genome sequencing of a single cell of the widely distributed marine  
1590 subsurface Dehalococcoidia, phylum Chloroflexi, *ISME J.*, 8, 383–397, 2013.
- 1591 Whiticar, M. J.: Carbon and hydrogen isotope systematics of bacterial formation and oxidation of  
1592 methane, *Chem. Geol.*, 161(1-3), 291–314, 1999.
- 1593 Wobus, A., Bleul, C., Maassen, S., Scheerer, C., Schuppler, M., Jacobs, E. and Röske, I.: Microbial  
1594 diversity and functional characterization of sediments from reservoirs of different trophic state.,  
1595 *FEMS Microbiol. Ecol.*, 46(3), 331–347, 2003.
- 1596 Yamada, C., Kato, S., Kimura, S., Ishii, M. and Igarashi, Y.: Reduction of Fe(III) oxides by  
1597 phylogenetically and physiologically diverse thermophilic methanogens., *FEMS Microbiol. Ecol.*,  
1598 89(3), 637–645, 2014.
- 1599 Yamada, T. and Sekiguchi, Y.: Cultivation of Uncultured Chloroflexi Subphyla: Significance and  
1600 Ecophysiology of Formerly Uncultured Chloroflexi “Subphylum I” with Natural and  
1601 Biotechnological Relevance, *Microbes Environ.*, 24(3), 205–216, 2009.
- 1602 Yamada, T., Sekiguchi, Y., Hanada, S., Imachi, H., Ohashi, A., Harada, H. and Kamagata, Y.:  
1603 *Anaerolinea thermolimos* sp. nov., *Levilinea saccharolytica* gen. nov., sp. nov. and *Leptolinea*  
1604 *tardivitalis* gen. nov., sp. nov., novel filamentous anaerobes, and description of the new classes

- 1605 Anaerolineae classis nov. and Caldilineae classis nov. in the, *Int. J. Syst. Evol. Microbiol.*, 56(Pt 6),  
1606 1331–1340, 2006.
- 1607 Yan, Z., Song, N., Cai, H., Tay, J.-H. and Jiang, H.: Enhanced degradation of phenanthrene and  
1608 pyrene in freshwater sediments by combined employment of sediment microbial fuel cell and  
1609 amorphous ferric hydroxide., *J. Hazard. Mater.*, 199-200, 217–225, 2012.
- 1610 Ye, W., Liu, X., Lin, S., Tan, J., Pan, J., Li, D. and Yang, H.: The vertical distribution of bacterial  
1611 and archaeal communities in the water and sediment of Lake Taihu., *FEMS Microbiol. Ecol.*, 70(2),  
1612 107–120, 2009.
- 1613 Yoshida, N., Takahashi, N. and Hiraishi, A.: Phylogenetic Characterization of a Polychlorinated-  
1614 Dioxin- Dechlorinating Microbial Community by Use of Microcosm Studies, , 71(8), 4325–4334,  
1615 2005.
- 1616 Zhu, B., van Dijk, G., Fritz, C., Smolders, A. J. P., Pol, A., Jetten, M. S. M. and Ettwig, K. F.:  
1617 Anaerobic oxidization of methane in a minerotrophic peatland: enrichment of nitrite-dependent  
1618 methane-oxidizing bacteria., *Appl. Environ. Microbiol.*, 78(24), 8657–8665, 2012.
- 1619
- 1620