Biogeosciences Discuss., 11, C4145–C4148, 2014 www.biogeosciences-discuss.net/11/C4145/2014/ © Author(s) 2014. This work is distributed under the Creative Commons Attribute 3.0 License.



BGD

11, C4145-C4148, 2014

Interactive Comment

Interactive comment on "Effects of experimental nitrogen deposition on peatland carbon pools and fluxes: a modeling analysis" by Y. Wu et al.

Anonymous Referee #1

Received and published: 6 August 2014

Wu and colleages tackle an important problem within the global carbon cycle. In their model study they analyze the behavior of peatland ecosystems under potential N fertilization with a sophisticated model. Their finding is well presented and concluded. The problem description and wording of the highly divers and complex peatland processes is comprehensible. I thus suggest the paper for publication after some minor corrections as suggested below.

General:

One of the novelties of the paper I consider the test of the GEPmax dependence on leaf N content. In my view this deserves an additional exposition in the introduction and being mentioned in the abstract.

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



It is hard to judge how robust the response of the different PFTs to N fertilization actualy is. Granath et al., 2009 suggest that some Sphagnum species behave differently and can actually adapt to higher leaf N content. Would a scenario with mosses having a higher max N tolerance in the N factor on GEPmax than in the original model also be possible? In this case mosses would simply be outcompeted by graminoids/shrubs through growth and light competition. Can you exclude this possibility?

Reading this model study I often found myself looking up the PEATBOG model paper (Wu and Blodau, 2013) in order to find the corresponding equation. I don't think this is a bad thing per se, regarding the complexity of the problem, but repeating the main model equations for a given sensitivity could greatly improve the visibility for the reader. My suggestions are e.g. the model eqs. for (i) the dependence of GEPmax on leaf N content, (ii) the C/N effect on ER and (iii) the competition for N uptake.

Specific:

- p. 10274, I. 17: Please add global estimates/modelling studies for future peatland C storage changes under N fertilization and dynamic peatland vegetation change, e.g. results from Spahni et al., 2013.
- p. 10277, l. 14: missing "g N" units
- p. 10277, l. 28: Please be careful with the wording. If you define GEP (gross ecosystem production) as photosynthesis, I assume you mean gross carbon assimilation during photosynthesis. Later on photosynthesis is shown beeing dependent on leaf N content. But there are two parts of N allocation as you explain 10 pages later: once to photosynthetically active processes and once to biomass growth. Please be more precise in general and early on, when you use the term "photosynthesis" as this is a major part of the paper.
- p. 10277, I. 28: To my understanding and according to Figure 1 this sentence is wrong, it should be GEP+ER = NEE and not ER+NEE = GEP.

BGD

11, C4145-C4148, 2014

Interactive Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- p. 10279, l. 5: Do 10000 years reflect the basal age of Mere Bleue Bog? Is the model in a steady state regarding net ecosystem carbon accumulation?
- p. 10279, l. 20: typo "leaf area index"
- p. 10283, I. 5: I can't follow the last statements. Do sphagnum and vascular plant biomass have different properties that directly affect the rate of respiration in the model? Please clarify.
- p. 10287, l. 9: Is the C/N increase because of the shift in vegetation composition?
- p. 10288, l. 26: Just to get it right, for the comparsions of the modifications 1,2,3 you do not correct GEP, ER, NEE anymore as you did in Fig. 1?
- p. 10290, l. 3: Where can I find supplementary Figures?
- p. 10290, l. 8: Use 'Mer Bleue' instead of 'MB' or define it at the first occurance of the name in the text.
- p. 10291, I. 18: How does the model handle competition for N uptake exactly? Assuming a bog ecosystem with limited N avilability: is there a priority rule for PFTs accessing nutrients? Would that change in the case for N fertilization and thus increased GEP?

Tables:

- Table 3 caption is missing units: g C m-2?

Figures:

- Fig. 1 caption has a typos: "The green dotted lines in? represent weekly averaged CO2 flux ...", and "Note that P was K not constrained in the model. "Please correct.
- Fig. 4: Figure shows "FPT" instead of "EPT"
- Fig. 8: Please write out 'sh' and 'tr' in the line description. I guess 'sh' means shoot not shrub. Also correct 'shurb'.

BGD

11, C4145-C4148, 2014

Interactive Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



References:

Granath, G., Strengbom, J., Breeuwer, A., Heijmans, M. M., Berendse, F., and Rydin, H.: Photosynthetic performance in Sphagnum transplanted along a latitudinal nitrogen deposition gradient, Oecologia, 159, 705–715, doi:10.1007/s00442-008-1261-1, 2009.

Wu, Y. and Blodau, C.: PEATBOG: a biogeochemical model for analyzing coupled carbon and nitrogen dynamics in northern peatlands, Geosci. Model Dev., 6, 1173-1207, doi:10.5194/gmd-6-1173-2013, 2013.

Spahni, R., Joos, F., Stocker, B. D., Steinacher, M., and Yu, Z. C.: Transient simulations of the carbon and nitrogen dynamics in northern peatlands: from the Last Glacial Maximum to the 21st century, Clim. Past, 9, 1287-1308, doi:10.5194/cp-9-1287-2013, 2013.

Interactive comment on Biogeosciences Discuss., 11, 10271, 2014.

BGD

11, C4145-C4148, 2014

Interactive Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

