

Interactive comment on “Comparison of seven packages that compute ocean carbonate chemistry” by J. C. Orr et al.

J. C. Orr et al.

james.orr@lsce.ipsl.fr

Received and published: 14 August 2014

Response to Short Comment by Dr. Ernie Lewis and Dr. Douglas Wallace

We thank Drs. Lewis and Wallace for their comments on this manuscript. Those are repeated in gray, whereas our response is in black.

We read with interest the manuscript comparing the various packages that calculate parameters of the CO₂ system in seawater and would like to commend the authors for their authoritative comparison of these programs. We found the review insightful, balanced, and informative, and were amused to see that conversions among pH scales and pressure corrections remain a source of mystery, error, and ambiguity, as

C4395

they have for the last 20-odd years. We have one minor comment on the manuscript: that the programming language for the original CO₂SYS be listed as QBasic rather than DOS (which is the operating system under which the program runs); this would make description of the original CO₂SYS more parallel to other versions for which languages such as C, Fortran, Excel, etc. are listed.

We will improve the description of CO₂SYS in the revised manuscript, mentioning that it is written in QBasic, not DOS. Thank you for this catching this glitch.

As authors of the original version of CO₂SYS, upon which most of the other programs were based, we thought that some historical perspective might be of interest. We also highlight some lesser known features of the project that we consider to be of continued utility. The version of CO₂SYS used in the manuscript by Orr et al. as the standard for comparison is the MATLAB version. It is one of the versions that took the original CO₂SYS code written and made available by us and translated by others into Excel or MATLAB (the manuscript mentions that there are four versions of CO₂SYS). Neither of us were actively involved in translating CO₂SYS into other languages, although we worked with others in some instances to fix errors in those versions. It is worth noting that the original CO₂SYS, and earlier programs from which it evolved, were based on DOE (1994), Handbook of Methods for the Analysis of the Various Parameters of the Carbon Dioxide System in Sea Water, version 2, eds. A. G. Dickson and C. Goyet, ORNL/CDIAC-74, and a later version of this handbook, under Andrew Dickson's oversight, remains the ultimate standard of "best practices" for this topic.

This comment raises an important point about the lineage of the translations of CO₂SYS into Excel and Matlab, which will do our best to relay in the revised manuscript. We will also state that CO₂SYS was based on Dickson and Goyet (1994).

C4396

The original CO2SYS is still widely used, and to this day we receive several emails each month with questions or comments on it. It has held up to the test of time very well, and to the best of our knowledge it is error-free (the one item we would change is to allow more spaces for outputs; for instance, a value for fCO₂ of 12,345.6 μ atm would be written "%12345.6" with the "%" denoting that an insufficient number of spaces were allowed to print the output). Although it is perhaps a bit dated at this point—there are more recent sets of constants that are not included, and it is difficult to run on current operating systems (although we are able to run it on the 64-bit version of Windows 7 and on Mac OS-X using DOSBox)—the original version has some useful features that we think should be included in any similar, follow-on program.

The online help feature in the original CO2SYS, albeit perhaps a bit crude by today's standards, explains the program, describes the various options, and lists references for the constants used in the program. There is also an accompanying report (available at <http://cdiac.ornl.gov/oceans/co2rprt.html>) that describes the program and provides a list of typographic errors in the relevant literatures (this will be discussed further below). The list of options in CO2SYS is comprehensive and covers most if not all of what would have been of any utility at the time the program was written: the choice of any two of the four CO₂ system parameters (total alkalinity, total dissolved inorganic carbon, pH, and either fCO₂ or pCO₂); the choice of fCO₂ or pCO₂; the choice of either of two definitions of total alkalinity; the choice of four pH scales; the choice of eight sets of CO₂ dissociation constants (including a freshwater choice and the GEOSECS choice, which would now be of historical interest only); the choice of two constants for dissociation of sulfate; and the choice of both input and output temperature and pressure (which readily allows for conversion between in situ and standard reporting conditions). The original CO2SYS also runs in single-input mode or batch mode (with either tab-separated or comma-separated inputs).

C4397

The CO2SYS report (Lewis and Wallace, 1998) proved valuable for our comparison and was cited heavily in the Discussion paper. We used CO2SYS's batch mode for our comparison of CO2SYS with its translations into Excel and Matlab.

Other features of CO2SYS that would (in our opinion) be useful for any similar program today involve the quantities calculated. The following quantities are reported in CO2SYS at both input and output conditions (i.e., temperature and pressure): the other two CO₂ system parameters; contributions to the alkalinity from various species; speciation of CO₂ (i.e., dissolved CO₂, carbonate, and bicarbonate); fCO₂, pCO₂, and xCO₂ (the latter in dry air at 1 atm); saturation of both calcite and aragonite; the Revelle factor; the value of pH on all four pH scales; and the values of the first and second dissociation constants of CO₂ (K₁ and K₂), and of the dissociation constants for borate and the hydroxyl ion (KB and KW) on all four pH scales. Additionally, the program in single input mode calculates and lists the sensitivities of the two calculated CO₂ system parameters to uncertainties in the two input parameters; salinity; both input and output conditions (i.e., temperature and pressure); and K₁, K₂, and the solubility of CO₂ in seawater (K₀). We feel that this last feature, what we call "partials," is one of the most useful attributes of the program for teaching purposes and is especially important, as it readily illustrates how accurately the calculated quantities can be determined.

It is a valuable feature to be able to compute calculated variables both at input and output temperatures, and it remains a feature that is only offered by CO2SYS and its derivatives. The partials feature offered by the original CO2SYS program has unfortunately been lost in the CO2SYS derivatives. Those calculations will be critical to recover for those who wish to do error propagation or compute a range of buffer factors.

The original CO2SYS is a combination of earlier programs CO2SYSTEM.EXE,

C4398

FCO2TCO2.EXE, PHTCO2.EXE, and CO2BTCH.EXE that were released in May, 1995 (this was back in the days when program names were limited to eight characters, with a limit of three characters for the extension). To the best of our knowledge, no serious errors have been found in these programs (the fits listed for the dissociation constants are not valid for extrapolation to salinity 0, but as the programs were designed for the oceanic CO₂ system, this wasn't an issue). CO2SYSTM (for which there were versions 1.00, 1.01, and 1.02) calculated any two of the four CO₂ system parameters from the other two, allowed for a choice of four sets of CO₂ dissociation constants, and included sensitivity of the outputs to the two CO₂ input system parameters, temperature and salinity, and K₀, K₁, and K₂; speciation of CO₂ (i.e., dissolved, bicarbonate, and carbonate); and contributions to the alkalinity at both input and output conditions. However, all inputs and outputs were on the total pH scale using Dickson's value for the dissociation constant of sulfate. FCO2TCO2.EXE and PHTCO2.EXE were similar, but the former is for fCO₂ or pCO₂ and the latter is for pH. Both use total dissolved inorganic carbon and either pH or fCO₂ (or pCO₂) as inputs and allow choice of carbonate constants, fCO₂ or pCO₂ as the desired variable, and input and output temperatures. The program then calculates the other CO₂ system variables at both input and output temperatures and sensitivities of these variables to input temperature.

Thank you for this historical perspective. We may well add some of it to the description of CO2SYS in the revised manuscript.

Program CO2SYS.EXE replaced and extended the above four programs. At the time, there were no publicly available programs to perform these types of calculations, and the few in-house programs had no conformity on pH scales, CO₂ dissociation constants, etc. Version 0.00 (February, 1997) was preliminary and the pressure corrections to the carbonate solubilities were incorrect. Version 1.02 (March, 1997)

C4399

contained incorrect pressure corrections to the pH scale conversions (although this has only a minor effect on outputs in most cases). Version 1.03 (May, 1997) is believed to be correct. Version 1.04 (May, 1997) and Version 1.05 (October, 1997) differ from Version 1.03 only cosmetically (typos corrected, more efficient codes, etc.), and no differences exist between calculations performed using these versions and Version 1.03. The original CO2SYS was written in QBASIC, which we chose because it ran on any 80x86 computer with the DOS operating system by typing the program name on the command line. We intended that the program could be used for decades: in our infinite wisdom, despite the advent of early versions of Windows, we felt that DOS would be the operating system of the future and we placed our bets on this. In our defense, it was probably the best choice at the time: an executable file could be created and distributed and it would run on any DOS-based computer without the necessity to install anything else (the same cannot be said for MATLAB or other programs, which require more than merely an executable to run). The program did not run on Macs, although they were not as common back then. Computers were very different back then, in ways that are probably difficult for younger programmers to understand. CO2SYS had to be broken into two parts and compiled separately, then combined. Lest this give the impression that it was a large program, consider the following text from the program: "PROGRAMMER'S NOTE: This program is DANGEROUSLY close to the DOS-imposed 64K limit : : : Don't make any unnecessary changes or the limit will be exceeded."

Predicting the future is dangerous, especially when its about computers. The fact that CO2SYS still runs today is an accomplishment.

As stated above, there was a large section in the CO2SYS report on typographic errors in the literature. One common feature in oceanographic papers not seen in other fields is the use of check values for expressions for dissociation constants and other

C4400

quantities; i.e., the value of the expression at a specified set of conditions (salinity, temperature, pressure). These check values are of immense help in determining if coefficients are listed correctly in the manuscript and if they are incorporated correctly in computer codes. Errors in listed coefficients or if their transcription can cost hours of time, in addition to leading to incorrect values. Thus, a compilation of such errors would be very useful for the community. We had compiled such a list while writing CO2SYS, which we were originally hesitant about presenting publicly, for several obvious reasons. However, Rik Wanninkhof convinced us that making this list available would be a service to the community, and we think that it has been. Considering how often such typographic errors continue to appear in the literature, we recommend that the outdated version on the CDIAC website (<http://cdiac.ornl.gov/oceans/co2rprt.html#typos>) be updated and made available. Perhaps this manuscript will prompt those who have updated CO2SYS or similar programs by including more recent constants will make those available and someone will come forward to collate and publish them.

The list of typographical errors provided in Lewis and Wallace (1998) has been a great service to the community. We used that for this comparison and when developing individual packages (seacarb and mocsy). Updating that list should be a high priority for the field.

We thank Drs. Lewis and Wallace for this historical perspective, which we hope package developers can take advantage of to make improvements in the future.

References

- Dickson, A. G. and Goyet, C.: Handbook of methods for the analysis of the various parameters of the carbon dioxide system in seawater; version 2, CDIAC-74, ORNL, 187 pp., 1994.
Lewis, E. and Wallace, D. W. R.: Program Developed for CO₂ System Calculations, C4401

ORNL/CDIAC-105, Carbon Dioxide Inf. Anal. Cent., Oak Ridge Natl. Lab., Oak Ridge, Tenn., 38 pp., 1998.

Interactive comment on Biogeosciences Discuss., 11, 5327, 2014.