

Interactive comment on “Components of near-surface energy balance derived from satellite soundings – Part 1: Net available energy” by K. Mallick et al.

Anonymous Referee #1

Received and published: 27 August 2014

The paper of Mallick et al (2014) presents a simple and potentially interesting approach for estimating midday net radiation and net available energy at the Earth' surface ($Rn - G$) from satellite sounders. The method has been applied to data acquired twice daily by AIRS and MODIS sounders, obtained or aggregated at 1° resolution and monthly averaged, and extensively evaluated with the help of ground observations of turbulent heat fluxes from 30 FLUXNET stations. This validation allowed the authors to draw conclusions on the quality of the estimates for each biome considered.

The paper is within the scope of BG journal, globally well written, and the results well presented in a concise style that the readers may appreciate. However, the paper intends to address multiple objectives at the same time (a new methodology to compute

C4681

surface heat capacity, application with satellite data and generation of a dataset for use in an evapotranspiration model presented in a second paper). This makes the paper not very focused on clear objectives, and sometimes do not address fully each of them properly. The paper would probably gain in focusing on one objective only (the methodology, for example), or merge the two papers intended for the series. From my point of view, the method to derive $Rn-G$, and in particular surface heat capacity, is particularly interesting as such, although some assumptions need to be checked, as well as the extensive validation at FLUXNET sites. In that perspective, I think that the paper should be more focused on the method, which is the most innovative part of the study. I would therefore strongly recommend to first validate the new method at FLUXNET sites using ground observations of the surface radiation components as input, before applying it with satellite data, as the latter part has already, but partially, been done in other studies with a high degree of success (eg Verstraeten et al (2005) with NOAA-AVHRR for instantaneous net radiation). An alternative would be to integrate Paper 1 (this paper) into Paper 2 (paper on the Bowen ratio), as it would justify better some choices made in the methodology (monthly time scale, combination of AIRS and MODIS data).

Besides, this recommendation, I have a couple of fundamental questions on choices undertaken in this study.

1 Questions on the choice of specific satellite data for this study.

- After reading the paper, it is not clear to me why a combination of AIRS and MODIS data has been chosen. Is there any reason to prefer the use of AIRS data in this context compared to MODIS (as described in Peng et al, 2013) ? Certainly, the use of AIRS data make the validation task much more difficult because of the huge difference in spatial footprint, and this choice imposes the results to be concentrated on 1 time slot per day, 13:30 local time. Other satellites could possibly have been used to have more passes per day.

C4682

- A monthly time scale has been chosen for the study. The perspective of this choice is not exposed. One may wonder why daily or weekly time scale has not been selected, as it could be at least equally or more interesting to study the daily time scale for monitoring.

2 Questions on the assumptions.

- I found the method to derive net available energy interesting, and certainly very promising for further applications. However, the paper would gain in credibility if all the assumptions were checked to be realistic in using ground observations. The assumptions to be checked are: the symmetry of the T_s difference over a month, $Rn - G = 0$ at 1:30 AM LT (Although there is already an indication of it in the paper, an information of the distribution of the $Rn - G$ at night may be useful).
- It is assumed that the estimates at 1° can be directly compared with point scale observations at the surface, which is an uncertain hypothesis. The authors themselves point out in the paper that the scale mismatch can be a source of discrepancy in their results, but without quantifying this effect. I would suggest to first verify the validity of the new methodology to compute $Rn-G$ with ground observations only (if possible, or at least with satellite data at finer scale), and then to apply it at global scale with satellite soundings. This would certainly help in both acknowledging the accuracy of the new method and better understand the effect of scale mismatch.

3 On the structure and text of the paper.

- The time scale should be mentioned in the title, the abstract and the introduction.

C4683

Table 1. RMSD [$W.m^{-2}$] on net radiation found in the litterature.

Instantaneous					Daily averages			Monthly averages	Instantaneous monthly averages
5-45	74	19.20	23/39	58-142	60	37.72/32	33.38	44	74-126
Verstraeten et al, 2005	Bisht et al, 2005	Cai et al, 2007	Bisht and Bras, 2010	Huang et al, 2014	Bisht et al, 2005	Huang et al, 2014	Peng et al, 2013	Jin et al, 2011	This study
NOAA	MODIS								AIRS/MODIS

- The satellite datasets should be described first to interpret correctly the methods as they imposes constrains, or the methods should be reformulated to avoid relying on AIRS and MODIS specific products, leaving their description for after.
- In the discussion section, the authors give a series of references to support the quality of their estimations. From my point of view, I found difficult to state the quality of the proposed estimations compared to the other studies as such: some references concern instantaneous values, others daily averages, some of them on very limited samples, but all with satellites soundings at finer scale (from MODIS, GOES). If I add three other references I found (Verstraeten et al, 2005; Jin et al, 2011; Peng et al, 2013), it is still difficult to know how to compare to the RMSD found in this study with others (See Table below). The suggestion of point 2 would certainly help the readers to apprehend the effect of scale discrepancy with the error due to the methods, assumptions. In addition, the authors could explain how to compare the different results from other studies with theirs.

C4684

- The last paragraph of the discussion section should be moved (and reformulated) to Introduction, as it justifies some choices, and make a link to the second paper.
- The conclusion section should state the conclusion based on the results obtained, the potential applicability to other sensors and perspectives. A part of the conclusion section (first paragraph) should be moved to the discussion section, as limitations of the method are discussed. A clear conclusion should be stated here. Applicability to other sensors is briefly mentioned by citing other satellite missions, but sometimes sounds a bit far stretched, without the authors giving a clue on how to proceed (for example, how to apply the method initially tested for monthly averages once a day to get 30 min estimations of Rn-G ?). Therefore, I would suggest either to limit the sensors list to which the method can be applied, or give a short explanation on how to proceed (the option I would recommend).

4 Typos and additional references

- p11828, eq 1: $\phi = \lambda E + H$
- p11832, eq 8: $-\lambda E(t)$
- Verstraeten W., Veroustraete, F., and Feyen, J., 2005: Estimating evapotranspiration of European forests from NOAA-imagery at satellite overpass time: Towards an operational processing chain for integrated optical and thermal sensor data products, *Remote Sensing of Environment*, 96 (2), 256-276.
- Peng, J., Liu, Y., Zhao, X., and Loew, A. (2013): Estimation of evapotranspiration from MODIS TOA radiances in the Poyang Lake basin, China, *Hydrol. Earth Syst. Sci.*, 17, 1431-1444.

C4685

- Jin, Y., Randerson, J. T., and Goulden, M. L. (2011): Continental-scale net radiation and evapotranspiration estimated using MODIS satellite observations, *Remote Sensing of Environment*, 115, 2302-2319.

Interactive comment on *Biogeosciences Discuss.*, 11, 11825, 2014.

C4686