

1 **Enhanced viral production and virus-mediated mortality of**  
2 **bacterioplankton in a natural iron-fertilized bloom event**  
3 **above the Kerguelen Plateau**

4

5 **Andrea Malits<sup>1,2</sup>, Urania Christaki<sup>3</sup>, Ingrid Obernosterer<sup>4,5</sup> and Markus G.**

6 **Weinbauer<sup>1,2</sup>**

7

8

9 <sup>1</sup> Sorbonne Universités, UPMC Univ Paris 06, UMR 7093, LOV, Observatoire océanographique,  
10 F-06230, Villefranche/mer, France

11 <sup>2</sup> CNRS, UMR 7093, LOV, Observatoire océanographique, F-06230, Villefranche/mer, France

12 <sup>3</sup> INSU-CNRS, UMR 8187 LOG, Laboratoire d'Océanologie et des Géosciences, Université du  
13 Littoral, ULCO, 32 avenue Foch, F-62930 Wimereux, France

14 <sup>4</sup> Sorbonne Universités, UPMC Univ Paris 06, UMR 7621, LOMIC, Observatoire  
15 océanographique, F-66650 Banyuls/mer, France

16 <sup>5</sup> CNRS, UMR 7621, LOMIC, Observatoire océanographique, F-66650 Banyuls/mer, France

17

18

19 | Correspondence to: A. Malits ([amalits@cadic-conicet.gob.ar](mailto:amalits@cadic-conicet.gob.ar))

20

21 | Running title: Enhanced viral production in iron-fertilized waters

22

1 **Abstract**

2 Above the Kerguelen Plateau in the Southern Ocean natural iron fertilization sustains a large  
3 phytoplankton bloom over three months during austral summer. During the KEOPS1 project  
4 (KErguelen Ocean and Plateau compared Study1) we sampled this phytoplankton bloom during  
5 its declining phase along with the surrounding HNLC waters to study the effect of natural iron  
6 fertilization on the role of viruses in the microbial food web. Bacterial and viral abundances were  
7 1.7 and 2.1 times, respectively, higher within the bloom than in HNLC waters. Viral production  
8 and virus-mediated mortality of bacterioplankton was 4.1 and 4.9 times, respectively, higher in  
9 the bloom, while the fraction of infected cells (FIC) and the fraction of lysogenic cells (FLC)  
10 showed no significant differences between environments. The present study suggests viruses to  
11 be more important for bacterial mortality within the bloom and dominate over grazing of  
12 heterotrophic nanoflagellates (HNF) during the late bloom phase. As a consequence, at least at a  
13 late bloom stage, viral lysis shunts part of the photosynthetically fixed carbon in iron-fertilized  
14 regions into the dissolved organic matter (DOM) pool with potentially less particulate organic  
15 carbon transferred to larger members of the food web or exported.

16

Andrea Malits 10/6/14 3:09 P.M.  
Eliminado: protozoan

1

## 2 1 Introduction

3 A quarter of a century ago the importance of viruses as the most abundant biological entity in the  
4 oceans (Bergh et al., 1989) and their role in the material and energy cycles have been recognized  
5 (Proctor and Fuhrman, 1990; Suttle et al., 1990). Shortly afterwards, Smith et al. (1992)  
6 conducted the first study on viral distribution and their relationship to bacteria in the Southern  
7 Ocean. Since then studies on viral abundance and production or infectivity in the cold high  
8 latitude marine environments remained limited or are only recently accumulating (Bird et al.,  
9 1993; Brussaard et al., 2008b; Evans and Brussaard, 2012; Evans et al., 2009; Guixa-Boixereu et  
10 al., 2002; Higgins et al., 2009; Manganelli et al., 2009; Marchant et al., 2000; Payet and Suttle,  
11 2008, 2013; Smith et al., 1992; Steward et al., 1996; Strzepek et al., 2005; Weinbauer et al., 2009a).  
12 These observations demonstrate that viruses are ecologically as important in these cold  
13 environments as in other world's oceans.

14 Viral lysis of cells converts particulate organic matter into dissolved and colloidal organic matter,  
15 reduces the carbon flow to higher trophic levels and increases the residence time of carbon and  
16 mineral nutrients in the euphotic zone (Fuhrman, 1999). By this process, called the "viral shunt"  
17 (Wilhelm and Suttle, 1999), heterotrophic bacteria are supplied with substrate, which finally  
18 increases respiration (Bonilla-Findji et al., 2008; Middelboe and Lyck, 2002). This could reduce  
19 the efficiency of the biological carbon pump, i.e. the process which transforms inorganic to  
20 organic carbon, part of which is then transferred to the deep ocean (Suttle, 2007). The relative  
21 significance of viral lysis and protistan grazing can strongly vary on temporal and spatial scales  
22 (Boras et al., 2009; Fuhrman and Noble, 1995). This has also been shown for cold marine  
23 environments (Boras et al., 2010; Guixa-Boixereu et al., 2002; Steward et al., 1996; Wells and  
24 Deming, 2006).

25 In about one-third of the World Ocean, including the sub-arctic northeast Pacific, the equatorial  
26 Pacific and the Southern Ocean, phytoplankton growth is limited by available iron resulting in  
27 excess phosphorus and nitrogen (Martin and Fitzwater, 1988). In these high nutrient-low

Andrea Malits 9/10/14 2:56 P.M.

Eliminado: living

1 chlorophyll (HNLC) regions, bacterioplankton are thought to be the key player of the « microbial  
2 ferrous wheel » (Kirchman, 1996), i.e. the uptake and remineralization of iron. Bacterioplankton  
3 contain more than twice iron per carbon units than eukaryotic phytoplankton, and they can  
4 thereby store up to 50% of the biogenic iron in the HNLC ocean (Tortell et al., 1996).

5 Viral activity has a potential impact on nutrient regeneration. Typically, nutrients released as a  
6 result of viral lysis are thought to be organically complexed, which may facilitate their use by  
7 marine plankton (Poorvin et al., 2004; Rue and Bruland, 1997). Iron released by viral lysis can  
8 account for more than 10% of ambient Fe concentrations (Gobler et al., 1997) and thus  
9 potentially relieve its limitation in depleted environments. Furthermore, marine viruses may serve  
10 as nuclei for iron adsorption and precipitation, and thus represent a significant reservoir of iron in  
11 seawater (Daughney et al., 2004). Despite their key role, viruses are hardly included in iron  
12 enrichment studies. These experiments were originally stimulated by the “iron hypothesis”  
13 (Martin, 1990) which assigns iron a paramount role in controlling ocean productivity and  
14 consequently atmospheric carbon dioxide concentrations. Only two out of 13 iron fertilization  
15 experiments so far performed (Secretariat of the Convention on Biological Diversity, 2009)  
16 report on viral abundance and activity (Higgins et al., 2009; Weinbauer et al., 2009a). Both  
17 studies from the Subarctic and Southern Ocean, respectively, found that viral production was  
18 significantly enhanced after iron fertilization.

19 Above the Kerguelen Plateau in the Southern Ocean, the largest HNLC ocean, a large  
20 phytoplankton bloom occurs annually during austral summer. The continuous supply of Fe and  
21 major nutrients from below has been shown to sustain this massive bloom (Blain et al., 2007).  
22 The region off Kerguelen provides the opportunity to study natural iron fertilization in the  
23 Southern Ocean and to compare it to blooms induced by mesoscale Fe additions. Within the  
24 KEOPS1 project (Kerguelen Ocean and Plateau compared Study1, 2005-2007), we sampled the  
25 phytoplankton bloom above the Kerguelen plateau during its late successional stage (~ 3<sup>rd</sup> month)  
26 along with the surrounding HNLC waters. The aim of the present study was to assess the role of  
27 viruses within the microbial food web affected by natural Fe fertilization and to elucidate the

1 possible implications for the final destiny of organic carbon. For this purpose, we measured viral  
2 production, the fraction of infected cells (FIC), lysogeny and estimated bacterial mortality  
3 through viral lysis in the bloom and surrounding HNLC waters.  
4

## 1 **2 Material and Methods**

### 2 **2.1 Description of the study site**

3 Sampling was performed in the Indian sector of the Southern Ocean above the Kerguelen Plateau  
4 (49°-53° S and 72°-78° E) in austral summer (18 January – 13 February 2005) on board the *R/V*  
5 *Marion Dufresne* in the framework of the project KEOPS (Blain et al., 2008). We sampled a  
6 large phytoplankton bloom dominated by diatoms from its peak to its decline (Mosseri et al.,  
7 2008) (Figure 1). Satellite images dated the onset of this bloom more than two month before its  
8 first visit (Blain et al., 2007). Hydrographic conditions are described in detail in Park et al.  
9 (2008). Dissolved Fe concentrations in the surface mixed layer were low and similar on and off  
10 the plateau ( $0.09 \pm 0.03$  nM) but increased with depth above the plateau reaching a mean  
11 maximum of 0.35 nM at 500m. This strong vertical gradient in combination with physical  
12 features such as internal waves and tidal activity sustained the phytoplankton bloom above the  
13 plateau (Blain et al., 2007).

### 14 **2.2 Sampling strategy**

15 Water was collected using General Oceanics 12L Niskin bottles mounted on a rosette with a Sea  
16 Bird SBE19 plus CTD sensor for salinity, temperature and oxygen from 2-3 depths (within and  
17 below the surface mixed layer) at the following stations to cover the centre and borders of each of  
18 three transects (A, B, C): A3, A11, B1, B5, B11, C3 and C11 (Figure 1, Table 1). The stations A3  
19 and C11 were considered as the most contrasting stations and sampled repeatedly. The first  
20 sampling of station A3 (A3-1) was done during the peak of the bloom, and about two weeks later,  
21 station A3 was re-sampled at a fourth visit (A3-4) during the decline of the bloom. Station B5  
22 was situated within a new phytoplankton bloom above the Kerguelen plateau (Obermosterer et al.,  
23 2008). Station A11 was located in iron-fertilized waters. The annually occurring spring bloom  
24 developed prior to our visit, explaining the low concentrations of chl *a* at this site (Table 1).  
25 Stations B11, C3 and C11 were in HNLC waters off the Kerguelen plateau and the latter was  
26 sampled twice (Table 1).

27 Total chlorophyll *a* (Chl *a* and divinyl-Chl *a*) was measured by High Performance Liquid

1 Chromatography (HPLC, Van Heukelem and Thomas, 2001, Uitz et al., 2009).

### 2 **2.3 Enumeration of viruses and prokaryotes**

3 Subsamples (2 mL) were fixed with glutaraldehyde (0.5% final concentration), incubated for 15-  
4 30 minutes at 4°C, subsequently frozen in liquid nitrogen and stored at -80°C. Within a few days  
5 samples were thawed and viral particles and bacteria were stained with SYBR Green I  
6 (Molecular Probes) and quantified using a FACScalibur (Becton and Dickinson) flow-cytometer  
7 after dilution with TE buffer (10mM Tris, 1 mM EDTA, ph = 8). For viruses an optimized  
8 protocol by Brussaard (2004) was followed. Viruses and prokaryotes were determined in plots of  
9 90° light scatter (SSC) and green DNA fluorescence. Differences in the green fluorescence and  
10 side scatter signature in the cytometric plot allowed to separate prokaryotes with low nucleic acid  
11 content (LNA) from prokaryotes with high nucleic acid content (HNA) as previously described  
12 by Gasol et al. (1999). Similarly, different size classes of viruses were distinguished on the basis  
13 of green fluorescence. Abundances were calculated by the used flow rate. Flow-cytometric  
14 assessment of viral abundance may encompass particles other than viruses such as bacterial  
15 vesicles (Biller et al., 2014). However, since bacterial and viral parameters were related  
16 significantly (Table 4), a potential overestimation of viral abundances did probably not bias the  
17 conclusions of the study.

18 To convert bacterial abundance (BA) to biomass we used a conversion factor of 12.4 fg C cell<sup>-1</sup>  
19 for oceanic prokaryotes (Fukuda et al., 1998).

### 20 **2.3 Bacterial Production**

21 The incorporation of <sup>3</sup>H leucine into protein (Smith and Azam, 1992) was used to estimate the  
22 production of heterotrophic bacteria (BP). At each depth, 1.5 mL duplicate samples and a  
23 trichloroacetic acid (TCA)-killed control were incubated with a mixture of L-[4,5 -<sup>3</sup>H] leucine  
24 (Amersham, 160 Ci mmol<sup>-1</sup>) and nonradioactive leucine added at final concentrations of 7 nM  
25 and 13 nM for the upper 100 m, and 13 nM and 7 nM for the 100-200 m depth layer. Samples  
26 were incubated in the dark at the ambient temperature of the depth where samples were collected.  
27 The incubation time (2-3h) was tested to satisfy linear incorporation with time. We checked by

1 concentration kinetics (2.5, 5, 10, 20 and 40 nM), at 3 stations inside and outside the bloom at 5  
2 m and 175 m depths that there was no isotopic dilution. The theoretical conversion factor of 1.55  
3 kg of C mol<sup>-1</sup> was used to convert leucine incorporation rates to prokaryotic carbon production  
4 (Kirchman, 1993).

#### 5 **2.4 Viral production, the fraction of infected cells and the fraction of lysogenic** 6 **cells**

7 Lytic viral production (VP<sub>l</sub>), the fraction of infected cells (FIC), induced viral production (VP<sub>i</sub>)  
8 and the fraction of lysogenic cells (FLC) were estimated using the virus reduction approach  
9 (VRA, Weinbauer et al., 2009b; Wilhelm et al., 2002). The rationale behind VRA is to reduce  
10 viral abundance in order to stop new viral infection. Thus, the viruses produced originate from  
11 already infected cells. Briefly, bacteria from 200 mL raw seawater were concentrated using a  
12 tangential flow system with a peristaltic pump (Watson-Marlow 323) equipped with a 0.2 μm  
13 cartridge (VIVAFLOW 50). To obtain virus-free seawater, the 0.2 μm pore-size ultrafiltrate was  
14 passed through a 100kDalton cartridge (VIVAFLOW 50). The bacterial concentrates were  
15 brought up to the original volume with virus-free seawater and incubated in duplicate 50 mL  
16 Falcon tubes in the dark at ± 2°C *in situ* temperature for 24 hours. Lysogeny was estimated by  
17 adding mitomycin C (SigmaChemical Co, No. M-0503, final concentration 1 μg mL<sup>-1</sup>) to  
18 duplicate 50mL Falcon tubes in order to induce the lytic cycle in lysogens; untreated duplicate  
19 samples served as controls (Paul and Weinbauer, 2010). Subsamples (2mL) for viral and bacterial  
20 abundance from each incubation were taken immediately (t<sub>0</sub> samples) and every 3-4 hours, fixed  
21 with glutaraldehyde (0.5% final concentration), incubated for 15-30 minutes at 4°C, subsequently  
22 frozen in liquid nitrogen and stored at -80°C until enumeration using a flow-cytometer as  
23 described above. VP was calculated as

$$24 \quad VP_l = (V_2 - V_1) / (t_2 - t_1) \quad (1)$$

25 where V<sub>1</sub> and V<sub>2</sub> are viral abundances and t<sub>1</sub> and t<sub>2</sub> the elapsed time. Dividing the number of  
26 produced phages by an estimated burst size (BS, i.e. the number of phages released during the



1 lysis of a single host) yields the number of lysed cells and thus gives an estimation of FIC  
2 (Weinbauer et al., 2002). FIC was calculated as

$$3 \quad FIC = 100 \times ([V_2 - V_1] / BS / BA) \quad (2)$$

4 where BA is the bacterial abundance at  $t_0$ . The difference in phage abundance between lysogeny  
5 treatment and the control is the number of induced phages, which is divided by BS to estimate the  
6 fraction of lysogenic cells (FLC). FLC was calculated as

$$7 \quad FLC = 100 \times ([V_{MC} - V_C] / BS / BA) \quad (3)$$

8 where  $V_{MC}$  and  $V_C$  is the maximum difference in viral abundance at corresponding time points in  
9 control and mitomycin C treatments, respectively. Calculations were performed for each replicate  
10 separately.

## 11 **2.5 Contact rates**

12 The rates of contact (R, number  $\text{mL}^{-1} \text{d}^{-1}$ ) between viruses and bacteria were calculated by using  
13 the following equations (Murray and Jackson, 1992).

$$14 \quad R = Sh \times 2\pi d \times D_V \times VA \times BA, \quad (4)$$

15 where  $Sh$  is the Sherwood number [1.06 for a bacterial community with 10% motile cells;  
16 (Wilhelm et al., 1998)],  $d$  is the diameter of the target,  $VA$  and  $BA$  are the abundances of viruses  
17 and bacteria, respectively, and  $D_V$  is the diffusivity of viruses.

$$18 \quad D_V = k \times T / (3 \times \pi \times \mu \times d_V) = 5 \times 10^{-8} \text{ cm}^2 \text{ s}^{-1}, \quad (5)$$

19 where  $k$  is the Boltzmann constant ( $1.38 \times 10^{-23} \text{ J K}^{-1}$ ),  $T$  is the *in situ* temperature ( $\sim 275 \text{ K}$ ),  $\mu$  is  
20 the viscosity of water ( $\text{Pascal s}^{-1}$ ) and  $d_V$  is the diameter of the viral capsid ( $\sim 60 \text{ nm}$ ). The contact  
21 rates were divided by *in situ* bacterial abundance to estimate the number of contacts per cell on a  
22 daily basis.

## 23 **2.6 Bacterial mortality**

24 To obtain the rate of cell lysis, viral production corrected for *in situ* bacterial abundance was  
25 divided by an estimated BS following the approach of Wells and Deming (2006), i.e. dividing the  
26 number of viruses produced during the first hours of incubation by the concomitant decline of

1 bacterial abundance. The number of lysed bacteria was converted into carbon by the factor of  
2 12.4 fg C cell<sup>-1</sup> (Fukuda et al., 1998). The fraction of bacterial mortality through viral lysis  
3 (VMM) was calculated following the Model by Binder (1999).

$$4 \quad VMM = FIC/LN(2) \times (1 - 0.186 - FIC) \quad (6)$$

## 5 **2.7 Statistics**

6 Normal distribution of data was checked using the Shapiro-Wilk *W*-test. Differences between  
7 different trophic situations were analyzed by the Kruskal-Wallis test for non parametric data and  
8 by one-way ANOVA for normally distributed data. Spearman rank correlation for non normally  
9 distributed data was applied. Significance was considered for  $P < 0.05$ .

10

## 1 **3 Results**

### 2 **3.1 Bacterial and viral abundances**

3 From surface water down to 200m, BA was on average 1.7 fold higher within the Fe-fertilized  
4 ( $3.9 \times 10^5 \text{ mL}^{-1}$ ) than in HNLC waters ( $2.4 \times 10^5 \text{ mL}^{-1}$ , Kruskal-Wallis test,  $P < 0.0001$ , Table 2;  
5 Figure 2). Similarly, viral abundance (VA) averaged  $9.9 \times 10^6 \text{ mL}^{-1}$  at the Fe-fertilized stations  
6 and was twice as high than in the HNLC environments ( $4.7 \times 10^6 \text{ particles mL}^{-1}$ , Kruskal-Wallis  
7 test,  $P < 0.05$ , Table 2). VA ranged from  $3.1 - 14.2 \times 10^6 \text{ mL}^{-1}$  with the highest values found at  
8 the main bloom station A3 and the lowest value detected in the deep layer of the HNLC station  
9 B11. Viruses were homogeneously distributed with depth at the HNLC stations. The virus to  
10 bacteria ratio (VBR) ranged from 11 to 34 and averaged 21 without significant differences  
11 between stations or trophic situations.

### 12 **3.2 Contact rates**

13 Contact rates were significantly higher at the Fe-fertilized stations than in HNLC waters  
14 (Kruskal-Wallis test,  $P < 0.05$ , Table 2). At the Fe-fertilized stations, on average  $29.4 \pm 11.1$   
15 viruses contacted a bacterial cell per day while in the HNLC waters contact rates were  $14.2 \pm 4.4$   
16 viruses  $\text{cell}^{-1} \text{ d}^{-1}$  with the highest values at the bloom station A3 and the lowest at the HNLC  
17 station B11 in accordance to the highest and lowest viral abundances, respectively (see Figure 2).

### 18 **3.3 Bacterial production, viral production, fraction of infected cells, lysogeny**

19 Bacterial production ranged from  $0.1 - 0.7 \mu\text{gC L}^{-1} \text{ d}^{-1}$  at the HNLC stations and from  $0.1 - 2.5$   
20  $\mu\text{gC L}^{-1} \text{ d}^{-1}$  at the Fe-fertilized stations (Table 2). The highest values were found throughout the  
21 depth profile of the main bloom station A3-1 and the lowest values were measured between 150  
22 and 200m at the HNLC stations. Despite the wide range of values, BP was on average four times  
23 higher at the Fe-fertilized stations than at the HNLC stations (Kruskal-Wallis test,  $P < 0.0001$ ,  
24 Table 2).

1 Initial virus abundance in the VRA was  $45 \pm 25\%$  (11 – 88%) of *in situ* abundance. Pre-filtration  
2 through  $0.2\mu\text{m}$  and 100kDa cartridge to obtain virus-free water and bacterial concentrate and  
3 adding back resulted on average in recovery efficiencies of  $26\% \pm 18\%$  (5-83%).  
4 Lytic viral production ( $\text{VP}_l$ ), corrected for *in situ* bacterial abundance averaged  $59.0 \times 10^6 \text{ mL}^{-1}\text{d}^{-1}$   
5 in the naturally Fe-fertilized patch compared to  $14.5 \times 10^6 \text{ mL}^{-1} \text{ d}^{-1}$  in the HNLC environments.  
6 This 4.1 fold difference was significant (Kruskal-Wallis test,  $P < 0.05$ , Table 2). Induced viral  
7 production ( $\text{VP}_i$ ) was detected in 4 out of 9 stations (3 fertilized and 1 HNLC stations, Table 3)  
8 and averaged  $44.8 \pm 44.2 \times 10^6 \text{ mL}^{-1} \text{ d}^{-1}$  (Table 2).  $\text{VP}_i$  at the main bloom station A3 at 50m  
9 increased from the first visit ( $15.6 \times 10^6 \text{ mL}^{-1} \text{ d}^{-1}$ ) to the fourth visit ( $105.6 \times 10^6 \text{ mL}^{-1} \text{ d}^{-1}$ ) by a  
10 factor of 6.8, when the decline of the bloom was sampled. BS estimates ranged from 36 to 261  
11 with mean values of  $115 \pm 74$  in the bloom and  $139 \pm 77$  in the HNLC waters.  
12 Although FIC values at the Fe-fertilized stations almost doubled that in HNLC waters, this  
13 difference between environments was not significant (Kruskal-Wallis test, Table 2). Average  
14 values for duplicate assays ranged from 4% to 47% (average: 22%) in fertilized waters and from  
15 3% to 23% (average: 12%) in HNLC waters. Lysogenic infection of bacterioplankton could be  
16 detected only in 7 out of 15 lysogenic phage induction essays and ranged from 1-31% in  
17 fertilized waters and from 1-4% in the HNLC environment.  
18 At the fertilized stations, on average  $5.4 \pm 4.1 \times 10^5$  bacteria  $\text{mL}^{-1} \text{ d}^{-1}$  were lysed, 5 times more  
19 than at the HNLC stations ( $1.1 \pm 0.6 \times 10^5$  bacteria  $\text{mL}^{-1} \text{ d}^{-1}$ ,  $P < 0.05$ , Kruskal-Wallis test, Table 2).  
20 The resulting viral mediated loss of bacterial standing stock was on average  $44 \pm 24 \%$  per day in  
21 the HNLC waters and more than twice as high at the fertilized stations although this was not  
22 significant ( $104 \pm 76 \%$   $\text{d}^{-1}$ , Kruskal-Wallis test, Table 2). The fraction of bacterial mortality  
23 through viral lysis (VMM) following the model by Binder (1999) averaged  $72 \pm 72\%$  in the  
24 bloom and  $27 \pm 19\%$  at the HNLC sites (Kruskal-Wallis, *ns*, Table 2).

### 25 **3.4 Relation between the different parameters**

26 Spearman rank correlation coefficients  $\rho$  for chlorophyll a, viral and bacterial parameters from  
27 HNLC and bloom stations are shown in Table 4. BA and BP correlated positively throughout

1 trophic situations but only in HNLC waters, BA and BP increased with chl $a$ . In the fertilized  
2 waters VA correlated positively with BP while in HNLC waters VP increased with BA. Only in  
3 these waters VP<sub>1</sub> correlated significantly and positively with the fraction of infected cells (Table  
4 4).  
5

## 1 **4 Discussion**

2 Viruses were the dominant mortality factor of bacteria during the late stage of a phytoplankton  
3 bloom induced by natural iron fertilization in the Southern Ocean (second visit to A3) but  
4 accounted for a small part of bacterial mortality within a new bloom (station B5, Table 3).  
5 Additionally, observations from the early bloom phase showed that heterotrophic nanoflagellates  
6 | [\(HNF\)](#) dominated the loss of BP, and viruses accounted for only 10% of bacterial mortality  
7 (Christaki et al., 2014). These seasonal dynamics point to a switch from an efficient functioning  
8 of the microbial food web during the onset of the phytoplankton bloom to a microbial food web  
9 where organic carbon is mainly processed by the viral shunt. The increase in viral mediated  
10 release of dissolved organic carbon over time has important consequences for the fate of part of  
11 the photosynthetically fixed carbon and reduces its transfer to higher trophic levels and export.

12

### 13 **4.1 Comparison of viral data with high latitude marine environments**

14 Viral production rates in the present study match well the data obtained from the Australian  
15 Sector of the Southern Ocean (Evans et al., 2009) and are within the range of VP rates from an  
16 iron induced bloom in the subarctic Pacific (Higgins et al., 2009). However, our VP rates are high  
17 when compared to data from an artificial iron-fertilization experiment in the Southern Ocean  
18 (Weinbauer et al., 2009a) or those from other high latitude marine environments, i.e. the Arctic  
19 Sea (Steward et al., 1996; Boras et al., 2010) (Table 5). Differences between studies could be due  
20 to spatio-temporal variations of VP, however, it is also conceivable that differences between  
21 methods (Helton et al., 2005; Weinbauer et al., 2009a; Winget et al., 2005) have contributed to the  
22 variability of reported VP data.

23 In the present study, the burst size averaged 128 throughout the experiments. This value is high  
24 compared to two studies from the Southern Ocean where measured BS was about 40 (Strzepek et  
25 al., 2005; Weinbauer et al., 2009a) and to a study in early spring above and off the Kerguelen  
26 plateau where BS evaluated with TEM observations varied from 6 to 88 (mean $\pm$ SD, 22 $\pm$ 15,  
27 Christaki et al., 2014). These different BS could be inherent to the study regions or due to the

1 used method, i.e. estimating BS by an increase in VA and a decrease of BA in the VRA (Wells  
2 and Deming, 2006) which can result in increases of BP and thus potentially increase VP (Helton  
3 et al., 2005; Weinbauer et al., 2009a; Winget et al., 2005). However, Steward et al. found BS as  
4 high as 270 for areas of high productivity in the Chukchi Sea and studies from the North Sea  
5 have reported 100 phages produced per lysed bacterium (Bratbak et al., 1992).

#### 6 7 **4.2 Viruses in HNLC waters versus a phytoplankton bloom induced by natural** 8 **iron fertilization**

9 Viral distribution during the late stage of the phytoplankton bloom above the Kerguelen plateau  
10 and its relation to the bacterial hosts (e.g. VBR) and phytoplankton biomass is extensively  
11 reported, discussed and compared to existing data from similar regions in Brussard et al. (2008b).  
12 During the late bloom average viral abundance at the bloom stations was twice as high as in  
13 HNLC waters (Brussaard et al., 2008b), while during the early bloom viral abundance remained  
14 unaffected (Christaki et al., 2014). Data from mesoscale Fe fertilization experiments showed that  
15 viral abundance inside the fertilized patch were higher (Weinbauer et al., 2009a) or not  
16 substantially different from outside (Higgins et al., 2009). The authors of the latter study  
17 explained the lack of differences between inside and outside the fertilized patch with the time lag  
18 of the microbial response to the induced bloom, since viral abundance and production were only  
19 increasing at the end of their observations (day 12 after iron fertilization). This observation is in  
20 line with the increase in viral abundance and activity on a seasonal scale in the Kerguelen bloom  
21 (Christaki et al., 2014).

22 The present study observed a mature bloom and could thus track a period with a more  
23 pronounced microbial response. The 4 times higher viral production at the naturally Fe-fertilized  
24 study sites compares well to the 3 fold increase in phage production after an induced bloom  
25 through iron addition (Weinbauer et al., 2009a). Interestingly, Christaki et al. (2014) reported  
26 higher VP rates already at the early bloom stages. Thus, there is a trend of higher viral production  
27 in the iron-fertilized bloom compared to the surrounding HNLC waters consistent with existing

1 | data on iron fertilization (Weinbauer et al., 2009a). Complementary, within the bloom, HNE did  
2 | not seem to control enhanced bacterial production rates while in HNLC waters heterotrophic  
3 | HNE consumed 95% of bacterial production (Christaki et al., 2008). These studies suggest that  
4 | there is a switch towards viral lysis dominating in the bloom situations. More generally, this is in  
5 | accordance with previous studies across environments which showed viral influence to be more  
6 | important in more eutrophic waters (Weinbauer et al., 1993;Steward et al., 1996), particularly in  
7 | the cold environments such as the Arctic (Steward et al., 1996) or the Southern Ocean, where  
8 | Guixa-Boixereu et al. (2002) found that viruses were responsible for the entire bacterial  
9 | mortality. The high viral induced mortality in the bloom could also be a reason for low biomass  
10 | accumulation, despite the high BP. We calculated carbon release rates through viral lysis in two  
11 | ways; first, based on VP, and second, based on VMM related to FIC by a Model of Binder (1999)  
12 | (Table 6). Independent of the absolute values, which were one order of magnitude higher in the  
13 | former than in the latter way (Table 6), C-release through viral lysis was 5-8 times higher in the  
14 | Fe-fertilized than in the surrounding HNLC waters.

15 | The percentages of lysogens (i.e. bacteria containing temperate viruses) were more variable in the  
16 | fertilized (0-31 %) than in the HNLC waters (0-4%) but not significantly different between  
17 | environments. Consistent with our study, Weinbauer et al. (2009a) did not find differences inside  
18 | and outside the iron-enriched patch during a fertilization experiment in the Southern Ocean. The  
19 | proportion of the lysogenized bacterial population can vary extensively, for example, from 1.5 to  
20 | 11.4% in the Gulf of Mexico (Weinbauer and Suttle, 1996), from 4 to 38% in the Canadian  
21 | Arctic Shelf (Payet and Suttle, 2013) and from 0 to 100% in Tampa Bay, Florida (Williamson et  
22 | al., 2002). According to conceptual models, lysogeny should occur in environments, where the  
23 | contact rate between infective phage and hosts is too low to sustain the lytic life style (Paul et al.,  
24 | 2002), e.g. in deep sea where host abundance was low and lysogeny was highest in an across  
25 | system study (Weinbauer et al., 2003) or when system productivity is low (Payet and Suttle,  
26 | 2013). Apparently, this was not the case in the present study as the fraction of lysogenic cells was  
27 | not different between trophic situations. It was suggested that enhanced growth causes temperate

Andrea Malits 10/6/14 3:17 P.M.

**Eliminado:** heterotrophic flagellates

Andrea Malits 10/6/14 3:17 P.M.

**Eliminado:** nanoflagellates



1 viruses to enter the lytic cycle (Wilson and Mann 1997). Both, filtration and incubation could  
2 have stimulated bacterial production in the virus reduction approach (Weinbauer et al., 2009a)  
3 and consequently induced prophages in the mitomycin C treatment controls. Additionally, it has  
4 to be stressed that mitomycin C used as an inducing agent of lysogens in the natural bacterial  
5 communities may not induce all prophages and be toxic to some bacteria (Paul, 2008;Paul and  
6 Weinbauer, 2010). Thus, the apparent low incidence of lysogenic infection, particularly in HNLC  
7 waters might be an artifact. However, it could also be that the study period was not long enough  
8 to induce potential changes of lysogenic infection. In addition, our study provides no evidence  
9 that lysogens were induced by relieving iron addition.

#### 11 **4.3 Role of viruses for sustaining phytoplankton productivity by Fe-supply**

12 Bacteria store about 50% of the biogenic iron in HNLC areas (Tortell et al., 1996) and the mode  
13 of bacterial mortality will affect the way of Fe regeneration and bioavailability (Kirchman,  
14 1996;Mioni et al., 2005;Strzepek et al., 2005). While viral lysis liberates organically complexed  
15 iron, which may be assimilated rapidly, grazing mainly sets free inorganic Fe (Gobler et al.,  
16 1997;Poorvin et al., 2004). Assimilation studies with a model heterotrophic bacterium  
17 demonstrated that Fe in the virus mediated cell lysates was more bioavailable than the  
18 siderophores produced by the same cells supporting the importance of virus-mediated Fe  
19 regeneration in marine surface waters (Poorvin et al., 2011). We calculated Fe release rates in  
20 two ways; first, based on VP, and second, based on VMM related to FIC by a Model of Binder  
21 (1999). The former resulted in average iron regeneration rates due to viral lysis of bacteria of  
22 4.18 and 0.86 pMol Fe d<sup>-1</sup> in fertilized and HNLC waters, respectively, while the latter resulted in  
23 more realistic values ranging from 0.03 pM d<sup>-1</sup> to 1.58 pM d<sup>-1</sup> (average: 0.42 ± 0.49 pM d<sup>-1</sup>) in  
24 iron-fertilized and from 0.004 pM d<sup>-1</sup> to 0.12 pM d<sup>-1</sup> (average: 0.05 ± 0.05 pM d<sup>-1</sup>) in HNLC  
25 waters (Table 6). These values are similar to those found in the Southern Ocean (Evans and  
26 Brussaard, 2012) and an iron-induced bloom (ibid., Weinbauer et al., 2009a) but low compared to  
27 other studies. Poorvin et al. (2004) reported Fe regeneration rates of 19.2-75.5 pM d<sup>-1</sup> in HNLC

1 waters off Peru, and Strzepek et al. (2005) found a high range over two orders of magnitude of  
2  $0.4\text{--}28 \text{ pM d}^{-1}$  in HNLC waters SE of New Zealand. Fe regeneration rates are calculated from  
3 viral induced bacterial loss, which is inversely related to burst size. When taking into account that  
4 the calculated burst size in the present study was five times higher than the assumed BS in the  
5 study of Poorvin et al. (2004), the values in the present study compare well to data on Fe  
6 regeneration through viral activity from artificial fertilization experiments and other  
7 environments.

8 Significantly more iron was released by viral lysis within the naturally Fe-fertilized bloom than at  
9 the HNLC stations ( $P < 0.05$ , Kruskal-Wallis, Table 6). The concentration of dissolved iron in  
10 the surface mixed layer on and off the Kerguelen plateau were typical for the open Southern  
11 Ocean and averaged  $90 \pm 34 \text{ pM}$  (Blain et al., 2007) and the estimated biogenic iron pool at the  
12 main bloom station equaled  $80 \pm 9 \text{ pM}$  (Sarhou et al., 2008). Taking into account the total Fe  
13 demand of the producers within the bloom of  $6.04 \pm 0.62 \text{ pM d}^{-1}$  (Sarhou et al., 2008), the  
14 remobilization of iron through viral lysis above the Kerguelen plateau following the Model by  
15 Binder (1999) accounts for up to 26% of the demand of the producers and this appears to be a non-  
16 negligible iron source for sustaining plankton productivity.

17

#### 18 **4.4 Implications for carbon cycling and sequestration**

19 Enrichment of bacterial biomass and production in the naturally fertilized bloom in the present  
20 study ranged from  $287$  to  $797 \text{ mg C m}^{-2}$  and from  $23.5 \text{ mg C m}^{-2} \text{ d}^{-1}$  to  $304 \text{ mg C m}^{-2} \text{ d}^{-1}$ ,  
21 respectively (Christaki et al., 2008). Bacterial abundance and production are often correlated with  
22 viral abundance and production. Thus, elevated bacterial activity in the (natural or induced)  
23 bloom could explain the enhanced viral abundance and production found in previous *in situ* Fe  
24 enrichment studies (Arrieta et al., 2000; Higgins et al., 2009; Weinbauer et al., 2009a).

25 The finding of higher viral lysis rates of bacteria in the sites of natural Fe fertilization, where  
26 HNF grazing could only explain a small fraction of bacterial mortality (Christaki et al., 2008) has  
27 important implications for the carbon cycling. Due to enhanced viral lysis, less carbon will be

Andrea Malits 10/6/14 3:19 P.M.

Eliminado: protozoan

1 transferred to larger members of the food web but becomes again part of the DOM pool  
2 (Middelboe et al., 1996). This viral shunt should result in elevated bacterial production and  
3 respiration, thus more CO<sub>2</sub> would be produced and less carbon sequestered. Experimental studies  
4 indicate that most of the lysis products belong to the labile fraction of DOM and are consequently  
5 rapidly degraded (Weinbauer et al., 2011). By the transformation of bacterial biomass into DOM,  
6 viruses have the effect of retaining carbon and nutrients in the photic zone (Suttle, 2007). Thus,  
7 viral lysis of bacteria could short-circuit the biological pump (Brussaard et al., 2008a).

8 However, there are other possible scenarios. For example, microbial activity converts part of the  
9 organic matter into recalcitrant DOM (RDOM) that is resistant to microbial utilization and can  
10 persist in the interior of oceans for up to thousand of years. The detailed role of viral lysis in this  
11 new concept of the microbial carbon pump (MCP) (Jiao et al., 2010) is still poorly known.  
12 However, a compilation of data suggests that viral lysis increases the DOM pool and the ratio of  
13 recalcitrant vs labile organic matter (Weinbauer et al., 2011). Thus, enhanced viral lysis of  
14 bacteria due to Fe fertilization could result in an enhanced carbon sequestration not related to the  
15 biological pump.

16 Rates of bacterial production ([<sup>3</sup>H] leucine incorporation) and respiration (< 0.8 μm size-fraction)  
17 were 5-6 times higher in the bloom at Station A3 than those in surrounding HNLC waters  
18 indicating that heterotrophic bacteria within the bloom processed a significant portion of primary  
19 production with most of it being rapidly respired (Obernosterer et al., 2008) fueling the CO<sub>2</sub> pool.  
20 This scenario is coherent with the finding of small particulate organic carbon export fluxes to  
21 depth necessary for long-term sequestration (de Baar et al., 2005; Street and Paytan, 2005),  
22 despite the role of iron in regulating primary productivity. However, most *in situ* mesoscale iron  
23 enrichment experiments so far performed in the HNLC regions did not last long enough to follow  
24 the termination of the bloom (Buesseler and Boyd, 2003; Smetacek et al., 2012). In the present  
25 study, we sampled a bloom in its late successional stage and could thereby track the fate of fixed  
26 carbon by an iron-fertilized phytoplankton bloom. Figure 4 shows a simple sketch to highlight  
27 the importance of each compartment of the microbial food web in the transfer of organic material

1 in an Fe-fertilized bloom compared to HNLC waters. Sequestration of material in viruses,  
2 bacteria and dissolved matter may lead to stronger retention of nutrients in the euphotic zone in  
3 systems with high viral lysis rates of bacteria, because more material remains in these small non-  
4 sinking forms. This could be of major importance for large-scale iron fertilization of ocean  
5 regions as a means of enhancing the ability of the ocean to store anthropogenic CO<sub>2</sub> and mitigate  
6 21<sup>st</sup> century climate change.

## 7 **5 Conclusions**

8 Enhanced bacterial production following the iron-fertilized phytoplankton bloom induced a  
9 switch from grazing to viral lysis as major mechanisms causing bacterial mortality. This could  
10 change the carbon flow through the microbial food web. We suggest that enhanced viral lysis of  
11 bacteria short-circuits the biological pump but potentially prime the microbial carbon pump.

12

## 13 **Acknowledgements**

14 We thank the chief scientists (S. Blain and B. Quéguiner) for the possibility to participate in this  
15 cruise, the captain and crew of R.V. Marion Dufresne for their efficient assistance during work at  
16 sea and the colleagues for help on board. The financial support was provided by the European  
17 Union in the framework of the BASICS project (EVK3-CT-2002-00078), by the French Research  
18 program of the INSU-CNRS PROOF, the French Polar Institute (IPEV) and a spanish grant from  
19 the ministry of education (SB2010-0079) to A.M.

## 20 **References**

1 Arrieta, J. M., Weinbauer, M. G., and Herndl, G. J.: Interspecific variability in sensitivity to uv  
2 radiation and subsequent recovery in selected isolates of marine bacteria, *Appl Environ*  
3 *Microbiol*, 66, 1468-1473, 2000.

4 Bergh, O., Børshheim, K. Y., Bratbak, G., and Heldal, M.: High abundance of viruses found in  
5 aquatic environments, *Nature*, 340, 467-468, 1989.

6 Biller, S. J., Schubotz, F., Roggensack, S. E., Thompson, A. W., Summons, R. E., and Chisholm,  
7 S. W.: Bacterial vesicles in marine ecosystems, *Science*, 343, 183-186, 343/6167/183 [pii]  
8 10.1126/science.1243457, 2014.

9 Binder, B.: Reconsidering the relationship between virally induced bacterial mortality and  
10 frequency of infected cells, *Aquat. Microb. Ecol.*, 18, 207-215, 1999.

11 Bird, D. F., Maranger, R., and Karl, D.: Palmer lter: Aquatic virus abundances near the antarctic  
12 peninsula, *Antarct. J. U.S.*, 28, 234-235, 1993.

13 Blain, S., Queguiner, B., Armand, L., Belviso, S., Bombled, B., Bopp, L., Bowie, A., Brunet, C.,  
14 Brussaard, C., Carlotti, F., Christaki, U., Corbiere, A., Durand, I., Ebersbach, F., Fuda, J. L.,  
15 Garcia, N., Gerringa, L., Griffiths, B., Guigue, C., Guillerm, C., Jacquet, S., Jeandel, C., Laan, P.,  
16 Lefevre, D., Lo Monaco, C., Malits, A., Mosseri, J., Obernosterer, I., Park, Y. H., Picheral, M.,  
17 Pondaven, P., Remenyi, T., Sandroni, V., Sarthou, G., Savoye, N., Scouarnec, L., Souhaut, M.,  
18 Thuiller, D., Timmermans, K., Trull, T., Uitz, J., van Beek, P., Veldhuis, M., Vincent, D.,  
19 Viollier, E., Vong, L., and Wagener, T.: Effect of natural iron fertilization on carbon  
20 sequestration in the southern ocean, *Nature*, 446, 1070-1074, 2007.

21 Blain, S., Queguiner, B., and Trull, T.: The natural iron fertilization experiment keeps (kerguelen  
22 ocean and plateau compared study): An overview, *Deep Sea Res. Part II: Topical Studies in*  
23 *Oceanography*, 55, 559-565, 2008.

24 Bonilla-Findji, O., Malits, A., Lefevre, D., Rochelle-Newall, E., Lemeë, R., Weinbauer, M. G.,  
25 and Gattuso, J.-P.: Viral effects on bacterial respiration, production and growth efficiency:  
26 Consistent trends in the southern ocean and the mediterranean sea, *Deep Sea Res. Part II: Topical*  
27 *Studies in Oceanography*, 55, 790-800, 2008.

28 Boras, J., Sala, M., Arrieta, J., Sà, E., Felipe, J., Agustí, S., Duarte, C., and Vaqué, D.: Effect of  
29 ice melting on bacterial carbon fluxes channelled by viruses and protists in the arctic ocean, *Polar*  
30 *Biol.*, 33, 1695-1707-1707, 10.1007/s00300-010-0798-8, 2010.

31 Boras, J. A., Sala, M. M., Vazquez-Dominguez, E., Weinbauer, M. G., and Vaque, D.: Annual  
32 changes of bacterial mortality due to viruses and protists in an oligotrophic coastal environment  
33 (nw mediterranean), *Environ Microbiol*, 11, 1181-1193, EMI1849 [pii] 10.1111/j.1462-  
34 2920.2008.01849.x, 2009.

35 Bratbak, G., Heldal, M., Thingstad, T. F., Riemann, B., and Haslund, O. H.: Incorporation of  
36 viruses into the budget of microbial c-transfer. A first approach, *Mar. Ecol. Prog. Ser.*, 83, 273-  
37 280, 1992.

38 Brussaard, C. P. D., Wilhelm, S. W., Thingstad, F., Weinbauer, M. G., Bratbak, G., Heldal, M.,  
39 Kimmance, S. A., Middelboe, M., Nagasaki, K., Paul, J. H., Schroeder, D. C., Suttle, C. A.,  
40 Vaque, D., and Wommack, K. E.: Global-scale processes with a nanoscale drive: The role of  
41 marine viruses, *Isme J*, 2, 575-578, ismej200831 [pii] 10.1038/ismej.2008.31, 2008a.

42 Brussaard, C. P. D.: Optimization of procedures for counting viruses by flow cytometry, *Appl.*  
43 *Environ. Microbiol.*, 70, 1506-1513, 2004.

44 Brussaard, C. P. D., Timmermans, K. R., Uitz, J., and Veldhuis, M. J. W.: Virioplankton  
45 dynamics and virally induced phytoplankton lysis versus microzooplankton grazing southeast of  
46 the kerguelen (southern ocean), *Deep Sea Res. II*, 55, 752-765, 2008b.

- 1 Buesseler, K. O., and Boyd, P. W.: Climate change. Will ocean fertilization work?, *Science*, 300,  
2 67-68, 2003.
- 3 Christaki, U., Obernosterer, I., Van Wambeke, F., Veldhuis, M., Garcia, N., and Catala, P.:  
4 Microbial food web structure in a naturally iron fertilized area in the southern ocean (kerguelen  
5 plateau), *Deep Sea Res. II*, 55, 706-719, 2008.
- 6 Christaki, U., Lefèvre, D., Georges, C., Colombet, J., Catala, P., Courties, C., Sime-Ngando, T.,  
7 Blain, S., and Obernosterer, I.: Microbial food web dynamics during spring phytoplankton  
8 blooms in the naturally iron-fertilized kerguelen area (southern ocean), *Biogeosciences Discuss.*,  
9 11, 6985-7028, 10.5194/bgd-11-6985-2014, 2014.
- 10 Daughney, C. J., Chatellier, X., Chan, A., Kenward, P., Fortin, D., Suttle, C. A., and Fowle, D.  
11 A.: Adsorption and precipitation of iron from seawater on a marine bacteriophage (pwh3a-p1),  
12 *Mar. Chem.*, 91, 101-115, 2004.
- 13 de Baar, H. J. W., Boyd, P. W., Coale, K. H., Landry, M. R., Tsuda, A., ASSMY, P., Bakker, D.  
14 C. E., Bozec, Y., Barber, R. T., Brzezinski, M. A., Buesseler, K. O., Boye, M., Croot, P. L.,  
15 Gervais, F., Gorbunov, M. Y., Harrison, P. J., Hiscock, M. R., Laan, P., Lancelot, C., Law, C. S.,  
16 Levasseur, M., Marchetti, A., Millero, F. J., Nishioka, J., Nojiri, Y., van Oijen, T., Riebesell, U.,  
17 Rijkenberg, M. J. A., Saito, H., Takeda, S., Timmermans, K. L., Veldhuis, M. J. W., Waite, A.  
18 M., and Wong, C. S.: Synthesis of iron fertilization experiments: From the iron age in the age of  
19 enlightenment, *J. Geophys. Res.*, 110, C09S16, doi:10.1029/2004JC002601, 2005.
- 20 Evans, C., Pearce, I., and Brussaard, C. P. D.: Viral-mediated lysis of microbes and carbon  
21 release in the sub-antarctic and polar frontal zones of the Australian southern ocean, *Environ.*  
22 *Microbiol.*, 11, 2924-2934, 2009.
- 23 Evans, C., and Brussaard, C. P. D.: Regional variation in lytic and lysogenic viral infection in the  
24 southern ocean and its contribution to biogeochemical cycling, *Appl. Environ. Microbiol.*, 78,  
25 6741-6748, 10.1128/aem.01388-12, 2012.
- 26 Fuhrman, J. A., and Noble, R. T.: Viruses and protists cause similar bacterial mortality in coastal  
27 seawater, *Limnol. Oceanogr.*, 40, 1236-1242, 1995.
- 28 Fuhrman, J. A.: Marine viruses and their biogeochemical and ecological effects, *Nature*, 399,  
29 541-548, 1999.
- 30 Fukuda, R., Ogawa, H., Nagata, T., and Koike, I. I.: Direct determination of carbon and nitrogen  
31 contents of natural bacterial assemblages in marine environments, *Appl Environ Microbiol*, 64,  
32 3352-3358, 1998.
- 33 Gasol, J. M., Zweifel, U. L., Peters, F., Fuhrman, J. A., and Hagstrom, A.: Significance of size  
34 and nucleic acid content heterogeneity as measured by flow cytometry in natural planktonic  
35 bacteria, *Appl Environ Microbiol*, 65, 4475-4483, 1999.
- 36 Gobler, C. J., Hutchins, D. A., Fisher, N. S., Coper, E. M., and Sanudo-Wilhelmy, S. A.:  
37 Release and bioavailability of c, n, p, se., and fe following viral lysis of a marine chrysophyte,  
38 *Limnol. Oceanogr.*, 42, 1492-1504, 1997.
- 39 Guixa-Boixereu, N., Vaqué, D., Gasol, J. M., Sánchez-Cámara, J., and Pedrós-Alió, C.: Viral  
40 distribution and activity in antarctic waters, *Deep-Sea Res.*, 49, 827-845, 2002.
- 41 Higgins, J. L., Kudo, I., Nishioka, J., Tsuda, A., and Wilhelm, S. W.: The response of the virus  
42 community to the seeds ii mesoscale iron fertilization, *Deep Sea Res. Part II: Topical Studies in*  
43 *Oceanography SEEDS II: The Second Subarctic Pacific Iron Experiment for Ecosystem*  
44 *Dynamics Study*, 56, 2788-2795, 2009.
- 45 Jiao, N., Herndl, G. J., Hansell, D. A., Benner, R., Kattner, G., Wilhelm, S. W., Kirchman, D. L.,  
46 Weinbauer, M., G., Luo, T., Chen, F., and Azam, F.: Microbial production of recalcitrant

1 dissolved organic matter: Long-term carbon storage in the global ocean, *Nature Reviews*  
2 *Microbiology*, 8, 593-599, 2010.

3 Kirchman, D. L.: Leucine incorporation as a measure of biomass production by heterotrophic  
4 bacteria, in: *Handbook of methods in aquatic microbial ecology*, edited by: Kemp, P. F., B.F.  
5 Sherr, E.B. Sherr and J.J. Cole, Lewis Publishers, Boca Raton, FL, 509-512, 1993.

6 Kirchman, D. L.: Microbial ferrous wheel, *Nature*, 383, 303-304, 1996.

7 Manganelli, M., Malfatti, F., Samo, T. J., Mitchell, B. G., Wang, H., and Azam, F.: Major role of  
8 microbes in carbon fluxes during austral winter in the southern drake passage, *PLoS ONE*, 4,  
9 e6941, doi:10.1371/journal.pone.0006941, 2009.

10 Marchant, H., Davidson, A., Wright, S., and Glazebrook, J.: The distribution and abundance of  
11 viruses in the southern ocean during spring, *Antarct. Sci.*, 12, 414-417, 2000.

12 Martin, J. H., and Fitzwater, S. E.: Iron deficiency limits phytoplankton growth in the north-east  
13 pacific subarctic, *Nature*, 331, 341-343, 1988.

14 Martin, J. H.: Glacial interglacial co<sub>2</sub> change: The iron hypothesis, *Paleoceanography*, 5, 1-13,  
15 1990.

16 Middelboe, M., Jorgensen, N., and Kroer, N.: Effects of viruses on nutrient turnover and growth  
17 efficiency of noninfected marine bacterioplankton, *Appl. Environ. Microbiol.*, 62, 1991-1997,  
18 1996.

19 Middelboe, M., and Lyck, P. G.: Regeneration of dissolved organic matter by viral lysis in  
20 marine microbial communities, *Aquat. Microb. Ecol.*, 27, 187-194, 2002.

21 Mioni, C. E., Poorvin, L., and Wilhelm, S. W.: Virus and siderophore-mediated transfer of  
22 available Fe between heterotrophic bacteria: Characterization using an Fe-specific bioreporter,  
23 *Aquat. Microb. Ecol.*, 41, 233-245, 2005.

24 Mosseri, J., Quéguiner, B., Armand, L., and Cornet-Barthaux, V.: Impact of iron on silicon  
25 utilization by diatoms in the southern ocean: A case study of Si/N cycle decoupling in a naturally  
26 iron-enriched area, *Deep Sea Res. Part II: Topical Studies in Oceanography*, 55, 801-819, 2008.

27 Murray, A. G., and Jackson, A. G.: Viral dynamics: A model of the effects of size, shape, motion  
28 and abundance of single-celled planktonic organisms and other particles, *Mar. Ecol. Prog. Ser.*,  
29 89, 103-116, 1992.

30 Obernosterer, I., Christaki, U., Lefevre, D., Catala, P., Van Wambeke, F., and Lebaron, P.: Rapid  
31 bacterial mineralization of organic carbon produced during a phytoplankton bloom induced by  
32 natural iron fertilization in the southern ocean, *Deep Sea Res. II*, 55, 777-789, 2008.

33 Park, Y.-H., Roquet, F., Durand, I., and Fuda, J.-L.: Large-scale circulation over and around the  
34 northern kerguelen plateau, *Deep Sea Res. Part II: Topical Studies in Oceanography*, 55, 566-  
35 581, <http://dx.doi.org/10.1016/j.dsr2.2007.12.030>, 2008.

36 Paul, J. H., Sullivan, M. B., Segall, A. M., and Rohwer, F.: Marine phage genomics, *Comp*  
37 *Biochem Physiol B Biochem Mol Biol*, 133, 463-476, 2002.

38 Paul, J. H.: Prophages in marine bacteria: Dangerous molecular time bombs or the key to survival  
39 in the seas?, *Isme J*, 2, 579-589, 2008.

40 Paul, J. H., and Weinbauer, M. G.: Detection of lysogeny in marine environments, in: *Manual of*  
41 *aquatic viral ecology*, edited by: Suttle, C., Wilhelm, S. W., and Weinbauer, M. G., ASLO, 1-8,  
42 2010.

43 Payet, J. P., and Suttle, C. A.: Physical and biological correlates of virus dynamics in the  
44 southern beaufort sea and amundsen gulf, *J. Mar. Syst. Sea ice and life in a river-influenced*  
45 *arctic shelf ecosystem*, 74, 933-945, 2008.

- 1 Payet, J. P., and Suttle, C. A.: To kill or not to kill: The balance between lytic and lysogenic viral  
2 infection is driven by trophic status, *Limnol. Oceanogr.*, 58, 465-474, 10.4319/lo.2013.58.2.0465,  
3 2013.
- 4 Poorvin, L., Rinta-Kanto, J. M., Hutchins, D. A., and Wilhelm, S. W.: Viral release of iron and  
5 its bioavailability to marine plankton, *Limnol. Oceanogr.*, 49, 1734-1741, 2004.
- 6 Poorvin, L., Sander, S. G., Velasquez, I., Ibanami, E., LeCleir, G. R., and Wilhelm, S. W.: A  
7 comparison of Fe bioavailability and binding of a catechol siderophore with virus-mediated  
8 lysates from the marine bacterium *Vibrio alginolyticus* pwh3a, *J. Exp. Mar. Biol. Ecol.*, 399, 43-  
9 47, 2011.
- 10 Proctor, L. M., and Fuhrman, J. A.: Viral mortality of marine bacteria and cyanobacteria, *Nature*,  
11 343, 60-62, 1990.
- 12 Rue, E. L., and Bruland, K. W.: The role of organic complexation on ambient iron chemistry in  
13 the equatorial Pacific ocean and the response of a mesoscale iron addition experiment, *Limnol.*  
14 *Oceanogr.*, 43, 901-910, 1997.
- 15 Sarthou, G., Vincent, D., Christaki, U., Obernosterer, I., Timmermans, K. R., and Brussaard, C.  
16 P. D.: The fate of biogenic iron during a phytoplankton bloom induced by natural fertilization:  
17 Impact of copepod grazing, *Deep Sea Res. II*, 55, 734-751, 2008.
- 18 Smetacek, V., Klaas, C., Strass, V. H., Assmy, P., Montresor, M., Cisewski, B., Savoye, N.,  
19 Webb, A., d'Ovidio, F., Arrieta, J. M., Bathmann, U., Bellerby, R., Berg, G. M., Croot, P.,  
20 Gonzalez, S., Henjes, J., Herndl, G. J., Hoffmann, L. J., Leach, H., Losch, M., Mills, M. M.,  
21 Neill, C., Peeken, I., Rottgers, R., Sachs, O., Sauter, E., Schmidt, M. M., Schwarz, J.,  
22 Terbruggen, A., and Wolf-Gladrow, D.: Deep carbon export from a southern ocean iron-fertilized  
23 diatom bloom, *Nature*, 487, 313-319,  
24 [http://www.nature.com/nature/journal/v487/n7407/abs/nature11229.html#supplementary-](http://www.nature.com/nature/journal/v487/n7407/abs/nature11229.html#supplementary-information)  
25 [information](http://www.nature.com/nature/journal/v487/n7407/abs/nature11229.html#supplementary-information), 2012.
- 26 Smith, D. C., and Azam, F.: A simple, economical method for measuring bacterial protein  
27 synthesis rates in seawater using <sup>3</sup>H-leucine, *Mar. Microb. Food Webs*, 6, 107-114, 1992.
- 28 Smith, D. C., Steward, G. F., and Azam, F.: Virus and bacteria abundances in the Drake passage  
29 during January and August 1991, *Antarct. J. U.S.*, 27, 125-127, 1992.
- 30 Steward, G. F., Smith, D. C., and Azam, F.: Abundance and production of bacteria and viruses in  
31 the Bering and Chukchi seas, *Mar. Ecol. Prog. Ser.*, 131, 287-300, 1996.
- 32 Street, J. H., and Paytan, A.: Iron, phytoplankton growth, and the carbon cycle, *Met Ions Biol*  
33 *Syst*, 43, 153-193, 2005.
- 34 Strzepek, R. F., Maldonado, M. T., Higgins, J. L., Hall, J., Safi, K., Wilhelm, S. W., and Boyd,  
35 P.: Spinning the "ferrous wheel": The importance of the microbial community in an iron budget  
36 during the FeCycle experiment, *Global Biogeochem. Cy.*, 19, GB4S26,  
37 doi:10.1029/2005GB002490, 2005.
- 38 Suttle, C. A., Chan, A. M., and Cottrell, M. T.: Infection of phytoplankton by viruses and  
39 reduction of primary productivity, *Nature*, 347, 467-469, 1990.
- 40 Suttle, C. A.: Marine viruses--major players in the global ecosystem, *Nat Rev Microbiol*, 5, 801-  
41 812, 2007.
- 42 Tortell, P. D., Maldonado, M. T., and Price, N. M.: The role of heterotrophic bacteria in iron-  
43 limited ocean ecosystems, *Nature*, 383, 330-332, 1996.
- 44 Van Heukelem, L., and Thomas, C. S.: Computer-assisted high-performance liquid  
45 chromatography method development with applications to the isolation and analysis of  
46 phytoplankton pigments, *J. Chromatogr. A*, 910, 31-49, 2001.



1 Weinbauer, G., Chen, F., and Wilhelm, A. W.: Virus-mediated redistribution and partitioning of  
2 carbon in the global oceans, in: *Microbial carbon pump*, edited by: Jiao, N., Azam, F., and  
3 Sanders, S., Science, 2011.

4 Weinbauer, M., Winter, C., and Höfle, M.: Reconsidering transmission electron microscopy  
5 based estimates of viral infection of bacterioplankton using conversion factors derived from  
6 natural communities, *Aquat. Microb Ecol*, 27, 103-110, 2002.

7 Weinbauer, M. G., Fuks, D., and Peduzzi, P.: Distribution of viruses and dissolved DNA along a  
8 coastal trophic gradient in the northern adriatic sea, *Appl. Environ. Microbiol.*, 59, 4074-4082,  
9 1993.

10 Weinbauer, M. G., and Suttle, C. A.: Potential significance of lysogeny to bacteriophage  
11 production and bacterial mortality in coastal waters of the gulf of mexico, *Appl. Environ.*  
12 *Microbiol.*, 62, 4374-4380, 1996.

13 Weinbauer, M. G., Brettar, I., and Höfle, M. G.: Lysogeny and virus-induced mortality of  
14 bacterioplankton in surface, deep, and anoxic marine waters, *Limnol. Oceanogr.*, 48, 1457-1465,  
15 2003.

16 Weinbauer, M. G., Arrieta, J. M., Griebler, C., and Herndl, G. J.: Enhanced viral production and  
17 infection of bacterioplankton during an iron induced phytoplankton bloom in the southern ocean,  
18 *Limnol. Oceanogr.*, 54, 774-784, 2009a.

19 Weinbauer, M. G., Rowe, J. M., and Wilhelm, S. W.: Determining rates of virus production in  
20 aquatic systems by the virus reduction approach, in: *Manual of aquatic viral ecology, limnology*  
21 *and oceanography methods*, edited by: Suttle, C., Wilhelm, S. W., and Weinbauer, M. G., 1-8,  
22 2009b.

23 Wells, L. E., and Deming, J. W.: Significance of bacterivory and viral lysis in bottom waters of  
24 franklin bay, canadian arctic, during winter, *Aquat. Microb. Ecol.*, 43, 209-221, 2006.

25 Wilhelm, S., Brigden, S., and Suttle, C.: A dilution technique for the direct measurement of viral  
26 production: A comparison in stratified and tidally mixed coastal waters, *Microb. Ecol.*, 43, 168-  
27 173, 2002.

28 Wilhelm, S. W., Weinbauer, M. G., Suttle, C. A., and Jeffrey, W. H.: The role of sunlight in the  
29 removal and repair of viruses in the sea, *Limnol. Oceanogr.*, 43, 586-592, 1998.

30 Wilhelm, S. W., and Suttle, C. A.: Virus and nutrient cycles in the sea, *BioScience*, 49, 781-787,  
31 1999.

32 Williamson, S. J., Houchin, L. A., McDaniel, L., and Paul, J. H.: Seasonal variation in lysogeny  
33 | as depicted by prophage induction in tampa bay, florida, *Appl. Environ. Microbiol.*, 68, 4307-  
34 4314, 2002.

35  
36  
37

1 Table 1. Date, location, mixed layer depth (Zm) and physicochemical characteristics of all  
 2 sampled stations.  
 3

Date	Station	latitude	longitude	Water type	Zm (m)	Sampling depth (m)	T ° C	Salinity	Chl <i>a</i> $\mu\text{g L}^{-1}$ *
1/19/05	A3-1	50°38'S	72°05'E	+ Fe	52	10	3.5	33.9	0.94
1/19/05	A3-1	50°38'S	72°05'E	+ Fe	52	50	3.3	33.9	1.72
1/19/05	A3-1	50°38'S	72°05'E	+ Fe	52	100	3	33.9	1.38
1/20/05	A11	49°09'S	74°00'E	+ Fe	44	10	3.8	33.9	0.41
1/20/05	A11	49°09'S	74°00'E	+ Fe	44	75	3.3	33.9	0.52
1/20/05	A11	49°09'S	74°00'E	+ Fe	44	200	1.6	34.1	0.21
1/26/05	C11-1	51°39'S	78°00'E	- Fe	73	10	1.9	33.8	0.19
1/26/05	C11-1	51°39'S	78°00'E	- Fe	73	80	1.6	33.8	0.29
1/26/05	C11-1	51°39'S	78°00'E	- Fe	73	200	1.3	34.2	0.01
1/29/05	B11	50°30'S	77°00'E	- Fe	59	10	2.2	33.8	0.11
1/29/05	B11	50°30'S	77°00'E	- Fe	59	120	0.5	33.8	0.24
1/29/05	B11	50°30'S	77°00'E	- Fe	59	200	0.2	34.1	0.03
2/1/05	B5	51°06'S	74°36'E	+ Fe	84	60	2.8	33.9	1.54
2/1/05	B5	51°06'S	74°36'E	+ Fe	84	100	2.6	33.9	1.39
2/2/05	B1	51°30'S	73°00'E	+ Fe	59	60	3.3	33.9	1.29
2/2/05	B1	51°30'S	73°00'E	+ Fe	59	100	2.7	33.9	1.04
2/4/05	A3-4	50°39'S	72°05'E	+ Fe	80	50	3.6	33.9	1.48
2/4/05	A3-4	50°39'S	72°05'E	+ Fe	80	150	1.7	33.9	1.54
2/6/05	C11-2	51°39'S	78°00'E	- Fe	20	60	1.6	33.8	0.26
2/6/05	C11-2	51°39'S	78°00'E	- Fe	20	100	0.6	33.9	0.20
2/9/05	C3	52°43'S	74°49'E	- Fe	42	60	2.5	33.9	0.19
2/9/05	C3	52°43'S	74°49'E	- Fe	42	100	1.9	33.9	0.17

4 T, temperature in ° Celsius; chl *a*, total chlorophyll *a*, +Fe, iron-fertilized, -Fe, HNLC waters  
 5 \* data are from Uitz et al. 2009  
 6

1 Table 2. Average  $\pm$  SD values of viral and bacterial parameters from the iron-fertilized and  
 2 HNLC stations in the upper 200m water layer and results from one-way ANOVA for normally  
 3 distributed data and Kruskal Wallis test for nonparametric data. Ranges are given in parenthesis.  
 4 The average ratio between the two environments is shown and significant differences are  
 5 indicated.

Parameters	Fe-fertilized stations	HNLC stations	ratio
BA mL <sup>-1</sup>	<b>3.9 <math>\pm</math> 0.9 (1.9-5.3) x 10<sup>5</sup></b>	<b>2.4 <math>\pm</math> 0.7 (1.3-3.8) x 10<sup>5</sup></b>	<b>1.7 ***</b>
BP $\mu$ gC L <sup>-1</sup> d <sup>-1</sup>	<b>1.1 <math>\pm</math> 0.7 (0.1-2.5)</b>	<b>0.3 <math>\pm</math> 0.2 (0.1-0.7)</b>	<b>4.1 ***</b>
VA mL <sup>-1</sup>	<b>9.9 <math>\pm</math> 3.6 (3.4-14.2) x 10<sup>6</sup></b>	<b>4.7 <math>\pm</math> 1.4 (3.1-7.4) x 10<sup>6</sup></b>	<b>2.1 *</b>
VP <sub>1</sub> mL <sup>-1</sup> d <sup>-1</sup>	<b>59.0 <math>\pm</math> 47.1 (9.9-117.9) x 10<sup>6</sup></b>	<b>14.5 <math>\pm</math> 7.4 (6.0-25.6) x 10<sup>6</sup></b>	<b>4.1 *</b>
VP <sub>1</sub> mL <sup>-1</sup> d <sup>-1</sup> †	50.9 $\pm$ 46.4 (2.8-125.5) x 10 <sup>6</sup>	13.9 x 10 <sup>6</sup>	3.7
FIC %	22 $\pm$ 17 (4-47)	12 $\pm$ 7 (3-23)	1.8
FLC %††	10 $\pm$ 14 (1-31)	3 $\pm$ 2 (1-4)	4.0
Prophage replication rate mL <sup>-1</sup> d <sup>-1</sup> ††	18.1 $\pm$ 29.2 (0.6-61.5) x 10 <sup>3</sup>	1.0 $\pm$ 1.2 (0.2-2.4) x 10 <sup>3</sup>	18.5
R cell <sup>-1</sup> d <sup>-1</sup>	<b>29.4 <math>\pm</math> 11.1 (10.3-43.0)</b>	<b>14.2 <math>\pm</math> 4.4 (9.3-22.4)</b>	<b>2.1 *</b>
lysed bacteria mL <sup>-1</sup> d <sup>-1</sup>	<b>5.4 <math>\pm</math> 4.1 (0.8-10.3) x 10<sup>5</sup></b>	<b>1.1 <math>\pm</math> 0.6 (0.4-2.1) x 10<sup>5</sup></b>	<b>4.9 *</b>
VMM %	72 $\pm$ 72 (8-202)	27 $\pm$ 19 (6-58)	2.6

7 VA, viral abundance, VP<sub>1</sub>, lytic viral production, VP<sub>i</sub>, induced viral production, FIC, fraction of  
 8 infected cells, FLC, fraction of lysogenic cells, BA, bacterial abundance, BP, bacterial  
 9 production, R, viral contacts per cell and day, VMM, virus-mediated bacterial mortality.

10 \* P < 0.05, \*\* P < 0.001, \*\*\* P < 0.0001

11 † detected in 6 out of 15 essays, only one in HNLC waters

12 †† detected in 7 out of 15 essays

Table 3. *In situ* BP and viral parameters from all virus reduction experiments.

Station	Water type	Depth m	BP $\mu\text{gC L}^{-1} \text{d}^{-1}$	VP <sub>1</sub> $10^6 \text{mL}^{-1} \text{d}^{-1}$	FIC %	FLC %	VMM %
A3-1	+ Fe	10	2.5	16.7	12	31	25
A3-1	+ Fe	50	1.9	15.6	6	6	12
A3-1	+ Fe	100	2.4	56.4	10	ND	19
A3-4	+ Fe	50	1.2	105.6	34	ND	106
A3-4	+ Fe	150	0.3	82.4	36	ND	115
B1	+ Fe	60	1.7	117.9	41	ND	147
B1	+ Fe	100	0.2	115.6	47	3	202
B5	+ Fe	100	1.1	11.2	4	1	8
A11	+ Fe	200	0.3	9.9	7	ND	14
B11	- Fe	10	0.2	16.3	14	ND	29
B11	- Fe	120	0.3	25.6	23	1	58
B11	- Fe	200	0.1	6.0	11	3	24
C3	- Fe	60	0.2	20.1	6	no exp	11
C3	- Fe	100	0.4	16.7	22	no exp	55
C11-1	- Fe	10	0.4	11.2	8	ND	17
C11-1	- Fe	80	0.7	9.6	9	4	17
C11-1	- Fe	200	0.1	7.5	6	ND	12
C11-2	- Fe	60	0.3	25.1	20	no exp	47
C11-2	- Fe	100	0.2	6.4	3	no exp	6

VPI, lytic viral production; FIC, fraction of infected cells; FLC, fraction of lysogenic cells; VMM, viral-mediated bacterial mortality; ND, Not detectable; no exp, no lysogen induction essay

Table 4. Nonparametric Spearman rank correlation matrix for chlorophyll *a*, bacterial and viral parameters from the fertilized (n=8-9, except for BP-BA-chla: 36-41) and HNLC stations (n=10, except for BP-BA-chla: 23-31). Bold numbers are significant r-values (\*  $P < 0.05$ , \*\*  $P < 0.001$ , \*\*\*  $P < 0.0001$ ).

	Chl <i>a</i>	BA	BP	VA	VP <sub>1</sub>
Fe-fertilized					
BA	0.209				
BP	0.243	<b>0.633***</b>			
VA	0.357	0.548	<b>0.762*</b>		
VP <sub>1</sub>	0.083	0.476	-0.050	0.310	
FIC	-0.183	0.333	-0.267	0.095	<b>0.900**</b>
HNLC					
BA	<b>0.688*</b>				
BP	<b>0.380*</b>	<b>0.635*</b>			
VA	0.164	0.576	0.515		
VP	0.426	<b>0.746*</b>	0.406	0.273	
FIC	0.168	0.304	0.310	-0.249	0.608

BA, bacterial abundance; BP, bacterial production; VA, viral abundance; VP<sub>1</sub>, lytic viral production; FIC, fraction of infected cells

Table 5. Comparison of viral abundance (VA) and production (VP), virus-mediated bacterial mortality (VMM) and % loss of bacterial production (% BP) and standing stock per day (% SS d<sup>-1</sup>) with literature data from other polar/subpolar environments.

Location	Depth (m)	Method	VA (10 <sup>9</sup> L <sup>-1</sup> )	VP (10 <sup>9</sup> L <sup>-1</sup> d <sup>-1</sup> )	VMM (10 <sup>8</sup> L <sup>-1</sup> d <sup>-1</sup> )	% BP	% SS d <sup>-1</sup>	Source
SO: Fe-fertilized	0-150	VRA	3.4-14.2 (9.9 ± 3.6)	9.9-117.9 (59.0 ± 47.1)	0.8-10.3 (5.4 ± 4.1)	8-202 (72)	104	present study
SO: HNLC	0-200	VRA	3.1-7.4 (4.7 ± 1.4)	6.0-25.6 (14.5 ± 7.4)	0.4-2.1 (1.1 ± 0.6)	6-58 (27)	44	present study
Antarctic	0-100	VDR	1-74 (13±10.4)			>100		Guixa-Boixereu et al., 2002
SO-Subantarctic	10	VRA	6.1-26	17.5-216.3	3.6-43.3	43-63	40-130	Evans et al., 2009
SO	5-200	VRA	0.5-7.6	0.4-16	0-8.7		0-72	Evans and Brussaard, 2012
SO: Fe patch	10-150	VRA	2.3-7(4.3±5.5)	0.9-3.6(1.9±0.5)		41-172 (104)*		Weinbauer et al., 2009
SO: HNLC	10-150	VRA	1.4-2.5(2.1±2)	0.3-0.8(0.6±0.1)		14-70 (39)*		Weinbauer et al., 2009
Arctic	0-10	TEM	2.5-36	0.2-4.6 (2)		2-36 (13)		Steward et al., 1996
North waters	0-200	TEM	1.36-5.55(3.3±1.6)	0.1-1.3			6-28	Middelboe et al., 2002
Arctic	0-230	VDA	1.4-4.5(2.8±1.3)	0.1-1.9	0.28-0.72			Wells and Deming, 2006
Subarctic Fe patch	0-10	TEM/VRA	40.5	30-200	90±25		7.4	Higgins et al., 2009
Subarctic outside	0-10	TEM/VRA	35.7	30-200	25.8±6.1		7.2	Higgins et al., 2009
Arctic	0-100	VRA	0.32-7.28	0.1-4.2		2-24 (9)	2-30	Boras et al., 2010
Canadian Arctic Shelf	2-56	VRA	2.7-27	0.03-7.7	0.02-4.3	31-156	1.4-29	Payet and Suttle, 2013

SO: Southern Ocean, VDR: viral decay rates, TEM: Frequency of visibly infected cells by transmission electron microscopy, VDA: Virus dilution approach, VRA: virus reduction approach. \*using BP in the VRA

Table 6. C and Fe release rates ( $L^{-1} d^{-1}$ ) through viral lysis calculated from VP ( $12.4 \text{ fg C cell}^{-1}$ , (Fukuda et al., 1998) and from FIC following the Model by Binder (1999) using bacterial iron quota of  $7.5 \mu\text{Mol Fe mol C}^{-1}$  (Tortell et al., 1996). Averages are given in parenthesis.

	Release based on VP		Release based on FIC	
	pmol Fe $L^{-1} d^{-1}$	$\mu\text{mol C } L^{-1} d^{-1}$	pmol Fe $L^{-1} d^{-1}$	$\mu\text{mol C } L^{-1} d^{-1}$
Fertilized stations	0.60-7.97 (4.18 $\pm$ 3.15)	0.08-1.06 (0.56 $\pm$ 0.42)	0.03-1.58 (0.42 $\pm$ 0.49)	0.003-0.21 (0.06 $\pm$ 0.07)
HNLC stations	0.28-1.60 (0.86 $\pm$ 0.43)	0.04-0.21 (0.11 $\pm$ 0.06)	0.004-0.12 (0.05 $\pm$ 0.05)	0.001-0.02 (0.01 $\pm$ 0.01)
ratio	<b>4.9*</b>	<b>4.9*</b>	<b>7.9*</b>	<b>7.9*</b>

\*Values are significantly higher in the Fe-fertilized than in the HNLC stations (Kruskal-Wallis,  $P < 0.05$ )

Figure 1. Real-time satellite images of chlorophyll during the KEOPS cruise dating from the first sampling of station A3 (19/2/2005), (MODIS results provided by CSIRO marine research) and overlaid transects and sampled stations.

Figure 2. Depth profiles of bacterial (A) and viral abundance (B) in the Kerguelen study area. Full symbols indicate Fe-fertilized sites, open symbols indicate HNLC waters.

Figure 3. Lytic viral production from the Fe-fertilized (A) and HNLC (B) stations. Values are the averages of duplicates and error bar indicate the minimum and maximum values. When not visible, error bars are within the width of the line.

Figure 4. Simple sketch of the carbon and nutrient flow through the microbial food web in the Fe- fertilized (left) and HNLC waters (right). Arrow thickness represents the relative importance of factors controlling the size of each pool of the microbial food web.