Biogeosciences Discuss., 11, C6335–C6337, 2014 www.biogeosciences-discuss.net/11/C6335/2014/

© Author(s) 2014. This work is distributed under the Creative Commons Attribute 3.0 License.



**BGD** 

11, C6335-C6337, 2014

Interactive Comment

## Interactive comment on "Ocean acidification accelerates dissolution of experimental coral reef communities" by S. Comeau et al.

S. Comeau et al.

steve.comeau@csun.edu

Received and published: 31 October 2014

Comment 1: "First, I need to highlight the fact that I am not a coral biologist and have no practical experience with the used methods. As a consequence, my report focusses on general experimental procedures. The most interesting part of the manuscript is the use of artificial "controlled" com- munities to evaluate the impact of ocean acidification on net calcification. This is an under-used methodological approach and a nice alternative to natural analogs and/or in situ mesocosms with the potential to better understand the mechanisms behind the response at the community level. From my limited expertise, I do not see any major problem with the manuscript and only have a few comments/questions:"

Response 1: We thank Sam Dupont for his useful comments and we are glad that C6335

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



someone outside of the coral scientific community finds our approach interesting.

Comment 2: - Can you provide some information on the natural variability experienced by these communities in the field? That would help putting the tested scenarios into context.

Response 2: Back reef communities in Moorea are exposed to relatively constant pH with diel variations of  $\sim$  0.1 unit. This information is provided in the manuscript lines 122-124" To match the natural diel variation in pH in the back reef of Moorea (Hofmann et al., 2011; Comeau et al., 2014a) pH was maintained 0.1 unit lower at night (from 18:00 to 6:00) than during the day."

Comment 3: - Please provide a Table with the seawater chemistry over the course of the experiment. Also include statistics to demonstrate that you have significant differences between treatments. Provide graphs if necessary to illustrate the variability.

Response 3: As suggested a table (Table 1) presenting the carbonate chemistry is now included in the manuscript. Statistics on the pCO2 treatments are now provided lines 170-176 " Mean pCO2 in the four flumes during the 8-week incubation was 456  $\pm$  21  $\mu$ atm and 451  $\pm$  21  $\mu$ atm in the ambient treatments, and 1329  $\pm$  28  $\mu$ atm and 1306  $\pm$  41  $\mu$ atm in the high pCO2 treatments ( $\pm$  SE, n = 42). pCO2 differed between treatments (repeated measure ANOVA, F1,232 = 734.38, p < 0.001), but there was no difference within treatments (F2,232 = 0.16, p = 0.852). Communities were maintained in conditions within the flumes that were super-saturated with respect to aragonite, as  $\Omega$ arag  $\sim$  3.5 under ambient conditions, and  $\sim$  1.6 in the high pCO2 treatment. ". However, we do not believe that a graph representing the carbonate chemistry will be insightful since variability of pCO2 was very limited (se  $\sim$  30  $\mu$ atm, Table 1).

Comment 4: Full statistics should be better presented in the results (provide model, F and p values for tested parameters (pCO2 and replicate).

Response 4: As suggested more details on the statistics are now presented.

## **BGD**

11, C6335-C6337, 2014

Interactive Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



Comment 5: - At the end of the Results, it is claimed that coralline algae experienced high mortality. Information on "health" of all tested species should be provided and presented at the beginning of the result section.

Response 5: Thank you for these recommendations, a "health" section is now included in the Results, lines 177-182 " No Pocillopora spp. and Montipora spp. colonies died during the 8-week treatments, but 10% of the Porites pooled across flumes died by the end of the experiment, regardless of treatment, because of an outbreak of corallivorous nudibranchs feeding on this taxon (Phestilla spp.). Coralline algae ( $\sim$  70%) died at the end of the incubation, which was likely due to sediment abrasion. No difference in mortality or bleaching was observed between treatments for corals and calcified algae."

Interactive comment on Biogeosciences Discuss., 11, 12323, 2014.

## **BGD**

11, C6335-C6337, 2014

Interactive Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

