

This discussion paper is/has been under review for the journal Biogeosciences (BG).  
Please refer to the corresponding final paper in BG if available.

# Seasonal survey of the composition and degradation state of particulate organic matter in the Rhone River using lipid tracers

M.-A. Galeron<sup>1</sup>, R. Amiraux<sup>1</sup>, B. Charriere<sup>2</sup>, O. Radakovitch<sup>3</sup>, P. Raimbault<sup>1</sup>,  
N. Garcia<sup>1</sup>, V. Lagadec<sup>1</sup>, F. Vaultier<sup>1</sup>, and J.-F. Rontani<sup>1</sup>

<sup>1</sup>Aix-Marseille University, Mediterranean Institute of Oceanography (MIO), 13288, Marseille, Cedex 9; Université du Sud Toulon-Var, 83957, CNRS-INSU/IRD UM 110, France

<sup>2</sup>Centre de Formation et de Recherche sur l'Environnement Méditerranéen (CEFREM, UMR CNRS 5110), Bât. U, Université de Perpignan, Via Domitia (JPVD), 66860 Perpignan, France

<sup>3</sup>CEREGE UM34, Aix Marseille Université CNRS, IRD, CDF, 13545 Aix-en-provence, France

Received: 1 September 2014 – Accepted: 15 September 2014 – Published: 2 October 2014

Correspondence to: M.-A. Galeron (marie-aimee.galeron@mio.osupytheas.fr)

Published by Copernicus Publications on behalf of the European Geosciences Union.

**Survey of composition and degradation state of POM in the Rhone River using lipid tracers**

M.-A. Galeron et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Abstract

Lipid tracers including fatty acids, hydroxyacids, *n*-alkanols, sterols and triterpenoids were used to determine the origin and fate of suspended particulate organic matter (POM) collected in the Rhone River (France). This seasonal survey (April 2011 to May 2013) revealed a year-round strong terrigenous contribution to the plant-derived particulate organic matter (POM), with significant algal inputs observed in March and attributed to phytoplanktonic blooms likely dominated by diatoms. Major terrigenous contributors to our samples are gymnosperms, and more precisely their roots and stems, as evidenced by the presence of high proportions of  $\omega$ -hydroxydocosanoic acid (a suberin biomarker). The high amounts of coprostanol detected clearly show that the Rhone River is significantly affected by sewage waters.

Specific sterol degradation products were quantified and used to assess the part of biotic and abiotic degradation of POM within the river. Plant-derived organic matter appears to be mainly affected by photo-oxidation and autoxidation (free radical oxidation), while organic matter of human origin, evidenced by the presence of coprostanol, is clearly more prone to bacterial degradation. Despite the involvement of an intense autoxidation-inducing homolytic cleavage of peroxy bonds, a significant proportion of hydroperoxides is still intact in higher plant debris. These compounds could affect the degradation of terrestrial material by inducing an intense autoxidation upon its arrival at sea.

## 1 Introduction

With an average water discharge of  $1700 \text{ m}^3 \text{ s}^{-1}$  (Ludwig et al., 2009), the Rhone is the main freshwater contributor to the Mediterranean Sea. Based on various studies, the mean annual flux of suspended matter is about  $6.7 \times 10^6 \text{ t yr}^{-1}$  over the period 1967–2008 (Eyrolle et al., 2012), and the annual particulate organic carbon (POC) flux was estimated to be  $1.94 \pm 1.09 \times 10^5 \text{ t yr}^{-1}$  (Sempéré et al., 2000). The labile POC fraction,

BGD

11, 14197–14237, 2014

### Survey of composition and degradation state of POM in the Rhone River using lipid tracers

M.-A. Galeron et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

---

**Survey of composition and degradation state of POM in the Rhone River using lipid tracers**

M.-A. Galeron et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

calculated as the carbon contribution from sugars and amino acids to Total POC, was estimated to be between 3 and 27 % depending on the amount of SPM (Sempéré et al., 2000). The Rhone River represents 14 and 10 % of fresh water and particulate carbon input to the Mediterranean Sea, respectively (Ludwig et al., 2009). Given the importance of such a contribution, it is crucial to study the provenance and composition of the organic matter found in the Rhone, prior to its arrival in the Mediterranean Sea.

Up until recently, it was widely considered that the terrestrial particulate organic matter (TPOM) flowing into oceans through rivers was refractory to degradation (biotic and abiotic), since it was constituted of terrestrial plant debris previously degraded during transport (de Leeuw et Largeau, 1993; Wakeham et Canuel, 2006). However, coastal sediments have shown very little trace of a terrestrial OM signature (Hedges and Keil, 1995; McKee et al., 2004), which implies that either global carbon fluxes and budgets are wrong, or, and it is more likely, that terrestrial POM undergoes a rapid and intense degradation upon its arrival at sea (Hedges et al., 1997). The belief that terrestrial POM is refractory to all sorts of decomposition has been recently challenged through studies in the Mediterranean Sea (Bourgeois et al., 2011) and Northern Canada (Rontani et al., 2014b), showing that well-preserved TPOM is heavily degraded upon leaving river systems. This is why we intend to study the chemical structure and degradation state of TPOM in the Rhone River, in order to better understand how such an intense degradation can be possible at sea, while it was not during freshwater transport.

In comparison with bulk geochemical analyses, where  $\delta^{13}\text{C}$  terrestrial end-member determination is precarious,  $\delta^{13}\text{C}$  signatures of organic matter at a molecular level can be more specific (Tolosa et al., 2013). Unfortunately, carbon isotopic data in freshwater ecosystems is not always source-specific, because freshwater phytoplankton and terrestrial plants can often produce similar  $\delta^{13}\text{C}$  signatures (Cloern et al., 2002). Many studies used biomarkers such as lipids, cutins or waxes in sediment or soil samples in order to determine the specific contribution of plants in organic matter (Amelung et al., 2008; Simpson et al., 2008). Sterols (steroidal alcohols) have specific structural features that can be linked to a restricted number of organisms (Volkman, 1986), and can

## Survey of composition and degradation state of POM in the Rhone River using lipid tracers

M.-A. Galeron et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

be used to determine the main contributors to a pool of organic matter. Their degradation products, such as stanols, 3,6-diols, or triols, can also inform on the type of degradation undergone (Christodoulou et al., 2009; Rontani et al., 2009). On the other hand, cuticular waxes and cutins can be linked to relatively specific higher plant groups (Mueller et al., 2012), allowing an estimation of the contribution of terrestrial plants to the OM.

There are no studies dealing with the composition of the plant-derived particulate organic matter carried along the Rhone River in the literature. **It would be logical** to think that it is mainly constituted of terrestrially-produced particulate organic matter travelling with the runoff, and we propose to confirm this hypothesis here, while also identifying other sources of OM, sometimes significantly aquatic/planktonic. Although limited research has been done on freshwater plankton dynamics in the Rhone, it has been found that phytoplankton communities in the middle Rhone suffer a high turnover (Fruget et al., 2001) implying an important potamoplankton input in the particulate organic matter flowing down the river.

Using specific lipids and their degradation products, here we report for the first time on the different sources of OM in the Rhone River and discuss the main degradation processes affecting this OM.

## 2 Materials and methods

### 2.1 Study area and Sampling

The Rhone River is 816 km long and has a drainage area of 97 800 km<sup>2</sup> (Fig. 1). The Rhone being a river prone to seasonal flooding, samples span a wide range of water flow conditions in order to be as representative as possible (Fig. 2).

To monitor temporal variations of suspended particulate matter (SPM), POC, nutrients and lipidic biomarkers in the Rhone River, 30 surface water and particle samples

were collected between April 2011 and May 2013 at the Rhone River reference estuarine station of Arles, 40 km upstream from the river mouth (Fig. 1).

The high frequency study of the nutrients and particulate matter input by the Rhone to the Mediterranean Sea has been carried out since 2010 in the framework of the national program MOOSE (Mediterranean Ocean Observing System for the Environment). Monitoring was undertaken at the Arles station (Fig. 1). Sampling included filtering between 100 and 400 mL of water on GF/F glass fiber filters, as well as particles collection using a high-speed centrifuge device (CEPA Z61) coated with teflon to avoid metal contamination. For suspended matter determination, water samples were filtered on pre-weighted GF/F filters. After drying, filters were weighted to determine the suspended particulate matter (SPM) content (in  $\text{mg L}^{-1}$ ). Non-weighted filters were used to quantify Particulate Organic Carbon (POC) contents: the filters were acidified with 50–100  $\mu\text{L}$  0.5 N sulfuric acid and dried overnight at 60 °C. POC contents were determined using high combustion (900 °C) procedure on a CN Integra mass spectrometer (Sercon). Filtrates were used to analyze inorganic nutrients (nitrate, nitrite, phosphate, silicate) using the automated colorimetric method described in Aminot and K erouel (2007). Uncertainty of SPM is estimated to be 0.05  $\text{mg L}^{-1}$  while the analytical error for POC determination is 5  $\mu\text{M}$  (both errors were calculated during method validation, using replicates, and take into account scale and spectrometer precision and detection limits). Chlorophyll concentration was quantified using methanol extraction according to Raimbault et al. (2004). Samples (filters, filtrates and particles) were immediately frozen at  $-20\text{ }^\circ\text{C}$  until analysis.

## 2.2 Chemical treatment of the samples

Prior to analysis all samples were thawed and their water content was measured through the weighting of humid vs. oven-dried particles: dry particles (g)/humid particles (g) · 100. The suspended particles collected (between 200 and 450 mg per sample) first needed to be reduced with  $\text{NaBH}_4$  and saponified.  $\text{NaBH}_4$ -reduction of hydroperoxides to alcohols that are amenable to

BGD

11, 14197–14237, 2014

### Survey of composition and degradation state of POM in the Rhone River using lipid tracers

M.-A. Galeron et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Survey of composition and degradation state of POM in the Rhone River using lipid tracers**

M.-A. Galeron et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

gas chromatography-electron impact mass spectrometry (GC-EIMS) is essential for estimating the importance of photo-oxidative and autoxidative degradation in natural samples (Marchand and Rontani, 2001). Without this preliminary treatment, these labile compounds can be thermally cleaved during alkaline hydrolysis or GC analysis and thus be overlooked during conventional organic geochemical studies. Lipids and their degradation products in the resulting total lipid extracts (TLEs) were then quantified by GC-EIMS. All manipulations were carried out using foil-covered vessels in order to exclude photochemical artifacts. It is well known that metal ions can promote autoxidation during hot saponification (Pokorny, 1987). The prior reduction of hydroperoxides with NaBH<sub>4</sub> allowed us to avoid such autoxidation artifacts during the alkaline hydrolysis.

Particles were placed in methanol (MeOH) (15 mL) and hydroperoxides were reduced to the corresponding alcohols with excess NaBH<sub>4</sub> (70 mg; 30 min at 20 °C). During this treatment, ketones are also reduced and the possibility of some ester cleavage cannot be excluded.

Saponification was carried out on reduced samples. After NaBH<sub>4</sub> reduction, 15 mL water and 2.24 g KOH were added and the mixture directly saponified by refluxing for 2 h. After cooling, the content of the flask was acidified with HCl (pH 1) and extracted (3×) with dichloromethane (DCM). The combined DCM extracts were concentrated to give the TLE. After solvent evaporation, residues were taken up in 300 μL of pyridine/N,O-bis(trimethylsilyl)trifluoroacetamide (BSTFA; Supelco; 2 : 1, v/v) and silylated for 1 h at 50 °C to convert OH-containing compounds to TMSi-ether or ester derivatives. After evaporation to dryness under a stream of N<sub>2</sub>, the derivatized residues were taken up in a mixture of ethyl acetate/BSTFA (to avoid desilylation of fatty acids) for GC-EIMS analysis. It should be noted that under these conditions steran-3β,5α,6β-triols were silylated only at C3 and C6 and thus need to be analyzed with great care (Rontani et al., 2014b).

A different treatment was used to quantify hydroperoxides and their ketonic and alcoholic degradation products. The samples were extracted three times with chloroform-

MeOH-H<sub>2</sub>O (1 : 2 : 0.8, v/v/v) using ultrasonication. The supernatant was separated by centrifugation at 3500G for 9 min. To initiate phase separation, purified H<sub>2</sub>O was added to the combined extracts to give a final volume ratio of 1 : 1 (v/v). The upper aqueous phase was extracted three times with DCM and the combined DCM extracts were filtered and the solvent removed via rotary evaporation. The residue obtained after extraction was dissolved in 4 mL of DCM and separated in two equal subsamples. After evaporation of the solvent, degradation products were obtained for the first subsample after acetylation (inducing complete conversion of hydroperoxides to the corresponding ketones, Mihara and Tateba, 1986) and saponification and for the second after reduction with NaBD<sub>4</sub> and saponification. Comparison of the amounts of alcohols present after acetylation and NaBD<sub>4</sub> reduction made it possible to estimate the proportion of hydroperoxides and alcohols present in the samples, while after NaBD<sub>4</sub>-reduction deuterium labelling allowed to estimate the proportion of ketones really present in the samples (Marchand and Rontani, 2003).

### 2.3 GC-EIMS analyses

Lipids and their oxidation products were quantified using an Agilent 7850-A gas chromatograph connected to an Agilent 7000-QQQ mass spectrometer. The following conditions were employed: 30m × 0.25mm (i.d.) fused silica column coated with HP-5MS (Agilent; film thickness: 0.25 μm); oven programmed from 70 to 130 °C at 20 °C min<sup>-1</sup>, then to 250 °C at 5 °C min<sup>-1</sup> and then to 300 °C at 3 °C min<sup>-1</sup>; carrier gas (He), 1.0 bar; injector (splitless), 250 °C; injector (on column), 50 °C; electron energy, 70 eV; source temperature, 230 °C; quadrupole temperature, 150 °C; *m/z* 40–700; collision energy, ranging from 5 to 15 eV; collision flow, 1.5 mL min<sup>-1</sup> (N<sub>2</sub>); quench flow, 2.25 mL min<sup>-1</sup> (He); cycle time, 0.2 s. An on-column injector was used for the analysis of sterol degradation products and a splitless injector for the analysis of FA degradation products. Compounds were assigned by comparison of retention times and mass spectra with those of standards and quantified (calibration with external standards) with GC-EIMS. For low concentrations, or in the case of co-elutions, quantification was achieved using

## Survey of composition and degradation state of POM in the Rhone River using lipid tracers

M.-A. Galeron et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





selected ion monitoring (SIM) or Multiple Reaction Monitoring (MRM). The main characteristic mass fragment ions used to quantify degradation products of sterols have been described previously (Christodoulou et al., 2009; Rontani et al., 2011). Using replicates, the analytical standard error for lipid quantification (from preparation to integration) was estimated to be 14 % (Standard Error = Standard Deviation/ $\sqrt{n}$  for  $n$  replicates).

### 3 Results and discussion

#### 3.1 Water discharge, suspended particulate matter (SPM) and particulate organic carbon (POC)

Water flow patterns in the Rhone show pronounced seasonal variations predominantly influenced by 3 factors that can induce maximum discharge (Vivian, 1989): snow melting (May–June), intense rains in the south basin (October–November) and rains over the entire basin (January–March). During our sampling period the daily flow rate fluctuated between 360 (8 October 2011) and 5200 m<sup>3</sup> s<sup>-1</sup> (19 May 2013; Fig. 2). Suspended Particulate Matter (SPM) and POC concentrations (Fig. 2) ranged from 6.5 to 1381.3 mg L<sup>-1</sup> and from 17.9 to 1383.3 μM C, respectively.

The temporal evolution of suspended particulate matter (SPM) clearly followed the same pattern than water discharge (Fig. 2), which is typical for river systems. The samples collected on flood dates (November 2011, May 2013, liquid discharges above 3000 m<sup>3</sup> s<sup>-1</sup>) display the highest SPM content and POC values (Fig. 2). POC expressed as a percentage of SPM, ranged from 0.8 to 11.6 %, and tends to be lower when water flows are highest (Sempéré et al., 2000)

**BGD**

11, 14197–14237, 2014

## Survey of composition and degradation state of POM in the Rhone River using lipid tracers

M.-A. Galeron et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





## 3.2 Use of lipid tracers to estimate the composition and origin of POM collected in the Rhone River

### 3.2.1 Sterols

Based on a literature review, cholest-5-en-3 $\beta$ -ol (cholesterol), 24-ethylcholest-5-en-3 $\beta$ -ol (sitosterol if the C-24 stereochemistry is 24 $\alpha$ ), cholesta-5,24-dien-3 $\beta$ -ol (desmosterol), 24-methylcholesta-5,24(28)-dien-3 $\beta$ -ol (24-methylenecholesterol), 24-methylcholesta-5,22-dien-3 $\beta$ -ol (brassicasterol and/or epi-brassicasterol depending on C-24 stereochemistry) and 24-methylcholest-5-en-3 $\beta$ -ol (campesterol) have been selected as tracers of the origin of POM. Apart from cholesterol, which can originate from a wide number of sources, all the other quantified sterols are relatively source-specific. Sitosterol constitutes the major sterol in higher plants, even though it can also be found in diatoms (Volkman, 1986), and is often used to trace terrestrial organic matter in lacustrine and marine systems (e.g. Meyers and Ishiwatari, 1993). Desmosterol is mainly found in algae (Volkman, 1986). 24-Methylenecholesterol is mainly found in diatoms, more particularly in the *Thalassiosira* and *Skeletonema* genera in the marine realm (Volkman, 2003). Epi-brassicasterol is mostly found in algae (Volkman, 1986).

All samples are dominated by sitosterol and cholesterol, with proportions being on average three times higher than those of the other sterols, apart from the 6 March 2012 sample (Table 1). This sample exhibits a rather different profile, dominated by desmosterol (0.122  $\mu\text{g mg}^{-1}$  (dry weight)) and 24-methylenecholesterol (0.096  $\mu\text{g mg}^{-1}$  (dry weight)), with a strong contribution from brassicasterol (0.026  $\mu\text{g mg}^{-1}$  (dry weight)) compared to all other samples. All these sterols have been considered to be planktonic markers, and have been summed to compose the planktonic sterol fraction shown in Fig. 3a. This fraction forms the major part of the total sterol fraction of the March 2012 sample, evidencing a phytoplanktonic bloom event. Given the nature of the sterols involved (desmosterol, methylene-cholesterol), diatoms seem to be major contributors (Rampen et al., 2010). This concurs with the drops in silica observed at the same period (MOOSE data, unpublished). A we also see an increase, although less important,

### Survey of composition and degradation state of POM in the Rhone River using lipid tracers

M.-A. Galeron et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Survey of composition and degradation state of POM in the Rhone River using lipid tracers

M.-A. Galeron et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

in the proportion of planktonic sterols in the Spring 2013 samples, this type of planktonic event is probably a yearly spring occurrence but our sampling frequency was not adapted to study blooms that can appear and disappear in a matter of days. Sitosterol has been previously identified in marine algae, and more widely as the major sterol in higher terrestrial plants (Volkman, 1986, 2003). However the clear increase observed here in March 2012 demonstrates that it is also present in potamoplankton (Fig. 3a). The presence of cholesterol can indicate a zooplankton contribution (Volkman, 1986), but it is also often found in freshwater algae (Volkman et al., 1981; Gagosian et al., 1983), and can also evidence human impacts (Sicre et al., 1993). The relatively high proportions of  $5\beta$ (H)-cholestan- $3\beta$ -ol (coprostanol) detected in the samples investigated (see Sect. 3.3.2) clearly show that in the Rhone River cholesterol mainly arises from waste water inputs (Brown and Wade, 1984).

We also looked at the ratio sitosterol/campesterol (Table 1) as an indicator of the terrigenous vs. diatom origin of our POM. While many plants have a sitosterol/campesterol ratio of less than 4 (Volkman, 1986), Nishimura and Koyama (1977) reported values ranging from 11.5 (*Pinus densiflora*) to 31 (holly *Ilex pedunculosa*). Dachs et al. (1998) identified the threshold of 1 as the limit under which samples are dominated by OM of aquatic origin, and above which sources of OM are terrigenous. In our case, all the samples present a sitosterol/campesterol ratio above 1, mainly between 2 and 8 (up to 12). Three samples show a ratio near 1 (July 2011 and 2012 and March 2012; Table 1), a low value which confirms the presence of diatoms in the plankton since freshwater diatoms exhibit sitosterol/campesterol ratios close to 1.0 (Ponomarenko et al., 2004).

### 3.2.2 Terrestrial vascular plant biomarkers

The contribution of terrestrial vascular plants to our samples is also evidenced by the presence of the triterpenoids betulin (lup-20(29)-ene- $3\beta$ ,28-diol), oleanolic ( $3\beta$ -hydroxyolean-12-en-28-oic) and ursolic ( $3\beta$ -hydroxyurs-12-en-28-oic) acids (Razboršek et al., 2008) as well as components of cuticular waxes

**Survey of composition and degradation state of POM in the Rhone River using lipid tracers**

M.-A. Galeron et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

(16,8-11-dihydroxyhexadecanoic,  $\omega$ -hydroxyhexadecanoic,  $\omega$ -hydroxyoleic and 18-hydroxyoctadec-9-enoic acids) (Kolattukudy, 1980a) (Fig. 3b). Betulin has been proposed as a tracer for paper birch (Fine et al., 2001), a common species along the Rhone River, while oleanolic and ursolic acids are widely distributed in terrestrial higher plants (Liu, 1995). The amount of cuticular waxes is variable amongst samples, between 0.02 and  $3.8 \mu\text{g mg}^{-1}$  (dry weight), with the highest in the 5 April 2011, 2 May 2013 and 4 November 2011 samples ( $3.8$ ,  $2.2$  and  $1.7 \mu\text{g mg}^{-1}$  respectively), which happen to be flood dates. It is clear that floods, during which higher water flows are coupled to surface runoff, collect and carry more terrestrial plant leaf debris, and hence increase the amount of cuticular waxes found in our samples. The yearly variations in quantity are probably due to the fact that waxes (linear compounds) are more easily degraded by bacteria than cyclic structures such as sterols or triterpenoids (Atlas and Bartha, 1992).

The betulin and sitosterol concentrations are significantly correlated in most of our samples ( $r = 0.67$  between sitosterol and betulin, on 29 samples,  $p$  value =  $3 \times 10^{-5}$ , excluding the 6 March 2012 sample due to its out-of-range phytoplanktonic profile), thus reinforcing the idea that in the Rhone River sitosterol mainly results from terrestrial higher plant inputs. However, at the time of the spring bloom a significant part of this sterol seems to derive from potamoplankton (Fig. 3a).

Another ratio commonly used to attest to the terrigenous origin of compounds is the Terrigenous-to-Aquatic ratio (TAR, Bourbonnière and Meyers, 1996). Here we used the  $\text{TAR}_{(\text{AL})}$  as calculated by Van Dongen et al. (2008) for  $n$ -alkanols:  $(C_{26} + C_{28}) / (C_{16} + C_{18})$ . The  $\text{TAR}_{(\text{AL})}$  in our samples is always above 1, and clearly indicates a strong terrigenous contribution to the suspended particulate matter found in the lower Rhone. The average chain length of  $n$ -alkanols, a proxy positively correlated to the abundance of higher plant debris (Van Dongen et al., 2008), ranged from 26 to 22 across all samples, also attesting to the strong contribution of terrestrial vascular plants. The long-chain even-numbered  $n$ -alcohol profiles show a strong contribution of  $C_{22}$  and  $C_{28}$   $n$ -alkanols. Compared with those previously described in the literature (Diefendorf et al.,

2011), this characteristic suggests a strong gymnosperm contribution, which concurs with the low amounts of long-chain *n*-alkanes detected.

### 3.2.3 Chlorophyll

The available data on chlorophyll *a* (MOOSE database, only available for 2012 and 2013) shows a content variability between 0.9 (10 October and 6 November 2012) and 14.0 (3 April 2012) mg m<sup>-3</sup> (Fig. 4a). Chlorophyll *a* is frequently used as a proxy for photosynthetic organisms and the variation observed here is consistent with the hypothesis of a yearly phytoplanktonic spring bloom, with a larger magnitude for the 2012 event.

### 3.2.4 Fatty acids

A number of saturated linear fatty acids have been found in our samples, with C<sub>16:0</sub> and C<sub>18:0</sub> being the most abundant (Table 2). These fatty acids are not specific, and can stem from a number of sources including terrestrial vascular plants, algae and bacteria (Volkman et al., 1981). Longer-chain saturated fatty acids (between C<sub>20</sub> and C<sub>28</sub>) with a strong even-chain predominance, which are specific to the epicuticular waxes of vascular plants (Kunst and Samuels, 2003), are scarcely present in our samples, with C<sub>20</sub> and C<sub>22</sub> being the most abundant relative to the others: over 98 % (on average) of the total of long-chain (C<sub>20</sub>–C<sub>28</sub>) saturated fatty acids across all samples.

**Polyunsaturated fatty acids** are present in very low proportions in our samples, apart from the 6 March 2012 and 12 March 2013 samples where they contributed to 44 and 40 % of total fatty acids (Fig. 3c). **These high contributions support the presence of a high proportion of fresh algal material in these samples.**

If we compare the average chain length (ACL) of fatty acids in our samples with that of *n*-alkanols, it appears clearly that the ACL of fatty acids is lower, with an average of 16.7 across samples, against 23.9 for alcohols. It is widely accepted that fatty acids are more prone to bacterial degradation than other lipids (Wakeham, 1995), and long-

BGD

11, 14197–14237, 2014

## Survey of composition and degradation state of POM in the Rhone River using lipid tracers

M.-A. Galeron et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



chain fatty acids tend to be degraded more efficiently by a number of bacteria (Novak and Carlson, 1970). Such a bacterial degradation could explain the ACL difference between *n*-alkanols and fatty acids, and the lack of terrestrial higher-plant fatty acids in our samples, while other markers for higher plants (such as waxes or betulin) are present in large quantities. This is reinforced by the fact that we also find a relatively high proportion of vaccenic acid in our samples, a specific marker for bacterial activity (Sicre et al., 1988).

### 3.2.5 Hydroxyacids

Hydroxyacids contents were low during the period studied but some samples (18 July 2011, 4 November 2011 and 16 January 2012) exhibited high amounts of C<sub>22</sub>  $\omega$ -hydroxyacid and small quantities of C<sub>20</sub> and C<sub>24</sub> homologues (Table 2). These compounds are generally considered to be suberin markers when found in soil (Nierop, 1998; Otto et al., 2005), even though they have been found in leaves and stems of higher plants as well (Mendez-Millan et al., 2010). Suberin is a cell wall component of cork cells, and is mainly found in bark, woody stems, and roots (Kolattukudy, 1980a). Given the geographical location of our sampling point, we can consider this compound to be a marker of industrial activity, probably associated to a paper paste mill being implanted less than 3 km upstream. The mill uses mainly local conifers (collected within a 250 km radius according to the company, Paper Excellence, 2014), more particularly *Pinus halepensis*, *Pinus nigra*, and *Pinus sylvestris* (Etude AGRESTE, 2011) and is authorized to reject a certain amount of waste water in the river. This is consistent with our findings that the POM is dominated by gymnosperms. *Pinus* species also display a sitosterol/campesterol ratio comprised between 5 and 10 (Conner et al., 1980) in line with most of our samples as well. This **harmless** industrial contamination could explain the large contribution of gymnosperms to our POM, and implies that we consider the extra input of terrestrial plant matter that will be released, and degraded, at sea.

As a summary, the overall lipid composition of the Rhone River SPM is characterized by major terrestrial higher plant components (mainly derived from gymnosperms)

## Survey of composition and degradation state of POM in the Rhone River using lipid tracers

M.-A. Galeron et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



with episodic, but significant, contributions from freshwater algal material (probably dominated by diatoms) in the spring. Despite the strong concentration of industries along this river, SPM appears to be very weakly contaminated by petroleum hydrocarbons, but is strongly impacted by the local paper mill and wastewater discharges (see Sect. 3.2.2).

### 3.3 Use of lipid tracers to estimate the degradation state of POM from the Rhone River

If they can inform us on the origin of organic matter in natural environments, lipid biomarkers are also invaluable in helping us estimate biotic and abiotic alterations of organic matter and determine what are the main processes involved in its degradation. Products resulting from the degradation of sterols, chlorophyll, monounsaturated fatty acids and hydroxyacids are among the most useful and specific tracers.

#### 3.3.1 Chlorophyll

The absorption of light by some compounds, called photosensitizers, in the presence of oxygen (regardless of these compounds being endogenous or not) causes an oxidation whose effects, chemical or biological, are mostly adverse (Spikes and Straight, 1967). Photosensitizers induce chemical reactions via the absorption of light that would not occur in their absence. Photosensitizers (Sens) are involved in indirect photo-oxidative processes: they have 2 systems of electronically excited states,  $^1\text{Sens}$  and  $^3\text{Sens}$ . The triplet state is much longer lived than the singlet state, which is the initial product issued from light absorption. Indirect photo-oxidation (photo-sensitized oxidation) can be intense during the senescence of phototrophic organisms (Rontani, 2012) due to the presence of chlorophyll, which is a very efficient photosensitizer (Foote, 1976) capable of generating singlet oxygen particularly reactive towards unsaturated cellular components (Type II photoprocesses). Chlorophyll may be also directly photodegraded by solar light (Nelson, 1993). Direct photodegradation of chlorophyll and Type II photo-

BGD

11, 14197–14237, 2014

### Survey of composition and degradation state of POM in the Rhone River using lipid tracers

M.-A. Galeron et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



oxidation of unsaturated cellular components can be thus considered two competitive photo-processes.

In the photic layer of aquatic environments, photo-oxidation has long been considered a major degradation process for phytoplankton chlorophyll pigments (Lorenzen, 1967; Vernet, 1991). Since we have no marker stable and specific enough for chlorophyll tetrapyrrolic ring photodegradation, we used the CPPI (Chlorophyll Phytyl side chain Photodegradation Index) for the in-situ determination of the rate of photodegradation of chlorophyll (Cuny et al., 1999). Indeed, the photodegradation of the chlorophyll phytyl side chain produces 3-methylidene-7,11,15-trimethylhexadecan-1,2-diol (phytyldiol), specific of Type II chlorophyll photodegradation and widespread in the environment (Cuny and Rontani, 1999). The CPPI, (phytyldiol : phytol molar ratio) can be linked, through a mathematical model, to the global quantity of photodegraded chlorophyll (Cuny et al., 1999),

The photodegradation rate of chlorophyll fluctuates greatly (Fig. 4b). The 6 March 2012 and 12 March 2013 samples display very low rates (2.6 and 9.8% respectively), in line with our identification of planktonic blooms in March 2012 and 2013. Such blooms result in an increase of fresh chlorophyll inputs, with intact phytyl side chains, and thus in a decrease of CPPI. The dips in chlorophyll photodegradation rates can help us identify blooms, or at least periods when the input of chlorophyll is higher. The summer 2012 samples (July and September) also display a low photodegradation rate ranging from 8.8 to 13.9% while their amount of planktonic sterols increases slightly.

### 3.3.2 $\Delta^5$ -sterols

$\Delta^5$ -sterols possess structural features that can be restricted to a limited number of organisms (Volkman, 1986, 2003). Moreover, biotic and abiotic degradation processes result to specific functionalizations of their cyclic skeleton (De Leeuw and Baas, 1986), which are very useful to estimate the relative importance of these processes (Christodoulou et al., 2009; Rontani et al., 2009). Consequently, degradation products

BGD

11, 14197–14237, 2014

## Survey of composition and degradation state of POM in the Rhone River using lipid tracers

M.-A. Galeron et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





## Survey of composition and degradation state of POM in the Rhone River using lipid tracers

M.-A. Galeron et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

of  $\Delta^5$ -sterols constitute excellent biomarkers for tracing diagenetic transformations of specific organisms (Mackenzie et al., 1982). Using these tracers to evaluate the relative influence of different degradation processes requires that their removal rate (by further degradation) is comparable to that of the parent  $\Delta^5$ -sterol. Although each sterol and its degradation products may be potentially totally mineralized by marine bacteria, we assume that they should exhibit similar reactivity towards bacterial degradation.

Aerobic bacterial hydrogenation may convert  $\Delta^5$ -sterols to  $5\alpha$ (H)-stanols,  $5\alpha$ (H)-stanones and ster-4-en-3-ones (Gagosian et al., 1982; De Leeuw and Baas, 1986; Wakeham, 1989). During the treatment undergone by our samples ( $\text{NaBH}_4$  reduction),  $5\alpha$ (H)-stanones and ster-4-en-3-ones are respectively converted in  $5\alpha$ (H)-stanols and ster-4-en-3-ols; these compounds hence constitute useful markers of bacterial degradation of sterols. To evaluate the proportion of biological degradation of cholesterol, and to better trace human impacts on the OM found in the Rhone, we also included coprostanol and epicoprostanol in its biodegradation products. Sterol biodegradation percentages were estimated using Eqs. (1a) and (1b). Coprostanol ( $5\beta$ (H)-cholestan- $3\beta$ -ol) is a stanol that arises from the anaerobic microbial degradation of cholesterol in the digestive tracts of higher land mammals, including man (Martin et al., 1973). Epicoprostanol ( $5\beta$ (H)-cholestan- $3\alpha$ -ol) is not a major sterol in human faeces, but it is often used as an indicator of sewage treatment (McCalley et al., 1981a).

$$\text{Sitosterol biodegradation \%} = \frac{[\text{sitostanol}]}{[\text{sitosterol}]} \cdot 100 \quad (1a)$$

$$\text{Cholesterol biodegradation \%} = \frac{[\text{cholestanol} + \text{coprostanol} + \text{epicoprostanol}]}{[\text{cholesterol}]} \cdot 100 \quad (1b)$$

Free radical autoxidation yields mainly  $\Delta^5$ - $3\beta$ - $7\alpha/\beta$ -hydroperoxides and in smaller quantities  $5,6$ -epoxysterols and  $3\beta,5\alpha,6\beta$ -trihydroxysterols.  $3\beta,5\alpha,6\beta$ -trihydroxysterols were chosen as tracers of autoxidation (Christodoulou et al., 2009; Rontani et al., 2009) and the sterol autoxidation percentage was estimated using

Eq. (2) based on autoxidation rate constants calculated by Morrissey and Kiely (2006).

$$\text{Sterol autoxidation \%} = \frac{([\Delta^3\beta, 5\alpha, 6\beta\text{-trihydroxysterols}] \cdot 2.4)}{[\text{sitosterol or cholesterol}]} \cdot 100 \quad (2)$$

<sup>1</sup>O<sub>2</sub>-mediated photo-oxidation (Type II photoprocesses) yields mainly  $\Delta^6$ -5 $\alpha$ -hydroperoxides and to a lower extent  $\Delta^4$ -3 $\beta$ -6 $\alpha/\beta$ -hydroperoxides.  $\Delta^6$ -5 $\alpha$ -hydroperoxides are unstable and are converted very easily to the non-specific 7-hydroperoxides, so they were discarded as markers of photo-oxidation. Although produced in lesser amounts,  $\Delta^4$ -3 $\beta$ -6 $\alpha/\beta$ -hydroperoxides, which are relatively stable and highly specific, have been chosen as tracers of photo-oxidation processes and quantified after NaBH<sub>4</sub> reduction to the corresponding diols. The percentage of sterol photo-oxidation was estimated using Eq. (3) (Christodoulou et al., 2009), based on the ratio  $\Delta^4$ -6 $\alpha/\beta$ -hydroperoxides/ $\Delta^6$ -5 $\alpha$ -hydroperoxides found in biological membranes (0.30) (Korytowski et al., 1992).

$$\text{Sterol photo-oxidation \%} = \frac{([\Delta^4\text{-}3\beta\text{-}6\alpha/\beta\text{-dihydroxysterols}] \cdot (1 + 0.3)/0.3)}{[\text{sitosterol or cholesterol}]} \cdot 100 \quad (3)$$

Here, values are expressed in proportions relative to the amount of remaining parent sterol in the sample. A total percentage of over 100% hence only means that degradation products were present in larger quantities than their associated parent sterol.

The results of the evaluation of sitosterol and cholesterol degradation processes are shown in Fig. 5. The most highly degraded samples sitosterol-wise were the ones from 18 July 2011, 26 June 2012 and 22 January 2013. Interestingly, cholesterol degradation shows a completely different trend. When looking at the type of degradation undergone by  $\Delta^5$ -sterols, it also appears clearly that, if auto- and photo-oxidation processes are the major drivers of sitosterol oxidation, biodegradation is the major player in cholesterol degradation. Hedges and Keil (1995) hinted that sterols associated with waxy higher plant material might not be as prone to enzymatic degradation as other sterols,

**Survey of composition and degradation state of POM in the Rhone River using lipid tracers**

M.-A. Galeron et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Survey of composition and degradation state of POM in the Rhone River using lipid tracers**

M.-A. Galeron et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

which would explain why sitosterol is only weakly biodegraded in our samples. Indeed, even though we showed that sitosterol is also produced during spring phytoplanktonic bloom, it is mainly derived from higher plants in most of the samples investigated. The dominance of coprostanol relative to cholestanol attests to a strong contribution of human or animal faeces to the cholesterol present in the Rhone River. Here we used two different ratios in order to better understand of how human activities and waste waters affect the Rhone waters. The first ratio, used to determine if waters are affected by waste water inputs and runoff has been previously used by Writer et al. (1995), and is calculated as follows: coprostanol/(cholesterol + cholestanol). These authors have defined a threshold of 0.06 above which samples are considered affected by wastewater inputs and runoff from pastures and feedlots, and below which they can be considered pristine. The second ratio is the epicoprostanol : coprostanol ratio proposed by Mudge and Seguel (1999), in order to illustrate the level of treatment undergone by wastewaters. The higher the ratio, the more treated the waters are, as epicoprostanol is only present in traces in primary sludge, while it is present in much higher quantities in treated sludge (McCalley et al., 1981a). Both ratios are represented in Fig. 6 and they show that the Rhone is undeniably affected by wastewaters and, to a much lower extent given the historical land use around the Rhone, pasture runoff. However, we noted that both the ratio and the threshold used are unofficial and in no way are proof of a large-scale contamination or pollution. We only highlight here the non-pristine state of the Rhone waters, which is logical given the level of urbanization along the river. The epicoprostanol : coprostanol ratio also shows fluctuations in the level of treatment of wastewaters, and fluctuates with flow rates and precipitation levels.

During senescence, unsaturated higher plant lipids (and notably  $\Delta^5$ -sterols) may be photodegraded (type II photo-oxidation), with chlorophyll acting as a sensitizer (Rontani et al., 1996). Sitosterol present in higher plant phytodetritus should thus have been intensely photodegraded on land. However, the photo-oxidation rate estimates appeared to be relatively low compared to that of autoxidation and biodegradation (Fig. 5a). This is probably due to an intense free-radical-driven breakdown of hydroperoxides pro-

duced during photo-oxidation (Rontani et al., 2003). The photo-oxidation percentages displayed here are thus certainly underestimated.

The presence of large amounts of 24-ethylcholestan-3 $\beta$ ,5 $\alpha$ ,6 $\beta$ -triol in most of the samples indicates that autoxidation plays an important role in the degradation of sitosterol (Fig. 5a). Autoxidation (spontaneous free radical reaction of organic compounds with O<sub>2</sub>), which has been largely ignored until now in the environment, seems to play a key role in the degradation of sitosterol (Fig. 5a) and thus of higher plant material carried by the Rhone River. This assumption was well supported by the detection of significant proportions of compounds deriving from betulin autoxidation (Data not shown). Recently, it has been demonstrated that autoxidation plays a key role in the degradation of terrestrial (Rontani et al., 2014b) and marine (Rontani et al., 2014a) vascular plant debris in seawater. There is clearly a growing body of evidence suggesting that autoxidation reactions can strongly impact the preservation of particulate organic matter in the environment and should be considered carefully alongside other removal processes such as biodegradation when constructing carbon cycles and evaluating carbon budgets. The lowest autoxidation rates observed in samples from 6 March 2012 and 12 March 2013 may be attributed to the phytoplanktonic bloom events, with high inputs of fresh material. While there is variability in the amount and type of degradation undergone by sterols in the sampled particulate matter, it is evident that sitosterol and cholesterol behave very differently when being degraded.

### 3.3.3 Unsaturated fatty acids and cuticular waxes

Fatty-acid sensitivity to photo- and autoxidation is intrinsically linked to their number of double bonds (Frankel, 1998), and we will logically be only looking at unsaturated fatty acids here. Unfortunately, oxidation products of polyunsaturated fatty acids (PUFA) are not stable enough to be used to monitor PUFA degradation. In contrast, photo- and autoxidation products of mono-unsaturated fatty acids (allylic hydroperoxyacids) are much more stable, and can be used (after NaBH<sub>4</sub>-reduction to the corresponding hydroxyacids) as tracers of the abiotic oxidation processes affecting POM (Marchand

BGD

11, 14197–14237, 2014

## Survey of composition and degradation state of POM in the Rhone River using lipid tracers

M.-A. Galeron et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and Rontani, 2001). Free-radical-mediated oxidation (autoxidation) processes can be easily discriminated against photo-oxidation processes thanks to the specific *cis* allylic hydroperoxyacids specifically produced by autoxidative processes (Marchand and Rontani, 2001).

5 Samples only displayed small amounts of oxidation products of oleic acid (not quantified), probably due to the fact that unsaturated fatty acids and their degradation products are very labile and easily metabolized by bacteria (Marchand et al., 2005). Despite this degradation, the profiles obtained by GC-MS (exhibiting relatively high proportions of *cis* oxidation products) allowed us to confirm the important role played by autoxidation in the degradation of POM in the Rhone River.

10 Cutin is present in cuticles covering all aerial parts of higher plants. It is constituted of biopolyesters mainly composed of hydroxy fatty acids. Long-chain *n*-alkanoic,  $\omega$ -hydroxy, dihydroxy, trihydroxy and epoxy-hydroxy acids constitute the major aliphatic monomers (Kolattukudy, 1980a). It was previously demonstrated that Type II photooxidation processes act on some unsaturated cutin monomers such as  $\omega$ -hydroxyoctadec-9-enoic acid ( $\omega$ -hydroxyoleic acid) during the senescence of higher plants (Rontani et al., 2005).  $^1\text{O}_2$  reacts with the carbon-carbon double bond, and leads to the formation of a hydroperoxide at each unsaturated carbon. Due to the involvement of allylic rearrangements, Type II photosensitized oxidation of  $\omega$ -hydroxyoleic acid results (after  $\text{NaBH}_4$ -reduction of hydroperoxides to the corresponding alcohols) in the formation of isomeric allylic 18,(8-11)-dihydroxyoctadecanoic acids, with a *trans* double bond. These compounds constitute interesting specific tracers of higher plant material photo-oxidation. Autoxidation of this compound was never studied, but by analogy with oleic acid oxidation the autoxidative formation of specific *cis* allylic hydroperoxyacids was expected.

25 Significant amounts of allylic 18,(8-11)-dihydroxyoleic acid, with *cis* and *trans* double bonds have been effectively detected in some (not all) samples analyzed attesting to the involvement of auto- and photo-oxidation of higher plant material (Table 2 and Fig. 5). The high proportions of *cis* isomers observed confirmed the domi-

## Survey of composition and degradation state of POM in the Rhone River using lipid tracers

M.-A. Galeron et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

nance of autoxidation processes. Some samples (16 January 2012, 26 June 2012 and 18 July 2011) even displayed larger amounts of oxidation products than  $\omega$ -hydroxyoleic acid, which evidences the importance of degradative processes on this compound. The previously discussed yearly variability in cuticular wax content in our samples explains some of these results.

### 3.3.4 Hydroperoxide stability in SPM

It was previously proposed that photochemically-produced hydroperoxides could induce intense autoxidation processes in the marine environment (Rontani et al., 2014a). Hydroperoxides resulting from photo-oxidation processes may undergo: (i) heterolytic cleavage catalyzed by protons (Frimer, 1979) and (ii) homolytic cleavage induced by transition metal ions (Pokorny, 1987) or UVR (Horspool and Armesto, 1992). Homolytic cleavage of hydroperoxides would lead to the formation of alkoxy radicals, which can then: (i) abstract a hydrogen atom from another molecule to give alcohols, (ii) lose a hydrogen atom to yield ketones, or (iii) undergo  $\beta$ -cleavage reactions affording volatile products. During the  $\text{NaBH}_4$ -reduction, hydroperoxides and ketones were reduced to the corresponding alcohols. The sum of the corresponding hydroperoxides, ketones and alcohols was evaluated under the form of alcohols. Application of a different treatment allowed us to specifically quantify hydroperoxides, alcohols and ketones (remaining in cuticular waxes, phytol, oleic acid, sitosterol and cholesterol oxidation products) (Fig. 7).

Clearly, the proportion of remaining hydroperoxides was highest in the case of sterols, with 49.4 and 31.3 % respectively for 3,6- and 3,7-diols of sitosterol, and 51.5 and 33.5 % for 3,6- and 3,7-diols of cholesterol, against less than 20 % (17.3 %) for cutins, 12.0 % for oleic acid, and 6.6 % for phytol. Standard error was calculated based on all the results obtained (Standard Error = Standard Deviation/ $\sqrt{n}$  for  $n$  samples). These results clearly indicate that despite the involvement of an intense free radical oxidation (autoxidation) inducing homolytic cleavage of peroxy bonds, a significant proportion of hydroperoxides is still intact in POM of the Rhone River. This proportion

## Survey of composition and degradation state of POM in the Rhone River using lipid tracers

M.-A. Galeron et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



reaches 10% of the parent residual compound in the case of sitosterol and 5% in the case of cholesterol. Probably due to high compartmentalization effects, preservation of these compounds seems to be enhanced in higher plant debris. It was recently proposed that homolytic cleavage of photochemically-produced hydroperoxides in riverine POM could be catalyzed by some redox-active metal ions released from SPM in the mixing zone of riverine and marine waters (Rontani et al., 2014b). Due to the presence of significant amounts of hydroperoxides in higher plant residues, the involvement of intensive autoxidation of this material in the Rhone estuary is thus likely.

## 4 Conclusions

Based on a two-year study of the particulate organic matter content of the Rhone River, the main freshwater supplier to the Mediterranean Sea, lipid analyses evidenced a strong contribution from terrestrial vascular plants, most notably gymnosperms. Phytoplanktonic cyclical inputs were also observed and identified as seasonal blooms. The observed amounts of sterols of diatom origin and steep drops in silica observed at the time of the blooms led us to suspect diatoms to be major contributors. Specific compounds like suberin markers, coprostanol and epicoprostanol allowed us to identify anthropic impacts on POM composition (paper mill discharging wastewater and treated waste water discharge).

Using specific lipidic degradation products, we were able to identify for the first time the part that bacterial degradation, autoxidation and photo-oxidation play in organic matter degradation in a Mediterranean river. The study of lipid oxidation products showed that autoxidation, which has been largely underestimated until now in the environment, plays a major role in the degradation of plant-derived organic matter: autoxidation-produced degradation products are nearly half as important as the remaining amount of sitosterol (25% on average across all samples), while biodegradation-produced compounds only represent about a quarter (15% on average across all samples). Photo-oxidation degradation products, although probably underes-

BGD

11, 14197–14237, 2014

### Survey of composition and degradation state of POM in the Rhone River using lipid tracers

M.-A. Galeron et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





## Survey of composition and degradation state of POM in the Rhone River using lipid tracers

M.-A. Galeron et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



5 timated, only stand at 10 % (on average across all samples) of the remaining sitosterol amounts. While autoxidation processes appeared to play only a minor role during the degradation of sitosterol in the cold Mackenzie River (Arctic) (Rontani et al., 2014b), it is worth questioning the role that temperature plays in OM degradation, and wondering  
 10 if it influences one degradation process over the others: do the temperatures found in the Rhone River, warmer than those of the Mackenzie, favor autoxidation? However, the low autoxidation state in the Mackenzie River could also be attributed to the presence of significant proportions of fresh sitosterol-producing phytoplanktonic species (Tolosa et al., 2013). Due to the lack of specificity of this sterol, it is clear that identification of autoxidation products of more specific tracers (such as betulin or amyri-  
 20 n) is absolutely necessary to monitor the degradation of higher plant material in rivers and oceans more precisely.

We also noted that cholesterol offers a very different degradation profile, undergoing little autoxidation, but mainly influenced by biotic degradation processes. Cholesterol is non-specific and can come from a number of sources, which is why it is not used as a tracer of the origin or the general degradation state of POM. However, its bacterial degradation products, including coprostanol and epicoprostanol, are good markers of sewage contamination and sewage treatment, and clearly indicated here that the Rhone waters were affected by wastewaters.

25 Finally, we have shown that even through the degradative processes impacting the TPOM flowing down the Rhone, there is still an important quantity of hydroperoxides remaining in the OM, which in turn should favor autoxidation upon the OM arrival at sea.

*Acknowledgements.* This work is a contribution to the Labex OT-Med (n° ANR-11-LABX-0061) funded by the French Government “Investissements d’Avenir” program of the French National Research Agency (ANR) through the A\*MIDEX project (n° ANR-11-IDEX-0001-02). It was also supported by the LEFE-CYBER (Les Enveloppes Fluides et l’Environnement) national program, as part of the MORTIMER (Matière ORganique Terrestre rejetée par les fleuves et les rivières en MER) research program. It was undergone as part of the transversal research axis DEBAT  
 30 of the Mediterranean Institute of Oceanography, Marseille, France.

Additional data provided by “MOOSE” (Mediterranean Oceanic Observing System for the Environment) with the support of the “Agence de l’Eau Rhone-Méditerranée-Corse”. Special thanks to Michel Fornier for providing samples

## References

- 5 AGRESTE: En Provence-Alpes-Côte d’Azur, la Forêt est Sous-Exploitée, Rapport du Ministère de l’Agriculture, de l’Alimentation, de la Pêche, de la Ruralité et de l’Aménagement du Territoire, 2011.
- Amelung, W., Brodowski, S., Sandhage-Hofmann, A., and Bol, R.: Combining biomarker with stable isotope analyses for assessing the transformation and turnover of soil organic matter, in: *Advances in Agronomy*, edited by: Sparks, D. L., Academic Press, chap. 6, 155–250, 10 2008.
- Aminot, A. and Kérouel, R.: *Dosage Automatique des Nutriments dans les Eaux Marines: Méthodes en Flux Continu*, Editions Quae, 2007.
- Atlas, R. M. and Bartha, R.: Hydrocarbon biodegradation and oil spill bioremediation, in: *Advances in Microbial Ecology*, edited by: Marshall, K. C., Springer, US, 287–338, 1992.
- 15 Bourbonniere, R. A. and Meyers, P. A.: Sedimentary geolipid records of historical changes in the watersheds and productivities of Lakes Ontario and Erie, *Limnol. Oceanogr.*, 41, 352–359, 1996.
- Bourgeois, S., Pruski, A. M., Sun, M.-Y., Buscail, R., Lantoiné, F., Kerhervé, P., Vétion, G., Rivière, B., and Charles, F.: Distribution and lability of land-derived organic matter in the surface sediments of the Rhône prodelta and the adjacent shelf (Mediterranean Sea, France): a multi proxy study, *Biogeosciences*, 8, 3107–3125, doi:10.5194/bg-8-3107-2011, 2011.
- 20 Brown, R. C. and Wade, T. L.: Sedimentary coprostanol and hydrocarbon distribution adjacent to a sewage outfall, *Water Res.*, 18, 621–632, 1984.
- 25 Cauwet, G., Gadel, F., de Souza Sierra, M. M., Donard, O., and Ewald, M.: Contribution of the Rhône River to organic carbon inputs to the northwestern Mediterranean Sea, *Cont. Shelf Res.*, 10, 1025–1037, 1990.
- Cloern, J. E., Canuel, E. A., and Harris, D.: Stable carbon and nitrogen isotope composition of aquatic and terrestrial plants of the San Francisco Bay estuarine system, *Limnol. Oceanogr.*, 30 47, 713–729, 2002.

## Survey of composition and degradation state of POM in the Rhone River using lipid tracers

M.-A. Galeron et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



**Survey of composition and degradation state of POM in the Rhone River using lipid tracers**

M.-A. Galeron et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Christodoulou, S., Marty, J.-C., Miquel, J.-C., Volkman, J. K., and Rontani, J.-F.: Use of lipids and their degradation products as biomarkers for carbon cycling in the northwestern Mediterranean Sea, *Mar. Chem.*, 113, 25–40, 2009.
- Conner, A. H., Nagasampagi, B. A., and Rowe, J. W.: Terpenoid and other extractives of western white pine bark, *Phytochemistry*, 19, 1121–1131, 1980.
- Cuny, P. and Rontani J.-F.: On the widespread occurrence of 3-methylidene-7,11,15-trimethylhexadecan-1,2-diol in the marine environment: a specific isoprenoid marker of chlorophyll photodegradation, *Mar. Chem.*, 65, 155–165, 1999.
- Cuny, P., Romano, J.-C., Beker, B., and Rontani, J.-F.: Comparison of the photodegradation rates of chlorophyll chlorin ring and phytol side chain in phytodetritus: is the phytyldiol versus phytol ratio (CPPI) a new biogeochemical index?, *J. Exp. Mar. Biol. Ecol.*, 237, 271–290, 1999.
- Dachs, J., Bayona, J. M., Fowler, S. W., Miquel, J.-C., and Albaigés, J.: Evidence for cyanobacterial inputs and heterotrophic alteration of lipids in sinking particles in the Alboran Sea (SW Mediterranean), *Mar. Chem.*, 60, 189–201, 1998.
- De Leeuw, J. W. and Baas, M.: Early-stage diagenesis of steroids, *Meth. Geoch.*, 24, 101–123, 1986.
- De Leeuw, J. W. and Largeau, C.: A review of macromolecular organic compounds that comprise living organisms and their role in kerogen, coal, and petroleum formation, in: *Organic Geochemistry*, edited by: Engel, M. H. and Macko, S. A., Springer, US, 23–72, 1993.
- Diefendorf, A. F., Freeman, K. H., Wing, S. L., and Graham, H. V.: Production of *n*-alkyl lipids in living plants and implications for the geologic past, *Geochim. Cosmochim. Ac.*, 75, 7472–7485, 2011.
- Eyrolle, F., Radakovitch, O., Raimbault, P., Charmasson, S., Antonelli, C., Ferrand, E., Aubert, D., Raccasi, G., Jacquet, S., and Gurriaran, R.: Consequences of hydrological events on the delivery of suspended sediment and associated radionuclides from the Rhône River to the Mediterranean Sea, *J. Soils Sediments*, 12, 1479–1495, 2012.
- Fine, P. M., Cass, G. R., and Simoneit, B. R. T.: Chemical characterization of fine particle emissions from fireplace combustion of woods grown in the Northeastern United States, *Environ. Sci. Technol.*, 35, 2665–2675, 2001.
- Foote, C. S.: Photosensitized oxidation and singlet oxygen: consequences in biological systems, *Free Radicals in Biology*, 2, 85–133, 1976.
- Frankel, E. N.: *Lipid Oxidation*, The Oily, Dundee, Scotland, 129–160, 1998.

**Survey of composition and degradation state of POM in the Rhone River using lipid tracers**

M.-A. Galeron et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Frimer, A. A.: The reaction of singlet oxygen with olefins: the question of mechanism, *Chem. Rev.*, **79**, 359–387, 1979.
- Fruget, J.-F., Centofanti, M., Dessaix, J., Olivier, J.-M., Druart, J.-C., and Martinez, P.-J.: Temporal and spatial dynamics in large rivers: example of a long-term monitoring of the middle Rhone River, *Ann. Limnol.-Int. J. Lim.*, **37**, 237–251, 2001.
- Gagosian, R. B., Zafiriou, O. C., Peltzer, E. T., and Alford, J. B.: Lipids in aerosols from the tropical North Pacific: temporal variability, *J. Geophys. Res.*, **87**, 11133–11144, 1982.
- Gagosian, R. B., Nigrelli, G. E., and Volkman, J. K.: Vertical transport and transformation of biogenic organic compounds from a sediment trap experiment off the coast of Peru, in: *Coastal Upwelling its Sediment Record*, edited by: Suess, E. and Thiede, J., Springer, US, 241–272, 1983.
- Hedges, J. I. and Keil, R. G.: Sedimentary organic matter preservation: an assessment and speculative synthesis, *Mar. Chem.*, **49**, 81–115, 1995.
- Hedges, J. I., Keil, R. G., and Benner, R.: What happens to terrestrial organic matter in the ocean?, *Org. Geochem.*, **27**, 195–212, 1997.
- Horspool, W. and Armesto, D.: *Organic Photochemistry: a Comprehensive Treatment*, Ellis Horwood, 1992.
- Kolattukudy, P. E.: Cutin, suberin, and waxes, in: *Biochemistry of Plants: a Comprehensive Treatise*, 1980a.
- Korytowski, W., Bachowski, G. J., and Girotti, A. W.: Photoperoxidation of cholesterol in homogeneous solution, isolated membranes, and cells: comparison of the 5 $\alpha$ , and 6 $\beta$ -hydroperoxides as indicators of singlet oxygen intermediacy, *Photochem. Photobiol.*, **56**, 1–8, 1992.
- Kunst, L. and Samuels, A. L.: Biosynthesis and secretion of plant cuticular wax, *Prog. Lipid Res.*, **42**, 51–80, 2003.
- Liu, J.: Pharmacology of oleanolic acid and ursolic acid, *J. Ethnopharmacol.*, **49**, 57–68, 1995.
- Lorenzen, C. J.: Vertical distribution of chlorophyll and phaeo-pigments?: Baja California, *Deep Sea Research and Oceanographic Abstracts*, **14**, 735–745, 1967.
- Ludwig, W., Dumont, E., Meybeck, M., and Heussner, S.: River discharges of water and nutrients to the Mediterranean and Black Sea: major drivers for ecosystem changes during past and future decades?, *Prog. Oceanogr.*, **80**, 199–217, 2009.
- Mackenzie, A. S., Brassell, S. C., Eglinton, G., and Maxwell, J. R.: Chemical fossils: the geological fate of steroids, *Science*, **217**, 491–504, 1982.

---

**Survey of composition and degradation state of POM in the Rhone River using lipid tracers**

M.-A. Galeron et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Marchand, D. and Rontani, J.-F.: Characterisation of photo-oxidation and autoxidation products of phytoplanktonic monounsaturated fatty acids in marine particulate matter and recent sediments, *Org. Geochem.*, 32, 287–304, 2001.
- Marchand, D. and Rontani, J.-F.: Visible light-induced oxidation of lipid components of purple sulfur bacteria: a significant process in microbial mats, *Org. Geochem.*, 34, 61–79, 2003.
- Marchand, D., Marty, J.-C., Miquel, J.-C., and Rontani, J.-F.: Lipids and their oxidation products as biomarkers for carbon cycling in the northwestern Mediterranean Sea: results from a sediment trap study, *Mar. Chem.*, 95, 129–147, 2005.
- Martin, W. J., Subbiah, M. T. R., Kottke, B. A., Birk, C. C., and Naylor, M. C.: Nature of fecal sterols and intestinal bacterial flora, *Lipids*, 8, 208–215, 1973.
- McCalley, D. V., Cooke, M., and Nickless, G.: Effect of sewage treatment on faecal sterols, *Water Res.*, 15, 1019–1025, 1981.
- McKee, B. A., Aller, R. C., Allison, M. A., Bianchi, T. S., and Kineke, G. C.: Transport and transformation of dissolved and particulate materials on continental margins influenced by major rivers: benthic boundary layer and seabed processes, *Cont. Shelf. Res.*, 24, 899–926, 2004.
- Mendez-Millan, M., Dignac, M.-F., Rumpel, C., Rasse, D. P., and Derenne, S.: Dynamics of shoot vs. root C assessed by natural <sup>13</sup>C abundance of their biomarkers, in: EGU General Assembly Conference Abstracts, 9968 pp., 2010.
- Meyers, P. A. and Ishiwatari, R.: Lacustrine organic geochemistry – an overview of indicators of organic matter sources and diagenesis in lake sediments, *Org. Geochem.*, 20, 867–900, 1993.
- Mihara, S. and Tateba, H.: Photosensitized oxygenation reactions of phytol and its derivatives, *J. Org. Chem.*, 51, 1142–1144, 1986.
- Morrissey, P. A. and Kiely, M.: Oxysterols: formation and biological function, in: *Advanced Dairy Chemistry*, vol. 2: Lipids, edited by: Fox, P. F. and McSweeney, P. L. H., Springer, US, 641–674, 2006.
- Mudge, S. M. and Seguel, C. G.: Organic contamination of San Vicente Bay, Chile, *Mar. Pollut. Bull.*, 38, 1011–1021, 1999.
- Mueller, K. E., Polissar, P. J., Oleksyn, J., and Freeman, K. H.: Differentiating temperate tree species and their organs using lipid biomarkers in leaves, roots and soil, *Org. Geochem.*, 52, 130–141, 2012.

---

**Survey of composition and degradation state of POM in the Rhone River using lipid tracers**

M.-A. Galeron et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Nelson, J. R.: Rates and possible mechanism of light-dependent degradation of pigments in detritus derived from phytoplankton, *J. Mar. Res.*, 51, 155–179, 1993.
- Nierop, K. G. J.: Origin of aliphatic compounds in a forest soil, *Org. Geochem.*, 29, 1009–1016, 1998.
- 5 Nishimura, M. and Koyama, T.: The occurrence of stanols in various living organisms and the behavior of sterols in contemporary sediments, *Geochim. Cosmochim. Ac.*, 41, 379–385, 1977.
- Novak, J. T. and Carlson, D. A.: The kinetics of anaerobic long chain fatty acid degradation, *J. Water Pollut. Con. F.*, 42, 1932–1943, 1970.
- 10 Otto, A., Shunthirasingham, C., and Simpson, M. J.: A comparison of plant and microbial biomarkers in grassland soils from the Prairie Ecozone of Canada, *Org. Geochem.*, 36, 425–448, 2005.
- Paper Excellence: Fibre Excellence Tarascon, available at: <http://www.paperexcellence.com/mills/fibre-excellence-tarascon/> (last access: 14 April 2014), 2014.
- 15 Pokorny, J.: Major factors affecting the autoxidation of lipids, in: *Autoxidation of Unsaturated Lipids*, 141–206, 1987.
- Ponomarenko, L. P., Stonik, I. V., Aizdaicher, N. A., Orlova, T. Y., Popovskaya, G. I., Pomazkina, G. V., and Stonik, V. A.: Sterols of marine microalgae *Pyramimonas cf. cordata* (Prasinophyta), *Attheya ussurensis* sp. nov. (Bacillariophyta) and a spring diatom bloom from Lake Baikal, *Comp. Biochem. Phys. B*, 138, 65–70, 2004.
- 20 Raimbault, P., Lantoine, F., and Neveux, J.: Dosage rapide de la chlorophylle *a* et des phéopigments a par fluorimétrie après extraction au méthanol, comparaison avec la méthode classique d'extraction à l'acétone, *Océanis S. D.*, 30, 189–205, 2004.
- Rampen, S. W., Abbas, B. A., Schouten, S., and Damsté, J. S. S.: A comprehensive study of sterols in marine diatoms (Bacillariophyta): Implications for their use as tracers for diatom productivity, *Limnol. Oceanogr.*, 55, 91–105, 2010.
- 25 Razboršek, M. I., Voncina, D. B., Dolecek, V., and Voncina, E.: Determination of oleanolic, betulinic and ursolic acid in lamiaceae and mass spectral fragmentation of their trimethylsilylated derivatives, *Chroma*, 67, 433–440, 2008.
- 30 Rontani, J.-F.: Photo-and free radical-mediated oxidation of lipid components during the senescence of phototrophic organisms, *Senescence*, Intech, Rijeka, 3–31, 2012.
- Rontani, J.-F., Cuny, P., and Grossi, V.: Photodegradation of chlorophyll phytyl chain in senescent leaves of higher plants, *Phytochemistry*, 42, 347–351, 1996.

**Survey of composition and degradation state of POM in the Rhone River using lipid tracers**

M.-A. Galeron et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Rontani, J.-F., Rabourdin, A., Marchand, D., and Aubert, C.: Photochemical oxidation and autoxidation of chlorophyll phytyl side chain in senescent phytoplanktonic cells: potential sources of several acyclic isoprenoid compounds in the marine environment, *Lipids*, 38, 241–254, 2003.

5 Rontani, J.-F., Rabourdin, A., Pinot, F., Kandel, S., and Aubert, C.: Visible light-induced oxidation of unsaturated components of cutins: a significant process during the senescence of higher plants, *Phytochemistry*, 66, 313–321, 2005.

Rontani, J.-F., Zabeti, N., and Wakeham, S. G.: The fate of marine lipids: biotic vs. abiotic degradation of particulate sterols and alkenones in the Northwestern Mediterranean Sea, *Mar. Chem.*, 113, 9–18, 2009.

10 Rontani, J.-F., Zabeti, N., and Wakeham, S. G.: Degradation of particulate organic matter in the equatorial Pacific Ocean: biotic or abiotic?, *Limnol. Oceanogr.*, 56, 333–349, 2011.

Rontani, J.-F., Vaultier, F., and Bonin, P.: Biotic and abiotic degradation of marine and terrestrial higher plant material in intertidal surface sediments from Arcachon Bay (France): a lipid approach, *Mar. Chem.*, 158, 69–79, 2014a.

15 Rontani, J.-F., Charrière, B., Sempéré, R., Doxaran, D., Vaultier, F., Vonk, J. E., and Volkman, J. K.: Degradation of sterols and terrigenous organic matter in waters of the Mackenzie Shelf, Canadian Arctic, *Org. Geochem.*, 75, 61–73, 2014b.

Sempéré, R., Charrière, B., Van Wambeke, F., and Cauwet, G.: Carbon inputs of the Rhône River to the Mediterranean Sea: biogeochemical implications, *Global Biogeochem. Cy.*, 14, 669–681, 2000.

20 Sicre, M.-A., Paillasseur, J.-L., Marty, J.-C., and Saliot, A.: Characterization of seawater samples using chemometric methods applied to biomarker fatty acids, *Org. Geochem.*, 12, 281–288, 1988.

25 Sicre, M.-A., Tian, R. C., Broyelle, I., and Saliot, A.: Aquatic distribution of 4-desmethyl sterols in the Chang Jiang Estuary, China, *Mar. Chem.*, 42, 11–24, 1993.

Simpson, M. J., Otto, A., and Feng, X.: Comparison of solid-state carbon-13 nuclear magnetic resonance and organic matter biomarkers for assessing soil organic matter degradation, *Soil Sci. Soc. Am. J.*, 72, 268–276, 2008.

30 Spikes, J. D. and Straight, R.: Sensitized photochemical processes in biological systems, *Annu. Rev. Phys. Chem.*, 18, 409–436, 1967.

Tolosa, I., Fiorini, S., Gasser, B., Martín, J., and Miquel, J. C.: Carbon sources in suspended particles and surface sediments from the Beaufort Sea revealed by molecular lipid biomarkers



## Survey of composition and degradation state of POM in the Rhone River using lipid tracers

M.-A. Galeron et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

and compound-specific isotope analysis, Biogeosciences, 10, 2061–2087, doi:10.5194/bg-10-2061-2013, 2013.

Van Dongen, B. E., Zencak, Z., and Gustafsson, Ö: Differential transport and degradation of bulk organic carbon and specific terrestrial biomarkers in the surface waters of a sub-arctic brackish bay mixing zone, Mar. Chem., 112, 203–214, 2008.

Vernet, M.: Phytoplankton dynamics in the Barents Sea estimated from chlorophyll budget models, Polar Res., 10, 129–146, 1991.

Vivian, H.: Hydrological changes of the Rhône River, in: Historical Change of Large Alluvial Rivers: Western Europe, John Wiley & Sons, New York, 57–77, 1989.

Volkman, J. K.: A review of sterol markers for marine and terrigenous organic matter, Org. Geochem., 9, 83–99, 1986.

Volkman, J. K.: Sterols in microorganisms, Appl. Microbiol. Biot., 60, 495–506, 2003.

Volkman, J. K., Smith, D. J., Eglinton, G., Forsberg, T. E. V., and Corner, E. D. S.: Sterol and fatty acid composition of four marine haptophycean algae, J. Mar. Biol. Assoc. UK, 61, 509–527, 1981.

Wakeham, S. G.: Reduction of stenols to stanols in particulate matter at oxic–anoxic boundaries in sea water, Nature, 342, 787–790, 1989.

Wakeham, S. G.: Lipid biomarkers for heterotrophic alteration of suspended particulate organic matter in oxygenated and anoxic water columns of the ocean, Deep-Sea Res. Pt. I, 42, 1749–1771, 1995.

Wakeham, S. G. and Canuel, E. A.: Degradation and preservation of organic matter in marine sediments, in: Marine Organic Matter: Biomarkers, Isotopes and DNA, Volkman, J. K., Springer, Berlin, Heidelberg, 295–321, 2006.

Writer, J. H., Leenheer, J. A., Barber, L. B., Amy, G. L., and Chapra, S. C.: Sewage contamination in the upper Mississippi River as measured by the fecal sterol, coprostanol, Water Res., 29, 1427–1436, 1995.

**Table 1.** Percentages of sterols (relative to the sum of sterols quantified), and sitosterol : campesterol ratio in the samples investigated.

Date	Sito-sterol	Chole-sterol	Desmo-sterol	Brassica-sterol	Methylene-Cholesterol	Campe-sterol	Sitosterol : Campesterol ratio
5 Apr 2011	74.4	12.7	0.4	4.9	1.2	6.4	11.6
18 Jul 2011	21.2	45.5	2.6	16.1	3.3	11.3	1.9
4 Nov 2011	54.1	23.8	1.9	10.0	2.2	8.0	6.7
5 Nov 2011	60.2	21.4	1.2	7.7	1.7	7.9	7.6
7 Nov 2011	56.7	28.6	0.4	6.1	1.3	6.9	8.2
14 Nov 2011	34.6	45.7	0.9	10.0	1.5	7.4	4.7
19 Dec 2011	52.5	29.1	0.5	7.5	1.9	8.5	6.2
16 Jan 2012	39.2	43.6	1.3	6.7	1.8	7.4	5.3
6 Mar 2012	17.6	12.4	28.8	6.2	22.6	12.4	1.4
17 Apr 2012	36.5	39.3	4.7	8.2	2.9	8.4	4.4
2 May 2012	38.4	41.4	2.1	9.5	1.9	6.7	5.7
22 May 2012	32.5	43.6	4.4	10.3	2.6	6.5	5.0
11 Jun 2012	35.0	43.5	2.5	9.6	1.6	7.7	4.5
26 Jun 2012	24.7	54.9	2.6	10.0	1.8	6.0	4.1
25 Jul 2012	21.6	37.5	3.9	20.8	4.4	11.7	1.8
5 Sep 2012	22.6	46.6	3.4	15.4	3.3	8.6	2.6
19 Sep 2012	21.1	44.3	3.8	19.1	3.0	8.7	2.4
3 Oct 2012	27.4	46.7	2.3	11.9	1.9	9.8	2.8
16 Oct 2012	42.9	38.9	1.4	7.8	1.4	7.6	5.7
6 Nov 2012	44.9	36.9	2.0	7.3	1.8	7.1	6.3
17 Dec 2012	66.3	22.8	1.0	3.5	1.2	5.2	12.7
10 Jan 2013	46.9	34.8	2.5	7.1	1.5	7.2	6.5
22 Jan 2013	36.0	48.4	1.4	6.9	1.3	6.1	5.9
4 Feb 2013	50.0	39.2	0.8	4.0	0.8	5.1	9.8
13 Feb 2013	55.4	30.7	2.0	4.8	1.3	5.9	9.4
12 Mar 2013	35.1	27.6	14.5	9.0	7.6	6.2	5.7
21 Mar 2013	37.4	29.0	11.1	8.9	6.6	7.2	5.2
17 Apr 2013	35.6	45.6	2.6	7.0	2.5	6.7	5.3
2 May 2013	59.0	24.5	1.1	6.8	1.7	6.8	8.7
13 May 2013	49.7	38.6	0.0	4.6	1.1	6.1	8.2
Average	41.0	35.9	3.6	8.9	3.0	7.6	5.9
Standard deviation	14.6	10.7	5.6	4.2	4.0	1.8	2.8

## Survey of composition and degradation state of POM in the Rhone River using lipid tracers

M.-A. Galeron et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Table 2a.** Fatty acid content of the different samples ( $\mu\text{g mg}^{-1}$  (dw)) (PUFA: poly-unsaturated fatty acids).



	2011						
	5 Apr	18 Jul	4 Nov	5 Nov	7 Nov	14 Nov	19 Dec
C <sub>14:0</sub>	0.34	0.83	0.41	0.08	0.01	0.19	0.23
C <sub>15:0</sub>	0.20	0.20	0.16	0.06	0.01	0.10	0.08
C <sub>15:1</sub>	–	–	–	–	–	0.03	–
Iso-C <sub>15:0</sub>	0.16	0.42	0.21	0.04	–	0.14	0.10
Anté-iso-C <sub>15:0</sub>	0.11	0.20	0.16	0.03	–	0.10	0.10
C <sub>16:0</sub>	3.11	4.26	3.25	1.35	0.25	2.16	1.90
C <sub>16:1</sub>	0.42	1.45	0.94	0.13	0.01	0.64	0.34
C <sub>16:2</sub>	–	–	–	–	–	–	–
C <sub>16:3</sub>	–	0.04	–	–	–	–	–
C <sub>16:4</sub>	–	–	–	–	–	–	–
C <sub>17:0</sub>	0.07	0.10	0.06	0.02	0.01	0.05	0.03
C <sub>18:0</sub>	0.71	0.90	0.57	0.29	0.13	0.51	0.46
C <sub>18:1<math>\omega</math>9</sub> (Oleic)	0.62	0.66	0.81	0.02	0.04	0.29	0.41
C <sub>18:1<math>\omega</math>7</sub> (Vaccenic)	0.25	0.49	0.35	0.09	0.03	0.24	0.26
C <sub>18:2</sub>	0.25	0.12	0.31	–	0.01	0.05	0.15
C <sub>18:3</sub>	–	–	–	–	–	–	–
C <sub>18:4</sub>	–	–	–	–	–	–	–
C <sub>20:0</sub>	0.16	0.09	0.15	0.03	0.02	0.04	0.07
C <sub>20:5</sub>	0.93	0.06	0.06	0.17	0.07	–	0.20
C <sub>22:0</sub>	0.23	0.08	0.15	0.03	0.02	0.03	0.10
C <sub>22:1</sub>	–	–	–	–	–	–	–
C <sub>22:2</sub>	–	–	–	–	–	–	0.01
C <sub>24:0</sub>	0.08	0.01	0.03	0.01	0.01	0.01	0.04
C <sub>26:0</sub>	–	–	–	–	–	–	0.01
$\omega$ -hydroxy-C <sub>16:0</sub>	0.34	0.83	0.41	0.08	0.01	0.19	0.23
$\omega$ -hydroxy-C <sub>18:1<math>\omega</math>9</sub>	0.20	0.20	0.16	0.06	0.01	0.10	0.08
18,(8-11)-dihydroxy-C <sub>16:0</sub>	–	–	–	–	–	0.03	–
$\omega$ -C <sub>16:0</sub> diacid	0.16	0.42	0.21	0.04	–	0.14	0.10
$\omega$ -hydroxy-C <sub>22:0</sub>	0.11	0.20	0.16	0.03	–	0.10	0.10
% PUFA	15.5	2.28	4.89	7.15	13.22	1.25	7.83

**Survey of composition and degradation state of POM in the Rhone River using lipid tracers**

M.-A. Galeron et al.

[Title Page](#)

[Abstract](#)   [Introduction](#)

[Conclusions](#)   [References](#)

[Tables](#)   [Figures](#)

[◀](#)   [▶](#)

[◀](#)   [▶](#)

[Back](#)   [Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



**Table 2b.** Fatty acid content of the different samples ( $\mu\text{g mg}^{-1}$  (dw)) (PUFA: poly-unsaturated fatty acids).

	2012													
	16 Jan	6 Mar	17 Apr	2 May	22 May	11 Jun	26 Jun	25 Jul	5 Sep	19 Sep	3 Oct	16 Oct	6 Nov	17 Dec
C <sub>14:0</sub>	0.39	14.97	0.47	0.05	0.06	0.10	0.11	0.08	0.53	0.06	0.03	0.04	0.02	0.02
C <sub>15:0</sub>	0.13	0.49	0.15	0.04	0.03	0.06	0.04	0.04	0.09	0.03	0.02	0.04	0.02	0.02
C <sub>15:1</sub>	–	0.07	–	–	–	0.01	–	–	–	–	–	–	–	–
Iso-C <sub>15:0</sub>	0.20	0.10	0.11	0.02	0.01	0.04	0.02	0.02	0.19	0.02	0.02	0.03	–	–
Anté-iso-C <sub>15:0</sub>	0.19	0.08	0.09	0.02	0.01	0.03	0.02	0.02	0.09	0.02	0.02	0.03	0.01	–
C <sub>16:0</sub>	2.97	23.54	3.20	1.10	1.18	1.52	1.17	1.71	3.02	1.82	0.13	0.15	0.80	0.83
C <sub>16:1</sub>	0.69	30.53	0.82	0.13	0.16	0.14	0.13	0.42	0.84	0.34	1.44	0.20	0.04	0.02
C <sub>16:2</sub>	–	7.98	–	–	–	–	–	0.09	0.17	–	–	–	–	–
C <sub>16:3</sub>	–	15.41	–	–	–	–	–	–	0.18	–	–	–	–	–
C <sub>16:4</sub>	–	5.39	0.04	–	–	–	–	–	–	–	–	–	–	–
C <sub>17:0</sub>	0.05	–	0.04	0.01	0.02	0.02	0.01	0.03	0.04	0.03	0.04	0.04	0.01	0.01
C <sub>18:0</sub>	0.79	1.55	0.73	0.25	0.53	0.39	0.24	0.40	0.45	0.50	0.41	0.63	0.39	0.33
C <sub>18:1</sub> (Oleic)	0.53	1.20	0.55	0.12	0.25	0.14	0.01	0.44	0.42	0.55	0.27	0.35	0.16	0.10
C <sub>18:1</sub> ω7 (Vaccenic)	0.27	–	0.25	0.06	0.16	0.10	0.04	0.24	0.21	0.26	0.19	0.30	0.08	0.04
C <sub>18:2</sub>	0.11	–	0.15	0.02	0.05	0.02	–	0.08	0.06	0.11	0.06	0.07	0.03	0.18
C <sub>18:3</sub>	–	–	0.11	–	–	–	–	–	–	–	–	–	–	–
C <sub>18:4</sub>	–	7.18	–	–	–	–	–	–	–	–	–	–	–	–
C <sub>20:0</sub>	0.12	0.08	0.06	0.02	0.04	0.03	–	0.03	0.03	0.04	0.03	0.99	0.03	0.03
C <sub>20:5</sub>	–	22.69	0.06	–	0.05	0.05	0.01	0.08	0.08	0.11	–	–	–	–
C <sub>22:0</sub>	0.01	0.06	0.04	0.02	0.04	0.03	0.01	0.02	0.08	0.04	0.02	0.10	–	0.03
C <sub>22:1</sub>	–	0.13	–	–	–	–	–	–	–	–	–	–	–	–
C <sub>23:0</sub>	–	–	–	–	–	–	–	–	–	–	–	–	–	–
C <sub>24:0</sub>	–	–	0.01	0.01	0.01	0.01	–	0.01	–	0.01	–	0.03	0.01	–
C <sub>26:0</sub>	–	–	–	–	–	–	–	–	–	–	–	–	–	–
ω-hydroxy-C <sub>16:0</sub>	0.39	14.97	0.47	0.05	0.06	0.10	0.11	0.08	0.53	0.06	0.03	0.04	0.02	0.02
ω-hydroxy-C <sub>18:1</sub> ω9	0.13	0.49	0.15	0.04	0.03	0.06	0.04	0.04	0.09	0.03	0.02	0.04	0.02	0.02
18,(8-11)-dihydroxy-C <sub>16:0</sub>	–	0.07	–	–	–	0.01	–	–	–	–	–	–	–	–
ω-C <sub>16:0</sub> diacid	0.20	0.10	0.11	0.02	0.01	0.04	0.02	0.02	0.19	0.02	0.02	0.03	–	–
ω-hydroxy-C <sub>22:0</sub>	0.19	0.08	0.09	0.02	0.01	0.03	0.02	0.02	0.09	0.02	0.02	0.03	0.01	–
% PUFA	1.73	44.62	5.39	0.87	4.12	2.62	0.43	6.77	7.69	5.57	2.08	2.26	1.95	11.36

## Survey of composition and degradation state of POM in the Rhone River using lipid tracers

M.-A. Galeron et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Table 2c.** Fatty acid content of the different samples ( $\mu\text{g mg}^{-1}$  (dw)) (PUFA: poly-unsaturated fatty acids).

	10 Jan	22 Jan	4 Feb	13 Feb	2013 12 Mar	21 Mar	17 Apr	2 May	13 May
C <sub>14:0</sub>	0.29	0.03	0.03	0.03	0.39	0.86	0.28	0.07	0.03
C <sub>15:0</sub>	0.14	0.03	0.02	0.02	0.08	0.14	0.13	0.05	0.02
C <sub>15:1</sub>	–	–	–	–	–	0.02	–	–	–
Iso-C <sub>15:0</sub>	0.12	–	–	0.01	0.02	0.08	0.09	0.03	–
Anté-iso-C <sub>15:0</sub>	0.11	0.01	–	0.01	0.03	0.08	0.08	0.03	–
C <sub>16:0</sub>	3.39	0.93	1.11	0.73	2.89	4.45	3.08	2.32	0.46
C <sub>16:1</sub>	0.52	0.06	0.05	0.06	1.60	2.22	0.67	0.17	0.01
C <sub>16:2</sub>	–	–	–	–	–	0.53	0.06	–	–
C <sub>16:3</sub>	–	–	–	–	0.90	0.99	0.09	–	–
C <sub>16:4</sub>	–	–	–	–	0.51	0.50	–	–	–
C <sub>17:0</sub>	0.06	0.01	0.02	0.01	0.03	0.06	0.07	–	–
C <sub>18:0</sub>	1.23	0.24	0.44	0.25	0.63	1.19	1.01	0.79	0.08
C <sub>18:1<math>\omega</math>9</sub> (Oleic)	0.19	0.07	0.26	0.19	0.85	1.48	0.91	0.46	0.02
C <sub>18:1<math>\omega</math>7</sub> (Vaccenic)	0.79	0.03	0.07	0.06	0.22	0.52	0.36	0.14	0.01
C <sub>18:2</sub>	0.34	0.01	0.03	0.03	0.22	0.44	0.19	–	–
C <sub>18:3</sub>	–	–	–	–	–	–	–	–	–
C <sub>18:4</sub>	–	–	–	–	0.61	0.58	–	–	–
C <sub>20:0</sub>	0.12	0.01	0.01	0.01	0.05	0.12	0.12	0.07	–
C <sub>20:5</sub>	–	–	–	–	2.27	1.98	0.30	–	–
C <sub>22:0</sub>	0.15	0.01	0.01	0.01	0.03	0.10	0.14	0.10	–
C <sub>22:1</sub>	–	–	–	–	–	–	–	–	–
C <sub>23:0</sub>	–	–	–	–	–	–	–	–	–
C <sub>24:0</sub>	0.04	–	–	–	0.01	0.02	0.04	0.02	–
C <sub>26:0</sub>	–	–	–	–	–	–	–	–	–
$\omega$ -hydroxy-C <sub>16:0</sub>	0.29	0.03	0.03	0.03	0.39	0.86	0.28	0.07	0.03
$\omega$ -hydroxy-C <sub>18:1<math>\omega</math>9</sub>	0.14	0.03	0.02	0.02	0.08	0.14	0.13	0.05	0.02
18,(8-11)-dihydroxy-C <sub>16:0</sub>	–	–	–	–	–	0.02	–	–	–
$\omega$ -C <sub>16:0</sub> diacid	0.12	–	–	0.01	0.02	0.08	0.09	0.03	–
$\omega$ -hydroxy-C <sub>22:0</sub>	0.11	0.01	–	0.01	0.03	0.08	0.08	0.03	–
% PUFA	4.49	0.85	1.61	2.37	39.73	30.67	8.38	0.00	0.00

## Survey of composition and degradation state of POM in the Rhone River using lipid tracers

M.-A. Galeron et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶


Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Figure 1.** Rhone River mouth area and sampling location: “Arles”. gR and pR mean grand Rhone and petit Rhone which are the two arms of the river.

**Survey of composition and degradation state of POM in the Rhone River using lipid tracers**

M.-A. Galeron et al.

[Title Page](#)

[Abstract](#)   [Introduction](#)

[Conclusions](#)   [References](#)

[Tables](#)   [Figures](#)

[◀](#)   [▶](#)

[◀](#)   [▶](#)

[Back](#)   [Close](#)

[Full Screen / Esc](#)


[Printer-friendly Version](#)

[Interactive Discussion](#)



## Survey of composition and degradation state of POM in the Rhone River using lipid tracers

M.-A. Galeron et al.



**Figure 2.** Water flow ( $\text{m}^3 \text{s}^{-1}$ ), suspended particulate matter ( $\text{mg L}^{-1}$ ) and particulate organic carbon (POC,  $\mu\text{M}$ ) of Rhone waters on our sampling dates (MOOSE data). Samples span a wide range of water flow conditions.




## BGD

11, 14197–14237, 2014

## Survey of composition and degradation state of POM in the Rhone River using lipid tracers


M.-A. Galeron et al.



**Figure 3.** Sitosterol and planktonic sterols (A), cuticular waxes and betulin (B), and polyunsaturated fatty acid (PUFA) (C) contents of the different samples. Standard error shown was estimated to be 14 % (see Sect. 2.3). Phytoplanktonic blooms are evidenced by the spike in planktonic sterols, and while the other tracers show the terrigenous origin of the POM sampled, the ubiquitous nature of sitosterol is made clear.

## Survey of composition and degradation state of POM in the Rhone River using lipid tracers


M.-A. Galeron et al.



**Figure 4.** Chlorophyll *a* levels (MOOSE data) and percentage of photodegradation of chlorophyll (calculated using CPPI, Cuny et al., 1999) in the Rhone waters on and around sample dates.

## Survey of composition and degradation state of POM in the Rhone River using lipid tracers


M.-A. Galeron et al.



**Figure 5.** Biotic and abiotic degradation of sitosterol (A) and cholesterol (B) in the different samples. Full error shown here incorporates the 14 % analytical standard error estimated for lipid quantification for all terms of the equations used. Sitosterol and cholesterol clearly have very different degradation patterns.

## Survey of composition and degradation state of POM in the Rhone River using lipid tracers

M.-A. Galeron et al.



**Figure 6.** Coprostanol:(cholesterol + cholestanol) and epicoprostanol:coprostanol ratios of the different samples. Full error shown here incorporates the 14 % analytical standard error estimated for lipid quantification for all terms of the ratios. Contamination threshold is 0.06 (See Sect. 3.3.2).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Survey of composition and degradation state of POM in the Rhone River using lipid tracers

M.-A. Galeron et al.



**Figure 7.** Relative percentages of intact hydroperoxides and their ketonic and alcoholic degradation products measured in the case of  $\omega$ -hydroxyoleic (cuticular waxes) and oleic acids, phytol, sitosterol and cholesterol oxidation products. Standard error was calculated based on the results obtained for all samples.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion