

Interactive comment on “Use of Ra isotopes to deduce rapid transfer of sediment-derived inputs off Kerguelen” by V. Sanial et al.

V. Sanial

virginie.sanial@legos.obs-mip.fr

Received and published: 17 December 2014

The authors thank W. Geibert for his relevant and constructive remarks.

The stations have different names but are indeed the names chosen by the project and consequently cannot be changed. UW stations were specific to the Ra study and refer to the surface samples collected “under way” using the ship intake.

All the ^{224}Ra and ^{223}Ra activities used in this study are excess ^{224}Ra and ^{223}Ra activities. The ^{224}Ra activities are corrected for the ^{224}Ra supported by ^{228}Th and the ^{223}Ra activities are corrected for the ^{223}Ra activities supported by ^{227}Ac . We clarified this in the section 2.3.2 (Sample analysis section) by adding “The ^{224}Ra activities are corrected for the ^{224}Ra supported by ^{228}Th and the ^{223}Ra activities are corrected

C7425

for the ^{223}Ra supported by ^{227}Ac . The ^{224}Ra and ^{223}Ra activities discussed here refer to the Ra activities corrected for the Ra supported by their parent isotopes.”

The Leclaire Rise is a topographic feature located west of the Kerguelen Islands (see Figure below; van Beek et al., 2008). Because it is located at ca 350 m depth (Weis and Frey, 2002), an impact of this topographic feature on the surface waters implies i) release of chemical elements from the sediments and ii) vertical mixing that would transport the chemical elements towards surface waters. Internal tides may promote such vertical transport of chemical elements in this area. In addition, because the Leclaire Rise is located north of the Polar Front (Park et al., 2014), the presence of Ra isotopes in the investigated area (east of the Polar Front) also suggests a transport across the front. If vertical mixing transports Ra in surface waters above the Leclaire Rise, it cannot be completely excluded that the Ra enrichments observed south of the Kerguelen Islands along the Polar Front (e.g. UW-23; UW-24; Figure 4) or south of the Polar Front (e.g. UW-15, UW-16 and potentially A3; Figure 4) are partly explained by an input associated with this topographic feature. However, sample UW-14 collected in surface waters lying above this rise (bottom depth at 342 m) does not show significant ^{223}Ra and ^{224}Ra activities and only low ^{228}Ra activity, which suggests slight impact of this topographic feature on the Ra activities of surface waters. Station R-2, which is located east of the Leclaire Rise - but south of the Polar Front - shows slight enrichments in ^{223}Ra and ^{224}Ra in surface waters (0.016 and 0.057 dpm/ 100 L, respectively), which could suggest an impact - but relatively minor when considering the Ra activities - in surface waters downstream of this plateau. Fe and other trace metal concentrations (REE, Mn, Al) do not show significant trends in surface waters at R-2, which indicates that the Leclaire Rise has a minor impact on surface waters downstream of this plateau (Bowie et al., 2014; van der Merwe et al., 2014; Grenier et al., in prep.). In contrast, these latter studies suggest that the shallow sediments of the Leclaire rise may impact waters in the 200-500 m depth interval downstream of the rise.

C7426

Bowie, A. R., van der Merwe, P., Trull, T., Qu  rou  , F., Fourquez, M., Planchon, F., Sarthou, G., and Blain, S.: Iron budgets for three distinct biogeochemical sites around the Kerguelen plateau (Southern Ocean) during the natural fertilization experiment KEOPS-2, *Biogeosciences Discuss.*, 2014.

Park, Y.-H., Durand, I., Kestenare, E., Rougier, G., Zhou, M., d' Ovidio, F., Cott  , C. and Lee, J.-H.: Polar Front around the Kerguelen Islands: An up-to-date determination and associated circulation of surface/subsurface waters, *J. Geophys. Res. Oceans*, 119(10), 6575–6592, doi:10.1002/2014JC010061, 2014.

Van Beek, P., Bourquin, M., Reyss, J.-L., Souhaut, M., Charette, M. A. and Jeandel, C.: Radium isotopes to investigate the water mass pathways on the Kerguelen Plateau (Southern Ocean), *Deep Sea Res. Part II Top. Stud. Oceanogr.*, 55(5-7), 622–637, doi:10.1016/j.dsr2.2007.12.025, 2008.

Van der Merwe, P., Bowie, A. R., Qu  rou  , F., Armand, L., Blain, S., Chever, F., Davies, D., Dehairs, F., Planchon, F., Sarthou, G., Townsend, A. T. and Trull, T.: Sourcing the iron in the naturally-fertilised bloom around the Kerguelen Plateau: particulate trace metal dynamics, *Biogeosciences Discuss*, 11(9), 13389–13432, doi:10.5194/bgd-11-13389-2014, 2014.

Weis, D. and Frey, F. A.: Submarine Basalts of the Northern Kerguelen Plateau: Interaction Between the Kerguelen Plume and the Southeast Indian Ridge Revealed at ODP Site 1140, *J. Petrol.*, 43(7), 1287–1309, doi:10.1093/petrology/43.7.1287, 2002.

Interactive comment on *Biogeosciences Discuss.*, 11, 14023, 2014.

C7427




Fig. 1.

C7428