

AUTHOR RESPONSE.

Ref.: Ms. Bg-2014-324. Title: *Riparian and in-stream controls on nutrient concentrations and fluxes in a headwater forested stream.*

Authors: Bernal, S., Lupon, A., Ribot, M., Sabater, F., Martí, Eugènia.

Dear Tom Battin and referees of *Biogeosciences*,

We would like to express our most sincere gratitude for your thoughtful comments and suggestions on the manuscript *Riparian and in-stream controls on nutrient concentrations and fluxes in a headwater forested stream*. We feel flattered that you consider our study *really clever* (R1), *challenging and attractive* (R2), and *pioneering* and that *it could potentially be an important contribution of our understanding of to what extent in-stream biogeochemical processes influence stream chemistry* (R3).

This is the sort of question we have been asking ourselves for a long time, and we carefully thought about how to conduct an empirical study to assess it at a relevant ecological scale. We agree with you that we have generated a “*large data set to tell a big story*” and that the paper needed to be improved and clarified in order to guide the readers through it. We have done a big effort in that sense from the introduction to the discussion following your suggestions. Moreover, we agree that the hydrological part of the study needed to be carefully reconsidered and we have reanalysed the data set thoughtfully to solve the cavities highlighted by you.

We are especially grateful to R3. First, for taking the time and patience to dive through the manuscript. Second, for the clarity in explaining what he/she felt was missing, wrong, or right. Third, for giving us clues that were essential to improve the paper and bring it, from our modest point of view, to a higher level. Our feeling is that the overall history was becoming crystal clear as we started calculating water and solute budgets at the whole-reach scale; just like if we had changed our googles. We felt a little bit overwhelmed at the beginning because this new set of results outcompeted part of the data analysis we proposed in the former version of the manuscript. Yet, we strongly believe that these new results have enormously contributed to improve the

manuscript because now we can sharply address our primeval question and further, they help us to solve several of the points highlighted not only by R3 but also by R1 and R2.

We provide below our detailed responses to the referees' comments. Your comments are listed in italics and our responses are inserted after each comment. In almost all cases, we have done what you have suggested. If we have revised things in a different way, or disagreed, we have stated why. If your suggestion was not applicable anymore in the new version of the manuscript, we have stated so.

We look forward hearing from you soon, and wish you a Happy New Year!

Susana Bernal

Center for Advanced Studies (CEAB-CSIC)

C/Accés a la Cala Sant Francesc 14,

17300, Blanes, Spain

Email: sbernal@ceab.csic.es

CC: Anna Lupon, Miquel Ribot, Francesc Sabater, and Eugènia Martí

Author Response to Reviewer #1

This manuscript uses a large synoptic stream and riparian nutrient dataset to tease out relative contribution of riparian zone and in-stream biotic processes on downstream nutrient fluxes. Very little research has addressed this question, thus the authors describe a novel finding. Their approach is really clever and the authors have a great dataset to test these ideas. However I have a lot of questions and suggestions to guide a revision.

This paper was a bit tough for me to follow. Part of it is that the authors are putting together a large dataset to tell a big story; I commend them for this. But this approach means that they need some sort of road map or conceptual guide for the readers to follow through the manuscript. Ways to do this may include some sort of conceptual model (see e.g. the approach taken Payn et al.), a results section that better describes the findings and how the data support these findings. I note that the discussion was much clearer in reporting the results than was the results section. It may also be possible to leave out some parts that do not add to the story the authors want to tell. E.g. the hillslope data in the discussion were not well integrated in the rest of the story.

*The authors used linear regression and GLM to examine change along the 15 sites of the reach. That probably works well to estimate if e.g. something doubled, but tests of significance will be hampered by the fact that these sites are not independent of each other. Processes at upstream sites determine those at downstream sites. In the case of nutrients it is likely the same molecules. Statistical test should consider this fact, and if they don't then the authors need to justify that point. I regret to not being an expert in spatial statistics, but something along the line of a generalized least squares regression may help. See, e.g., Ives and Zhu, *Ecol. Apps.* 16:2-32 for possible ideas on what, if anything, to do.*

*I am having trouble with the idea of only focusing net groundwater input when in fact there may be new groundwater entering a stream reach with concomitant losses. Thus net will be zero, but there could be a lot of new solute entering the stream via ground- water. See Payn et al *WRR VOL. 45, W11427, doi:10.1029/2008WR007644.**

Answer: Many thanks for your positive comments! We agree with the reviewer in that we are putting together a large dataset and that we could have done a better job when explaining the results and the overall message of the study. We are happy because we

have received excellent inputs from the three reviewers and we are convinced that the new version of the manuscript reads better and more smoothly than the former one.

Following the suggestions from the three reviewers, we have included a schematic representation of our mass balance approach, where nutrient fluxes in and out a stream segment are shown (new Figure 3). Moreover, we have included a more detailed explanation of our objectives and expectations in the last paragraph of the introduction (sort of road map), so that the reader is now better guided from the very beginning (pp 5-lines 106-116).

In addition, we have excluded several results from the former version that were not adding much to the overall aim of the study (namely, coefficients of variation, area-specific nutrient exports, GLM model, and hillslope data). On the other hand, we have rewritten the results and discussion emphasizing the results (both former and new) that do help to understand the influence of riparian groundwater and in-stream nutrient cycling on stream nutrient chemistry (namely, comparison between stream and riparian groundwater chemistry, tributary inputs, and in-stream net nutrient uptake rates at the segment and whole-reach scale).

We agree with the reviewer that the statistical analysis of this sort of data set needs to be done carefully because contiguous points may not be independent of each other. We considered several possibilities before applying GLM models and finally, we thought that including *distance* explicitly in the model could partially deal with the problem of co-linearity between distance and other explanatory variables. We are neither experts in statistical modelling and there might be better options than this one, as we have learned from the literature you suggested. In any case, we have finally decided to exclude it from the new version of the manuscript. The main reason why we did this statistical analysis was to put into the same picture in-stream processes and riparian groundwater inputs. However, as highlighted by R3, there are other ways much more simple and elegant to do so, such as to calculate nutrient budgets at the whole reach-scale and from there, calculate the relative contribution of each component to total stream nutrient fluxes (which was also a request from R2).

Regarding hydrological groundwater inputs, we agree with the reviewer that concomitant gross hydrological gains and losses may happen within each stream segment and thus, only considering the net groundwater inputs to each segment is an oversimplification of the complex hydrological interactions that occur at the riparian-stream interface. Unfortunately, we do not have enough data to calculate gross gains

and losses as in Payn et al. (2009). Nevertheless, we do now consider this limitation within the paper and we refer to net groundwater inputs more appropriately throughout the text and figures. Moreover, we have reanalyzed our hydrological data set thoughtfully following suggestions from you and R3 in order to get a better understanding of the stream-riparian hydrology in our study site (see our responses to your specific comments).

Specific comments

598-5 unclear what “which. . .” modifies **Answer:** Clarification not needed now.

599-7 considerably important. **Answer:** OK.

25 See the now classic work of Lowrance et al. on riparian controls of watershed nutrient export. **Answer:** Thanks. Lowrance et al. 2007 included (pp 3-lines 62).

601-6 recast to downstream-most site. **Answer:** OK from here on.

No comma after both. **Answer:** OK from here on.

602-3 two separate findings in one sentence.

Answer: The sentence: “The riparian zone covers 6% of the catchment area on average the slope is < 10%”, reads now: “The riparian zone is relatively flat (slope < 10%), and it covers 6% of the catchment area” (pp 7-lines 138).

602-19 composed of. **Answer:** OK

602-22 wetted width. **Answer:** OK

603-22 were used. **Answer:** OK

604-4 If using the molybdate blue technique on filtered samples, then one analyzes soluble reactive P (SRP) of which phosphate is one component.

Answer: OK, we do now refer to SRP rather than to PO_4^{3-} throughout the text, tables, and figures.

604-20. This method assumes no loss of water along the reach, i.e., the stream is solely gaining. Is this point true for this stream?

Answer: As mentioned before, we agree with the reviewer that considering only net groundwater inputs to each segment is an oversimplification of the complex hydrological interactions that occur at the riparian-stream interface. We do now state this limitation in the M&M (pp 10-lines 232-234).

As you noted, this stream is not solely gaining water. We now address explicitly this point in two ways following suggestions from you and R3. First, we have calculated the cumulative area-specific net groundwater inputs along the reach by summing up net groundwater inputs (Q_{gw}) from the upstream-most site to each of the downstream segments and dividing it by the cumulative catchment area (pp 10,11-lines 235-238; new Figure 4b).

Second, we have examined whether the reach was either a net gaining or a net losing stream by comparing concurrent hydrological gains and losses over the whole reach for each sampling date. We have considered that stream segments exhibiting $Q_{gw} > 0$ contributed to hydrological gains ($\Sigma Q_{gw} > 0$) while segments with $Q_{gw} < 0$ contributed to hydrological losses ($\Sigma Q_{gw} < 0$) (pp 11-lines 239-248). We have carefully thought about whether refer to these water fluxes as *gross* hydrological losses/gains or *whole-reach cumulative net* hydrological losses/gains. Following the suggestion from R3, we have finally considered that at this spatial scale (reach-level) these were *gross* hydrological gains and losses. Again, this approach is an oversimplification of the hydrology at the stream-riparian interface but we got to the conclusion that the question of whether water fluxes can be considered gross or net is, at the end, a question of spatial resolution. In any case, we would be happy to change the terms we are using to refer to these fluxes if you consider that some other terminology would be more accurate.

These new results indicate that, on average, the stream was a net gaining reach though gross hydrological gains exceed gross losses over the reach during two sampling dates (new Figure 2c and d). Further, the stream was a net gaining reach along the first 1.5 km and the last 0.5 km, while hydrological losses were evident along the intermediate 2 km (new Figure 4b) (pp 15-lines 342-248).

604-21. *So divided by watershed area, correct? Unclear as written.*

Answer: Yes. We have added “*by dividing instantaneous discharge by catchment area*” (pp 10-lines 220-221). Thanks.

605-9. *It seems like one could get a lot more information if the data were not averaged for each period but done separately using a multilevel approach.*

Answer: The GLM has been excluded from the new version of the manuscript. Thanks for the suggestion anyway.

606-4. *But Table 3 reports a p-value from the ANOVA, which is not a goodness of fit (as I understand it). I am not sure why to report a P-value between the differences in these models anyway.*

Answer: The GLM has been excluded from the new version of the manuscript. Thanks for the suggestion anyway.

606-10. *Up to now all of the stats are based on parametric distribution. Why a switch to a non-parametric test here?*

Answer: Thanks for noticing. In the former version, we forgot to include that the reason why we switch to non-parametric test is because solute concentrations were not normally distributed, as usual. We have added the following sentence in M&M: “*A non-parametric test was used because solute concentrations were not normally distributed (Shapiro-Wilk test, $p < 0.01$ for all studied solutes) (Zar, 2010)*” (pp 11-lines 254-256).

Furthermore, we agree with the reviewer that concentrations and fluxes may or may not change linearly over space and thus, we do now consider linear and nonlinear models when examining the longitudinal pattern of discharge and nutrient concentrations (pp 10-lines 223-226).

606-15. *I usually think of a CI as a confidence interval. This common use of CI may be confusing when redefined as chemical index. Chemical index is by itself not very meaningful, so there may be a better phrase (and therefore acronym) to use.*

Answer: Agree. We were struggling for a while trying to find a better acronym and obviously, we did not succeed.

Considering comments from you and R2, we have decided to use dissolved oxygen concentration (DO) instead of the chemical index. DO is easy to interpret and it provides the same information.

608-2. *What about segments that are gaining and losing at the same time, where the nutrient concentration in the groundwater inflow is different than the outflow, which we would assume is the same concentration as the streamwater.*

Answer: We agree with the reviewer that some assumptions are needed when applying a mass balance approach, especially because concomitant gross gains and losses were not measured within each stream segment. Note, however, that even if we would have been able to measure those fluxes, we would have had to deal with uncertainty too. For instance, we could have never been sure if gross hydrological gains along each stream segment do actually have the same chemical signature than riparian groundwater measured in the two wells installed in each segment (at the top and at the bottom). Also, we could not know either whether the chemical signal of gross water losses is actually similar to that measured in the stream water column.

In this study, we have characterized to some extent riparian groundwater chemistry along the reach. This is one of the novelties of our design in comparison with other synoptic studies, and we believe that this information, despite being limited, is extremely helpful to constrain the uncertainty associated to the water chemistry of riparian groundwater inputs. Otherwise, we might have had to make even stronger assumptions regarding groundwater chemistry, and the uncertainty associated to F_{sw} would have been even larger. Nevertheless, we completely agree with you that we need to acknowledge the limitations of our approach and be careful when explaining and discussing these results.

In the new version of the manuscript, we are more explicit about the assumptions made when calculating nutrient groundwater fluxes (F_{gw}) (pp 13-lines 288-294). Moreover, we now state that reliable measurements of riparian groundwater inputs are difficult to obtain and that by installing only 15 piezometers along the reach we might be missing part of the intrinsic variation of riparian groundwater chemistry along the study reach. However, and despite its limitations, we highlight that sampling near-stream riparian groundwater can help to constrain the uncertainty associated to this water source and provide more reliable estimations of in-stream net nutrient uptake (pp 25-lines 600-608).

608-18 *SE estimated by averaging over what, time? Also, what is the frequency of F_{sw} not equal to 0 by chance using this technique? It seems to me that many would be not significant, but that would require a little explanation.*

Answer: Yes, we agree that some additional explanation was needed. Thanks for noticing. In the new version of the manuscript, however, we have calculated the upper and lower limit of F_{sw} based on the empirical uncertainty associated with Q (please, see our responses to your specific comment on this regard). Further, we now assume that F_{sw} was undistinguishable from 0 when its upper and lower limit contained zero (pp 13-lines 307-308)

We believe that with this approach we are more coherent because the uncertainty associated to water and nutrient fluxes is based on the same grounds throughout the study. Note that this approach is more conservative than the previous one, in the sense that the proportion of cases for which in-stream gross uptake and release counterbalance each other ($F_{sw} = 0$) is higher (please compare former and new figure 6). However, the main results are the same ones: (1) the occurrence of in-stream NO_3^- release increases along the reach, (2) in-stream gross uptake tends to prevail over release for NH_4^+ , and (3) the frequency of $F_{sw} > 0$, < 0 , and $= 0$ does not show any clear pattern for SRP.

608-22. But there could be groundwater input in a losing segment.

Answer: Yes, that's right. However, even if we may have had data on gross hydrological gains and losses, we would have had to consider the difference between the two (that is net groundwater inputs) to calculate F_{in} . Or in other words, it may not be accurate to consider gross hydrological gains in F_{in} without simultaneously considering that part of this water is simultaneously lost towards the riparian zone and thus, does not contribute ultimately to downstream export.

Taking into account your point, we do now clearly state that F_{in} as is the sum of input fluxes from upstream, tributaries and *net* riparian groundwater inputs when $Q_{gw} > 0$ (pp 14-lines 320-321).

608-24. Putting the R code and data into an appendix will help readers to replicate this work in the future. This is a really valuable dataset.

Answer: The GLM has been excluded from the new version of the manuscript and thus, this R code is not in the paper anymore. This data set is part of the grounds of the thesis of Anna Lupon who is in her last year of PhD. We have in mind to add these data as Supplementary Material or Appendix as soon as she starts publishing the chapters of her thesis. Hopefully, this will be soon. Thanks for the suggestion.

609-3 Units for this area specific discharge simplify to units of length per time (say mm/d) which is common in hydrological literature and therefore less confusing for some to read. **Answer:** Ok. We have transformed water fluxes to mm d^{-1} .

609-15 The Cl data and the Q data in this paragraph are not really well linked, i.e. I am not sure what collective finding they support.

Answer: Agree. We have reorganized this section of the results. In the new version, we introduce first results related to water fluxes and then, those related to Cl^- concentrations and fluxes. Thanks.

609-20 What is the error in measuring Q via using dilution gaging? Given that there is always some measurement error, streams will always be gaining or losing when calculated as a strict difference. How much does a stream have to gain or lose to detect a difference above the measurement error?

Answer: This is a good point. Thanks. We have calculated the uncertainty associated to our empirical estimates of Q by selecting pairs of slug additions performed under similar water depth conditions (difference in water level at the thalweg < 1 mm). To do so, we have benefited of a data set with 126 slug additions used to calibrate the water level sensor from a parallel study at the Font del Regàs stream. We have considered the relative difference in Q between these pairs of data as an empirical uncertainty estimator. There were 11 pairs of data that fall within our criterion, and the relative difference in Q between them was pretty small (1.9%).

In the new version of the manuscript, we have included the estimation of this uncertainty (pp 8,9-lines 183-188). Moreover, we have considered this uncertainty for calculating the upper and lower limits of water fluxes and posteriorly, the uncertainty of nutrient fluxes. The uncertainty associated to stream and groundwater fluxes is now included in the results (see new Figure 2 and 7).

610-23. I assume that these predictor variables do not covary among themselves?

Answer The GLM has been excluded from the new version of the manuscript.

610-27. ok so then why fit a straight line to the data?

Answer: Agree. We do now describe the longitudinal pattern of NO_3^- concentrations more accurately: “During the vegetative period, stream NO_3^- showed a U-shaped

pattern: concentrations decreased along the first 1.5 km, remained constant along the following 1 km, and increased by 60% along the last km of the reach” (pp 16-lines 368-370).

612-9 Makes sense given the chemical index uses NH₄ as part of its calculation.

Answer: After taking into account comments from you and R2, we have decided to use dissolved oxygen concentration as a proxy of the redox conditions in stream water and groundwater rather than the chemical index (pp 17-lines 399-401).

613-1 This paragraph is problematic in that it is repeating things from the introduction. Then it describes the significance of the findings before stating what the main findings are. I note that the results were a difficult place for me to understand the main finding that was mostly a description of the data. In any case this paragraph can be safely deleted. Better, given the results section, would be to summarize the main findings.

Answer: Thanks for the suggestion. This paragraph has been deleted and we have rewritten the discussion to start going directly to the point.

614 8-12. This clear statement of the findings are restating results.

Answer: We agree with the three reviewers about that. The reanalysis of the hydrological data set motivated by your comments and suggestions has shed new light on what drives the longitudinal pattern of stream Cl⁻ concentration. Briefly, we have found that Cl⁻ concentration was higher at the tributaries than at the main stream, especially during the vegetative period. Permanent tributaries comprised ca. 50% of the catchment area and were the main contributors to stream discharge (56%). Therefore, hydrological mixing with water from tributaries could partially explain the 40% increase in stream Cl⁻ concentrations observed along the reach.

In addition, riparian groundwater inputs could also explained the longitudinal pattern exhibited by stream Cl⁻ concentration because this compartment contributed substantially to stream discharge (26%) and also exhibited higher Cl⁻ concentration than stream water during the two periods.

These new results and the associated discussion are included in the new version of the manuscript (results section 4.1 and pp 19-line 447-455)

615-6 By saying .. "where the N₂ fixers are highest" implies that the N₂ fixer may control NO₃, but later in the paragraph we are reminded that there was no relationship with N₂ fixers. I would recast to avoid creating confusion here and instead simply state the most plausible mechanism up front, and not one found to be not plausible.

Answer: Agree. We have rewritten this part of the discussion that is now focused on the empirical evidences we have. First, and following the suggestion from R3, we highlight that the flux of NH₄⁺ from riparian groundwater is not large enough to sustain in-stream NO₃⁻ release during the vegetative period along the last 0.7 km of the reach (pp 22-lines 511-515). Then, we discuss that an additional source of N could be increased in-stream nitrification at the valley bottom based on previous studies showing that leaf litter from riparian trees, and especially from N₂-fixing species, can enhance in-stream nutrient cycling. Thus, we propose that large stocks of leaf litter from alder and black locust combined with warm temperatures could enhance in-stream mineralization and nitrification during the vegetative period. This process could explain the marked increase in stream NO₃⁻ release observed at the lowest part of the catchment (pp 22-lines 515-522).

In addition, we consider other alternative explanations such as anthropogenic sources and autotrophic N fixation as suggested by R3, which seem not likely given the empirical evidences we have. Finally, we end up concluding that although the sharp increase in NO₃⁻ availability observed at the end of the reach could be related to the presence of black locust, further research is needed to test this hypothesis (pp 22,23-lines 523-543).

615-16 lowest part of the catchment. **Answer:** OK.

616-18 New results in the discussion?

Answer: Following R1 and R3 suggestions, we have deleted this part of the discussion as well as former Figure 8.

617-20. Is that because NO₃ concentrations are higher than SRP and NH₄?

Answer: Good point. To answer your question, we have analyzed whether differences in F_{sw} and $|F_{sw} \cdot x / F_{in}|$ were statistically different among nutrients. Differences were not statistically significant for F_{sw} , but they were significant for $|F_{sw} \cdot x / F_{in}|$. These results

suggest that differences in $|F_{sw} \cdot x / F_{in}|$ among nutrients were mainly because input fluxes showed large differences in concentrations that tend to be 20-fold lower for NH_4^+ and SRP than NO_3^- , as you pointed out (pp 21-lines 480-483).

617-22. *What is meant by “cycled more efficiently” Longer uptake length? Lower v_f ?*

Answer: Thanks. We now refer to *in-stream gross uptake velocity* (pp 20-lines 476-478).

618-21 *“if we are to understand”?* **Answer:** Yes. Thanks.

618-23. *This sentence says that instream processing is important but manifests itself at a small spatial scale than riparian processes?*

Answer: We have been thinking thoroughly about the implications of our results, and we believe that the new version of the manuscript states more clearly the take home message that emerges from this study (new conclusions section).

Briefly, our study highlights that the stream had a strong potential to transform nutrients and that in-stream processes were substantially contributing to either increase or decrease stream nutrient export to downstream ecosystems. However, the influence of in-stream processes on stream nutrient concentration and fluxes did not systematically translate into longitudinal patterns because changes in nutrient concentration along the reach were the combination of both in-stream cycling and nutrient inputs from terrestrial sources (pp 24,25-lines 576-599).

619-1. *Ok a great way to end, but I could use bit more explanation here.*

Answer: Agree. We have carefully thought about the bottom line and implications derived from our results, as mentioned in our earlier response. One of the major implications of our study is that the assessment of both in-stream cycling and terrestrial nutrient sources is crucial if we are to understand the contribution of in-stream processes to stream nutrient dynamics at relevant ecological scales. Please read the new conclusions section for more detailed explanations.

Table 1. Reporting an SE implies normally distributed data, yet there is a non-parametric test used. The equation for CI is not how it is described in the text.

Answer: Following the reviewer suggestion, we do now provide the median and the interquartile range in Tables 1, 3, and 4. The chemical index has been substituted by dissolved oxygen concentration.

Table 3. Likelihood is relative likelihood, correct?

Answer: The GLM along with former Table 3 have been excluded from the new version of the manuscript.

Fig 2. The X-axis looks like it is plotted categorically vs. numerically as a function of distance. It seems to me that plotting numerically would be clearer because the sites are not equally spaced. Stream width and % sand should be on separate plots.

Answer: Following the reviewer suggestion, the X-axis in Figure S1 (former Figure 2) is now plotted numerically, and stream wetted width and the percentage of sand in streambed are plotted in two separated plots. In alias of simplicity and to condense the study, note that this figure is now in Supplementary Materials.

Fig. 5. A straight line seems to be a poorly fitting model for a U-shaped pattern of nitrate concentration and flux.

Answer: Following the reviewer suggestion, we do now use lineal and nonlinear models. Moreover, we refer specifically to a U-shaped pattern when describing the longitudinal pattern of stream nitrate concentrations in the results section.

Fig 7 legend, bottom “post-hoc”. **Answer:** OK. In any case, this figure has been excluded.

Author Response to Reviewer #2

Presented data and descriptions include interesting and important information. The aim and approach to evaluate the riparian and in-stream controls presented in this manuscript is challenging and attractive. Therefore, the document itself is valuable, even though this is a report of a case study at particular middle scale catchments. However, there are several weaknesses for considering a publication as an original article in the current version. I hope following comments will help the authors to improve the contents and descriptions.

Answer: Thanks for your positive comments. Yes, this study only reports results for a particular headwater catchment. Yet, as pointed out by R3, it is worth noticing that the experimental design we have conducted is not commonly found in the literature and in this sense, our study can provide valuable information to the reader of *Biogeosciences*. In this new version of the manuscript, we have worked throughout the results and the text in order to highlight the novelties and uniqueness of our study. We hope that you find the paper improved.

Comments:

1. As the authors stated in the first paragraph of the discussion section, the novel point of this paper was statistically quantitative analysis of sources of nutrients in the stream water. However, in the conclusion section and abstract, their descriptions on relative contribution of riparian and in-stream processes was still very qualitative. More quantitative expressions on relative contributions were favorable.

Answer: Thanks for noticing, this is a very important point. Following the suggestion made by you and R3, we now quantify the relative contribution of riparian groundwater and in-stream cycling to stream nutrient fluxes for the whole reach. By going that, we are able to provide quantitative estimates of the contribution of riparian groundwater, in-stream release, upstream, and tributaries to stream inputs fluxes for each sampling date. Moreover, we are able to quantify in-stream net nutrient retention at the whole-reach scale (see new Table and Figure 7, and associated results).

2. P11600, L17-20: This type of hypothesis has been commonly recognized by many researchers in the field of river ecosystems. Not only your proposal.

Answer: That's right. Following your suggestion, we now indicate that some authors have proposed that nutrient concentrations should decline in the downstream direction if in-stream net uptake is high enough and changes in riparian groundwater inputs are relatively small (Brookshire et al., 2009), while other have shown that nutrient concentrations are patchy and highly variable along the stream as a result of spatial patterns in upwelling and in-stream nutrient processing (Dent and Grimm, 1999) (pp 4,5-lines 87-98).

3. *In the methods section, it would be helpful for readers to show a conceptual diagram or picture examples expressing the spatial distributions and scales of riparian forests (zone), riparian-stream interface (zone?).*

Answer: Thanks for your suggestion. As already mentioned in our general response to R1, we have added a schematic representation of our mass balance approach. This scheme is intended to clearly define fluxes in and out a stream segment (new Figure 3). Following the suggestion from R3, we have also simplified our conceptualization of the stream-riparian interface. As depicted in the new Figure 3, there are only two compartments: the stream and the riparian zone, and by definition biogeochemical processes can only occur either inside (stream) or outside (riparian zone) the boundary.

4. *How does the climatic seasons correspond to the vegetated/dormant seasons? Is the vegetative season rainy season or high flow season? And also, doesn't the seasonal variation in discharge rate affect those of the nutrient concentrations and fluxes?*

Answer: In order to answer your question as well as those from R3, we have included more information regarding the seasonality of precipitation and discharge during the study period.

First, the temporal pattern of P and Q can be seen in new Figure 2 (panels a and b). These data comes from our permanent sampling station which is equipped with a water sensor and an automatic sampler. Second, we do now provide information about the seasonality of rainfall during the study period which was minima in summer and highest in spring (pp 6-lines 127-129). Moreover, we specify that rainfall was similar between the dormant and vegetative period (pp 10-lines 213-214). This was so because the spring 2011 was rainy and this rain compensate the following dry summer.

Finally, we also provide information about the seasonality of stream discharge at base flow conditions which was low in summer and peaked in spring (pp 7-lines 160). High-temporal resolution data showed that stream discharge was higher during the dormant than during the vegetative (Table R1). However, this seasonality was not reflected in our synoptic study likely because our sample size was too low. In addition, the vegetative period included dates with high flow (spring 2011) and low flow (summer 2011) which likely balanced each other (Table R1).

Table R1. Median and interquartile range [25th, 75th percentile] of area-specific discharge (mm) at the downstream-most site. Data from this study and from the permanent sampling station installed at the valley bottom is shown. The number of cases is shown in parenthesis for each group. Different letters indicate statistically significant differences between either calendar seasons or periods (Wilcoxon paired rank sum test, $p < 0.01$).

	This study	Permanent Station
Autumn 2010	0.46 (2)	0.41 [0.28, 0.73] ^B (91)
Winter 2011	0.36 (2)	0.44 [0.40, 0.57] ^B (91)
Spring 2011	0.88 (2)	0.79 [0.62, 1.07] ^A (91)
Summer 2011	0.26 (2)	0.33 [0.20, 0.38] ^C (91)
Autumn 2011	0.70 (2)	0.42 [0.17, 1.88] ^A (91)
Vegetative	0.41 [0.24, 0.59] ^A (7)	0.37 [0.20, 0.62] ^B (214)
Dormant	0.43 [0.35, 0.64] ^A (4)	0.46 [0.40, 1.55] ^A (151)

Regarding your last question, our data showed no seasonality in stream nutrient concentrations, at least for nitrate which was the nutrient exhibiting the highest concentrations (Table R2).

Table R2: Mean and standard deviation of nitrate concentration at the downstream-most site. Data is from this study (only mean) and from our permanent station. The number of cases is shown in parenthesis for each group. No differences were found between calendar seasons for the permanent station data set (Wilcoxon paired rank sum test, $p < 0.01$).

	Nitrate (mg N l^{-1})	
	This study	Permanent Station
Autumn 2010	0.27 (2)	0.17±0.03 (20)
Winter 2011	0.22 (2)	0.21±0.09 (71)
Spring 2011	0.13 (2)	0.17±0.02 (80)
Summer 2011	0.17 (2)	0.17±0.02 (89)
Autumn 2011	0.22 (2)	0.23±0.12 (89)

5. P11602, L14-16: *In the discussion section, the authors discussed about the influences of the N₂ fixing trees and the N dynamics of soil microbes on the nutrient input from the riparian zone to the stream. Those processes are usually most active at the organic horizon and the near surface part of the mineral soils. Was the sampling conditions which made the influences from those soil horizons minimal appropriate to investigate the direct effect of those N dynamics? Related to this question, in order to elucidate the controls by the biogeochemical processes of riparian zones comprehensively, relative degree of impact under the low flow condition should be estimated compare to that under the high flow or storm conditions.*

Answer: Thanks for your suggestion. The reviewer is right in that the contribution of different water sources may change between base flow and storm flow conditions. However, the present study was designed to understand in-stream nutrient cycling during base flow conditions, when its contribution could be more relevant and thus easier to tease out from other factors as indicated in the M&M section.

As we already mentioned in our specific responses to R1, we have rewritten this part of the discussion to avoid creating confusion. We have focused on the most plausible mechanisms that could explain the observed increase in stream NO₃⁻ concentration and the high frequency of in-stream NO₃⁻ release at the valley bottom during the vegetative period. We still invoke the potential role of black locust, but mostly as a source of edible leaf litter to the stream, which could enhance in-stream nitrification during warm periods. We have been more cautious on our rationale and we do now explicitly say that further research is needed to test the hypothesis that black locust alter stream nutrient dynamics (pp 22,23-lines 10-543).

6. *What was the specific benefit to use the CI value (considering the NH₄ + concentration) for evaluating redox condition instead of DO?*

Answer: Following your suggestion, we do now use dissolved oxygen concentration (DO) instead of the chemical index.

7. *A table for summarizing definition of each variable (Q_{gw}, Q_{spf}, F_{spf}, C_{sw}, etc.) would be very helpful for readers.*

Answer: As we already mentioned in our earlier responses to your point #3, we have included a schematic representation of the mass balance (new Figure 3). The

conceptual model indicates the two compartments considered: the stream and the riparian zone, as well as fluxes in and out the segment (F_{ef} , F_{top} , F_{bot} , F_{gw} , F_{sw}). In addition, we have indicated in the caption how nutrient fluxes were calculated (Q x C). We believe that in this manner the reader can get a quick idea of the variables considered in the mass balance, which is more illustrative than a glossary table.

8. *More detailed explanations of stream water sampling should be needed. Because, solute concentrations are often different with parts of cross section. Effect of the riparian groundwater might be evaluated differently at the bank side and the center of the stream.*

Answer: In the new version of the manuscript, we have specified that stream water samples were collected at the thalweg of the watercourse (pp 8-line 168). Moreover, we acknowledge that our characterization of riparian groundwater, despite valuable, is limited because riparian groundwater inputs can be highlight variable over space and it is complicate to determine the chemical signature of the groundwater that really enters the stream (pp 25, 600-603).

9. *Sub-chapter 4.3 included some discussions based on observed results. Some contents in this section might be categorized into the discussion chapter.*

Answer: After rereading carefully this subsection, I'm afraid to say that we do not really get which particular contents you are referring to.

Our impression is that we are not including any discussion into the results section, though at some point we add some clarifications such as that the stream was acting as a net source or sink for a particular nutrient. Not sure if this is what you meant, but in any case we believe that this is a result in itself as the considered variables were already defined using these terms in the M&M section.

10. *The first paragraph of the Chapter 5 was redundant, and some contents overlapped with the introductory section.*

Answer: Agree. Thanks for noticing. This paragraph has been deleted.

11. *P11613, L25- P11614, L1: The authors cited the paper Asano et al. (2009) to show an example indicating a reduction in the variability of solute concentrations as catchment size increases. Then, they stated that their case did not show the decrease*

in the CV of stream solute concentration along the reach. However, the discussions in Asano et al. (2009) did not tell the variability of solute concentrations decrease with the distance from the headwater in a SINGLE PARTICULAR stream. The “variability” in their paper indicated a variability of solute concentrations AMONG THE MULTIPLE DIFFERENT STREAMS WITH SAME CATCHMENT SCALE. The discussion was totally different with that in this manuscript.

Answer: Thanks for noticing. We do not refer to this paper anymore in the new version of the manuscript because, as suggested by R1, we should focus on the results that really contribute to solve our question, and the longitudinal patterns of CV of nutrient concentrations were not very helpful in this sense.

12. *P11614, L407: It is generally accepted that a major source of phosphate in stream is chemical weathering of the bedrocks. And also, phosphate in the vegetated floor and surface soils are usually recycled tightly within the plant-soil internal cycling. If the authors want to conclude that the controlling factor for stream phosphate concentration was the “hill slope water source”, they had to explain how different the hill slope water (groundwater?) input was from the water input from the riparian zones. Citations of Asano et al. (2009) was not appropriate, because they did not discuss the sources of solutes using riparian-hill slope comparison. Their concept was that the stream solute concentrations could be explained the mixing of shallow and deep groundwaters both from hillslopes.*

Answer: Thanks for your comment. See our previous response.

Following suggestions by the three reviewers, we do now focus on whole-reach mass balance calculations to examine the relative contribution of different sources to stream nutrient fluxes. Regarding SRP, our calculations indicate that tributaries and riparian groundwater were the major contributors of this nutrient to the stream. In-stream processes mostly contributed to increase rather than decrease stream fluxes of SRP (see new Table 4 and new Figure 7).

13. *P11614, L8-14: The authors concluded that the drop in the specific discharge and the Cl⁻ concentration along the reach were due to increase of evapotranspiration. This explanation might be doubtful. Could the evapotranspiration rate vary significantly within a few-km scale? Did vegetation type change from headwater to valley bottom significantly?*

Answer: We agree with the three reviewers that this point needed to be better discussed.

As already mentioned to R1, a more careful look to our data set highlighted that the longitudinal increase in Cl^- concentrations could be explained by hydrological mixing with tributaries and riparian groundwater. These two sources contributed substantially to stream discharge (> 50% and 26%, respectively), and both exhibited higher Cl^- concentrations than the stream, in particular during the vegetative period. Please, see more details about the changes made in our response to R1, as well as in results section 4.1 and pp 19,20-line 447-455.

14. *P11615, L4-L28: The discussions of this part were confusing. It seemed that the longitudinal pattern of NO_3^- concentration had two phases: decrease from (0-3000 m) and increase (3000-3700 m). But, the ratio of the N_2 -fixing trees increased monotonically from headwater to valley bottom. To explain the longitudinal pattern of NO_3^- concentration sufficiently, they should show the more consistent logic to apply to both decreasing and increasing phases with certain evidences. Probably, there might be two different controllers*

Answer: Agree. Thanks for noticing. As already mentioned in our responses to R1, we now describe the U-shaped pattern of stream NO_3^- concentrations (pp 16-lines 368-370). Moreover, we now acknowledge two 2 different drivers of stream NO_3^- concentrations. First, as included in our responses to your comment #5, we have rewritten this discussion paragraph and now we refer specifically to the increase in stream NO_3^- concentrations observed at the lowest part of catchment. We now consider several alternative explanations that could be behind such pattern, being one of them the predominance of N_2 -fixing species along the last 0.7 km (pp 22,23-lines 515-543). Second, we discuss that the decreasing pattern exhibited by NO_3^- concentrations along the first 1.5 km could not be explained by in-stream processes alone because its contribution to reduce stream NO_3^- fluxes was too low. Thus, the observed longitudinal pattern resulted from the combination of both in-stream nutrient cycling and hydrological mixing with riparian groundwater and tributary inputs (pp 24-lines 556-570).

15. *P11615, L17-19: On English expression at the following part: "...we could not establish any positive relationship between the proportion of N_2 -fixing tree species*

and stream NO₃- concentrations along the reach.” Any relationships can not be “established” by researchers. They can only observe and find the relationships.

Answer: We have substituted “*establish*” by “*find*”

Author Response to Reviewer #3

This is a pioneering and potentially an important contribution to the long standing question of whether, how, and to what extent in-stream biogeochemical processes influence stream chemistry. The question has sparked debate over whether in-stream processes alter chemistry enough to compromise the interpretation of stream exports as a measure of terrestrial watershed losses, and therefore whether inferences based on the small watershed concept need to be reassessed. It has been suggested that if in-stream processes are important enough to make a difference, then this should be revealed by longitudinal concentration gradients reflecting the removal or addition of nutrients to the water column. And, in fact, a number of studies have linked net uptake to declining downstream concentrations. But because nutrients enter (and leave) the stream laterally along its length, the notion a net uptake necessarily generates a declining concentration gradient is a simplistic fallacy. (See Eq. 4 of Brookshire et al. 2009 Ecol. 90:293, which is correct but misinterpreted by those same authors).

The work under review demonstrates (1) that in-stream processes may strongly affect stream chemistry and yet leave concentrations longitudinally uniform (or otherwise varying), as did Meyer and Likens (1979 Ecol. 60:1255), and (2) the magnitude of biotic effects can be inferred using a mass balance approach that fully accounts for lateral inputs and losses. The authors estimated the effect on stream chemistry of uptake or release by taking the difference between inputs and outputs assessed on each of 15 segments of a 3700 m reach of reach. The assessments were synoptic, repeated 11 times over a year and a half. No one has done this before, at least on this scale. The approach has limitations. We do not get an annual budget and a “sink” is simply the instantaneous difference of inputs and outputs: We do not see whether the nutrient is accumulating, being transformed, or being lost to the atmosphere. However, we do see detailed spatial resolution, revealing a surprising degree of spatial heterogeneity and, wherever there are lateral losses as well as gains, mass balance even at the whole-reach scale, cannot be assessed without it.

I use the caveat “potentially” because I think the presentation is weak in several ways and falls far short of what this amazing data set could support. The major problem, I think, is that the authors are trapped by the misconception that in-stream processes should produce longitudinal gradients. They often do, but they don't necessarily and there are good reasons why they wouldn't. I think this could be a very

powerful paper if directly challenged the misconception, and showed how streams can strongly affect concentrations without generating longitudinal gradients. Much more on this below. I have trouble understanding the basic message or messages of the paper, find that important information is missing, see a number of inconsistencies in presentation, and suspect that there may be some major errors. I hope all these can be resolved because this is important work.

Answer: Many thanks for your constructive comments! Your review is plenty of good advices and helpful suggestions. Overall, we have learned and enjoyed while working on it. As we mentioned in our letter to the editor, it has been like changing googles!

We have done a great effort to clarify the message, we have added the information that was missing and we believe that the inconsistencies are resolved (see our detailed responses below). Considering how fast and mercantilist is becoming science in the last years, at least, at this end, we sincerely appreciate the time you have spent to get through the paper and explain us how to improve it. For us it has been one of those rare and beautiful moments, in which doing science and the whole peer review process is worth and meaningful. Thank you.

First, here is what I consider missing:

(1) A whole reach budget, or budgets (by period, or sampling date). The whole-reach budget summarizes the big picture and will, in the long-run, be an essential, citable result of the paper. Moreover, what happens segment by segment must add up to the whole reach, so it provides closure. A budget would consist of upstream inputs, tributary inputs, groundwater inputs, groundwater losses, instream source-or-sink, and downstream export, i.e., the same as your segment (Eq. 2) budgets, for water, chloride, and nutrients, reported in mass/time (not area-specific).

Answer: Great suggestion! The paper now includes water and nutrient budgets at the whole-reach scale for each sampling date. These calculations are now an essential result of the paper because they allow quantifying the relative contribution of different sources to either increase or decrease stream nutrient fluxes, which was one of our aims. See new Table 4 and Figure 7 for a quick overview.

The consequences of following your advice have cascaded down because these results outcompete an important part of the analysis proposed in our former version, such as the examination of the CV of stream nutrient concentration, the longitudinal

pattern of area-specific nutrient fluxes, and the GLM. These results did not help much to address our objectives as pointed out by R1. We have rewritten the M&M, results and discussion section according to these changes (see more detailed changes below).

Overall, calculating the budgets at the catchment scale is a beautiful idea, and we feel that these new results bring the paper to a higher level.

(2) The map shows many unmeasured tributaries that account for roughly half as much contributing area as the measured tributaries (a table of sub-basin areas would be helpful). The way inputs from the unmeasured tributaries were incorporated into the mass balance should be described. If they were dry, this should be so stated.

Answer: Yes, thanks for noticing. You are correct: we only collected water samples from the four tributaries carrying water, at least during the dates we went to the field.

In addition, and motivated by some of your comments below, we have added a fifth tributary in this new version of the manuscript. This was a small effluent (catchment area $< 0.4 \text{ km}^2$) that drained through the inhabited area at the valley bottom. It was covered with bushes which prevented us from successfully conducting slug additions (and thus include it in our budgets). This was the reason why we did not include these data in the former version. However, stream chemistry from this tributary can be valuable to evaluate the potential influence of human impacts on stream water chemistry (see our responses to your comment in this regard).

This info together with their drainage area is now included in M&M (pp 9-lines 179-194) See also improvements in the caption of Figure 1.

(3) The seasonality of rainfall and flow should be provided. Knowing that dormant and vegetative season flows were similar is not sufficient because we would expect increasing baseflows throughout the dormant season and declining flows throughout the vegetative season.

Answer: Following suggestions by you and R2, we have added more detailed information about the seasonality of rainfall and discharge during the study period (please see our responses to R2 in this regard).

Briefly, high-temporal resolution data from our permanent sampling station showed that stream discharge at base flow conditions was lower in summer than in winter/spring. Yet, this was not reflected in our synoptic study likely because the sampling size was too low and high discharge in spring compensate low summer flows

(see Table R1). Please, see more details in our response to your *temporal pattern* comment.

(4) We need more information on groundwater gains and losses, especially the gross inputs and losses over the reach. The assessment of groundwater contribution requires both inputs and losses, not simply a net inflow. Also needed is an explanation of the groundwater input shown in Fig 3. c. If it is cumulative with watershed area, as is Q_{spf} in Fig. 3a, then Fig. 3 c tells us that, over the whole reach, there was no net gain from groundwater, yet the longitudinal increase in flow (L/s, not area-specific) was substantially greater than explained by measured tributary inputs.

Answer: As already mentioned in our responses to R1, we have followed your suggestion and calculated gross hydrological gains and losses over the entire reach for each sampling date (see new Figure 2 and related results in section 4.1). Please, find more details in our specific responses to R1.

The former Figure 3c showed mean net groundwater inputs for each segment, no cumulative inputs. We agree, however, that plotting cumulative area-specific net groundwater inputs along the reach (new Figure 4b) is more helpful and easy to compare to area-specific discharge in Figure 4a (which is a cumulative variable by definition).

(5) We need a more complete presentation of the F_{sw} data, longitudinal and temporal. These are the real contribution of the paper, but we see only the whole-study averages, standard errors, and a graph of the frequency of positive and negative estimates.

Answer: That's right. Thanks for your suggestion. These results have been emphasized in the new version of the manuscript. We do now show:

- (1) medians and IQR of F_{sw} at two spatial scales (segment and whole reach) (new Table 3),
- (2) medians and IQR relative contribution of different sources (upstream, riparian groundwater, tributaries, and in-stream release) to stream solute fluxes (new Table 4),
- (3) the longitudinal variation of the frequency of $F_{sw} > 0$, < 0 , and $= 0$ (Figure 6), and
- (4) the temporal pattern of F_{sw} (whole reach-scale) together with its contribution to reduce stream nutrient fluxes (new Figure 7).

In addition, here we include the longitudinal variation of F_{sw} for individual segments (Figure R1). For the three nutrients, F_{sw} shows high spatial variation and

longitudinal patterns, if any, are difficult to visualize. This is why we believe that showing the longitudinal variation of the frequencies (sort of “stream modes” as used in animal ecology) could be more informative to the reader.




Figure R1. In-stream net nutrient uptake (F_{sw}) for individual segments along the reach for (a) nitrate, (b) ammonium, and (c) SRP during the study period. Circles are means and whiskers are standard error for the dormant (white) and vegetative (black) period. The square at the right end of each plot is the mean F_{sw} calculated for the whole reach. $F_{sw} > 0$ indicates that gross uptake prevail over release. $F_{sw} < 0$ indicates the opposite. The zero line is shown as a reference.

(6) *We need an acknowledgment of the human activity in the lower part of the catchment. I see buildings and agricultural or horticultural activity on Google Earth. As much as 15 hectares of riparian area may be disturbed. This much activity could account for the higher N and P concentrations in the lower-most segments. If you are sure that the activity is inconsequential, you should explain this.*

Answer: Right. Thanks for raising this point. The catchment has a low population density (< 1 person/km²) which is concentrated at the valley bottom (info now included in pp 6-lines 133-135). Human settlements consist in few summer residences with some orchards, and one permanent residence with few field crops for personal consumption

(pp 9-lines 192-194). There is also a small water bottling plant between the last sampling sites (this is spring water that emerges from deep, old groundwater layers, which are disconnected from the annual hydrological water cycle).

As you point out, the potential influence of human activity on stream nutrient chemistry cannot be rule out. As already mentioned in our responses, your comment has motivated us to rescue some stream water samples collected from a small tributary that drained though the inhabited area of the catchment. We could not measure stream discharge at this site but we have characterized stream water chemistry which can now be seen in Figures 4 and 5. These data suggest that anthropogenic sources were unlikely behind the increase in nitrate concentrations observed at the valley bottom because DIN concentrations at this human-altered tributary were low. In contrast, this tributary showed high SRP concentrations (multiple-fold higher than in the main stream), and thus could partially explain the increase in SRP observed at the valley bottom. However, its discharge should have had to be ca. 4 times higher than expected for its drainage area ($< 0.4 \text{ km}^2$) to explain the observed changes in concentration (pp 22,23-lines 523-531).

In addition, we have checked out some isotopic N data available from past studies performed at this site. In particular, data from a survey carried out in 2010 at the downstream-most site of the reach showed that N isotopic signatures ($\delta^{15}\text{N}$ in ‰) were -0.44 ± 3.70 and 0.30 ± 0.54 for $^{15}\text{N-NH}_4^+$ and $^{15}\text{N-NO}_3^-$, respectively (mean \pm SE, $n = 4$ for each DIN species). These values are well bracketed by data reported for pristine streams, including upstream reaches at our study site and other streams worldwide (Pastor et al. 2013; Pastor et al. 2014; Peipoch et al. 2012). Thus, these data do not support either that the increase in DIN concentration at the downstream reaches was a consequence of human activities at the valley bottom.

(7) Fig. 5 shows no tributary data for the vegetative season. Tributaries were the major fluvial input, so these data are critical to understanding what was influencing concentrations.

Answer: You're right! The water budgets at the whole reach-scale indicate that tributaries were contributing $> 50\%$ to stream discharge. In addition, tributaries showed contrasting Cl^- concentration between the dormant and vegetative period. This info is crucial to understand the longitudinal pattern of this hydrological tracer.

In the former version, plotted data from tributaries were not separated by seasons. Now, we do so for Cl⁻ (Figure 4c) and nutrients (Figure 5). In addition, we pay more attention to tributaries' stream water chemistry in the results section.

(8) Correct units for $|F_{sw}/F_{in}|$. These would be inverse meters (like a longitudinal loss rate) except that the values are so high that I assume that F_{sw} is actually $F_{sw} \cdot x$.

Answer: That's right. Thanks for noticing! We have corrected this mistake throughout the text, tables, and figures.

The message: *I see two contradictory messages emerging from this paper. One is that in-stream biogeochemical processes can be evaluated from mass balance considerations and are large. The other is that the signal from these processes is overwhelmed by other factors—inputs, losses, riparian processes, and landscape features—so that the stream chemistry doesn't reveal what the biogeochemical processes are doing. Both can't be true. Your equation (2) -the statement of mass balance—says that the stream chemistry does reveal biogeochemical processes. It does not make sense to say that “other factors...may overcome the effect of...processing” (598:6), or that “longitudinal trends...were not consistent with...biogeochemical processes” (598:19, 618:25), or that “sources...can offset the effect of nutrient cycling on stream water chemistry” (619:5). If an in-stream process removes some nutrient then it unequivocally affects stream concentration, and the effect can be measured (the point of Eq. 2). It is not overcome, made inconsistent, or offset by other factors. The problem is the implicit equating of an effect on concentration with the presence of a longitudinal gradient. This shows up most clearly in the first sentence of the abstract and as a “hypothesis” in the discussion (614:22). More careful wording would resolve the problem: Clearly state that you are not talking about an effect on concentration but about the expectation that an effect shows up as a longitudinal gradient. Your work shows that a longitudinal gradient does not equate to an effect on concentration. It also shows that biogeochemical processes have a strong effect on concentration—that require careful mass balance to see.*

One way to clarify the message might be to focus on how in-stream processes can be large without creating a concentration gradient. The basis for this lies in Brookshire et al's (2009) equation 4: Biological uptake draws the concentration

groundwater that enters the stream laterally (presuming that it is higher than the streamwater concentration) down to the concentration of the streamwater that entered the reach. Thus the streamwater concentration shows no gradient, but is maintained lower than the concentration of its sources. Below I question whether your reported in-stream processes (F_{sw}) are too high, but regardless, it would be instructive to drill down on the high values, asking whether these were associated with segment-specific concentration declines or high groundwater inputs.

The confusion in message extends to the treatment of riparian and landscape factors as variables that explain stream concentrations. The glm model leaves out F_{sw} despite your finding that in-stream processes affect ammonium and phosphate fluxes by 29 and 30% in a single segment. Instead it includes D (the gradient) making the implicit assumption that in-stream processes can only be represented by a gradient. The image I get is that of two demons—in-stream processes and riparian characteristics—competing for control of stream concentrations. But you already know how much each is controlling, from the mass balance calculations, and don't need statistics for that. The riparian and landscape variables can affect concentrations either through affecting inputs (tributary and groundwater fluxes) or by affecting in-stream processes (e.g., wider stream increases uptake per unit distance, or opens the canopy to primary production). Because the inputs have been measured, the analysis should separate these pathways: one analysis for riparian concentrations (or inputs) and one for in-stream processing (F_{sw}). For example, terrestrial nitrogen-fixation should be related to its effects on riparian concentrations.

Answer: We agree that we had to improve the bottom line of the paper and explain clearly what our results mean. Your comments in this regard are smart and have been very helpful. Following your suggestion, we have reconsidered the wording throughout the text to avoid creating confusion and unclear messages.

In the introduction, we now explicitly address that some authors have proposed that nutrient concentrations should decline in the downstream direction if in-stream net uptake is high enough (Brookshire et al., 2009). This declining pattern is not systematically observed in reach-scale studies, which could bring us to the conclusion that terrestrial inputs are the major driver of stream water chemistry because in-stream gross uptake and release may counterbalance each other most of the time (Brookshire et

al., 2009). However, previous synoptic studies have revealed that nutrient concentrations are patchy and highly variable along the stream as a result of spatial patterns in upwelling and in-stream nutrient processing (Dent and Grimm, 1999). Thus, in-stream nutrient cycling could be substantial but not necessarily lead to longitudinal increases or declines in nutrient concentration, a question that probably needs to be addressed at spatial scales larger than few hundred meters (pp 4,5-lines 87-98).

The new version of the manuscript emphasizes the results (both former and new) showing that in-stream processes are relevant and contribute substantially to stream nutrient fluxes. Namely, (1) poor correlations between stream and riparian groundwater nutrients, (2) large $|F_{sw} \cdot x/F_{in}|$ values, and (3) relative contribution of in-stream processes to either increase (i.e., $|F_{sw} \cdot x < 0|/F_{in}$) or decrease (i.e. $F_{sw} \cdot x > 0/F_{in}$) stream nutrient fluxes.

The comparison of these results with the longitudinal patterns in stream nutrient concentrations highlights that in-stream processes can be important without generating longitudinal gradients in nutrient concentration, as you point out. The case of NH_4^+ is paradigmatic: concentrations were consistently lower in the stream than in riparian groundwater and $F_{sw} > 0$ predominated (gross uptake > release). Yet, stream NH_4^+ concentrations showed small longitudinal variation likely because in-stream net uptake balances the elevated inputs from riparian groundwater. Thus, our results challenge the idea that stream nutrient concentrations should decrease in the downstream direction when in-stream processes are efficient in taking up nutrient from receiving waters, as highlighted by you (pp 23,24-lines 544-556).

In the conclusions section, we now remark that dissolved nutrients underwent biogeochemical transformation while travelling along the stream and further, in-stream processes contributed substantially to modify stream nutrient fluxes. In addition, we found that the influence of in-stream processes on stream nutrient concentrations and fluxes did not translate into gradual changes in nutrient concentration along the reach because the longitudinal pattern in nutrient concentration resulted from the combination of both in-stream cycling and nutrient inputs from terrestrial sources. Thus, one of the implications of our study is that the assessment of these two sources of variation of stream nutrient chemistry is crucial to understand the contribution of in-stream processes to stream nutrient dynamics at relevant ecological scales (pp 24,25-lines 576-599).

Finally, as mentioned in our previous responses, we have decided to eliminate the GLM analysis from the paper. As you suggest, calculating whole-reach budgets is a more simple and elegant way to put in-stream cycling and riparian groundwater inputs into the same picture and thus, we believe that these new results are more helpful to address the overall aim of the study.

In-stream process estimates: *The F_{sw} and $|F_{sw}/F_{sin}|$ values for ammonium and phosphate are surprisingly large (except, perhaps for nitrate), so large that I suspect an error. What follows is an explanation of why I suspect an error. I may be wrong, but either way, I think the paper should provide the data and considerations needed to answer the questions I raise here—because other readers would surely raise them as well.*

F_{sw} for ammonium is reported to average 0.6 ± 0.2 (SE) $\mu\text{g}/\text{m}/\text{s}$ which, for the 3700-m reach is a net uptake of 2220 $\mu\text{g}/\text{s}$. The inputs to the reach from upstream are roughly 200 $\mu\text{g}/\text{s}$ and the downstream export is about 800 $\mu\text{g}/\text{s}$, for a net export of 600 $\mu\text{g}/\text{s}$. Thus an input of 2800 $\mu\text{g}/\text{s}$ from lateral (tributary and groundwater) sources is needed to balance the budget. The net inflow of water from these sources is about 70 L/s, so the average concentration of the lateral sources would have to be about 40 $\mu\text{g}/\text{L}$. This is far higher than the reported concentrations for both tributary and groundwater, especially tributary which supplies most of the water. The needed concentration may be lower than 40 $\mu\text{g}/\text{L}$ if there is a large exchange of groundwater but this exchange would have to be quite large. Thus I suspect that the F_{sw} of 0.6 $\mu\text{g}/\text{m}/\text{s}$ is erroneously high. Based on the data available in the paper I calculate that the average F_{sw} could be no higher than 0.2 $\mu\text{g}/\text{m}/\text{s}$, even allowing for substantial additional gains through groundwater exchange. (This conclusion could change if vegetative-season tributary inputs, missing from Fig. 5, were much higher than in the dormant season). Also, the mean F_{sw} of 0.6 $\mu\text{g}/\text{m}/\text{s}$ translates to an areal net uptake (U) of about 0.27 $\mu\text{g}/\text{m}^2/\text{s}$, which seems implausibly high, at least for a mean. It is far greater than the median of 0.03 $\mu\text{g}/\text{m}^2/\text{s}$ reported by von Shiller et al. 2011. And it is about half the gross uptake typical of second and third order reaches (Ensign and Doyle 2006 JGR 111:G04009), which we expect to be much higher than net uptake because retention in stream biota occurs on a timescale of days or weeks.

My concerns are similar for phosphorus, for which the mean Fsw = 0.2 ± 0.4 ug/m/s. In this case the large SE indicates that the net uptake is not statistically significant, yet a net uptake is claimed (598:16). Using the whole reach mass balance approach that I used for ammonia above, I find that the average FSW of 0.2 ug/m/s is reasonable only for the dormant period, and only if the high downstream concentrations are ignored (in which case P concentrations are longitudinally uniform). If the downstream concentrations are included, then I estimate FSW at -0.20, i.e., a net P release. This matches your lower error bound, but because this is simple arithmetic and not statistical sampling, your average should be correct. The large apparent release of P comes from the lower 3-4 stations. If we consider only the reach upstream of 3000 m, then I estimate a net uptake of about 0.1 ug/m/s. I did not check the vegetative period, but unless the tributary concentrations were much higher than in the dormant period, there is no way that there could be a significant net uptake in the vegetative period, over the whole reach or even the upper 3000 m.

For nitrate, I'd say that your negative Fsw is entirely driven by the high downstream concentrations in the vegetative period. I get a net uptake for the dormant period, and a net uptake for both periods in the upper 3000 m.

One source of error may be in the measurements. The mass balance calculations depend on accurate measures of both concentration and flow and depend on small differences in sometimes small numbers. The errors are made worse by multiplying these estimates together. Some evaluation of the role such errors should be included. Error may also come from averaging Fsw (assuming you have divided by x). The segments are unequally weighted with the fourth segment contributing only one-quarter as much as it should.

Answer: Tones of thanks for taking the time and patience to calculate the nutrient budgets. Your exercise has been extremely useful for us and we have enjoyed and learn while following your rationale.

Your back-of-the-envelope calculations have helped us to sort out why inputs and outputs were not matching. Our main mistake was using an erroneous set of distances between sampling sites. In the former version, all segments summed up to 2400 m but the reach is 3691 m long! It's a pity we did not realize before, and it's great that you highlight this issue. This mistake would have been obvious to us if we would have calculated budgets for the whole reach from the very beginning. Now, we have rechecked the whole data base and recalculated nutrient budgets for individual

segments. Further, we have calculated budgets for whole reach; these new results are included in the manuscript. As shown in Table R3, there is a good match between inputs and outputs.

Table R3. Mean values of whole-reach nutrient fluxes for the study period. Nutrient fluxes are as in new Figure 3. Nutrient inputs to the reach are upstream sources (F_{top}), net riparian groundwater inputs (F_{gw}) and tributaries (F_{ef}). Nutrient downstream export is F_{bot} . In-stream processes can either contribute to input ($F_{sw} \cdot x < 0$) or output ($F_{sw} \cdot x > 0$) stream nutrient fluxes. For the three nutrients, mismatches between mean inputs and outputs were < 4%.

	Flux ($\mu\text{g/s}$)		
	N-NH ₄ ⁺	N-NO ₃ ⁻	SRP
F_{top}	237	6863	170
F_{bot}	1058	23633	1530
$F_{sw} \cdot x$	1362	-2045	-337
F_{gw}	1719	9122	476
F_{ef}	556	6261	609
Inputs	2512	24291	1592
Outputs	2420	23633	1530
dif (%)	3.7	2.7	3.9

Overall, the recalculated in-stream net nutrient fluxes are close to your back-of-the-envelope calculations. Regarding NH₄⁺, median F_{sw} during the study period was 0.17 and 0.2 $\mu\text{g N m}^{-1} \text{s}^{-1}$ by segment and whole reach, respectively (new Table 3). If we consider a mean wetted of 240 m and a reach length of 3691 m (8860 m² of streambed area), the estimated in-stream net uptake areal rate is 0.03 $\mu\text{g N m}^{-2} \text{s}^{-1}$ which is in the range of values reported for streams nearby our study site (-0.4 to 0.5 $\mu\text{g N m}^{-2} \text{s}^{-1}$, von Schiller et al., 2011).

Regarding SRP, median F_{sw} was -0.06 and 0 $\mu\text{g P m}^{-1} \text{s}^{-1}$ by segment and whole reach, respectively (new Table 3). This result suggests that in-stream gross uptake and release tended to counterbalanced each other, or at least that positive and negative values of F_{sw} were of similar magnitude. We have calculated budgets separately for the first 3 km (sampling sites from 1 to 10) and the last 0.7 km (sampling sites from 11 to 15). This last 0.7 km is the section that could be most influenced by human activities and its riparian zone is the most affected by the invasion of black locust. As you pointed out, these factors could be behind the increased stream NO₃⁻ and SRP concentrations at end of the reach. As required, here we show F_{sw} separately for these two sections of the reach (Figure R2). If nutrient budgets are calculated in this way, the obtained values

follow your predication in that $F_{sw,SRP}$ was higher for the upper than for the lower section of the reach (0.005 vs -0.28 $\mu\text{g P m}^{-1} \text{s}^{-1}$), though differences were not significant (Wilcoxon paired rank sum test, $p = 0.1$).

Regarding NO_3^- , median F_{sw} was -0.43 and -0.97 $\mu\text{g N m}^{-1} \text{s}^{-1}$ by segment and whole-reach, respectively. This rate equals to -0.07 $\mu\text{g N m}^{-2} \text{s}^{-1}$, a value within the range reported by von Schiller et al. (2011) (-3.5 to 5 $\mu\text{g N m}^{-2} \text{s}^{-1}$). As you predicted, F_{sw,NO_3} was higher for the upper than for the lower section of the reach (0.61 vs. -3 $\mu\text{g N m}^{-1} \text{s}^{-1}$). In this case, differences were statistically significant, at least for a $p < 0.05$ (Wilcoxon signed-rank test, signed-rank = 19.5, $df = 9$, $p = 0.049$). These differences occurred mainly during the vegetative period, and no statistical differences were found for the dormant period, as you expected. Thus, you were correct in that in-stream net uptake for NO_3^- prevailed over release for the upper-most segments, though its contribution to reduce stream NO_3^- flux was relatively small 10% [0%, 20%] (median [IQR]) compared to that of NH_4^+ 42% [24%, 62%] (Figure R2). We have we decided not to include all these results in the paper to avoid redundancies, though we refer to these values when discussing about the influence of in-stream processes on stream NO_3^- concentrations and fluxes (pp 24-lines 558-565).

Finally, we agree that we should take into account the uncertainty associated to these measurements. Following suggestions by you and R1, we have calculated the empirical uncertainty associated to stream discharge (see more detailed explanations in our responses to R1), and we have considered it when calculating water and nutrient fluxes and the frequency of cases for which $F_{sw} > 0$, < 0 , and $= 0$. In addition, we have analyzed whether differences in F_{sw} and $|F_{sw} \cdot x / F_{IN}|$ were statistically significant between nutrients which is also a way to consider the uncertainty associated to these estimations. In any case, we would like to highlight that differences in F_{sw} are pretty similar when calculated either by segment or whole reach, indicating that our estimates are reliable despite the fact that F_{sw} for individual segments may be unequally weighted.




Figure R2. Whole-reach in-stream nutrient uptake (F_{sw}) for the first 3-km (gray) and the last 0.7-km section of the reach (blue) for (a) nitrate, (b) ammonium, and (c) SRP. Whiskers are the uncertainty associated with the estimation of stream discharge from slug tracer additions. $F_{sw} > 0$ indicates that gross uptake prevail over release. $F_{sw} < 0$ indicates the opposite. For those cases for which $F_{sw} > 0$, the contribution of in-stream net nutrient uptake to reduce stream nutrient fluxes ($F_{sw} \cdot x > 0/F_{in}$, in %) is shown (same color code as for F_{sw}).

Groundwater and the riparian interface: *The conceptual framework is not entirely clear. The use of mass balance requires a boundary within which processes are measured (as F_{sw}). For clarity I would suggest the following: The boundary is the riparian-stream edge, the stream channel is inside, and the riparian zone is outside. Fluxes pass through the stream edge, but no processes can occur “at the stream edge” (cf.617:3), only on one side or the other. In re-reading, I see that your suggestion that the apparent downstream nitrate source represents nitrification of riparian ammonium that occurs “at the riparian-stream edge” (617:14). From the standpoint of the Eq. 2, this process occurs inside the boundary, i.e., in the stream channel, the nitrogen having crossed the boundary as ammonium. So you could define “stream-edge” this way, clarifying that it is inside the riparian-stream boundary. If this nitrification actually*

explains the downstream increase in nitrate concentration, you should be able to demonstrate that Fgw of ammonium was sufficient to account for the negative Fsw of nitrate in those segments.

As you formulated the mass balance, all groundwater fluxes to and from the stream pass through the riparian zone. If hillslope groundwater is to explain phosphorus concentrations in the stream (616:1-20), it must pass through the riparian and would have been measured there. The possibility that hillslope water flows bypasses the riparian, e.g., via underflow, to reach the stream is not consistent with Eq. 2. That is, to invoke hillslope groundwater you must acknowledge that the mass balance is incorrect. Regardless of how you handle the hillslope question, you should explicitly address the question of whether the riparian well samples correctly characterized the groundwater that reached the stream.

Is there an explanation as to why the chloride differed between periods even though the flow did not? At 614:13 you suggest that evapotranspiration (ET) may explain the higher chloride in the vegetative period, which makes sense except that the greater ET should also have reduced flow. Higher precipitation during the vegetative season, if it occurs, could compensate for the additional ET, explaining the similar flows.

The 40% longitudinal increase in chloride seems higher than can be explained by the decrease in area specific discharge and hence by ET. In support of the ET suggestion you state that chloride was higher in the main stream than in the tributaries (614:11). But Fig. 3b shows the opposite: Tributary chloride was higher than the mainstream and this, rather than ET, appears to explain much of the increase.

Answer: As mentioned in our earlier responses, we agree that the conceptualization adopted in this study needs to be clearly defined. Consequently, we have included a schematic representation of our mass balance approach (new Figure 3). Following your suggestion, we have kept our conceptualization as clear and simple as possible, defining two compartments: the stream and the riparian zone. Moreover, we have removed ambiguous wording throughout the manuscript such as referring to biogeochemical transformation at the riparian-stream edge, which does not match with our conceptual model, as highlighted by you.

Following your suggestion, we have calculated whether increased stream NO_3^- concentration at the valley bottom during the vegetative period could be explained by nitrification of NH_4^+ entering from riparian groundwater. Our results suggest additional

sources of nitrate within the stream because the flux of NH_4^+ from riparian groundwater, $F_{gw,NH4/x} \sim 2 \mu\text{g N m}^{-1} \text{ s}^{-1}$, was not large enough to sustain in-stream NO_3^- release, $|F_{sw,NO3} < 0| \sim 10 \mu\text{g N m}^{-1} \text{ s}^{-1}$ (pp22-lines 511-515). Discussion about these potential additional sources can be read in pp 22,23-lines 515-543 (see below our responses to your specific comments in this regard).

We agree with you, as well as with R1 and R2, that considering the influence of hillslope groundwater on stream water chemistry independently of riparian groundwater is confusing and further it violates the assumptions underlying the mass balance. In the new version of the manuscript, we have deleted this part from the discussion as well as former Figure 8. Further, the whole reach-scale nutrient budgets have shed new light about catchment sources contributing to stream SRP. According to our mass balance calculations, tributaries and riparian groundwater were the major contributors. In-stream processes further contributed to increase stream fluxes (new Table 4 and Figure 7).

As already mentioned, the concluding remarks do now explicitly address the issue of whether water samples from riparian wells correctly characterized the groundwater that reached the stream (pp 25,26-lines 600-608).

As mentioned in our responses to R1 and R2, the easiest explanation for the increase in stream Cl^- during the vegetative period is that concentrations were higher in tributaries than at the main stream, in particular during the vegetative period (the former interpretation was wrong). According to whole reach budgets, permanent tributaries were the major contributor to stream discharge (56%) and therefore, hydrological mixing with water from tributaries could partially explain the 40% increase in stream Cl^- concentrations observed along the reach as well as the higher concentrations during the vegetative period. In addition, riparian groundwater inputs could also contribute to these patterns because this compartment was an important source of water to the stream (26%) and it exhibited higher Cl^- concentration than stream water during through the year. These new results and the associated discussion are included in the new version of the manuscript (results section 4.1 and pp 19,20-lines 447-455).

Regarding the seasonality in P and Q, a more detailed look to data collected from our permanent sampling station has highlighted that both P and Q followed a seasonal pattern with minima in summer 2011, and further that Q at base flow conditions was higher during the dormant than during the vegetative period (Table R1). Yet, P was similar between the vegetative and dormant period mainly because spring 2011 was rainy and this rain counterbalanced the following dry summer. In addition,

similar Q between the dormant and vegetative period for the synoptic survey were likely a consequence of the small sample size and the fact that the highest (April 2011) and the lowest flows (September 2011) fall within the vegetative period.

Temporal variation: *The absence of a temporal perspective (beyond the dormant/vegetative classification) is surprising. Most streams exhibit distinct seasonal variations in both flow and nutrient chemistry, and these patterns have proven critical to understanding biotic influences on stream chemistry (e.g., Mulholland et al.'s many papers on Walker Branch). If neither precipitation nor flow vary with season, this should be clearly stated. Otherwise we should at least be informed of the seasonal patterns and, ideally, your analysis would make use of them. You do state that dormant and vegetative flows were similar. However, if precipitation is uniform we would expect flow to reach a minimum at the end of the vegetative period and a maximum at the end of the dormant period, with the result that the average of the two seasons is the same.*

Answer: The reviewer is right. Our synoptic survey was not large enough to allow splitting the data set by calendar seasons, and by splitting the data between vegetative and dormant period the seasonality was screened in some cases, such as the case of Q.

As mentioned in p-#4 (R2), our high-temporal resolution data indicated that stream discharge was highest in spring and autumn 2011 and lowest in summer 2011 (Table R1). Following the suggestion of both reviewers, we now provide information about the temporal variation in precipitation and stream discharge in the text as well as in new Figure 2.

Page:line-referenced comments:

610:22. *“According to the glm results, this decrease was positively related to basal area...” etc. These are the opposite of what is given in Table 3, which shows how these factors would relate to an increase, not a decrease. According to Table 3, basal area increases nitrate, but % N-fixers decreases it. In the GLM the independent variables clearly suffer from collinearity, as they all vary consistently with downstream direction. Collinearity leads to erratic regression coefficients, which may explain some of the seemingly odd relationships.*

Answer: Yes, right. As pointed out by R1, the statistical analysis of this sort of data set requires very careful consideration. This has been one of the reasons why we have decided to delete the GLM analysis from the new version of the manuscript.

614:17. “...suggesting...a gaining reach”? You should know from your flow measurements whether it was or was not gaining.

Answer: Yes, that’s right. As already mentioned, we have reanalyzed our data set in order to improve the hydrological characterization of the study reach. Please, see our previous responses to you and R1 in this regard. New results are shown in Figure 2c and d, and Figure 4b. The new discussion on these results can be read in pp 19-lines 439-444).

614:23. ” We found a decreasing longitudinal pattern of stream NO₃⁻ concentrations, though only during the dormant period.” Fig. 5 clearly shows that nitrate decreased in both seasons above 3000 m. Don’t let the regression model obscure reality.

Answer: That’s right. We now clearly state the U-shaped pattern exhibited by stream NO₃⁻ concentrations in the results section. In the discussion, we indicate that stream NO₃⁻ concentrations showed a decreasing longitudinal pattern along the first 1.5 km (regardless of the period) (pp24 –lines 558-561).

614:27-615:2 I disagree. It makes little sense to ignore the clear change in nitrate gradient that occurred at 3000 m, in both periods. It looks to me like in-stream processes were important for the downstream decline above 3000 m. I would say $|F_{sw}/F_{in}|$, at 11% (per segment, right?), is large. Positive values were much more frequent than negative values in the 11 segments above 3000 m. Tributary concentrations were low, so also contributed to the decline. You should calculate how much was instream and how much was dilution.

Answer: Thanks for raising this point. We agree that some further work was needed. Following the suggestions from the three reviewers, the U-shaped pattern exhibited by stream NO₃⁻ concentrations is now clearly stated.

We have calculated separately the nutrient budgets for the upper and lower sections of the reach (first 3 km vs. last 0.7 km, respectively). As included in our earlier responses, F_{sw,NO_3} was positive for the upper than for the lower section of the reach (0.61 vs. -3 $\mu\text{g N m}^{-1}\text{s}^{-1}$) (Wilcoxon signed-rank test, Signed-rank = 19.5, df = 9, p =

0.049). Thus, you are essentially right in that in-stream net uptake prevailed over release for NO_3^- along the upper section and in-stream processes could contribute to reduce stream nitrate flux. According to our calculations, this contribution could explain 10% [0%, 20%] of the reduction in stream NO_3^- flux along the first 3 km of the reach.

In addition, hydrological mixing with water from tributaries could also contribute to decrease stream NO_3^- concentrations because NO_3^- was lower in the tributaries. We have calculated the expected stream NO_3^- concentration if the upstream-most tributary would have explained the change in stream NO_3^- concentration between the 4th and the 5th sampling sites. For the 11 sampling dates, the relative difference between measured and expected NO_3^- concentration ranges from -45 to +18% (mean = -12%). Considering the uncertainty associated to these calculations, there were only 2 out of 11 dates for which hydrological mixing could explain 100% of the change in NO_3^- concentration. In the other 9 cases, the combination of in-stream net uptake and dilution were likely responsible for the observed longitudinal decrease. We have not included all these calculations in the manuscript, though we now clearly state in the discussion that both in-stream cycling and hydrological mixing could explain the observed declining pattern for NO_3^- (pp 24-lines 558-567).

615: 18 The lack of correlation with N_2 -fixing species does not rule out their possible influence because there was an opposite nitrate gradient in the region upstream of 3000 that would have obscured the correlation in the lower reaches.

Answer: Yes, right. Following suggestions from the three reviewers, we have rewritten this part of the discussion. In particular, we highlight that the flux of NH_4^+ from riparian groundwater is not large enough to sustain in-stream NO_3^- release during the vegetative period along the last 700 m of the reach (pp 22-lines 511-515). Among several potential explanations for such additional source of stream N, we pointed out leaf litter from riparian trees, and especially from N_2 -fixing species, that combined with warm temperatures could enhance in-stream mineralization and nitrification during the vegetative period (pp 22,23-lines 515-522).

615: 22 Release of nitrate from the streambed raises the question of the ultimate source of the nitrogen. Simple storage and release cannot produce a net release; the high concentrations at one time would have to be balanced by low concentrations at another. On 616:14-18 you suggest that the source may be nitrified ammonium. Perhaps that

should be mentioned here. Another possibility is instream N fixation. See Finley et al. 2011 Ecol. 92:140 who saw a sharp downstream increase due to Nfixation. Also, as mentioned above, I am concerned about human sources.

Answer: Right. We have reorganized the discussion section as suggested. After noticing that the flux of NH_4^+ from riparian groundwater is not high enough to sustain in-stream NO_3^- release along the last 700 m, we propose several additional sources: (1) in-stream nitrification enhanced by large stocks of riparian leaf litter (from N_2 -fixing species in particular), (2) human sources of dissolved N (3) autotrophic N fixation. The feasibility of these different sources is discussed in light of the evidences we have (pp22,23-lines 515-537).

616:1 Stream fluxes did not decrease downstream; what decreased was area-specific flux. The absolute nutrient fluxes increased greatly.

Answer: Right, thanks. This sentence has been deleted from the discussion, though.

616:3 "...concentrations increased from the top to the bottom of the reach for all nutrients (except for NO_3 during the dormant period." Nitrate did not increase in the vegetative period either, and ammonium did not increase in the vegetative period.

Answer: Right, thanks. This sentence has been deleted from the discussion.

617:13 "especially during the vegetative period" should read "although only in the vegetative period".

Answer: Yes. Thanks (pp22-line 507).

617:25-28. This sentence does not quite make sense. The first part refers to "predominant" in-stream processes while the second part effectively states that there was not a net uptake or release along the reach, i.e., that there was no "predominant" process. As I have argued, it is fallacious to say that a longitudinal trend is consistent or in-consistent with in-stream processing, as it is a matter of balance with lateral inputs. I think that for nitrate above 3000 m, you had both net uptake and a declining trend although low tributary concentrations may also have contributed to the decline. For ammonium and phosphate above 3000 m you had net uptake and no decline because the uptake drew down the concentrations of lateral inputs, i.e., uptake balanced lateral inputs. In my view this is a common case in streams where there is a net uptake.

However, this is hard to document from the literature, Meyer and Likens (1979) and Alexander et al. (2009 Biogeochemistry 93:91) being the only two examples I can readily find. This is why your study is so important.

Answer: Yes, thanks. We have rewritten this paragraph following several of the points highlighted in your review. As already included in earlier parts of this letter, (1) we don't refer to "predominant" in-stream processes because F_{sw} was highly variable as it is now explicitly included in the discussion (pp 20-line 470), (2) we have calculated nutrient budgets separately for the upper and lower part of the reach, (3) we have calculated the contribution of tributaries to stream nutrient fluxes for the whole-reach, and (4) we have calculated whether riparian groundwater fluxes of NH_4^+ at the end of the reach could sustain the measured in-stream NO_3^- release.

With these new pieces of information, we now discuss that the declining pattern exhibited by NO_3^- , especially along the first 1.5-km could be explained by both in-stream net NO_3^- uptake (ca. 10%) and hydrological mixing with water from tributaries (pp 24-lines 558-567). Conversely, the marked increase in stream NO_3^- along the 700 km may be explained by in-stream mineralization and nitrification of organic matter accumulated within the stream. Human sources of DIN and autotrophic N fixation seem not plausible explanations (pp22,23-lines 515-537).

As you pointed out, our results for NH_4^+ indicated that in-stream net uptake balanced elevated inputs from riparian groundwater. Thus, the case of NH_4^+ is paradigmatic because it convincingly shows that in-stream processes can strongly affect stream nutrient chemistry and downstream nutrient export without generating longitudinal gradients (pp 23-lines 544-553).

For SRP, the importance of in-stream net uptake above 3 km is not so clear. If nutrient budgets are calculated separately for the upper and lower sections, $F_{sw,SRP}$ is higher for the first 3 km than for the last 0.7 km (0.005 vs $-0.28 \mu\text{g P m}^{-1} \text{s}^{-1}$). However, these differences are not significant (Wilcoxon paired rank sum test, $p = 0.1$), and $F_{sw,SRP}$ is close to zero even for the first 3-km of the reach. Along the last 0.7- km, mixing with the downstream-most tributary could partially explain the increase in stream SRP concentrations. Yet, on average, the discharge at this tributary should be 4 folds higher than expected for its drainage area to fully explain the observed increase in SRP (pp 22,23-lines 526-529).

618:14 delete “through” (an apparent typo) and “profound” (exaggeration).

Answer: Ok. Thanks.

619:19 “can offset the effect of” should read “must be taken into account when interpreting longitudinal gradients”.

Answer: Agree. Thanks. We have reworded the text to avoid this sort of confusion in our message.

REFERENCES

- Brookshire, E.N.J., Valett, H.M., and Gerber, S.G. Maintenance of terrestrial nutrient loss signatures during in-stream transport. *Ecology*, 90, 293-299, 2009.
- Covino, T.P., McGlynn, B.L., and Baker, M. Separating physical and biological nutrient retention and quantifying uptake kinetics from ambient to saturation in successive mountain stream reaches. *J. Geophys. Res.*, 115, G04010, doi:10.1029/2009/JG001263, 2010.
- Dent, C.L., and Grimm, N.B. Spatial heterogeneity of stream water nutrient concentrations over successional time. *Ecology*, 80, 2283-2298, 1999.
- Lowrance, R., Altier, L. S., Newbold, J. D., Schnabel, R. R., Groffman, P. M., Denver, J. M., Correl, D. L., Gilliam, J. W., Robinson, J. L., Brinsfield, R. B., Staver, K. W., Locas, W., and Todd, A. H. Water quality functions of riparian forest buffers in Chesapeake Bay watersheds. *Environ. Manag.*, 21, 687-712, 1997.
- Pastor, A., Peipoch, M., Cañas, L., Chappuis, E., Ribot, M., Gacia, E., Riera, J. L., Martí, E., and Sabater, F. Nitrogen stable isotopes in primary uptake compartments across streams differing in nutrient availability. *Environ. Sci. & Tech.* 47:10155-10162 doi:10.1021/es400726e, 2013.
- Pastor, A., Riera, J. L., Peipoch, M., Cañas, L., Ribot, M., Gacia, E., Martí, E., and Sabater, F. Temporal variability of nitrogen stable isotopes in primary uptake compartments in four streams differing in human impacts. *Environ. Sci. & Tech.*, 48:6612-6619 doi:10.1021/es405493k, 2014.
- Payn, R. A., Gooseff, M. N., McGlynn, B. L., Bencala, K. E., and Wondzell, S. M. Channel water balance and exchange with subsurface flow along a mountain headwater stream in Montana, United States. *Water Resour. Res.*, 45, W11427, doi:10.1029/2008WR007644, 2009.
- Peipoch, M., Martí, E., and Gacia, E. Variability in delta N-15 natural abundance of basal resources in fluvial ecosystems: a meta-analysis *Freshw Sci* 31:1003-1015 doi:10.1899/11-157.1, 2012.
- von Schiller, D., Bernal, S., and Martí, E. A comparison of two empirical approaches to estimate in-stream net nutrient uptake. *Biogeosciences*, 8, 875-882, 2011.
- Zar, J. H. *Biostatistical analysis*. Prentice-Hall/Pearson, Ed. 5th Edition. Upper Saddle River, N.J., 2010.