

## ***Interactive comment on “Oceanic N<sub>2</sub>O emissions in the 21st century” by J. Martinez-Rey et al.***

**Anonymous Referee #1**

Received and published: 12 January 2015

This paper presents a model simulation, using NEMO-PISCES, aimed at predicting how oceanic N<sub>2</sub>O emissions and storage will change over the next century in the face of decreasing export production, increasing water column stratification, and declining interior O<sub>2</sub> content. I enjoyed the introductory discussion, which raised important issues and motivated the current study in a compelling way. My main concern is whether the framework of the 2 major N<sub>2</sub>O production pathways used here, O<sub>2</sub>-independent ammonia oxidation, and the low O<sub>2</sub> pathway at levels < 5 μmol/L is adequate to describe the complexity of the oceanic N<sub>2</sub>O cycle, especially considering that the extent of oxygen minimum zones in the global ocean is poorly captured by NEMO-PISCES. There is essentially no discussion of nitrifier denitrification, which can be important at more modestly depleted O<sub>2</sub> levels well above 5 μmol/L and may be responsible for the bulk of oceanic N<sub>2</sub>O production. For those who believe that much of oceanic N<sub>2</sub>O production occurs in and around OMZs (e.g., see work by Codispoti), the P.TEMP and

C8023

P.OMZ formulations are unsatisfactory as independent parameterizations that encompass the full range of possible future oceanic N<sub>2</sub>O response. Both parameterizations are heavily weighted toward nitrification, with at least 75% of total N<sub>2</sub>O production occurring via ammonia oxidation. As a result, there is a lack of significant variability in some aspects of the results, e.g., in Figure 1d.

That said, given the current state of knowledge, the authors have done a reasonable job with the information and modeling tools available, and it seems unreasonable to insist upon a complete overhaul of the modeling approach. I therefore recommend publication with minor editorial revisions, aimed primarily at acknowledging the uncertainty associated with the potentially incomplete and overly simplified representation of the oceanic N<sub>2</sub>O cycle in the model. In particular, I would like to see some discussion of the fact that the current model is unable to predict what might happen to future N<sub>2</sub>O emissions if much of N<sub>2</sub>O production does indeed occur in association with the OMZs. While the Conclusion does acknowledge some of these points already, they could be emphasized more strongly throughout the paper.

Below are some more detailed comments.

p16705, line 10. N<sub>2</sub>O is destroyed about 90% by photolysis, 10% by O(1D), but not really by the OH radical.

p16705, line 13. Change “atmosphere that caused” to “atmosphere, which has caused”

p16706, line 2. The most recent of these citations is from 2004. It would be good to include more recent work, e.g., by Westley, Farias, Frame, etc.)

p16707, line 24. Please support this statement with a reference: “Ocean warming might increase the rate of N<sub>2</sub>O production during nitrification”

p16707, line 29. “could substantially affect denitrification and the N<sub>2</sub>O production.” Better as something like, “could substantially affect N<sub>2</sub>O production via both nitrifier denitrification and classic denitrification.”

C8024

P16708, line 1. Instead of “Models” it might be better to use a more specific term like “Ocean biogeochemistry models”

P16709, line 10. A concern about the Zamora et al. analysis, which is used to justify the near-linear N<sub>2</sub>O yield ( $\gamma$ ) in the P.TEMP formulation, is that this analysis was based on  $\Delta$ N<sub>2</sub>O vs. AOU relationships at depth, representing the integrated effects of N<sub>2</sub>O production and O<sub>2</sub> consumption in old water parcels. However, in the NEMO-PISCES model, the relationship is applied to  $JN_2O = f(JO_2)$ , i.e., the instantaneous production and consumption rates, which may be significantly more nonlinear. Further, the Zamora analysis excluded all data above 150m depth, but this may be where the bulk of N<sub>2</sub>O production is actually occurring, i.e., at the base of the euphotic zone, much of which may quickly ventilate to the atmosphere (see, e.g., Popp et al., GBC, vol.16, no.4, 2002). Please acknowledge or discuss this point.

p16711 line 15. “We assume a constant atmospheric N<sub>2</sub>O concentration of 284 ppb in all simulations.” It would be good to add a clause clarifying that this value is only slightly above the natural, preindustrial N<sub>2</sub>O concentration. Also, perhaps explain why 284 ppb was chosen, considering that this paper deals with 21st Century projections, in which N<sub>2</sub>O may rise well above 325 ppb, approaching 350 or even 400 ppb.

P16712, lines 1-4. “This assumption is based on growing evidence that nitrification is the dominant pathway of N<sub>2</sub>O production on a global scale, based on estimations considering N<sub>2</sub>O production along with water mass transport (Freing et al., 2012).” I don’t think this can be taken as an accepted fact. Other lines of evidence, e.g., based on isotopes, suggest that denitrification (including nitrifier denitrification) is responsible for most N<sub>2</sub>O production (e.g., Park et al., Nature Geoscience, DOI: 10.1038/NNGEO1421, 2012.)

p16712-13, Section 3.1. Some further discussion of model shortcomings would be useful. Figure 1 shows a tendency to overestimate the N<sub>2</sub>O flux in the North Atlantic and to underestimate the N<sub>2</sub>O flux in hot spots of N<sub>2</sub>O production such as the ETSP

C8025

and ETNP. The Nevison et al., 1995 map, which is used to evaluate the NEMO results, also tends to underestimate the flux in the ETSP and ETNP, due to lack of surface pN<sub>2</sub>O data in these regions in the original Weiss dataset, but even so, captures substantially higher N<sub>2</sub>O emissions from the ETNP than the NEMO model, as shown in Figure 1d. (See Nevison et al., GBC, vol. 18, 2004 for further discussion.) Collectively, the NEMO results could be interpreted to show an overestimate of N<sub>2</sub>O production from widely distributed nitrification (i.e., ammonia oxidation) sources and an underestimate of N<sub>2</sub>O production from nitrifier denitrification and denitrification sources in lower O<sub>2</sub> regions.

p16714, line 9. “P.OMZ shows a good correlation with the observations” doesn’t seem like an accurate statement. The shape of the depth profile is considerably off from MEMENTO, although the maximum values in the 500-900 m depth range are in fairly good agreement.

p16714, line 11 and subsequent discussion. “Below1500m, both parameterizations simulate too high N<sub>2</sub>O compared to the observations.” An alternative explanation from those given is that the coefficient assigned to N<sub>2</sub>O production at high O<sub>2</sub> is too high.

P16714, line 22-23. Neither/nor should be either/or

Figure 3 and Section 3.2. It seems from this analysis, esp. the bar graph comparing to WOA, that NEMO-PISCES doesn’t capture any of the OMZs in the world oceans – there is almost no volume with O<sub>2</sub> < 50  $\mu$ mol/L !! This is mentioned only briefly as a “deficient representation of the OMZs” in a way that downplays the potential scope of the problem. Given that the jury is still out on the question of how important the OMZs are to global N<sub>2</sub>O production, the lack of OMZs in NEMO-PISCES raises serious questions about whether this model can be trusted to predict N<sub>2</sub>O emissions in the present let alone the future. This problem needs further discussion, and it would be good to provide a global map either in the supplement or main text of the O<sub>2</sub> concentration at the depth of the water column O<sub>2</sub> minimum (or else at some appropriate fixed depth),

C8026

comparing model to WOA.

P16716, lines 5-7 "In particular, the P.TEMP parameterization projects a larger enhancement of the flux than P.OMZ at the BUS, whereas the emissions in the Southern Ocean are enhanced in the P.OMZ parameterization." Please explain further why this happens, since the Southern Ocean is far removed from the OMZs.

P16717, line 3 "As the N<sub>2</sub>O production in THIS pathway" I am confused about which pathway is being discussed. I presume high, but this is unclear as written.

P16718, line 7-9, "Overall these changes are negative, and happen to nearly completely compensate the increase in production in the OMZs, resulting in the near constant global N<sub>2</sub>O production by the low-O<sub>2</sub> production pathway up to year 2100" Yes, but please put this in the context that NEMO-PISCES strongly underestimates the global volume of the OMZs.

p. 16718, section 4.2.2. Please state the absolute value of the inventory to put these changes into context.

p. 16719, lines 6-8. This sentence seems at odds with Figure 7, in which inventory is mainly increasing while production decreases. If this is not the case, then please explain more clearly in the caption whether a bar to left of center = decrease and a bar to right of center = increase (which is what I assumed for lack of other information).

p. 16719, lines 11-14, This sentence also seems at odds with Figure 7. "Figure 7 shows how almost all the relevant changes in N<sub>2</sub>O production and storage are related to low-latitude processes, with little or no contribution from changes in polar regions."

p. 16720, discussion of box model. It would be helpful to provide a better explanation of why this model is presented and whether it's really worth including in the paper. What questions does it address that cannot be answered with the 3D NEMO-PISCES model? Also, in Figure 8, what criteria are used to define the range of the box model parameters? Are some 3D models really predicting decreases of up to 80% in mixing?

C8027

P16723, Section 6. I found this section confusing and am not sure it adds to the value of the paper. The back-of-the-envelope calculations presumably reflect the indirect result of temperature on stratification and export production, but they also could be interpreted as a direct response of N<sub>2</sub>O production as a function of temperature, given the formulation of P.TEMP. Overall, the calculation is fraught with so much uncertainty that it in my opinion should be deleted.

P16724, line 27. For balance, it might be worth mentioning that other studies (e.g., Suthof, GBC, Vol 15., no.3, 2001.) have explained ice core variations in N<sub>2</sub>O with mechanisms driven primarily by changes in OMZ-related production.

p. 16740, Figure 6 caption. Please provide more details on the MLD 5m change criteria. Is hatching drawn when the summertime mixed layer depth, the annual mean depth or some other time average changes by 5m?

P16742 Figure 8. Please explain in the figure caption what the x's are.

---

Interactive comment on Biogeosciences Discuss., 11, 16703, 2014.

C8028