

1 **Soil carbon sequestration by three perennial legume pastures is greater in deeper**  
2 **soil layers than in the surface soil**

3 Xiao-Kang Guan<sup>A</sup>, Neil C. Turner<sup>BC</sup>, Lei Song<sup>D</sup>, Yan-Jie Gu<sup>B</sup>, Tong-Chao Wang<sup>A\*</sup>,  
4 Feng-Min Li<sup>BC\*</sup>

5 <sup>A</sup> Collaborative Innovation Center of Henan Grain Crops, Agronomy College of  
6 Henan Agricultural University, Zhengzhou, Henan 450002, China

7 <sup>B</sup> State Key Laboratory of Grassland Agro-Ecosystem, Institute of Arid Agroecology,  
8 School of Life Sciences, Lanzhou University, Lanzhou 730000, China

9 <sup>C</sup> The UWA Institute of Agriculture and Centre for Plant Genetics and Breeding, The  
10 University of Western Australia, M080, Perth, WA 6009, Australia

11 <sup>D</sup> Medical College of Northwest University for Nationalities, Lanzhou 730030, China

12 \* Corresponding author.

13

14 No. of table: 2

15 No. of figures: 5

16

17 Corresponding authors: Tong-Chao Wang (Prof.), Feng-Min Li (Prof.)

18 Tel/Fax: +86 371 63978418, +86 931 8912848

19 E-mail: [wtcwrn@126.com](mailto:wtcwrn@126.com), [fmli@lzu.edu.cn](mailto:fmli@lzu.edu.cn)

20

## 21 **ABSTRACT**

22 Soil organic carbon (SOC) plays a vital role as both a sink for and source of  
23 atmospheric carbon. Revegetation of degraded arable land in China is expected to  
24 increase soil carbon sequestration, but the role of perennial legumes on soil carbon  
25 stocks in semiarid areas has not been quantified. In this study, we assessed the effect  
26 of alfalfa (*Medicago sativa* L.) and two locally-adapted forage legumes, bush clover  
27 (*Lespedeza davurica* S.) and milk vetch (*Astragalus adsurgens* Pall.) on the SOC  
28 concentration and SOC stock accumulated annually over a 2-m soil profile. The  
29 results showed that the concentration of SOC in the bare soil decreased slightly over  
30 the 7 years, while 7 years of legume growth substantially increased the concentration  
31 of SOC over the 0-2.0 m soil depth. Over the 7-year growth period the SOC stocks  
32 increased by 24.1 Mg C ha<sup>-1</sup>, 19.9 Mg C ha<sup>-1</sup> and 14.6 Mg C ha<sup>-1</sup> under the alfalfa,  
33 bush clover and milk vetch stands, respectively, and decreased by 4.2 Mg C ha<sup>-1</sup> in the  
34 bare soil. The sequestration of SOC in the 1-2 m depth of the soil accounted for 79, 68  
35 and 74 % of the SOC sequestered in the 2 m deep soil profile under alfalfa, bush  
36 clover and milk vetch, respectively. Conversion of arable land to perennial legume  
37 pasture resulted in a significant increase in SOC, particularly at soil depths below 1 m.

38 **Keywords:** Soil organic carbon, SOC stocks, SOC sequestration, perennial legumes  
39 pasture.

## 40 **1. Introduction**

41 Concerns about global warming and increasing atmospheric greenhouse gas  
42 concentrations (CO<sub>2</sub>, CH<sub>4</sub>, and N<sub>2</sub>O) have led to questions on the role of soils as a  
43 source or sink for carbon. Excluding carbonated rocks, soils constitute the largest  
44 surface carbon pool, approximately 1500 Gt, equivalent to almost twice that in the  
45 terrestrial biomass and three times that in the atmosphere (IPCC, 2000). Globally,  
46 soil cultivation has resulted in the loss of more than 40 Pg C, at a rate of about 1.6 Pg  
47 C year<sup>-1</sup>, to the atmosphere during the 1990s (Smith, 2008). Chinese agricultural soils  
48 have also lost 30-50% or more of the soil carbon pool (Lal, 2004a).

49 Soil organic carbon (SOC) is a significant component of the global carbon stocks  
50 (Chen et al., 2008). Globally, 24% of the SOC stock has been lost through the  
51 conversion of forest to cropland (He et al., 2011) and 59% through the conversion of  
52 pasture to cropland (Zhu, 1994). Fortunately, the loss of SOC can be slowed down by  
53 implementing crop management practices such as conservation tillage (Lal, 2004b;  
54 Puget and Lal, 2005), converting degraded arable land to perennial grassland (Gentile  
55 et al., 2005), using diverse rotations, and introducing legume and grass mixtures into  
56 the rotation (Lal, 2002, 2004b, c).

57 In the USA, the revegetation of highly-erodible cropland or other environmentally-  
58 sensitive areas to resource-conserving vegetation for a period of 10 to 15 years  
59 increased the SOC content in the upper 3 m of soil at average rate of 1.1 Mg C ha<sup>-1</sup>  
60 year<sup>-1</sup> (Osborn, 1993). This conservation reserve program (CRP) also significantly  
61 increased the soil C pool (Staben et al., 1997) and provided multiple benefits both  
62 environmentally and economically (Munson et al., 2012; Wu and Lin, 2010). Like the

63 CRP program in USA, a program of soil and water conservation, namely “Grain for  
64 Green” was implemented on the Loess Plateau of China in 1999 to alleviate land  
65 degradation. The program of eco-environmental revegetation focused on the recovery  
66 of damaged ecosystems (Wang et al., 2010) by the use of perennial vegetation to  
67 control soil erosion, increase the stocks of SOC and prevent the occurrence of dry  
68 layers in the loess soils (Fu et al., 2010). Alfalfa (*Medicago sativa* L.) has been widely  
69 grown on the Loess Plateau to increase livestock production and improve water-use  
70 efficiency and soil fertility through high forage production, and for its ability to  
71 decrease soil erosion and fix atmospheric N (Guan et al., 2013). Additionally,  
72 locally-adapted legume species such as bush clover (*Lespedeza davurica* S.) and milk  
73 vetch (*Astragalus adsurgens* Pall.) have been widely grown as cover crops or  
74 windbreaks to protect the soil from water and wind erosion in arid and semiarid  
75 regions of northern China (Wang, 2003; Xu et al., 2006). The “Grain for Green”  
76 program has reduced wind and water erosion of marginal arable land and is expected  
77 to significantly contribute to soil C sequestration. Recent studies on the Loess Plateau  
78 have investigated and estimated the changes in SOC stocks in the top 1 m of soil as a  
79 result of revegetation of regional watersheds (Fu et al., 2010; Wang et al., 2011; Yan  
80 et al., 2007; Zhang et al., 2013). However, deep-rooted perennial legumes may  
81 penetrate deeper in the soil profile than 1 m, likely underestimating the SOC-  
82 sequestration potential of these forage legumes in northwest China (Smith et al.,  
83 2015).

84 The objective of this research was to assess the effect of alfalfa and two locally-  
85 adapted forage legumes, bush clover and milk vetch, on the SOC concentration and  
86 SOC stock accumulated annually for 7 years over a 2-m soil profile. The SOC content  
87 in the 2-m soil profile was measured at the end of each growing season to quantify the  
88 SOC concentration and stock under the locally-adapted forage legumes and alfalfa,  
89 and to provide specific information for estimating the SOC sequestration potential of  
90 an important agricultural area. The hypothesis tested was that long-term growth of  
91 deep-rooted perennial legumes will increase soil organic C, provide a feed source for  
92 animals and provide a sink for atmospheric carbon.

## 93 **2. Materials and methods**

### 94 **2.1 Experimental site description**

95 The experiment was conducted at the Changwu Agro-ecological Experiment Station  
96 on the Loess Plateau (35°12'N, 107°40'E), Shaanxi province, China, from 2004 to  
97 2010. The level site is located at 1220 m above sea level. The climate is semiarid,  
98 with an annual mean temperature of 9.1 °C and a mean annual precipitation of 579  
99 mm (1979-2003) with rainfall concentrated in the period from July to September.  
100 Precipitation and temperature data were recorded at the Changwu Meteorological  
101 Station, 20 m from the experimental site. The groundwater table is 50 – 80 m below  
102 the soil surface, making it unavailable for plant growth. Prior to the establishment of  
103 this experiment, the site was planted to winter wheat for many (at least 20) years. For  
104 winter wheat production, the site was ploughed to a depth of 0.3 m twice a year, after

105 harvest in early July and again in September before sowing; only wheat stubble was  
106 returned to the soil, but 108 kg N ha<sup>-1</sup> and 276 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> of fertilizer was applied  
107 each year before sowing. In 2003, after the winter wheat was harvested, the site  
108 remained fallow for 280 days to allow moisture accumulation over the winter before  
109 the legumes were sown in May 2004.

110 Soil at the experimental site belongs to the Loess series. The texture in the top 5 m is  
111 a uniform silty clay loam (haplic greyxems, FAO-UNESCO, 1988), with a mean sand,  
112 silt, and clay content of 3.5, 65.6, and 30.9%, respectively. The soil physical  
113 characteristics do not significantly change in the upper 5 m. The measured average  
114 bulk density of the soil in the upper 2 m is 1.31 g cm<sup>-3</sup>, does not change with depth,  
115 and the top 0.3 m contained 1.55% total organic matter, 0.106% nitrogen, and 0.095%  
116 available phosphate prior to the commencement of the experiment in 2004.

## 117 **2.2 Treatments and measurement of aboveground biomass**

118 Twelve experiment plots, each 4 m by 3 m, were established in early May 2004 with  
119 one of three forage legume species, milk vetch (*Astragalus adsurgens* Pall.), alfalfa  
120 (*Medicago sativa* L.) and bush clover (*Lespedeza davurica* S.), and an unplanted  
121 control. Each legume species was grown as an evenly-spaced monoculture (but  
122 assumed a more patchy distribution with time) at a seeding density of 25 plants m<sup>-2</sup>,  
123 weeds were removed from all plots by hand. The plots were adjacent to each other.  
124 During the experimental period from 2004 to 2010, there was no irrigation or other  
125 form of supplementary water and the plots were not fertilized. The plants were not

126 inoculated, but relied on the naturally-occurring root nodule bacteria from previous  
127 growth of the three species on the experimental station. **Lack of** nodulation was not  
128 observed to be a problem. Treatments were completely randomized in three replicate  
129 blocks.

130 Each year from 2005-2010, measurements of aboveground biomass production for  
131 each legume were taken at the end May, July and September (in 2004 only one cut  
132 was made in September) by cutting the plants at ground level with hand-held shears in  
133 a randomly-selected 1 m × 1 m quadrat within each plot, but avoiding border areas. At  
134 the same time, the rest of the plot was also cut at the same height and the forage  
135 removed. The oven-dry weight was determined after drying at 105 °C for 0.5 h and  
136 then further dried at 75 °C for 48 h (Guan et al., 2013).

### 137 **2.3 Soil sampling and analysis**

138 Soil samples were taken with a cylindrical steel corer (diameter 40 mm and height  
139 200 mm) at two random positions in each plot which were combined into one  
140 composite sample per plot before analysis. Each plot was sampled from the surface to  
141 2 m deep at depths of 0-0.3, 0.3-0.6, 0.6-1.0, 1.0-1.5 and 1.5-2.0 m before sowing on  
142 10 May 2004 and at the end of each growing season (29 October) from 2004 to 2010.  
143 The soil samples were air-dried, roots and organic debris removed, ground and sieved  
144 through a 2 mm sieve, then stored at room temperature before analyzing the SOC.

145 The concentration of SOC (in g kg<sup>-1</sup>) was measured using the wet dichromate  
146 oxidation procedure (Moinuddin and Khanna-Chopra, 2004). Briefly, a 0.5 g soil

147 sample was digested with 5 mL of 1N K<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub> and 5 mL of concentrated H<sub>2</sub>SO<sub>4</sub> at  
148 150°C for 0.5 h, followed by titration of the digest with standardized FeSO<sub>4</sub>.

## 149 **2.4 SOC stock calculation and statistical analyses**

150 Soil organic C stock was calculated as Eq (1):

$$151 \quad C_{\text{stock}} = SOC \times \rho \times H \times 10 \quad (1)$$

152 where SOC is the SOC concentration (g kg<sup>-1</sup>) in each soil layer, ρ is the bulk density  
153 (g cm<sup>-3</sup>) of the soil, and H is the depth of each layer.

154 The data were analyzed by analysis of variance (ANOVA), and means were compared  
155 using the LSD at  $P < 0.05$  to characterize the differences among treatments. PROC  
156 GLM (General Linear Model) in the Statistical Analysis System (SAS Institute, Cary,  
157 NC, version 8.02) software was used to assess the temporal changes in SOC stock and  
158 the rate and amount of SOC sequestered.

## 159 **3. Results**

### 160 **3.1 Meteorological conditions**

161 The average monthly mean temperature from June to August, the primary growth  
162 period for the legumes, was about 20 °C. Monthly mean temperatures were about 1 °C  
163 warmer than the long-term mean throughout the experimental period (Fig. 1). Over  
164 the experimental period, the total annual precipitation varied from 470 mm in 2006 to  
165 583 mm in 2010, and was below the long-term mean in all years except 2010 when  
166 the rainfall was similar to the long-term mean (Fig. 1). Rainfall from July to



167 September accounted for 55-60% of total annual precipitation, while rainfall in the  
168 legume-growing season (from April to October) was about 90% (range from 84% in  
169 2009 to 96% in 2005) of the total annual precipitation (Fig. 1).

### 170 **3.2 Aboveground forage biomass production**

171 The results of the aboveground biomass production over the seven years have been  
172 reported by Guan et al. (2013). Briefly, the annual production of milk vetch increased  
173 from 2.2 t ha<sup>-1</sup> in the first year to 14.3 t ha<sup>-1</sup> in 2006 and then decreased, alfalfa  
174 increased from 2.3 t ha<sup>-1</sup> in the first year to a maximum of 22.2 t ha<sup>-1</sup> in 2006 and then  
175 decreased, while bush clover increased from 0.2 t ha<sup>-1</sup> in the first year to 7.8 t ha<sup>-1</sup> in  
176 2009 and did not decrease significantly thereafter (Table 1). Total aboveground  
177 biomass production over the experimental period was highest in alfalfa at 91 t ha<sup>-1</sup>  
178 (equivalent to 43 Mg C ha<sup>-1</sup> assuming a C to dry weight ratio of 0.475, Magnussen  
179 and Reed, 2004) compared to 56 t ha<sup>-1</sup> (27 Mg C ha<sup>-1</sup>) in milk vetch and 42 t ha<sup>-1</sup> (20  
180 Mg C ha<sup>-1</sup>) in bush clover (Table 1).

### 181 **3.3 SOC concentration over the soil profile**

182 The legumes significantly ( $P < 0.001$ ) increased the SOC concentration at each soil  
183 depth, and this effect varied with legume species and experimental year (Table 2).  
184 The initial concentration of SOC in May 2004 decreased with increasing soil depth  
185 (Fig. 2). In the upper 0-0.3 m of soil, the initial SOC concentration was  $8.0 \pm 0.03$  g kg<sup>-1</sup>  
186 <sup>1</sup>, while it was only  $3.3 \pm 0.27$  g kg<sup>-1</sup> in the 1.5-2.0 m soil layer (Fig. 2). Comparison of  
187 the SOC concentration between the initial values on 10 May 2004 and those at the end

188 of the experimental period in October 2010 showed that the concentration of SOC in  
189 the bare soil decreased slightly over the 7 years, while 7 years of legume growth  
190 substantially increased the concentration of SOC over whole 2 m soil depth. There  
191 were large increases in the concentration of SOC at 0.6-1.0 m, 1.0-1.5 m and 1.5-2.0  
192 m soil depth and a small, but significant, increase in the upper 0.3 m of the soil in  
193 bush clover, but not in milk vetch and alfalfa. No significant changes were observed  
194 after 7 years at the 0.3-0.6 m depth (Fig. 2).

### 195 **3.4 SOC stock over the experimental period**

196 SOC stock was calculated by converting SOC concentration to the amount of SOC  
197 per soil layer per unit area. The SOC stock in 2004 varied from  $21.4 \pm 0.85 \text{ Mg C ha}^{-1}$   
198 in the 0.3-0.6 m soil layer to  $32 \pm 0.14 \text{ Mg C ha}^{-1}$  and  $32 \pm 0.68 \text{ Mg C ha}^{-1}$  at 0-0.3 and  
199 1.0-1.5 m depth respectively (Fig.3). In the bare soil, the SOC stock decreased at all  
200 depths across the experimental period, but only decreased significantly at  $-0.36 \text{ Mg C}$   
201  $\text{ha}^{-1} \text{ year}^{-1}$  ( $P < 0.05$ ) in the 1.5-2.0 m layer (Fig. 3d), presumably from the decay and  
202 turnover of the wheat roots accumulated over the many years of wheat production  
203 prior to the planting of the legumes. In the legume plots, the SOC stock increased  
204 linearly with time (2004-2010) in the 0-0.3 m, 0.6-1.0 m, 1.0-1.5 m and 1.5-2.0 m soil  
205 layers, but not in the 0.3-0.6 m soil layer (Fig. 3). The change in SOC stock over the 7  
206 years was greatest at soil depths below 1.0 m in all three species and was greatest in  
207 the alfalfa plots with rates of  $1.35 \text{ Mg C ha}^{-1} \text{ year}^{-1}$  at a depth of 1.0-1.5 m ( $P < 0.001$ )  
208 (Fig. 3d), and  $1.39 \text{ Mg C ha}^{-1} \text{ year}^{-1}$  at a depth of 1.5-2.0 m ( $P < 0.001$ ) (Fig. 3e). The

209 highest accumulation of SOC stock occurred at a depth of 1.0-1.5 m in bush clover  
210 where it averaged  $1.58 \text{ Mg C ha}^{-1} \text{ year}^{-1}$  ( $P < 0.001$ ) (Fig. 3d).

211 Over the full 0-2.0 m depth, the SOC stock at the beginning of the experiment in 2004  
212 was  $137 \text{ Mg C ha}^{-1}$ , decreased to  $133 \text{ Mg C ha}^{-1}$  in the bare soil plots, while it  
213 increased to 151, 157 and  $161 \text{ Mg C ha}^{-1}$  under the milk vetch, bush clover and alfalfa  
214 stands, respectively, by the end of the experiment in 2010 (Fig. 4). The SOC stock  
215 increased more under the stand of alfalfa than milk vetch, but there was no significant  
216 difference between alfalfa and bush clover (Fig. 4). When calculated over the full 2-m  
217 soil layer, over the 7-year growth period, the SOC stocks increased by  $24.1 \text{ Mg C ha}^{-1}$ ,  
218  $19.9 \text{ Mg C ha}^{-1}$  and  $14.6 \text{ Mg C ha}^{-1}$  under the alfalfa, bush clover and milk vetch  
219 stands, respectively, and decreased by  $4.2 \text{ Mg C ha}^{-1}$  under bare soil (Fig. 5). In the  
220 1.0-2.0-m soil layer the stocks of SOC increased by  $19.1 \text{ Mg C ha}^{-1}$ ,  $13.6 \text{ Mg C ha}^{-1}$   
221 and  $10.8 \text{ Mg C ha}^{-1}$ , under the alfalfa, bush clover and milk vetch stands, respectively,  
222 that is, by 79, 68 and 74% of the increases over the whole soil profile (Fig. 5).

#### 223 4. Discussion

224 While the unplanted plots lost SOC over the 7 years of the study, particularly over the  
225 first 5 years, all three forage legumes more-than countered this loss, so that over the 7  
226 years the SOC increased on average by  $2.1 \text{ Mg C ha}^{-1} \text{ y}^{-1}$ ,  $2.8 \text{ Mg C ha}^{-1} \text{ y}^{-1}$  and  $3.4$   
227  $\text{Mg C ha}^{-1} \text{ y}^{-1}$ , to sequester  $15 \text{ Mg C ha}^{-1}$ ,  $20 \text{ Mg C ha}^{-1}$  and  $24 \text{ Mg C ha}^{-1}$  under the  
228 milk vetch, bush clover and alfalfa stands, respectively (Fig. 5). The legumes  
229 increased the SOC primarily at depth (Fig. 2) with between 68 and 79% of the SOC

230 being sequestered below 1 m. While the three perennial forage legumes have been  
231 shown to cause a significant decrease in soil water at depth in a semiarid environment  
232 (Guan et al., 2013), presumably as a consequence of root activity, the high proportion  
233 of the sequestration of SOC at depths below 1 m observed in this study was  
234 unexpected.

235 An increase in SOC stock is usually associated with the production, turnover,  
236 sloughing off of epidermal cells, and exudation of soluble carbon compounds by the  
237 roots, particularly the turnover and exudation by fine roots (Luo et al., 1995). A high  
238 rate of turnover of fine roots and a high rate of exudation of carbon by the roots  
239 influences the stability of plant C in soil and the accumulation of SOC (Shahzad et al.,  
240 2015). Although the root biomass was not measured in this study, root biomass is  
241 usually associated with aboveground biomass. Assuming an average shoot:root ratio  
242 of 1.3:1 in the three species (Chen and Nie, 1978; Cheng et al., 2004; Fan et al., 2015;  
243 Sun et al., 2001) over the 7-year life of the pastures, the 91 t ha<sup>-1</sup> of aboveground  
244 biomass (43 Mg C ha<sup>-1</sup>) produced would result in the production of 33 Mg C ha<sup>-1</sup> by  
245 the roots in alfalfa, significantly higher than the measured increase in SOC stock in  
246 the upper 2 m of the soil (24 Mg C ha<sup>-1</sup>). The estimated production of SOC by the  
247 roots of milk vetch from the aboveground biomass was also higher than that measured  
248 under milk vetch (estimated 20 Mg C ha<sup>-1</sup>, measured 15 Mg C ha<sup>-1</sup>), but the estimated  
249 SOC was lower than the measured SOC in bush clover (estimated 15 Mg C ha<sup>-1</sup>,  
250 measured 20 Mg C ha<sup>-1</sup>) over the 7-year period. Thus the actual accumulation of SOC  
251 was about 70% of that estimated from the aboveground biomass in alfalfa and milk

252 vetch, but was 33% higher than that estimated in bush clover. The differences in SOC  
253 between the estimated and measured values in alfalfa and milk vetch were presumably  
254 the result of losses by respiration by the roots and associated soil microbial  
255 populations. The observation that the measured values of SOC were higher than those  
256 estimated from aboveground biomass may indicate that the shoot:root ratio was less  
257 than 1.3:1 in bush clover (a shoot:root ratio of 1:1 would make the estimated and  
258 measured values the same), or the bush clover had a greater proportion and turnover  
259 of fine roots than milk vetch and alfalfa, resulting in a greater accumulation of SOC  
260 (Shahzad et al., 2015). Sun et al. (2001) reported that the fine roots (root diameter <  
261 0.5 mm) of bush clover accounted for 42% of total root biomass in 0-0.3 m soil layer,  
262 while the fine roots of milk vetch were only 25% of total root biomass (Chen and Nie,  
263 1978).

264 The accumulation of SOC by bush clover was particularly high in the upper 0.3 m of  
265 the soil (Fig. 2, Fig. 3). This accumulation of SOC in the upper soil layer may be  
266 attributable to the high accumulation of legume residues and litter (Zhou et al., 2006),  
267 or due to the proliferation and turnover of roots in this surface layer. Nevertheless, the  
268 sequestration of SOC in the upper 0.3 m of the soil in this study was significantly  
269 lower than Zhang et al. (2009) who reported that the SOC stocks in upper 0.3 m of the  
270 soil increased by 16 Mg C ha<sup>-1</sup> in 10 years from the conversion of a wet reed meadow  
271 to an irrigated alfalfa pasture in the Hexi Corridor of north-west China. This suggests  
272 that well-managed legume pastures in areas with higher precipitation and with

273 appropriate fertilizer use could sequester significantly more SOC than in the present  
274 unirrigated and unfertilized legumes growing in a semiarid environment.

275 An unexpected result from this study was the greater increase in SOC at soil depths  
276 from 1 – 2 m than above 1 m, accounting for 79, 68 and 74% of SOC sequestered  
277 through the whole 2 m of soil under alfalfa, bush clover and milk vetch, respectively.  
278 This suggests the presence of roots deep in the soil profile. Indeed, alfalfa and milk  
279 vetch have taproots that can penetrate to 6.8 and 7.6 m, respectively, in loess soils in 6  
280 years (Cheng et al., 2005; Cheng et al., 2004), and the water extraction patterns  
281 measured on the plots used in this study clearly showed the presence of roots  
282 throughout the upper 2 m of the soil profile (Guan et al., 2013). In bush clover in  
283 which the proportion of SOC below 1 m was smaller than in the other two species, the  
284 taproot predominates in the 0-0.3 m soil layer with coarse roots (root diameter > 2  
285 mm) accounting for 48% of the total root biomass, and fine roots predominating  
286 below 0.3 m (Cheng et al., 2005; Cheng et al., 2004; Sun et al., 2001). Although  
287 alfalfa extracted more water below 1 m than the other two species, the water  
288 extraction patterns do not suggest a greater root presence below 1 m in milk vetch  
289 than bush clover (Guan et al., 2013). We suggest that the greater increase in SOC at  
290 depth may be associated with a greater proliferation and turnover of fine roots at  
291 depth, or alternatively may reflect the movement down the profile of soluble C  
292 compounds from the roots as a result of the movement of water after heavy rainfall  
293 events.

294 The conversion of arable land that had been growing crops for many years to  
295 perennial legume pasture resulted in a significant increase in SOC, particularly at soil  
296 depths below 1 m. All three legume species increased the SOC in the top 2 m of the  
297 soil profile, but the increase was greatest in alfalfa and least in milk vetch. While the  
298 production of aboveground biomass was least in bush clover, the SOC sequestration  
299 in the soil profile was not significantly different from alfalfa, indicating that carbon  
300 sequestration in the soil is not **simply** associated with aboveground biomass  
301 production in a system in which the forage is removed for animal feed, as in the  
302 present study. **Further study of** root biomass production, turnover of fine roots and  
303 exudation of carboxylic acids and other carbon compounds by the roots **of** the legume  
304 species would be a valuable step in understanding the differences in carbon  
305 sequestration **by** the three species.

### 306 **Acknowledgements**

307 This research was supported by grants (No. 31470496 and No. 31471452) from the  
308 National Natural Science Foundation of China, the National Key Technology R&D  
309 Program of China (Projects 2012BAD04B07 and 2012BAD14B08) and the  
310 Innovation Team Program of the Ministry of Education of China (No. IRT\_13R26).  
311 This research was also supported by the Fundamental Research Funds for the Central  
312 Universities (lzujbky-2015-ct02), and the Program of Introducing Talents of  
313 Discipline to Universities ('111 Project 2007B051'). We are grateful to Xiao-Hong  
314 Zhang for her help in the laboratory. NCT thanks the Centre for Plant Genetics and  
315 Breeding and the UWA Institute of Agriculture at the University of Western Australia

316 for financial support. This work is a part of the doctoral dissertation of the first author,

317 Dr Xiao-Kang Guan, at Lanzhou University.

318



319 **References**

- 320 Chen, B. and Nie, C.: Research of *Astragalus adsurgens* Pall. root system, Journal of Gansu  
321 Agricultural University, 2, 71-75, 1978. (in Chinese with English abstract)
- 322 Chen, H., Shao, M., and Li, Y.: Soil desiccation in the Loess Plateau of China, Geoderma,  
323 143, 91–100, 2008.
- 324 Cheng, J., Wan, H. E., and Wang, J.: Alfalfa growth and its relation with soil water status in  
325 loess hilly and gully region, The Journal of Applied Ecology, 16, 435-438, 2005. (in Chinese  
326 with English abstract)
- 327 Cheng, J. M., Wan, H. E., Wang, J., and Yong, S. P.: Over depletion and recovery of soil  
328 moisture on *Astragalus adsurgens* grasslands in the loess hilly-gully region, Acta Ecologica  
329 Sinica, 24, 2979-2983, 2004. (in Chinese with English abstract)
- 330 Fan, J. W., Du, Y. L., Turner, N. C., Wang, B. R., Fang, Y., Xi, Y., Guo, X. R., and Li, F. M.:  
331 Changes in root morphology and physiology to limited phosphorus and moisture in a locally-  
332 selected cultivar and an introduced cultivar of *Medicago sativa* L. growing in alkaline soil,  
333 Plant & Soil, 392, 215-226, 2015.
- 334 Fu, X., Shao, M., Wei, X., and Horton, R.: Soil organic carbon and total nitrogen as affected  
335 by vegetation types in Northern Loess Plateau of China, Geoderma, 155, 31-35, 2010.
- 336 Gentile, R. M., Martino, D. L., and Entz, M. H.: Influence of perennial forages on subsoil  
337 organic carbon in a long-term rotation study in Uruguay, Agriculture, Ecosystems &  
338 Environment, 105, 419-423, 2005.
- 339 Guan, X. K., Zhang, X. H., Turner, N. C., Xu, B. C., and Li, F. M.: Two perennial legumes  
340 (*Astragalus adsurgens* Pall. and *Lespedeza davurica* S.) adapted to semiarid environments are  
341 not as productive as lucerne (*Medicago sativa* L.), but use less water, Grass and Forage  
342 Science, 68, 469-478, 2013.
- 343 He, Z., Huang, C., Zheng, H., Zhou, J., Pang, J., Li, X., and Wang, L.: Holocene loess and its  
344 deposition dynamics in the upper reaches of the Huaihe River, J. Geogr. Sci., 21, 561-573,  
345 2011.
- 346 IPCC: Land-use, land-use change, and forestry. In: Land use, land-use change, and forestry: a  
347 special report of the intergovernmental panel on climate change, Watson, R. T., Noble, I. R.,  
348 Bolin, B. R., Ravindranath, N. H., Verardo, D. J., and Dokken, D. J. (Eds.), Cambridge  
349 University Press, U.K., 2000.
- 350 Lal, R.: Offsetting China's CO<sub>2</sub> Emissions by Soil Carbon Sequestration, Climatic Change,  
351 65, 263–275, 2004a.
- 352 Lal, R.: Soil carbon dynamics in cropland and rangeland, Environmental Pollution, 116, 353-  
353 362, 2002.
- 354 Lal, R.: Soil carbon sequestration impacts on global climate change and food security,  
355 Science, 304, 1623-1627, 2004b.
- 356 Lal, R.: Soil carbon sequestration to mitigate climate change, Geoderma, 123, 1-22, 2004c.
- 357 Luo, Y., Meyerhoff, P. A., and Loomis, R. S.: Seasonal patterns and vertical distributions of  
358 fine roots of alfalfa (*Medicago sativa* L.), Field Crops Research, 40, 119-127, 1995.
- 359 Mangnussen, S. and Reed, D.: Modeling for Estimation and Monitoring, Available at  
360 <http://www.fao.org/forestry/17109/en/>, 2004.

361 Moinuddin and Khanna-Chopra, R.: Osmotic Adjustment in Chickpea in Relation to Seed  
362 Yield and Yield Parameters, *Crop Science*, 44, 449-455, 2004.

363 Munson, S., Lauenroth, W., and Burke, I.: Soil carbon and nitrogen recovery on semiarid  
364 Conservation Reserve Program lands, *Journal of Arid Environments*, 79, 25-31, 2012.

365 Osborn, T.: The conservation reserve program: Status, future and policy options, *Journal of*  
366 *Soil and Water Conservation*, 48, 271-279, 1993.

367 Puget, P. and Lal, R.: Soil organic carbon and nitrogen in a Mollisol in central Ohio as  
368 affected by tillage and land use, *Soil and Tillage Research*, 80, 201-213, 2005.

369 Shahzad, T., Chenu, C., Genet, P., Barot, S., Perveen, N., Mougin, C., and Fontaine, S.:  
370 Contribution of exudates, arbuscular mycorrhizal fungi and litter depositions to the  
371 rhizosphere priming effect induced by grassland species, *Soil Biology and Biochemistry*, 80,  
372 146-155, 2015.

373 Smith, P.: Land use change and soil organic carbon dynamics, *Nutrient Cycling in*  
374 *Agroecosystems*, 81, 169-178, 2008.

375 Smith, P., House, J. I., Bustamante, M., Sobocká J., Harper, R., Pan, G., West, P., Clark, J.,  
376 Adhya, T., Rumpel, C., Paustian, K., Kuikman, P., Cotrufo, M. F., Elliott, J. A., McDowell,  
377 R., Griffiths, R. I., Asakawa, S., Bondeau, A., Jain, A. K., Meersmans, J., and Pugh, T. A. M.:  
378 Global Change Pressures on Soils from Land Use and Management, *Global Change Biology*,  
379 doi: 10.1111/gcb.13068, 2015. n/a-n/a, 2015.

380 Staben, M., Bezdicek, D., Fauci, M., and Smith, J.: Assessment of soil quality in conservation  
381 reserve program and wheat-fallow soils, *Soil Science Society of America Journal*, 61, 124-  
382 130, 1997.

383 Sun, Q., Han, J., Gui, R., Li, Z., and Liu, G.: Biomass in *Lespedeza dahurica* S., Grassland of  
384 China, 04, 21-26, 2001. (in Chinese with English abstract)

385 Wang, G.: Differences in leaf  $\delta^{13}\text{C}$  among four dominant species in a secondary succession  
386 sere on the Loess Plateau of China, *Photosynthetica*, 41, 525-531, 2003.

387 Wang, Y., Fu, B., Lü, Y., and Chen, L.: Effects of vegetation restoration on soil organic  
388 carbon sequestration at multiple scales in semi-arid Loess Plateau, China, *Catena*, 85, 58-66,  
389 2011.

390 Wang, Y., Shao, M. a., and Shao, H.: A preliminary investigation of the dynamic  
391 characteristics of dried soil layers on the Loess Plateau of China, *Journal of Hydrology*, 381,  
392 9-17, 2010.

393 Wu, J. and Lin, H.: The effect of the conservation reserve program on land values, *Land*  
394 *Economics*, 86, 1-21, 2010.

395 Xu, B., Gichuki, P., Shan, L., and Li, F.: Aboveground biomass production and soil water  
396 dynamics of four leguminous forages in semiarid region, northwest China, *South African*  
397 *Journal of Botany*, 72, 507-516, 2006.

398 Yan, H., Cao, M., Liu, J., and Tao, B.: Potential and sustainability for carbon sequestration  
399 with improved soil management in agricultural soils of China, *Agriculture, Ecosystems &*  
400 *Environment*, 121, 325-335, 2007.

401 Zhang, C., Liu, G., Xue, S., and Sun, C.: Soil organic carbon and total nitrogen storage as  
402 affected by land use in a small watershed of the Loess Plateau, China, *European Journal of*  
403 *Soil Biology*, 54, 16-24, 2013.

404 Zhang, T., Wang, Y., Wang, X., Wang, Q., and Han, J.: Organic carbon and nitrogen stocks  
405 in reed meadow soils converted to alfalfa fields, *Soil and Tillage Research*, 105, 143-148,  
406 2009.  
407 Zhou, Z., Sun, O. J., Huang, J., Li, L., Liu, P., and Han, X.: Soil carbon and nitrogen stores  
408 and storage potential as affected by land-use in an agro-pastoral ecotone of northern China,  
409 *Biogeochemistry*, 82, 127-138, 2006.  
410 Zhu, X.: The formation and significance of loess-soil struction profile, *Journal of Soil and*  
411 *Water Conservation*, 8, 1-9, 1994. (in Chinese with English abstract)

412

413 **Figure Captions**


414 **Fig. 1.** Mean monthly temperature and precipitation from 2004-2010 and the long  
415 term mean at the experimental site at Changwu Agricultural Research Station,  
416 Shaanxi Province, China.

417 **Fig. 2.** Concentration of soil organic carbon (SOC) with depth in May 2004 (IV) and  
418 in October 2010 under three forage legumes: milk vetch, alfalfa and bush clover, and  
419 bare soil (CK). Bars give + one standard error of the mean ( $n = 3$ ).

420 **Fig. 3.** Change with stand age in soil organic carbon amount (stock) per hectare at soil  
421 depths of 0-0.3m (a), 0.3-0.6m (b), 0.6-1.0m (c), 1.0-1.5m (d) and 1.5-2.0m (e) under  
422 milk vetch, alfalfa, bush clover and bare soil (CK). Note the soil layers vary in depth.  
423 Data are means  $\pm$  one standard error of the mean ( $n = 3$ ) when larger than the symbol.  
424 Linear regressions fitted when significant and fitted regressions given.

425 **Fig. 4.** The soil organic carbon amount (stock) under milk vetch, alfalfa, bush clover  
426 and under bare soil (CK) over the upper 2 m of the soil profile. The lower case letters  
427 indicate significant differences ( $P < 0.05$ ) between forage types and bare soil (CK)  
428 within a year. IV denotes initial value, the soil organic carbon stock in May 2004.  
429 Bars give + one standard error of the mean ( $n = 3$ ).


430 **Fig. 5.** Change in soil organic carbon amount (stock) in different soil layers under  
431 milk vetch, alfalfa, bush clover and bare soil (CK) from May 2004 to October 2010.  
432 Different letters indicate significant differences ( $P < 0.05$ ) between total carbon stocks.



433

434 Figure 1


435



436

437 Figure 2


438



439

440 Figure 3

441




442

443 Figure 4

444





445

446 Figure 5

447 **Table 1. Annual aboveground biomass production of the three legume species,**  
 448 **milk vetch, alfalfa and bush clover, from 2004 to 2010. Adopted from (Guan et al.**  
 449 **(2013)) and used with permission.**

Year	Aboveground biomass production (t ha <sup>-1</sup> )		
	Milk vetch	Alfalfa	Bush clover
2004	2.2Ac	2.3Ad	0.2Bd
2005	14.1Ba	20.2Aa	5.3Cc
2006	14.3Ba	22.2Aa	7.8Ca
2007	6.8Bb	9.3Ac	6.4Bbc
2008	5.6Bb	13.4Ab	7.3Bab
2009	7.2Bb	12.4Ab	7.8Ba
2010	5.8Bb	10.8Aabc	7.4Bab
2004-2010 Mean	8.0B	13.0A	6.0C
2004-2010 Total	56.0B	90.7A	42.1C

450 Data in each column with a different lower-case letter are significantly different ( $P <$   
 451  $0.05$ ) and data in each row with a different capital letter are significantly different ( $P$   
 452  $< 0.05$ ).

453

454 **Table 2. Results of the ANOVA for soil organic carbon concentration as affected**  
 455 **by legume species, soil depth and experimental year. The bare soil plot is**  
 456 **considered as a legume species in the analysis. The SAS PROC ANOVA was**  
 457 **used in the analysis.**

Factors	df	<i>F</i> value	<i>Pr</i> > <i>F</i>
Species	3	38.52	0.0003
Depth	4	1649.40	<0.0001
Year	7	31.68	<0.0001
Species*Depth	12	5.65	<0.0001
Species*Year	21	5.96	<0.0001
Depth*Year	28	3.20	<0.0001
Species*Depth*Year	84	0.95	0.6053

458