

1 **Permafrost coverage, watershed area and season control of dissolved carbon**  
2 **and major elements in western Siberia rivers**

3  
4  
5 **O.S. Pokrovsky<sup>1\*</sup>, R.M. Manasypov<sup>2,3</sup>, S. Loiko<sup>2</sup>, L.S. Shirokova<sup>1,3</sup>, I.A. Krickov<sup>2</sup>,**  
6 **B.G. Pokrovsky<sup>4</sup>, L.G. Kolesnichenko<sup>2</sup>, S.G. Kopysov<sup>2</sup>, V.A. Zemtsov<sup>2</sup>,**  
7 **S.P. Kulizhsky<sup>2</sup>, S.N. Vorobyev<sup>2</sup>, S.N. Kirpotin<sup>2</sup>**

8  
9 <sup>1</sup> GET UMR 5563 CNRS, University of Toulouse, 14 Avenue Edouard Belin, 31400 Toulouse,  
10 France, [oleg@get.obs-mip.fr](mailto:oleg@get.obs-mip.fr)

11 <sup>2</sup> BIO-GEO-CLIM Laboratory, Tomsk State University, Lenina av., 36, Tomsk, Russia

12 <sup>3</sup> Institute of Ecological Problem of the North, 23 Nab Severnoi Dviny, RAS, Arkhangelsk, Russia

13 <sup>4</sup> Geological Institute, Russian Academy of Science, Pyzhevskiy per., 7c1, Moscow, Russia

14  
15  
16 Submitted to *Biogeosciences*, Special issue

17  
18 Keywords: *organic, cations, silica, permafrost, peat, groundwater, soil,*

19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29

## 30 ABSTRACT

31 Analysis of organic and inorganic carbon (DOC and DIC, respectively), pH, Na, K, Ca, Mg,  
32 Cl, SO<sub>4</sub> and Si in ~100 large and small rivers (< 10 km<sup>2</sup> to ≤ 150,000 km<sup>2</sup>) of western Siberia  
33 sampled in winter, spring, and summer over a more than 1500 km latitudinal gradient allowed  
34 establishing main environmental factors controlling the transport of river dissolved components in  
35 this environmentally important region, comprising continuous, discontinuous, sporadic and  
36 permafrost-free zones. There was significant latitudinal trend consisting in general decrease of  
37 DOC, DIC, SO<sub>4</sub>, and major cation (Ca, Mg, Na, K) concentration northward, reflecting the interplay  
38 between groundwater feeding (detectable mostly in the permafrost-free zone, south of 60°N) and  
39 surface flux (in the permafrost-bearing zone). The northward decrease of concentration of inorganic  
40 components was strongly pronounced both in winter and spring, whereas for DOC, the trend of  
41 concentration decrease with latitude was absent in winter, and less pronounced in spring flood than  
42 in summer baseflow. The most significant decrease of K concentration from the southern (< 59°N) to  
43 the northern (61-67°N) watersheds occurs in spring, during intense plant litter leaching. The  
44 latitudinal trends persisted for all river watershed size, from < 100 to > 10,000 km<sup>2</sup>. Environmental  
45 factors are ranked by their increasing effect on DOC, DIC, δ<sup>13</sup>C<sub>DIC</sub>, and major elements in western  
46 Siberian rivers as the following: watershed area < season < latitude. Because the degree of the  
47 groundwater feeding is different between large and small rivers, we hypothesize that, in addition to  
48 groundwater feeding of the river, there was a significant role of surface and shallow subsurface flow  
49 linked to plant litter degradation and peat leaching. We suggest that plant litter- and topsoil derived-  
50 DOC adsorbs on clay mineral horizons in the southern, permafrost-free and discontinuous/sporadic  
51 permafrost zone but lacks the interaction with minerals in the continuous permafrost zone. It can be  
52 anticipated that, under climate warming in western Siberia, the maximal change will occur in small  
53 (< 1000 km<sup>2</sup> watershed) rivers DOC, DIC and ionic composition and this change will be mostly  
54 pronounced in summer.

## 55           **1. Introduction**

56           The Western Siberia Lowland (WSL) can be considered as one of the most vulnerable  
57 permafrost-bearing territories with respect to ongoing climate change, due to (i) the dominance of  
58 discontinuous, sporadic and intermittent permafrost coverage rather than continuous and  
59 discontinuous permafrost of Central and Eastern Siberia and the Canadian High Arctic, (ii) its flat  
60 area and high impact of flooding and thermokarst development and most importantly (iii) its high  
61 stock of ancient and recent organic carbon in the form of partially frozen peat deposits. Due to the  
62 importance of the boreal and sub-arctic continental zones in the Earth's carbon cycle and the high  
63 vulnerability of circumpolar zones to the climate warming, the majority of conducted works has been  
64 devoted to the biogeochemistry of organic carbon and sediments in large rivers of the Russian boreal  
65 circumpolar zone (Gordeev et al., 1996, 2004; Moran and Woods, 1997; Lobbes et al., 2000; Dittmar  
66 and Kattner, 2003; Gebhardt et al., 2004; Lobbes et al., 2000; Cooper et al., 2008; Magritsky, 2010,  
67 Nikanorov et al., 2010a, b; Holmes et al., 2000, 2001, 2012; Pokrovsky et al., 2010; Feng et al.,  
68 2013). While these studies have allowed for the quantification of the carbon and major element  
69 delivery fluxes from the continent to the Arctic ocean, the mechanisms responsible for carbon and  
70 metals mobilization from the soil/groundwater to the rivers remain very poorly understood. The  
71 WSL offers a unique site to test various hypotheses of element sources and to reveal related  
72 mechanisms as it presents the full gradient of the permafrost coverage, climate and vegetation over  
73 homogeneous sedimentary basement rock, essentially peat soil, flat orography and similar annual  
74 precipitation. Taking advantage of these features, in their pioneering studies, Frey et al (2007a, b)  
75 and Frey and Smith (2005) provided a first-order assessment of the relative contributions of shallow  
76 surface water and deep groundwater to small western Siberia rivers. Their study was conducted  
77 during the summer baseflow season presenting the largest contrast between permafrost-free and  
78 permafrost-affected rivers. This allowed them to conclude that climate warming should shift the

79 permafrost-affected part of the region from surface-feeding to groundwater feeding while the  
80 permafrost-free zone may remain unaffected.

81           However, unlike many regions of the world, the boreal and subarctic river regions exhibit  
82 extreme seasonal variations in discharge and chemical elements concentrations (see Voronkov et al.,  
83 1966; Gordeev and Sidorov, 1993; Gordeev et al., 1996; Gislason et al., 1996; Gaillardet et al., 2003;  
84 Rember and Trefry, 2004; Zakharova et al., 2005, 2007; Bagard et al., 2011, 2013; Prokushkin et al.,  
85 2011; Guo et al., 2004b, 2007; Olefeldt and Roulet, 2012; Voss et al., 2015). The quantitative  
86 description of these systems, therefore, requires an understanding of how weathering rates and  
87 riverine fluxes of major and trace elements as well as their main carrier (organic carbon) vary  
88 seasonally. High seasonality implies significant variations in the source of the elements in river flow  
89 over the year, which is further accentuated by high variability of the depth of the active layer and  
90 relevant contributions of mineral soil weathering and the leaching of the soil organic horizon. As  
91 such, the chemistry of fluxes on the seasonal scale depends on the relative role of mineral dissolution  
92 vs. plant litter (organic soil) leaching. Although several recent studies have used isotopic techniques  
93 in an attempt to resolve the sources of elements in subarctic rivers (Engstrom et al., 2010; Keller et  
94 al., 2010; Pokrovsky et al., 2013a; Mavromatis et al., 2014), the relative contributions of mineral and  
95 plant litter/organic soil components remain poorly constrained particularly for boreal watersheds.

96           The purpose of the present work is to improve our understanding of western Siberia river  
97 transport of organic and inorganic carbon and major elements (Ca, Mg, K, Si) via studying numerous  
98 watersheds across the 1500-km latitudinal profile during three main hydrological seasons: winter  
99 baseflow, spring flood and summer-autumn period (Zakharova et al., 2014). As a working  
100 hypothesis, we assume, following the previous works of Frey et al. (2007 a, b) that the permafrost  
101 controls riverine chemical composition via regulating the degree of (i) groundwater feeding and (ii)  
102 leaching of elements from unfrozen (active) soil layers. Because groundwaters in the permafrost  
103 zone are discharged to the river via unfrozen taliks underneath the river bed (Anisimova, 1981;

104 Bagard et al., 2011, 2013), it can be suggested that the impact of groundwaters via taliks will be  
105 mostly visible **in** large rivers, as it is also known from the geocryological studies of the WSL (Fotiev,  
106 1989, 1991). As a result, the contrast in groundwater-related element concentration between rivers of  
107 different latitude is expected to be the largest during winter baseflow. This is especially true in the  
108 WSL, exhibiting highly homogeneous, extremely flat topography and similar lithological cover  
109 (peat, sand and silt). Therefore, the first objective of this study was to test the effect of river size  
110 (watershed area) on inorganic river water components across the permafrost gradient. The second  
111 objective was to assess the effect of the permafrost coverage on DOC, DIC and its isotopic  
112 composition in rivers during different seasons. Specifically, during spring flood when the majority of  
113 soil layer is frozen only surface flux should be important and the concentrations should reflect the  
114 degree of DOC and element leaching from the plant litter. The largest contrast between rivers of  
115 different size is therefore expected in August, whereas the spring flood should exhibit the lowest  
116 differences in terms of DOC transport by rivers of different climate and permafrost zones. Finally,  
117 the third objective of this study was first-order assessment of the major river constituent  
118 concentration across the 2000-km latitudinal profile. Here, we expect, in accordance with general  
119 knowledge of DOC and major cation concentration and export fluxes dependence on temperature,  
120 vegetation and permafrost distribution (White and Blum, 1995; Dessert et al., 2003; Gaillardet et al.,  
121 2003; Millot et al., 2003; Oliva et al., 2003; Smedberg et al., 2006; Frey and McClelland, 2009;  
122 Prokushkin et al., 2011; Beaulieu et al., 2012; Tank et al., 2012a, b; Olefeldt et al., 2014), a gradual  
123 or stepwise decrease of all river water constituents northward, from permafrost-free to discontinuous  
124 and continuous permafrost zone. Verifying the correctness of these research statements should allow  
125 for the quantitative prediction of the degree of river water composition modification in response to  
126 changing environmental conditions, notably the increase in the thickness of active (unfrozen) layer,  
127 increasing the winter discharge and augmenting plant biomass and productivity.

128

129 **2. Study site and Methods**

130 *2.1. Geographical setting*

131 Western Siberia Lowland (WSL) is the world second largest flooding territory, after the  
132 Amazon's Varzea. The rivers (mainly the tributaries of the Ob, Pur, and Taz) drain Pleistocene sands  
133 and clays, covered by thick (1 to 3 m) peat and enclose three main zones of the boreal biome, taiga,  
134 forest-tundra and tundra. **Approximate coverage of studied territory by sand, peat and clay deposits**  
135 **in the first 3 m soil layer is shown in Fig. 1. Note that the peat is always dominant on the watershed**  
136 **divides and bog zones whereas the sand is abundant along the river valleys.** Quaternary clays, sands,  
137 and silts ranging in thickness from several meters to 200-250 m have alluvial, lake-alluvial and,  
138 rarely, aeolian origin south of 60°N and fluvio-glacial and lake-glacial origin north of 60°N. The  
139 older, Paleogene and Neogene, rocks are rarely exposed on the surface and are represented by sands,  
140 alevrolites and clays, where carbonate material is present as concretions of individual shells  
141 (Geologicheskoe Stroenie, 1958). The climate is humid semi-continental with a mean annual  
142 temperature (MAT) ranging from -0.5°C in the south (Tomsk region) to -9.5°C in the north  
143 (Yamburg). The annual precipitation increases from 550 mm at the latitude of Tomsk to 650-700 mm  
144 at Nojabrsk and further decreases to 600 mm at the lower reaches of the Taz River. The annual river  
145 runoff gradually increases northward, from 160-220 mm y<sup>-1</sup> in the permafrost-free region to 280-320  
146 mm y<sup>-1</sup> in the Pur and Taz river basins located in the discontinuous to continuous permafrost zone  
147 (Nikitin and Zemtsov, 1986). A detailed physico-geographical, hydrology, lithology and soil  
148 description can be found in earlier works (Botch et al., 1995; Smith et al., 2004; Frey and Smith,  
149 2005, 2007; Frey et al., 2007a, b; Beilman et al., 2009) and in our recent limnological and  
150 pedological studies (Shirokova et al., 2013; Manasypov et al., 2014, 2015; Stepanova et al., 2015). A  
151 detailed map of studied region together with main permafrost provenances and river runoff in the  
152 WSL is given in **Fig. 1** and the list of sampled rivers grouped by watershed size and season is  
153 presented in **Table 1**. **Permafrost zonation in the WSL** shown in this figure is based on extensive

154 geocryological work in this region (Baulin et al., 1967; Gruzlov and Trofimov, 1980; Baulin, 1985;  
155 Liss et al., 2001). The hydrological parameters of the WSL rivers based to calculate a runoff contour  
156 line are described in the **Supplement 1**.

157  
158 *2.2. Chemical and isotope analyses and statistical treatment*

159 Altogether, 96 river samples were collected in early June 2013 (spring flood), August 2013  
160 and 2014 (summer baseflow), October 2013 (autumn) and February 2014 (winter baseflow) along  
161 the 1500 km latitudinal gradient (Table 1). All sampled rivers of western Siberian Lowland belong to  
162 the Kara Sea basin. **Seasonal sampling covered a full gradient** from south to north, except the month  
163 of October which was sampled only in rivers south of 60°N (12 rivers in total). The watershed area  
164 of sampled rivers ranged from 2 to 150,000 km<sup>2</sup>, not considering the Ob River in its medium course  
165 zone. Collected water samples were immediately filtered in pre-washed 30 mL PP Nalgene® flasks  
166 through single-use Minisart filter units (Sartorius, acetate cellulose filter) with a diameter of 25 mm  
167 and a pore size of 0.45 µm. The first 20 to 50 mL of filtrate were discarded. Filtered solutions for  
168 cation analyses were acidified (pH ~ 2) with ultrapure double-distilled HNO<sub>3</sub> and stored in pre-  
169 washed HDPE bottles. The preparation of bottles for sample storage was performed in a clean bench  
170 room (ISO A 10,000). Filtered samples for DOC, DIC, UV<sub>280 nm</sub> **absorbance** and anions were stored  
171 in the refrigerator maximum 3 weeks before the analyses. The effect of storage for DOC, DIC and  
172 optical measurements in boreal waters was found to be within the uncertainty of analysis (Ilina et al.,  
173 2014). Blanks were performed to control the level of pollution induced by sampling and filtration.  
174 The DOC blanks of filtrate never exceeded 0.1 mg/L which is quite low for the organic-rich river  
175 waters sampled in this study (i.e., 10–60 mg/L DOC). pH was measured in the field using a  
176 combined electrode calibrated against NIST buffer solutions (pH of 4.00 and 6.86 at 25°C). The  
177 accuracy of pH measurements was ±0.02 pH units. DOC and DIC were analyzed using a Carbon  
178 Total Analyzer (Shimadzu TOC VSCN) with an uncertainty better than 3%. Special calibration of the

179 instrument for analysis of both form of dissolved carbon in organic-rich, DIC-poor waters was  
180 performed as described elsewhere (Prokushkin et al., 2011). Major anions (Cl, SO<sub>4</sub>) concentrations  
181 were measured by ion chromatography (HPLC, Dionex ICS 2000) with an uncertainty of 2%. The  
182 UV- absorbance at 280 nm is used as a proxy for aromatic C and source of DOM in the river water. It  
183 was measured using 1 cm quartz cuvette in a CARY-50 UV-VIS spectrophotometer (Bruker, UK).  
184 Major cations (Ca, Mg, Na, K) and Si were determined with an ICP-MS Agilent ce 7500 with In and  
185 Re as internal standards and 3 various external standards, placed each 10 samples in a series of river  
186 water. Approximately 30% of samples were analyzed for Ca, Mg and Na concentration using atomic  
187 absorption spectroscopy (flame) with an uncertainty of 2 %. Reasonable and non-systematic  
188 agreement (between 5 and 10%) with the results of ICP MS analyses was achieved. Aqueous Si  
189 concentrations were also determined colorimetrically (molybdate blue method) with an uncertainty  
190 of 1% using a Technicon automated analyzer. The SLRS-5 (Riverine Water Reference Material for  
191 Trace Metals certified by the National Research Council of Canada) was used to check the accuracy  
192 and reproducibility of each analysis (Yeghicheyan et al., 2014).

193 The <sup>13</sup>C in dissolved inorganic carbon was analyzed in filtered river water sampled in bubble-  
194 free sealed glass bottles by gas chromatography and isotope mass spectrometry, using Delta V  
195 Advantage and Finnigan Gas-Bench-II in order to determine δ<sup>13</sup>C<sub>DIC</sub> (per mil relative to V-PDB;  
196 Fritz and Fontes, 1980). For these measurements, 0.1 mg of 100% H<sub>3</sub>PO<sub>4</sub> was added to the  
197 borosilicate vial and flushed with He (purity of 7.0) for 400 s. Afterwards, 1 mL of the sample was  
198 injected into the vial and shaken for 36 h at 24°C. Standard samples of C-O-1 and NBS-19 were  
199 routinely analyzed to test the accuracy of our measurements; typically, a disagreement of less than  
200 0.3 ‰ between the measured and certified values was observed, with a total estimated measurement  
201 uncertainty of ± 0.2 ‰.

202 The concentration of carbon and major elements in rivers were treated using the least squares  
203 method, Pearson correlation and one-way ANOVA (SigmaPlot version 11.0/Systat Software, Inc).



204 The ANOVA was used to reveal the differences between different permafrost zones. It was carried  
205 out using Dunn's method because each sampling period contained a different number of rivers.  
206 Regressions and power functions were used to examine the relationships between the dissolved  
207 component concentrations and the watershed area, river discharge, average latitude of the watershed  
208 and seasons. Comparison of DOC and major element concentration in rivers sampled in three main  
209 permafrost zones (continuous, discontinuous and permafrost-free regions), during all seasons and of  
210 different watershed size class was conducted using the non-parametric H-criterion Kraskal-Wallis  
211 test. First, we separated the watershed into four main classes encompassing all studied rivers (except  
212 Ob): < 100 km<sup>2</sup>, 100 to 1000 km<sup>2</sup>, 1000 to 10,000 km<sup>2</sup> and > 10,000 km<sup>2</sup>. We considered three main  
213 seasons in six different ranges of latitude (56 to 58°N, 58 to 60°N, 60 to 62°N, 62 to 64°N, 64 to  
214 66°N and 66 to 68°N). We checked for the variation of measured parameters of each watershed size  
215 as a function of latitude, separately in each season. In addition, a generalized assessment of the role  
216 of permafrost type and abundance on river water chemical composition was possible via separating  
217 all the sampled watersheds into three categories according to the permafrost distribution in WSL:  
218 permafrost-free, discontinuous and continuous permafrost.

219

### 220 3. Results

221 Results of major element analysis in rivers are listed in **Table S1** of the Supplement and the  
222 main results of statistical treatment are listed in **Table S2**. Based on the Kraskal-Wallis H statistics,  
223 the differences between the seasons and between different latitudes were found to be significantly  
224 higher (p-level < 0.0001) for most elements than the difference between watershed size classes,  
225 within each seasons and within each latitude range. This is illustrated for pH, DOC, DIC and  $\delta^{13}\text{C}_{\text{DIC}}$   
226 in **Figs 2, 3, 4 and 5**, respectively which show the measured value as a function of latitude for  
227 different watershed classes, individually for each main season. Similar plots for major cations (Ca,  
228 Mg, K) and Si are given in supplementary figures **S1, S2, S3 and S4**, respectively. The latitudinal

229 coverage of October was too small to be presented in these figures; however, the October data of 12  
230 rivers were used for statistical treatment and for assessing the permafrost impact. There is a clear and  
231 significant trend of concentration with latitude; the differences between different latitude ranges are  
232 significant at  $p < 0.0001$  for all elements, and at  $p < 0.05$  for Si. The effect of the watershed size on  
233 river water chemical composition in summer, winter and spring is much smaller than that of latitude  
234 ( $9 < H < 12$ ,  $p < 0.05$  and  $20 < H < 50$ ,  $p < 0.001$ , respectively). Considering all rivers  
235 simultaneously, the effect of the season is clearly seen at  $p < 0.001$  for all elements except DOC; the  
236 latter, however, is also statistically significant ( $H = 10.6$ ,  $p = 0.014$ ). Considering full data set of all  
237 seasons and watershed sizes, we distinguished three geographical zones in terms of the permafrost  
238 abundance: continuous, discontinuous and absent. For most river water parameters (pH, DIC, DOC,  
239 major anions and cations) the differences between three zones are significant ( $30 < H < 95$ , p-level  $<$   
240  $0.001$ ). Si concentration exhibited lower but statistically significant differences between different  
241 zones ( $H = 9.5$ ,  $p = 0.0086$ )

242         Considering all seasons and watershed sizes revealed a significant decrease of pH, Ca and Mg  
243 northward with the largest changes occurring at the beginning of discontinuous permafrost coverage  
244 (**Fig. S5 A, B and C**, respectively). The DOC and DIC also decrease in concentration with the  
245 increase in the degree of permafrost coverage (**Fig. S6 A, B**, respectively) whereas the isotopic  
246 composition of the DIC becomes progressively more negative northward (from ca. -15 ‰ in the  
247 permafrost-free zone to -20...-25 ‰ in the continuous permafrost zone, **Fig. S6 C**). In contrast, the  
248 effect of the permafrost on Si concentration is not clearly seen; the scatter of the data between  
249 different seasons and watersheds does not allow tracing any significant trend (not shown).

250         The optical properties of DOC remain essentially constant throughout the full range of  
251 watershed sizes, latitudes and seasons (**Fig. S7**). The largest variation of specific  $UV_{280\text{ nm}}$   
252 absorbance occurred in winter, when several DOC-rich waters from the southern (permafrost-free)  
253 part of WSL demonstrated quite low concentration of aromatic (colored) compounds.

254 **4. Discussion**

255 *4.1. Effect of latitude (permafrost and vegetation) on major cation, Si and DIC mobilization*  
256 *from the soil profile and groundwater to the river*

257 From general knowledge of environmental control on carbon and major element fluxes in  
258 rivers of the Russian subarctic (Prokushkin et al., 2011; Pokrovsky et al., 2012) and other boreal and  
259 subarctic regions (Laudon et al., 2004; Petrone et al., 2006; Walvoord and Striegl, 2007; Jantze et al.,  
260 2013; Giesler et al., 2014), we anticipate a decrease of most element concentration, including DOC,  
261 northward regardless of the season and the river size in the WSL due to (1) a decrease of chemical  
262 weathering intensity with the temperature, well demonstrated for igneous rocks such as basalts  
263 (Dessert et al., 2003) and granites (Oliva et al., 2003); (2) decrease of the thickness of peat deposits  
264 in total and the active soil (peat) layer in particular (Liss et al., 2001; Beilman et al., 2009; Stepanova  
265 et al., 2015 and references therein); (3) a decrease of plant biomass and related plant litter stock on  
266 the surface of the soils (Tyrtikov, 1979; Frey and Smith, 2007); (4) a shortening of the unfrozen  
267 period of the year and (5) a decrease of the degree of groundwater feeding (Romanovsky, 1983;  
268 Nikitin and Zemtsov, 1986; Fotiev, 1991). The factors capable of enhancing element export fluxes in  
269 northern (permafrost-bearing) rivers relative to southern (permafrost-free) rivers of the WSL are (1)  
270 the decrease of dissolved organic matter (DOM) respiration by heterotrophs in the water and soil  
271 column and thus the increasing removal of allochthonous DOC from the soil to the river (Striegl et  
272 al., 2005); (2) the increase of DOC and related element leaching from plant litter and topsoil  
273 (Pokrovsky et al., 2005; Giesler et al., 2006; Fraysse et al., 2010) during more pronounced massive  
274 freshet event or summer high flow (Michel and Vaneverdingen, 1994; McClelland et al., 2006;  
275 White et al., 2007); (3) the decrease of DOM retention (adsorption) on the mineral soil horizon  
276 because clay horizon is typically frozen in the north (Kawahigashi et al., 2004); 4) the decrease of  
277 authigenic clay and allophane mineral formation in the soil horizons (Targulian, 1971).

278 At the current, rather limited, stage of knowledge of mineral, organic soil horizons and plant  
279 biomass chemical composition and reactivity across the WSL, only a few environmental factors can  
280 be quantitatively tested based on river water chemical analyses. In the case of the dominance of  
281 groundwater feeding of the river, the decrease of element concentrations from water-rock interaction  
282 whose transport is not limited by availability of DOM (Ca, Mg, DIC) is expected to be most  
283 pronounced in winter, when the groundwater feeding is maximal (see Walvoord and Striegl (2007)  
284 for the Yukon River basin example). Moreover, in the permafrost-bearing zone during winter  
285 baseflow, one should expect significant difference in element concentration in winter between small  
286 rivers (weakly or not affected by taliks) and large rivers (essentially fed by taliks), as it is known  
287 from local geocryological conditions (Baulin et al., 1967; Romanovsky, 1983; Fotiev, 1989, 1991;  
288 Ivanov and Beshentsev, 2005). In spring, when the active layer is very thin and the majority of the  
289 soil column is frozen, the export from the watershed is dominated by surface flow and thus the  
290 difference in groundwater-related element concentration between (i) small and large rivers and (ii)  
291 north and south should be minimal. However, the abovementioned hypotheses are not supported by  
292 DIC, Ca and Mg concentrations observed in rivers (**Fig. 4, S1 and S2**). First, the DIC concentrations  
293 decrease between permafrost-free and discontinuous/continuous permafrost zones is a factor of  $15\pm 5$   
294 in winter (Fig. 4 A) and a factor of  $60\pm 10$  in spring (Fig. 4 B). Similarly, the decrease of Ca and Mg  
295 concentrations between south of  $59^{\circ}\text{N}$  and  $62\text{-}66^{\circ}\text{N}$  zones is  $\times 10$  in winter and  $\times 20\text{-}30$  in May. In  
296 fact, it is the spring period which exhibits the highest contrast in element concentrations between the  
297 south and the north. Second, for the latitude concentrations gradient from south to north, the relative  
298 DIC, Ca and Mg concentration change between large ( $1000\text{-}10,000\text{ km}^2$  and  $> 10,000\text{ km}^2$ ) and small  
299 ( $< 100\text{ km}^2$ ) rivers in winter is not statistically significant ( $p > 0.05$ ).

300 However, a systematic decrease of Ca concentration in the WSL rivers northward (Fig. S1,  
301 S5 B) is consistent with general decrease of Ca concentration in soil ecosystems as illustrated in Fig.  
302 S8. An order of magnitude decrease of Ca concentration in mineral horizons of SWL peat columns

303 occurred between 55 and 66°N (Stepanova et al., 2015). On a smaller scale, a **three times** decrease of  
304 exchangeable Ca concentration in alluvial soils of the Ob basin from 56 to 60°N was reported  
305 (Izerskaya et al., 2014). These observations **confirm a strong control of lithology and soil weathering**  
306 **on Ca concentration** in both deep and surface soil horizons and vegetation, which finally determines  
307 **the extent** of Ca transport via surface flux to the river.

308 North of 66°N, concentrations of Ca, Mg and sulfate increase relatively to their concentration  
309 at 62-66°N of discontinuous permafrost zone. This is especially pronounced during the summer  
310 period (Fig. S1 C, S2 C). We do not exclude here the influence of marine sedimentary deposits  
311 containing salts in the deep part of the mineral soil profile below the peat layer. These deposits are  
312 described in the low reaches of Taz and Pur rivers, based on sedimentary cores extracted during  
313 extensive drilling of the territory (Liss et al., 2001). This influence, however, cannot be  
314 unequivocally evidenced because (i) DIC concentrations also increase in summer, north of 66°N,  
315 although DIC is not likely to be affected by marine deposits, and (ii) chloride, an efficient marker of  
316 sea salts, is not increasing in the north (not shown).

317 The isotopic composition of DIC confirms the general features of DIC and cation  
318 concentration (Fig. 5). The groundwater feeding by taliks in winter is highly uniform over 10 degrees  
319 of latitude with the value of  $\delta^{13}\text{C}_{\text{DIC}}$  being equal to  $-15 \pm 5$  ‰, reflecting both carbonate/silicate  
320 weathering and a buildup of  $\text{CO}_2$  with a stronger respiratory signal (Finlay, 2003; Striegl et al., 2001;  
321 Giesler et al., 2014; Rinta et al., 2015). During this period, the variability of  $\delta^{13}\text{C}_{\text{DIC}}$  is the highest in  
322 small ( $< 100 \text{ km}^2$ ) watersheds, but no trend of isotopic composition with latitude could be evidenced  
323 at  $p < 0.05$  (Fig. 5 A). This isotopic signature is preserved in spring for southern ( $< 60^\circ\text{N}$ ) watersheds  
324 whereas in permafrost-affected regions,  $\delta^{13}\text{C}_{\text{DIC}}$  decreases to c.a.  $-25 \dots -20$  ‰ regardless of the river  
325 size and the type and the abundance of the permafrost (Fig. 5b). Such low values in the permafrost-  
326 affected zone could not anymore represent the influence of carbonate/silicate rock weathering by soil  
327  $\text{CO}_2$  and likely reflect direct microbial processing of soil and sedimentary organic matter (Waldron et

328 al., 2007; Giesler et al., 2013), with the DIC isotopic signature similar to that of organic carbon in  
329 western Siberia subarctic topsoil ( $-26\pm 2\text{‰}$ , Gentsch et al., 2015) and the Ob river organic sediments  
330 ( $-25$  to  $-27\text{‰}$ , Guo et al., 2004a).

331 A plausible explanation for the  $\delta^{13}\text{C}_{\text{DIC}}$  seasonal variation being mostly pronounced in the  
332 permafrost zone can be that microbial mineralization of dissolved organic carbon occurs most  
333 efficiently during the springtime, when significant amounts of fresh organic matter from ground  
334 vegetation are leached by melted snow. Higher bioavailability of vegetation leachates relative to  
335 more refractory soil humic and fulvic acids is known from studies in other temperate (van Hees et  
336 al., 2005) and boreal (Wickland et al., 2007) regions. The lack of  $\delta^{13}\text{C}_{\text{DIC}}$  decrease in spring relative  
337 to winter in the permafrost-free zone may stem from (i) significant input of the carbonate/silicate  
338 rock-hosted groundwaters during full period of the year in the south, or (ii) the different nature of  
339 DOM in the south, where the more refractory organic matter originated from peat leaching is less  
340 subjected to microbial processing compared to fresh vegetation leachates in the north, where the peat  
341 soil in spring is frozen. One has also take into account that the DIC concentrations in spring are a  
342 factor of 30 lower in the permafrost-bearing region relative to permafrost-free region (Fig 4 b). As  
343 such, a relatively small input of microbially-respired  $\text{CO}_2$  will be significantly more visible in the  
344  $\delta^{13}\text{C}_{\text{DIC}}$  value of the northern rivers compared to that of the southern rivers.

345 The variation of  $\delta^{13}\text{C}_{\text{DIC}}$  along the permafrost/latitude gradient helps to better explain the  
346 origin of DIC in rivers in contrasting permafrost zones. Consistent with a progressive decrease of the  
347 groundwater feeding of rivers northward (Nikitin and Zemtsov, 1986; Frey et al., 2007b), we observe  
348 a distinct trend of  $\delta^{13}\text{C}_{\text{DIC}}$  with the latitude during spring period, reflecting the shift of DIC origin  
349 from groundwater in the south to plant litter degradation and soil respiration in surface waters north  
350 of  $62^\circ\text{N}$  (Fig. 5 B). In winter, the  $\delta^{13}\text{C}_{\text{DIC}}$  is rather constant within the full latitudinal profile (Fig.  
351 5A) confirming the dominant role of carbonate/silicate mineral weathering by atmospheric and soil  
352  $\text{CO}_2$  dissolved in the groundwaters. Indeed, hydrological studies in the WSL revealed that the

353 groundwater feeding of small ( $< 10,000 \text{ km}^2$  watershed) rivers decreases from 20-30% in the  
354 discontinuous and sporadic/isolated part of the WSL to 3-6% in the northern, continuous permafrost  
355 zone (Novikov et al., 2009). These numbers agree with estimations based on RHS data of large  
356 western Siberian rivers (Nadym, Pur and Taz) and the left tributaries of the Yenisei River (Dubches,  
357 Elogyi and Turukhan; Nikitin and Zemtsov, 1986). According to more recent evaluations of Frey et  
358 al. (2007b), the groundwater contribution to summertime period river chemical composition ranges  
359 between 30 and 80% for the rivers located between 56 and 58°N.

360 Consistent with these findings, the pH values of 7 to 7.5 in the southern rivers observed both  
361 in winter and spring (Fig. 2 A, B) are indicative of carbonate/silicate rock input. The spring acid  
362 pulse, well established in other permafrost-free boreal regions (Buffam et al., 2007), is not at all  
363 pronounced in the south of the WSL but becomes clearly visible in the permafrost-affected, northern  
364 regions where the spring-time pH decreases to 5...6 (Fig. 2 B). This illustrates the more important  
365 role of plant litter and moss leaching in the permafrost-bearing zone on solute export from the  
366 watershed. In addition, the dominance of sands north of 62°N (Liss et al., 2001) may allow low  
367 molecular weight (LMW) organic acids migrate to the river from the soil profile. In the southern,  
368 permafrost-free zone, the dominating clays underneath the peat can adsorb acidic LMW organic  
369 compounds and thus do not allow the acid pulse to be clearly visible.

370 The increase of pH in summer relative to spring period is again less visible in the south than  
371 in the north (Fig. 2 C) and may reflect the persisting role of bedrock dissolution as well as the change  
372 of the river feeding regime, from top soil and vegetation in the north to the peat soil column leaching  
373 in the south. The summertime increase of river water pH north of 60°N, in the forest-tundra and  
374 tundra zone may be linked to (i) enhanced photosynthesis in rivers of the north due to better  
375 insolation and less forest shading and (ii) mobilization of DOM and other solutes from soil  
376 depressions rather than from watershed divides. The depressions are subjected to intense rinsing  
377 during the spring seasons, when the majority of soluble acidic compounds are flushed from the litter

378 and O<sub>e</sub> horizon. These mechanisms are evidenced from studies of the hydrological balance of frozen  
379 bogs performed in the northern part of studied territory (Novikov et al., 2009). In contrast, the  
380 watershed divides contain significant amounts of organic litter and release organic acids only in  
381 spring, when they are covered by temporary ponds of melted snow (see Manasyrov et al., 2015).  
382 This hydrological scheme of river water feeding is based on the seasonal multiannual observations  
383 on frozen bogs of the north of WSL (Novikov et al., 2009), although the chemical nature of DOM  
384 mobilized from different parts of the watershed remains unknown.

385 The importance of plant litter and ground vegetation leaching as element sources in western  
386 Siberian rivers can be assessed from the comparison of K concentrations as a function of latitude  
387 during different seasons (Fig. S 3). The most significant decrease of K concentration from the  
388 southern (< 59°N) to the northern (61-67°N) watersheds occurs in spring, during intense plant litter  
389 leaching. Regardless of latitude, K concentration follows the order spring > winter > summer with  
390 the highest concentrations, up to 2500 ppb, recorded in permafrost-free region. Given that the other  
391 cations, possibly originating from the water-mineral interaction at some depth, do not exhibit such  
392 high concentration in spring, we interpret the spring-time K “pulse” as indicative of plant litter  
393 leaching in productive taiga zone. This “pulse” is much less visible in the permafrost zone due to  
394 significantly lower biomass and primary productivity of forest-tundra and tundra biomes compared  
395 to the taiga of the WSL (Tyrtikov, 1979; Liss et al., 2001).

396 Despite significant variability of Si concentrations among rivers of various sizes across the  
397 latitude profile (Fig. S 4), the concentrations in the permafrost zone are not lower than those in the  
398 south of the WSL. Results of a previous study of WSL rivers during summer show that Si  
399 concentration are weakly dependent on latitude (Frey et al., 2007), as also confirmed in this work for  
400 spring flood and winter baseflow period. Given that (i) the dominance of permafrost north of 64°N  
401 implies very low groundwater feeding (4 to 6% of the annual discharge, see Nikitin and Zemtsov,  
402 1986; Novikov, 2009) and (ii) the upper part of the soil profile including its seasonally frozen and



403 unfrozen parts is mostly peat rather than silicate mineral sediments, the role of groundwater – silicate  
404 rock interaction in Si supply to northern rivers should be quite low. Therefore, we hypothesize that  
405 elevated concentrations of Si in northern rivers are due to peat leaching and degradation. A depletion  
406 of Si in rivers of the southern part of the WSL may be due to Si retained by abundant bog and forest  
407 vegetation. This is consistent with general setting of the WSL, recovering from the last glaciation  
408 (Liss et al., 2001), with contemporary peat accumulation in the south and old frozen peat  
409 thawing/degradation in the north.

410

#### 411 *4.2. DOC concentration across 1500 km latitude transect of variable permafrost coverage*

412 Results of organic carbon concentration in western Siberia rivers collected over various  
413 seasons of the year generally confirm the pioneering findings of Frey and Smith (2005). The strong  
414 statistically significant ( $p < 0.05$ ) contrast in DOC concentration between permafrost-free,  
415 discontinuous and continuous permafrost zone persists over the course of the year and each season  
416 except probably winter (Fig. 3 and Fig. S6 A). This difference is also seen in  $\delta^{13}\text{C}_{\text{DIC}}$  value among all  
417 three zones (Fig. S6 C) suggesting, on the annual scale, a more significant contribution of microbial  
418 processing of plant and soil organic carbon to  $\text{HCO}_3$  and  $\text{CO}_2$  of the river water in the permafrost-  
419 bearing zone compared to the permafrost-free zone.

420 In accordance with the conclusion reached by Frey and Smith (2005), the variation in  
421 hydrology may play a limited role in DOC variability and export from the watershed of WSL rivers.  
422 The gradient in DOC concentrations along the latitudinal profile remains similar between spring flood  
423 and summer baseflow (Fig. 3 b and c). Although the winter period does not exhibit such a clear  
424 difference between permafrost-free and permafrost-affected regions (Fig. 3 a), the contribution of the  
425 winter discharge to the annual flux of DOC is between 10 and 15% and as such does not significantly  
426 affect annual export of DOC from the watersheds.

427 In contrast, the gradient of organic carbon concentration along the latitudinal profile in spring  
428 will be mostly controlled by the difference in plant litter stock subjected to leaching by melted snow.  
429 As such, one would not expect any significant difference between large and small rivers at otherwise  
430 similar runoff, vegetation and bog coverage. This is partially confirmed by the similarity of the  
431  $UV_{280\text{ nm}} - \text{DOC}$  slope, corresponding to similar degree of DOM humification, among different  
432 seasons and latitudinal positions (Fig. S7). The uniform distribution of  $UV_{280}$  absorbance witnesses  
433 the main control of DOC by allochthonous (terrestrial) input from peat and/or ground vegetation  
434 leachates. The exceptions are the rivers Vasyugan (No 21), Shegarka (No 4) and Vatinsky Egan (No  
435 34) exhibiting low  $UV_{280\text{nm}}$  at high [DOC] (Fig. S7). These rivers are potentially affected by oil  
436 production sites and may contain some uncolored products of hydrocarbon oxidation in the  
437 underground waters.

438 Overall, results on western Siberia rivers generally confirm the conclusion of Finlay et al.  
439 (2006) on (i) the lack of groundwater contribution to streamflow in arctic watersheds and (ii) that  
440 river DOC dynamics are driven essentially by processes occurring at the soil surface. However we  
441 doubt the importance of large DOC pool production under very cold conditions as the main reason of  
442 sustained high concentration of DOC at snowmelt suggested by Finlay et al. (2006). Indeed, the plant  
443 litter degradation in winter, even in the warmest scenario, is minimal and does not contribute  
444 significantly to annual litter leaching (Bokhorst et al., 2010, 2013). Instead, we suggest fast plant  
445 litter and ground vegetation leaching in spring, at the very beginning of the snow melt. Such a fast  
446 enrichment in DOC and colored organic compounds of surface water depressions, on the order of  
447 several hours, has been observed in the discontinuous permafrost zone in early June (Manasypov et  
448 al., 2015). Significant release of DOC and nutrients from flooded ground vegetation in the southern  
449 part of WSL is also known (Vorovyev et al., 2015).

450 An unexpected result of the study of western Siberian watersheds is the lack of the  
451 enrichment in DOC of small headwater streams, in contrast to what was reported for Scandinavian

452 rivers and streams (Agren et al., 2007, 2014 and references therein). In WSL, especially in the  
453 northern, permafrost-affected zone, the small ( $< 100 \text{ km}^2$ ) streams yielded DOC concentration that  
454 were not statistically higher ( $p > 0.05$ ) than those of larger rivers, neither in spring flood nor in  
455 summer. A number of factors can be responsible for the observed difference between permafrost-free  
456 European and permafrost-bearing Siberian watersheds. In the north of western Siberia, the microbial  
457 processing of DOM in large rivers may be weakly pronounced. This is confirmed by the observation  
458 that the degree of light C isotope enrichment (lowering  $\delta^{13}\text{C}_{\text{DIC}}$ ) in spring is independent ( $p > 0.05$ )  
459 of the size of the river (Fig. 5 B), and, correspondently, of the water residence time on the watershed.  
460 According to Kawahigashi et al. (2004), the DOM in northern, permafrost-affected tributaries of the  
461 Yenisey River was significantly less biodegradable than that in southern tributaries. This may  
462 contribute to better preservation of DOM in the stream yielding its independence of the water travel  
463 time. Small watersheds of western Siberia exhibit a runoff and average slope very similar to that of  
464 the large rivers, given very flat orographic context of WSL. This contrasts the mountain regions of  
465 Sweden and Alaska where the headwater streams may exhibit higher runoff and thus higher export of  
466 the dissolved constituents. Finally, the riparian zone, very important for regulation DOC stock and  
467 export in small streams draining glacially-formed terrain of NW Europe (Dick et al., 2014;  
468 Kuglerova et al., 2014), is much less pronounced in western Siberia, where generally flat, frequently  
469 flooded areas dominate the watershed profile.

470         The elevated DOC concentrations in continuous permafrost zone, especially north of  $67^\circ\text{N}$   
471 observed in the present study (Fig. 3 B, C) are consistent with previous results showing that, at  
472 otherwise similar factors, the permafrost areas are a greater source of DOC than the areas with  
473 seasonal frost (Carey et al., 2003). In permafrost areas, meltwater travels through organic-rich layers  
474 in the form of so called supra-permafrost flow, as opposed to areas without impermeable permafrost  
475 table. In the latter, the infiltration of organic-rich surface waters to deep mineral layer and DOC  
476 sorption on clay minerals may occur thus decreasing the overall export of DOC (see Smedberg et al.,

477 2006 for discussion). Given the dominance of peat rather than minerals within the active (unfrozen)  
478 layers of soil profile, the difference between permafrost-free and permafrost-affected zones is even  
479 more accentuated in western Siberia.

480 A sketch of typical soil profiles of western Siberia in the permafrost-free and permafrost-  
481 bearing zone presenting DOC mobilization pathways from the soil to the river in the end of active  
482 period is illustrated in **Fig. 6**. The two cross sections shown in this figure are highly representative  
483 for two most contrasting cases of soil and watershed flux formation, corresponding to dark  
484 coniferous taiga in the permafrost-free zone and dwarf shrubs with green mosses of tundra and  
485 forest-tundra in frozen peatlands of continuous permafrost zone; both sites are located at the  
486 watershed divide. The detailed position of soil horizons and their attribution to FAO is based on  
487 available literature data (Tyrtikov, 1973, 1979; Liss et al., 2001; Pavlov and Moskalenko, 2002) and  
488 our recent investigations of the region (Loiko et al., 2015; Stepanova et al., 2015). We hypothesize  
489 that plant litter- and topsoil derived-DOC adsorbs on clay mineral horizons in the southern,  
490 permafrost-free and discontinuous/sporadic permafrost zone but lacks the interaction with minerals  
491 in the continuous permafrost zone. This assumption corroborates results found during another  
492 latitudinal river transect of Siberia, along the Yenisey river and its left tributaries draining peatlands  
493 of the WSL (Kawahigashi et al., 2004, 2006): the northern tributaries exhibited significantly higher  
494 DOC concentrations than the southern tributaries of this river. Specifically, given the significant  
495 thickness of the peat even in the northernmost part of the WSL and the active layer thickness of < 50  
496 to 80 cm (30 cm on mounds and 80 to 150 m in troughs and depressions, Tyrtikov (1973, 1979),  
497 Baulin et al. (1967), Baulin (1985), Khrenov (2011), Novikov et al. (2009)), even in the region of  
498 continuous permafrost development, peat soil interstitial solutions might not enter in contact with the  
499 mineral soil horizon and thus will not decrease their DOC concentration during migration from the  
500 soil column to the river along the permafrost impermeable layer (**Fig. 6**).

501 Therefore, in the permafrost zone, the DOC export is strongly controlled by DOC residence  
502 time and water travel pathway through organic topsoil and lichens, moss and litter leaching vs. peat  
503 and mineral layer leaching (Fig. 6 B). In this case, it is only the thickness of the unfrozen peat and  
504 local permafrost coverage that control the DOC export from the soil to the river. As a result, DOC  
505 concentration in the streams will be weakly dependent on the watershed size and seasons. It follows  
506 that DOC export from peat soils by medium size ( $n \cdot 10,000 - n \cdot 100,000 \text{ km}^2$ ) rivers located entirely  
507 in the permafrost zone may be higher than that of the larger rivers, crossing permafrost-free regions.  
508 This hypothesis is supported by available information on DOC yield by rivers of the WSL. Thus,  
509 rivers Taz ( $s = 150,000 \text{ km}^2$ ), Pur ( $112,000 \text{ km}^2$ ) and Nadym ( $64,000 \text{ km}^2$ ), entirely located in the  
510 discontinuous permafrost zone, exhibit 1.9, 2.1, and  $4.4 \text{ t km}^{-2} \text{ y}^{-1}$  DOC yield, respectively (Gordeev  
511 et al., 1996 and calculated based on data of the RHS). This is significantly higher than the value  
512 suggested for the Ob River ( $1.2 \text{ t km}^{-2} \text{ y}^{-1}$ , Gordeev et al., 1996).

513

#### 514 *4.3. Possible evolution of chemical composition and fluxes of western Siberian rivers under* 515 *climate change scenarios*

516 The most likely scenario of the climate change in western Siberia consists of shifting the  
517 permafrost boundary further north and increase of the active layer thickness (Pavlov and  
518 Moskalenko, 2002; Frey, 2003; Romanovsky et al., 2010; Vasiliev et al., 2011; Anisimov et al.,  
519 2013). The permafrost boundary change, equivalent to the northward shift of the river latitudes, may  
520 decrease maximum 2 fold the DOC concentrations of the most northern rivers due to the change of  
521 continuous to discontinuous permafrost. The thickness of the active layer is projected to increase  
522 more than 30% during this century across the tundra area in the Northern Hemisphere (Anisimov et  
523 al., 2002; Stendel and Christensen, 2002; Dankers et al., 2011). In the WSL, this increase will be  
524 most dramatic in the north, where the peat deposits are thinner than those in the discontinuous  
525 permafrost zone (Botch et al., 1995; Liss et al., 2001; Novikov et al., 2009; Kremenetsky et al.,

526 2004). Assuming a short-term (hundred years) scenario in the WSL, we hypothesize that the main  
527 consequences of this increase will be the involvement of upper clay horizon and sand/silts in water  
528 pathways within the soil profile. As a result, the DOC originated from the upper peat layer leaching  
529 and plant litter degradation will be retained on mineral surfaces and in the clay interlayers (Kaiser et  
530 al., 2007; Oosterwoud et al., 2010; Mergelov and Targulian, 2011; Gentsch et al., 2015). To which  
531 degree this change of water pathways in the soil column may affect the other dissolved components  
532 cannot be predicted. However, this effect for inorganic solutes is expected to be lower than that for  
533 DOC, given much lower affinity of  $\text{HCO}_3^-$ , major cations and Si to clay surfaces and the lack of  
534 unweathered (primary) silicate rocks underneath peat soil column. Nevertheless, the possibility of  
535 leaching of inorganic components from the mineral layers should be considered. For example, DOC  
536 export exceeded DIC export in a tributary of the Yukon River during high flow, whereas DIC  
537 predominated during low flow and the DIC yields increased with decreasing permafrost extend  
538 (Dornblaser and Striegl, 2015). Unfortunately, no time series on hydrochemistry of rivers of  
539 continuous permafrost development, north of  $64^\circ\text{N}$ , are available to test the hypothesis of the impact  
540 of climate change on a possible decreasing DOC flux from frozen peatlands and the DOC/DIC  
541 change due to ongoing decrease of permafrost protection of mineral layer from adsorbing DOC.

542         Important modifications linked to the climate change in boreal and subarctic zones concern  
543 the change of the hydrological regime (Karlsson et al., 2015), in particular the increase of the winter  
544 baseflow (Yang et al., 2004; Ye et al., 2009; Serreze et al., 2000) due to the increase of the  
545 groundwater feeding (Frey et al., 2007a,b; Walvoord and Striegl, 2007; Rowland et al., 2010;  
546 Walvoord et al., 2012), coupled with the increase of the overall precipitation and, consequently,  
547 water runoff (Peterson et al., 2002, McClelland et al., 2006). Here, we argue that 10 to 30%  
548 modification in the annual runoff will be within the variation of the DOC and cation concentrations  
549 between watersheds of various sizes observed in the present study and as such will not significantly  
550 affect the export fluxes of river water constituents.

551 To which degree the ongoing DOC concentration and flux rise in rivers, linked to climate  
552 change and/or acidification as reported in Western Europe and Canada (Worrall et al., 2004; Porcal  
553 et al., 2009) can be applied to the WSL is unknown. However, we did not observe any significant  
554 (i.e., > 30%) change of DOC fluxes over past 30 to 40 years neither in the boreal non-permafrost  
555 pristine region of NW Russia (Severnaya Dvina River, Pokrovsky et al., 2010), nor in the Central  
556 Siberian, continuous permafrost rivers of the Yenisei basin (Pokrovsky et al., 2005). Moreover, a  
557 decrease of DOC fluxes in the Yukon River was reported and suggested to be linked to enhanced  
558 mineralization of DOC by biota (Striegl et al., 2005). Note also that the more recent evaluation of  
559 the Ob River DOC discharge using flow-weighted concentration of 9.4 mg/L measured in 2003-2007  
560 (Cooper et al., 2008) gives a flux of 1.3 t C/km<sup>2</sup>/y, well comparable with earlier estimate of 1.2 t  
561 C/km<sup>2</sup>/y, based on the RHS data of 1950-1990 (Gordeev et al., 1996).

562 The increase of vegetation productivity reported for Arctic river basins (Sturm et al., 2001,  
563 Tape et al., 2006; Kirilyanov et al., 2012) will most likely proportionally increase the spring-time K  
564 flux due to its leaching from plant litter but likely decrease the summer –time Si flux, especially in  
565 the permafrost-bearing regions. The increase of vegetation density in the next decades to centuries  
566 may produce a transient uptake of Si by growing vegetation in the discontinuous permafrost zone  
567 during summer period. However, this potential decrease of Si export flux may be outweighed by the  
568 increasing release of Si from previously frozen mineral horizons and as such the overall modification  
569 of the Si concentration and riverine flux in discontinuous/continuous permafrost zone may be smaller  
570 than those projected by simple latitudinal shift.

571

## 572 *5. Conclusions*

573 An unexpected result of the present study was rather low sensitivity of DOC, DIC, cations  
574 and Si concentration and fluxes to the size of the river. The season also played a secondary role in  
575 determining element concentration pattern. The most important governing parameter for

576 concentrations of dissolved river water components was the latitude, allowing us to distinguish  
577 between permafrost-free, discontinuous and continuous permafrost regions. A northward decrease of  
578 DIC and dissolved cation (Ca, Mg) concentrations in the WSL rivers was mostly pronounced during  
579 spring flood. It was consistent with general trend of soil cations (such as Ca) concentration decrease  
580 from the south to the north, reported for peat, moss and mineral layer.

581 Both seasonal and latitudinal patterns of DOC and DIC concentrations in WSL rivers are  
582 consistent with previous observations that in continuous permafrost zone of frozen peat bogs, the  
583 underlining mineral layer is not reactive being protected by the permafrost so that the major part of  
584 the active layer is located within the organic (peat), not mineral matrix. The variation of  $\delta^{13}\text{C}_{\text{DIC}}$   
585 along the permafrost/latitude gradient is consistent with a progressive decrease of the groundwater  
586 feeding of rivers northward, reflecting the shift of DIC origin from groundwater in the south to plant  
587 litter degradation and soil respiration in surface waters north of 62°N. In winter, the  $\delta^{13}\text{C}_{\text{DIC}}$  is rather  
588 constant within the full latitudinal profile confirming the dominant role of carbonate/silicate mineral  
589 weathering by atmospheric and soil  $\text{CO}_2$  dissolved in the groundwaters.

590 Because the thickness of the unfrozen peat and local permafrost coverage essentially control  
591 the DOC export from the soil to the river, the DOC concentration in the streams is weakly dependent  
592 on the watershed size and seasons. It follows that DOC export from peat soils by medium size (<  
593 100,000 km<sup>2</sup>) rivers located entirely in the permafrost zone may be higher than that of the larger  
594 rivers, crossing permafrost-free regions. Assuming a short-term (hundred years) climate warming  
595 scenario in the WSL, we hypothesize that the increase of the active layer thickness will bring about  
596 the involvement of upper clay horizon and sand/silts in water pathways within the soil profile. As a  
597 result, the DOC export in permafrost-affected watersheds may decrease whereas the export of DIC  
598 and major cations will increase. Enhanced non-stationary uptake of Si by growing vegetation in the  
599 permafrost-bearing zone may attenuate expected increase of its riverine concentration linked to  
600 progressive involvement of thawed mineral horizons.



601 **Acknowledgements**

602 We acknowledge support from a BIO-GEO-CLIM grant from the Ministry of Education and Science  
603 of the Russian Federation and Tomsk State University (No 14.B25.31.0001) and partial support from  
604 RSF grant No 15-17-10009 “Evolution of thermokarst ecosystems” awarded to LS (25%).

605

606 **Authors contribution:** O. Pokrovsky designed the study and wrote the paper; R. Manasyrov, I.  
607 Kritzkov, S. Vorobyev performed sampling, analysis of major cations and their interpretation; S.  
608 Vorobyev and S. Kirpotin were responsible for the choice of sampling objects and statistical  
609 treatment; S. Loiko and S. Kulizhsky provided the background information on soil, peat, and  
610 permafrost active layer; L. Shirokova was in charge of DOC, DIC and anion measurements and their  
611 interpretation; BP performed <sup>13</sup>C measurements and their interpretation; L. Kolesnichenko provided  
612 GIS-based interpretation, mapping and identification of river watersheds; S. Kopysov and V.  
613 Zemtsov performed all primary hydrological data collection, their analysis and interpretation. All 12  
614 authors participated in field expeditions. Each co-author have seen and approved the final paper and  
615 contributed to writing the manuscript.

616

617 **References**

- 618 Agren, A., Buffam, I., Jansson, M., and Laudon, H.: Importance of seasonality and small streams for  
619 the landscape regulation of dissolved organic carbon export, *J. Geophys. Res.*, 112, G03003,  
620 doi:10.1029/2006JG000381, 2007.
- 621 Agren, A. M., Buffam, I., Cooper, D. M., Tiwari, T., Evans, C. D., and Laudon, H.: Can the  
622 heterogeneity in stream dissolved organic carbon be explained by contributing landscape  
623 elements? *Biogeosciences*, 11, 1199–1213, 2014.
- 624 Anisimov, O. A., Shiklomanov, N. I., and Nelson, F. E.: Variability of seasonal thaw depth in  
625 permafrost regions: A stochastic modeling approach, *Ecol. Model.*, 153, 217–227, 2002.
- 626 Anisimov, O., Kokorev, V., and Zhil'tsova, Y.: Temporal and spatial patterns of modern climatic  
627 warming: Case study of Northern Eurasia, *Climatic Change*, 118, 871–883, 2013.
- 628 Anisimova, N. P.: *Cryohydrochemical Features of Permafrost Zone*, Nauka, 1981 (in Russian).
- 629 Bagard, M. L., Chabaux, F., Pokrovsky, O. S., Prokushkin, A. S., Viers, J., Dupré, B., and Stille, P.  
630 Seasonal variability of element fluxes in two Central Siberian rivers draining high latitude  
631 permafrost dominated areas, *Geochim. Cosmochim. Ac.*, 75, 3335–3357, 2011.
- 632 Bagard, M. L., Schmitt, A. D., Chabaux, F., Pokrovsky, O. S., Viers, J., Stille, P., Labolle, F., and

633 Prokushkin, A. S.: Biogeochemistry of stable Ca and radiogenic Sr isotopes in larch-covered  
634 permafrost-dominated watersheds of Central Siberia, *Geochim. Cosmochim. Ac.*, 114, 169–  
635 187, 2013.

636 Baulin, V. V.: Permanently frozen rocks of oil and gaz producing regions of the USSR, Moscow,  
637 Nauka, 176 pp, 1985 (in Russian).

638 Baulin, V. V., Belopukhova, E. B., Dubikov, G. I., Shmelev, L. M.: Geocryological conditions of  
639 western Siberia Lowland, Moscow, Nauka, 214 pp, 1967 (in Russian).

640 Beaulieu, E., Godderis, Y., Donnadieu, Y., Labat, D., and Roelandt, C.: High sensitivity of the  
641 continental-weathering carbon dioxide sink to future climate change, *Nature Climate Change*,  
642 2, 346–349, 2012.

643 Beilman, D. W., MacDonald, G. M., Smith, L. C., and Reimer, P. J.: Carbon accumulation in  
644 peatlands of West Siberia over the last 2000 years, *Global Biogeochem. Cy.*, 23, GB1012,  
645 doi:10.1029/2007GB003112, 2009.

646 Botch, M. S., Kobak, K. I., Vinson, T. S., and Kolchugina, T. P.: Carbon pools and accumulation in  
647 peatlands of the former Soviet Union, *Global Biogeochem. Cy.*, 9, 37–46, doi:  
648 10.1029/94GB03156, 1995.

649 Brown, J., Ferrians Jr., O. J., Heginbottom, J. A., and Melnikov, E. S.: Circum-Arctic Map of  
650 Permafrost and Ground Ice Conditions. National Snow and Ice Data Center/World Data Center  
651 for Glaciology, Boulder, CO, USA, Digital media, 2002.

652 Bokhorst, S., Bjerke, J. W., Melillo, J., Callaghan, T. V., and Phoenix, G. K.: Impacts of extreme  
653 winter warming events on litter decomposition in a sub-Arctic heathland, *Soil Biol. Biochem.*,  
654 42, 611–617, 2010.

655 Bokhorst, S., Metcalfe, D. B., and Wardle, D. A.: Reduction in snow depth negatively affects  
656 decomposers but impact on decomposition rates is substrate dependent, *Soil Biol. Biochem.*,  
657 62, 157–164, 2013.

658 Buffam, I., Laudon, H., Temnerud, J., Mörth, C.-M., and Bishop, K.: Landscape-scale variability of  
659 acidity and dissolved organic carbon during spring flood in a boreal stream network, *J.*  
660 *Geophys. Res.*, 112, G01022, doi:10.1029/2006JG000218, 2007.

661 Carey, S. K.: Dissolved organic carbon fluxes in a discontinuous permafrost subarctic alpine  
662 catchment, *Permafrost Periglac.*, 14, 161–171, 2003.

663 Cooper, L. W., McClelland, J. W., Holmes, R. M., Raymond, P. A., Gibson, J. J., Guay, C. K., and  
664 Peterson, B. J.: Flow-weighted values of runoff tracers ( $\delta^{18}\text{O}$ , DOC, Ba, alkalinity) from the  
665 six largest Arctic rivers, *Geophys. Res. Lett.*, 35, L18606, doi:10.1029/2008GL035007, 2008.

666 Dankers, R., Burke, E. J., and Price, J.: Simulation of permafrost and seasonal thaw depth in the  
667 JULES land surface scheme, *Cryosphere*, 5, 773–790, 2011.

668 Dessert, C., Dupré, B., Gaillardet, J., Francois, L. M., and Allégre, C. J.: basalt weathering laws and  
669 the impact of basalt weathering on the global carbon cycle, *Chem. Geol.*, 202, 257–273, 2003.

670 Dick, J. J., Tetzlaff, D., Birkel, C., and Soulsby, C.: Modelling landscape controls on dissolved  
671 organic carbon sources and fluxes to streams, *Biogeochemistry*, doi: 10.1007/s10533-014-  
672 0046-3, 2014.

673 Dittmar, Th. and Kattner, G.: The biogeochemistry of the river and shelf ecosystem of the Arctic  
674 Ocean: a review, *Mar. Chem.*, 83, 103–120, 2003.

675 Dornblaser, M. M. and Striegl, R. G.: Switching predominance of organic versus inorganic carbon  
676 exports from an intermediate-size subarctic watershed, *Geophys. Res. Lett.*, 42, 386–394, 2015.

677 Engström, E., Rodushkin, I., Ingri, J., Baxter, D., Ecke, F., Österlund, H., and Öhlander, B.:  
678 Temporal isotopic variations of dissolved silicon in a pristine boreal river, *Chem. Geol.*, 271,  
679 142–152, 2010.

680 FAO, Guidelines for soil description. 4<sup>th</sup> edition. Rome. FAO, 2006.

681 Feng, X. J., Vonk, J. E., van Dongen, B. E., Gustafsson, O., Semiletov, I. P., Dudarev, O. V., Wang,  
682 Z. H., Montlucon, D. B., Wacker, L., and Eglinton, T.I.: Differential mobilization of terrestrial  
683 carbon pools in Eurasian Arctic river basins, *P. Natl. Acad. Sci. USA*, 110, 14168–14173,  
684 2013.

685 Finlay, J. C.: Controls of streamwater dissolved inorganic carbon dynamics in a forested watershed,  
686 *Biogeochemistry*, 62, 231–252, doi:10.1023/A:1021183023963, 2003.

687 Finlay, J., Neff, J., Zimov, S., Davydova, A., and Davydov, S.: Snowmelt dominance of dissolved  
688 organic carbon in high-latitude watersheds: Implications for characterization and flux of river  
689 DOC, *Geophys. Res. Lett.*, 33, L10401, doi:10.1029/2006GL025754, 2006.

690 Fotiev, C. M. : Taliks and their formations (Taliki and zakonomernosti ix formirovaniya), In :  
691 *Geocryology of the USSR, Western Siberia*, Ed : E.D. Ershov, Moscow, Nedra, pp. 72-84,  
692 1989. (in Russian)

693 Fotiev, C. M. : Formation of taliks of Western Siberia, In : *Permanently frozen rocks and cryogenic*  
694 *processes*, Moscow, Nauka, pp. 71-78, 1991. (in Russian).

695 Fraysse, F., Pokrovsky, O. S., and Meunier, J.-D.: Experimental study of terrestrial plant litter  
696 interaction with aqueous solutions, *Geochim. Cosmochim. Ac.*, 74, 70–84, 2010.

697 Frey, K. E. and Smith, L. C.: Recent temperature and precipitation increases in West Siberia and  
698 their association with the Arctic Oscillation, *Polar Res.*, 22, 287–300, 2003.

699 Frey, K. E., McClelland, J. W., Holmes, R. M., and Smith, L. C.: Impacts of climate warming and

700 permafrost thaw on the riverine transport of nitrogen and phosphorus to the Kara Sea, J.  
701 Geophys. Res., 112, G04S58, doi: 10.1029/2006JG000369, 2007a.

702 Frey, K. E., Siegel, D. I., and Smith, L. C.: Geochemistry of west Siberian streams and their potential  
703 response to permafrost degradation, Water Resour. Res., 43, W03406,  
704 doi:10.1029/2006WR004902, 2007b.

705 Frey, K. E. and Smith, L.C.: Amplified carbon release from vast West Siberian peatlands by 2100,  
706 Geophys. Res. Lett., 32, L09401, doi:10.1029/2004GL022025, 2005.

707 Frey, K. E. and Smith, L. C.: How well do we know northern land cover? Comparison of four global  
708 vegetation and wetland products with a new ground-truth database for West Siberia, Global  
709 Biogeochem. Cy., 21, GB1016, doi:10.1029/2006GB002706, 2007.

710 Frey, K. E. and McClelland, J. W.: Impacts of permafrost degradation on arctic river  
711 biogeochemistry, Hydrol. Process., 23, 169–182, 2009.

712 Fritz, P. and Fontes, J. C.: Handbook of environmental isotope geochemistry, v. 1, Elsevier, 1980.

713 Gaillardet, J., Millot, R., and Dupré, B.: Chemical denudation rates of the western Canadian orogenic  
714 belt: the Stikine terrane, Chem. Geol., 201, 257–279, 2003.

715 Gebhardt, A. C., Gaye-Haake, B., Unger, D., Lahajnar, N., and Ittekkot, V.: Recent particulate  
716 organic carbon and total suspended matter fluxes from the Ob and Yenisei Rivers into the Kara  
717 Sea (Siberia), Mar. Geol., 207, 225–245, 2004.

718 Gentsch, N., Mikutta, R., Alves, R. J. E., Barta, J., Capek, P., Gitte, A., Hugelius, G., Kuhry, P.,  
719 Lashchinskiy, N., Palmtag, J., Richter, A., Santrucková, H., Schneckner, J., Shibistova, O.,  
720 Urich, T., Wild, B., and Guggenberger, G.: Storage and transformation of organic matter  
721 fractions in cryoturbated permafrost soils across the Siberian Arctic, Biogeosciences Discuss.,  
722 12, 2697–2743, 2015.

723 Geological composition of the USSR, v. 1, Stratigraphy, Moscow, Gostoptekhizdat, 588 pp., 1958  
724 (in Russian).

725 Giesler, R., Högberg, M. N., Strobel, B. W., Richter, A., Nordgren, A., and Högberg, P.: Production  
726 of dissolved organic carbon and low-molecular weight organic acids in soil solution driven by  
727 recent tree photosynthate, Biogeochemistry, 84, 1–12, 2006.

728 Giesler, R., Mörth, C.-M., Karlsson, J., Lundin, E. J., Lyon, S. W., and Humborg, C.: Spatiotemporal  
729 variations of pCO<sub>2</sub> and δ<sup>13</sup>C-DIC in subarctic streams in northern Sweden, Global  
730 Biogeochem. Cy., 27, 176–186, doi:10.1002/gbc.20024, 2013.

731 Giesler, R., Lyon, S. W., Mörth, C.-M., Karlsson, J., Karlsson, E. M., Jantze, E. J., Destouni, G., and  
732 Humborg, C.: Catchment-scale dissolved carbon concentrations and export estimates across six

733 subarctic streams in northern Sweden, *Biogeosciences*, 11, 525–537, doi:10.5194/bg-11-525-  
734 2014, 2014.

735 Gislason, S. R., Arnorsson, S., and Armannsson, H.: Chemical weathering of basalt as deduced from  
736 the composition of precipitation, rivers and rocks in Southwest Iceland: effect of runoff, age of  
737 rocks and vegetative/glacial cover, *Am. J. Sci.*, 296, 837–907, 1996.

738 Gordeev, V. V. and Sidorov, I. S.: Concentrations of major elements and their outflow into the  
739 Laptev Sea by the Lena River, *Mar. Chem.*, 43, 33–46, 1993.

740 Gordeev, V. V., Martin, J.-M., Sidorov, I. S., and Sidorova, M. V.: A reassessment of the Eurasian  
741 river input of water, sediment, major elements, and nutrients to the Arctic Ocean, *Am. J. Sci.*,  
742 296, 664–691, 1996.

743 Gordeev, V. V., Rachold, V., and Vlasova, I. E.: Geochemical behavior of major and trace elements  
744 in suspended particulate material of the Irtysh river, the main tributary of the Ob river, Siberia,  
745 *Appl. Geochem.*, 19, 593–610, 2004.

746 Gruzdov, A. V., Trofimov, V. T.: Abundance and temperature of permanently frozen and seasonally  
747 frozen rocks, In: *Cryogenic composition and ice content of permanently frozen rocks of*  
748 *Western Siberian Plate*, Moscow, Izd-vo Moscow State University, p. 29-57, 1980. (in  
749 Russian)

750 Guo, L., Semiletov, I., Gustafsson, O., Ingri, J., Andersson, P., Dudarev, O., and White, D.:  
751 Characterization of Siberian Arctic coastal sediments: Implications for terrestrial organic  
752 carbon export, *Global Biogeochem. Cy.*, 18, GB1036, doi:10.1029/2003GB002087, 2004a.

753 Guo, L., Zhang, J. Z., and Guéguen, C.: Speciation and fluxes of nutrients (N, P, Si) from the upper  
754 Yukon River, *Global Biogeochem. Cy.*, 18, GB1038, doi:10.1029/2003GB002152, 2004b.

755 Guo, L., Ping, C. L., and MacDonald, R.W.: Mobilization pathways of organic carbon from  
756 permafrost to Arctic rivers in a changing climate, *Geophys. Res. Lett.*, 34, L13603,  
757 doi:10.1029/2007GL030689, 2007.

758 Holmes, R. M., Peterson, B. J., Gordeev, V. V., Zhulidov, A. V., Meybeck, M., Lammers, R. B., and  
759 Vörösmarty, C. J.: Flux of nutrients from Russian rivers to the Arctic Ocean: Can we establish  
760 a baseline against which to judge future changes? *Water Resour. Res.*, 36, 2309–2320, 2000.

761 Holmes, R. M., Peterson, B. J., Zhulidov, A. V., Gordeev, V. V., Makkaveev, P. N., Stunzhas, P. A.,  
762 Kosmenko, L. S., Köhler, G. H., and Shiklomanov, A. I.: Nutrient chemistry of the Ob' and  
763 Yenisey Rivers, Siberia: results from June 2000 expedition and evaluation of long-term data  
764 sets, *Mar. Chem.*, 75, 219–227, 2001.

765 Holmes, R. M., McClelland, J. W., Peterson, B. J., Tank, S. E., Bulygina, E., Eglinton, T. I.,  
766 Gordeev, V. V., Gurtovaya, T. Y., Raymond, P. A., Repeta, D. J., Staples, R., Striegl, R. G.,

767 Zhulidov, A. V., and Zimov, S. A.: Seasonal and annual fluxes of nutrients and organic matter  
768 from large rivers to the Arctic Ocean and surrounding seas, *Estuar. Coast.*, 35, 369–382,  
769 doi:10.1007/s12237-011-9386-6, 2012.

770 Ilina, S. M., Drozdova, O. Y., Lapitskiy, S. A., Alekhin, Y. V., Demin, V. V., Zavgorodnyaya, Y. A.,  
771 Shirokova, L. S., Viers, J., Pokrovsky, O. S.: Size fractionation and optical properties of dissolved  
772 organic matter in the continuum soil solution-bog-river and terminal lake of a boreal watershed,  
773 *Org. Geochem.*, 66, 14–24, 2014.

774 Ivanov, Yu. K. and Beshentsev, V. A.: Paleo-geographical aspects of formation of chemical  
775 composition of fresh underground waters in Yamalo-Nenets region, *Lithosphere*, No 4, 188–196,  
776 2005.

777 Izerskaia, L. A., Vorobyev, S. N, Vorobyeva, T. E., Kolesnichenko, L. G, Zakharchenko, A. V.: The  
778 concentration of Mn, Cu, Zn, Co, B, Sr, Cd and Pb in alluvial soils of the Ob River (forest-steppe,  
779 southern taiga and central taiga), *Internat. J. Environ. Stud.*, 71(5), 691–697, 2014.

780 Jantze, E. J., Lyon, S. W., and Destouni, G.: Subsurface release and transport of dissolved carbon in a  
781 discontinuous permafrost region, *Hydrol. Earth Syst. Sc.*, 17, 3827–3839, doi:10.5194/hess-17-  
782 3827-2013, 2013.

783 Kaiser, C., Meyer, H., Biasi, C., Rusalimova, O., Barsukov, P., and Richter, A.: Conservation of soil  
784 organic matter through cryoturbation in arctic soils in Siberia, *J. Geophys. Res.*, 112, 9–17, 2007.

785 Karlsson, J. M., Jaramillo, F., and Destouni, G.: Hydro-climatic and lake change patterns in Arctic  
786 permafrost and non-permafrost areas, *J. Hydrol.*, in press, 2015.

787 Kawahigashi, M., Kaiser, K., Kalbitz, K., Rodionov, A., and Guggenberger, G.: Dissolved organic  
788 matter in small streams along a gradient from discontinuous to continuous permafrost, *Glob.  
789 Change Biol.*, 10, 1576–1586, doi:10.1111/j.1365-2486.2004.00827.x, 2004.

790 Kawahigashi, M., Kaiser, K., Rodionov, A., and Guggenberger, G.: Sorption of dissolved organic  
791 matter by mineral soils of the Siberian forest tundra. *Global Change Biol.*, 12, 1868–1877,  
792 doi:10.1111/j.1365-2486.2006.01203.x, 2006.

793 Keller, K., Blum, J. D., and Kling, G. W.: Stream geochemistry as an indicator of increasing  
794 permafrost thaw depth in an arctic watershed, *Chem. Geol.*, 273, 76–81, 2010.

795 Kirdyanov, A. V., Hagedorn, F., Knorre, A. A., Fedotova, E. V., Vaganov, E. A., Naurzbaev, M. M.,  
796 Moiseev, P. A., and Rigling, A.: 20th century tree-line advance and vegetation changes along  
797 an altitudinal transect in Putorana Mountains, northern Siberia, *Boreas*, 41, 56–67, 2012.

798 Khrenov V. Ya.: Soils of cryolithozone of western Siberia: Morphology, physico-chemical properties  
799 and geochemistry, Nauka, Moscow, 214 pp., 2011 (in Russian).

800 Kremenetski, K. V., Velichko, A. A., Borisova, O. K., MacDonald, G. M., Smith, L. C., Frey, K. E.,  
801 and Orlova, L. A.: Peatlands of the West Siberian Lowlands: Current knowledge on zonation,  
802 carbon content, and Late Quaternary history, *Quaternary Sci. Rev.*, 22, 703–723, 2003.

803 Kuglerová, L., Jansson, R., Ågren, A., Laudon, H., and Malm-Renöfält, B.: Groundwater discharge  
804 creates hotspots of riparian plant species richness in a boreal forest stream network, *Ecology*  
805 95, 715–725. <http://dx.doi.org/10.1890/13-0363.1>, 2014.

806 Laudon, H., Köhler, S., and Buffam, I.: Seasonal TOC export from seven boreal catchments in  
807 northern Sweden, *Aquat. Sci.*, 66, 223–230, 2004.

808 Liss, O. L., Abramova, L. I., Avetov, N. A., Berezina, N. A., Inisheva, L. I., Kurnishnikova, T. V.,  
809 Sluka, Z. A., Tolpysheva, T. Yu., and Shvedchikova, N. K.: Wetland systems of West Siberia  
810 and their importance for nature conservation, Grif i K publisher, Tula, Russia, 584 pp., 2001,  
811 ISBN 5-8125-0150-5 (in Russian).

812 Lobbes, J. M., Fitznar, H. P., and Kattner, G.: Biogeochemical characteristics of dissolved and  
813 particulate organic matter in Russian rivers entering the Arctic Ocean, *Geochim. Cosmochim.*  
814 *Ac.*, 64, 2973–2983, 2000.

815 Loiko, S. V., Geras'ko, L. I., Kulizhskii, S. P., Amelin, I. I., Istigechev, G. I.: Soil cover patterns in the  
816 northern part of the area of aspen-fir taiga in the southeast of Western Siberia, *Eurasian Soil Sci.*,  
817 48(4), 359–372, DOI: 10.1134/S1064229315040067, 2015.

818 Manasyrov, R. M., Pokrovsky, O. S., Kirpotin, S. N., and Shirokova, L. S.: Thermokarst lake waters  
819 across the permafrost zones of western Siberia, *Cryosphere*, 8, 1177–1193, doi:10.5194/tc-8-  
820 1177-2014, 2014.

821 Manasyrov, R. M., Vorobyev, S. N., Loiko, S. V., Kritzkov, I. V., Shirokova, L. S., Shevchenko, V. P.,  
822 Kirpotin, S. N., Kulizhsky, S. P., Kolesnichenko, L. G., Zemtsov, V. A., Sinkinov, V. V., and  
823 Pokrovsky, O. S.: Seasonal dynamics of organic carbon and metals in thermokarst lakes from the  
824 discontinuous permafrost zone of western Siberia, *Biogeosciences*, 12, 3009–3028,  
825 doi:10.5194/bg-12-3009-2015, 2015.

826 Mavromatis, V., Prokushkin, A. S., Pokrovsky, O. S., Viers, J., and Korets, M. A.: Magnesium isotopes  
827 in permafrost-dominated Central Siberian larch forest watersheds, *Geochim. Cosmochim. Ac.*,  
828 147, 76–89, 2014.

829 McClelland, J. W., Déry, S. J., Peterson, B. J., Holmes, R. M., and Wood, E. F.: A pan-Arctic evaluation  
830 of changes in river discharge during the latter half of the 20<sup>th</sup> century, *Geophys. Res. Lett.*, 33,  
831 L06715, 2006.

832 Mergelov, N. and Targulian, V.: Accumulation of organic matter in the mineral layers of permafrost-  
833 affected soils of coastal lowlands in East Siberia, *Eurasian Soil Sci.*, 44, 249–260, 2011.

834 Methodical recommendations for determination of hydrological parameters under insufficient number  
835 of observations, Rota-print GNS AANII, St. Petersburg, 66 pp., 2007 (in Russian).

836 Michel, F. A. and Vaneverdingen, R. O.: Changes in hydrologic regimes in permafrost regions due to  
837 climate change, *Permafrost Periglac. Proc.*, 5, 191–195, 1994.

838 Millot, R., Gaillardet, J., Dupré, B., and Allègre, C. J.: Northern latitude chemical weathering rates:  
839 Clues from the Mackenzie River Basin, Canada, *Geochim. Cosmochim. Ac.*, 67, 1305–1329,  
840 2003.

841 Moran, S. B. and Woods, W. L.: Cd, Cr, Cu, Ni and Pb in the water column and sediments of the Ob-  
842 Irtysh Rivers, Russia, *Mar. Pollut. Bull.*, 35, 270–279, 1997.

843 Nikanorov, A. M., Bryzgalo, V. A., Kosmenko, L. S., and Reshetnyak, O. S.: The role of chemical  
844 river runoff in the anthropogenic transformation of the state of the aquatic environment in the  
845 Enisei mouth area, *Water Resour.*, 37, 471–480, 2010a.

846 Nikanorov, A. M., Smirnov, M. P., and Klimenko, O. A.: Long-term trends in total and  
847 anthropogenic discharge of organic and biogenic substances by Russian rivers into the Arctic  
848 and Pacific Seas, *Water Resour.*, 37, 361–371, 2010b.

849 Nikitin, S. P. and Zemtsov, V. A.: The variability of hydrological parameters of western Siberia,  
850 Nauka, Novosibirsk, 204 pp., 1986 (in Russian).

851 Novikov, S. M., Moskvina, Y. P., Trofimov, S. A., Usova, L. I., Batuev, V. I., Tumanovskaya, S. M.,  
852 Smirnova, V. P., Markov, M. L., Korotkevich, A. E., and Potapova, T. M.: Hydrology of bog  
853 territories of the permafrost zone of western Siberia, *BBM publ. House*, St. Petersburg, 535  
854 pp., 2009 (in Russian).

855 Olefeldt, D., and Roulet, N. T.: Effects of permafrost and hydrology on the composition and  
856 transport of dissolved organic carbon in a subarctic peatland complex, *J. Geophys. Res.*, 117,  
857 G01005, doi:10.1029/2011JG001819, 2012.

858 Olefeldt, D., Persson, A., and Turetsky, M. R.: Influence of the permafrost boundary on dissolved  
859 organic matter characteristics in rivers within the Boreal and Taiga plains of western Canada,  
860 *Environ. Res. Lett.*, 9, Art No 035005, doi:10.1088/1748-9326/9/3/035005, 2014.

861 Oliva, P., Viers, J., and Dupré, B.: Chemical weathering in granitic environments, *Chem. Geol.*, 202,  
862 225–256, 2003.

863 Oosterwoud, M. R., Temminghoff, E. J. M., and van der Zee, S. E. A. T. M.: Quantification of DOC  
864 concentrations in relation with soil properties of soils in tundra and taiga of Northern European  
865 Russia, *Biogeosciences Discuss.*, 7, 3189–3226. 2010.

866 Pavlov, A. V. and Moskalenko, N. G.: The thermal regime of soils in the north of Western Siberia,  
867 *Permafrost Periglac.*, 13, 43–51, doi:10.1002/ppp.409, 2002.



868 Peterson, B. J., Holmes, R. M., McClelland, J. W., Vorosmarty, C. J., Lammers, R. B., Shiklomanov,  
869 A. I., Shiklomanov, I. A., and Rahmstorf, S.: Increasing river discharge to the Arctic Ocean,  
870 *Science*, 298, 2171–2173, 2002.

871 Petrone, K. C., Jones, J. B., Hinzman, L. D., and Boone, R. D.: Seasonal export of carbon, nitrogen,  
872 and major solutes from Alaskan catchments with discontinuous permafrost, *J. Geophys. Res.*,  
873 111, G02020, doi:10.1029/2005JG000055, 2006.

874 Pokrovsky, O. S., Schott, J., Kudryavtzev, D. I., and Dupré, B.: Basalts weathering in Central Siberia  
875 under permafrost conditions, *Geochim. Cosmochim. Ac.*, 69, 5659–5680, 2005.

876 Pokrovsky, O. S., Schott, J., and Dupre, B.: Trace element fractionation and transport in boreal rivers  
877 and soil porewaters of permafrost-dominated basaltic terrain in Central Siberia, *Geochim.*  
878 *Cosmochim. Ac.*, 70, 3239–3260, 2006.

879 Pokrovsky, O. S., Viers, J., Shirokova, L. S., Shevchenko, V. P., Filipov, A. S., and Dupré, B.:  
880 Dissolved, suspended, and colloidal fluxes of organic carbon, major and trace elements in  
881 Severnaya Dvina River and its tributary, *Chem. Geol.*, 273, 136–149, 2010.

882 Pokrovsky, O. S., Viers, J., Dupré, B., Chabaux, F., Gaillardet, J., Audry, S., Prokushkin, A. S.,  
883 Shirokova, L. S., Kirpotin, S. N., Lapitsky, S. A., and Shevchenko, V. P.: Biogeochemistry of  
884 carbon, major and trace elements in watersheds of Northern Eurasia drained to the Arctic Ocean:  
885 The change of fluxes, sources and mechanisms under the climate warming prospective, *C.R.*  
886 *Geosci.*, 344, 663–677, 2012.

887 Pokrovsky, O. S., Reynolds, B. C., Prokushkin, A. S., Schott, J., and Viers, J.: Silicon isotope variations  
888 in Central Siberian rivers during basalt weathering in permafrost-dominated larch forests, *Chem.*  
889 *Geol.*, 355, 103–116, 2013a.

890 Pokrovsky, O. S., Shirokova, L. S., Kirpotin, S. N., Kulizhsky, S. P., and Vorobiev, S. N.: Impact of  
891 western Siberia heat wave 2012 on greenhouse gases and trace metal concentration in thaw lakes  
892 of discontinuous permafrost zone, *Biogeosciences*, 10, 5349–5365, 2013b.

893 Porcal, P., Koprivnjak, J.-F., Molot, L. A., and Dillon, P. J.: Humic substaces – part 7: the  
894 biogeochemistry of dissolved organic carbon and its interactions with climate change, *Environ.*  
895 *Sci. Pollut. R.*, 16, 714–726, 2009.

896 Prokushkin, A. S., Pokrovsky, O. S., Shirokova, L. S., Korets, M. A., Viers, J., Prokushkin, S. G.,  
897 Amon, R., Guggenberger, G., and McDowell, W. H.: Sources and export fluxes of dissolved  
898 carbon in rivers draining larch-dominated basins of the Central Siberian Plateau, *Environ. Res.*  
899 *Let.*, 6, 045212, doi:10.1088/1748-9326/6/4/045212, 2011.

900 Rember, R. D. and Trefry, J. H.: Increased concentrations of dissolved trace metals and organic  
901 carbon during snowmelt in rivers of the Alaskan Arctic, *Geochim. Cosmochim. Ac.*, 68, 477–  
902 489, 2004.

903 Rinta, P., Bastviken, D., van Hardenbroek, M., Kankaala, P., Leuenberger, M., Schilder, J., Stötter,  
904 T., and Heiri, O.: An inter-regional assessment of concentrations and  $\delta^{13}\text{C}$  values of methane  
905 and dissolved inorganic carbon in small European lakes, *Aquat Sci.*, doi:10.1007/s00027-015-  
906 0410-y, 2015.

907 Romanovsky, N.N. : *Underground waters of the cryolithozone*, Moscow, Moscow State University  
908 Publ. House, 232 pp, 1983.

909 Romanovsky, V.E., Drozdov, D.S. Oberman, N.G., Malkova G.V., Kholodov A.L., Marchenko, S.S.,  
910 Moskalenko, N.G., Sergeev D.O., Ukraintseva, N.G., Abramov A.A., Gilichinsky, D.A., and  
911 Vasiliev, A.A.: Thermal State of Permafrost in Russia, *Permafrost Periglacial Proc.*, 21, 136-  
912 155, 2010.

913 Serreze, M. C., Walsh, J. E., Chapin, E., Osterkamp, T., Dyugero, M., Romanovsky, V., Oechel, W.  
914 C., Morison, J., Zhang, T., and Barry, R. G.: Observation evidence of recent change in the  
915 northern high-latitude environment, *Climatic Change*, 46, 159–207, 2000.

916 Shirokova, L. S., Pokrovsky, O. S., Kirpotin, S. N., Desmukh, C., Pokrovsky, B. G., Audry, S., and  
917 Viers, J.: Biogeochemistry of organic carbon,  $\text{CO}_2$ ,  $\text{CH}_4$ , and trace elements in thermokarst  
918 water bodies in discontinuous permafrost zones of Western Siberia, *Biogeochemistry*, 113,  
919 573–593, 2013.

920 Smith, L. C., Macdonald, G. M., Velichko, A. A., Beilman, D. W., Borisova, O. K., Frey, K. E.,  
921 Kremenetsky, K. V., and Sheng, Y.: Siberian peatlands as a net carbon sink and global methane  
922 source since the early Holocene, *Science*, 303, 353–356, 2004.

923 Smedberg, E., Mörth, C. M., Swaney, D. P., and Humborg, C.: Modeling hydrology and silicon-  
924 carbon interactions in taiga and tundra biomes from a landscape perspective: Implications for  
925 global warming feedback, *Global Biogeochem. Cy.*, 20, GB2014, doi:10.1029/2005GB002567,  
926 2006.

927 Stendel, M. and Christensen, J. H.: Impact of global warming on permafrost conditions in a coupled  
928 GCM, *Geophys. Res. Lett.*, 29, 10-1–10-4, 2002.

929 Stepanova, V. M., Pokrovsky, O. S., Viers, J., Mironycheva-Tokareva, N. P. Kosykh, N. P., and  
930 Vishnyakova, E. K.: Major and trace elements in peat profiles in Western Siberia: impact of  
931 the landscape context, latitude and permafrost coverage, *Appl. Geochem.*, 53, 53–70, 2015.

932 Striegl, R. B., Kortelainen, P., Chanton, J. P., Wickland, K. P., Bugna, G. C., and Rantakari, M.:  
933 Carbon dioxide partial pressure and  $^{13}\text{C}$  content of north temperate and boreal lakes at spring  
934 ice melt, *Limnol. Oceanogr.*, 46, 941–945, 2001.

935 Striegl, R. G., Aiken, G. R., Dornblaser, M. M., Raymond, P. A., and Wickland, K. P.: A decrease in  
936 discharge-normalized DOC export by the Yukon River during summer through autumn,  
937 *Geophys. Res. Lett.*, 32, L21413, doi:10.1029/2005GL024413, 2005.

938 Sturm, M., Racine, C., and Tape, K.: Increasing shrub abundance in the Arctic, *Nature*, 411, 546–  
939 547, 2001.

940 Svod pravil SP 33-101-2003. Determination of basic calculated hydrological parameters. Official  
941 release, Gosstroy, Moscow, Russia, 73 pp., 2004.

942 Tank, S. E., Raymond, P. A., Striegl, R. G., McClelland, J. W., Holmes, R. M., Fiske, G. J., and  
943 Peterson, B. J.: A land-to-ocean perspective on the magnitude, source and implication of DIC  
944 flux from major Arctic rivers to the Arctic Ocean, *Global Biogeochem. Cy.*, 26, GB4018,  
945 doi:10.1029/2011GB004192, 2012.

946 Tank, S. E., Frey, K. E., Striegl, R. G., Raymond, P. A., Holmes, R. M., McClelland, J. W., and  
947 Peterson, B. J.: Landscape-level controls on dissolved carbon flux from diverse catchments of  
948 the circumboreal, *Global Biogeochem. Cy.*, 26, GB0E02, doi:10.1029/2012GB004299, 2012.

949 Tape, K., Sturm, M., and Racine, C.: The evidence for shrub expansion in Northern Alaska and the  
950 Pan-Arctic, *Glob. Change Biol.*, 12, 686–702, doi:10.1111/j.1365-2486.2006.01128.x., 2006.

951 Targulian, V. O.: Soil Formation and Weathering in Cold Humid Regions (On Massive-Crystalline  
952 and Sandy Polymictic Rocks), Nauka, Moscow, 1971 (in Russian).

953 Tyrtikov, A. P.: Thawing of soils in tundra of western Siberia, in: Popov A.I. (Ed.) Natural  
954 environment of western Siberia, Issue 3, Izd-vo MG, Moscow, 160–169, 1973 (in Russian).

955 Tyrtikov, A. P.: Dynamics of vegetation coverage and permafrost development in western Siberia,  
956 Moscow, Nauka, 114 pp, 1979 (in Russian).

957 van Hees, P.A.W., Jones, D.L., Finlay, R., Godbold, D.L., Lundstrom, U.S.: 2005. The carbon we do  
958 not see—the impact of low molecular weight compounds on carbon dynamics and respiration  
959 in forest soils: a review, *Soil Biol Biochem.*, 37, 1–13, 2005.

960 Vasiliev, A. A., Streletskaia, I. D., Shirokov, R. S., and Oblogov, G. E.: Evolution of cryolithozone  
961 of coastal zone of western Yamal during climate change, *Kriosfera Zemli*, 2, 56–64, 2011 (in  
962 Russian).

963 Vorobyev, S. N., Pokrovsky, O. S., Kirpotin, S. N., Kolesnichenko, L. G., Shirokova, L. S.,  
964 Manasyrov, R. M.: Flood zone biogeochemistry of the Ob' River middle course, *Applied*  
965 *Geochem.*, 63, 133-145, 2015.

- 966 Voronkov, P. P., Sokolova, O. K., Zubareva, V. I., and Naidenova, V. I.: Hydrochemical features of  
967 local discharge during spring flood from the soil coverage of European territory of the USSR,  
968 Trudy GGI (Proceedings of State Hydrological Institute), 137, 3–57, 1966 (in Russian).
- 969 Voss, B. M., Peucker-Ehrenbrink, B., Eglinton, T. I., Spencer, R. G. M., Bulygina, E., Galy, V.,  
970 Lamborg, C. H., Ganguli, P. M., Montluçon, D. B., Marsh, S., Gillies, S. L., Fanslau, J., Epp,  
971 A., and Luymes, R.: Seasonal hydrology drives rapid shifts in the flux and composition of  
972 dissolved and particulate organic carbon and major and trace ions in the Fraser River, Canada,  
973 *Biogeosciences*, 12, 5597–5618, doi:10.5194/bg-12-5597-2015, 2015.
- 974 Waldron, S., Scott, M. E., and Soulsby, C.: Stable isotope analysis reveals lower-order river  
975 dissolved inorganic carbon pools are highly dynamic, *Environ. Sci. Technol.*, 41, 6156–6162,  
976 doi:10.1021/es0706089, 2007.
- 977 Walvoord, M. A., Voss, C. I., and Wellman, T. P.: Influence of permafrost distribution on  
978 groundwater flow in the context of climate-driven permafrost thaw: Example from Yukon Flats  
979 Basin, Alaska, United States, *Water Res. Research*, 48, W07524, doi:  
980 10.1029/2011WR011595, 2012.
- 981 Walvoord, M. A. and Striegl, R. G.: Increased groundwater to stream discharge from permafrost  
982 thawing in the Yukon River basin: Potential impacts on lateral export of carbon and nitrogen, *J.*  
983 *Geophys. Res.*, 34, L12402, doi:10.1029/2007GL030216, 2007.
- 984 White, A. F. and Blum, A. E.: Effects of climate on chemical weathering in watersheds, *Geochim.*  
985 *Cosmochim. Ac.*, 59, 1729–1747, 1995.
- 986 White, D., Hinzman, L., Alessa, L., Cassano, J., Chambers, M., Falkner, K., Francis, J., Gutowski Jr.,  
987 W. J., Holland, M., Holmes, R. M., Huntington, H., Kane, D., Kliskey, A., Lee, C.,  
988 McClelland, J., Peterson, B., Rupp, T. S., Straneo, F., Steele, M., Woodgate, R., Yang, D.,  
989 Yoshikawa, K., and Zhang T.: The arctic freshwater system: Changes and impacts, *J. Geophys.*  
990 *Res.*, 112, G04S54, doi:10.1029/2006JG000353, 2007.
- 991 Wickland, K. P., Neff, J. C., Aiken, G. R.: Dissolved organic carbon in Alaskan boreal forest:  
992 sources, chemical characteristics, and biodegradability, *Ecosystems*, doi: 10.1007/s10021-  
993 007-9101-4, 2007.
- 994 Worrall, F., Harriman, R., Evans, C. D., Watts, C. D., Adamson, J., Neal, C., Tipping, E., Burt, T.,  
995 Grieve, I., Monteith, D., Naden, P. S., Nisbet, T., Reynolds, B., and Stevens, P.: Trends in  
996 dissolved organic carbon in UK rivers and lakes, *Biogeochemistry*, 70, 369–402, 2004.
- 997 Yang, D., Ye, B., and Shiklomanov, A.: Discharge characteristics and changes over the Ob River  
998 watershed in Siberia, *J. Hydrometeorol.*, 5, 595–610, 2004.

999 Ye, B., Yang, D., Zhang, Z., and Kane, D. L.: Variation of hydrological regime with permafrost  
1000 coverage over Lena basin in Siberia, *J. Geophys. Res.*, 114, D07102, 2009.

1001 Yeghicheyan, D., Bossy, C., Bouhnik Le Coz, M., Douchet, Ch., Granier, G., Heimbürger, A.,  
1002 Lacan, F., Lanzanova, A., Rousseau, T. C. C., Seidel, J.-L., Tharaud, M., Candaudap, F.,  
1003 Chmeleff, J., Cloquet, C., Delpoux, S., Labatut, M., Losno, R., Pradoux, C., Sivry, Y., and  
1004 Sonke, J. E.: A Compilation of Silicon, Rare Earth Element and Twenty-One other Trace  
1005 Element Concentrations in the Natural River Water Reference Material SLRS-5 (NRC-CNRC),  
1006 *Geostand. Geoanal. Res.*, 37, 449–467, doi:10.1111/j.1751-908X.2013.00232.x, 2013.

1007 Zakharova, E. A., Pokrovsky, O. S., Dupré, B., and Zaslavskaya, M. B.: Chemical weathering of  
1008 silicate rocks in Aldan Shield and Baikal Uplift: insights from long-term seasonal  
1009 measurements of solute fluxes in rivers, *Chem. Geol.*, 214, 223–248, 2005.

1010 Zakharova, E. A., Pokrovsky, O. S., Dupré, B., Gaillardet, J., and Efimova, L.: Chemical weathering  
1011 of silicate rocks in Karelia region and Kola peninsula, NW Russia: Assessing the effect of rock  
1012 composition, wetlands and vegetation, *Chem. Geol.*, 242, 255–277, 2007.

1013 Zakharova, E. A., Kouraev, A. V., Rémy, F., Zemtsov, V. A., and Kirpotin, S.N.: Seasonal  
1014 variability of the Western Siberia wetlands from satellite radar altimetry, *J Hydrol.*, 512, 366-  
1015 378, 2014. DOI: <http://dx.doi.org/10.1016/j.jhydrol.2014.03.002>

1016  
1017  
1018  
1019

Table 1. List of sampled rivers, their watershed area and annual runoff. The codes under the months identify the sampling sites listed in Table S1. The annual runoff was calculated following the approach of Frey et al (2007b) as explained in Supplement 1.

Number on the map	Season				N	E	River	watersheds, km <sup>2</sup>	Annual runoff, mm/y
	June	August	October	February					
1	<b>RJ-1</b>		<b>R-1</b>	<b>RF1</b>	56°31'48"	84°09'44"	Ob'	423100	207
2	<b>RJ-3</b>		<b>R-3</b>	<b>RF2</b>	56°46'19.5"	83°57'35.7"	Prud	61.5	44.8
3	<b>RJ-2</b>		<b>R-2</b>		56°43'15.0"	83°55'35.1"	Chybyr'	8.14	44.8
4	<b>RJ-4; R-10</b>		<b>R-4</b>	<b>RF3</b>	57°06'39.2"	83°54'41.1"	Shegarka	12000	58.3
5	<b>RJ-5</b>		<b>R-5</b>	<b>RF4</b>	57°19'20.7"	83°55'53.8"	Brovka	320	63.4
6		<b>BL-3</b>			56°54'39,1"	82°33'33,3"	Cherniy Klyuch	32	168
7		<b>BL-2</b>			57°02'23,75"	82°04'02,44"	Bakchar	3197	96.1
8	<b>RJ-6</b>			<b>RF5</b>	57°36'43.3"	83°37'02.1"	Malyi Tatosh	302	63.4
9	<b>RJ-7</b>		<b>R-7</b>	<b>RF6</b>	57°37'17.3"	83°31'53.3"	Bolshoy Tatosh	1020	74.6
10	<b>RJ-8</b>		<b>R-8</b>	<b>RF7</b>	57°52'26.8"	83°11'29.9"	Chemondaevka	177	63.4
11	<b>RJ-9</b>		<b>R-9</b>	<b>RF8</b>	57°58'45.7"	82°58'32.2"	Sugotka	275	63.4
12	<b>RJ-10</b>	<b>RA-23</b>		<b>RF9</b>	58°04'20.8"	82°49'19.7"	Chaya	27200	95.6
13	<b>RJ-11</b>			<b>RF10</b>	58°23'16.8"	82°11'39.0"	Tatarkin Istok	58.6	33.4
14	<b>RJ-12</b>		<b>R-12</b>		58°24'38.0"	82°08'46.0"	Istok	12.3	127
15	<b>RJ-13</b>	<b>RA-22</b>	<b>R-13</b>	<b>RF11</b>	58°26'06.9"	82°05'43.6"	Shudelka	3460	211
16	<b>RJ-14</b>		<b>R-14</b>	<b>RF12</b>	58°33'03.1"	81°48'44.3"	Chigas	689	180
17		<b>BL-9</b>			58°32'05,8"	80°51'26,8"	Karza	473	148
18		<b>BL-6</b>			58°37'29,9"	81°06'09,0"	Sochiga	510	148
19	<b>RJ-15</b>	<b>RA-21; BL-4</b>	<b>R-17</b>	<b>RF64</b>	58°42'34.5"	81°22'22.0"	Parabel	25500	131
20	<b>RJ-58</b>	<b>BL-5</b>	<b>R-15</b>	<b>RF65</b>	58°40'46.5"	84°27'56.6"	Vyalovka	117	127
21				<b>RF63</b>	58°59'37"	80°34'00"	Vasyugan	63780	177
22				<b>RF62</b>	59°41'01,6"	77°44'33,9"	Kornilovskaya	190	133
23				<b>RF61</b>	59°44'09,2"	77°26'06"	Levyi Il'yas	253	133
24				<b>RF60</b>	60°08'43"	77°16'53"	Koltogorka	220	155.4
25				<b>RF58</b>	60°30'19"	76°58'57"	Sosninskii Yegan	732	199

Number on the map	Season				N	E	River	watersheds, km <sup>2</sup>	Annual runoff, mm/y
	June	August	October	February					
26	<b>RJ-16</b>	<b>BL-36</b>			60°40'28.8"	77°31'29.4"	Ob'	773200	216
27		<b>BL-35</b>			60°44'10,9"	77°22'55,9"	Medvedka	7	173
28		<b>BL-34</b>			60°45'58,5"	77°26'12,6"	Saim	26	173
29		<b>BL-33</b>			60°47'29,3"	77°19'13,5"	Mishkin Saim	32	173
30		<b>BL-32</b>			60°49'32,3"	77°13'46,3"	Alenkin Egan	44	173
31		<b>BL-31</b>			60°50'43,6"	77°05'03,0"	Kaima	31	173
32		<b>BL-30</b>			60°55'41,0"	76°53'49,3"	Vakh	75090	298
33	<b>RJ-23</b>			<b>RF53</b>	61°34'27.4"	77°46'35.4"	Mokhovaya	1260	192.3
34	<b>RJ-17</b>	<b>BL-29; RA-20</b>		<b>RF57</b>	61°11'52.7"	75°25'20.2"	Vatinsky Egan	3190	287
35		<b>BL-28</b>			61°12'19,5"	75°23'06,5"	Er-Yakh	9.35	173
36	<b>RJ-18</b>	<b>RA-19; BL-27</b>			61°19'41.2"	75°04'0.3"	Ur'evskii Egan	359	272
37	<b>RJ-19; R-9</b>	<b>BL-26</b>		<b>RF56</b>	61°26'13,6"	74°47'39,7"	Agan	27600	291
38	<b>RJ-20</b>	<b>RA-18</b>		<b>RF55</b>	61°27'17.3"	74°40'23.3"	Kottym'egan	7.18	192
40	<b>RJ-21</b>			<b>RF54</b>	61°29'46.6"	74°15'30.3"	Segut-Yagun	3.37	192
41	<b>RJ-22</b>			<b>RF13</b>	61°29'11.1"	74°09'42.9"	Vach-Yagun	1.79	192
42	<b>RJ-24</b>			<b>RF52</b>	61°50'28.6"	70°50'28.2"	Vachinguriyagun	9.52	192
43	<b>RJ-25</b>			<b>RF14</b>	61°58'05.1"	73°47'03.4"	Lyukh-Yagun	21.6	192
44				<b>RF51</b>	61°59'39"	73°47'39"	Limpas	1648	320
45	<b>RJ-26; R-7; R-8</b>	<b>RA-17</b>		<b>RF50</b>	62°07'50,0"	73°44'05,6"	Tromyegan	10770	263
46	<b>RJ-57</b>			<b>RF49</b>	62°33'39,8"	74°00'29,5"	Pintyr'yagun	33.5	192
47	<b>RJ-56</b>	<b>BL-25</b>		<b>RF48</b>	62°37'08.4"	74°10'15.9"	Petriyagun	9.65	192
48	<b>RJ-54; R-6</b>	<b>BL-24</b>		<b>RF47</b>	63°38'23,4"	74°10'52"	Kirill-Vys'yagun	598	225
49	<b>RJ-55</b>	<b>BL-23</b>		<b>RF46</b>	62°43'09.9"	74°13'45.9"	Ai-Kirill-Vys'yagun	24.0	192
50		<b>BL-22</b>		<b>RF45</b>	63°11'19,3"	74°36'25,5"	Pyrya-Yakha	82	194
51		<b>RA-14</b>			63°11'40.68"	74°38'16.92"	Itu-Yakha	250	194

Number on the map	Season				N	E	River	watersheds, km <sup>2</sup>	Annual runoff, mm/y
	June	August	October	February					
52		RA-13			63°10'3.48"	74°45'16.32"	Nekhtyn-Pryn	96	194
53		RA-4			63°10'4.68"	76°28'19.08"	Nyudya-Pidya-Yakha	79.5	194
54		RA-12			63°9'31.38"	75°3'2.58"	Ponto-Yakha	19	194
55		RA-11			63°9'39.84"	75°09'10.86"	Velykh-Pelykh-Yakha	170	194
56		RA-10			63°13'12.06"	75°38'52.26"	Yangayakha	88	194
57		RA-9			63°13'25.2"	76°5'23.04"	Tlyatsayakha	43	194
58		RA-8			63°13'3.66"	76°15'24.6"	Chukusamal	121	194
59		RA-3; RA-7			63°46'22.92"	76°25'28.86"	Vyngapur	1979	324
60		RA-6			63°12'43.38"	76°21'27.66"	Goensapur	11	194
61		RA-5			63°12'45.96"	76°24'1.32"	Denna	15	194
62		RA-15			63°8'34.02"	74°54'29.1"	Nyudya-Itu-Yakha	32	194
63	RJ-53; R-5	RA-16; BL-21		RF38	63°22'01.6"	74°31'53.2"	Kamgayakha	175	194
64	RJ-52; R-4	BL-19		RF39	63°36'48.2"	74°35'28.6"	Khatytayakha	34.6	194
65	RJ-51	BL-18		RF40	63°40'41.8"	74°35'20.7"	Pulpuyakha	281	194
66	RJ-50; R-3	BL-17		RF41	63°49'58.0"	74°39'02.5"	Khanupiyakha	74	194
67	RJ-29; R-2	BL-16		RF42; RF37	63°51'23.4"	75°08'05.6"	Kharucheiyakha	820	292
68	R-1; Z-55; RJ-28	BL-20; RA-2; BL-15		RF43	63°49'54.2"	75°22'47.1"	Pyakupur	9880	324
69	RJ-27; Z-86	BL-14; RA-1		RF44	63°47'04.5"	75°37'06.8"	Lymbyd'yakha	115	194
70		BL-13			63°43'37.9"	75°59'04.1"	Chuchi-Yakha	1396	292
71	RJ-32				64°12'08.4"	75°24'28.4"	Ngarka-Tyde-Yakha	59.9	186
72	RJ-30			RF36	64°06'50.7"	75°14'17.3"	Ngarka-Varka-Yakha	67.1	186
73	RJ-31				64°09'06.4"	75°22'18.1"	Apoku-Yakha	18.8	186
74	RJ-33	RY 14-49		RF35	64°17'31.9"	75°44'33.4"	Etu-Yakha	71.6	186
75	RJ-34				64°19'10.1"	76°08'26.7"	Varka-Yakha	105	186



Number on the map	Season				N	E	River	watersheds, km <sup>2</sup>	Annual runoff, mm/y
	June	August	October	February					
76		<b>RY 14-48</b>			64°23'30,6"	76°19'50,1"	Khaloku-Yakha	53	186
77	<b>RJ-35</b>			<b>RF34</b>	64°26'05.2"	76°24'37.0"	Kharv'-Yakha	46.4	186
78	<b>RJ-36</b>	<b>RY 14-47</b>		<b>RF33</b>	64°32'07.9"	76°54'21.3"	Seryareyakha	15.2	186
79	<b>RJ-37</b>	<b>RY 14-46</b>		<b>RF32</b>	64°40'14.0"	77°05'27.2"	Purpe	5110	309
80	<b>RJ-38</b>				64°55'55.1"	77°56'08.2 "	Aivasedapur	26100	309
81	<b>RJ-39</b>			<b>RF31</b>	65°06'48.8"	77°47'58.8"	Tydylyakha	7.46	185
82	<b>RJ-40</b>	<b>RY 14-45</b>		<b>RF30</b>	65°12'17.6"	77°43'49.8"	Tydyotta	12.0	309
83	<b>RJ-41</b>	<b>RY 14-44</b>		<b>RF29</b>	65°23'34.1"	77°45'46.7"	Ponie-Yakha	78.9	185
84	<b>RJ-42</b>	<b>RY 14-43</b>		<b>RF28</b>	65°41'51.1"	78°01'05.0"	Yamsovey	4030	309
85		<b>RY 14-42</b>			65°46'34,5"	78°08'25,8"	Khiroyakha	183	185
86	<b>RJ-43</b>			<b>RF27</b>	65°47'48.6"	78°10'09.0"	Almayakha	106	185
87	<b>RJ-45</b>			<b>RF25</b>	65°58'54"	77°34'05"	Yude-Yakha	42.4	185
88	<b>RJ-46</b>			<b>RF26</b>	65°59'05.7"	77°40'52.6"	Tadym-Yakha	39.9	185
89	<b>RJ-44</b>	<b>RY 14-41</b>			65°57'05.5"	78°18'59.1"	Pur	112000	298
90	<b>RJ-49</b>	<b>RT2 14-32</b>			65°59'14.7"	78°32'25.2"	Malaya Khadyr-Yakha	512	278
91	<b>RJ-48</b>	<b>RT2 14-31</b>			66°17'10.8"	79°15'06.1"	Ngarka Khadyta-Yakha	1970	277
92		<b>RT2 14-30</b>			66°59'20,9"	79°22'30,5"	Malokha Yakha	157	208
93		<b>RT2 14-29</b>			67°10'54,8"	78°51'04,5"	Nuny-Yakha	656	312
94	<b>RJ-47</b>	<b>RT2 14-40</b>			67°22'13.28"	79°00'25,9"	Taz	150000	330
95				<b>RF21</b>	67°24'39"	76°21'12"	Khadutte	5190	346




Fig.1. Map of the study site with permafrost boundaries (Brown et al., 2002; <http://portal.inter-map.com> (NSIDC)), runoff contour lines (Nikitin and Zemtsov, 1986) and sampling points along the latitudinal transect of rivers Ob, Pur and Taz basin. The numbers of the sampling sites are listed in Table 1.












Fig. 1, continued. Detailed map of the four rectangles in Fig 1 a.




**Figure 2.** Decrease of river water pH with the increase of the latitude during winter (A), spring (B) and summer (C). The spring acid pulse is seen only in permafrost-affected rivers north of 62°N (B), and the scatter of the values is maximal during summer (C). The variability among different watershed size is smaller than that between the seasons and within the latitude gradient. Diamonds, squares, triangles and circles represent watershed of size < 100 km<sup>2</sup>, 100 to 1000 km<sup>2</sup>, 1000 to 10,000 km<sup>2</sup>, and > 10,000 km<sup>2</sup>, respectively.



**Figure 3.** Decrease of DOC with latitude during winter (A), spring (B) and summer (C). The latitudinal trend is significant at  $p < 0.05$ . Considering all seasons together, the differences between different watershed sizes are not statistically significant ( $p > 0.05$ ). The symbols are the same as in Fig. 2.




**Figure 4.** Significant decrease of DIC with latitude during winter (A), spring (B) and summer (C). Note the logarithmic scale on concentration in all three plots. The symbols represent different size of the watershed, see Fig. 2.



**Figure 5.** The variation of  $\delta^{13}\text{C}_{\text{DIC}}$  with latitude during winter (A) and spring (B) for watershed of different size. The symbols are the same as in Fig. 2. Isotopically-light DIC is observed in permafrost-affected zone during spring, suggesting intensive respiration of soil or plant litter carbon (Ob river sediments are from -25 to -27‰, Guo et al., 2004a).



Forest watershed, 57°N, August  
Dark coniferous taiga




**Fig. 6.** Scheme of DOC pathways within the soil profile and to the river, (A): in forest watershed of the south, permafrost-free zone (57°N). Soil horizons (FAO, 2006): 1, O (Mor, forest litter); 2, Medium-decomposed peat (He) transforming into strongly decomposed peat (Ha) in the bottom layer; 3, Mollic humic horizon (A); 4, ABg surface horizons with stagnic properties; 5, Bg middle stagnic horizon; 6, Cgk carbonate-bearing clays and clay loam.



# Frozen bog watershed 67°N, August

**B**



**Fig. 6, continued. (B):** DOC pathways in frozen bog peatlands of continuous permafrost (67°N). Soil horizons (FAO, 2006): 7, weakly decomposed peat (Hi); 8, partially decomposed peat (He); 9, humic horizons (AH); 10, cryoturbated frozen stagnic horizon (Bgf); 11, frozen stagnic horizon (BCgf); 12 sedimentary deposits (Cf).

In the south, DOC is retained by clay and deep in the soil profile, by clay loam with carbonates. In the north, the active layer depth does not exceed the overall thickness of the peat and thus the leachate of ground vegetation and peat layer do not meet mineral horizons during their transit to the river.