

Dear author,

By studying comments from reviewers and your responses to them, I have judged that your manuscript will be acceptable after minor revision. Although, as was pointed out by both reviewers, the idea “primary productivity is influenced by freshwater content” is not new, the manuscript provides valuable information on regional and interannual variability in primary productivity in the Chukchi Sea. Based on this view, the title “The potential effects of freshwater content on the primary production in the Chukchi Sea” may be too strong and I would recommend to change it to “Primary production in the Chukchi Sea with potential effects of freshwater content”.

→ As the editor recommended, we changed the title (in lines 1-3, page 1).

Your response to some of reviewer #3's comments are not adequate and need to be reconsidered before the final acceptance. Please see below.

Reviewer #3's comment: “this is the Russian Exclusive Economic Zone(20 miles), not territorial waters, which extend only 12 miles from shore”

Your reply to this comment missed the point. Territorial waters is defined as a belt of coastal waters extending at most 12 miles from the coastline. Is your station really within 12 miles from the Russian coastline? If not, please change “including the territorial waters of the Russian Federation” (line 82) to “including coastal waters of...” or “including the Exclusive Economic Zone of ...”

→ As the editor commented, our stations are including the Russian Exclusive Economic Zone (not territorial waters). Thus, we changed this sentence to “including the Exclusive Economic Zone of the Russian Federation (in line 83, page 6).

Reviewer #3's comment: “These are inventories of nutrients, not concentrations”

I agree with the reviewer #3. Once concentrations of nutrient were integrated, the obtained value is inventory or amount, not concentration any more. Please change “concentration” to “inventory” for values with the unit “mol m⁻²”.

→ As the editor commented, we changed “concentration” to “inventory” (in lines 188-

197, pages 11-12).

Reviewer #3's comment: "I do not follow what mechanism is being invoked for replenishment of nutrients from deep waters (e.g., Canada Basin)."

You have added (e.g., Canada Basin) to the text, but I suppose that what you want to mean here is a replenishment of nutrients below the surface layer, not from "deep waters". I would suggest to modify the sentence to clarify the meeting to read "These inputs of freshwater presumably influenced the nutrient reservoir and its replenishment from deeper layer by altering stratification of the water column."

→ As the editor mentioned, we meant here is a replenishment of nutrients below the surface layer, not from "deep waters". Thus, we changed this sentence based on the editor suggested (in lines 336-337, page 20).

Reviewer #3's comment: "comparisons to other productivity measurements are incomplete without careful consideration of the influence of seasonality and location of sampling—high productivity in the Chukchi Sea is rather localized...."

Difference in sampling dates is discussed in the revised manuscript and clearly presented in Figure 9. However, location of previous studies other than RUSALCA cruises are not indicated. As productivity is largely different between regions, location information is essential to judge Figure 9. Please add a description of locations of previous studies in the text and/or Figure 9.

→ We marked location information in Figure 10 (not Figure 9).

Figure 9. What are open circle and open square with a cross in Figure 9 (a) and (b)? Please mention in the caption.

→ Actually, they should be same closed symbols in each Figure 9 (a) and (b). We think that some open symbols (in Figure 9 (a) and (b)) might be induced by the conversion from PPT file to PDF file. Thus, we input a new Figure 9 (with only closed symbols).

Figure 10, Hill et al. (2005)-> Hill and Cota (2005)?

→ Since "Hill et al. (2005)" in an earlier version of our manuscript was not a proper reference, we corrected "Hill and Cota (2005)" in the revised manuscript.

1 ~~The potential effects of fresh water content on the primary~~

2 ~~production in the Chukchi Sea with potential effects of~~

3 ~~freshwater content~~

4

5 Mi Sun Yun^a, Terry E. Whitledge^b, Dean Stockwell^b, SeungHyun Son^c, Jang Han Lee^a, Jung Woo

6 Park^a, Da Bin Lee^a, Jinku Park^a, Sang H. Lee^{a,*}

7

8

9 ^a Department of Oceanography, Pusan National University, 30, Jangjeon-dong, Geumjeong-gu, Busan

10 609-735, Korea

11 ^b Institute of Marine Science, University of Alaska Fairbanks, Fairbanks, AK 99775-7220, USA

12 ^c CIRA, Colorado State University, Fort Collins, Colorado, USA

13

14

15

16 *Corresponding author. Tel.: +82 51 510 2256; fax: +82 51 581 2963.

17 E-mail address: sanglee@pusan.ac.kr

18

19 **Abstract**

20 The *in situ* primary production rates and various environmental variables were investigated in the

21 Chukchi Sea during the RUSALCA expedition, which was conducted in 2012, to identify the current

22 status of primary production. A ^{13}C - ^{15}N dual tracer technique was used to measure the daily primary

23 production rates, which ranged from 0.02 to 1.61 $\text{g C m}^{-2} \text{ d}^{-1}$ (mean \pm SD = $0.42 \pm 0.52 \text{ g C m}^{-2} \text{ d}^{-1}$). The

24 primary production rates showed large regional differences, with the southern region ($0.66 \pm 0.62 \text{ g C m}^{-2}$

25 d^{-1}) producing approximately five times as much as the northern region ($0.14 \pm 0.10 \text{ g C m}^{-2} \text{ d}^{-1}$), which

26 was primarily due to the differences in phytoplankton biomasses induced by regional nutrient conditions.

27 The primary production rates in the Chukchi Sea were averaged using data acquired during the three

28 different RUSALCA expeditions (2004, 2009, and 2012) as $0.33 \text{ g C m}^{-2} \text{ d}^{-1}$ (SD = $0.40 \text{ g C m}^{-2} \text{ d}^{-1}$),

29 which was significantly lower than previously reported rates. In addition to strong seasonal and

30 interannual variations in primary production, recent decreases in the concentrations of major inorganic

31 nutrients and chlorophyll *a* could be among the reasons for the recent low primary production in the

32 Chukchi Sea because the primary production is mainly affected by nutrient concentration and

33 phytoplankton biomass. The nutrient inventory and primary production appear to be largely influenced by

34 the freshwater content (FWC) variability in the region due to the significant relationships between FWC,

35 nitrate concentrations ($r = 0.54, p < 0.05$) and primary production rates ($r = 0.56, p < 0.05$). Moreover, we

36 found highly significant relationships between the nutrient levels and the primary production rates ($r =$

37 0.75, $p < 0.001$). In conclusion, the primary production in the Chukchi Sea is primarily controlled by
38 nutrient availability which is strongly related to the FWC variability. Our results imply that the predicted
39 increase in freshwater accumulation might cause a decrease in primary production by lowering the
40 nutrient inventory in the euphotic zone of the Chukchi Sea.

41

42 **Keywords:**

43 Phytoplankton, Primary production, Chukchi Sea, Freshwater content, Arctic

44

45

46 **1. Introduction**

47 Over recent years, the Arctic Ocean has undergone drastic changes in the extent and thickness of sea
48 ice (Stroeve et al., 2008; Comiso et al., 2008; Kwok et al., 2009; Overland and Wang, 2013). The
49 continuing loss of sea ice may result in changes to various physical and chemical environmental
50 conditions in the Arctic Ocean. For example, the loss in sea ice cover allows more sunlight to enter the
51 surface layer of the Arctic Ocean, which results in a longer growing season for phytoplankton growth
52 (Arrigo et al., 2008; Ardyna et al., 2014). Stroeve et al. (2014) reported that the arctic melt season has
53 lengthened at a rate of 5 days decade⁻¹ from 1979 to 2013, due to later autumn freeze-up. In accordance
54 with their findings, Ardyna et al. (2014) documented the development of a second bloom in the Arctic
55 Ocean during the fall, which coincides with the delayed freeze-up and the increased exposure to wind
56 stress.

57 However, the loss in sea ice can cause an increase in the input of freshwater (McPhee et al., 2009). In
58 fact, the freshwater volumes in the Canada and Makarov Basins increased by of 8500 km³ in 2008 due to
59 increased sea ice melting and river discharge (McPhee et al., 2009; Rabe et al., 2011). This phenomenon
60 can enhance the stratification in the upper ocean (Yamamoto-Kawai et al., 2009) and consequently reduce
61 vertical mixing, thereby preventing nutrient inputs from deep waters to the euphotic zone. In fact,
62 McLaughlin and Carmack (2010) found a deepening of the nutricline due to the accumulation of surface
63 freshwater in the Canada Basin.

64 In the Chukchi Sea as inflow shelf, there was an increased volume flux of 50% in 2011 (~ 1.1Sv)

65 relative to 2001 (~ 0.7 Sv), which was accompanied by increases in heat and freshwater fluxes (Woodgate

66 et al., 2012). Though the volume flux may vary both seasonally and annually under the influence of the

67 local wind fields, the recent increases in freshwater fluxes in the region may have important implications

68 for phytoplankton in terms of nutrient availability for their growth (Woodgate et al., 2005a, 2005b, 2006).

69 Thus, it is important to identify how phytoplankton respond to these environmental changes in the region

70 in terms of production and/or community structure. According to Li et al. (2009), the phytoplankton

71 community has changed under the freshening and stratifying condition in the Canada Basin. Notably, the

72 abundance of small phytoplankton (< 2 μ m) has increased, whereas the abundance of large phytoplankton

73 (2-20 μ m) has decreased. Yun et al. (2014) also found that compared with previous reports, the small

74 phytoplankton were more abundant on the Chukchi Sea shelf, which is dominated by low nutrients and

75 freshening conditions. Therefore, the changes in recent phytoplankton production under the rapidly

76 changing environmental conditions need to be monitored because the changes in phytoplankton

77 production could have important implications for understanding ecosystem changes in the Arctic Ocean.

78 In order to understand climate and ecosystem change in the Pacific Arctic Ocean which is a region

79 summer sea ice cover was declining dramatically (Crane and Ostrovskiy, 2015), the RUSALCA (Russian-

80 American Long-term Census of the Arctic) expedition, which is a joint US-Russian research program,

81 started from 2004 as multidisciplinary investigations in the Bering and Chukchi Seas. Three RUSALCA

82 expeditions (2004, 2009, and 2012) provided a good opportunity for continuous measurements of the
83 primary production in the entire Chukchi Sea, including the ~~territorial waters~~Exclusive Economic Zone
84 of the Russian Federation. The 2004 RUSALCA expedition was conducted from 8 to 24 August, 2004
85 (Lee et al., 2007). The 2009 RUSALCA expedition was executed from 1 to 30 September, 2009 (Yun et
86 al., 2014). The 2012 RUSALCA expedition was carried out from 27 August to 16 September, 2012. This
87 study is part of the 2012 RUSALCA expedition.

88 In this study, we addressed the regional characteristics of primary production by examining the main
89 driving factors responsible for the regional variability in the Chukchi Sea based on measurements taken in
90 2012. In addition, we investigated the recent trends in primary production in the Chukchi Sea based on
91 the results of the three RUSALCA expeditions (2004, 2009, and 2012) in the Chukchi Sea. Finally, we
92 emphasized the potential effects of freshwater accumulation on the primary production in the Chukchi
93 Sea because changing amounts and distributions of freshwater content could lead to changes in the
94 primary production rates.

95

96 **2. Materials and methods**

97 *2.1. Study area and sampling*

98 The RUSALCA expedition in 2012 was conducted onboard the Russian vessel *Professor Khromov* in
99 the Chukchi Sea from 27 August to 16 September. The study area was comprised of several sections

100 between the Bering Strait and the vicinity of Herald Canyon (Fig. 1). To understand the regional
101 characteristics of primary production, the study area was divided into two geographic regions
102 (northern/southern) following Yun et al. (2014). The northern region consisted of stations in the vicinity
103 of Herald Canyon (CEN and HC sections) (Fig. 1). The stations in the Chukchi South and Cape Lisburne
104 (CS and CL sections) were included in the southern region. Most of the bathymetric depths in the entire
105 study area were quite shallow, with a mean of 55 m (SD = \pm 11 m). Between the production stations, the
106 depth of euphotic zone from the surface to 1% light depth varied between 20 and 46 m, with a mean of 29
107 \pm 10 m (Table 1).

108 Oceanographic/biological samples were taken from a total of 54 conductivity-temperature-depth
109 (CTD) stations. The vertical profiles of water temperature and salinity were obtained using a Sea-Bird
110 model SBE911plus CTD profiler. Water samples were collected with a stainless-steel rosette sampler that
111 was equipped with 21 10-liter bottles at every CTD station. The data from the previous RUSALCA
112 expeditions (in 2004 and 2009) were included to understand the recent trends in primary production in the
113 Chukchi Sea.

114

115 2.2. Physical and chemical variables

116 The stratification index of the water column ($\Delta\sigma_t$) (in kg m^{-3}) was determined as the difference in $\Delta\sigma_t$
117 values between the surface and the bottom depth according to Yun et al. (2014). The surface mixed layer

118 (Z_m) was defined as the depth at which the density (sigma- t) gradient was 0.05 kg m^{-3} higher than the
119 surface density, as in Coupel et al. (2015). The depth of the euphotic zone (Z_{eu}) in this study was defined
120 as the depth receiving 1% of the surface PAR value, as in Lee et al. (2007) and Yun et al. (2014), and was
121 obtained from a Biospherical QSP-2300 PAR sensor (Biospherical Instruments Inc.) that was lowered
122 with the CTD/rosette sampler. The nitracline (Z_{nit}) was determined as the depth at which the nitrate
123 gradient was greater than $0.1 \mu\text{M m}^{-1}$ according to the definition of Coupel et al. (2015).

124

125 *2.3. Fresh Water Content (FWC)*

126 To assess the surface water freshening, the freshwater content (FWC) was calculated following
127 Carmack et al. (2008):

128

$$129 FWC = \int_{Z_{lim}}^0 (1 - S(z)/S_{ref}) dz$$

130 where S and S_{ref} are the *in situ* and reference salinities, respectively, and Z_{lim} is the depth where S equals
131 S_{ref} (34.8 on the practical salinity scale). We used a reference salinity of 34.8 following Aagaard and
132 Carmack (1989) to computing freshwater since it has been considered as the mean salinity for the Arctic
133 Ocean.

134

135 *2.4. Nutrient concentration measurements*

136 The discrete water samples used in measuring the nutrient concentrations were obtained from 5 to 9
137 different depths depending on the water depths. The dissolved inorganic nutrient concentrations
138 (nitrite+nitrate, ammonium, phosphate, and silicate) were analyzed onboard immediately after collection
139 using an automated nutrient analyzer (ALPKEM RFA model 300) following the method of Whitledge et
140 al. (1981).

141

142 *2.5. Chlorophyll a concentration measurements*

143 The water samples used for measuring the chlorophyll *a* concentration were obtained from 4 to 7
144 different depths at most stations. The water samples were filtered through Whatman GF/F filters (24 mm),
145 and the filters were then kept frozen until analysis in the laboratory. The filters were subsequently
146 extracted in a 3:2 mixture of 90% acetone and DMSO in a freezer for 24 h, followed by centrifugation
147 (Shoaf and Lium, 1976). The chlorophyll *a* concentrations were measured using a Turner Designs model
148 10-AU fluorometer, which was calibrated using commercially available preparations of purified
149 chlorophyll *a* (Turner Designs, USA). The methods and calculations used to determine the chlorophyll *a*
150 concentrations followed the procedure of Parsons et al. (1984).

151

152 *2.6. In situ primary production measurements*

153 The water samples used to measure primary production were collected at six photic depths (100, 50,

154 30, 12, 5, and 1% penetration of the surface irradiance, PAR). At 11 selected morning stations, the *in situ*
155 primary productions of phytoplankton were measured using a ^{13}C - ^{15}N dual tracer technique (Lee and
156 Whitlege, 2005; Lee et al., 2007). This method could be useful for distinguish the relative importance of
157 nitrate and ammonium as nitrogen sources for the cell and population (Dugdale and Goering, 1967). We
158 followed the same analytical procedure of Lee et al. (2007) and Yun et al. (2014) to the measure primary
159 production to consistently compare the primary production levels determined in the three studies. Briefly,
160 heavy isotope-enriched (98-99%) carbon ($\text{NaH}^{13}\text{CO}_3$), nitrate (K^{15}NO_3), and ammonium ($^{15}\text{NH}_4\text{Cl}$)
161 substrates were inoculated in polycarbonate bottles (1 L) and then incubated on deck in a large
162 polycarbonate incubator cooled with running surface seawater under natural light conditions. After
163 approximately 4 to 5 h of incubation , all samples were filtered using pre-combusted (450°C, 4 h) glass
164 fiber filters (Whatman GF/F; diameter = 25 mm). After HCl fume treatment, the samples were sent to the
165 Alaska Stable Isotope Laboratory of the University of Alaska, Fairbanks, USA. The abundances of ^{13}C
166 and ^{15}N and the total amounts of particulate organic carbon (POC) and nitrogen (PON) were determined
167 using a Thermo Finnigan Delta+XL mass spectrometer. Finally, the carbon and nitrogen production rates
168 were calculated based on Hama et al. (1983) and Dugdale and Goering (1967), respectively.
169

170 **3. Results**

171 *3.1. Physical conditions*

172 The surface temperature (T_{sur}) varied from -2 to 9 °C in the study area in 2012 (Fig. 2a). The higher
173 temperatures were found in the eastern side of the southern Chukchi Sea due to the strong influence of the
174 Alaskan Coastal Water (warmer and less saline). The freezing temperatures were observed in the vicinity
175 of the Herald Canyon and gradually decreased toward the northward. At the surface, the salinity varied
176 between 21 and 33 psu. The surface salinity (S_{sur}) was considerably lower in the southwestern side
177 compared with the northeastern side of the southern Chukchi Sea (Fig. 2b). The stratification index ($\Delta\sigma_t$)
178 in the study area ranged from 0.7 to 9.7 kg m^{-3} , with a mean of $3.8 \pm 2.2 \text{ kg m}^{-3}$. The stratification in the
179 southern region was higher than in the northern region (Fig. 2c). The general distribution of the
180 stratification index was similar to that of surface salinity because it tended to be high in areas where
181 surface salinity was low. The surface mixed layer (Z_m) was thinner than 15 m over the entire study area
182 (Fig. 2d). In the study area, the depths of nitracline (Z_{nit}) ranged from 2.5 m to 35 m (Fig. 2e), with a
183 mean nitracline depth of $12.8 \pm 7.7 \text{ m}$.

184

185 3.2. Nutrient distribution

186 Since the mean depths of euphotic zone in this study was about 30 m, the distribution of ambient
187 nutrient concentrations integrated from surface to 30 m of the water column is shown in Fig. 3. The
188 ~~concentrations inventory~~ of nitrite+nitrate ranged from 21.51 to 355.43 mmol m^{-2} , whereas the
189 ammonium ~~concentration inventory~~ ranged from 15.36 to 109.51 mmol m^{-2} (Figs. 3a and 3b). High

190 nitrite+nitrate ~~econcentrations-inventory~~ that exceeded 200 mmol m⁻² were observed at the center of the
191 CL section (Fig. 3a). The ~~ambient-concentrations-inventories~~ of these nutrients in the southern region
192 (134.15 ± 98.41 mmol m⁻² for nitrite+nitrate and 61.22 ± 20.55 mmol m⁻² for ammonium, respectively)
193 were approximately two times higher than their ~~econcentrations-inventories~~ in the northern region (75.01 ±
194 52.01 mmol m⁻² for nitrite+nitrate and 40.49 ± 20.69 mmol m⁻² for ammonium) (see Table 2). The
195 ~~concentration-inventory~~ of phosphate in the study area was fairly uniform, with a mean of 24.03 ± 8.30
196 mmol m⁻² (Fig. 3c). The silicate ~~concentration-inventory~~ was generally higher in the southern region than
197 in the northern region (Fig. 3d).

198

199 3.3. Chlorophyll *a* concentration

200 The distribution of the chlorophyll *a* concentration in the upper 30 m (i.e., mean depth of euphotic
201 zone in this study) of the entire study area is shown in Fig. 4. High chlorophyll *a* concentrations of over
202 80 mg m⁻² were observed in the western side of the CL section (from st. CL5 to st. CL8), and low
203 chlorophyll *a* concentrations were shown in the western side of the CS section (Fig. 4). The highest
204 concentration (286.4 mg m⁻²) was obtained at station CL8. Over the entire study area, the mean
205 chlorophyll *a* concentration integrated from the surface to 30 m was 42.7 mg m⁻² (SD = ± 57.4 mg m⁻²).
206 The average concentrations were 21.7 mg m⁻² (SD = ± 19.6 mg m⁻²) and 54.5 mg m⁻² (SD = ± 67.7 mg m⁻²)
207 for the northern and southern regions, respectively.

208

209 *3.4. Primary production rates*

210 Overall, the hourly carbon production rates integrated over the euphotic zone from six light depths

211 ranged from 1.1 to 108.6 mg C m⁻² h⁻¹, with a mean of 27.7 mg C m⁻² h⁻¹ (SD = 34.7 mg C m⁻² h⁻¹). The

212 highest primary production rates were found at station CL8 (108.6 mg C m⁻² h⁻¹) followed by station

213 CL5A (82.1 mg C m⁻² h⁻¹) (Fig. 5). In the northern region, the carbon production rates ranged from 1.1 to

214 18.7 mg C m⁻² h⁻¹, with a mean of 9.0 mg C m⁻² h⁻¹ (SD = ± 6.4 mg C m⁻² h⁻¹). In comparison, the average

215 rates in the southern region were approximately five times higher than the average rates in the northern

216 region (43.3 ± 41.7 mg C m⁻² h⁻¹).

217 The vertically integrated nitrate production rates ranged from 0.14 to 18.77 mg NO₃ m⁻² h⁻¹, with a

218 mean of 2.72 mg N m⁻² h⁻¹ (SD = ± 5.51 mg N m⁻² h⁻¹), whereas the ammonium production rates ranged

219 from 1.16 mg NH₄ m⁻² h⁻¹ to 16.16 mg NH₄ m⁻² h⁻¹, with a mean of 4.66 mg NH₄ m⁻² h⁻¹ (SD = ± 4.38 mg

220 NH₄ m⁻² h⁻¹) (Fig. 6). The total nitrogen (nitrate+ammonium) production rates ranged from 1.31 mg N m⁻

221 ² h⁻¹ to 34.94 mg N m⁻² h⁻¹, with a mean of 7.38 mg N m⁻² h⁻¹ (SD = ± 9.71 mg N m⁻² h⁻¹). At most stations

222 except for stations of CL8 and CS8R, the ammonium production rates were generally higher than the

223 nitrate production rates (Fig. 6). The average nitrate production rate was 0.41 mg NO₃ m⁻² h⁻¹ (SD = ±

224 0.51 mg NO₃ m⁻² h⁻¹) in the northern region, whereas the average nitrate production rate for the southern

225 region was 4.64 mg NO₃ m⁻² h⁻¹ (SD = ± 7.13 mg NO₃ m⁻² h⁻¹). In comparison, the average ammonium

226 production rates for the northern and southern regions were $2.56 \text{ mg NH}_4 \text{ m}^{-2} \text{ h}^{-1}$ ($SD = \pm 1.74 \text{ mg NH}_4$
227 $\text{m}^{-2} \text{ h}^{-1}$) and $6.41 \text{ mg NH}_4 \text{ m}^{-2} \text{ h}^{-1}$ ($SD = \pm 5.28 \text{ mg NH}_4 \text{ m}^{-2} \text{ h}^{-1}$), respectively.

228

229 *3.5. Statistical analysis of environmental variables according to geographic regions*

230 One-way analysis of variance (ANOVA) was performed to assess significant regional differences in
231 the environmental and biological variables of the two geographic regions (i.e., northern and southern).

232 One-way ANOVA revealed significant regional differences for some of the environmental and biological
233 variables in the study area (Table 2). The temperature and salinity of the surface were significantly

234 different due to the effects of various water masses in the region. The stratification also exhibited a
235 significant regional variability due to the higher accumulation of freshwater in the southern region ($p <$

236 0.05). However, the mean mixed layer depths were not significantly different, with means of 7.6 m ($SD =$
237 $\pm 2.8 \text{ m}$) and 8.4 m ($SD = \pm 2.4 \text{ m}$) for the northern and southern regions, respectively (Table 2). The

238 mean depths of nitrcline were similar between the regions, although there were differences between the
239 stations. The ambient nutrient concentrations of the upper 30 m showed highly significant differences,

240 with higher concentrations in the southern region, although the phosphate concentration was not
241 significantly different between the regions (Fig. 3 and Table 2). In addition, the chlorophyll *a*

242 concentrations were significantly different ($p < 0.05$), with a value that was approximately two times
243 higher in the southern region than in the northern region.

244

245 *3.6. FWC distribution*

246 To understand the potential effects of recent changes in the FWC on the primary production in the

247 Chukchi Sea, the FWC data obtained from the three RUSALCA expeditions were used for a comparison.

248 In 2012, the FWC in the study area were ranged from 2.1 to 8.5 m, with a mean of 4.5 m (SD = 1.2 m)

249 (Fig. 7a). The strongest freshwater accumulation was observed in the western side of the CS section and

250 north of the Herald Canyon (FWC = 6.7-8.5 m), whereas the lowest freshwater accumulation was

251 observed at the center of the CL section in the southern region (FWC = 2.8-3.7 m) (Fig. 7a). The FWC in

252 2009 ranged from 2.6 to 11.8 m, with a mean of 5.1 m (SD = \pm 1.7 m) (Fig. 7b). The mean value in 2009

253 was a little higher than that of 2012 due to the high accumulation of FWC from the East Siberian Sea and

254 the region north of Herald Canyon (Fig. 7b). In 2009, the FWC in the southern region was evenly

255 distributed with an accumulation of below 6 m. In 2004, the mean FWC was 4.7 ± 1.3 m and ranged from

256 2.0 to 9.9 m (Fig. 7c). Unlike the observations from 2012 and 2009, the FWC in the southern region in

257 2004 indicated a low accumulation in the western side and a progressive increase in FWC toward the

258 eastern side (Fig. 7c).

259

260 **4. Discussion**

261 *4.1. Regional carbon and nitrogen production rates in 2012*

262 In this study, there were large differences in the carbon and nitrogen production rates the between
263 southern and northern regions (Figs. 5, 6 and Table 2). The average rate of carbon production in the
264 southern region was about five times higher than that of the northern region (Fig. 5 and Table 2).
265 Similarly, the total nitrogen (nitrate+ammonium) production rates were approximately four times higher
266 in the southern region than in the northern region (Fig. 6). In particular, the regional differences were
267 much higher for the nitrate production rate than the ammonium production rate (Table 2). We also found
268 that the carbon production rates obtained from all of the RUSALCA expeditions (2004, 2009 and 2012)
269 showed highly significant differences between the regions ($p < 0.001$, $n = 43$) (data not shown).
270 The regional differences in phytoplankton production rates may have resulted from the different
271 environmental conditions, as revealed the statistical analysis (Table 2). Especially, the different nutrient
272 conditions and thereby different phytoplankton biomasses may be an important reason for the regional
273 differences in the production rates of phytoplankton, since there was a positive relationship between the
274 ambient nutrient concentrations (nitrate) and the chlorophyll *a* concentrations integrated from surface to
275 30 m ($r = 0.6468$, $p < 0.0001$, $n = 41$) (Fig. 8a). Moreover, we found that the carbon, nitrate and
276 ammonium production rates were significantly correlated with the chlorophyll *a* concentration ($r = 0.9234$,
277 $r = 0.9641$ and $r = 0.9798$, $p < 0.0001$, $n=11$, respectively) (Fig. 8b). Even though the regional differences
278 in temperature are quite similar to that in primary production rates, there was no significant relationship
279 between temperature and primary production rates in this study. According to Gosselin et al. (1997), the

280 latitudinal variability in the phytoplankton production and biomass were primarily regulated by changes
281 in the surface ice cover and the depths of the surface mixed layer, which determine the amount of light
282 available to the phytoplankton in the water column. However, this was not the case in our study, as the
283 mixed-layer depths were not significantly different between the southern and northern regions of the
284 Chukchi Sea (Table 2).

285 The production/biomass ratio (P/B ratio), which was calculated by dividing the daily carbon
286 production rate ($\text{mg C m}^{-2} \text{ d}^{-1}$) by the integrated chlorophyll *a* concentration (mg chl m^{-2}), in the southern
287 region ($9.61 \pm 4.26 \text{ mg C (mg chl-a)}^{-1} \text{ d}^{-1}$) was somewhat higher than the P/B ratio in the northern region
288 ($5.46 \pm 1.27 \text{ mg C (mg chl-a)}^{-1} \text{ d}^{-1}$). This result indicated better carbon production efficiency by the
289 phytoplankton in the southern region. Therefore, the regional differences in the primary production rates
290 may have been affected by different production efficiencies in addition to the different phytoplankton
291 biomasses induced under different nutrient conditions.

292

293 *4.2. Primary production rate in 2012 compared to the previous RUSALCA expeditions*

294 Based on a 15-hour photo period in the Chukchi Sea (Hansell and Goering 1990; Lee et al. 2007; Yun
295 et al. 2014) and the hourly carbon production rates measured in this study, in 2012, the daily carbon
296 production rates integrated from the surface to 1 % light depth ranged from 0.02 to $1.61 \text{ g C m}^{-2} \text{ d}^{-1}$. The
297 daily carbon production rate in 2012 (mean \pm SD = $0.42 \pm 0.52 \text{ g C m}^{-2} \text{ d}^{-1}$), which was averaged from the

298 values from all the productivity stations, was quite similar to the daily carbon production rate of 2004
299 (mean \pm SD = 0.41 ± 0.53 g C m $^{-2}$ d $^{-1}$) reported by Lee et al. (2007). The production rates (mean \pm SD =
300 0.26 ± 0.24 g C m $^{-2}$ d $^{-1}$) obtained in 2009 and presented by Yun et al. (2014) were significantly lower than
301 those from 2012 and 2004, which is believed to be due to the different sampling times among the three
302 cruises because the seasonal variation in primary productivity is quite large in this region (Springer and
303 McRoy 1993; Wang et al. 2005; Hill et al., 2013). These differences in the primary production rates
304 obtained by the three cruises also may have been due to interannual variations in primary productivity in
305 the Chukchi Sea, as Hirawake et al. (2012) used satellite remote sensing data obtained from 2002 to 2010
306 to show that the Chukchi Sea experiences strong interannual variation in August and September.

307 In 2012, the average daily carbon production rates were 0.66 g C m $^{-2}$ d $^{-1}$ (SD = ± 0.62 g C m $^{-2}$ d $^{-1}$) in
308 the southern region and 0.14 g C m $^{-2}$ d $^{-1}$ (SD = ± 0.10 g C m $^{-2}$ d $^{-1}$) in the northern region. By comparison,
309 the average daily carbon production rates in the southern and northern regions were 0.57 g C m $^{-2}$ d $^{-1}$ (SD
310 = ± 0.64 g C m $^{-2}$ d $^{-1}$) and 0.16 g C m $^{-2}$ d $^{-1}$ (SD = ± 0.18 g C m $^{-2}$ d $^{-1}$) in 2004, respectively, and 0.38 g C m $^{-2}$
311 d $^{-1}$ (SD = ± 0.26 g C m $^{-2}$ d $^{-1}$) and 0.14 g C m $^{-2}$ d $^{-1}$ (SD = ± 0.16 g C m $^{-2}$ d $^{-1}$) in 2009, respectively. From
312 the regional comparisons, we found that the pattern of primary production in the Chukchi Sea is largely
313 different depending on regions. The primary production rates in the northern region were consistently low,
314 since the regionally low nutrient conditions and phytoplankton biomass. Thus, they were not largely
315 changed among the three cruises. In contrast, the primary production rates for the southern region were

316 considerably variable among the three cruises, although they including seasonal and interannual
317 variations. Since this study revealed that the nutrient is an important factor in controlling primary
318 production, the recent change in primary production for the southern region could be induced by changes
319 in nutrient conditions in the region. The changes in freshwater inputs in the region may have been closely
320 related to the nutrient and primary production variability (detailed in section 4.3).

321

322 *4.3. The effects of FWC on the nutrients and primary production in the southern Chukchi Sea*

323 FWC plays an important role in determining the nutrient distribution/inventory and, therefore, the
324 availability of nutrients for phytoplankton growth in the Arctic Ocean. Coupel et al. (2015) showed that
325 the strong freshening of the Canada Basin resulted in the deepening of the nitracline, which had a
326 negative impact on primary production. In addition, Yun et al. (2014) reported that the low primary
327 production rate in the Chukchi Sea could be due to the decreases in the nutrient and chlorophyll *a*
328 concentrations that resulted from the increased input of fresh waters. In 2012, we found that the
329 freshwater had strongly accumulated in the western side of the southern Chukchi Sea and especially in the
330 CS section (Fig. 7a) due to an inflow of fresh Siberian Coastal Water or sea ice meltwater. This could
331 have resulted in the low primary production rates observed in the western region and the CS section of the
332 southern Chukchi Sea (Fig. 5). In contrast, relatively high production rates were observed in the center of
333 the CL section, the region with the lowest accumulation of freshwater (Figs. 5 and 7a). The strong inflow

334 of Siberian Coastal Water from the East Siberian Sea into the Chukchi Sea was also found in 2009,

335 though it was not detected in 2004 (Figs. 7b and 7c). These inputs of freshwater presumably influenced

336 the nutrient reservoir and its replenishment from deep ~~er waters~~ ~~layers by altering stratification of the water~~

337 ~~column (e.g. Canada Basin)~~ (Coupel et al., 2015), eventually driving the observed changes in primary

338 production in the region. Based on data obtained from southern region during three cruises, we found that

339 FWC had a significant negative effect on the nitrate concentrations ($r = 0.5363, p < 0.05$) and primary

340 production rates ($r = 0.5645, p < 0.05$) (Figs. 9a and 9b). As a result, the primary production rates in the

341 Chukchi Sea could be highly significantly correlated with the nitrate concentrations ($r = 0.7482, p < 0.001$)

342 (Fig. 9c). Therefore, we might conclude that the primary production in the Chukchi Sea could be

343 primarily controlled by nutrient availability related to FWC variability, as reported in previous studies

344 conducted in different regions of the Arctic Ocean (Tremblay and Gagnon, 2009; Tremblay et al., 2002,

345 2006, Coupel et al., 2015). However, the influence of ocean circulations should be examined further

346 because the ocean circulation such as pacific inflow and Beaufort Gyre can redistribute the amount of

347 freshwater (Giles et al., 2012), eventually leading to regional differences in FWC (Giles et al., 2012;

348 Morison et al., 2012). Additionally, we need to consider the local wind field, as the spatial distribution of

349 FWC is largely dependent on the wind and is controlled by atmospheric pressure patterns (Anderson et al.,

350 2011).

351

352 4.4. Current status of the primary production in the Chukchi Sea

353 To understand the recent status of primary production in the Chukchi Sea, the *in situ* measurements of
354 primary production in the region in recent years were plotted with those from the previous studies in
355 decades ago (Fig. 10). The average carbon production rate from the three RUSALCA cruises in the
356 Chukchi Sea was $0.33 \text{ g C m}^{-2} \text{ d}^{-1}$ ($SD = 0.40 \text{ g C m}^{-2} \text{ d}^{-1}$). In addition, Hill and Cota (2005) reported that
357 the mean daily production rate during the initial ice breakup was $0.8 \text{ g C m}^{-2} \text{ d}^{-1}$ in 2002. The daily
358 production rates obtained by Lee et al. (2012) and Lee et al. (2013) were 0.54 and $0.86 \text{ g C m}^{-2} \text{ d}^{-1}$,
359 respectively (Fig. 10). Even though the different sea ice condition could affect difference in the
360 productivity, these recent measurements of primary production (Hill and Cota 2005; Lee et al., 2007,
361 2012 and 2013; Yun et al., 2014; This study) showed significantly lower rates compared with the
362 previously reported rates from the region (Hameedi, 1978; Korsak, 1992; Zeeman, 1992) (t -test, $p < 0.01$).
363 Moreover, it is obviously shown a decreasing trend of primary production ($r = 0.7689$, $p < 0.01$) (Fig. 10).
364 This is very interesting because primary production could be expected to increase in the region due to the
365 increased light availability to the phytoplankton. For example, based on satellite ocean color data, Arrigo
366 et al. (2008) found large increases in the annual net primary production on the continental shelves of the
367 Chukchi Sea as well as Siberian and Laptev seas due to increased open water areas and longer growing
368 seasons. However, the *in situ* measurements of primary production in recent years (Hill and Cota 2005;
369 Lee et al., 2007, 2012 and 2013; Yun et al., 2014; This study) have shown consistently lower primary

370 production rates compared to those in previous studies.

371 The strong seasonal and interannual variation in the region could be suggested for reason causing the

372 low primary production, as discussed above. Hill et al. (2013) found that the seasonal variations in

373 primary production in the southern Chukchi Sea peaked in July and then progressively declined in August

374 and September. In fact, the lowest primary production rates given by Yun et al. (2014) were obtained from

375 the late summer season (i.e., from 1 to 30, September, 2009) compared with the rates found in the present

376 study (from 30 August to 14 September, 2012) or in Lee et al. (2007) (from 11 to 22 August, 2004). In

377 comparison, previous studies (Hameedi, 1978; Korsak, 1992; Zeeman, 1992) included the measurements

378 obtained from July to August (Fig. 10). However, their measurements just starting from the end of July

379 were mostly done during August (Korsak, 1992; Zeeman, 1992). Although recent measurements from the

380 three RUSALCA cruises (2004, 2009 and 2012) may not have reflected the highest values (i.e., July) of

381 primary production, the measurements from Hill and Cota (2005) or Lee et al. (2012 and 2013) include

382 the values in the mid-July and early August. Therefore, the recent low rates of primary production might

383 be reflected by decreasing trend rather than results of seasonal and interannual variations.

384 More plausible reason for the recent low primary production in the Chukchi Sea could be due to the

385 decreased concentrations of nutrients and chlorophyll *a*. According to Whittlestone and Lee (unpublished

386 data), in recent years, there have been significant decreases of 30-50% in nutrient concentrations and

387 approximately 40% in the integrated chlorophyll *a* concentration in the Bering Strait and the Chukchi Sea.

388 Based on the significant relationships between primary production and the nutrient and FWC (discussed
389 in section 4.3), the recent decrease in nutrient and chlorophyll a concentrations may have been closely
390 related to the changes in freshwater inputs in the region. According to Serreze et al. (2006), there was
391 recently larger import of freshwater through the Bering Strait compared with previous estimates.
392 Therefore, the recent decreases in the concentrations of major inorganic nutrients and chlorophyll *a* may
393 have resulted in lower primary production rates in the Chukchi Sea.

394 Recently, the freshwater content in the Arctic Ocean, which includes river discharge, pacific water
395 inflow through the Bering Strait, sea ice melt water and net precipitation (Jones et al., 2008), has
396 increased over the past few decades. If the increased freshwater content in the Chukchi Sea are
397 continuously observed, the Chukchi Sea might have become less productive region compared with
398 previous decades.

399

400 **5. Conclusions**

401 This study reported the regional characteristics of primary production in the Chukchi Sea and recent
402 trend of primary production based on *in situ* measurements. The different nutrient conditions and
403 phytoplankton biomass could be an important reason for the regional differences in the production rates
404 of phytoplankton. Based on comparison between previous studies in decades ago and recent
405 measurements, we found that recent primary production in the Chukchi Sea showed a decreasing trend.

406 The changes in freshwater inputs in the region may have been closely related to the nutrient and primary
407 production variability. Although Coupel et al. (2015) reported that the recent freshening of the Arctic
408 Ocean does not significantly affect primary production in the Chukchi shelf based on comparison with
409 measurements in the deep Canada Basin, our results showed that the freshwater variability in the Chukchi
410 Sea has had a large influence on the recent changes in primary production by controlling the nutrient
411 inventory. If the increased freshwater inflow persists, the primary production in the region will
412 considerably decrease, ultimately resulting in changes in the regional characteristics of primary
413 production. However, a large interannual variability of primary production remains despite the statistical
414 significance observed in this study. Therefore, more measurements under various environmental
415 conditions are needed to better understand the recent variations in the primary production in the Chukchi
416 Sea. In particular, there could be some changes in the phytoplankton community structures because the
417 smaller cells benefit more than the larger cells under increased freshening conditions (Li et al., 2009).

418

419 **Acknowledgments**

420 We thank the captain and crew of the *Professor Khromov* for their outstanding assistance during the cruise.
421 We gratefully acknowledge the physical oceanographers from Woods Hole Oceanographic Institute (M.
422 Swartz, S. Mills and J. Pietro), who provided the CTD data. We also appreciate NOAA for supporting the
423 RUSALCA program. This study was financially supported by the “2015 Post-Doc. Development Program”

424 of Pusan National University, Korea. This work was also supported by grants from the Korea-Polar Ocean

425 in Rapid Transition (K-PORT; PM14040) program funded by the Ministry of Oceans and Fisheries, Korea.

426

427 **References**

428

429 Anderson, L.G. Björk, G., Jutterström, S., Pipko, I., Shakhova, N., Semiletov, I., and Wählström, I.: East
430 Siberian Sea, an Arctic region of very high biogeochemical activity, *Biogeosciences*, 8, 1745–1754,
431 doi:10.5194/bg-8-1745-2011, 2011.

432 Ardyna, M., Babin, M., Gosselin, M., Devred, E., Rainville, L., and Tremblay, J.-É.: Recent Arctic Ocean
433 sea ice loss triggers novel fall phytoplankton blooms, *Geophys. Res. Lett.*, 41, 6207–6212,
434 doi:10.1002/2014GL061047, 2014.

435 Arrigo, K.R., van Dijken, G., and Pabi, S.: Impact of a shrinking Arctic ice cover on marine primary
436 production, *Geophys. Res. Lett.*, 35, L19603, doi:10.1029/2008GL035028, 2008.

437 Carmack, E., McLaughlin, F., Yamamoto-Kawai, M., Itoh, M., Shimada, K., Krishfield, R., and
438 Proshutinsky, A.: Freshwater storage in the Northern Ocean and the special role of the Beaufort
439 Gyre, in: *Arctic-Subarctic Ocean Fluxes: Defining the Role of the Northern Seas in Climate*, edited
440 by: Dickson, R. R., Meincke, J., and Rhines, P., Springer, New York, 145 – 169, 2008.

441 Comiso, J.C., Parkinson, C.L., Gersten, R., and Stock, L.: Accelerated decline in the Arctic sea ice cover,
442 *Geophys. Res. Lett.*, 35, L01703, doi:10.1029/2007GL031972, 2008.

443 Coupel, P., Ruiz-Pino, D., Sicre, M.A., Chen, J.F., Lee, S.H., Schiffrene, N., Li, H.L., and Gascard, J.C.:
444 The impact of freshening on phytoplankton production in the Pacific Arctic Ocean, *Prog. Oceanogr.*,

445 131, 113-125, 2015.

446 Crane, K., and Ostrovskiy, A.: Introduction to the special issue: Russian-American Long-term Census of

447 the Arctic (RUSALCA). *Oceanography*, 28(3), 18(3), <http://dx.doi.org/10.5670/oceanog.2015.54>,

448 2015.

449 Dugdale, R.C., and Goering, J.J.: Uptake of new and regenerated forms of nitrogen in primary

450 productivity, *Limnol. Oceanogr.*, 12, 196-206, 1967.

451 Giles, K.A., Laxon, S.W., Ridout, A.L., Wingham, D.J., and Bacon, S.: Western Arctic Ocean freshwater

452 storage increased by wind-driven spin-up of the Beaufort Gyre, *Nat. Geosci.*, 5, 194–197, 2012.

453 Gosselin, M., Levasseur, M., Wheeler, P.A., Horner, R.A., and Booth, B.C.: New measurements of

454 phytoplankton and ice algal production in the Arctic Ocean. *Deep-Sea Res. Part II*, 44, 1623–1644,

455 1997.

456 Hama, T., Miyazaki, T., Ogawa, Y., Iwakuma, T., Takahashi, M., Otsuki, A., and Ichimura, S.:

457 Measurement of photosynthetic production of a marine phytoplankton population using a stable ^{13}C

458 Isotope, *Mar. Biolo.*, 73, 31-36, 1983.

459 Hameedi, M.J.: Aspects of water column primary productivity in the Chukchi Sea during summer, *Mar.*

460 *Biol.*, 45, 37–46, 1978.

461 Hansell, D.A., and Goering, J.J.: Pelagic nitrogen flux in the northern Bering Sea, *Cont. Shelf Res.*, 10,

462 501–519, 1990.

463 Hill, V., and Cota, G.F.: Spatial patterns of primary production on the shelf, slope and basin of the western

464 Arctic in 2002, *Deep-Sea Res. Part II*, 52, 3344-3354, 2005.

465 Hill, V.J., Matrai, P.A., Olson, E., Suttles, S., Steele, M., Codispoti, L.A., and Zimmerman, R.C.:

466 Synthesis of integrated primary production in the Arctic Ocean: II. In situ and remotely sensed

467 estimates, *Prog. Oceanogr.*, 110, 107–125, <http://dx.doi.org/10.1016/j.pocean.2012.11.005>, 2013.

468 Hirawake, T., Shinmyo, K., Fujiwara, A., and Saitoh, S.: Satellite remote sensing of primary productivity

469 in the Bering and Chukchi Seas using an absorption-based approach, *ICES J. Mar.Sci.*, 69, 1194–

470 1204, 2012.

471 Jones, E.P., Anderson, L.G., Jutterstrom, S., Mintrop, L., and Swift, J.H.: Pacific freshwater, river water

472 and sea ice meltwater across Arctic Ocean basins: results from the 2005 Beringia Expedition, *J.*

473 *Geophys. Res.*, 113, C08012, 2008.

474 Korsak, M.N.: Primary production of organic matter, in: *Results of the Third Joint US–USSR Bering and*

475 *Chukchi Seas Expedition (BERPAC): Summer 1988*, edited by: Nagel, P.A., US Fish and Wildlife

476 Service, Washington, DC, 215–218, 1992.

477 Kwok, R., Cunningham, G. F., Wensnahan, M., Rigor, I., Zwally, H. J., and Yi, D.: Thinning and volume

478 loss of Arctic sea ice: 2003–2008, *J. Geophys. Res.*, 114, C07005, doi:10.1029/2009JC005312.,

479 2009.

480 Lee, S.H., and Whittle, T.E.: Primary and new production in the deep Canada Basin during summer

481 2002, *Polar Biol.*, 28, 190-197, 2005.

482 Lee, S.H., Whittlestone, T.E., and Kang, S.H.: Recent carbon and nitrogen uptake rates of phytoplankton in

483 Bering Strait and the Chukchi Sea, *Cont. Shelf Res.*, 27, 2231-2249, 2007.

484 Lee S.H., Joo, H.M., Liu, Z., Chen, J., and He, J.: Phytoplankton productivity in newly opened waters of

485 the western Arctic Ocean, *Deep Sea Res. Part II*, 81–84, 18–27, 2012.

486 Lee, S.H., Yun, M.S., Kim, B.K., Saitoh, S., Kang, C.-K., Kang, S.H., and Whittlestone, T.E.: Latitudinal

487 carbon productivity in the Bering and Chukchi Seas during the summer in 2007, *Cont. Shelf Res.*, 59,

488 28-36, 2013.

489 Li, W.K.W., McLaughlin, F.A., Lovejoy, C., and Carmack, E.C.: Smallest algae thrive as the Arctic

490 Ocean freshens, *Science*, 326, 539, doi:10.1126/science.1179798, 2009.

491 McLaughlin, F.A., and Carmack, E.C.: Deepening of the nutricline and chlorophyll maximum in the

492 Canada Basin interior, 2003-2009, *Geophys. Res. Lett.*, 37, L24602, doi:10.1029/2010GL045459,

493 2010.

494 McPhee, M.G., Proshutinsky, A., Morison, J.H., Steele, M., and Alkire, M.B.: Rapid change in freshwater

495 content of the Arctic Ocean, *Geophys. Res. Lett.*, 36, L10602, 2009.

496 Morison, J., Kwok, R., Peralta-Ferriz, C., Alkire, M., Rigor, I., Andersen, R., and Steele, M.: Changing

497 Arctic Ocean freshwater pathways, *Nature*, 481, 66–70, 2012.

498 Overland, J.E., and Wang, M.: When will the summer Arctic be nearly sea ice free?, *Geophys. Res.*

499 Lett., 40, 2097-2101, DOI: 10.1002/grl.50316, 2013.

500 Parsons, T.R., Maita, Y., and Lalli, C.M.: A manual of chemical and biological methods for seawater

501 analysis, Pergamon Press, New York, pp 173, 1984.

502 Proshutinsky, A., Krishfield, R., Timmermans, M.-L., Toole, J., Carmack, E., McLaughlin, F., Williams,

503 W.J., Zimmermann, S., Itoh, M., and Shimada, K.: Beaufort Gyre freshwater reservoir: state and

504 variability from observation, *J. Geophys. Res.*, 114, doi:10.1029/2008JC005104, 2009.

505 Rabe, B., Karcher, M., Schauer, U., Toole, J.M., Krishfield, R.A., Pisarev, S., Kauker, F., Gerdes, R., and

506 Kikuchi, T.: An assessment of Arctic Ocean freshwater content changes from the 1990s to the 2006–

507 2008 period, *Deep-Sea Res. Part I*, 58, 173–185, 2011.

508 Serreze, M.C., Barrett, A.P., Slater, A.G., Woodgate, R.A., Aagaard, K., Lammers, R.B., Steele, M.,

509 Moritz, R., Meredith, M., and Lee, C.M.: The large-scale freshwater cycle of the Arctic, *J. Geophys.*

510 *Res.*, 111, C11010, 2006.

511 Shoaf, W.T., and Liim, B.W.: Improved extraction of chlorophyll-a and -b from algae using dimethyl

512 sulfoxide, *Limnol. Oceanogr.*, 21, 926–928, 1976.

513 Springer, A.M., and McRoy, C.P.: The paradox of pelagic food webs in the northern Bering Sea—III.

514 Patterns of primary production, *Cont. Shelf Res.*, 13, 575–599, 1993.

515 Stroeve, J., Serreze, M., Drobot, S., Gearheard, S., Holland, M., Maslanik, J., Meier, W., and Scambos, T.:

516 Arctic sea ice extent plummets in 2007, *EOS Trans., AGU*, 89(2), 13–14, 2008.

517 Stroeve, J. C., Markus, T., Boisvert, L., Miller, J., and Barrett, A.: Changes in Arctic melt season and
518 implications for sea ice loss, *Geophys. Res. Lett.*, 41, 1216–1225, doi:10.1002/2013GL058951, 2014.

519 Tremblay, J.-E., and Gagnon, J.: The effects of irradiance and nutrient supply on the productivity of
520 Arctic waters: a perspective on climate change, *Nato Sci Peace Secur*, http://dx.doi.org/10.1007/978-1-4020-9460-6_7, 73-93, 2009.

521 Tremblay, J.E., Gratton, Y., Fauchot, J., and Price, N.M.: Climatic and oceanic forcing of new, net, and
522 diatom production in the North Water, *Deep-Sea Res. Part II*, 49, 4927–4946, 2002.

523 Tremblay, J.E., Michel, C., Hobson, K.A., Gosselin, M., and Price, N.M.: Bloom dynamics in early
524 opening waters of the Arctic Ocean, *Limnol. Oceanogr.*, 51, 900–912, 2006.

525 Wang, J., Cota, G.F., and Comiso, J.C.: Phytoplankton in the Beaufort and Chukchi Seas: Distribution,
526 dynamics, and environmental forcing, *Deep-Sea Res. Part II*, 52, 3355-3368, 2005.

527 Whitledge, T.E., Malloy, S.C., Patton, C.J., and Wirick, C.D.: Automated nutrient analysis in seawater.
528 Brookhaven National Laboratory Technical Report BNL 51398, 1981.

529 Woodgate, R.A., Aagaard, K., and Weingartner, T.J.: A year in the physical oceanography of the Chukchi
530 Sea: Moored measurements from autumn 1990–1991, *Deep-Sea Res. Part II*, 52, 3116–3149,
531 doi:10.1016/j.dsr2.2005.10.016, 2005a.

532 Woodgate, R.A., Aagaard, K., and Weingartner, T.J.: Monthly temperature, salinity, and transport
533 variability of the Bering Strait throughflow, *Geophys. Res. Lett.*, 32, L04601,

535 doi:10.1029/2004GL021880, 2005b.

536 Woodgate, R.A., Aagaard, K., Weingartner, T.J.: Interannual changes in the Bering Strait fluxes of
537 volume heat and freshwater between 1991 and 2004, *Geophys. Res. Lett.*, 33, L15609,
538 doi:10.1029/2006GL026931, 2006.

539 Woodgate, R.A., Weingartner, T.J., and Lindsay, R.: Observed increases in Bering Strait oceanic fluxes
540 from the Pacific to the Arctic from 2001 to 2011 and their impacts on the Arctic Ocean water
541 column, *Geophys. Res. Lett.*, 39, L24603, doi:10.1029/2012GL054092, 2012.

542 Yamamoto-Kawai, M., McLaughlin, F.A., Carmack, E.C., Nishino, S., Shimada, K., and Kurita, N.:
543 Surface freshening of the Canada Basin, 2003-2007: River runoff versus sea ice meltwater, *J.*
544 *Geophys. Res.*, 114, C00A05, doi:10.1029/2008JC005000, 2009.

545 Yun, M.S., Whitledge, T.E., Gong, M., and Lee, S.H.: Low primary production in the Chukchi Sea shelf,
546 2009, *Cont. Shelf Res.*, 76, 1-11, 2014.

547 Zeeman, S.I.: The importance of primary production and CO₂, in: Results of the Third Joint US-USSR
548 Bering and Chukchi Seas Expedition (BERPAC): Summer 1988, edited by: Nagel, P.A., US Fish and
549 Wildlife Service, Washington, DC, 39-49, 1992.

Figure Legends

Fig. 1. Locations of sampling stations during the 2012 RUSALCA expedition in the Chukchi Sea. The primary production rates were measured at the stations identified by blue circles. The st. CS8R location represents a revisit to st. CS8.

Fig. 2. The distribution of surface temperature [T_{sur}] $^{\circ}\text{C}$ (a), surface salinity [S_{sur}] (b), stratification index [$\Delta\sigma_t$] $[\text{kg m}^{-3}]$ (c), surface mixed layer depth [Z_m] $[\text{m}]$ (d), and nitracline depth [Z_{nit}] $[\text{m}]$ (e) during the 2012 RUSALCA .

Fig. 3. The distributions of the integrated concentrations of ambient nitrite+nitrate [NO_2+NO_3] $[\text{mmol m}^{-2}]$ (a), ammonium [NH_4] $[\text{mmol m}^{-2}]$ (b), phosphate [PO_4] $[\text{mmol m}^{-2}]$ (c), and silicate [SiO_4] $[\text{mmol m}^{-2}]$ (d) from surface to 30 m during the 2012 RUSALCA.

Fig. 4. The chlorophyll *a* concentrations integrated from surface to 30 m [mg m^{-2}] during the 2012 RUSALCA.

Fig. 5. Hourly carbon uptake rates [$\text{mg C m}^{-2} \text{h}^{-1}$] integrated from the surface to 1% light depth during the 2012 RUSALCA.

Fig. 6 Hourly nitrate and ammonium uptake rates [$\text{mg N m}^{-2} \text{h}^{-1}$] integrated from the surface to 1% light depth during the 2012 RUSALCA.

Fig. 7. The distributions of Fresh Water Content (FWC in m) in the Chukchi Sea in 2012 (a), 2009 (b) and 2004 (c).

Fig. 8. Relationships between nitrate concentrations (mmol m^{-2}) and chlorophyll a concentrations (mg m^{-2}) integrated from surface to 30 m (a) ($n = 41$); chlorophyll a concentrations (mg m^{-2}) and daily carbon ($\text{g C m}^{-2} \text{d}^{-1}$) and nitrogen production rate ($\text{mg N m}^{-2} \text{d}^{-1}$) over the euphotic zones (b) ($n = 11$). All data obtained

during the 2012 RUSALCA.

Fig. 9. Relationships between FWC (m) and nitrate concentrations (mmol m^{-2}) (a); FWC (m) and daily primary production rate ($\text{g C m}^{-2} \text{ d}^{-1}$) (b); nitrate concentrations (mmol m^{-2}) and daily primary production rate ($\text{g C m}^{-2} \text{ d}^{-1}$) (c). All data obtained from southern region during the three RUSALCA cruises.

Fig. 10. A recent trend of primary production based on *in situ* carbon uptake measurements (^{13}C or ^{14}C) in the Chukchi Sea. All Chukchi Sea represents southern and northern Chukchi Sea.

Table 1 Location, water depth (m) and euphotic depth (Z_{eu}) for primary productivity stations in the Chukchi Sea in 2012.

Region	Station	Date (mm/dd/yr)	Location		Depth (m)	Z_{eu} (m)
			Latitude ($^{\circ}$ N)	Longitude ($^{\circ}$ W)		
Northern	CEN4	09/05/12	69.9828	-175.6857	63	34
	CEN1A	09/06/12	70.7085	-178.2988	38	20
	HC2	09/07/12	70.9000	-175.0127	74	36
	HC26	09/08/12	71.7878	-174.3945	55	46
	G12	09/11/12	71.3980	-171.2597	55	46
Southern	CS8	08/30/12	67.4312	-169.6030	51	24
	CS17	09/01/12	68.2983	-167.0418	40	22
	CL5A	09/02/12	68.6407	-170.9423	59	20
	CL3R	09/12/12	69.0048	-168.9000	57	26
	CL8	09/13/12	67.8692	-172.5482	53	24
	CS8R	09/14/12	67.4312	-169.6030	51	26

Table 2 Summary of one-way analysis of variance (ANOVA) for environmental variables in two geographic regions of the Chukchi Sea in 2012. The mean values (ranges in parentheses) and their significant differences (> or <) between northern and southern regions are given for surface temperature (T_{sur}), surface salinity (S_{sur}), stratification index ($\Delta\sigma_t$), surface mixed layer depth (Z_m), nitracline depth (Z_{nit}), fresh water content (FWC), nitrite+nitrate concentration of the upper 30 m (NO_2+NO_3), ammonium concentration of the upper 30 m (NH_4), phosphate concentration of the upper 30 m (PO_4), silicate concentration of the upper 30 m (SiO_4) and chlorophyll *a* concentration of the upper 30 m (Chl-a).

*: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$, ns: not significant. n = 52. Also given are mean and range values for carbon production (CP), nitrate production (NP) and ammonium production (AP) (all $\text{mg C or N m}^{-2} \text{ d}^{-1}$). n=11.

Variables	Northern	Southern	
T_{sur} ($^{\circ}\text{C}$)	0.62 (-1.33 ~ 4.13)	<***	3.89 (1.60 ~ 8.53)
S_{sur}	29.27 (27.30 ~ 32.04)	>*	27.48 (21.48 ~ 32.35)
$\Delta\sigma_t$ (kg m^{-3})	3.15 (0.79 ~ 5.34)	<*	4.47 (0.71 ~ 9.71)
Z_m (m)	7.6 (4.0 ~ 14.0)	ns	8.4 (4.0 ~ 14.0)
Z_{nit} (m)	13.0 (2.5 ~ 30.0)	ns	12.6 (2.5 ~ 35.0)
NO_2+NO_3 (mmol m^{-2})	75.01 (21.51 ~ 218.22)	<*	134.15 (21.82 ~ 355.43)
NH_4 (mmol m^{-2})	40.49 (15.36 ~ 86.93)	<**	61.22 (28.54 ~ 109.51)
PO_4 (mmol m^{-2})	22.19 (5.43 ~ 34.26)	ns	25.95 (8.30 ~ 43.57)
SiO_4 (mmol m^{-2})	245.49 (104.79 ~ 800.49)	<***	410.86 (129.17 ~ 669.94)
Chl-a (mg/m^{-2})	21.7 (2.2 ~ 69.3)	<*	54.5 (3.1 ~ 286.4)
CP ($\text{mg C m}^{-2} \text{ d}^{-1}$)	134.7 (16.3 ~ 280.7)		649.1 (151.3 ~ 1628.9)

NP (mg N m ⁻² d ⁻¹)	6.1 (2.2 ~ 19.9)	69.7 (4.5 ~ 281.6)
AP (mg N m ⁻² d ⁻¹)	38.4 (17.4 ~ 83.6)	96.2 (45.0 ~ 242.4)



