

Comments to the Author:

Thank you for your quick response.

Please make final corrections as follows.

There are still many "concentrations" for values with unit "mol m<sup>-2</sup>". Same for Chla with "mg m<sup>-2</sup>". At least I found this in lines 35, 239, 240, 242, 339, 341, Table 2, captions for Figure 8 and Figure 9, label for x-axis in Figure 9.

Please change these to "inventory", "integrated concentration", or "content in upper 30m" etc..

→ As the editor suggested, we changed all (in lines 34, 35, 197-204, 237-240, 272-273, 275, 337, 339, 341, Table 2, captions for Figure 4, Figure 8 and Figure 9, labels for x and y-axis in Figure 8a, label for x-axis in Figure 8b, label for y-axis in Figure 9a, label for x-axis in Figure 9c).

Study area is now indicated in Figure 10, but "all Chukchi Sea" is an exaggeration and "Chukchi Sea" is no need to say. Please mark S, N and S and N in the Figure and explain in the caption: "Study area of each study is indicated by N or S for northern or southern Chukchi Sea, respectively."

→ As the editor suggested, we changed figure 10 and its caption.

1 **Primary production in the Chukchi Sea with potential effects**  
2 **of freshwater content**

3

4 **Mi Sun Yun<sup>a</sup>, Terry E. Whittedge<sup>b</sup>, Dean Stockwell<sup>b</sup>, SeungHyun Son<sup>c</sup>, Jang Han Lee<sup>a</sup>, Jung Woo**  
5 **Park<sup>a</sup>, Da Bin Lee<sup>a</sup>, Jinku Park<sup>a</sup>, Sang H. Lee<sup>a,\*</sup>**

6

7

8 <sup>a</sup> Department of Oceanography, Pusan National University, 30, Jangjeon-dong, Geumjeong-gu, Busan

9 609-735, Korea

10 <sup>b</sup> Institute of Marine Science, University of Alaska Fairbanks, Fairbanks, AK 99775-7220, USA

11 <sup>c</sup> CIRA, Colorado State University, Fort Collins, Colorado, USA

12

13

14

15 \*Corresponding author. Tel.: +82 51 510 2256; fax: +82 51 581 2963.

16 E-mail address: [sanglee@pusan.ac.kr](mailto:sanglee@pusan.ac.kr)

17

18 **Abstract**

19 The *in situ* primary production rates and various environmental variables were investigated in the  
20 Chukchi Sea during the RUSALCA expedition, which was conducted in 2012, to identify the current  
21 status of primary production. A <sup>13</sup>C-<sup>15</sup>N dual tracer technique was used to measure the daily primary  
22 production rates, which ranged from 0.02 to 1.61 g C m<sup>-2</sup> d<sup>-1</sup> (mean ± SD = 0.42 ± 0.52 g C m<sup>-2</sup> d<sup>-1</sup>). The  
23 primary production rates showed large regional differences, with the southern region (0.66 ± 0.62 g C m<sup>-2</sup>  
24 d<sup>-1</sup>) producing approximately five times as much as the northern region (0.14 ± 0.10 g C m<sup>-2</sup> d<sup>-1</sup>), which  
25 was primarily due to the differences in phytoplankton biomasses induced by regional nutrient conditions.  
26 The primary production rates in the Chukchi Sea were averaged using data acquired during the three  
27 different RUSALCA expeditions (2004, 2009, and 2012) as 0.33 g C m<sup>-2</sup> d<sup>-1</sup> (SD = 0.40 g C m<sup>-2</sup> d<sup>-1</sup>),  
28 which was significantly lower than previously reported rates. In addition to strong seasonal and  
29 interannual variations in primary production, recent decreases in the concentrations of major inorganic  
30 nutrients and chlorophyll *a* could be among the reasons for the recent low primary production in the  
31 Chukchi Sea because the primary production is mainly affected by nutrient concentration and  
32 phytoplankton biomass. The nutrient inventory and primary production appear to be largely influenced by  
33 the freshwater content (FWC) variability in the region due to the significant relationships between FWC,  
34 nitrate ~~concentrations-inventory~~ (r = 0.54, p < 0.05) and primary production rates (r = 0.56, p < 0.05).  
35 Moreover, we found highly significant relationships between the nutrient ~~levels-inventory~~ and the primary

36 production rates ( $r = 0.75$ ,  $p < 0.001$ ). In conclusion, the primary production in the Chukchi Sea is  
37 primarily controlled by nutrient availability which is strongly related to the FWC variability. Our results  
38 imply that the predicted increase in freshwater accumulation might cause a decrease in primary  
39 production by lowering the nutrient inventory in the euphotic zone of the Chukchi Sea.

40

41 ***Keywords:***

42 Phytoplankton, Primary production, Chukchi Sea, Freshwater content, Arctic

43

44

45 **1. Introduction**

46 Over recent years, the Arctic Ocean has undergone drastic changes in the extent and thickness of sea  
47 ice (Stroeve et al., 2008; Comiso et al., 2008; Kwok et al., 2009; Overland and Wang, 2013). The  
48 continuing loss of sea ice may result in changes to various physical and chemical environmental  
49 conditions in the Arctic Ocean. For example, the loss in sea ice cover allows more sunlight to enter the  
50 surface layer of the Arctic Ocean, which results in a longer growing season for phytoplankton growth  
51 (Arrigo et al., 2008; Ardyna et al., 2014). Stroeve et al. (2014) reported that the arctic melt season has  
52 lengthened at a rate of 5 days decade<sup>-1</sup> from 1979 to 2013, due to later autumn freeze-up. In accordance  
53 with their findings, Ardyna et al. (2014) documented the development of a second bloom in the Arctic  
54 Ocean during the fall, which coincides with the delayed freeze-up and the increased exposure to wind  
55 stress.

56 However, the loss in sea ice can cause an increase in the input of freshwater (McPhee et al., 2009). In  
57 fact, the freshwater volumes in the Canada and Makarov Basins increased by of 8500 km<sup>3</sup> in 2008 due to  
58 increased sea ice melting and river discharge (McPhee et al., 2009; Rabe et al., 2011). This phenomenon  
59 can enhance the stratification in the upper ocean (Yamamoto-Kawai et al., 2009) and consequently reduce  
60 vertical mixing, thereby preventing nutrient inputs from deep waters to the euphotic zone. In fact,  
61 McLaughlin and Carmack (2010) found a deepening of the nutricline due to the accumulation of surface  
62 freshwater in the Canada Basin.

63 In the Chukchi Sea as inflow shelf, there was an increased volume flux of 50% in 2011 (~ 1.1Sv)  
64 relative to 2001 (~ 0.7 Sv), which was accompanied by increases in heat and freshwater fluxes (Woodgate  
65 et al., 2012). Though the volume flux may vary both seasonally and annually under the influence of the  
66 local wind fields, the recent increases in freshwater fluxes in the region may have important implications  
67 for phytoplankton in terms of nutrient availability for their growth (Woodgate et al., 2005a, 2005b, 2006).  
68 Thus, it is important to identify how phytoplankton respond to these environmental changes in the region  
69 in terms of production and/or community structure. According to Li et al. (2009), the phytoplankton  
70 community has changed under the freshening and stratifying condition in the Canada Basin. Notably, the  
71 abundance of small phytoplankton (< 2  $\mu\text{m}$ ) has increased, whereas the abundance of large phytoplankton  
72 (2-20  $\mu\text{m}$ ) has decreased. Yun et al. (2014) also found that compared with previous reports, the small  
73 phytoplankton were more abundant on the Chukchi Sea shelf, which is dominated by low nutrients and  
74 freshening conditions. Therefore, the changes in recent phytoplankton production under the rapidly  
75 changing environmental conditions need to be monitored because the changes in phytoplankton  
76 production could have important implications for understanding ecosystem changes in the Arctic Ocean.

77 In order to understand climate and ecosystem change in the Pacific Arctic Ocean which is a region  
78 summer sea ice cover was declining dramatically (Crane and Ostrovskiy, 2015), the RUSALCA (Russian-  
79 American Long-term Census of the Arctic) expedition, which is a joint US-Russian research program,  
80 started from 2004 as multidisciplinary investigations in the Bering and Chukchi Seas. Three RUSALCA

81 expeditions (2004, 2009, and 2012) provided a good opportunity for continuous measurements of the  
82 primary production in the entire Chukchi Sea, including the Exclusive Economic Zone of the Russian  
83 Federation. The 2004 RUSALCA expedition was conducted from 8 to 24 August, 2004 (Lee et al., 2007).  
84 The 2009 RUSALCA expedition was executed from 1 to 30 September, 2009 (Yun et al., 2014). The 2012  
85 RUSALCA expedition was carried out from 27 August to 16 September, 2012. This study is part of the  
86 2012 RUSALCA expedition.

87 In this study, we addressed the regional characteristics of primary production by examining the main  
88 driving factors responsible for the regional variability in the Chukchi Sea based on measurements taken in  
89 2012. In addition, we investigated the recent trends in primary production in the Chukchi Sea based on  
90 the results of the three RUSALCA expeditions (2004, 2009, and 2012) in the Chukchi Sea. Finally, we  
91 emphasized the potential effects of freshwater accumulation on the primary production in the Chukchi  
92 Sea because changing amounts and distributions of freshwater content could lead to changes in the  
93 primary production rates.

94

## 95 **2. Materials and methods**

### 96 *2.1. Study area and sampling*

97 The RUSALCA expedition in 2012 was conducted onboard the Russian vessel *Professor Khromov* in  
98 the Chukchi Sea from 27 August to 16 September. The study area was comprised of several sections

99 between the Bering Strait and the vicinity of Herald Canyon (Fig. 1). To understand the regional  
100 characteristics of primary production, the study area was divided into two geographic regions  
101 (northern/southern) following Yun et al. (2014). The northern region consisted of stations in the vicinity  
102 of Herald Canyon (CEN and HC sections) (Fig. 1). The stations in the Chukchi South and Cape Lisburne  
103 (CS and CL sections) were included in the southern region. Most of the bathymetric depths in the entire  
104 study area were quite shallow, with a mean of 55 m (SD =  $\pm 11$  m). Between the production stations, the  
105 depth of euphotic zone from the surface to 1% light depth varied between 20 and 46 m, with a mean of 29  
106  $\pm 10$  m (Table 1).

107 Oceanographic/biological samples were taken from a total of 54 conductivity-temperature-depth  
108 (CTD) stations. The vertical profiles of water temperature and salinity were obtained using a Sea-Bird  
109 model SBE911*plus* CTD profiler. Water samples were collected with a stainless-steel rosette sampler that  
110 was equipped with 21 10-liter bottles at every CTD station. The data from the previous RUSALCA  
111 expeditions (in 2004 and 2009) were included to understand the recent trends in primary production in the  
112 Chukchi Sea.

113

## 114 2.2. *Physical and chemical variables*

115 The stratification index of the water column ( $\Delta\sigma_t$ ) (in  $\text{kg m}^{-3}$ ) was determined as the difference in  $\Delta\sigma_t$   
116 values between the surface and the bottom depth according to Yun et al. (2014). The surface mixed layer



117 ( $Z_m$ ) was defined as the depth at which the density ( $\sigma_t$ ) gradient was  $0.05 \text{ kg m}^{-3}$  higher than the  
118 surface density, as in Coupel et al. (2015). The depth of the euphotic zone ( $Z_{eu}$ ) in this study was defined  
119 as the depth receiving 1% of the surface PAR value, as in Lee et al. (2007) and Yun et al. (2014), and was  
120 obtained from a Biospherical QSP-2300 PAR sensor (Biospherical Instruments Inc.) that was lowered  
121 with the CTD/rosette sampler. The nitracline ( $Z_{nit}$ ) was determined as the depth at which the nitrate  
122 gradient was greater than  $0.1 \mu\text{M m}^{-1}$  according to the definition of Coupel et al. (2015).

123

### 124 2.3. Fresh Water Content (FWC)

125 To assess the surface water freshening, the freshwater content (FWC) was calculated following  
126 Carmack et al. (2008):

$$127 \quad FWC = \int_{z_{lim}}^0 (1 - S(z)/S_{ref}) dz$$

128

129 where  $S$  and  $S_{ref}$  are the *in situ* and reference salinities, respectively, and  $Z_{lim}$  is the depth where  $S$  equals  
130  $S_{ref}$  (34.8 on the practical salinity scale). We used a reference salinity of 34.8 following Aagaard and  
131 Carmack (1989) to computing freshwater since it has been considered as the mean salinity for the Arctic  
132 Ocean.

133

### 134 2.4. Nutrient concentration measurements

135 The discrete water samples used in measuring the nutrient concentrations were obtained from 5 to 9  
136 different depths depending on the water depths. The dissolved inorganic nutrient concentrations  
137 (nitrite+nitrate, ammonium, phosphate, and silicate) were analyzed onboard immediately after collection  
138 using an automated nutrient analyzer (ALPKEM RFA model 300) following the method of Whitedge et  
139 al. (1981).

140

#### 141 *2.5. Chlorophyll a concentration measurements*

142 The water samples used for measuring the chlorophyll *a* concentration were obtained from 4 to 7  
143 different depths at most stations. The water samples were filtered through Whatman GF/F filters (24 mm),  
144 and the filters were then kept frozen until analysis in the laboratory. The filters were subsequently  
145 extracted in a 3:2 mixture of 90% acetone and DMSO in a freezer for 24 h, followed by centrifugation  
146 (Shoaf and Lium, 1976). The chlorophyll *a* concentrations were measured using a Turner Designs model  
147 10-AU fluorometer, which was calibrated using commercially available preparations of purified  
148 chlorophyll *a* (Turner Designs, USA). The methods and calculations used to determine the chlorophyll *a*  
149 concentrations followed the procedure of Parsons et al. (1984).

150

#### 151 *2.6. In situ primary production measurements*

152 The water samples used to measure primary production were collected at six photic depths (100, 50,

153 30, 12, 5, and 1% penetration of the surface irradiance, PAR). At 11 selected morning stations, the *in situ*  
154 primary productions of phytoplankton were measured using a  $^{13}\text{C}$ - $^{15}\text{N}$  dual tracer technique (Lee and  
155 Whitedge, 2005; Lee et al., 2007). This method could be useful for distinguish the relative importance of  
156 nitrate and ammonium as nitrogen sources for the cell and population (Dugdale and Goering, 1967). We  
157 followed the same analytical procedure of Lee et al. (2007) and Yun et al. (2014) to the measure primary  
158 production to consistently compare the primary production levels determined in the three studies. Briefly,  
159 heavy isotope-enriched (98-99%) carbon ( $\text{NaH}^{13}\text{CO}_3$ ), nitrate ( $\text{K}^{15}\text{NO}_3$ ), and ammonium ( $^{15}\text{NH}_4\text{Cl}$ )  
160 substrates were inoculated in polycarbonate bottles (1 L) and then incubated on deck in a large  
161 polycarbonate incubator cooled with running surface seawater under natural light conditions. After  
162 approximately 4 to 5 h of incubation , all samples were filtered using pre-combusted (450°C, 4 h) glass  
163 fiber filters (Whatman GF/F; diameter = 25 mm). After HCl fume treatment, the samples were sent to the  
164 Alaska Stable Isotope Laboratory of the University of Alaska, Fairbanks, USA. The abundances of  $^{13}\text{C}$   
165 and  $^{15}\text{N}$  and the total amounts of particulate organic carbon (POC) and nitrogen (PON) were determined  
166 using a Thermo Finnigan Delta+XL mass spectrometer. Finally, the carbon and nitrogen production rates  
167 were calculated based on Hama et al. (1983) and Dugdale and Goering (1967), respectively.

168

### 169 **3. Results**

#### 170 *3.1. Physical conditions*

171 The surface temperature ( $T_{sur}$ ) varied from -2 to 9 °C in the study area in 2012 (Fig. 2a). The higher  
172 temperatures were found in the eastern side of the southern Chukchi Sea due to the strong influence of the  
173 Alaskan Coastal Water (warmer and less saline). The freezing temperatures were observed in the vicinity  
174 of the Herald Canyon and gradually decreased toward the northward. At the surface, the salinity varied  
175 between 21 and 33 psu. The surface salinity ( $S_{sur}$ ) was considerably lower in the southwestern side  
176 compared with the northeastern side of the southern Chukchi Sea (Fig. 2b). The stratification index ( $\Delta\sigma_t$ )  
177 in the study area ranged from 0.7 to 9.7 kg m<sup>-3</sup>, with a mean of  $3.8 \pm 2.2$  kg m<sup>-3</sup>. The stratification in the  
178 southern region was higher than in the northern region (Fig. 2c). The general distribution of the  
179 stratification index was similar to that of surface salinity because it tended to be high in areas where  
180 surface salinity was low. The surface mixed layer ( $Z_m$ ) was thinner than 15 m over the entire study area  
181 (Fig. 2d). In the study area, the depths of nitracline ( $Z_{nit}$ ) ranged from 2.5 m to 35 m (Fig. 2e), with a  
182 mean nitracline depth of  $12.8 \pm 7.7$  m.

183

### 184 3.2. Nutrient distribution

185 Since the mean depths of euphotic zone in this study was about 30 m, the distribution of ambient  
186 nutrient concentrations integrated from surface to 30 m of the water column is shown in Fig. 3. The  
187 inventory of nitrite+nitrate ranged from 21.51 to 355.43 mmol m<sup>-2</sup>, whereas the ammonium inventory  
188 ranged from 15.36 to 109.51 mmol m<sup>-2</sup> (Figs. 3a and 3b). High nitrite+nitrate inventory that exceeded 200

189 mmol m<sup>-2</sup> were observed at the center of the CL section (Fig. 3a). The inventories of these nutrients in the  
190 southern region (134.15 ± 98.41 mmol m<sup>-2</sup> for nitrite+nitrate and 61.22 ± 20.55 mmol m<sup>-2</sup> for ammonium,  
191 respectively) were approximately two times higher than their inventories in the northern region (75.01 ±  
192 52.01 mmol m<sup>-2</sup> for nitrite+nitrate and 40.49 ± 20.69 mmol m<sup>-2</sup> for ammonium) (see Table 2). The  
193 inventory of phosphate in the study area was fairly uniform, with a mean of 24.03 ± 8.30 mmol m<sup>-2</sup> (Fig.  
194 3c). The silicate inventory was generally higher in the southern region than in the northern region (Fig.  
195 3d).

196

### 197 3.3. Chlorophyll *a* ~~concentration-content~~

198 The distribution of the chlorophyll *a* ~~concentration-content~~ in the upper 30 m (i.e., mean depth of  
199 euphotic zone in this study) of the entire study area is shown in Fig. 4. High chlorophyll *a* ~~concentrations~~  
200 ~~content~~ of over 80 mg m<sup>-2</sup> were observed in the western side of the CL section (from st. CL5 to st. CL8),  
201 and low chlorophyll *a* ~~concentrations-content~~ were shown in the western side of the CS section (Fig. 4).  
202 The highest ~~concentration-content~~ (286.4 mg m<sup>-2</sup>) was obtained at station CL8. Over the entire study area,  
203 the mean chlorophyll *a* ~~concentration-content~~ ~~integrated from the surface to in upper~~ 30 m was 42.7 mg m<sup>-2</sup>  
204 <sup>2</sup> (SD = ± 57.4 mg m<sup>-2</sup>). The average ~~concentrations-contents~~ were 21.7 mg m<sup>-2</sup> (SD = ± 19.6 mg m<sup>-2</sup>) and  
205 54.5 mg m<sup>-2</sup> (SD = ± 67.7 mg m<sup>-2</sup>) for the northern and southern regions, respectively.

206

207 *3.4. Primary production rates*

208 Overall, the hourly carbon production rates integrated over the euphotic zone from six light depths  
209 ranged from 1.1 to 108.6 mg C m<sup>-2</sup> h<sup>-1</sup>, with a mean of 27.7 mg C m<sup>-2</sup> h<sup>-1</sup> (SD = 34.7 mg C m<sup>-2</sup> h<sup>-1</sup>). The  
210 highest primary production rates were found at station CL8 (108.6 mg C m<sup>-2</sup> h<sup>-1</sup>) followed by station  
211 CL5A (82.1 mg C m<sup>-2</sup> h<sup>-1</sup>) (Fig. 5). In the northern region, the carbon production rates ranged from 1.1 to  
212 18.7 mg C m<sup>-2</sup> h<sup>-1</sup>, with a mean of 9.0 mg C m<sup>-2</sup> h<sup>-1</sup> (SD = ± 6.4 mg C m<sup>-2</sup> h<sup>-1</sup>). In comparison, the average  
213 rates in the southern region were approximately five times higher than the average rates in the northern  
214 region (43.3 ± 41.7 mg C m<sup>-2</sup> h<sup>-1</sup>).

215 The vertically integrated nitrate production rates ranged from 0.14 to 18.77 mg NO<sub>3</sub> m<sup>-2</sup> h<sup>-1</sup>, with a  
216 mean of 2.72 mg N m<sup>-2</sup> h<sup>-1</sup> (SD = ± 5.51 mg N m<sup>-2</sup> h<sup>-1</sup>), whereas the ammonium production rates ranged  
217 from 1.16 mg NH<sub>4</sub> m<sup>-2</sup> h<sup>-1</sup> to 16.16 mg NH<sub>4</sub> m<sup>-2</sup> h<sup>-1</sup>, with a mean of 4.66 mg NH<sub>4</sub> m<sup>-2</sup> h<sup>-1</sup> (SD = ± 4.38 mg  
218 NH<sub>4</sub> m<sup>-2</sup> h<sup>-1</sup>) (Fig. 6). The total nitrogen (nitrate+ammonium) production rates ranged from 1.31 mg N m<sup>-2</sup>  
219 h<sup>-1</sup> to 34.94 mg N m<sup>-2</sup> h<sup>-1</sup>, with a mean of 7.38 mg N m<sup>-2</sup> h<sup>-1</sup> (SD = ± 9.71 mg N m<sup>-2</sup> h<sup>-1</sup>). At most stations  
220 except for stations of CL8 and CS8R, the ammonium production rates were generally higher than the  
221 nitrate production rates (Fig. 6). The average nitrate production rate was 0.41 mg NO<sub>3</sub> m<sup>-2</sup> h<sup>-1</sup> (SD = ±  
222 0.51 mg NO<sub>3</sub> m<sup>-2</sup> h<sup>-1</sup>) in the northern region, whereas the average nitrate production rate for the southern  
223 region was 4.64 mg NO<sub>3</sub> m<sup>-2</sup> h<sup>-1</sup> (SD = ± 7.13 mg NO<sub>3</sub> m<sup>-2</sup> h<sup>-1</sup>). In comparison, the average ammonium  
224 production rates for the northern and southern regions were 2.56 mg NH<sub>4</sub> m<sup>-2</sup> h<sup>-1</sup> (SD = ± 1.74 mg NH<sub>4</sub>

225  $\text{m}^{-2} \text{h}^{-1}$ ) and  $6.41 \text{ mg NH}_4 \text{ m}^{-2} \text{ h}^{-1}$  ( $\text{SD} = \pm 5.28 \text{ mg NH}_4 \text{ m}^{-2} \text{ h}^{-1}$ ), respectively.

226

### 227 3.5. Statistical analysis of environmental variables according to geographic regions

228 One-way analysis of variance (ANOVA) was performed to assess significant regional differences in  
229 the environmental and biological variables of the two geographic regions (i.e., northern and southern).

230 One-way ANOVA revealed significant regional differences for some of the environmental and biological  
231 variables in the study area (Table 2). The temperature and salinity of the surface were significantly

232 different due to the effects of various water masses in the region. The stratification also exhibited a

233 significant regional variability due to the higher accumulation of freshwater in the southern region ( $p <$

234  $0.05$ ). However, the mean mixed layer depths were not significantly different, with means of  $7.6 \text{ m}$  ( $\text{SD} =$

235  $\pm 2.8 \text{ m}$ ) and  $8.4 \text{ m}$  ( $\text{SD} = \pm 2.4 \text{ m}$ ) for the northern and southern regions, respectively (Table 2). The

236 mean depths of nitracline were similar between the regions, although there were differences between the

237 stations. The ambient nutrient ~~concentrations-inventory~~ of the upper 30 m showed highly significant

238 differences, with higher ~~concentrations-inventory~~ in the southern region, although the phosphate

239 ~~concentration-inventory~~ was not significantly different between the regions (Fig. 3 and Table 2). In

240 addition, the chlorophyll *a* ~~concentrations-contents~~ were significantly different ( $p < 0.05$ ), with a value

241 that was approximately two times higher in the southern region than in the northern region.

242

243 3.6. FWC distribution

244 To understand the potential effects of recent changes in the FWC on the primary production in the  
245 Chukchi Sea, the FWC data obtained from the three RUSALCA expeditions were used for a comparison.  
246 In 2012, the FWC in the study area were ranged from 2.1 to 8.5 m, with a mean of 4.5 m (SD = 1.2 m)  
247 (Fig. 7a). The strongest freshwater accumulation was observed in the western side of the CS section and  
248 north of the Herald Canyon (FWC = 6.7-8.5 m), whereas the lowest freshwater accumulation was  
249 observed at the center of the CL section in the southern region (FWC = 2.8-3.7 m) (Fig. 7a). The FWC in  
250 2009 ranged from 2.6 to 11.8 m, with a mean of 5.1 m (SD = ± 1.7 m) (Fig. 7b). The mean value in 2009  
251 was a little higher than that of 2012 due to the high accumulation of FWC from the East Siberian Sea and  
252 the region north of Herald Canyon (Fig. 7b). In 2009, the FWC in the southern region was evenly  
253 distributed with an accumulation of below 6 m. In 2004, the mean FWC was  $4.7 \pm 1.3$  m and ranged from  
254 2.0 to 9.9 m (Fig. 7c). Unlike the observations from 2012 and 2009, the FWC in the southern region in  
255 2004 indicated a low accumulation in the western side and a progressive increase in FWC toward the  
256 eastern side (Fig. 7c).

257

258 **4. Discussion**

259 *4.1. Regional carbon and nitrogen production rates in 2012*

260 In this study, there were large differences in the carbon and nitrogen production rates the between



261 southern and northern regions (Figs. 5, 6 and Table 2). The average rate of carbon production in the  
262 southern region was about five times higher than that of the northern region (Fig. 5 and Table 2).  
263 Similarly, the total nitrogen (nitrate+ammonium) production rates were approximately four times higher  
264 in the southern region than in the northern region (Fig. 6). In particular, the regional differences were  
265 much higher for the nitrate production rate than the ammonium production rate (Table 2). We also found  
266 that the carbon production rates obtained from all of the RUSALCA expeditions (2004, 2009 and 2012)  
267 showed highly significant differences between the regions ( $p < 0.001$ ,  $n = 43$ ) (data not shown).

268 The regional differences in phytoplankton production rates may have resulted from the different  
269 environmental conditions, as revealed the statistical analysis (Table 2). Especially, the different nutrient  
270 conditions and thereby different phytoplankton biomasses may be an important reason for the regional  
271 differences in the production rates of phytoplankton, since there was a positive relationship between the  
272 ambient nutrient ~~concentrations-inventory~~ (nitrate) and the chlorophyll *a* ~~concentrations-integrated from~~  
273 ~~surface-to-content in upper~~ 30 m ( $r = 0.6468$ ,  $p < 0.0001$ ,  $n = 41$ ) (Fig. 8a). Moreover, we found that the  
274 carbon, nitrate and ammonium production rates were significantly correlated with the chlorophyll *a*  
275 ~~concentration-content in upper 30 m~~ ( $r = 0.9234$ ,  $r = 0.9641$  and  $r = 0.9798$ ,  $p < 0.0001$ ,  $n=11$ ,  
276 respectively) (Fig. 8b). Even though the regional differences in temperature are quite similar to that in  
277 primary production rates, there was no significant relationship between temperature and primary  
278 production rates in this study. According to Gosselin et al. (1997), the latitudinal variability in the

279 phytoplankton production and biomass were primarily regulated by changes in the surface ice cover and  
280 the depths of the surface mixed layer, which determine the amount of light available to the phytoplankton  
281 in the water column. However, this was not the case in our study, as the mixed-layer depths were not  
282 significantly different between the southern and northern regions of the Chukchi Sea (Table 2).

283 The production/biomass ratio (P/B ratio), which was calculated by dividing the daily carbon  
284 production rate ( $\text{mg C m}^{-2} \text{d}^{-1}$ ) by the integrated chlorophyll *a* concentration ( $\text{mg chl-a m}^{-2}$ ), in the southern  
285 region ( $9.61 \pm 4.26 \text{ mg C (mg chl-a)}^{-1} \text{d}^{-1}$ ) was somewhat higher than the P/B ratio in the northern region  
286 ( $5.46 \pm 1.27 \text{ mg C (mg chl-a)}^{-1} \text{d}^{-1}$ ). This result indicated better carbon production efficiency by the  
287 phytoplankton in the southern region. Therefore, the regional differences in the primary production rates  
288 may have been affected by different production efficiencies in addition to the different phytoplankton  
289 biomasses induced under different nutrient conditions.

290

#### 291 4.2. Primary production rate in 2012 compared to the previous RUSALCA expeditions

292 Based on a 15-hour photo period in the Chukchi Sea (Hansell and Goering 1990; Lee et al. 2007; Yun  
293 et al. 2014) and the hourly carbon production rates measured in this study, in 2012, the daily carbon  
294 production rates integrated from the surface to 1 % light depth ranged from 0.02 to 1.61  $\text{g C m}^{-2} \text{d}^{-1}$ . The  
295 daily carbon production rate in 2012 (mean  $\pm$  SD =  $0.42 \pm 0.52 \text{ g C m}^{-2} \text{d}^{-1}$ ), which was averaged from the  
296 values from all the productivity stations, was quite similar to the daily carbon production rate of 2004

297 (mean  $\pm$  SD =  $0.41 \pm 0.53$  g C m<sup>-2</sup> d<sup>-1</sup>) reported by Lee et al. (2007). The production rates (mean  $\pm$  SD =  
298  $0.26 \pm 0.24$  g C m<sup>-2</sup> d<sup>-1</sup>) obtained in 2009 and presented by Yun et al. (2014) were significantly lower than  
299 those from 2012 and 2004, which is believed to be due to the different sampling times among the three  
300 cruises because the seasonal variation in primary productivity is quite large in this region (Springer and  
301 McRoy 1993; Wang et al. 2005; Hill et al., 2013). These differences in the primary production rates  
302 obtained by the three cruises also may have been due to interannual variations in primary productivity in  
303 the Chukchi Sea, as Hirawake et al. (2012) used satellite remote sensing data obtained from 2002 to 2010  
304 to show that the Chukchi Sea experiences strong interannual variation in August and September.

305 In 2012, the average daily carbon production rates were  $0.66$  g C m<sup>-2</sup> d<sup>-1</sup> (SD =  $\pm 0.62$  g C m<sup>-2</sup> d<sup>-1</sup>) in  
306 the southern region and  $0.14$  g C m<sup>-2</sup> d<sup>-1</sup> (SD =  $\pm 0.10$  g C m<sup>-2</sup> d<sup>-1</sup>) in the northern region. By comparison,  
307 the average daily carbon production rates in the southern and northern regions were  $0.57$  g C m<sup>-2</sup> d<sup>-1</sup> (SD  
308 =  $\pm 0.64$  g C m<sup>-2</sup> d<sup>-1</sup>) and  $0.16$  g C m<sup>-2</sup> d<sup>-1</sup> (SD =  $\pm 0.18$  g C m<sup>-2</sup> d<sup>-1</sup>) in 2004, respectively, and  $0.38$  g C m<sup>-2</sup>  
309 d<sup>-1</sup> (SD =  $\pm 0.26$  g C m<sup>-2</sup> d<sup>-1</sup>) and  $0.14$  g C m<sup>-2</sup> d<sup>-1</sup> (SD =  $\pm 0.16$  g C m<sup>-2</sup> d<sup>-1</sup>) in 2009, respectively. From  
310 the regional comparisons, we found that the pattern of primary production in the Chukchi Sea is largely  
311 different depending on regions. The primary production rates in the northern region were consistently low,  
312 since the regionally low nutrient conditions and phytoplankton biomass. Thus, they were not largely  
313 changed among the three cruises. In contrast, the primary production rates for the southern region were  
314 considerably variable among the three cruises, although they including seasonal and interannual

315 variations. Since this study revealed that the nutrient is an important factor in controlling primary  
316 production, the recent change in primary production for the southern region could be induced by changes  
317 in nutrient conditions in the region. The changes in freshwater inputs in the region may have been closely  
318 related to the nutrient and primary production variability (detailed in section 4.3).

319

#### 320 *4.3. The effects of FWC on the nutrients and primary production in the southern Chukchi Sea*

321 FWC plays an important role in determining the nutrient distribution/inventory and, therefore, the  
322 availability of nutrients for phytoplankton growth in the Arctic Ocean. Coupel et al. (2015) showed that  
323 the strong freshening of the Canada Basin resulted in the deepening of the nitracline, which had a  
324 negative impact on primary production. In addition, Yun et al. (2014) reported that the low primary  
325 production rate in the Chukchi Sea could be due to the decreases in the nutrient and chlorophyll *a*  
326 concentrations that resulted from the increased input of fresh waters. In 2012, we found that the  
327 freshwater had strongly accumulated in the western side of the southern Chukchi Sea and especially in the  
328 CS section (Fig. 7a) due to an inflow of fresh Siberian Coastal Water or sea ice meltwater. This could  
329 have resulted in the low primary production rates observed in the western region and the CS section of the  
330 southern Chukchi Sea (Fig. 5). In contrast, relatively high production rates were observed in the center of  
331 the CL section, the region with the lowest accumulation of freshwater (Figs. 5 and 7a). The strong inflow  
332 of Siberian Coastal Water from the East Siberian Sea into the Chukchi Sea was also found in 2009,

333 though it was not detected in 2004 (Figs. 7b and 7c). These inputs of freshwater presumably influenced  
334 the nutrient reservoir and its replenishment from deeper layers by altering stratification of the water  
335 column (Coupel et al., 2015), eventually driving the observed changes in primary production in the region.  
336 Based on data obtained from southern region during three cruises, we found that FWC had a significant  
337 negative effect on the nitrate ~~concentrations-inventory~~ ( $r = 0.5363$ ,  $p < 0.05$ ) and primary production rates  
338 ( $r = 0.5645$ ,  $p < 0.05$ ) (Figs. 9a and 9b). As a result, the primary production rates in the Chukchi Sea  
339 could be highly significantly correlated with the nitrate ~~concentrations-inventory~~ ( $r = 0.7482$ ,  $p < 0.001$ )  
340 (Fig. 9c). Therefore, we might conclude that the primary production in the Chukchi Sea could be  
341 primarily controlled by nutrient ~~availability-inventory~~ related to FWC variability, as reported in previous  
342 studies conducted in different regions of the Arctic Ocean (Tremblay and Gagnon, 2009; Tremblay et al.,  
343 2002, 2006, Coupel et al., 2015). However, the influence of ocean circulations should be examined  
344 further because the ocean circulation such as pacific inflow and Beaufort Gyre can redistribute the  
345 amount of freshwater (Giles et al., 2012), eventually leading to regional differences in FWC (Giles et al.,  
346 2012; Morison et al., 2012). Additionally, we need to consider the local wind field, as the spatial  
347 distribution of FWC is largely dependent on the wind and is controlled by atmospheric pressure patterns  
348 (Anderson et al., 2011).

349

350 *4.4. Current status of the primary production in the Chukchi Sea*

351 To understand the recent status of primary production in the Chukchi Sea, the *in situ* measurements of  
352 primary production in the region in recent years were plotted with those from the previous studies in  
353 decades ago (Fig. 10). The average carbon production rate from the three RUSALCA cruises in the  
354 Chukchi Sea was  $0.33 \text{ g C m}^{-2} \text{ d}^{-1}$  (SD =  $0.40 \text{ g C m}^{-2} \text{ d}^{-1}$ ). In addition, Hill and Cota (2005) reported that  
355 the mean daily production rate during the initial ice breakup was  $0.8 \text{ g C m}^{-2} \text{ d}^{-1}$  in 2002. The daily  
356 production rates obtained by Lee et al. (2012) and Lee et al. (2013) were  $0.54$  and  $0.86 \text{ g C m}^{-2} \text{ d}^{-1}$ ,  
357 respectively (Fig. 10). Even though the different sea ice condition could affect difference in the  
358 productivity, these recent measurements of primary production (Hill and Cota 2005; Lee et al., 2007,  
359 2012 and 2013; Yun et al., 2014; This study) showed significantly lower rates compared with the  
360 previously reported rates from the region (Hameedi, 1978; Korsak, 1992; Zeeman, 1992) (*t*-test,  $p < 0.01$ ).  
361 Moreover, it is obviously shown a decreasing trend of primary production ( $r = 0.7689$ ,  $p < 0.01$ ) (Fig. 10).  
362 This is very interesting because primary production could be expected to increase in the region due to the  
363 increased light availability to the phytoplankton. For example, based on satellite ocean color data, Arrigo  
364 et al. (2008) found large increases in the annual net primary production on the continental shelves of the  
365 Chukchi Sea as well as Siberian and Laptev seas due to increased open water areas and longer growing  
366 seasons. However, the *in situ* measurements of primary production in recent years (Hill and Cota 2005;  
367 Lee et al., 2007, 2012 and 2013; Yun et al., 2014; This study) have shown consistently lower primary  
368 production rates compared to those in previous studies.

369 The strong seasonal and interannual variation in the region could be suggested for reason causing the  
370 low primary production, as discussed above. Hill et al. (2013) found that the seasonal variations in  
371 primary production in the southern Chukchi Sea peaked in July and then progressively declined in August  
372 and September. In fact, the lowest primary production rates given by Yun et al. (2014) were obtained from  
373 the late summer season (i.e., from 1 to 30, September, 2009) compared with the rates found in the present  
374 study (from 30 August to 14 September, 2012) or in Lee et al. (2007) (from 11 to 22 August, 2004). In  
375 comparison, previous studies (Hameedi, 1978; Korsak, 1992; Zeeman, 1992) included the measurements  
376 obtained from July to August (Fig. 10). However, their measurements just starting from the end of July  
377 were mostly done during August (Korsak, 1992; Zeeman, 1992). Although recent measurements from the  
378 three RUSALCA cruises (2004, 2009 and 2012) may not have reflected the highest values (i.e., July) of  
379 primary production, the measurements from Hill and Cota (2005) or Lee et al. (2012 and 2013) include  
380 the values in the mid-July and early August. Therefore, the recent low rates of primary production might  
381 be reflected by decreasing trend rather than results of seasonal and interannual variations.

382 More plausible reason for the recent low primary production in the Chukchi Sea could be due to the  
383 decreased concentrations of nutrients and chlorophyll *a*. According to Whitledge and Lee (unpublished  
384 data), in recent years, there have been significant decreases of 30-50% in nutrient concentrations and  
385 approximately 40% in the integrated chlorophyll *a* concentration in the Bering Strait and the Chukchi Sea.  
386 Based on the significant relationships between primary production and the nutrient and FWC (discussed

387 in section 4.3), the recent decrease in nutrient and chlorophyll *a* concentrations may have been closely  
388 related to the changes in freshwater inputs in the region. According to Serreze et al. (2006), there was  
389 recently larger import of freshwater through the Bering Strait compared with previous estimates.  
390 Therefore, the recent decreases in the concentrations of major inorganic nutrients and chlorophyll *a* may  
391 have resulted in lower primary production rates in the Chukchi Sea.

392 Recently, the freshwater content in the Arctic Ocean, which includes river discharge, pacific water  
393 inflow through the Bering Strait, sea ice melt water and net precipitation (Jones et al., 2008), has  
394 increased over the past few decades. If the increased freshwater content in the Chukchi Sea are  
395 continuously observed, the Chukchi Sea might have become less productive region compared with  
396 previous decades.

397

## 398 **5. Conclusions**

399 This study reported the regional characteristics of primary production in the Chukchi Sea and recent  
400 trend of primary production based on *in situ* measurements. The different nutrient conditions and  
401 phytoplankton biomass could be an important reason for the regional differences in the production rates  
402 of phytoplankton. Based on comparison between previous studies in decades ago and recent  
403 measurements, we found that recent primary production in the Chukchi Sea showed a decreasing trend.  
404 The changes in freshwater inputs in the region may have been closely related to the nutrient and primary




405 production variability. Although Coupel et al. (2015) reported that the recent freshening of the Arctic  
406 Ocean does not significantly affect primary production in the Chukchi shelf based on comparison with  
407 measurements in the deep Canada Basin, our results showed that the freshwater variability in the Chukchi  
408 Sea has had a large influence on the recent changes in primary production by controlling the nutrient  
409 inventory. If the increased freshwater inflow persists, the primary production in the region will  
410 considerably decrease, ultimately resulting in changes in the regional characteristics of primary  
411 production. However, a large interannual variability of primary production remains despite the statistical  
412 significance observed in this study. Therefore, more measurements under various environmental  
413 conditions are needed to better understand the recent variations in the primary production in the Chukchi  
414 Sea. In particular, there could be some changes in the phytoplankton community structures because the  
415 smaller cells benefit more than the larger cells under increased freshening conditions (Li et al., 2009).

416

#### 417 **Acknowledgments**

418 We thank the captain and crew of the *Professor Khromov* for their outstanding assistance during the cruise.  
419 We gratefully acknowledge the physical oceanographers from Woods Hole Oceanographic Institute (M.  
420 Swartz, S. Mills and J. Pietro), who provided the CTD data. We also appreciate NOAA for supporting the  
421 RUSALCA program. This study was financially supported by the “2015 Post-Doc. Development Program”  
422 of Pusan National University, Korea. This work was also supported by grants from the Korea-Polar Ocean



423 in Rapid Transition (K-PORT; PM14040) program funded by the Ministry of Oceans and Fisheries, Korea.

424

425 **References**

426

427 Anderson, L.G. Björk, G., Jutterström, S., Pipko, I., Shakhova, N., Semiletov, I., and Wählström, I.: East

428 Siberian Sea, an Arctic region of very high biogeochemical activity, *Biogeosciences*, 8, 1745–1754,

429 doi:10.5194/bg-8-1745-2011, 2011.

430 Ardyna, M., Babin, M., Gosselin, M., Devred, E., Rainville, L., and Tremblay, J.-É.: Recent Arctic Ocean

431 sea ice loss triggers novel fall phytoplankton blooms, *Geophys. Res. Lett.*, 41, 6207–6212,

432 doi:10.1002/2014GL061047, 2014.

433 Arrigo, K.R., van Dijken, G., and Pabi, S.: Impact of a shrinking Arctic ice cover on marine primary

434 production, *Geophys. Res. Lett.*, 35, L19603, doi:10.1029/2008GL035028, 2008.

435 Carmack, E., McLaughlin, F., Yamamoto-Kawai, M., Itoh, M., Shimada, K., Krishfield, R., and

436 Proshutinsky, A.: Freshwater storage in the Northern Ocean and the special role of the Beaufort

437 Gyre, in: *Arctic-Subarctic Ocean Fluxes: Defining the Role of the Northern Seas in Climate*, edited

438 by: Dickson, R. R., Meincke, J., and Rhines, P., Springer, New York, 145 – 169, 2008.

439 Comiso, J.C., Parkinson, C.L., Gersten, R., and Stock, L.: Accelerated decline in the Arctic sea ice cover,

440 *Geophys. Res. Lett.*, 35, L01703, doi:10.1029/2007GL031972, 2008.

441 Coupel, P., Ruiz-Pino, D., Sicre, M.A., Chen, J.F., Lee, S.H., Schiffrine, N., Li, H.L., and Gascard, J.C.:

442 The impact of freshening on phytoplankton production in the Pacific Arctic Ocean, *Prog. Oceanogr.*,

443 131, 113-125, 2015.

444 Crane, K., and Ostrovskiy, A.: Introduction to the special issue: Russian-American Long-term Census of  
445 the Arctic (RUSALCA). *Oceanography*, 28(3), 18(3), <http://dx.doi.org/10.5670/oceanog.2015.54>,  
446 2015.

447 Dugdale, R.C., and Goering, J.J.: Uptake of new and regenerated forms of nitrogen in primary  
448 productivity, *Limnol. Oceanogr.*, 12, 196-206, 1967.

449 Giles, K.A., Laxon, S.W., Ridout, A.L., Wingham, D.J., and Bacon, S.: Western Arctic Ocean freshwater  
450 storage increased by wind-driven spin-up of the Beaufort Gyre, *Nat. Geosci.*, 5, 194–197, 2012.

451 Gosselin, M., Levasseur, M., Wheeler, P.A., Horner, R.A., and Booth, B.C.: New measurements of  
452 phytoplankton and ice algal production in the Arctic Ocean. *Deep-Sea Res. Part II*, 44, 1623–1644,  
453 1997.

454 Hama, T., Miyazaki, T., Ogawa, Y., Iwakuma, T., Takahashi, M., Otsuki, A., and Ichimura, S.:  
455 Measurement of photosynthetic production of a marine phytoplankton population using a stable <sup>13</sup>C  
456 Isotope, *Mar. Biolo.*, 73, 31-36, 1983.

457 Hameedi, M.J.: Aspects of water column primary productivity in the Chukchi Sea during summer, *Mar.*  
458 *Biol.*, 45, 37–46, 1978.

459 Hansell, D.A., and Goering, J.J.: Pelagic nitrogen flux in the northern Bering Sea, *Cont. Shelf Res.*, 10,  
460 501–519, 1990.

461 Hill, V., and Cota, G.F.: Spatial patterns of primary production on the shelf, slope and basin of the western  
462 Arctic in 2002, *Deep-Sea Res. Part II*, 52, 3344-3354, 2005.

463 Hill, V.J., Matrai, P.A., Olson, E., Suttles, S., Steele, M., Codispoti, L.A., and Zimmerman, R.C.:  
464 Synthesis of integrated primary production in the Arctic Ocean: II. In situ and remotely sensed  
465 estimates, *Prog. Oceanogr.*, 110, 107–125, <http://dx.doi.org/10.1016/j.pocean.2012.11.005>, 2013.

466 Hirawake, T., Shinmyo, K., Fujiwara, A., and Saitoh, S.: Satellite remote sensing of primary productivity  
467 in the Bering and Chukchi Seas using an absorption-based approach, *ICES J. Mar.Sci.*, 69, 1194–  
468 1204, 2012.

469 Jones, E.P., Anderson, L.G., Jutterstrom, S., Mintrop, L., and Swift, J.H.: Pacific freshwater, river water  
470 and sea ice meltwater across Arctic Ocean basins: results from the 2005 Beringia Expedition, *J.*  
471 *Geophys. Res.*, 113, C08012, 2008.

472 Korsak, M.N.: Primary production of organic matter, in: Results of the Third Joint US–USSR Bering and  
473 Chukchi Seas Expedition (BERPAC): Summer 1988, edited by: Nagel, P.A., US Fish and Wildlife  
474 Service, Washington, DC, 215–218, 1992.

475 Kwok, R., Cunningham, G. F., Wensnahan, M., Rigor, I., Zwally, H. J., and Yi, D.: Thinning and volume  
476 loss of Arctic sea ice: 2003–2008, *J. Geophys. Res.*, 114, C07005, doi:10.1029/2009JC005312.,  
477 2009.

478 Lee, S.H., and Whitledge, T.E.: Primary and new production in the deep Canada Basin during summer

479 2002, *Polar Biol.*, 28, 190-197, 2005.

480 Lee, S.H., Whitley, T.E., and Kang, S.H.: Recent carbon and nitrogen uptake rates of phytoplankton in  
481 Bering Strait and the Chukchi Sea, *Cont. Shelf Res.*, 27, 2231-2249, 2007.

482 Lee S.H., Joo, H.M., Liu, Z., Chen, J., and He, J.: Phytoplankton productivity in newly opened waters of  
483 the western Arctic Ocean, *Deep Sea Res. Part II*, 81-84, 18-27, 2012.

484 Lee, S.H., Yun, M.S., Kim, B.K., Saitoh, S., Kang, C.-K., Kang, S.H., and Whitley, T.E.: Latitudinal  
485 carbon productivity in the Bering and Chukchi Seas during the summer in 2007, *Cont. Shelf Res.*, 59,  
486 28-36, 2013.

487 Li, W.K.W., McLaughlin, F.A., Lovejoy, C., and Carmack, E.C.: Smallest algae thrive as the Arctic  
488 Ocean freshens, *Science*, 326,539, doi:10.1126/science.1179798, 2009.

489 McLaughlin, F.A., and Carmack, E.C.: Deepening of the nutricline and chlorophyll maximum in the  
490 Canada Basin interior, 2003-2009, *Geophys. Res. Lett.*, 37, L24602, doi:10.1029/2010GL045459,  
491 2010.

492 McPhee, M.G., Proshutinsky, A., Morison, J.H., Steele, M., and Alkire, M.B.: Rapid change in freshwater  
493 content of the Arctic Ocean, *Geophys. Res. Lett.*, 36, L10602, 2009.

494 Morison, J., Kwok, R., Peralta-Ferriz, C., Alkire, M., Rigor, I., Andersen, R., and Steele, M.: Changing  
495 Arctic Ocean freshwater pathways, *Nature*, 481, 66-70, 2012.

496 Overland, J.E., and Wang, M.: When will the summer Arctic be nearly sea ice free?, *Geophys. Res.*

497 Lett., 40, 2097-2101, DOI: 10.1002/grl.50316, 2013.

498 Parsons, T.R., Maita, Y., and Lalli, C.M.: A manual of chemical and biological methods for seawater  
499 analysis, Pergamon Press, New York, pp 173, 1984.

500 Proshutinsky, A., Krishfield, R., Timmermans, M.-L., Toole, J., Carmack, E., McLaughlin, F., Williams,  
501 W.J., Zimmermann, S., Itoh, M., and Shimada, K.: Beaufort Gyre freshwater reservoir: state and  
502 variability from observation, *J. Geophys. Res.*, 114, doi:10.1029/2008JC005104, 2009.

503 Rabe, B., Karcher, M., Schauer, U., Toole, J.M., Krishfield, R.A., Pisarev, S., Kauker, F., Gerdes, R., and  
504 Kikuchi, T.: An assessment of Arctic Ocean freshwater content changes from the 1990s to the 2006–  
505 2008 period, *Deep-Sea Res. Part I*, 58, 173–185, 2011.

506 Serreze, M.C., Barrett, A.P., Slater, A.G., Woodgate, R.A., Aagaard, K., Lammers, R.B., Steele, M.,  
507 Moritz, R., Meredith, M., and Lee, C.M.: The large-scale freshwater cycle of the Arctic, *J. Geophys.*  
508 *Res.*, 111, C11010, 2006.

509 Shoaf, W.T., and Liem, B.W.: Improved extraction of chlorophyll-a and -b from algae using dimethyl  
510 sulfoxide, *Limnol. Oceanogr.*, 21, 926–928, 1976.

511 Springer, A.M., and McRoy, C.P.: The paradox of pelagic food webs in the northern Bering Sea—III.  
512 Patterns of primary production, *Cont. Shelf Res.*, 13, 575–599, 1993.

513 Stroeve, J., Serreze, M., Drobot, S., Gearheard, S., Holland, M., Maslanik, J., Meier, W., and Scambos, T.:  
514 Arctic sea ice extent plummets in 2007, *EOS Trans., AGU*, 89(2), 13–14, 2008.

515 Stroeve, J. C., Markus, T., Boisvert, L., Miller, J., and Barrett, A.: Changes in Arctic melt season and  
516 implications for sea ice loss, *Geophys. Res. Lett.*, 41, 1216–1225, doi:10.1002/2013GL058951, 2014.

517 Tremblay, J.-E., and Gagnon, J.: The effects of irradiance and nutrient supply on the productivity of  
518 Arctic waters: a perspective on climate change, *Nato Sci Peace Secur*, [http://dx.doi.org/10.1007/978-](http://dx.doi.org/10.1007/978-1-4020-9460-6_7)  
519 [1-4020-9460-6\\_7](http://dx.doi.org/10.1007/978-1-4020-9460-6_7), 73-93, 2009.

520 Tremblay, J.E., Gratton, Y., Fauchot, J., and Price, N.M.: Climatic and oceanic forcing of new, net, and  
521 diatom production in the North Water, *Deep-Sea Res. Part II*, 49, 4927–4946, 2002.

522 Tremblay, J.E., Michel, C., Hobson, K.A., Gosselin, M., and Price, N.M.: Bloom dynamics in early  
523 opening waters of the Arctic Ocean, *Limnol. Oceanogr.*, 51, 900–912, 2006.

524 Wang, J., Cota, G.F., and Comiso, J.C.: Phytoplankton in the Beaufort and Chukchi Seas: Distribution,  
525 dynamics, and environmental forcing, *Deep-Sea Res. Part II*, 52, 3355-3368, 2005.

526 Whitley, T.E., Malloy, S.C., Patton, C.J., and Wirrick, C.D.: Automated nutrient analysis in seawater.  
527 Brookhaven National Laboratory Technical Report BNL 51398, 1981.

528 Woodgate, R.A., Aagaard, K., and Weingartner, T.J.: A year in the physical oceanography of the Chukchi  
529 Sea: Moored measurements from autumn 1990–1991, *Deep-Sea Res. Part II*, 52, 3116–3149,  
530 doi:10.1016/j.dsr2.2005.10.016, 2005a.

531 Woodgate, R.A., Aagaard, K., and Weingartner, T.J.: Monthly temperature, salinity, and transport  
532 variability of the Bering Strait throughflow, *Geophys. Res. Lett.*, 32, L04601,



533 doi:10.1029/2004GL021880, 2005b.

534 Woodgate, R.A., Aagaard, K., Weingartner, T.J.: Interannual changes in the Bering Strait fluxes of  
535 volume heat and freshwater between 1991 and 2004, *Geophys. Res. Lett.*, 33, L15609,  
536 doi:10.1029/2006GL026931, 2006.

537 Woodgate, R.A., Weingartner, T.J., and Lindsay, R.: Observed increases in Bering Strait oceanic fluxes  
538 from the Pacific to the Arctic from 2001 to 2011 and their impacts on the Arctic Ocean water  
539 column, *Geophys. Res. Lett.*, 39, L24603, doi:10.1029/2012GL054092, 2012.

540 Yamamoto-Kawai, M., McLaughlin, F.A., Carmack, E.C., Nishino, S., Shimada, K., and Kurita, N.:  
541 Surface freshening of the Canada Basin, 2003-2007: River runoff versus sea ice meltwater, *J.*  
542 *Geophys. Res.*, 114, C00A05, doi:10.1029/2008JC005000, 2009.

543 Yun, M.S., Whitedge, T.E., Gong, M., and Lee, S.H.: Low primary production in the Chukchi Sea shelf,  
544 2009, *Cont. Shelf Res.*, 76, 1-11, 2014.

545 Zeeman, S.I.: The importance of primary production and CO<sub>2</sub>, in: *Results of the Third Joint US–USSR*  
546 *Bering and Chukchi Seas Expedition (BERPAC): Summer 1988*, edited by: Nagel, P.A., US Fish and  
547 Wildlife Service, Washington, DC, 39–49, 1992.

## Figure Legends

**Fig. 1.** Locations of sampling stations during the 2012 RUSALCA expedition in the Chukchi Sea. The primary production rates were measured at the stations identified by blue circles. The st. CS8R location represents a revisit to st. CS8.

**Fig. 2.** The distribution of surface temperature [ $T_{sur}$ ][°C] (a), surface salinity [ $S_{sur}$ ] (b), stratification index [ $\Delta\sigma_t$ ][kg m<sup>-3</sup>] (c), surface mixed layer depth [ $Z_m$ ][m] (d), and nitracline depth [ $Z_{nit}$ ][m] (e) during the 2012 RUSALCA .

**Fig. 3.** The distributions of the integrated concentrations of ambient nitrite+nitrate [ $NO_2+NO_3$ ][mmol m<sup>-2</sup>] (a), ammonium [ $NH_4$ ][mmol m<sup>-2</sup>] (b), phosphate [ $PO_4$ ][mmol m<sup>-2</sup>] (c), and silicate [ $SiO_4$ ][mmol m<sup>-2</sup>] (d) from surface to 30 m during the 2012 RUSALCA.

**Fig. 4.** The chlorophyll *a* ~~concentrations integrated from surface to~~ content in upper 30 m [mg m<sup>-2</sup>] during the 2012 RUSALCA.

**Fig. 5.** Hourly carbon uptake rates [mg C m<sup>-2</sup> h<sup>-1</sup>] integrated from the surface to 1% light depth during the 2012 RUSALCA.

**Fig. 6** Hourly nitrate and ammonium uptake rates [mg N m<sup>-2</sup> h<sup>-1</sup>] integrated from the surface to 1% light depth during the 2012 RUSALCA.

**Fig. 7.** The distributions of Fresh Water Content (FWC in m) in the Chukchi Sea in 2012 (a), 2009 (b) and 2004 (c).

**Fig. 8.** Relationships between nitrate ~~concentrations-inventory~~ (mmol m<sup>-2</sup>) and chlorophyll *a* ~~concentrations-content~~ (mg m<sup>-2</sup>) ~~integrated from surface to~~ in upper 30 m (a) (n = 41); chlorophyll *a* ~~concentrations-content~~ (mg m<sup>-2</sup>) and daily carbon (g C m<sup>-2</sup> d<sup>-1</sup>) and nitrogen production rate (mg N m<sup>-2</sup> d<sup>-1</sup>)

over the euphotic zones (b) (n = 11). All data obtained during the 2012 RUSALCA.

서식 있음: 강조 없음

**Fig. 9.** Relationships between FWC (m) and nitrate ~~concentrations-inventory~~ (mmol m<sup>-2</sup>) (a); FWC (m) and daily primary production rate (g C m<sup>-2</sup> d<sup>-1</sup>) (b); nitrate ~~concentrations-inventory~~ (mmol m<sup>-2</sup>) and daily primary production rate (g C m<sup>-2</sup> d<sup>-1</sup>) (c). All data obtained from southern region during the three RUSALCA cruises.

**Fig. 10.** A recent trend of primary production based on *in situ* carbon uptake measurements (<sup>13</sup>C or <sup>14</sup>C) in the Chukchi Sea. ~~All Chukchi Sea represents southern and northern Chukchi Sea~~ Study area of each study is indicated by N or S for northern and southern Chukchi Sea, respectively.

Table 1 Location, water depth (m) and euphotic depth ( $Z_{eu}$ ) for primary productivity stations in the Chukchi Sea in 2012.

Region	Station	Date (mm/dd/yr)	Location		Depth (m)	$Z_{eu}$ (m)
			Latitude ( $^{\circ}$ N)	Longitude ( $^{\circ}$ W)		
Northern	CEN4	09/05/12	69.9828	-175.6857	63	34
	CEN1A	09/06/12	70.7085	-178.2988	38	20
	HC2	09/07/12	70.9000	-175.0127	74	36
	HC26	09/08/12	71.7878	-174.3945	55	46
	G12	09/11/12	71.3980	-171.2597	55	46
Southern	CS8	08/30/12	67.4312	-169.6030	51	24
	CS17	09/01/12	68.2983	-167.0418	40	22
	CL5A	09/02/12	68.6407	-170.9423	59	20
	CL3R	09/12/12	69.0048	-168.9000	57	26
	CL8	09/13/12	67.8692	-172.5482	53	24
	CS8R	09/14/12	67.4312	-169.6030	51	26

Table 2 Summary of one-way analysis of variance (ANOVA) for environmental variables in two geographic regions of the Chukchi Sea in 2012. The mean values (ranges in parentheses) and their significant differences (> or <) between northern and southern regions are given for surface temperature ( $T_{sur}$ ), surface salinity ( $S_{sur}$ ), stratification index ( $\Delta\sigma_t$ ), surface mixed layer depth ( $Z_m$ ), nitracline depth ( $Z_{nit}$ ), fresh water content (FWC), nitrite+nitrate ~~concentration~~ inventory of the upper 30 m ( $NO_2+NO_3$ ), ammonium ~~concentration~~ inventory of the upper 30 m ( $NH_4$ ), phosphate ~~concentration~~ inventory of the upper 30 m ( $PO_4$ ), silicate ~~concentration~~ inventory of the upper 30 m ( $SiO_4$ ) and chlorophyll *a* ~~concentration~~ content of in the upper 30 m (Chl-*a*). \*:  $p < 0.05$ , \*\*:  $p < 0.01$ , \*\*\*:  $p < 0.001$ , ns: not significant.  $n = 52$ . Also given are mean and range values for carbon production (CP), nitrate production (NP) and ammonium production (AP) (all  $mg\ C\ or\ N\ m^{-2}\ d^{-1}$ ).  $n=11$ .

서식 있음: 강조 없음

서식 있음: 강조 없음

서식 있음: 강조 없음

서식 있음: 강조 없음

서식 있음: 강조 없음

서식 있음: 강조 없음

Variables	Northern		Southern
$T_{sur}$ ( $^{\circ}C$ )	0.62 (-1.33 ~ 4.13)	<***	3.89 (1.60 ~ 8.53)
$S_{sur}$	29.27 (27.30 ~ 32.04)	>*	27.48 (21.48 ~ 32.35)
$\Delta\sigma_t$ ( $kg\ m^{-3}$ )	3.15 (0.79 ~ 5.34)	<*	4.47 (0.71 ~ 9.71)
$Z_m$ (m)	7.6 (4.0 ~ 14.0)	ns	8.4 (4.0 ~ 14.0)
$Z_{nit}$ (m)	13.0 (2.5 ~ 30.0)	ns	12.6 (2.5 ~ 35.0)
$NO_2+NO_3$ ( $mmol\ m^{-2}$ )	75.01 (21.51 ~ 218.22)	<*	134.15 (21.82 ~ 355.43)
$NH_4$ ( $mmol\ m^{-2}$ )	40.49 (15.36 ~ 86.93)	<***	61.22 (28.54 ~ 109.51)
$PO_4$ ( $mmol\ m^{-2}$ )	22.19 (5.43 ~ 34.26)	ns	25.95 (8.30 ~ 43.57)
$SiO_4$ ( $mmol\ m^{-2}$ )	245.49 (104.79 ~ 800.49)	<***	410.86 (129.17 ~ 669.94)
Chl- <i>a</i> ( $mg/m^{-2}$ )	21.7 (2.2 ~ 69.3)	<*	54.5 (3.1 ~ 286.4)
CP ( $mg\ C\ m^{-2}\ d^{-1}$ )	134.7 (16.3 ~ 280.7)		649.1 (151.3 ~ 1628.9)

NP (mg N m <sup>-2</sup> d <sup>-1</sup> )	6.1 (2.2 ~ 19.9)	69.7 (4.5 ~ 281.6)
AP (mg N m <sup>-2</sup> d <sup>-1</sup> )	38.4 (17.4 ~ 83.6)	96.2 (45.0 ~ 242.4)

