

This discussion paper is/has been under review for the journal Biogeosciences (BG).  
Please refer to the corresponding final paper in BG if available.

# Characterizing Leaf Area Index (LAI) and Vertical Foliage Profile (VFP) over the United States

H. Tang<sup>1</sup>, S. Ganguly<sup>2</sup>, G. Zhang<sup>2</sup>, M. A. Hofton<sup>1</sup>, R. F. Nelson<sup>3</sup>, and R. Dubayah<sup>1</sup>

<sup>1</sup>Department of Geographical Sciences, University of Maryland, College Park, Maryland, USA

<sup>2</sup>Bay Area Environmental Research Institute (BAERI)/NASA Ames Research Center, Moffett Field, California, USA

<sup>3</sup>Biospheric Sciences Branch, Code 618, NASA Goddard Space Flight Center, Greenbelt, Maryland, USA

Received: 4 July 2015 – Accepted: 3 August 2015 – Published: 24 August 2015

Correspondence to: H. Tang (htang@umd.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

**BGD**

12, 13675–13710, 2015

## Characterizing Leaf Area Index (LAI) and Vertical Foliage Profile (VFP)

H. Tang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Abstract

Leaf area index (LAI) and vertical foliage profile (VFP) are among the important canopy structural variables. Recent advances in lidar remote sensing technology have demonstrated the capability of accurately mapping LAI and VFP over large areas. The primary objective of this study was to derive and validate a LAI and VFP product over the contiguous United States using spaceborne waveform lidar data. This product was derived at the footprint level from the Geoscience Laser Altimeter System (GLAS) using a biophysical model. We validated GLAS derived LAI and VFP across major forest biomes using airborne waveform lidar. The comparison results showed that GLAS retrievals of total LAI were generally accurate with little bias ( $r^2 = 0.67$ , bias =  $-0.13$ , RMSE =  $0.75$ ). The derivations of GLAS retrievals of VFP within layers was not as accurate overall ( $r^2 = 0.36$ , bias =  $-0.04$ , RMSE =  $0.26$ ), and these varied as a function of height, increasing from understory to overstory  $-0$  to  $5$  m layer:  $r^2 = 0.04$ , bias =  $0.09$ , RMSE =  $0.31$ ;  $10$  to  $15$  m layer:  $r^2 = 0.53$ , bias =  $-0.08$ , RMSE =  $0.22$ ; and  $15$  to  $20$  m layer:  $r^2 = 0.66$ , bias =  $-0.05$ , RMSE =  $0.20$ . Significant relationships were also found between GLAS LAI products and different environmental factors, in particular elevation and annual precipitation. In summary, our results provide a unique insight into vertical canopy structure distribution across North American ecosystems. This data set is a first step towards a baseline of canopy structure needed for evaluating climate and land use induced forest changes at continental scale in the future and should help deepen our understanding of the role of vertical canopy structure on terrestrial ecosystem processes across varying scales.

## 1 Introduction

Accurate measurements of three dimensional canopy structure and function play a key role in global carbon dynamics, climate feedbacks as well as biodiversity studies (Heimann and Reichstein, 2008; Loreau et al., 2001; Cramer et al., 2001; Schimel

BGD

12, 13675–13710, 2015

## Characterizing Leaf Area Index (LAI) and Vertical Foliage Profile (VFP)

H. Tang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





liage profile as a function of height is beyond the capability of passive optical remote sensing unless multiple look angles are used (Chopping et al., 2009). We argue that spaceborne lidar (light detection and ranging) technology provides a means of overcoming this limitation and of measuring vertical structure even over dense forests.

Lidar has proven effective at measuring three dimensional canopy structural information (Lefsky et al., 2002). Lidar measures the distance between a target and the sensor by the round-trip traveling time of an emitted laser pulse. It allows direct 3-D measurements of canopy structural components, including foliage, branch and trunk which then be used to estimate biophysical variables, such as canopy height and biomass (Drake et al., 2002; Saatchi et al., 2011; Los et al., 2012; Lefsky, 2010; Simard et al., 2011; Asner et al., 2012; Baccini et al., 2012; Strahler et al., 2008), as well as LAI and VFP (Morsdorf et al., 2006; Tang et al., 2012; Zhao et al., 2013).

Garcia et al. (2012) and Luo et al. (2013) demonstrated the possibility of deriving LAI and VFP data across different landscapes from Geoscience Laser Altimeter System (GLAS) on board of Ice, Cloud and land Elevation Satellite (ICESat). Tang et al. (2014a) derived LAI and VFP data from GLAS data, but using a physically based model rather than an empirical methodology. The use of a physical model greatly simplified application over large areas because site specific, statistical calibrations were not required. Further improvement of the model led to a GLAS LAI and VFP product over the entire state of California, USA (Tang et al., 2014b). However, there is still a need to further examine the relationship between vertical foliage distribution and lidar waveforms over even broader areas. Assessment of their relationship across different forest types and environmental gradients will not only strengthen our confidence in acquiring a potential global LAI and VFP measurement, but will also provide guidance on the design and science definition of future lidar missions such as the Global Ecosystem Dynamics Investigation (GEDI) (Dubayah et al., 2014).

The objective of this study is to characterize the continental scale variability of canopy structure across the United States using lidar observations from space. First, we implement our existing algorithm at the GLAS footprint level and compare the derived data

## BGD

12, 13675–13710, 2015

### Characterizing Leaf Area Index (LAI) and Vertical Foliage Profile (VFP)

H. Tang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



with LAI and VFP products from airborne lidar in different forest types. Next we map the aggregated LAI and VFP product according to different ecoregions and land cover types over CONUS. Finally we analyze the distribution of GLAS LAI across different environmental factors, including elevation and precipitation.

## 2 Methods

### 2.1 GLAS data

GLAS is a spaceborne, sampling waveform lidar sensor with the working wavelength in the near-infrared band (1064 nm). It emits laser pulses at a frequency of 40 Hz and records the energy reflected from both the ground surface and canopy in an approximately 65 m diameter footprint (Abshire et al., 2005). GLAS samples the Earth surface in transects with individual footprints separated by  $\sim 175$  m along track, and with between track spacing that varies as a function of latitude (e.g. 30 km spacing between tracks at the equator and 5 km spacing at  $80^\circ$  latitude (Brenner et al., 2012)). As a result of this sampling pattern, GLAS does not provide a wall-to-wall observation of forests. Its spatial allocation of laser footprints is best defined as a pseudo-systematic sampling or cluster sampling strategy (Stahl et al., 2011; Healey et al., 2012). To obtain a spatially continuous estimate of LAI at continental scale, footprint level GLAS data would need to be extrapolated using other remote sensing data (Dubayah et al., 2008; Lefsky, 2010), or can be mapped into appropriate geographic strata such as land cover types or ecoregions.

### 2.2 Retrieval of GLAS LAI and VFP

We collected a total of 1 100 498 cloud-free GLAS data from Campaigns GLA01 and GLA14 data over the contiguous United States from 2003 to 2007. GLA01 included the complete recorded waveform at a vertical resolution of 15 cm for land surface products, and GLA14 products were comprised of geographical information and various

## Characterizing Leaf Area Index (LAI) and Vertical Foliage Profile (VFP)

H. Tang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



parameters calculated from the waveform (Harding and Carabajal, 2005). Low energy shots (peak energy < 0.5 Volt) were excluded from data process for retrieval quality control because those waveforms were susceptible to noise contamination. Shots during leaf-off season (November to March) were also filtered out over deciduous forests and mixed forests. LAI and its profiles (0.15 m at vertical resolution) were initially calculated for GLAS footprints based on a Geometric Optical and Radiative Transfer (GORT) model (Ni-Meister et al., 2001), and further corrected for slope effects using an iterative method (Tang et al., 2014a). Canopy VFP were calculated from integration of footprint level LAI profiles at height intervals of 0 to 5, 5 to 10, 10 to 15 and 15 to 20 m.

### 2.3 Comparison data sets

We validated LAI and VFP data sets using an airborne lidar system, LVIS (Laser Vegetation Imaging Sensor). LVIS is a medium resolution (~ 25 m diameter) waveform scanning lidar system designed by NASA Goddard Space Flight Center (GSFC) (Blair et al., 1999). It can image the terrestrial surface across a 2 km wide swath and has been deployed to map many different forest structural parameters at regional scales across diverse biomes (Tang et al., 2012; Drake et al., 2002; Swatantran et al., 2012). We calculated both total LAI and VFP at 5 m height intervals from existing LVIS data using our physically based model, which has been validated using different types of field measurements (destructive sampling, LAI-2000 and hemispherical photos) (Tang et al., 2012, 2014a; Zhao et al., 2013). LVIS data used in this study included major forest types from eastern, central and western US, including Maine forests just north of Orono, Maine (2003), Sierra National Forest in California (2008), mixed forests along Baltimore/Washington corridor (2003) and the White River National Wildlife Refuge in Arkansas (2006). These LVIS datasets were all collected during leaf-on season.

We also included a 30 m resolution Landsat LAI map to examine the spatial distribution of GLAS total LAI. Landsat has the longest earth observation history at moderate resolution (30 m), and for decades has provided a consistent and unique measure-

BGD

12, 13675–13710, 2015

## Characterizing Leaf Area Index (LAI) and Vertical Foliage Profile (VFP)

H. Tang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ment of terrestrial ecosystems. The Landsat LAI map was produced using Global Land Survey (GLS) 2005 orthorectified Landsat data (Ganguly et al., 2012).

## 2.4 Analysis

The comparison between LVIS and GLAS was performed at the GLAS footprint level. LVIS shots falling within a 32.5 m radius from a GLAS shot center were selected. We filtered GLAS footprints to have a minimum of 3 coincident LVIS shots to increase the likelihood that the LVIS data covered a sufficient portion of the larger GLAS footprints. Both LAI and the 5 m interval VFP of LVIS shots were averaged onto each coincident GLAS footprint for comparison. We also made a footprint level comparison between GLAS LAI and the Landsat LAI map. A 3 × 3 window was applied to each GLAS footprint center to extract the averaged Landsat LAI pixels. Pixels with invalid values (e.g. retrieval failure or non-vegetation pixel) were excluded in the comparison. Agreements of different LAI datasets were assessed by coefficient of determination, bias and RMSE (Root Mean Square Error):

$$\text{bias} = \sum_{i=1}^n \frac{\text{GLAS}_i - \text{Ref}_i}{n} \quad (1)$$

$$\text{RMSE} = \sqrt{\frac{\sum_{i=1}^n (\text{GLAS}_i - \text{Ref}_i)^2}{n}} \quad (2)$$

In Eqs. (1) and (2),  $\text{GLAS}_i$  is GLAS LAI (or VFP) value at footprint level and  $\text{Ref}_i$  is that extracted from LVIS or Landsat.

Next, we aggregated the footprint level GLAS data into terrestrial ecoregions based on subset of a global map (Olson et al., 2001). Statistical analysis of total LAI and LAI strata (VFP aggregated at every 10 m height interval) was performed subsequently for each ecoregion. We also analyzed the GLAS LAI and VFP distribution across different environmental gradients throughout CONUS. GLAS footprints were categorized according to different environmental factors, including vegetation type, topographic data

BGD

12, 13675–13710, 2015

## Characterizing Leaf Area Index (LAI) and Vertical Foliage Profile (VFP)

H. Tang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





bias =  $-0.08$ , RMSE = 0.22), reaching a maximum at the top of the canopy (15 to 20 m,  $r^2 = 0.66$ , bias =  $-0.05$ , RMSE = 0.20).

The comparison between Landsat LAI and GLAS LAI had a much lower agreement than that of LVIS ( $r^2 = 0.18$ , bias = 0.18 and RMSE = 2.02) (Fig. 4). Even though the two data sets agreed well at lower LAI values, Landsat overestimated LAI at the middle range (from LAI values of 1 to 3) and then saturated above a value of about 4 to 5 against GLAS data.

### 3.2 Aggregated GLAS LAI and VFP within Ecoregions

We next mapped GLAS LAI across US ecoregions (Fig. 5). Highest LAI values were found along northern Pacific Coast while lowest values occurred in the basin and range province and the arid rains shadow region east of the Rocky Mountains. Northern California coastal forests (Pacific temperate rainforests) were found to have the highest mean LAI value of 5.24. In the eastern US, the mixed deciduous forests of the Appalachian–Blue Ridge province had the highest value of 3.95 while other ecogreions around north–south direction of Appalachian Mountains had similar LAI values around 3 ~ 4 (Table 1). Forest ecogreions with lowest LAI values (excluding desert, shrubland and grassland) were located in Arizona mountains forests (1.15) and Great Basin montane forests (0.90). Differences between these ecoregion-level LAI were significant based on a bonferroni adjusted  $t$  test, except for those among Willamette Valley forests, Appalachian–Blue Ridge forests, Puget lowland forests and Appalachian mixed mesophytic forests ( $p$  values  $> 0.05$ ).

LAI strata formed by VFP at each 10 m height interval were also averaged and mapped across the US (Fig. 6). We chose the 10 m height interval rather than that of 5 m because LAI strata aggregated at 10 m height interval represented a more accurate and reliable description of vertical canopy structure given the relatively lower measurement accuracy in the understory ( $< 5$  m) we found in comparison to LVIS data. Each strata showed a generally similar geographic pattern as that of total LAI with the decreasing trend from coast to interior lands, but the specific patterns among strata

BGD

12, 13675–13710, 2015

## Characterizing Leaf Area Index (LAI) and Vertical Foliage Profile (VFP)

H. Tang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



differed. Northwestern forests were observed to have the highest total LAI values as well as LAI strata values. Northern California coastal forests exhibited the largest total LAI value as well as highest foliage density under 20 m height, while British Columbia mainland coastal forests showed the highest foliage density (1.13) above 20 m height with a lower total LAI value (4.74).

The distribution of GLAS total LAI and profiles were examined across different land cover types (Fig. 7 and 8). Not surprisingly, forests were found to have a consistently greater value than non-forest biomes in both total LAI and its strata. For example, deciduous broadleaf forests had the highest value of total LAI (mean = 4.03) as well as that of middle and upper LAI strata (height > 10 m), while open shrubland showed the lowest total LAI values of 0.77. However, vertical LAI distributions of most forests and non-forests were similar with peak foliage density distributed around a height of 2 ~ 4 m. The only exception was deciduous broadleaf forest, of which most of leaves were distributed at middle-story level with a peak height at about 8 m. Its VFP values did not decrease significantly until reaching a height of 15 m.

### 3.3 GLAS LAI Distributions by Environmental Factors

A linear regression analysis between GLAS LAI and SRTM DEM showed that increasing altitude led to an overall decreasing, but non-monotonic, trend in LAI values ( $LAI = 3.60 - 0.686 \times \text{Elevation (km)}$ ,  $r^2 = 0.59$ , all  $P < 0.01$ ) (Fig. 9). GLAS LAI values increased with DEM at the elevation range from 0 to 750 m and 2000 to 3000 m. The variation in the LAI-DEM relationship agreed well with Forest Ratio ( $LAI = 0.112 + 3.18 \times \text{Forest Ratio}$ ,  $r^2 = 0.45$ ,  $P < 0.01$ ). Here Forest Ratio was, defined as the percentage of footprints classified as forests in total GLAS shots (forest and non-forest). A multiple linear regression analysis showed that about 87% of total variance could be explained by a simple combination of elevation groups and Forest Ratio values:  $LAI = 2.59 \times \text{Forest Ratio} - 0.595 \times \text{Elevation (km)} + 1.58$ .

We also analyzed GLAS LAI by 30 yr normal annual climate data using linear regression models (Fig. 10). It was observed that increasing precipitation signifi-

BDG

12, 13675–13710, 2015

## Characterizing Leaf Area Index (LAI) and Vertical Foliage Profile (VFP)

H. Tang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



cantly increased LAI values ( $\Delta\text{LAI} = 1.84$  per 1000 mm precipitation increase) but only at low and moderate precipitation levels ( $< 2400$  mm):  $\text{LAI} = 1.84 \times \text{precipitation (mm)} \times 10^{-3} + 0.774$ ,  $r^2 = 0.96$ ,  $\text{adj} - r^2 = 0.95$ ,  $P < 0.01$ . It contributed little when exceeding that threshold ( $\text{LAI} = 0.22 \times \text{precipitation (mm)} \times 10^{-3}$ ,  $r^2 = 0.40$ ,  $\text{adj} - r^2 = 0.30$ ,  $P = 0.09$ ), as we found no significant LAI increase among groups greater than 2400 mm using a bonferroni adjusted  $t$  test. GLAS LAI was also negatively but slightly correlated with minimum (maximum) vapor pressure deficit with a Pearson's correlation coefficient of  $-0.29$  ( $-0.15$ ). The correlation coefficients between GLAS LAI and annual mean/minimum/maximum temperature were even lower with values of 0.13, 0.18 and 0.08 respectively.

Finally, we applied multiple linear regression analysis to illustrate the combined environmental effects of altitude and precipitation the on distributions of LAI and VFP. The regression analyses were conducted at both GLAS footprint level and aggregated scale on altitude and precipitation groups. At footprint level, altitude and precipitation together explained about 30 % of variance of total LAI ( $\text{LAI} = 2.73 - 0.69 \times \text{Elevation (km)} + 0.58 \times \text{precipitation (mm)} \times 10^{-3}$ ,  $r^2 = 0.29$ ,  $\text{adj} - r^2 = 0.29$ ,  $P < 0.01$ ). However, their correlations with footprint level VFP (0–10 m, 10–20 m and  $> 20$  m height intervals) were not significant with  $r^2$  of 0.07, 0.12 and 0.08 respectively. At the aggregated scale, there was a better relationship between averaged LAI (VFP) values and environmental factors. The combination of altitude and precipitation can explain more than 60 % variance in both total LAI and VFP, but explains only about 36 % of variance on LAI for canopies less than 10 m height.

## 4 Discussion

In this study, we generated GLAS estimates of LAI and VFP across the United States, validated with an airborne lidar sensor, LVIS. Comparisons between LVIS and GLAS LAI and VFP estimates in different forest types across the United States show that GLAS generally provides accurate LAI and VFP estimates at footprint level. Consid-

# BGD

12, 13675–13710, 2015

## Characterizing Leaf Area Index (LAI) and Vertical Foliage Profile (VFP)

H. Tang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





## BGD

12, 13675–13710, 2015

## Characterizing Leaf Area Index (LAI) and Vertical Foliage Profile (VFP)

H. Tang et al.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)




[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


each other at all along the entire waveform due to the offset. However, integration at every 5 m creates a signal overlap of 4 m in each height bin with a maximum of 20 % measurement error. Thus there is ultimately a tradeoff between vertical resolution and VFP accuracy. It also explains the higher agreement of total LAI (essentially an integration of VFP over the entire canopy) in the comparison between LVIS and GLAS. Lastly, measurement of near-ground understory vegetation by GLAS is difficult. By default GLAS waveforms are processed by a Gaussian decomposition method to get an approximate fit comprised of a series of Gaussian functions where the last one usually represents the ground (Hofton et al., 2000). The upper tail of the ground Gaussian peak may be mixed with signals from lower understory, and their separation is problematic, especially over slopes. All of these factors, plus the nature of high complexity and heterogeneity in canopy understory (Aubin et al., 2000; Valladares and Niinemets, 2008), may help explain the lower agreement on understory VFP between LVIS and GLAS.

Comparison between GLAS and Landsat displayed a much lower agreement than that of LVIS, was somewhat biased, and showed clear signals of saturation beyond LAI values of about 5. This result, along with all previous studies (Tang et al., 2012, 2014b), clearly showed the non-saturation advantage of lidar data against passive remote sensing in observing high LAI forests. On the low end of LAI spectrum, GLAS values were lower as compared with Landsat. There are different factors (some in the LVIS comparison too) could possibly lead to their difference in LAI estimates such as geolocation errors of GLAS shots, observation scale difference (65 m vs. 30 m) and misclassifications from MODIS land cover types (mainly impacting the correction of clumping effect). But this underestimation should be largely due to the fact that GLAS may not be able to adequately capture LAI values of short grassland with limited vertical structure or areas of sparse canopy cover, whereas Landsat is able measure such areas based on their total spectral response (tree and grass).

Analysis of GLAS LAI and VFP across ecoregions displayed a reasonable and expected geographical distribution. The great advantage of lidar based estimates is that they can produce LAI vertical strata maps, providing a view of canopy variability across

ecosystem types over large areas. Specifically, we can identify the foliage concentrations at various vertical bins and at spatial resolutions of interest (Figs. 6 and 8 and Table 1). This approach may reduce errors that arise from assumptions of uniformly distributed foliage within canopy, and could potentially be a contribution towards continental scale ecological and biological studies of forest structure and dynamics.

LAI and VFP also varied across different landscapes represented by various land cover types. As expected, we found both total LAI and maximum value of foliage density significantly increase along the vegetation gradient described by the transition from shrubland to savanna to woody savanna to forests (Figs. 7 and 8). In particular, we found deciduous broadleaf forest showing a different pattern with its foliage more evenly distributed in understory and mid-story when compared with all other forests. Our results suggest the existence of canopy layering, and highlight the feasibility of quantifying these layers across landscapes (Whitehurst et al., 2013). Regardless, of whether the data are conceptualized as layers or as continuously varying profiles, they nonetheless provide the actual vertical structure, and thus should help refine current empirical assumptions about vegetation structure of different land cover types in current LAI inversion algorithms (e.g. MODIS) and in ecosystem models (Hurtt et al., 2010; Antonarakis et al., 2014).

Elevation and precipitation were found to be significantly correlated with LAI at both footprint level and across aggregated groupings by elevation and forest ratio. LAI decreased with elevation and this trend was consistent with previous studies (Luo et al., 2004; Moser et al., 2007; Pfeifer et al., 2012). Variations of the trend can be largely explained (about 45% of total variance) by the Forest Ratio (defined in Sect. 3.3). A combination of the two factors (elevation groups and Forest Ratio) explained almost 90% variance of average LAI spatial distribution. We also found a significant but nonlinear relationship between GLAS LAI and annual precipitation (Fig. 10). This non-linear relationship agrees with previous studies in the tropics (Pfeifer et al., 2014; Spracklen et al., 2012). However, we found no significant variation of GLAS LAI with either temperature or vapor pressure deficit variables. A combined effect of elevation and precipitation

## BGD

12, 13675–13710, 2015

### Characterizing Leaf Area Index (LAI) and Vertical Foliage Profile (VFP)

H. Tang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



explains about 30 % of LAI variation at GLAS footprint level, suggesting the natural complexity highly spatial variability of LAI distribution.

As a direct quantification of 3-D foliage distribution, GLAS LAI profiles are thus far the best representations of terrestrial ecosystem structure over broad geographical areas and suggest that ecological applications of these profiles are worth exploring. First, this data could refine large scale modeling of plant respiration and photosynthesis and consequently improve ecosystem modeling (Houghton, 2007). Previous studies have reported a potential 50 % underestimate of GPP values when vertical foliage stratification is not considered (Kotchenova et al., 2004; Sprintsin et al., 2012). A consistent, global data set of VFP should thus improve initialization of ecological models (Hurtt et al., 2004), and refine estimation of GPP, in conjunction with passive remote sensing data (Turner et al., 2006). Secondly, these profiles may be important descriptors of habitat as related to biodiversity and habitat quality. Many studies have confirmed the general relationship between species richness, habitat heterogeneity and forest structural complexity across different landscapes (Swatantran et al., 2012; Goetz et al., 2010; Schut et al., 2014; Ferger et al., 2014). The inclusion of LAI profiles provides spatially explicit vegetation structure data and may potentially improve current observations of species distribution at continental scale, e.g. for avian species (Sauer et al., 2008; Culbert et al., 2013), and lead to entirely new biodiversity metrics (e.g. see Huang et al., 2014). For example the concept of an “edge” has been traditionally defined as the boundary between forest and non-forest areas. LAI profiles provide a means of defining new edges based on differences in LAI as a function of height, so the edge is now the boundary between a rapid change in foliage density at a particular height.

## 5 Conclusion

Accurate representation of canopy vertical structure and its dynamics has long been recognized as a priority because it represents a key interface between terrestrial sur-

**BGD**

12, 13675–13710, 2015

## Characterizing Leaf Area Index (LAI) and Vertical Foliage Profile (VFP)

H. Tang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





## References

- Abdalati, W., Zwally, H. J., Bindschadler, R., Csatho, B., Farrell, S. L., Fricker, H. A., Harding, D., Kwok, R., Lefsky, M., Markus, T., Marshak, A., Neumann, T., Palm, S., Schutz, B., Smith, B., Spinhirne, J., and Webb, C.: The ICESat-2 laser altimetry mission, *Proc. IEEE*, 98, 735–751, 2009.
- Aber, J. D.: Foliage-height profiles and succession in northern hardwood forests, *Ecology*, 60, 18–23, 1979.
- Abshire, J. B., Sun, X., Riris, H., Sirota, J. M., McGarry, J. F., Palm, S., Yi, D., and Liiva, P.: Geoscience Laser Altimeter System (GLAS) on the icesat mission: on-orbit measurement performance, *Geophys. Res. Lett.*, 32, L21S02, doi:10.1029/2005GL024028, 2005.
- Abuelgasim, A. A., Fernandes, R. A., and Leblanc, S. G.: Evaluation of national and global lai products derived from optical remote sensing instruments over canada, *IEEE T. Geosci. Remote*, 44, 1872–1884, doi:10.1109/Tgrs.2006.874794, 2006.
- Antonarakis, A. S., Munger, J. W., and Moorcroft, P. R.: Imaging spectroscopy- and lidar-derived estimates of canopy composition and structure to improve predictions of forest carbon fluxes and ecosystem dynamics, *Geophys. Res. Lett.*, 41, 2535–2542, doi:10.1002/2013gl058373, 2014.
- Asner, G. P., Knapp, D. E., Boardman, J., Green, R. O., Kennedy-Bowdoin, T., Eastwood, M., Martin, R. E., Anderson, C., and Field, C. B.: Carnegie airborne observatory-2: increasing science data dimensionality via high-fidelity multi-sensor fusion, *Remote Sens. Environ.*, 124, 454–465, doi:10.1016/j.rse.2012.06.012, 2012.
- Asner, G. P., Mascaro, J., Anderson, C., Knapp, D. E., Martin, R. E., Kennedy-Bowdoin, T., van Breugel, M., Davies, S., Hall, J. S., Muller-Landau, H. C., Potvin, C., Sousa, W., Wright, J., and Bermingham, E.: High-fidelity national carbon mapping for resource management and redd +, *Carbon Balance and Management*, 8, 7, doi:10.1186/1750-0680-8-7, 2013.
- Aubin, I., Beaudet, M., and Messier, C.: Light extinction coefficients specific to the understory vegetation of the southern boreal forest, quebec, *Can. J. Forest Res.*, 30, 168–177, doi:10.1139/cjfr-30-1-168, 2000.
- Baccini, A., Goetz, S. J., Walker, W. S., Laporte, N. T., Sun, M., Sulla-Menashe, D., Hackler, J., Beck, P. S. A., Dubayah, R., Friedl, M. A., Samanta, S., and Houghton, R. A.: Estimated carbon dioxide emissions from tropical deforestation improved by carbon-density maps, *Nature Climate Change*, 2, 182–185, doi:10.1038/Nclimate1354, 2012.

### Characterizing Leaf Area Index (LAI) and Vertical Foliage Profile (VFP)

H. Tang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Characterizing Leaf Area Index (LAI) and Vertical Foliage Profile (VFP)

H. Tang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Baret, F., Hagolle, O., Geiger, B., Bicheron, P., Miras, B., Huc, M., Berthelot, B., Nino, F., Weiss, M., Samain, O., Roujean, J. L., and Leroy, M.: LAI, fapar and fcover cyclopes global products derived from vegetation – Part 1: Principles of the algorithm, *Remote Sens. Environ.*, 110, 275–286, doi:10.1016/j.rse.2007.02.018, 2007.

5 Baret, F., Weiss, M., Lacaze, R., Camacho, F., Makhmara, H., Pacholczyk, P., and Smets, B.: Geov1: LAI and fapar essential climate variables and fcover global time series capitalizing over existing products. Part1: Principles of development and production, *Remote Sens. Environ.*, 137, 299–309, doi:10.1016/j.rse.2012.12.027, 2013.

10 Blair, J. B., Rabine, D. L., and Hofton, M. A.: The Laser Vegetation Imaging Sensor (LVIS): a medium-altitude, digitization-only, airborne laser altimeter for mapping vegetation and topography, *ISPRS J. Photogramm.*, 54, 115–122, 1999.

15 Brenner, A. C., Zwally, H. J., Bentley, C. R., Csatho, B. M., Harding, D. J., Hofton, M. A., Minster, J.-B., Roberts, L., Saba, J. L., and Thomas, R. H.: The Algorithm Theoretical Basis Document for the Derivation of Range and Range Distributions from Laser Pulse Waveform Analysis for Surface Elevations, Roughness, Slope, and Vegetation Heights, technical report NASA/TM-2012-208641/Vol 7, GSFC.TM.7299.2012, Goddard Space Flight Center, Greenbelt, MD, USA, 2012.

20 Chopping, M., Nolin, A., Moisen, G. G., Martonchik, J. V., and Bull, M.: Forest canopy height from the multiangle imaging spectroradiometer (MISR) assessed with high resolution discrete return lidar, *Remote Sens. Environ.*, 113, 2172–2185, doi:10.1016/j.rse.2009.05.017, 2009.

Clark, D. B. and Kellner, J. R.: Tropical forest biomass estimation and the fallacy of misplaced concreteness, *J. Veg. Sci.*, 23, 1191–1196, doi:10.1111/j.1654-1103.2012.01471.x, 2012.

25 Cramer, W., Bondeau, A., Woodward, F. I., Prentice, I. C., Betts, R. A., Brovkin, V., Cox, P. M., Fisher, V., Foley, J. A., Friend, A. D., Kucharik, C., Lomas, M. R., Ramankutty, N., Sitch, S., Smith, B., White, A., and Young-Molling, C.: Global response of terrestrial ecosystem structure and function to CO<sub>2</sub> and climate change: results from six dynamic global vegetation models, *Glob. Change Biol.*, 7, 357–373, doi:10.1046/j.1365-2486.2001.00383.x, 2001.

30 Culbert, P. D., Radeloff, V. C., Flather, C. H., Kellendorfer, J. M., Rittenhouse, C. D., and Pidgeon, A. M.: The influence of vertical and horizontal habitat structure on nationwide patterns of avian biodiversity, *Auk*, 130, 656–665, doi:10.1525/auk.2013.13007, 2013.

## Characterizing Leaf Area Index (LAI) and Vertical Foliage Profile (VFP)

H. Tang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Deng, F., Chen, J. M., Plummer, S., Chen, M. Z., and Pisek, J.: Algorithm for global leaf area index retrieval using satellite imagery, *IEEE T. Geosci. Remote*, 44, 2219–2229, doi:10.1109/Tgrs.2006.872100, 2006.

Drake, J. B., Dubayah, R. O., Clark, D. B., Knox, R. G., Blair, J. B., Hofton, M. A., Chazdon, R. L., Weishampel, J. F., and Prince, S. D.: Estimation of tropical forest structural characteristics using large-footprint lidar, *Remote Sens. Environ.*, 79, 305–319, doi:10.1016/S0034-4257(01)00281-4, 2002.

Dubayah, R., Bergen, K., Hall, F., Hurtt, G., Houghton, R., Kelldorfer, J., Lefsky, M., Moorcroft, P., Nelson, R., and Saatchi, S.: Global vegetation structure from nasa's desdyni mission: an overview, in: *AGU Fall Meeting Abstracts*, 01, San Francisco, 15 December, B31H-01, 2008.

Dubayah, R., Goetz, S., Blair, J. B., Fatoyinbo, T., Hansen, M., Healey, S., Hofton, M., Hurtt, G., Kellner, J. R., Luthcke, S. B., and Swatantran, A.: The global ecosystem dynamics investigation, in: *American Geophysical Union, Fall Meeting 2014*, San Francisco, 15 December, U14A-07, 2014.

Ferger, S. W., Schleuning, M., Hemp, A., Howell, K. M., and Böhning-Gaese, K.: Food resources and vegetation structure mediate climatic effects on species richness of birds, *Global Ecol. Biogeogr.*, 23, 541–549, doi:10.1111/geb.12151, 2014.

Friedl, M. A., Sulla-Menashe, D., Tan, B., Schneider, A., Ramankutty, N., Sibley, A., and Huang, X. M.: Modis collection 5 global land cover: algorithm refinements and characterization of new datasets, *Remote Sens. Environ.*, 114, 168–182, doi:10.1016/j.rse.2009.08.016, 2010.

Ganguly, S., Nemani, R. R., Zhang, G., Hashimoto, H., Milesi, C., Michaelis, A., Wang, W. L., Votava, P., Samanta, A., Melton, F., Dungan, J. L., Vermote, E., Gao, F., Knyazikhin, Y., and Myneni, R. B.: Generating global leaf area index from landsat: algorithm formulation and demonstration, *Remote Sens. Environ.*, 122, 185–202, doi:10.1016/j.rse.2011.10.032, 2012.

Garcia, M., Popescu, S., Riano, D., Zhao, K., Neuenschwander, A., Agca, M., and Chuvieco, E.: Characterization of canopy fuels using ICESat/GLAS data, *Remote Sens. Environ.*, 123, 81–89, doi:10.1016/j.rse.2012.03.018, 2012.

Goetz, S., Steinberg, D., Dubayah, R., and Blair, B.: Laser remote sensing of canopy habitat heterogeneity as a predictor of bird species richness in an eastern temperate forest, USA, *Remote Sens. Environ.*, 108, 254–263, doi:10.1016/j.rse.2006.11.016, 2007.

## Characterizing Leaf Area Index (LAI) and Vertical Foliage Profile (VFP)

H. Tang et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Goetz, S. J., Steinberg, D., Betts, M. G., Holmes, R. T., Doran, P. J., Dubayah, R., and Hofton, M.: Lidar remote sensing variables predict breeding habitat of a neotropical migrant bird, *Ecology*, 91, 1569–1576, doi:10.1890/09-1670.1, 2010.

Gower, S. T. and Norman, J. M.: Rapid estimation of leaf area index in conifer and broad-leaf plantations, *Ecology*, 72, 1896–1900, 1991.

Harding, D. J. and Carabajal, C. C.: ICESat waveform measurements of within-footprint topographic relief and vegetation vertical structure, *Geophys. Res. Lett.*, 32, L21s10, doi:10.1029/2005gl023471, 2005.

Healey, S. P., Patterson, P. L., Saatchi, S., Lefsky, M. A., Lister, A. J., and Freeman, E. A.: A sample design for globally consistent biomass estimation using lidar data from the Geoscience Laser Altimeter System (GLAS), *Carbon Balance and Management*, 7, 10, doi:10.1186/1750-0680-7-10, 2012.

Heimann, M. and Reichstein, M.: Terrestrial ecosystem carbon dynamics and climate feedbacks, *Nature*, 451, 289–292, doi:10.1038/Nature06591, 2008.

Hofton, M. A., Minster, J. B., and Blair, J. B.: Decomposition of laser altimeter waveforms, *IEEE T. Geosci. Remote*, 38, 1989–1996, 2000.

Houghton, R. A.: Balancing the global carbon budget, *Annu. Rev. Earth Pl. Sc.*, 35, 313–347, doi:10.1146/annurev.earth.35.031306.140057, 2007.

Huang, Q. Y., Swatantran, A., Dubayah, R., and Goetz, S. J.: The influence of vegetation height heterogeneity on forest and woodland bird species richness across the united states, *Plos One*, 9, e103236, doi:10.1371/journal.pone.0103236, 2014.

Hurtt, G. C., Dubayah, R., Drake, J., Moorcroft, P. R., Pacala, S. W., Blair, J. B., and Fearon, M. G.: Beyond potential vegetation: combining lidar data and a height-structured model for carbon studies, *Ecol. Appl.*, 14, 873–883, 2004.

Hurtt, G. C., Fisk, J., Thomas, R. Q., Dubayah, R., Moorcroft, P. R., and Shugart, H. H.: Linking models and data on vegetation structure, *J. Geophys. Res.-Biogeo.*, 115, G00e10, doi:10.1029/2009jg000937, 2010.

Kotchenova, S. Y., Song, X. D., Shabanova, N. V., Potter, C. S., Knyazikhin, Y., and Myneni, R. B.: Lidar remote sensing for modeling gross primary production of deciduous forests, *Remote Sens. Environ.*, 92, 158–172, doi:10.1016/j.rse.2004.05.010, 2004.

Lee, S., Ni-Meister, W., Yang, W. Z., and Chen, Q.: Physically based vertical vegetation structure retrieval from icesat data: validation using LVIS in white mountain national forest, New



## Characterizing Leaf Area Index (LAI) and Vertical Foliage Profile (VFP)

H. Tang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



from year one of modis data, *Remote Sens. Environ.*, 83, 214–231, doi:10.1016/S0034-4257(02)00074-3, Pii S0034-4257(02)00074-3, 2002.

Ni-Meister, W., Jupp, D. L. B., and Dubayah, R.: Modeling lidar waveforms in heterogeneous and discrete canopies, *IEEE T. Geosci. Remote*, 39, 1943–1958, 2001.

5 Nilson, T.: A theoretical analysis of the frequency of gaps in plant stands, *Agr. Meteorol.*, 8, 25–38, 1971.

Olson, D. M., Dinerstein, E., Wikramanayake, E. D., Burgess, N. D., Powell, G. V. N., Underwood, E. C., D'amico, J. A., Itoua, I., Strand, H. E., Morrison, J. C., Loucks, C. J., Allnutt, T. F., Ricketts, T. H., Kura, Y., Lamoreux, J. F., Wettengel, W. W., Hedao, P., and Kassem, K. R.: Terrestrial ecoregions of the world: a new map of life on earth: a new global map of terrestrial ecoregions provides an innovative tool for conserving biodiversity, *BioScience*, 51, 933–938, doi:10.1641/0006-3568(2001)051[0933:teotwa]2.0.co;2, 2001.

10 Park, T., Kennedy, R. E., Choi, S. H., Wu, J. W., Lefsky, M. A., Bi, J., Mantooth, J. A., Myrneni, R. B., and Knyazikhin, Y.: Application of physically-based slope correction for maximum forest canopy height estimation using waveform lidar across different footprint sizes and locations: tests on LVIS and GLAS, *Remote Sensing*, 6, 6566–6586, doi:10.3390/Rs6076566, 2014.

Pereira, H. M., Ferrier, S., Walters, M., Geller, G. N., Jongman, R. H. G., Scholes, R. J., Bruford, M. W., Brummitt, N., Butchart, S. H. M., Cardoso, A. C., Coops, N. C., Dulloo, E., Faith, D. P., Freyhof, J., Gregory, R. D., Heip, C., Hoft, R., Hurtt, G., Jetz, W., Karp, D. S., McGeoch, M. A., Obura, D., Onoda, Y., Pettorelli, N., Reyers, B., Sayre, R., Scharlemann, J. P. W., Stuart, S. N., Turak, E., Walpole, M., and Wegmann, M.: Essential biodiversity variables, *Science*, 339, 277–278, doi:10.1126/science.1229931, 2013.

15 Pfeifer, M., Gonsamo, A., Disney, M., Pellikka, P., and Marchant, R.: Leaf area index for biomes of the eastern arc mountains: Landsat and spot observations along precipitation and altitude gradients, *Remote Sens. Environ.*, 118, 103–115, doi:10.1016/j.rse.2011.11.009, 2012.

Pfeifer, M., Lefebvre, V., Gonsamo, A., Pellikka, P., Marchant, R., Denu, D., and Platts, P.: Validating and linking the gimms leaf area index (LAI3g) with environmental controls in tropical africa, *Remote Sensing*, 6, 1973–1990, 2014.

20 PRISM: 30 yr normal precipitation: annual, in: 1981–2010, August 2013 ed., PRISM Climate Group, Oregon State University, Corvallis, OR, 2013.

Randerson, J. T., Hoffman, F. M., Thornton, P. E., Mahowald, N. M., Lindsay, K., Lee, Y. H., Nevison, C. D., Doney, S. C., Bonan, G., Stockli, R., Covey, C., Running, S. W., and Fung, I. Y.:

## Characterizing Leaf Area Index (LAI) and Vertical Foliage Profile (VFP)

H. Tang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Systematic assessment of terrestrial biogeochemistry in coupled climate-carbon models, *Glob. Change Biol.*, 15, 2462–2484, doi:10.1111/j.1365-2486.2009.01912.x, 2009.

Reuter, H. I., Nelson, A., and Jarvis, A.: An evaluation of void-filling interpolation methods for srtm data, *Int. J. Geogr. Inf. Sci.*, 21, 983–1008, doi:10.1080/13658810601169899, 2007.

5 Saatchi, S. S., Harris, N. L., Brown, S., Lefsky, M., Mitchard, E. T. A., Salas, W., Zutta, B. R., Buermann, W., Lewis, S. L., Hagen, S., Petrova, S., White, L., Silman, M., and Morel, A.: Benchmark map of forest carbon stocks in tropical regions across three continents, *P. Natl. Acad. Sci. USA*, 108, 9899–9904, doi:10.1073/pnas.1019576108, 2011.

10 Sauer, J. R., Hines, J. E., Fallon, J., Pardieck, K., Ziolkowski Jr., D., and Link, W.: The North American Breeding Bird Survey, Results and Analysis 1966–2007, version 5, Laurel, MD, USGS Patuxent Wildlife Research Center, 2008.

15 Schimel, D. S., House, J. I., Hibbard, K. A., Bousquet, P., Ciais, P., Peylin, P., Braswell, B. H., Apps, M. J., Baker, D., Bondeau, A., Canadell, J., Churkina, G., Cramer, W., Denning, A. S., Field, C. B., Friedlingstein, P., Goodale, C., Heimann, M., Houghton, R. A., Melillo, J. M., Moore, B., Murdiyarso, D., Noble, I., Pacala, S. W., Prentice, I. C., Raupach, M. R., Rayner, P. J., Scholes, R. J., Steffen, W. L., and Wirth, C.: Recent patterns and mechanisms of carbon exchange by terrestrial ecosystems, *Nature*, 414, 169–172, doi:10.1038/35102500, 2001.

20 Schut, A. G. T., Wardell-Johnson, G. W., Yates, C. J., Keppel, G., Baran, I., Franklin, S. E., Hopper, S. D., Van Niel, K. P., Mucina, L., and Byrne, M.: Rapid characterisation of vegetation structure to predict refugia and climate change impacts across a global biodiversity hotspot, *Plos One*, 9, e82778, doi:10.1371/journal.pone.0082778, 2014.

25 Shabanov, N. V., Huang, D., Yang, W. Z., Tan, B., Knyazikhin, Y., Myneni, R. B., Ahl, D. E., Gower, S. T., Huete, A. R., Aragao, L. E. O. C., and Shimabukuro, Y. E.: Analysis and optimization of the modis leaf area index algorithm retrievals over broadleaf forests, *IEEE T. Geosci. Remote*, 43, 1855–1865, doi:10.1109/Tgrs.2005.852477, 2005.

Simard, M., Pinto, N., Fisher, J. B., and Baccini, A.: Mapping forest canopy height globally with spaceborne lidar, *J. Geophys. Res.-Biogeo.*, 116, G04021, doi:10.1029/2011jg001708, 2011.

30 Spracklen, D. V., Arnold, S. R., and Taylor, C. M.: Observations of increased tropical rainfall preceded by air passage over forests, *Nature*, 489, 282–U127, doi:10.1038/Nature11390, 2012.

## Characterizing Leaf Area Index (LAI) and Vertical Foliage Profile (VFP)

H. Tang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Sprintsin, M., Chen, J. M., Desai, A., and Gough, C. M.: Evaluation of leaf-to-canopy upscaling methodologies against carbon flux data in north america, *J. Geophys. Res.-Biogeo.*, 117, G01023, doi:10.1029/2010jg001407, 2012.

Stahl, G., Holm, S., Gregoire, T. G., Gobakken, T., Naesset, E., and Nelson, R.: Model-based inference for biomass estimation in a lidar sample survey in hedmark county, norway, *Can. J. Forest Res.*, 41, 96–107, doi:10.1139/X10-161, 2011.

Strahler, A. H., Jupp, D. L. B., Woodcock, C. E., Schaaf, C. B., Yao, T., Zhao, F., Yang, X., Lovell, J., Culvenor, D., Newnham, G., Ni-Miester, W., and Boykin-Morris, W.: Retrieval of forest structural parameters using a ground-based lidar instrument (echidna<sup>®</sup>), *Can. J. Remote Sens.*, 34, S426–S440, doi:10.5589/m08-046, 2008.

Swatantran, A., Dubayah, R., Goetz, S., Hofton, M., Betts, M. G., Sun, M., Simard, M., and Holmes, R.: Mapping migratory bird prevalence using remote sensing data fusion, *Plos One*, 7, e28922, doi:10.1371/journal.pone.0028922, 2012.

Tang, H., Dubayah, R., Swatantran, A., Hofton, M., Sheldon, S., Clark, D. B., and Blair, B.: Retrieval of vertical lai profiles over tropical rain forests using waveform lidar at la selva, costa rica, *Remote Sens. Environ.*, 124, 242–250, doi:10.1016/j.rse.2012.05.005, 2012.

Tang, H., Brolly, M., Zhao, F., Strahler, A. H., Schaaf, C. L., Ganguly, S., Zhang, G., and Dubayah, R.: Deriving and validating leaf area index (LAI) at multiple spatial scales through lidar remote sensing: a case study in Sierra National Forest, CA, *Remote Sens. Environ.*, 143, 131–141, 2014a.

Tang, H., Dubayah, R., Brolly, M., Ganguly, S., and Zhang, G.: Large-scale retrieval of leaf area index and vertical foliage profile from the spaceborne waveform lidar (glas/icesat), *Remote Sens. Environ.*, 154, 8–18, doi:10.1016/j.rse.2014.08.007, 2014b.

Turner, D. P., Ritts, W. D., Cohen, W. B., Gower, S. T., Running, S. W., Zhao, M. S., Costa, M. H., Kirschbaum, A. A., Ham, J. M., Saleska, S. R., and Ahl, D. E.: Evaluation of modis npp and gpp products across multiple biomes, *Remote Sens. Environ.*, 102, 282–292, doi:10.1016/j.rse.2006.02.017, 2006.

Turner, W., Spector, S., Gardiner, N., Fladeland, M., Sterling, E., and Steininger, M.: Remote sensing for biodiversity science and conservation, *Trends Ecol. Evol.*, 18, 306–314, doi:10.1016/S0169-5347(03)00070-3, 2003.

Valladares, F. and Niinemets, U.: Shade tolerance, a key plant feature of complex nature and consequences, *Annu. Rev. Ecol. Evol. S.*, 39, 237–257, doi:10.1146/annurev.ecolsys.39.110707.173506, 2008.

## Characterizing Leaf Area Index (LAI) and Vertical Foliage Profile (VFP)

H. Tang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Whitehurst, A. S., Swatantran, A., Blair, J. B., Hofton, M. A., and Dubayah, R.: Characterization of canopy layering in forested ecosystems using full waveform lidar, *Remote Sensing*, 5, 2014–2036, doi:10.3390/Rs5042014, 2013.

5 Yang, W. Z., Tan, B., Huang, D., Rautiainen, M., Shabanov, N. V., Wang, Y., Privette, J. L., Huemmrich, K. F., Fensholt, R., Sandholt, I., Weiss, M., Ahl, D. E., Gower, S. T., Nemani, R. R., Knyazikhin, Y., and Myneni, R. B.: Modis leaf area index products: from validation to algorithm improvement, *IEEE T. Geosci. Remote*, 44, 1885–1898, doi:10.1109/Tgrs.2006.871215, 2006.

10 Zhao, F., Yang, X. Y., Strahler, A. H., Schaaf, C. L., Yao, T., Wang, Z. S., Roman, M. O., Woodcock, C. E., Ni-Meister, W., Jupp, D. L. B., Lovell, J. L., Culvenor, D. S., Newnham, G. J., Tang, H., and Dubayah, R. O.: A comparison of foliage profiles in the sierra national forest obtained with a full-waveform under-canopy evi lidar system with the foliage profiles obtained with an airborne full-waveform lvis lidar system, *Remote Sens. Environ.*, 136, 330–341, doi:10.1016/j.rse.2013.05.020, 2013.

15 Zhao, M. S., Heinsch, F. A., Nemani, R. R., and Running, S. W.: Improvements of the modis terrestrial gross and net primary production global data set, *Remote Sens. Environ.*, 95, 164–176, doi:10.1016/j.rse.2004.12.011, 2005.

## Characterizing Leaf Area Index (LAI) and Vertical Foliage Profile (VFP)

H. Tang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



**Table 1.** Ecoregions with highest total LAI values (unit:  $\text{m}^2 \text{m}^{-2}$ ).

Ecoregions	Total LAI Mean( $\pm$ SD)	LAI 0–10 m Mean( $\pm$ SD)	LAI 10–20 m Mean( $\pm$ SD)	LAI > 20 m Mean( $\pm$ SD)
Northern California coastal forests	5.24 $\pm$ 2.11	2.06 $\pm$ 1.32	1.67 $\pm$ 1.09	1.08 $\pm$ 1.15
Central Pacific coastal forests	5.00 $\pm$ 2.14	1.52 $\pm$ 1.61	1.10 $\pm$ 1.16	0.84 $\pm$ 1.25
British Columbia mainland coastal forests	4.74 $\pm$ 2.26	1.48 $\pm$ 1.31	1.23 $\pm$ 1.08	1.13 $\pm$ 1.13
Central and Southern Cascades forests	4.31 $\pm$ 2.34	1.06 $\pm$ 1.35	0.79 $\pm$ 1.02	0.64 $\pm$ 1.07
Klamath–Siskiyou forests	4.31 $\pm$ 2.31	1.26 $\pm$ 1.30	0.99 $\pm$ 1.07	0.73 $\pm$ 0.99
Willamette Valley forests	3.99 $\pm$ 2.24	0.73 $\pm$ 1.09	0.60 $\pm$ 0.89	0.75 $\pm$ 1.31
Appalachian–Blue Ridge forests	3.95 $\pm$ 2.03	1.04 $\pm$ 1.27	0.82 $\pm$ 0.99	0.47 $\pm$ 0.82
Puget lowland forests	3.91 $\pm$ 2.25	0.98 $\pm$ 1.39	0.71 $\pm$ 1.08	0.40 $\pm$ 0.81
Appalachian mixed mesophytic forests	3.86 $\pm$ 2.04	1.06 $\pm$ 1.29	0.77 $\pm$ 0.93	0.48 $\pm$ 0.83
North Central Rockies forests	3.67 $\pm$ 2.27	1.61 $\pm$ 1.55	0.84 $\pm$ 0.89	0.47 $\pm$ 0.72



**Figure 1.** A comparison between two lidar derived Leaf Area Index (LAI) datasets at different sites across the US ( $N = 318$ ), produced from the Laser Vegetation Imaging Sensor (LVIS) and the Geoscience Laser Altimeter System (GLAS) respectively. Each point represents a comparison at GLAS footprint while different colors and shapes indicate different sites (AR: White River National Wildlife Refuge in Arkansas; CA: Sierra National Forest in California; MD: Baltimore/Washington corridor in Maryland; ME: Maine forests to the north of Orono, Maine). The comparison produces  $r^2$  of 0.60, bias of  $-0.23$ , and RMSE of 0.83. Dashed line is the 1 : 1 line.

**Characterizing Leaf Area Index (LAI) and Vertical Foliage Profile (VFP)**

H. Tang et al.

[Title Page](#)

[Abstract](#) | [Introduction](#)

[Conclusions](#) | [References](#)

[Tables](#) | [Figures](#)

[◀](#) | [▶](#)

[◀](#) | [▶](#)


[Back](#) | [Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)





**Figure 2.** A comparison of Vertical Foliage Profile (VFP) density derived from LVIS and GLAS over different sites in the US (same sites as Fig. 1 but with  $N = 1272$ ). Each VFP point represents an integrated value of foliage density at 5 m height interval.

## BGD

12, 13675–13710, 2015

### Characterizing Leaf Area Index (LAI) and Vertical Foliage Profile (VFP)

H. Tang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## Characterizing Leaf Area Index (LAI) and Vertical Foliage Profile (VFP)

H. Tang et al.



**Figure 3.** Comparison between LVIS and GLAS VFP density integrated at every 5 m height interval (from ground to canopy top).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






**BGD**

12, 13675–13710, 2015

**Characterizing Leaf Area Index (LAI) and Vertical Foliage Profile (VFP)**

H. Tang et al.


**Figure 5.** GLAS LAI distributions by ecoregion.[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## BGD

12, 13675–13710, 2015

## Characterizing Leaf Area Index (LAI) and Vertical Foliage Profile (VFP)

H. Tang et al.




**Figure 6.** LAI strata distributions by WWF ecoregions. Despite similar total LAI values, the southeastern forests show different LAI values at stratified height intervals.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


## Characterizing Leaf Area Index (LAI) and Vertical Foliage Profile (VFP)

H. Tang et al.



**Figure 7.** Distribution of total GLAS LAI across different land cover types. The width of the boxes is proportional to the number of observations for each type ( $N$  = Evergreen Needleleaf: 45 207, Evergreen Broadleaf: 438, Deciduous Needleleaf: 123, Deciduous Broadleaf: 48 283, Mixed Forest: 62 053, Closed Shrubland: 4087, Open Shrubland: 7364, Woody Savanna: 43 536, Savanna: 3051). Notches show the approximate 95 % confidence interval of the median.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Characterizing Leaf Area Index (LAI) and Vertical Foliage Profile (VFP)

H. Tang et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶


[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)


[Interactive Discussion](#)



**Figure 8.** Averaged GLAS VFP for different land cover types across US: non-forest vegetation types (left) and forest types (right). Mean values are central lines within the color-filled 95 % CI envelope.

## Characterizing Leaf Area Index (LAI) and Vertical Foliage Profile (VFP)

H. Tang et al.



**Figure 9.** Distribution of GLAS LAI (left axis) and Forest Ratio – GLAS shots over forest divided by total shot numbers – (right axis). Overall, there is a decreasing trend of LAI values as elevation increases, but deviations occur from this trend that are associated with elevational variation in Forest Ratio.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## Characterizing Leaf Area Index (LAI) and Vertical Foliage Profile (VFP)

H. Tang et al.



**Figure 10.** Distribution of GLAS LAI as a function of precipitation. A linear regression analysis of LAI values averaged by precipitation groups shows an increasing trend up to areas of about 2400 mm (blue line). Beyond this value the rate of change slows considerably (magenta line) but the trend is only weakly significant ( $P = 0.09$ ).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

