

1 **Microbial communities and their predicted metabolic**  
2 **characteristics in deep fracture groundwaters of the crystalline**  
3 **bedrock at Olkiluoto, Finland**

4

5 **M. Bomberg<sup>1</sup>, T. Lamminmäki<sup>2</sup> and M. Itävaara<sup>1</sup>**

6 [1] VTT Technical Research Centre of Finland, P.O. Box 1000, FIN-02044 VTT, Finland

7 [2] Posiva Oy, Olkiluoto, 27160 Eurajoki, Finland

8 Correspondence to: M. Bomberg (malin.bomberg@vtt.fi)

9

10 **Abstract**

11 The microbial diversity in oligotrophic isolated crystalline Fennoscandian Shield bedrock  
12 fracture groundwaters is high but the core community has not been identified. Here we  
13 characterized the bacterial and archaeal communities in 12 water conductive fractures situated  
14 at depths between 296 m and 798 m by high throughput amplicon sequencing using the  
15 Illumina HiSeq platform. Between  $1.7 \times 10^4$  –  $1.2 \times 10^6$  bacterial or archaeal sequence reads  
16 per sample was obtained. These sequences revealed that up to 95% and 99% of the bacterial  
17 and archaeal sequences obtained from the 12 samples, respectively, belonged to only a few  
18 common species, i.e. the core microbiome. However, the remaining rare microbiome  
19 contained over 3 and 6 fold more bacterial and archaeal taxa. The metabolic properties of the  
20 microbial communities were predicted using PICRUSt. The approximate estimation showed  
21 that the metabolic pathways included commonly fermentation, fatty acid oxidation,  
22 glycolysis/gluconeogenesis, oxidative phosphorylation and methanogenesis/anaerobic  
23 methane oxidation, but carbon fixation through the Calvin cycle, reductive TCA cycle and the  
24 Wood-Ljungdahl pathway was also predicted. The rare microbiome is an unlimited source of  
25 genomic functionality in all ecosystems. It may consist of remnants of microbial communities  
26 prevailing in earlier environmental conditions, but could also be induced again if changes in  
27 their living conditions occur.

## 1 **1 Introduction**

2 Identifying and understanding the core microbiome of any given environments is of crucial  
3 importance for predicting and assessing environmental change both locally and globally  
4 (Shade and Handelsman, 2012). In a previous study (Bomberg et al., 2015) we showed by 454  
5 amplicon sequencing that the active microbial communities in Olkiluoto deep subsurface  
6 were strictly stratified according to aquifer water type. Nevertheless, more rigorous  
7 sequencing efforts and more samplings have shown that an archaeal core community  
8 consisting of the DeepSea Hydrothermal Vent Euryarchaeotal Group 6 (DHVEG-6), ANME-  
9 2D and Terrestrial Miscellaneous Group (TMEG) archaea may exist in the anaerobic deep  
10 groundwater of Olkiluoto (Miettinen et al., 2015). The bacterial core groups in Olkiluoto deep  
11 groundwater include at least members of the Pseudomonadaceae, Comamonadaceae and  
12 Sphingomonadaceae (Bomberg et al., 2014; 2015; Miettinen et al., 2015). The relative  
13 abundance of these main groups varies at different depths from close to the detection limit to  
14 over 90% of the bacterial or archaeal community (Bomberg et al., 2015; Miettinen et al.,  
15 2015). However, both the archaeal and the bacterial communities contain a wide variety of  
16 less abundant groups, which are distributed unevenly in the different water conductive  
17 fractures.

18 The rare biosphere is a concept describing the hidden biodiversity of an environment (Sogin  
19 et al., 2006). The rare biosphere consists of microbial groups that are ubiquitously distributed  
20 in nature but often present at low relative abundance and may thus stay below the limit of  
21 detection. Due to modern high throughput sequencing techniques, however, the hidden  
22 diversity of rare microbiota has been revealed. These microorganisms are the basis for  
23 unlimited microbial functions in the environment and upon environmental change specific  
24 groups can readily activate and become abundant. Access to otherwise inaccessible nutrients  
25 activate specific subpopulations in the bacterial communities within hours of exposure (Rajala  
26 et al., 2015) and enrich distinct microbial taxa at the expense of the original microbial  
27 community in the groundwater (Kutvonen, 2015). Mixing of different groundwater layers due  
28 to e.g. breakage of aquifer boundaries and new connection of separated aquifers may cause  
29 the microbial community to change and activate otherwise dormant processes. This has  
30 previously been shown by Pedersen et al. (2013), who indicated increased sulphate reduction  
31 activity when sulphate-rich and methane-rich groundwater mixed. The stability of deep  
32 subsurface microbial communities in isolated deep subsurface groundwater fractures are

1 assumed to be stable. However, there are indications that they may change over the span of  
2 several years as slow flow along fractures is possible (Miettinen et al., 2015; Sohlberg et al.,  
3 2015).

4 The microbial taxa present in an environment interact with both biotic and abiotic factors. In  
5 deep subsurface groundwater the biomass concentration is often low and the sampling efforts  
6 may not yield enough biomass for extensive metagenomic analysis of the microbial  
7 communities. Tools for predicting metabolic pathways may help to establish a consensus of  
8 the microbial metabolic characteristics present in an environment and the possible interactions  
9 of the microbial communities with the abiotic environment. Tools, such as PICRUSt (Langille  
10 *et al.*, 2013), allow us to estimate microbial metabolic functions based on NGS microbiome  
11 data. For example, Tsitko et al. (2014) showed that oxidative phosphorylation was the most  
12 important energy producing metabolic pathway throughout the 7 m depth profile of an  
13 *Acidobacteria*-dominated nutrient poor boreal bog. Cleary et al. (2015) showed that tropical  
14 mussel-associated bacterial communities could be important sources of bioactive compounds  
15 for biotechnology. This approach is nevertheless hampered by the fact that only little is so far  
16 known about uncultured environmental microorganisms and their functions and the PICRUSt  
17 approach is best applied for human microbiome for which it was initially developed (Langille  
18 *et al.*, 2013). However, metagenomic estimations may give important indications of novel  
19 metabolic possibilities even in environmental microbiome studies.

20 Using extensive high throughput amplicon sequencing in this study we aimed to identify the  
21 core microbiome in the deep crystalline bedrock fractures of Olkiluoto Island and also to  
22 study the rare microbiome. In addition, we aimed to estimate the prevailing metabolic  
23 activities that may occur in the deep crystalline bedrock environment of Olkiloto, Finland.

## 24 **2 Materials and methods**

### 25 **2.1 Background**

26 The Olkiluoto site has previously been extensively described (Posiva, 2013) and is only  
27 briefly described here. The Island of Olkiluoto situating on the west coast of Finland has  
28 approximately 60 drillholes drilled for research and monitoring purposes. Studies on the  
29 chemistry and microbiology of the groundwater have been on-going since the 1980s. The  
30 groundwater is stratified with a salinity gradient extending from fresh to brackish water to a  
31 depth of 30 m and the highest salinity concentration of 125 g L<sup>-1</sup> total dissolved solids (TDS)

1 at 1000 m depth (Posiva, 2013). The most abundant salinity causing cations are  $\text{Na}^{2+}$  and  $\text{Ca}^{2+}$   
2 and anions  $\text{Cl}^-$ . Between 100 and 300 m depths, the groundwater originates from ancient (pre-  
3 Baltic) seawater and has high concentrations of  $\text{SO}_4^{2-}$ . Below 300 m the concentration of  
4 methane in the groundwater increases and  $\text{SO}_4^{2-}$  is almost absent. A sulphate-methane  
5 transition zone (SMTZ), where sulphate-rich fluid replaces methane-rich fluid, is located at  
6 250 – 350 m depth. Temperature rises linearly with depth, from ca. 5 – 6 °C at 50 m to ca. 20  
7 °C at 1,000 m depth (Ahokas et al., 2008). The pH of the groundwater is slightly alkaline  
8 throughout the depth profile. Multiple drillholes intersect several groundwater-filled bedrock  
9 fractures, including larger hydrogeological zones such as HZ20 or HZ21 (Table 1). The  
10 bedrock of Olkiluoto consists mainly of micagneiss and pegmatitic granite type rocks (Kärki  
11 & Paulamäki, 2006). The *in situ* temperature at 300 m depth in the Olkiluoto bedrock is stable  
12 at approximately 10°C and increases linearly to approximately 16°C at 800 m depth (Sedighi  
13 et al., 2013).

14 This study focused on 12 groundwater samples from water conductive fractures situated  
15 between 296 m and 798 m below sea level bsl and originating from 11 different drillholes in  
16 Olkiluoto (Figure 1). The samples represented brackish sulphate waters and saline waters (as  
17 classified in Posiva, 2013). The samples were collected between December 2009 and January  
18 2013 (Table 1). The physicochemical parameters of the groundwater samples have been  
19 reported by reported by Miettinen et al. (2015), but have for clarity been collected here (Table  
20 1).

## 21 **2.2 Sample collection**

22 The collection of samples occurred between December 2009 and January 2013 (Table 1) as  
23 described previously (Bomberg et al., 2015; Miettinen et al., 2015; Sohlberg et al., 2015). The  
24 samples were obtained from 11 different permanently packered or open drillholes equipped  
25 with removable inflatable packers. The position and direction of the drillholes are indicated in  
26 Figure 1. Shortly, in order to obtain indigenous fracture fluids, the packer-isolated fracture  
27 zones were purged by removing stagnant drillhole water by pumping for a minimum of four  
28 weeks before the sample water was collected. The water samples were collected directly from  
29 the drillhole into an anaerobic glove box (MBRAUN, Germany) via a sterile, gas-tight poly  
30 acetate tube (8 mm outer diameter). Microbial biomass DNA extraction was concentrated  
31 from 1000 mL samples by filtration on cellulose acetate filters (0.2 µm pore size, Corning) by  
32 vacuum suction inside the glove box. The filters were immediately extracted from the

1 filtration funnels and frozen on dry ice in sterile 50 ml cone tubes (Corning). The frozen  
2 samples were transported on dry ice to the laboratory where they were stored at -80°C until  
3 use.

### 4 **2.3 Nucleic acid isolation**

5 Community DNA was isolated directly from the frozen cellulose-acetate filters with the  
6 PowerSoil DNA extraction kit (MoBio Laboratories, Inc., Solana Beach, CA), as previously  
7 described (Bomberg et al., 2015). Negative DNA isolation controls were included in the  
8 isolation protocol. The DNA concentration of each sample was determined using the  
9 NanoDrop 1000 spectrophotometer.

### 10 **2.4 Estimation of microbial community size**

11 The size of the microbial community was determined by epifluorescence microscopy of 4',6  
12 diamidino-2-phenylindole dihydrochloride (DAPI) (Sigma, MO, USA) stained cells as  
13 described in Purkamo et al. (2013). The size of the bacterial population was determined by  
14 16S rRNA gene targeted quantitative PCR (qPCR) as described by Tsitko et al. (2014) using  
15 universal bacterial 16S rRNA gene-targeting primers fD1 (Weisburg et al., 1991) and P2  
16 (Muyzer et al., 1993), which specifically target the V1- V3 region of the bacterial 16S rDNA  
17 gene. The size of the archaeal population in the groundwater was determined by using primers  
18 ARC344f (Bano *et al.*, 2004) and Ar744r (reverse compliment from Barns *et al.*, 1994)  
19 flanking the V4-V6 region of the archaeal 16S rRNA gene.

20 The qPCR reactions were performed in 10µL reaction volumes using the KAPA 2 × Syrb®  
21 FAST qPCR-kit on a LightCycler480 qPCR machine (Roche Applied Science, Germany) on  
22 white 96-well plates (Roche Applied Science, Germany) sealed with transparent adhesive  
23 seals (4titude, UK). Each reaction contained 2.5 µM of relevant forward and reverse primer  
24 and 1 µL DNA extract. Each reaction was run in triplicate and no-template control reactions  
25 were used to determine background fluorescence in the reactions.

26 The qPCR conditions consisted of an initial denaturation at 95 °C for 10 minutes followed by  
27 45 amplification cycles of 15 seconds at 95 °C, 30 seconds at 55 °C and 30 seconds at 72 °C  
28 with a quantification measurement at the end of each elongation. A final extension step of  
29 three minutes at 72 °C was performed prior to a melting curve analysis. This consisted of a  
30 denaturation step for 10 seconds at 95 °C followed by an annealing step at 65 °C for one

1 minute prior to a gradual temperature rise to 95 °C at a rate of 0.11 °C s<sup>-1</sup> during which the  
2 fluorescence was continuously measured. The number of bacterial 16S rRNA genes was  
3 determined by comparing the amplification result (Cp) to that of a ten-fold dilution series  
4 (10<sup>1</sup>-10<sup>7</sup> copies μL<sup>-1</sup>) of *Escherichia coli* (ATCC 31608) 16S rRNA genes in plasmid for  
5 bacteria and a dilution series of genomic DNA of *Halobacterium salinarum* (DSM 3754) for  
6 archaea. The lowest detectable standard concentration for the qPCRs was 10<sup>2</sup> gene  
7 copies/reaction. Inhibition of the qPCR by template tested by adding 2.17×10<sup>4</sup> plasmid copies  
8 containing fragment of the morphine-specific Fab gene from *Mus musculus* gene to reactions  
9 containing template DNA as described in Nyysönen *et al.* (2012). Inhibition of the qPCR  
10 assay by the template DNA was found to be low. The average Crossing point (Cp) value for  
11 the standard sample (2.17×10<sup>4</sup> copies) was 28.7 (± 0.4 sd), while for the DNA samples Cp  
12 was 28.65 - 28.91 (± 0.03-0.28 sd). Nucleic acid extraction and reagent controls were run in  
13 all qPCRs in parallel with the samples. Amplification in these controls was never higher than  
14 the background obtained from the no template controls.

## 15 **2.5 Amplicon library preparation**

16 This study is part of the Census of Deep Life initiative, which strives to obtain a census of the  
17 microbial diversity in deep subsurface environment by collecting samples around the world  
18 and sequencing the 16S rRNA gene pools of both archaea and bacteria. The extracted DNA  
19 samples were sent to the Marine Biological Laboratory in Woods Hole, MA, USA, for  
20 preparation for HiSeq sequencing using the Illumina technology. The protocol for amplicon  
21 library preparation for both archaeal and bacterial 16S amplicon libraries can be found at  
22 <http://vamms.mbl.edu/resources/faq.php>. Shortly, amplicon libraries for completely  
23 overlapping paired-end sequencing of the V6 region of both the archaeal and bacterial 16S  
24 rRNA genes were produced as previously described (Eren *et al.*, 2013). For the archaea,  
25 primers A958F and A1048R containing Truseq adapter sequences at their 5' end were used,  
26 and for the bacteria primers B967F and B1064R for obtaining 100 nt long paired end reads  
27 (<https://vamms.mbl.edu/resources/primers.php>). The sequencing was performed using a HiSeq  
28 1000 system (Illumina).

## 29 **2.6 Sequence processing and analysis**

30 Contigs of the paired end fastq files were first assembled with mothur v 1.32.1 (Schloss *et al.*,  
31 2009). Analyzes were subsequently continued using QIIME v. 1.8. (Caporaso *et al.*, 2010).

1 Only sequences with a minimum length of 50 bp were included in the analyses. The bacterial  
2 and archaeal 16S rRNA sequences were grouped into OTUs (97% sequence similarity) using  
3 both the open reference and closed reference OTU picking strategy and classified using the  
4 GreenGenes 13\_8 16S reference database (DeSantis et al., 2006). The core archaea and  
5 bacteria communities were identified from the OTU tables with the  
6 `compute_core_microbiome.py` function in QIIME using default values, with exception of the  
7 minimum number of samples where an OTU must be detected, which was set to 80%. The  
8 sequencing coverage was evaluated by rarefaction analysis and the estimated species richness  
9 and diversity indices were calculated. For comparable  $\alpha$ - and  $\beta$ -diversity analyses the data sets  
10 were normalized by random subsampling of 17,000 sequences/sample for archaea and  
11 140,000 sequences/sample for bacteria. Microbial metabolic pathways were estimated based  
12 on the 16S rRNA gene data from the closed OTU picking method using the PICRUSt  
13 software (Langille *et al.*, 2013) on the web based Galaxy application (Goecks et al., 2010;  
14 Blankenberg et al., 2010; Giardine et al., 2005). The predicted KO numbers were plotted on  
15 KEGG pathway maps (<http://www.genome.jp/kegg/>) separately for the bacterial and archaeal  
16 predicted metagenomes, with a threshold of a minimum of 100 genes in total estimated from  
17 all samples. The sequence data has been submitted to the Sequence Read Archive (SRA,  
18 <http://www.ncbi.nlm.nih.gov/sra>) under study SRP053854, Bioproject PRJNA275225.

## 19 **2.7 Statistical analyses and data visualization**

20 The similarity of the archaeal and bacterial communities between the different samples was  
21 tested by principal coordinates analysis (PCoA) using the Phyloseq package in R (McMurdie  
22 and Holmes, 2014; R Core Team, 2013). The analysis was performed using the raw OTU  
23 tables outputted by QIIME. In addition, a PCoA analysis showing the effect of library size on  
24 the ordination of the samples was calculated using `vegan` (Oksanen et al., 2016). The Bray-  
25 Curtis distance model was used for both analyses. The samples were hierarchically clustered  
26 in a UPGMA tree based on the raw OTU counts using the `heatmap` function of `phyloseq` in R.

27

## 1 **3 Results**

### 2 **3.1 Microbial community size**

3 The total number of microbial cells detected by epifluorescence microscopy of DAPI stained  
4 cells was between  $2.3 \times 10^4$  and  $4.2 \times 10^5$  cells mL<sup>-1</sup> groundwater (Figure 2, Table 1). The  
5 concentration of bacterial 16S rRNA gene copies mL<sup>-1</sup> varied between  $9.5 \times 10^3$  and  $7.0 \times 10^5$   
6 and that of the archaea  $2.6 \times 10^1$  and  $6.3 \times 10^4$  (Figure 2, Table 1).

### 7 **3.2 Sequence statistics, diversity estimates and sequencing coverage**

8 The number of bacterial v6 sequence reads from the 12 samples varied between  $1.4 - 7.8 \times$   
9  $10^5$  reads, with a mean sequencing depth of  $2.9 \times 10^5$  ( $\pm 1.8 \times 10^5$  standard deviation)  
10 reads/sample (Table 2). The archaeal v6 sequence reads ranged from  $0.17 - 12.1 \times 10^5$  reads  
11 with a mean sequencing depth of  $4.1 \times 10^5$  ( $\pm 3.5 \times 10^5$  standard deviation) reads/sample. The  
12 numbers of observed operational taxonomic units (OTUs) represented on average 82.6% ( $\pm$   
13 12.5%) of the Chao1- and 78.1 % ( $\pm 13.4\%$ ) of the ACE-estimated numbers of bacterial  
14 OTUs (Table 2ab). The archaeal communities were slightly better covered, with on average  
15 88.5% ( $\pm 11.5\%$ ) of the Chao1 and 84.8% ( $\pm 12.6\%$ ) of the ACE estimated number of OTUs  
16 detected. Shannon diversity index H', calculated from 140,000 and 17,000 random sequence  
17 reads per sample for the bacteria and archaea, respectively, was high for both bacterial and  
18 archaeal communities. High H' values and climbing rarefaction curves (Figure S1) indicated  
19 high diversity in the microbial communities in the different deep groundwater fracture zones  
20 of Olkiluoto. The bacterial H' was on average 13 ( $\pm 0.74$ ), ranging from 11 to 14 between the  
21 different samples. The archaeal H' was on average 11 ( $\pm 1.2$ ) ranging from 9 to 12 between  
22 the samples. A total of 468,684 archaeal and 301,458 bacterial OTUs were obtained in this  
23 study.

### 24 **3.3 Microbial communities**

25 From the bacterial v6 sequences 49 different bacterial Phyla were detected (Appendix 1).  
26 These phyla included 165 bacterial classes, 230 orders, 391 families and 651 genera. The  
27 greatest number of sequences, between 21.83% and 47.94% per sample, clustered into an  
28 undetermined bacterial group (Bacteria, Other). This may be due the fact that sequences of  
29 poorer quality may be difficult to classify, especially as the sequences are short, or these

1 sequences belong to thus far uncultured and unknown bacterial species, the so called  
2 microbial dark matter (Solden et al., 2016).

3 The archaea were represented by two identified phyla, the Euryarchaeota and the  
4 Crenarchaeota (Appendix 2). These included 21 classes, 38 orders, 61 families and 81 genera.  
5 Between 4.7% and 35.0% of the archaeal sequences of each sample were classified to  
6 unassigned Archaea, with a general increase in unassigned archaeal sequences with increasing  
7 depth.

8 The archaeal and bacterial core communities were determined as OTUs present in at least  
9 80% of the samples. Of the more than  $4.6 \times 10^5$  archaeal OTUs the core community consisted  
10 of 82 OTUs belonging to three archaeal orders, the E2 of the Thermoplasmatales, the  
11 Methanobacteriales and the Methanosarcinales (Figure 3). Additionally, a great proportion of  
12 the OTUs of the core community did not receive any other taxonomic identity than Archaea.  
13 The most common archaeal family of the core community was the ANME-2D belonging to  
14 the Methanosarcinales. The bacterial core community consisted of only 26 OTUs, compared  
15 to more than  $3.0 \times 10^5$  bacterial OTUs in total (Figure 3). These OTUs belonged to six  
16 different families, the Alteromonadales, Burkholderiales, SB-45, Sphingomonadales,  
17 Syntrophobacterales and Thiobacterales. In addition, a great portion of the core community  
18 OTUs were classified only as unassigned bacteria. The most abundant of the bacterial core  
19 community OTUs belonged to Thiobacteriaceae and Comamonadaceae. In both the archaeal  
20 and bacterial sequence data a great proportion of the sequence reads were only identified as  
21 archaea or bacteria, without more detailed taxonomic assignments. The core OTUs were  
22 distributed with different abundance in the different samples (Figure 3). Most of the OTUs  
23 detected were present in less than 20% of the samples (Figure 4).

#### 24 **3.4 Environmental parameters and the microbial communities**

25 The microbial community profiles of the different samples were clustered in a UPGMA tree  
26 based on the OTU tables and the Bray-Curtis distance model (Figure 5). The archaeal and  
27 bacterial communities clustered according to the OTUs detected in the samples, but not  
28 clearly according to any physicochemical parameter. In the PCoA analysis, however, the  
29 archaeal communities in water containing total dissolved solids (TDS) between 10,670 mg  
30 and 18,580 mg L<sup>-1</sup> clustered together (Figures 6a, S2a), but no similar clustering could be  
31 observed in the bacterial communities (Figures 6b, S2b). Nevertheless, Principal coordinate 1

1 determined 14.9% and coordinate 2 12.6% of the variance in the archaeal communities and  
2 20% and 17.2%, respectively, of the variance in the bacterial communities (Figure 6ab, S2cd).

### 3 **3.5 Predicted metabolic functions of the deep subsurface microbial communities**

4 The putative metabolic functions of the microbial communities at different depth was  
5 predicted using the PICRUSt software, which compares the identified 16S rRNA gene  
6 sequences to those of known genome sequenced species thereby estimating the possible gene  
7 contents of the uncultured microbial communities. The analysis is only an approximation, but  
8 may give an idea of the possible metabolic activities in the deep biosphere. In order to  
9 evaluate the soundness of the analysis a nearest sequenced taxon index (NSTI) for each of the  
10 bacterial and archaeal communities was calculated by PICRUSt. An NSTI value of 0 indicates  
11 high similarity to the closest sequenced taxon while NSTI=1 indicates no similarity. The  
12 NSTI of the bacterial communities at different depths varied between 0.045 in sample OL-  
13 KR44 and 0.168 in sample OL-KR13 (Table 3). The NSTI for archaea were much higher  
14 ranging from 0.141 in sample OL-KR9 at depth of 432 m and 0.288 in OL-KR44. This  
15 indicates that the metagenomic estimates are only indicative. The estimated microbial  
16 metabolism did not differ noticeably between the different depths (Figure 7a and b). The most  
17 important predicted metabolic pathways included membrane transport in both bacterial and  
18 archaeal communities. The most common pathways predicted for carbohydrate metabolism  
19 were the butanoate, propionate, glycolysis/gluconeogenesis and pyruvate metabolism  
20 pathways for the bacteria and glycolysis/gluconeogenesis and pyruvate metabolism pathways  
21 for the archaea (Figure 8). Glucose is predicted to be converted into pyruvate and further to  
22 Acetyl-CoA by both bacteria and archaea. The bacterial community may produce and utilize  
23 acetate. Both the bacterial and archaeal communities are predicted to fix carbon via the  
24 Wood-Ljungdal (WL), reverse Citric acid cycle (rTCA) and Calvin pathways. Methane is  
25 most likely produced from methylamines, CO<sub>2</sub> and methanol by the methanogenic archaea. In  
26 addition to the strong evidence of methanogenesis in the archaeal community the reverse  
27 methanogenesis, i.e. anaerobic methane oxidation by the ANME-2D archaea is possible.  
28 Based on the predicted metagenomes the bacterial community is not able to oxidize methane  
29 or hydrolyze methanol, but the methylotrophs present may use formic acid and  
30 trimethylamines.

31 The most abundant predicted energy metabolic pathway in the bacterial communities was the  
32 oxidative phosphorylation (Figure S3) while for the archaea the methane metabolism was the

1 most important (Figure 7, 8). Our analyses predicted utilization of propanoate and butanoate  
2 (Figure 7) by the bacterial communities as well as completely covered fatty acid biosynthesis  
3 and degradation pathways, which indicate that the bacterial community may be capable of  
4 fermentation (Figure S4a and b). The bacterial community is predicted to reduce nitrate both  
5 through dissimilatory nitrate reduction to ammonia and through denitrification to nitrous  
6 oxide (Figure S5). In addition, the predicted metagenomes indicate that nitrogen is fixed to  
7 ammonia by both archaea and bacteria. The ammonia is furthermore predicted to be used as  
8 raw material for L-glutamate synthesis (Figure S5). Sulfur metabolism was not a major  
9 pathway in either the bacterial or the archaeal communities according to the predicted number  
10 of genes. However, assimilatory sulphate reduction was indicated in both the bacterial and  
11 archaeal communities, while dissimilatory sulphate reduction and sulphur oxidation was  
12 indicated only in the bacterial communities (Figure S6).

13 Several amino acid synthesis pathways were predicted (Figure 7), of which the most  
14 prominent were the alanine, aspartate and glutamate synthesis, arginine and proline synthesis,  
15 cysteine and methionine synthesis, glycine, serine and threonine synthesis, phenylalanine,  
16 tyrosine and tryptophan synthesis and the valine, leucine and isoleucine synthesis pathways.

17 Different types of membrane transport (ABC transporters) were predicted where sulphate and  
18 iron (III) would be taken up by the bacteria and tungstate, molybdate, proline, zinc, cobalt  
19 and nickel by both archaea and bacteria (Figure S7). The estimated number of genes for both  
20 the purine and pyrimidine metabolism was more than two times higher in the archaeal  
21 community than in the bacterial community (Figure 7a and b).

22

## 23 **4 Discussion**

24 The phenotypic characteristics of the Fennoscandian Shield deep subsurface microbial  
25 communities are still largely unknown although specific reactions to introduced  
26 environmental stimulants have been shown (e.g. Pedersen et al., 2013; 2014; Rajala et al.,  
27 2015; Kutvonen 2015). Nevertheless, the connection of these microbial responses to specific  
28 microbial groups is still only in an early phase. Metagenomic and gene specific analyses of  
29 deep subsurface microbial communities have revealed prominent metabolic potential of the  
30 microbial communities, which appear to be associated with the prevailing lithology and  
31 physicochemical parameters (Nyysönen et al., 2014; Purkamo et al., 2015). It has also been  
32 shown with fingerprinting methods with ever increasing efficiency that the bacterial and

1 archaeal communities are highly diverse in the saline anaerobic Fennoscandian deep fracture  
2 zone groundwater (Bomberg et al., 2014; 2015; Nyysönen et al., 2012; 2014; Pedersen et al.,  
3 2014; Miettinen et al., 2015; Sohlberg et al., 2015). Nevertheless, the concentration of  
4 microbial cells in the groundwater is quite low (Figure 2, Table 1). In accordance with other  
5 Fennoscandian deep subsurface environments (Purkamo et al., 2016), most of the microbial  
6 communities at different depth in Olkiluoto bedrock fractures consist of bacteria. Archaea  
7 have in general been shown to constitute at most approximately 1% of the Fennoscandian  
8 deep bedrock groundwater (Purkamo et al., 2016). However, at specific depths in Olkiluoto  
9 (328 m, 423 m) the archaea contributed with over 50% of the estimated 16S rRNA gene pool  
10 (Table 1). The major archaeal group present at these depths were the ANME-2D archaea  
11 indicating that nitrate-mediated anaerobic oxidation of methane may be especially common  
12 (Haroon et al., 2013). The high abundance of archaea in Olkiluoto is special for this  
13 environment. Archaea have also been quantified from the Outokumpu deep scientific  
14 borehole (Purkamo et al., 2016), but unlike the situation in Olkiluoto the archaeal community  
15 was less than 1% of the total community at best.

16 Previously, using 454 amplicon sequencing, we have observed OTU numbers of  
17 approximately 800 OTUs per sample covering approximately 550 bacterial genera (or  
18 equivalent groups) and approximately 350 archaeal OTUs including approximately 80  
19 different genera (or equivalent groups) (Miettinen et al., 2015). Miettinen et al. (2015) defined  
20 the OTUs 97% sequence homology and the number of sequence reads per sample was at most  
21 in the range of  $10^4$ . In contrast, our sequence read numbers were 10- to 100-fold higher and  
22 the number of OTUs per sample in general 100-fold higher. This indicates that a greater  
23 sequencing depth increases the number of taxa detected from the subsurface environment and  
24 allows us a novel view of the previously rare biosphere. Nevertheless, in comparison to the  
25 high number of OTUs detected the number of identified genera, 651 and 81 bacterial and  
26 archaeal genera, respectively, seems low. On the other hand, this indicates that the sequencing  
27 depth has been sufficient to detect most of the prokaryotic groups present. Nevertheless, the  
28 obtained numbers of OTUs per sample in this study were huge (Table 2). This may reflect the  
29 high level of variability in the short sequence reads of the v6 region used in this study. Huse  
30 et al. (2008) discussed the problem with short sequence reads, because these reads very often  
31 match full-length 16S rRNA gene sequences belonging to several different taxa, and a precise  
32 taxonomic assignment of the short sequence read cannot be done. As shown in our study  
33 taxonomic assignments, such as 'Proteobacteria\_other' were common and may be due to

1 multiple matches for the individual sequence reads obtained in the identification step of the  
2 analysis. However, it may also be possible that the sequences represent novel microbial  
3 clades, the so called Microbial Dark Matter (Solden et al., 2016), that do not have  
4 representatives in the databases yet.

5 In general, the microbial communities at different depth grouped loosely into clusters (Figure  
6 6). Although no clear environmental factor seemed to drive the microbial communities at  
7 different depths, the core communities appeared to be more similar in samples from similar  
8 depths, especially in the bacterial communities (Figure 3). OTUs belonging to both sulphate  
9 reducers (*Desulfobacteraceae*) and sulphur oxidizers (*Thiobacteraceae*) were present in the  
10 bacterial core community. The archaeal core community consisted mostly of methane-  
11 oxidizing ANME-2D archaea. Interestingly, however, their abundance was higher in the  
12 deeper samples. Previous studies on the Finnish deep biosphere has shown that the microbial  
13 communities at different sites vary strongly from each other. Purkamo et al. (2015)  
14 investigated the bacterial and archaeal communities of different fracture zones of the  
15 Outokumpu deep scientific borehole and found that the majority of the bacterial populations  
16 at depths between 180 m and 500 m depth consist of Betaproteobacteria belonging to the  
17 *Commamonadaceae* and the archaeal communities consist of *Methanobacteriaceae* and  
18 *Methanoregula*.

19 The core communities, defined as OTUs present in all the studied samples, accounted for  
20 between 0.2% and 11.7% of the archaeal and 0.4% - 4.1% of the bacterial sequence reads,  
21 respectively. These proportions are surprisingly low and show that most of the archaeal and  
22 bacterial communities (83.3-99.8% and 95.5-99.6% of the archaeal and bacterial sequence  
23 reads, respectively) consist of taxa that are present in only specific samples. Nevertheless, the  
24 short read length and high sequence variability within the v6 region may over estimate this  
25 diversity. Nevertheless, on genus-level between 95 and 99% of the archaeal sequence reads  
26 fell with only 25 genera, which were present in all samples. Likewise for the bacterial  
27 communities, 80 – 97% of the sequence reads belonged to 95 bacterial genera that were  
28 detected in all samples. The number of OTUs found in at least 80% of the samples greatly  
29 outnumbered the number of OTUs present in all samples (i.e. the core community) both in the  
30 archaeal and the bacterial communities. The number of archaeal and bacterial OTUs present  
31 in at least 50% of the samples was only 800 and less than 600, respectively. OTUs present in  
32 at least 20% of all samples (i.e. two to three samples) consisted only of approximately 10,000

1 archaeal and 30,000 bacterial OTUs. Compared to the total number of 468,684 archaeal and  
2 301,458 bacterial OTUs detected in total in these samples, the proportion of rare OTUs  
3 present in only 1 or 2 samples is huge.

4 Our results agree with Sogin et al. (2006) and Magnabosco et al. (2014), who showed that a  
5 relatively small number of taxa dominate deep-sea water and deep groundwater habitats,  
6 respectively, but a rare microbiome consisting of thousands of taxonomically distinct  
7 microbial groups are detected at low abundances. What this means for the functioning of the  
8 deep subsurface is that the microbial communities have the capacity to respond and change  
9 due to changes in environmental conditions. For example, Pedersen et al. (2014) showed that  
10 by adding sulphate to the sulphate-poor but methane-rich groundwater in Olkiluoto the  
11 bacterial population changed over the span of 103 days from a non-SRB community to a  
12 community dominated by SRB. In addition, a change in the geochemical environment  
13 induced by H<sub>2</sub> and methane impacted the size, composition and functions of the microbial  
14 community and ultimately led to acetate formation (Pedersen et al., 2012; Pedersen, 2013;  
15 Pedersen et al., 2014).

16 The metabolic pathways predicted by PICRUSt are far from certain when uncultured and  
17 unculturable deep subsurface microbial communities are concerned. The NSTI values for both  
18 the bacterial and well as the archaeal communities were high indicating that closely related  
19 species to those found in our deep groundwater have yet to be sequenced. This is in  
20 accordance with Langille et al. (2013), who showed that environments containing a high  
21 degree of unexplored microbiota also tend to have high NSTI values. Staley et al. (2014) also  
22 showed in a comparison between PICRUSt and shot gun metagenomic sequencing of riverine  
23 microbial communities that PICRUSt may not be able to correctly assess rare biosphere  
24 functions. Nevertheless, Langille et al. (2013) showed that PICRUSt may predict the  
25 metagenomic content of a microbial community more reliably than shallow metagenomic  
26 sequencing. Although PICRUSt does not give as detailed results as metagenomics or genomic  
27 analyses may give, it is still a useful tool for predicting functions in microbial communities  
28 when the possibility for metagenomics analysis may be impossible, e.g. due to low biomass in  
29 the samples.

30 **Energy metabolism.** Deep subsurface environments are often declared energy deprived  
31 environments dominated by autotrophic microorganisms (Hoehler and Jorgensen, 2013).  
32 However, recent reports indicate that heterotrophic microorganisms play a greater role than

1 the autotrophic microorganisms in Fennoscandian deep crystalline subsurface environments  
2 (Purkamo et al., 2015). Heterotrophic communities with rich fatty acid assimilation strategies  
3 have been reported to fix carbon dioxide on the side of e.g. fermenting activities in order to  
4 replenish the intracellular carbon pool, which otherwise would be depleted. Wu et al. (2015)  
5 also found by metagenomic analyses that fermentation was a major metabolic activity in the  
6 microbial community of Swedish deep groundwater. Our results agree with Purkamo et al.  
7 (2015) that a greater proportion of the microbial community is involved in carbohydrate and  
8 fatty and organic acid oxidation than in fixation of inorganic carbon. Nevertheless,  
9 autotrophic carbon fixation pathways were predicted in the analysis with PICRUSt, indicating  
10 that both the archaeal and bacterial communities include autotrophic members, although these  
11 microorganisms might not be obligate autotrophs. It was also noted that even though evidence  
12 for methane oxidation could not be inferred from the PICRUSt predictions (no *pmoA* genes),  
13 the bacterial community may oxidize formate, which is in agreement with the findings  
14 reported by Wu et al. (2015).

15 Several carbon fixation pathways were predicted in the metagenomes, the Calvin cycle,  
16 reductive TCA (rTCA) cycle and Wood-Ljungdahl (WL) pathway. The WL-pathway is  
17 considered the most ancient autotrophic carbon fixation pathway in bacteria and archaea  
18 (Fuchs 1989, Martin et al. 2008, Berg et al. 2010; Hügler and Sievert, 2011) and was found in  
19 both the bacterial and the archaeal communities. In the archaeal community the Calvin cycle  
20 and the rTCA were especially pronounced in the samples from 296 m, 405 – 423 m and  
21 somewhat lower at 510 – 527 m depth. The bacterial communities are predicted to fix CO<sub>2</sub> at  
22 almost all depths with the exception of 405 m and 559 m depth. Nevertheless, our results  
23 agree with Nyysönen et al. (2014), who showed my metagenomic analysis that the microbial  
24 communities at different depth of the Outokumpu scientific deep drill hole may fix carbon in  
25 several ways, of which the rTCA, the WL pathway and the Calvin cycle were identified.  
26 Magnabosco et al. (2016) showed that the WL pathway was the dominating form of carbon  
27 fixation in metagenomes of 3 km deep Precambrian crust biospheres in South Africa. Dong et  
28 al. (2014) also suggested that microorganisms in low-energy deep subsurface environment  
29 may have several strategies for e.g. carbon fixation, as shown in the *Halomonas sulfidaeris*, in  
30 order to access as many resources as possible. The predicted methane metabolism (methane  
31 and methyl compound consumption) and oxidative phosphorylation were equally strong in the  
32 bacterial community. Sulphur metabolism was not predicted to be a common pathway for  
33 energy in either the archaeal or the bacterial communities. However, PICRUSt predicted

1 bacteria with either assimilative or dissimilative sulphate reduction to be present. Sulphur  
2 oxidation through the *sox* system was in general not predicted, but the *soxD* gene was  
3 predicted and oxidation of thiosulphate to sulphate may be possible (Figure S6). According to  
4 the predicted metagenomes, nitrate may be reduced both through dissimilatory nitrate  
5 reduction to ammonia and through denitrification to nitrous oxide by the bacteria. In addition,  
6 nitrogen may be fixed to ammonia by both archaea and bacteria. The ammonia may be used  
7 as raw material for L-glutamate synthesis.

8 Oxidative phosphorylation was predicted as one of the most prominent energy generating  
9 metabolic pathways in the bacterial community. This indicates that ATP is generated by  
10 electron transfer to a terminal electron acceptor, such as oxygen, nitrate or sulphate. In the  
11 archaeal community the oxidative phosphorylation was not as strongly indicated, but this may  
12 be due to missing data on archaeal metabolism in the KEGG database.

13 The main energy metabolism of the archaeal communities was predicted to be the  
14 methanogenesis, especially at 296 m and 405 m. Methanogenesis was common also at all  
15 other depths except 330 m – 347m, 415 m and 693 m – 798 m. Methane is produced from  
16 CO<sub>2</sub>-H<sub>2</sub> and methanol, and from acetate, although evidence for the acetate kinase enzyme was  
17 lacking. Methanogenesis from methylamines may also be possible, especially at 296 m and  
18 405 m. Methane oxidation using methane monooxygenases and methanol dehydrogenases does  
19 not occur in either bacterial or archaeal communities. It should be noted, however, that the  
20 ANME-2D archaea are likely to use the methanogenesis pathway in the reverse for oxidizing  
21 methane anaerobically to carbon dioxide (Haroon et al., 2013). The produced carbon dioxide  
22 may be fixed by the same archaea and turned in to acetate, which may serve as carbon  
23 substrate and electron donor and acceptor for a large variety of microorganisms in the  
24 groundwater.

25 **Carbohydrate metabolism.** Glycolysis/gluconeogenesis was one of the most common  
26 carbohydrate-metabolizing pathways predicted for both the archaeal and bacterial  
27 communities (Figure 8). Pyruvate from glycolysis is oxidized to acetyl-CoA by both archaea  
28 and bacteria and used in the TCA cycle. The TCA cycle provides for example raw material  
29 for many amino acids, such as lysine and glutamate. The butanoate and propanoate  
30 metabolisms were also common in the bacterial communities, indicating fermentative  
31 metabolism and capability of fatty acid oxidation.

1 **Amino acid metabolism.** The predicted metagenomes indicated that non-essential amino  
2 acids, such as alanine, aspartate and glutamate may be produced from ammonia and pyruvate  
3 or oxaloacetate especially in the archaeal populations. In the archaeal population proline  
4 appeared to be produced from glutamate. Despite the predicted low use of sulphate as energy  
5 source in the microbial communities sulphate and other sulphur compounds could be taken up  
6 for the production of the amino acids cysteine and methionine by both the archaeal and the  
7 bacterial communities. A higher predicted relative abundance of genes involved in aromatic  
8 amino acid synthesis (phenylalanine, tyrosine, tryptophane) was seen in the archaeal than in  
9 the bacterial communities. Both the archaeal and the bacterial communities were predicted to  
10 synthesise branched chained amino acids (isoleucine, leucine and valine), but only the  
11 bacteria degrade them. Especially proteobacteria have been shown to be able to use the  
12 branched chained amino acids (isoleucine, leucine and valine) and short chained fatty acids  
13 (acetate, butyrate, propionate) as sole energy and carbon source (Kazakov et al., 2009). The  
14 branched chained amino acids function as raw material in the biosynthesis of branched  
15 chained fatty acids, which regulate the membrane fluidity of the bacterial cell. In salt stress  
16 conditions, the proportion of branch-chained fatty acids in the membranes decreases.

17 **Membrane transport.** According to the predicted metagenomes, the microbial cells transport  
18 sulphate into the cell, but do not take up nitrate. Nitrogen is taken up as glutamate but not as  
19 urea. Iron is taken up by an Fe(III) transport system and an iron complex transport system in  
20 the bacterial communities, but generally only by the iron complex transport system in  
21 archaea. However, Fe(III) transport system may also exist in the archaeal communities at 405  
22 m to 423 m depth, where also some manganese/iron transport systems could be found.  
23 According to the metagenomes predictions, molybdate and phosphate is transported into the  
24 cell by molybdate and phosphate ATPases, respectively. Nickel is taken up mainly by a  
25 nickel/peptide transport system but also to some extent by a cobalt/nickel transport system.  
26 Zinc is taken up to some extent by a zinc transport system, but transport systems for  
27 manganese, manganese/iron, manganese/zinc/iron, or iron/zinc/copper were not predicted.  
28 Ammonia was predicted to be taken up by an Amt transport system.

29

## 30 **5 Conclusions**

31 The wide diversity of microbial groups in the deep Fennoscandian groundwater at the  
32 Olkiluoto site revealed that the majority of the microbial community present belong to only a

1 few microbial taxa while the greatest part of the microbial diversity is represented by low  
2 abundance and rare microbiome taxa. The core community was present in all tested samples  
3 from different depths, but the relative abundance of the different taxa varied in the different  
4 samples. Nevertheless, the proportion of OTUs found in only a small proportion (e.g. 20%) of  
5 the samples far surpassed the number of OTUs included in the core communities.  
6 Fermentation or oxidation of fatty acids was a common carbon cycling and energy harvesting  
7 metabolic pathways in the bacterial communities whereas the archaea may either produce or  
8 consume methane. Glycolysis/gluconeogenesis was predicted to be common in both the  
9 archaeal and bacterial communities. In addition both the bacterial and archaeal communities  
10 were predicted to contain several different common carbon fixation pathways, such as the  
11 Calvin cycle and the reductive TCA and the Wood-Ljungdahl pathway.

12

### 13 **Acknowledgements**

14 The Illumina sequencing data were made possible by the Deep Carbon Observatory's Census  
15 of Deep Life supported by the Alfred P. Sloan Foundation. Illumina sequencing was  
16 performed at the Marine Biological Laboratory (Woods Hole, MA, USA) and we are grateful  
17 for the assistance of Mitch Sogin, Susan Huse, Joseph Vineis, Andrew Voorhis, Sharon Grim,  
18 and Hilary Morrison at MBL and Rick Colwell, OSU. MB was supported by the Academy of  
19 Finland (project 261220).

20

## 1 **References**

- 2 Bano, N., Ruffin, S., Ransom, B. and Hollibaugh, J.T.: Phylogenetic composition of Arctic  
3 ocean archaeal assemblages and comparison with Antarctic assemblages. *Appl. Environ.*  
4 *Microbiol.* 70, 781-789, 2004.
- 5 Barns, S.M., Fundyga, R.E., Jeffries, M.W. and Pace, N.R.: Remarkable archaeal diversity  
6 detected in Yellowstone National Park hot spring environment. *Proc. Natl. Acad. Sci. USA*  
7 91, 1609-1613, 1994.
- 8 Berg, I.A., Kockelkorn, D., Ramos-Vera, W.H., Say, R.F., Zarzycki, J., Hügler, M., Alber,  
9 B.E. and Fuchs, G.: Autotrophic carbon fixation in archaea. *Nat. Rev. Microbiol.* 8, 447-  
10 460, 2010.
- 11 Blankenberg, D., Von Kuster, G., Coraor, N., Ananda, G., Lazarus, R., Mangan, M.,  
12 Nekrutenko, A. and Taylor, J.: Galaxy: a web-based genome analysis tool for  
13 experimentalists. *Curr. Protoc. Mol. Biol.*, 19, 19.10.1-21, 2010.
- 14 Bomberg, M., Nyysönen, M., Nousiainen, A., Hultman, J., Paulin, L., Auvinen, P. and  
15 Itävaara, M.: Evaluation of molecular techniques in characterization of deep terrestrial  
16 biosphere. *Open J Ecol* 4, 468-487, 2014.
- 17 Bomberg, M., Nyysönen, M., Pitkänen, P., Lehtinen, A. and Itävaara, M.: Active microbial  
18 communities inhabit sulphate-methane interphase in deep bedrock fracture fluids in  
19 Olkiluoto, Finland. *Biomed. Res. Int.*, Article ID 979530, 2015.  
20 <http://dx.doi.org/10.1155/2015/979530>.
- 21 Caporaso, J. G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F. D., Costello, E. K.,  
22 Fierer, N., Gonzalez Pena, A., Goodrich, J. K., Gordon, J. I., Huttley, G. A., Kelley, S. T.,  
23 Knights, D., Koenig, J. E., Ley, R. E., Lozupone, C. A., McDonald, D., Muegge, B. D.,  
24 Pirrung, M., Reeder, J., Sevinsky, J. R., Turnbaugh, P. J., Walters, W. A., Widmann, J.,  
25 Yatsunencko, T., Zaneveld, J. and Knight, R. QIIME allows analysis of high-throughput  
26 community sequencing data. *Nature Methods*, 7, 335-336, 2010.
- 27 Cleary, D. F., Becking, L. E., Polónia, A. R., Freitas, R. M. and Gomes, N. C.: Composition  
28 and predicted functional ecology of mussel-associated bacteria in Indonesian marine lakes.  
29 *Antonie van Leeuwenhoek*, 107, 821-834, 2015.
- 30 DeSantis, T. Z., Hugenholtz, P., Larsen, N., Rojas, M., Brodie, E. L., Keller, K., Huber, T.,  
31 Dalevi, D., Hu, P. and Andersen, G. L. Greengenes, a Chimera-Checked 16S rRNA Gene  
32 Database and Workbench Compatible with ARB. *Appl. Environ. Microbiol.*, 72, 5069-72,  
33 2006.
- 34 Dong, Y., Kumar, C.G., Chia, N., Kim, P.-J., Miller, P.A., Price, N.D., Cann, I.K.O., Flynn,  
35 T.M., Sanford, R.A., Krapac, I.G., Locke, R.A.II, Hong, P.-Y., Tamaki, H., Liu, W.-T.,  
36 Mackie, R.I., Hernandez, A.G., Wright, C.L., Mikel, M.A., Walker, J.L., Sivaguru, M.,  
37 Fried, G., Yannarell, A.C., Fouke, B.W. Halomonas sul- fidaeris-dominated microbial  
38 community inhabits a 1.8 km-deep subsurface Cambrian Sandstone reservoir. *Environ.*  
39 *Microbiol.* 16, 1695-1708, 2014.
- 40 Eren, A.M., Maignien, L., Sul, W.J., Murphy, L.G., Grim, S.L., Morrison, H.G. and Sogin,  
41 M.L. Oligotyping: differentiating between closely related microbial taxa using 16S rRNA  
42 gene data. *Meth. Ecol. Evol.*, 4, 1111-1119, 2013.
- 43 Fuchs, G.: Alternative pathways of autotrophic CO<sub>2</sub> fixation. In: *Autotrophic Bacteria*, ed.

- 1 HG Schlegel, B Bowien, pp. 365–82. Berlin: Springer, 1989.
- 2 Giardine, B., Riemer, C., Hardison, R.C., Burhans, R., Elnitski, L., Shah, P., Zhang, Y.,  
3 Blankenberg, D., Albert, I., Taylor, J., Miller, W., Kent, W.J., Nekrutenko, A.: Galaxy: a  
4 platform for interactive large-scale genome analysis. *Genome Res.*, 15, 1451-5, 2005.
- 5 Goecks, J., Nekrutenko, A., Taylor, J. and The Galaxy Team: Galaxy: a comprehensive  
6 approach for supporting accessible, reproducible, and transparent computational research in  
7 the life sciences. *Genome Biol.* 11, R86, 2010.
- 8 Hallbeck, L. and Pedersen, K.: Culture-dependent comparison of microbial diversity in deep  
9 granitic groundwater from two sites considered for a Swedish final repository of spent  
10 nuclear fuel. *FEMS Microbiol. Ecol.*, 81, 66-77, 2012.
- 11 Hammer, Ø., Harper, D.A.T. and Ryan, P.D.: PAST: Paleontological statistics software  
12 package for education and data analysis. *Palaeontologia Electronica* 4: 9pp. [http://palaeo-](http://palaeo-electronica.org/2001_1/past/issue1_01.htm)  
13 [electronica.org/2001\\_1/past/issue1\\_01.htm](http://palaeo-electronica.org/2001_1/past/issue1_01.htm), 2001.
- 14 Haroon, M.F., Hu, S., Shi, Y., Imelfort, M., Keller, J., Hugenholtz, P., Yuan, Z. and Tyson,  
15 G.W.: Anaerobic oxidation of methane coupled to nitrate reduction in a novel archaeal  
16 lineage. *Nature*, 500, 567-570, 2013.
- 17 Hoehler, T.M. and Jørgensen, B.B.: Microbial life under extreme energy limitation. *Nat Rev*  
18 *Microbiol* 11, 83-94, 2013.
- 19 Hügler, M. and Sievert, S. M.: Beyond the Calvin cycle: autotrophic carbon fixation in the  
20 ocean. *Mar. Sci.*, 3, 2011.
- 21 Huse, S.M., Dethlefsen, L., Huber, J.A., Welch, D.M., Relman, D.A. and Sogin, M.L.  
22 Exploring microbial diversity and taxonomy using SSU rRNA hypervariable tag sequencing.  
23 *PLoS Genet*, 4, p.e1000255, 2008.
- 24 Kärki, A. and Paulamäki, S.: Petrology of Olkiluoto. Posiva 2006-02, 2006.
- 25 Kazakov, A.E., Rodionov, D.A., Alm, E., Arkin, A.P., Dubchak, I. and Gelfand, M.S.:  
26 Comparative genomics of regulation of fatty acid and branched-chain amino acid utilization  
27 in proteobacteria. *J. Bacteriol.*, 191, 52-64, 2009.
- 28 Kutvonen, H.: Nitrogen-cycling bacteria in groundwater of the low and medium active nuclear  
29 waste repository in Olkiluoto, Finland (In Finnish) M.Sc. Thesis, Faculty of Agriculture and  
30 Forestry, University of Helsinki, Finland, 2015.
- 31 Langille, M. G. I., Zaneveld, J., Caporaso, J. G., McDonald, D., Knights, D., Reyes, J. A.,  
32 Clemente, J. C., Burkepille, D. E., Vega Thurder, R. L., Knight, R., Beiko, R. G. and  
33 Huttenhower, C.: Predictive functional profiling of microbial communities using 16S rRNA  
34 marker gene sequences. *Nature Biotechnol.*, 31, 814-821, 2013.
- 35 Magnabosco, C., Tekere, M., Lau, M. C., Linage, B., Kuloyo, O., Erasmus, M., Cason, E.,  
36 van Heerden, E., Borgonie, G. and Onstott, T. L. Comparisons of the composition and  
37 biogeographic distribution of the bacterial communities occupying South African thermal  
38 springs with those inhabiting deep subsurface fracture water. *Front. Microbiol*, 5, 10-3389,  
39 2014.
- 40 Magnabosco, C., Ryan, K., Lau, M.C., Kuloyo, O., Lollar, B.S., Kieft, T.L., van Heerden, E.  
41 and Onstott, T.C. A metagenomic window into carbon metabolism at 3 km depth in  
42 Precambrian continental crust. *ISME J.*, 10, 730-741, 2016.

- 1 Martin, W., Baross, J., Kelley, D. and Russell, M.J.: Hydrothermal vents and the origin of  
2 life. *Nat. Rev. Microbiol.* 6, 805–14, 2008.
- 3 McMurdie, P.J.; Holmes, S.: Shiny-phyloseq: Web Application for Interactive Microbiome  
4 Analysis with Provenance Tracking. *Bioinformatics* 31, 282–283, 2015.
- 5 Miettinen H., Bomberg, M., Nyysönen, M., Salavirta, H. Sohlberg, E. Vikman, M. and  
6 Itävaara, M.: The Diversity of Microbial Communities in Olkiluoto Bedrock Groundwaters  
7 2009–2013. Olkiluoto, Finland, Posiva Oy. Posiva Working Report 2015-12, 1-160, 2015.  
8 [www.posiva.fi](http://www.posiva.fi)
- 9 Muyzer, G., de Waal, E.C., and Uitterlinden, A.G.: Profiling of complex microbial  
10 populations by denaturing gradient gel electrophoresis analysis of polymerase chain  
11 reaction-amplified genes coding for 16S rRNA. *Appl. Environ. Microbiol.* 59, 695–700,  
12 1993.
- 13 Nyysönen, M., Bomberg, M., Kapanen, A., Nousiainen, A., Pitkänen, P. and Itävaara, M.:  
14 Methanogenic and sulphate-reducing microbial communities in deep groundwater of  
15 crystalline rock fractures in Olkiluoto, Finland. *Geomicrobiol. J.*, 29, 863–878, 2012.
- 16 Nyysönen, M., Hultman, J., Ahonen, L., Kukkonen, I.T., Paulin, L., Laine, P., Itävaara, M.  
17 and Auvinen, P.: Taxonomically and functionally diverse microbial communities in deep  
18 crystalline rocks of the Fennoscandian shield. *ISME J.*, 8, 126-138, 2014.
- 19 Oksanen, J.; Blanchet, F.G.; Friendly, M.; Kindt, R.; Legendre, P.; McGlinn, D.; Minchin,  
20 P.R.; O’Hara, R.B.; Simpson, G.L.; Solymos, P.; Stevens, M.H.H.; Szoecs, E.; Wagner, H.:  
21 vegan: Community Ecology Package. R package version 2.4-0. 2016. [http://CRAN.R-](http://CRAN.R-project.org/package=vegan)  
22 [project.org/package=vegan](http://CRAN.R-project.org/package=vegan).
- 23 Osburn, M.R., LaRowe, D.E., Momper, L.M. and Amend, J.P. Chemolithotrophy in the  
24 continental deep subsurface: Sanford Underground Research Facility (SURF), USA. *Front.*  
25 *Microbiol.*, 5, 610, 2014.
- 26 Pedersen, K., Arlinger, J., Eriksson, S., Hallbeck, A., Hallbeck, L. and Johansson, J.:  
27 Numbers, biomass and cultivable diversity of microbial populations relate to depth and  
28 borehole-specific conditions in groundwater from depths of 4–450 m in Olkiluoto, Finland.  
29 *ISME J.*, 2, 760-775, 2008.
- 30 Pedersen, K., Bengtsson, A., Edlund, J. and Eriksson, L.: Sulphate-controlled diversity of  
31 subterranean microbial communities over depth in deep groundwater with opposing  
32 gradients of sulphate and methane. *Geomicrobiol. J.*, 31, 617-631, 2014.
- 33 Pedersen, K.: Metabolic activity of subterranean microbial communities in deep granitic  
34 groundwater supplemented with methane and H<sub>2</sub>. *ISME J.*, 7, 839-849, 2013.
- 35 Pedersen, K.: Subterranean microbial populations metabolize hydrogen and acetate under in  
36 situ conditions in granitic groundwater at 450 m depth in the Äspö Hard Rock Laboratory,  
37 Sweden. *FEMS Microbiol. Ecol.*, 81, 217-229, 2012.
- 38 Posiva. Olkiluoto site description 2011. POSIVA 2011-02, 2013.
- 39 Purkamo, L., Bomberg, M., Nyysönen, M., Kukkonen, I., Ahonen, L., Kietäväinen, R.,  
40 Itävaara, M.: Retrieval and analysis of authentic microbial communities from packer-  
41 isolated deep crystalline bedrock fractures: evaluation of the method and time of sampling.  
42 *FEMS Microbiol. Ecol.*, 85, 324-337, 2013.

- 1 Purkamo, L., Bomberg, M., Nyysönen, M., Kukkonen, I., Ahonen, L. and Itävaara, M.:  
2 Heterotrophic communities supplied by ancient organic carbon predominate in deep  
3 Fennoscandian bedrock fluids. *Microb. Ecol.*, 69, 319-332, 2015.
- 4 R Core Team (2013). R: A language and environment for statistical computing. R Foundation  
5 for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0, <http://www.R-project.org/>.
- 6 Rajala, P., Bomberg, M., Kietäväinen, R., Kukkonen, I., Ahonen, L., Nyysönen, M.,  
7 Itävaara, M.: Deep subsurface microbes rapidly reactivate in the presence of C-1  
8 compounds. *Microorganisms*, 3, 17-33, 2015.
- 9 Schloss, P.D., Westcott, S.L., Ryabin, T., Hall, J.R., Hartmann, M., Hollister, E.B.,  
10 Lesniewski, R.A., Oakley, B.B., Parks, D.H., Robinson, C.J., Sahl, J.W., Stres, B.,  
11 Thallinger, G.G., Van Horn, D.J., Weber, C.F.: Introducing MOTHUR: open-source,  
12 platform-independent, community-supported software for describing and comparing  
13 microbial communities. *Appl. Environ. Microbiol.*, 75, 7537–41, 2009.
- 14 Sedighi, M., Bennett, D., Masum, S.A., Thomas, H.R., Johansson, E.: Analysis of temperature  
15 data at the Olkiluoto. Posiva Working Report 2013-58, 1-78, 2013
- 16 Shade, A. and Handelsman, J.: Beyond the Venn diagram: the hunt for a core microbiome.  
17 *Environ. Microbiol.*, 14, 4-12, 2012.
- 18 Sogin, M.L., Morrison, H.G, Huber, J.A, Welch, D.M., Huse, S.M., Neal, P.R., Arrieta, J.M.  
19 and Herndl, G.J.: Microbial diversity in the deep sea and the underexplored “rare  
20 biosphere”. *Proc. Natl. Acad. Sci. USA*, 103, 12115-12120, 2006.
- 21 Sohlberg, E., Bomberg, M., Miettinen, H., Nyysönen, M., Salavirta, H., Vikman, M. and  
22 Itävaara, M.: Revealing the unexplored fungal communities in deep groundwater of  
23 crystalline bedrock fracture zones in Olkiluoto, Finland. *Front. Microbiol.*, 6, 573, 2015.
- 24 Solden, L., Lloyd, K., Wrighton, K.: The bright side of microbial dark matter: lessons  
25 learned from the uncultivated majority. *Curr. Op. Microbiol.*, 31, 217-226, 2016.
- 26 Staley, C., Gould, T.J., Wang, P., Phillips, J., Cotner, J.B., Sadowsky, M.J. Core functional  
27 traits of bacterial communities in the Upper Mississippi River show limited variation in  
28 response to land cover. *Front. Microbiol.* 5, 414, 2014.
- 29 Tsitko, I., Lusa, M., Lehto, J., Parviainen, L., Ikonen, A.T.K, Lahdenperä, A.-M. and  
30 Bomberg, M. The variation of microbial communities in a depth profile of an acidic,  
31 nutrient-poor boreal bog in southwestern Finland. *Open J. Ecol.*, 4, 832-859, 2014.
- 32 Weisburg, W.G., Barns, S.M., Pelletier, D.A. and Lane, D.J. 16S ribosomal DNA  
33 amplification for phylogenetic study. *J. Bacteriol.*, 173, 697-703, 1991.
- 34 Wu, X., Holmfeldt, K., Hubalek, V., Lundin, D., Åström, M., Bertilsson, S. and Dopson, M.  
35 Microbial metagenomes from three aquifers in the Fennoscandian shield terrestrial deep  
36 biosphere reveal metabolic partitioning among populations. *ISME J.*, on line.

37

- 1 Table 1. Geochemical and microbiological measurements from 12 different water conductive fractures in the bedrock of Olkiluoto, Finland.
- 2 The different drillholes are presented at the top of the table. The data is compiled from Posiva (2013) and Miettinen et al. (2015). NPOC=non-purgable organic carbon, DIC=dissolved inorganic carbon, TDS=total dissolved solids, TNC=total number of cells.
- 3

Drillhole	OL-KR13	OL-KR6	OL-KR3	OL-KR23	OL-KR5	OL-KR49	OL-KR9	OL-KR9	OL-KR2	OL-KR1	OL-KR44	OL-KR29
Sampling date	3/11/2010	18/5/2010	29/8/2011	15/12/2009	16/10/2012	14/12/2009	31/10/2011	29/8/2011	27/1/2010	26/1/2010	15/1/2013	18/5/2010
Depth (m)	296	328	340	347	405	415	423	510	559	572	693	798
Alkalinity (mEq/L)	2.19	0.37	0.47	0.05	0.27	0.16	0.18	0.13	0.29	0.23	0.49	0.13
Ec (mS/m)	897	1832	1047	2190	2240	2670	2300	2960	4110	3770	6690	7820
pH	7.9	7.9	7.9	7.5	7.9	8.1	7.7	8.1	8.6	7.8	7.5	7.3
NPOC (mg L <sup>-1</sup> )	10	0	12	5.1	19	3	5.1	6.6	11	5	110	10
DIC (mg L <sup>-1</sup> )	27	4.1	4.1	3.9	0	3	3	0	3.75	3.75	6.5	81
HCO <sub>3</sub> (mg L <sup>-1</sup> )	134	22.6	25	17.1	16	9.8	11.6	7.3	17.7	14	30	424
N <sub>tot</sub> (mg L <sup>-1</sup> )	0.71	0	1.1	0.42	1.2	0.16	0.38	0.66	1.1	0.41	10	3.1
NH <sub>4</sub> <sup>+</sup> (mg L <sup>-1</sup> )	0.07	0.03	0.03	0	0	0	0.05	0	0.02	0.04	0.08	0.08
S <sub>tot</sub> (mg L <sup>-1</sup> )	31	130	12	1.7	1.7	0	4.8	0	0	0	4	0
SO <sub>4</sub> <sup>2-</sup> (mg L <sup>-1</sup> )	79.5	379	32	2.9	3	1.4	13.7	0.9	0.5	0.5	9.6	2
S <sub>2</sub> <sup>-</sup> (mg L <sup>-1</sup> )	5.1		0.38	0.62	2	0.02	0.36	0	0.02	0.13	0.02	0.02
Fe <sub>tot</sub> (mg L <sup>-1</sup> )	0.0042	0.0037	0.022	0.062	0.2	0.71	0.036	0.02	0	0.49	1.2	560
Fe(II) (mg L <sup>-1</sup> )	0	0	0.02	0.08	0.21	0.53	0.06	0.02	0.02	0.04	1.2	0.46
TDS (mg L <sup>-1</sup> )	4994	10670	5656	12710	12880	15900	13430	18580	25500	23260	37410	53210
K (mg L <sup>-1</sup> )	8.2	9.3	7.6	8.3	18	27	12	17	19	20	24	27
Mg (mg L <sup>-1</sup> )	35	77	17	55	68	19	32	41	18	52	33	136
Ca (mg L <sup>-1</sup> )	460	1100	290	2100	1750	2700	2260	2930	4600	3700	7680	10000
Cl (mg L <sup>-1</sup> )	2920	6230	3400	7930	7950	9940	8220	11500	15700	14600	22800	33500
Na (mg L <sup>-1</sup> )	1320	2800	1850	2530	2990	3110	2790	3970	4980	4720	6570	9150
TNC ml <sup>-1</sup>	4.2 × 10 <sup>5</sup>	1.0 × 10 <sup>5</sup>	2.4 × 10 <sup>5</sup>	2.5 × 10 <sup>5</sup>	2.1 × 10 <sup>5</sup>	1.5 × 10 <sup>4</sup>	na	2.9 × 10 <sup>4</sup>	5.9 × 10 <sup>4</sup>	8.7 × 10 <sup>4</sup>	5.5 × 10 <sup>4</sup>	2.3 × 10 <sup>4</sup>
16S qPCR ml <sup>-1</sup>												
bacteria	7.0 × 10 <sup>5</sup>	9.5 × 10 <sup>3</sup>	2.0 × 10 <sup>4</sup>	3.6 × 10 <sup>5</sup>	4.9 × 10 <sup>4</sup>	1.3 × 10 <sup>4</sup>	7.2 × 10 <sup>4</sup>	1.5 × 10 <sup>5</sup>	1.4 × 10 <sup>5</sup>	1.9 × 10 <sup>4</sup>	3.2 × 10 <sup>4</sup>	1.5 × 10 <sup>4</sup>
archaea	5.8 × 10 <sup>3</sup>	2.0 × 10 <sup>4</sup>	9.9 × 10 <sup>3</sup>	6.3 × 10 <sup>4</sup>	6.2 × 10 <sup>3</sup>	1.5 × 10 <sup>2</sup>	4.4 × 10 <sup>4</sup>	5.2 × 10 <sup>2</sup>	7.5 × 10 <sup>2</sup>	3.0 × 10 <sup>3</sup>	2.6 × 10 <sup>1</sup>	2.8 × 10 <sup>2</sup>

1 Table 2a. The total number of sequence reads, observed and estimated (Chao1, ACE) number of OTUs, number of singleton and doubleton  
2 OTUs, and Shannon diversity index per sample of the bacterial 16S rRNA gene data set. The analysis results are presented for both the total  
3 number of sequence reads per sample as well as for data normalized according to the sample with the lowest number of sequence reads, i.e.  
4 140,000 random sequences per sample.

Bacteria		All sequences						Normalized to 140,000 sequences					
Sample	Number of sequence reads	Observed OTUs	Chao1	ACE	Singletons	Doubletons	Shannon	Observed OTUs	Chao1	ACE	Singletons	Doubletons	Shannon
OLKR13/296m	786,346	79,527	87,188	91,360	18,025	21,203	13	37,045	74,288	84,530	22,445	6,762	13
OLKR6/328m	188,812	29,411	35,018	37,269	9,209	7,561	13	26,442	34,964	37,626	10,420	6,369	13
OLKR3/340m	345,433	52,381	53,238	54,961	5,789	19,557	14	39,309	57,793	64,021	19,287	10,061	13
OLKR23/347m	485,154	33,257	37,175	38,895	8,000	8,166	11	20,494	34,268	37,305	10,641	4,109	11
OLKR5/405m	141,886	40,445	70,520	78,340	22,166	8,167	14	40,145	70,288	78,232	22,086	8,090	14
OLKR49/415m	184,052	38,275	49,758	53,525	14,799	9,535	13	34,117	48,804	52,938	15,372	8,043	13
OLKR9/423m	175,295	36,412	44,452	47,571	12,357	9,494	14	33,596	44,496	48,161	13,489	8,345	14
OLKR9/510m	241,312	41,545	51,348	54,535	14,251	10,357	13	33,208	49,115	53,631	15,592	7,640	13
OLKR2/559m	257,789	45,456	72,269	78,325	22,550	9,481	13	32,600	62,318	69,573	19,071	6,118	12
OLKR1/572m	210,659	29,804	35,362	37,491	9,197	7,607	12	25,703	34,934	37,682	10,650	6,142	12
OLKR44/693m	303,058	31,410	31,589	32,188	2,005	11,200	12	25,937	33,448	36,295	10,346	7,124	12
OLKR29/798m	221,524	37,989	45,126	48,042	11,991	10,071	13	31,911	44,957	48,533	14,078	7,594	13

5

1 Table 2b. The total number of sequence reads, observed and estimated (Chao1, ACE) number of OTUs, number of singleton and doubleton  
 2 OTUs, and Shannon diversity index per sample of the archaeal 16S rRNA gene data set. The analysis results are presented for both the total  
 3 number of sequence reads per sample as well as for data normalized according to the sample with the lowest number of sequence reads, i.e.  
 4 17,000 random sequences per sample.


Sample	Number of sequence reads	All sequences						Normalized to 17,000 sequences					
		Observed OTUs	Chao1	ACE	Singletons	Doubletons	Shannon	Observed OTUs	Chao1	ACE	Singletons	Doubletons	Shannon
OLKR13/296m	507,373	27,111	29,516	30,699	5,835	7,076	10	3,957	13,380	15,062	2,867	435	10
OLKR6/328m	446,380	21,597	22,930	23,781	3,861	5,588	10	3,776	11,705	14,020	2,748	475	9
OLKR3/340m	271,699	25,491	32,299	34,231	9,205	6,221	11	4,955	15,044	17,238	3,546	622	10
OLKR23/347m	395,339	20,800	22,403	23,214	4,083	5,199	10	3,919	11,855	13,323	2,755	477	9
OLKR5/405m	769,026	21,127	22,235	23,078	3,515	5,574	9	2,596	10,114	10,078	1,852	227	9
OLKR49/415m	210,545	22,600	23,372	24,004	2,975	5,733	12	7,023	17,088	19,874	4,738	1,114	12
OLKR9/423m	697,360	22,014	22,527	23,082	2,381	5,520	9	3,180	9,617	10,586	2,224	383	9
OLKR9/510m	169,142	12,709	12,782	12,960	713	3,488	11	4,879	11,205	13,215	3,148	782	11
OLKR2/559m	100,101	15,359	24,950	27,026	7,840	3,203	11	5,119	14,497	16,488	3,548	670	11
OLKR1/572m	1,213,360	28,884	33,207	34,832	7,846	7,118	9	2,273	9,233	9,923	1,631	190	9
OLKR44/693m	17,716	6,436	8,748	9,750	2,890	1,805	12	6,325	8,743	9,804	2,921	1,763	12
OLKR29/798m	98,770	15,641	16,720	17,483	3,158	4,617	12	6,951	14,655	17,184	4,483	1,303	12

5


1 Table 3. The nearest sequenced taxon index (NSTI) values for the archaeal and bacterial  
 2 communities in the 12 different samples according to PICRUSt. The NSTI value describes the  
 3 sum of phylogenetic distances of each OTU to its nearest relative with a sequenced reference  
 4 genome, and measures substitutions per site in the 16S rRNA gene and the weighted the  
 5 frequency of the each OTU in a sample dataset. A higher NSTI value indicates greater  
 6 distance to the closest sequenced relatives of the OTUs in each sample.  
 7

Sample	Archaea	Bacteria
OLKR13/296m	0.19	0.17
OLKR6/328m	0.17	0.15
OLKR3/340m	0.20	0.12
OLKR23/347m	0.25	0.08
OLKR5/405m	0.18	0.11
OLKR49/415m	0.22	0.12
OLKR9/423m	0.14	0.12
OLKR9/510m	0.21	0.11
OLKR2/559m	0.21	0.16
OLKR1/572m	0.21	0.14
OLKR44/693m	0.21	0.05
OLKR29/798m	0.29	0.15


8  
 9  
 10




11  
 12 Figure 1. Map of Olkiluoto. The boreholes used in this study are marked with a turquoise  
 13 triangle and the attached black line depicts the direction of the borehole. (with courtesy of  
 14 Pöyry Oy, Nov 17<sup>th</sup>, 2015 by Eemeli Hurmerinta)  
 15




1  
2 Figure 2. The concentration of A) microbial cells  $\text{mL}^{-1}$  determined by epifluorescence  
3 microscopy and the estimated concentration of B) bacterial and C) archaeal 16S rRNA gene  
4 copies  $\text{mL}^{-1}$  groundwater determined by qPCR in water conductive fractures situated at  
5 different depths in the Olkiluoto bedrock.




6  
7 Figure 3. The core A) archaeal and B) bacterial community OTUs detected from at least 80%  
8 of the samples with heatmaps on the abundance of the C) archaeal and D) bacterial core  
9 community profiles. In A) and B) the OTUs are stacked in the columns according to the  
10 number of sequence reads, with the most abundant OTUs at the bottom of the columns. The  
11 OTU segments of the columns are colored according to the family to which they belong. Each  
12 Order is presented as a separate column for each sample.



1  
 2 Figure 4. The number of shared A) archaeal and B) bacterial OTUs in the 12 different  
 3 samples. The number of shared OTUs is shown on the Y-axis and the proportion of samples  
 4 on the X-axis.



5  
 6 Figure 5. A UPGMA cladogram clustering the samples based on the 1000 biggest OTUs of  
 7 the A) archaeal and B) bacterial OTU profile according to the Bray Curtis distance model.  
 8 Black indicates low abundance and red colour indicates high abundance. C) and D) show the  
 9 corresponding physicochemical parameters as shown in Table 1, with the lowest values in  
 10 green, medium values in yellow and high values in red.



1


2

3 Figure 6. Principal coordinates analysis (PCoA) on the whole OTU profiles of the different  
 4 samples based on Bray Curtis distance model for the A) archaeal and B) bacterial  
 5 communities. The points indicate water type, where the circle is for brackish sulphate-rich  
 6 water, the square is for brackish chloride-rich water and the diamond is for saline water. The

1 colouring of the points are according to concentration of total dissolved solids (TDS) as  
2 indicated in the upper left corner of figure A.


3

A.



4

B.




5

6 Figure 7. The relative abundance of predicted genes of the most abundant pathways identified  
7 in the A) archaeal and B) bacterial populations in the PICRUSt analysis. The pathways are


1 presented according to KEGG. The samples are ordered according to depth, with OL-  
 2 KR13/296m as innermost and OL-KR29/798m as the outermost sample.

3




4  
 5 Figure 8. The microbial carbon metabolism pathway according to KEGG. The predicted  
 6 genes combined from all samples were plotted on the map. Green arrows indicate enzymes  
 7 predicted only in the archaeal communities, red arrows indicate genes predicted only in the  
 8 bacterial communities, black arrows show enzymes predicted in both the archaeal and  
 9 bacterial communities.

1  
2  
3  
4 **Supplementary figures**



5  
6 Figure S1. The rarefaction curves of observed bacterial (left pane) and archaeal (right pane)  
7 OTUs in each sample generated on sequence data normalized to 140,000 reads for bacteria  
8 and 17,000 reads for archaea.


9



10  
11 Figure S2. The Principal Coordinates Analysis as affected by sample size for the A) archaea  
12 and B) bacteria. The sample positions on the plots are as described in Figure 6. E) and F)  
13 display the proportion of variance for 12 dimensions, of which 1 and 2 were used for plotting  
14 the PCoA graphs in A) and B) and in Figure 6.








1  
2 Figure S5. The predicted genes of enzymes included in the microbial nitrogen metabolism  
3 according to KEGG. Enzymes predicted from the bacterial communities are shown in red,  
4 archaeal communities in green and enzymes predicted from both archaeal and bacterial  
5 communities in green/red. Enzymes not predicted from either community are shown in blue  
6 or white.

7

8



1

2 Figure S6. The predicted genes of enzymes included in the microbial sulphur metabolism


3 according to KEGG. Enzymes predicted from the bacterial communities are shown in red,

4 archaeal communities in green and enzymes predicted from both archaeal and bacterial

5 communities in green/red. Enzymes not predicted from either community are shown in blue

6 or white.

7



1  
2  
3  
4  
5

Figure S7. The genes of ABC transporters predicted from the bacterial (pink), archaeal (green) or both (pink/green) communities. Genes not predicted in any of the communities are shown in blue.