

1 **Microbial communities and their predicted metabolic**
2 **characteristics in deep fracture groundwaters of the crystalline**
3 **bedrock at Olkiluoto, Finland**

4

5 **M. Bomberg¹, T. Lamminmäki² and M. Itävaara¹**

6 [1] VTT Technical Research Centre of Finland, P.O. Box 1000, FIN-02044 VTT, Finland

7 [2] Posiva Oy, Olkiluoto, 27160 Eurajoki, Finland

8 Correspondence to: M. Bomberg (malin.bomberg@vtt.fi)

9

10 **Abstract**

11 The microbial diversity in oligotrophic isolated crystalline Fennoscandian Shield bedrock
12 fracture groundwaters is great but the core community has not been identified. Here we
13 characterized the bacterial and archaeal communities in 12 water conductive fractures situated
14 at depths between 296 m and 798 m by high throughput amplicon sequencing using the
15 Illumina HiSeq platform. **Between 1.7×10^4 – 1.2×10^6 bacterial or archaeal sequence reads**
16 **per sample was obtained. These sequences revealed that up to 95% and 99% of the bacterial**
17 **and archaeal sequences obtained, respectively, belonged to only a few common species, i.e.**
18 **the core microbiome.** However, the remaining rare microbiome contained over 3 and 6 fold
19 more bacterial and archaeal taxa. The metabolic properties of the microbial communities were
20 predicted using PICRUSt. The approximate estimation showed that the metabolic pathways
21 included commonly fermentation, fatty acid oxidation, glycolysis/gluconeogenesis, oxidative
22 phosphorylation and methanogenesis/anaerobic methane oxidation, but carbon fixation
23 through the Calvin cycle, reductive TCA cycle and the Wood-Ljungdahl pathway was also
24 predicted. The rare microbiome is an unlimited source of genomic functionality in all
25 ecosystems. It may consist of remnants of microbial communities prevailing in earlier
26 environmental conditions, but could also be induced again if changes in their living conditions
27 occur.

1 **1 Introduction**

2 Identifying and understanding the core microbiome of any given environments is of crucial
3 importance for predicting and assessing environmental change both locally and globally
4 (Shade and Handelsman, 2012). In a previous study (Bomberg et al., 2015) we showed by 454
5 amplicon sequencing that the active microbial communities in Olkiluoto deep subsurface
6 were strictly stratified according to aquifer water type. Nevertheless, more rigorous
7 sequencing efforts and more samplings have shown that an archaeal core community
8 consisting of the DeepSea Hydrothermal Vent Euryarchaeotal Group 6 (DHVEG-6), ANME-
9 2D and Terrestrial Miscellaneous Group (TMEG) archaea may exist in the anaerobic deep
10 groundwater of Olkiluoto (Miettinen et al., 2015). The bacterial core groups in Olkiluoto deep
11 groundwater include at least members of the Pseudomonadaceae, Comamonadaceae and
12 Sphingomonadaceae (Bomberg et al., 2014; 2015; Miettinen et al., 2015). The relative
13 abundance of these main groups varies at different depths from close to the detection limit to
14 over 90% of the bacterial or archaeal community (Bomberg et al., 2015; Miettinen et al.,
15 2015). **However, both the archaeal and the bacterial communities contain a wide variety of**
16 **less abundant groups, which are distributed unevenly in the different water conductive**
17 **fractures.**

18 The rare biosphere is a concept describing the hidden biodiversity of an environment (Sogin
19 et al., 2006). The rare biosphere consists of microbial groups that are ubiquitously distributed
20 in nature but often present at low relative abundance and may thus stay below the limit of
21 detection. Due to modern high throughput sequencing techniques, however, the hidden
22 diversity of rare microbiota has been revealed. These microorganisms are the basis for
23 unlimited microbial functions in the environment and upon environmental change specific
24 groups can readily activate and become abundant. Access to otherwise inaccessible nutrients
25 activate specific subpopulations in the bacterial communities within hours of exposure (Rajala
26 et al., 2015) and enrich distinct microbial taxa at the expense of the original microbial
27 community in the groundwater (Kutvonen, 2015). Mixing of different groundwater layers due
28 to e.g. breakage of aquifer boundaries and new connection of separated aquifers may cause
29 the microbial community to change and activate otherwise dormant processes. This has
30 previously been shown by Pedersen et al. (2013), who indicated increased sulphate reduction
31 activity when sulphate-rich and methane-rich groundwater mixed. The stability of deep
32 subsurface microbial communities in isolated deep subsurface groundwater fractures are

1 assumed to be stable. However, there are indications that they may change over the span of
2 several years as slow flow along fractures is possible (Miettinen et al., 2015; Sohlberg et al.,
3 2015).

4 The microbial taxa present in an environment interact with both biotic and abiotic factors. In
5 deep subsurface groundwater the biomass concentration is often low and the sampling efforts
6 may not yield enough biomass for extensive metagenomic analysis of the microbial
7 communities. Tools for predicting metabolic pathways may help to establish a consensus of
8 the microbial metabolic characteristics present in an environment and the possible interactions
9 of the microbial communities with the abiotic environment. Tools, such as PICRUSt (Langille
10 *et al.*, 2013), allows us to estimate microbial metabolic functions based on NGS microbiome
11 data. For example, Tsitko et al. (2014) showed that oxidative phosphorylation was the most
12 important energy producing metabolic pathway throughout the 7 m depth profile of an
13 *Acidobacteria*-dominated nutrient poor boreal bog. Cleary et al. (2015) showed that tropical
14 mussel-associated bacterial communities could be important sources of bioactive compounds
15 for biotechnology. This approach is nevertheless hampered by the fact that only little is so far
16 known about uncultured environmental microorganisms and their functions and the PICRUSt
17 approach is best applied for human microbiome for which it was initially developed (Langille
18 *et al.*, 2013). However, metagenomic estimations may give important indications of novel
19 metabolic possibilities even in environmental microbiome studies.

20 Using extensive high throughput amplicon sequencing in this study we aimed to identify the
21 core microbiome in the deep crystalline bedrock fractures of Olkiluoto Island and also to
22 study the rare microbiome. In addition, we aimed to estimate the prevailing metabolic
23 activities that may occur in the deep crystalline bedrock environment of Olkiloto, Finland.

24 **2 Materials and methods**

25 **2.1 Background**

26 The Olkiluoto site has previously been extensively described (Posiva, 2013) and is only
27 briefly described here. The Island of Olkiluoto situating on the west coast of Finland has
28 approximately 60 drillholes drilled for research and monitoring purposes. Studies on the
29 chemistry and microbiology of the groundwater have been on-going since the 1980s. The
30 groundwater is stratified with a salinity gradient extending from fresh to brackish water to a
31 depth of 30 m and the highest salinity concentration of 125 g L⁻¹ total dissolved solids (TDS)

1 at 1000 m depth (Posiva, 2013). The most abundant salinity causing cations are Na^{2+} and Ca^{2+}
2 and anions Cl^- . Between 100 and 300 m depths, the groundwater originates from ancient (pre-
3 Baltic) seawater and has high concentrations of SO_4^{2-} . Below 300 m the concentration of
4 methane in the groundwater increases and SO_4^{2-} is almost absent. A sulphate-methane
5 transition zone (SMTZ), where sulphate-rich fluid replaces methane-rich fluid, is located at
6 250 – 350 m depth. Temperature rises linearly with depth, from ca. 5 – 6 °C at 50 m to ca. 20
7 °C at 1,000 m depth (Ahokas et al., 2008). The pH of the groundwater is slightly alkaline
8 throughout the depth profile. Multiple drillholes intersect several groundwater-filled bedrock
9 fractures, including larger hydrogeological zones such as HZ20 or HZ21 (Table 1). The
10 bedrock of Olkiluoto consists mainly of micagneiss and pegmatitic granite type rocks (Kärki
11 & Paulamäki, 2006). **The *in situ* temperature at 300 m depth in the Olkiluoto bedrock is stable**
12 **at approximately 10°C and increases linearly to approximately 16°C at 800 m depth (Sedighi**
13 **et al., 2013).**

14 This study focused on 12 groundwater samples from water conductive fractures situated at
15 between 296 m and 798 m below sea level bsl and originating from 11 different drillholes in
16 Olkiluoto (Figure 1). The samples represented brackish sulphate waters and saline waters (as
17 classified in Posiva, 2013). The samples were collected between December 2009 and January
18 2013 (Table 1). The physicochemical parameters of the groundwater samples have been
19 reported by reported by Miettinen et al. (2015), but have for clarity been collected here (Table
20 1).

21 **2.2 Sample collection**

22 The collection of samples occurred between December 2009 and January 2013 (Table 1) as
23 described previously (Bomberg et al., 2015; Miettinen et al., 2015; Sohlberg et al., 2015). The
24 samples were obtained from 11 different permanently packered or open drillholes equipped
25 with removable inflatable packers. The position and direction of the drillholes are indicated in
26 Figure 1. Shortly, in order to obtain indigenous fracture fluids, the packer-isolated fracture
27 zones were purged by removing stagnant drillhole water by pumping for a minimum of four
28 weeks before the sample water was collected. The water samples were collected directly from
29 the drillhole into an anaerobic glove box (MBRAUN, Germany) via a sterile, gas-tight poly
30 acetate tube (8 mm outer diameter). Microbial biomass DNA extraction was concentrated
31 from 1000 mL samples by filtration on cellulose acetate filters (0.2 µm pore size, Corning) by
32 vacuum suction inside the glove box. The filters were immediately extracted from the

1 filtration funnels and frozen on dry ice in sterile 50 ml cone tubes (Corning). The frozen
2 samples were transported on dry ice to the laboratory where they were stored at -80°C until
3 use.

4 **2.3 Nucleic acid isolation**

5 Community DNA was isolated directly from the frozen cellulose-acetate filters with the
6 PowerSoil DNA extraction kit (MoBio Laboratories, Inc., Solana Beach, CA), as previously
7 described (Bomberg et al., 2015). **Negative DNA isolation controls were included in the**
8 **isolation protocol.** The DNA concentration of each sample was determined using the
9 NanoDrop 1000 spectrophotometer.

10 **2.4 Estimation of microbial community size**

11 The size of the microbial community was determined by epifluorescence microscopy of 4',6
12 diamidino-2-phenylindole dihydrochloride (DAPI) (Sigma, MO, USA) stained cells as
13 described in Purkamo et al. (2013). The size of the bacterial population was determined by
14 16S rRNA gene targeted quantitative PCR (qPCR) as described by Tsitko et al. (2014) using
15 universal bacterial 16S rRNA gene-targeting primers fD1 (Weisburg et al., 1991) and P2
16 (Muyzer et al., 1993), which specifically target the V1- V3 region of the bacterial 16S rDNA
17 gene. The size of the archaeal population in the groundwater was determined by using primers
18 ARC344f (Bano *et al.*, 2004) and Ar744r (reverse compliment from Barns *et al.*, 1994)
19 flanking the V4-V6 region of the archaeal 16S rRNA gene.

20 The qPCR reactions were performed in 10µL reaction volumes using the KAPA 2 × Syrb®
21 FAST qPCR-kit on a LightCycler480 qPCR machine (Roche Applied Science, Germany) on
22 white 96-well plates (Roche Applied Science, Germany) sealed with transparent adhesive
23 seals (4titude, UK). Each reaction contained 2.5 µM of relevant forward and reverse primer
24 and 1 µL DNA extract. Each reaction was run in triplicate and no-template control reactions
25 were used to determine background fluorescence in the reactions.

26 The qPCR conditions consisted of an initial denaturation at 95 °C for 10 minutes followed by
27 45 amplification cycles of 15 seconds at 95 °C, 30 seconds at 55 °C and 30 seconds at 72 °C
28 with a quantification measurement at the end of each elongation. A final extension step of
29 three minutes at 72 °C was performed prior to a melting curve analysis. This consisted of a
30 denaturation step for 10 seconds at 95 °C followed by an annealing step at 65 °C for one

1 minute prior to a gradual temperature rise to 95 °C at a rate of 0.11 °C s⁻¹ during which the
2 fluorescence was continuously measured. The number of bacterial 16S rRNA genes was
3 determined by comparing the amplification result (Cp) to that of a ten-fold dilution series
4 (10¹-10⁷ copies μL⁻¹) of *Escherichia coli* (ATCC 31608) 16S rRNA genes in plasmid for
5 bacteria and a dilution series of genomic DNA of *Halobacterium salinarum* (DSM 3754) for
6 archaea. The lowest detectable standard concentration for the qPCRs was 10² gene
7 copies/reaction. Inhibition of the qPCR by template tested by adding 2.17×10⁴ plasmid copies
8 containing fragment of the morphine-specific Fab gene from *Mus musculus* gene to reactions
9 containing template DNA as described in Nyysönen *et al.* (2012). Inhibition of the qPCR
10 assay by the template DNA was found to be low. The average Crossing point (Cp) value for
11 the standard sample (2.17×10⁴ copies) was 28.7 (± 0.4 sd), while for the DNA samples Cp
12 was 28.65 - 28.91 (± 0.03-0.28 sd). Nucleic acid extraction and reagent controls were run in
13 all qPCRs in parallel with the samples. Amplification in these controls was never higher than
14 the background obtained from the no template controls.

15 **2.5 Amplicon library preparation**

16 This study is part of the Census of Deep Life initiative, which strives to obtain a census of the
17 microbial diversity in deep subsurface environment by collecting samples around the world
18 and sequencing the 16S rRNA gene pools of both archaea and bacteria. The extracted DNA
19 samples were sent to the Marine Biological Laboratory in Woods Hole, MA, USA, for
20 preparation for HiSeq sequencing using the Illumina technology. The protocol for amplicon
21 library preparation for both archaeal and bacterial 16S amplicon libraries can be found at
22 <http://vamps.mbl.edu/resources/faq.php>. Shortly, amplicon libraries for completely
23 overlapping paired-end sequencing of the V6 region of both the archaeal and bacterial 16S
24 rRNA genes were produced as previously described (Eren *et al.*, 2013). For the archaea,
25 primers A958F and A1048R containing Truseq adapter sequences at their 5' end were used,
26 and for the bacteria primers B967F and B1064R for obtaining 100 nt long paired end reads
27 (<https://vamps.mbl.edu/resources/primers.php>). The sequencing was performed using a HiSeq
28 1000 system (Illumina).

29 **2.6 Sequence processing and analysis**

30 Contigs of the paired end fastq files were first assembled with mothur v 1.32.1 (Schloss *et al.*,
31 2009). Analyzes were subsequently continued using QIIME v. 1.8. (Caporaso *et al.*, 2010).

1 Only sequences with a minimum length of 50 bp were included in the analyses. The bacterial
2 and archaeal 16S rRNA sequences were grouped into OTUs (97% sequence similarity) using
3 both the open reference and closed reference OTU picking strategy and classified using the
4 GreenGenes 13_8 16S reference database (DeSantis et al., 2006). The core archaea and
5 bacteria communities were identified from the OTU tables with the
6 `compute_core_microbiome.py` function in QIIME using default values, with exception of the
7 minimum number of samples where an OTU must be detected, which was set to 80%. The
8 sequencing coverage was evaluated by rarefaction analysis and the estimated species richness
9 and diversity indices were calculated. For comparable α - and β -diversity analyses the data sets
10 were normalized by random subsampling of 17,000 sequences/sample for archaea and
11 140,000 sequences/sample for bacteria. Microbial metabolic pathways were estimated based
12 on the 16S rRNA gene data from the closed OTU picking method using the PICRUSt
13 software (Langille *et al.*, 2013) on the web based Galaxy application (Goecks et al., 2010;
14 Blankenberg et al., 2010; Giardine et al., 2005). The predicted KO numbers were plotted on
15 KEGG pathway maps (<http://www.genome.jp/kegg/>) separately for the bacterial and archaeal
16 predicted metagenomes, with a threshold of a minimum of 100 genes in total estimated from
17 all samples. The sequence data has been submitted to the Sequence Read Archive (SRA,
18 <http://www.ncbi.nlm.nih.gov/sra>) under study SRP053854, Bioproject PRJNA275225.

19 **2.7 Statistical analyses and data visualization**

20 The similarity of the archaeal and bacterial communities between the different samples was
21 tested by principal coordinates analysis (PCoA) using the Phyloseq package in R (McMurdie
22 and Holmes, 2014; R Core Team, 2013). The analysis was performed using the raw OTU
23 tables outputted by QIIME. In addition, a PCoA analysis showing the effect of library size on
24 the ordination of the samples was calculated using `vegan` (Oksanen et al., 2016). The Bray-
25 Curtis distance model was used for both analyses. The samples were hierarchically clustered
26 in a UPGMA tree based on the raw OTU counts using the `heatmap` function of `phyloseq` in R.

27

1 **3 Results**

2 **3.1 Microbial community size**

3 The total number of microbial cells detected by epifluorescence microscopy of DAPI stained
4 cells was between 2.3×10^4 and 4.2×10^5 cells mL⁻¹ groundwater (Figure 2, Table 1). The
5 concentration of bacterial 16S rRNA gene copies mL⁻¹ varied between 9.5×10^3 and 7.0×10^5
6 and that of the archaea 2.6×10^1 and 6.3×10^4 (Figure 2, Table 1).

7 **3.2 Sequence statistics, diversity estimates and sequencing coverage**

8 The number of bacterial v6 sequence reads from the 12 samples varied between $1.4 - 7.8 \times$
9 10^5 reads, with a mean sequencing depth of 2.9×10^5 ($\pm 1.8 \times 10^5$ standard deviation)
10 reads/sample (Table 2). The archaeal v6 sequence reads ranged from $0.17 - 12.1 \times 10^5$ reads
11 with a mean sequencing depth of 4.1×10^5 ($\pm 3.5 \times 10^5$ standard deviation) reads/sample. The
12 numbers of observed operational taxonomic units (OTUs) represented on average 82.6% (\pm
13 12.5%) of the Chao1- and 78.1 % ($\pm 13.4\%$) of the ACE-estimated numbers of bacterial
14 OTUs (Table 2ab). The archaeal communities were slightly better covered, with on average
15 88.5% ($\pm 11.5\%$) of the Chao1 and 84.8% ($\pm 12.6\%$) of the ACE estimated number of OTUs
16 detected. Shannon diversity index H', calculated from 140,000 and 17,000 random sequence
17 reads per sample for the bacteria and archaea, respectively, was high for both bacterial and
18 archaeal communities. High H' values and climbing rarefaction curves (Figure S1) indicated
19 high diversity in the microbial communities in the different deep groundwater fracture zones
20 of Olkiluoto. The bacterial H' was on average 13 (± 0.74), ranging from 11 to 14 between the
21 different samples. The archaeal H' was on average 11 (± 1.2) ranging from 9 to 12 between
22 the samples. **A total of 468,684 archaeal and 301,458 bacterial OTUs were obtained in this**
23 **study.**

24 **3.3 Microbial communities**

25 From the bacterial v6 sequences 49 different bacterial Phyla were detected (Appendix 1).
26 These phyla included 165 bacterial classes, 230 orders, 391 families and 651 genera. The
27 greatest number of sequences, between 21.83% and 47.94% per sample, clustered into an
28 undetermined bacterial group (Bacteria, Other), which may be due the fact that sequences of
29 poorer quality may be difficult to classify, especially as the sequences are short.

1 The archaea were represented by two identified phyla, the Euryarchaeota and the
2 Crenarchaeota (Appendix 2). These included 21 classes, 38 orders, 61 families and 81 genera.
3 Between 4.7% and 35.0% of the archaeal sequences of each sample were classified to
4 unassigned Archaea, with a general increase in unassigned archaeal sequences with increasing
5 depth.

6 The archaeal and bacterial core communities were determined as OTUs present in at least
7 80% of the samples. Of the more than 4.6×10^5 archaeal OTUs the core community consisted
8 of 82 OTUs belonging to three archaeal orders, the E2 of the Thermoplasmatales, the
9 Methanobacteriales and the Methanosarcinales (Figure 3). Additionally, a great proportion of
10 the OTUs of the core community did not receive any other taxonomic identity than Archaea.
11 The most common archaeal family of the core community was the ANME-2D belonging to
12 the Methanosarcinales. The bacterial core community consisted of only 26 OTUs, compared
13 to more than 3.0×10^5 bacterial OTUs in total (Figure 3). These OTUs belonged to six
14 different families, the Alteromonadales, Burkholderiales, SB-45, Sphingomonadales,
15 Syntrophobacterales and Thiobacterales. In addition, a great portion of the core community
16 OTUs were classified only as unassigned bacteria. The most abundant of the bacterial core
17 community OTUs belonged to Thiobacteriaceae and Comamonadaceae. In both the archaeal
18 and bacterial sequence data a great proportion of the sequence reads were only identified as
19 archaea or bacteria, without more detailed taxonomic assignments. The core OTUs were
20 distributed with different abundance in the different samples (Figure 3). Most of the OTUs
21 detected were present in less than 20% of the samples (Figure 4).

22 **3.4 Environmental parameters and the microbial communities**

23 The microbial community profiles of the different samples were clustered in a UPGMA tree
24 based on the OTU tables and the Bray-Curtis distance model (Figure 5). The archaeal and
25 bacterial communities clustered according to the OTUs detected in the samples, but not
26 clearly according to any physicochemical parameter. In the PCoA plots, the samples clustered
27 into three groups (Figures 6a-d). However, no clear environmental factor was identified to
28 drive the communities. In these analyses the microbial communities, with 4.6×10^5 archaeal
29 and 3.0×10^5 bacterial OTUs, were the strongest clustering forces and thus even quite
30 different communities obtained similar coordinates on the PCoA plots. Coordinate 1
31 determined 14.9% and Coordinate 2 12.6% of the variance in the archaeal communities and
32 20% and 17.2%, respectively, of the variance in the bacterial communities (Figure 6e,f).

1 3.5 Predicted metabolic functions of the deep subsurface microbial communities

2 The putative metabolic functions of the microbial communities at different depth was
3 predicted using the PICRUSt software, which compares the identified 16S rRNA gene
4 sequences to those of known genome sequenced species thereby estimating the possible gene
5 contents of the uncultured microbial communities. The analysis is only an approximation, but
6 may give an idea of the possible metabolic activities in the deep biosphere. In order to
7 evaluate the soundness of the analysis a nearest sequenced taxon index (NSTI) for each of the
8 bacterial and archaeal communities was calculated by PICRUSt. An NSTI value of 0 indicates
9 high similarity to the closest sequenced taxon while NSTI=1 indicates no similarity. The
10 NSTI of the bacterial communities at different depths varied between 0.045 in sample OL-
11 KR44 and 0.168 in sample OL-KR13 (Figure 7). The NSTI for archaea were much higher
12 ranging from 0.141 in sample OL-KR9 at depth of 432 m and 0.288 in OL-KR44. This
13 indicates that the metagenomic estimates are only indicative. The estimated microbial
14 metabolism did not differ noticeably between the different depths (Figure 8a and b). The most
15 important predicted metabolic pathways included membrane transport in both bacterial and
16 archaeal communities. The most common pathways for carbohydrate metabolism were the
17 butanoate, propionate, glycolysis/gluconeogenesis and pyruvate metabolism pathways for the
18 bacteria and glycolysis/gluconeogenesis and pyruvate metabolism pathways for the archaea
19 (Figure 9). Glucose is converted into pyruvate and further to Acetyl-CoA by both bacteria and
20 archaea. The bacterial community may produce and utilize acetate. Both the bacterial and
21 archaeal communities fix carbon via the Wood-Ljungdal (WL) reverse Citric acid cycle
22 (rTCA) and Calvin pathways. Methane is produced from methylamines, CO₂ and methanol by
23 the methanogenic archaea. **In addition to the strong evidence of methanogenesis in the**
24 **archaeal community the reverse methanogenesis, i.e. anaerobic methane oxidation by the**
25 **ANME-2D archaea is possible.** Based on the predicted metagenomes the bacterial community
26 is not able to oxidize methane or hydrolyze methanol, but the methylotrophs present may use
27 formic acid and trimethylamines.

28 The most abundant energy metabolic pathway in the bacterial communities was the oxidative
29 phosphorylation (Figure S3) while for the archaea the methane metabolism was the most
30 important (Figure 8, 9). Utilization of propanoate and butanoate (Figure 8) by the bacterial
31 communities as well as well covered fatty acid biosynthesis and degradation pathways
32 indicate that the bacterial community is capable of fermentation (Figure S4a and b). Nitrate is

1 reduced both through dissimilatory nitrate reduction to ammonia and through denitrification
2 to nitrous oxide by the bacteria (Figure S5). In addition, nitrogen is fixed to ammonia by both
3 archaea and bacteria. The ammonia is then used as raw material for L-glutamate synthesis
4 (Figure S5). Sulfur metabolism was not a major pathway in either the bacterial or the archaeal
5 communities according to the predicted number of genes. However, assimilatory sulphate
6 reduction was indicated in both the bacterial and archaeal communities, while dissimilatory
7 sulphate reduction and sulphur oxidation was indicated only in the bacterial communities
8 (Figure S6).

9 Several amino acid synthesis pathways were predicted (Figure 8), of which the most
10 prominent were the alanine, aspartate and glutamate synthesis, arginine and proline synthesis,
11 cysteine and methionine synthesis, glycine, serine and threonine synthesis, phenylalanine,
12 tyrosine and tryptophan synthesis and the valine, leucine and isoleucine synthesis pathways.

13 Different types of membrane transport (ABC transporters) was identified where sulphate and
14 iron (III) were taken up by the bacteria and tungstate, molybdate, proline, zink, cobalt and
15 nickel was taken up by both archaea and bacteria (Figure S7). The estimated number of genes
16 for both the purine and pyrimidine metabolism was more than two times higher in the
17 archaeal community than in the bacterial community (Figure 8a and b).

18

19 **4 Discussion**

20 The phenotypic characteristics of the Fennoscandian Shield deep subsurface microbial
21 communities are still largely unknown although specific reactions to introduced
22 environmental stimulants have been shown (e.g. Pedersen et al., 2013; 2014; Rajala et al.,
23 2015; Kutvonen 2015). Nevertheless, the connection of these microbial responses to specific
24 microbial groups is still only in an early phase. Metagenomic and gene specific analyses of
25 deep subsurface microbial communities have revealed prominent metabolic potential of the
26 microbial communities, which appear to be associated with the prevailing lithology and
27 physicochemical parameters (Nyyssönen et al., 2014; Purkamo et al., 2015). It has also been
28 shown with fingerprinting methods with ever increasing efficiency that the bacterial and
29 archaeal communities are highly diverse in the saline anaerobic Fennoscandian deep fracture
30 zone groundwater (Bomberg et al., 2014; 2015; Nyyssönen et al., 2012; 2014; Pedersen et al.,
31 2014; Miettinen et al, 2015; Sohlberg et al., 2015). Nevertheless, the concentration of
32 microbial cells in the groundwater is quite low (Figure 2, Table 1). **In accordance with other**

1 Fennoscandian deep subsurface environments (Purkamo et al., 2016), most of the microbial
2 communities at different depth in Olkiluoto bedrock fractures consist of bacteria. Archaea
3 have in general been shown to constitute at most approximately 1% of the Fennoscandian
4 deep bedrock groundwater (Purkamo et al., 2016). However, at specific depths in Olkiluoto
5 (328 m, 423 m) the archaea contributed with over 50% of the estimated 16S rRNA gene pool
6 (Table 1). The major archaeal group present at these depths were the ANME-2D archaea
7 indicating that nitrate-mediated anaerobic oxidation of methane may be especially common
8 (Haroon et al., 2013). The high abundance of archaea in Olkiluoto is special for this
9 environment. Archaea have also been quantified from the Outokumpu deep scientific
10 borehole (Purkamo et al., 2016), but unlike the situation in Olkiluoto the archaeal community
11 was less than 1% of the total community at best.

12 Previously, using 454 amplicon sequencing, we have observed OTU numbers of
13 approximately 800 OTUs per sample covering approximately 550 bacterial genera (or
14 equivalent groups) and approximately 350 archaeal OTUs including approximately 80
15 different genera (or equivalent groups) (Miettinen et al., 2015). Miettinen et al. (2015) defined
16 the OTUs 97% sequence homology and the number of sequence reads per sample was at most
17 in the range of 10^4 . In contrast, our sequence read numbers were 10- to 100-fold higher and
18 the number of OTUs per sample in general 100-fold higher. This indicates that a greater
19 sequencing depth increases the number of taxa detected from the subsurface environment and
20 allows us a novel view of the so far hidden rare biosphere. Nevertheless, in comparison to the
21 high number of OTUs detected the number of identified genera, 651 and 81 bacterial and
22 archaeal genera, respectively, seems low. On the other hand, this indicates that the sequencing
23 depth has been sufficient to detect most of the prokaryotic groups present. Nevertheless, the
24 obtained numbers of OTUs per sample in this study were huge (Table 2). This may reflect the
25 high level of variability in the short sequence reads of the v6 region used in this study. As
26 discussed by Huse et al. (2008), short sequence reads very often match several different full-
27 length 16S rRNA reads. As shown in our study taxonomic assignments, such as
28 'Proteobacteria_other' were common and may be due to multiple matches for the individual
29 sequence reads obtained in the identification step of the analysis.

30 In general, the microbial communities at different depth grouped loosely into clusters (Figure
31 6). Although no clear environmental factor seemed to drive the microbial communities at
32 different depths, the core communities appeared to be more similar in samples from similar

1 depths, especially in the bacterial communities (Figure 3). OTUs belonging to both sulphate
2 reducers (*Desulfobacteraceae*) and sulphur oxidizers (*Thiobacteraceae*) were present in the
3 bacterial core community. The archaeal core community consisted mostly of methane-
4 oxidizing ANME-2D archaea. Interestingly, however, their abundance was higher in the
5 deeper samples. Previous studies on the Finnish deep biosphere has shown that the microbial
6 communities at different sites vary strongly from each other. Purkamo et al. (2015)
7 investigated the bacterial and archaeal communities of different fracture zones of the
8 Outokumpu deep scientific borehole and found that the majority of the bacterial populations
9 at depths between 180 m and 500 m depth consist of Betaproteobacteria belonging to the
10 *Commamonadaceae* and the archaeal communities consist of *Methanobacteriaceae* and
11 *Methanoregula*.

12 The core communities, defined as OTUs present in all the studied samples, accounted for
13 between 0.2% and 11.7% of the archaeal and 0.4% - 4.1% of the bacterial communities,
14 respectively. This is surprisingly low and indicates an obviously high microbial diversity in
15 these deep fracture water samples. Nevertheless, the short read length and high sequence
16 variability within the v6 region may affect these results. Nevertheless, genus-level
17 identification of the sequence data showed that between 95 – >99% of the archaeal sequence
18 reads fell in to 25 genera, which were present in all samples. Likewise for the bacterial
19 communities, 80 – 97% of the sequene reads belonged to 95 bacterial genera that were
20 detected in all samples. The number of OTUs that were not present in more than 80% of the
21 samples far outnumbered the number of core OTUs both in the archaeal and the bacterial
22 communities. Only approximately 800 archaeal OTUs and less than 600 bacterial OTUs were
23 present in at least 50% of the samples. Interestingly, 20% of the samples share only
24 approximately 10,000 archaeal and 30,000 bacterial OTUs, indicating that the proportion of
25 rare OTUs present in only 1 or 2 samples is huge.

26 Our results agree with Sogin et al. (2006) and Magnabosco et al. (2014), who showed that a
27 relatively small number of taxa dominate deep-sea water and deep groundwater habitats,
28 respectively, but a rare microbiome consisting of thousands of taxonomically distinct
29 microbial groups are detected at low abundances. What this means for the functioning of the
30 deep subsurface is that the microbial communities have the capacity to respond and change
31 due to changes in environmental conditions. For example, Pedersen et al. (2014) showed that
32 by adding sulphate to the sulphate-poor but methane-rich groundwater in Olkiluoto the

1 bacterial population changed over the span of 103 days from a non-SRB community to a
2 community dominated by SRB. In addition, a change in the geochemical environment
3 induced by H₂ and methane impacted the size, composition and functions of the microbial
4 community and ultimately led to acetate formation (Pedersen et al., 2012; Pedersen, 2013;
5 Pedersen et al., 2014).

6 The metabolic pathways predicted by PICRUSt are far from certain when uncultured and
7 unculturable deep subsurface microbial communities are concerned. The NSTI values for both
8 the bacterial and well as the archaeal communities were high indicating that closely related
9 species to those found in our deep groundwater have yet to be sequenced. This is in
10 accordance with Langille et al. (2013), who showed that environments containing a high
11 degree of unexplored microbiota also tend to have high NSTI values. Staley et al. (2014) also
12 showed in a comparison between PICRUSt and shot gun metagenomic sequencing of riverine
13 microbial communities that PICRUSt may not be able to correctly assess rare biosphere
14 functions. Nevertheless, Langille et al. (2013) showed that PICRUSt may predict the
15 metagenomic content of a microbial community more reliably than shallow metagenomic
16 sequencing. **Although PICRUSt does not give as detailed results as metagenomics or genomic
17 analyses may give, it is still a useful tool for predicting functions in microbial communities
18 when the possibility for metagenomics analysis may be impossible, e.g. due to low biomass in
19 the samples.**

20 **Energy metabolism.** Deep subsurface environments are often declared energy deprived
21 environments dominated by autotrophic microorganisms (Hoehler and Jorgensen, 2013).
22 However, recent reports indicate that heterotrophic microorganisms play a greater role than
23 the autotrophic microorganisms in Fennoscandian deep crystalline subsurface environments
24 (Purkamo et al., 2015). Heterotrophic communities with rich fatty acid assimilation strategies
25 have been reported to fix carbon dioxide on the side of e.g. fermenting activities in order to
26 replenish the intracellular carbon pool, which otherwise would be depleted. Wu et al. (2015)
27 also found by metagenomic analyses that fermentation was a major metabolic activity in the
28 microbial community of Swedish deep groundwater. Our results agree with Purkamo et al.
29 (2015) that a greater proportion of the microbial community is involved in carbohydrate and
30 fatty and organic acid oxidation than in fixation of inorganic carbon. Nevertheless,
31 autotrophic carbon fixation pathways were predicted in the analysis with PICRUSt, indicating
32 that both the archaeal and bacterial communities include autotrophic members, although these

1 microorganisms might not be obligate autotrophs. It was also noted that even though evidence
2 for methane oxidation could not be inferred from the PICRUSt predictions (no *pmoA* genes),
3 the bacterial community may oxidize formate, which is in agreement with the findings
4 reported by Wu et al. (2015).

5 Several carbon fixation pathways were predicted in the metagenomes, the Calvin cycle,
6 reductive TCA (rTCA) cycle and Wood-Ljungdahl (WL) pathway. The WL-pathway is
7 considered the most ancient autotrophic carbon fixation pathway in bacteria and archaea
8 (Fuchs 1989, Martin et al. 2008, Berg et al. 2010; Hügler and Sievert, 2011) and was found in
9 both the bacterial and the archaeal communities. In the archaeal community the Calvin cycle
10 and the rTCA were especially pronounced in the samples from 296 m, 405 – 423 m and
11 somewhat lower at 510 – 527 m depth. The bacterial communities are predicted to fix CO₂ at
12 almost all depths with the exception of 405 m and 559 m depth. **Nevertheless, our results**
13 **agree with Nyssönen et al. (2014), who showed my metagenomic analysis that the microbial**
14 **communities at different depth of the Outokumpu scientific deep drill hole may fix carbon in**
15 **several ways, of which the rTCA, the WL pathway and the Calvin cycle were identified.**
16 **Magnabosco et al. (2016) showed that the WL pathway was the dominating form of carbon**
17 **fixation in metagenomes of 3 km deep Precambrian crust biospheres in South Africa. Dong et**
18 **al. (2014) also suggested that microorganisms in low-energy deep subsurface environment**
19 **may have several strategies for e.g. carbon fixation, as shown in the *Halomonas sulfidaeris*, in**
20 **order to access as many resources as possible.** The predicted methane metabolism (methane
21 and methyl compound consumption) and oxidative phosphorylation were equally strong in the
22 bacterial community. Sulphur metabolism was not **predicted to be** a common pathway for
23 energy in either the archaeal or the bacterial **communities. However, PICRUSt predicted**
24 **bacteria** with either assimilative or dissimilative sulphate reduction **to be present. Sulphur**
25 **oxidation through the *sox* system was in general not predicted, but the *soxD* gene was**
26 **predicted and oxidation of thiosulphate to sulphate may be possible (Figure S6).** Nitrate is
27 reduced both through dissimilatory nitrate reduction to ammonia and through denitrification
28 to nitrous oxide by the bacteria. In addition, nitrogen is fixed to ammonia by both archaea and
29 bacteria. The ammonia is then used as raw material for L-glutamate synthesis.

30 Oxidative phosphorylation was one of the most prominent energy generating metabolic
31 pathways in the bacterial community. This indicates that ATP is generated by electron
32 transfer to a terminal electron acceptor, such as oxygen, nitrate or sulphate. In the archaeal

1 community the oxidative phosphorylation was not as strongly indicated, but this may be due
2 to missing data on archaeal metabolism in the KEGG database.

3 The main energy metabolism of the archaeal communities appeared to be the methanogenesis,
4 especially at 296 m and 405 m. Methanogenesis was common also at all other depths except
5 330 m – 347m, 415 m and 693 m – 798 m. Methane is produced from CO₂-H₂ and methanol,
6 and from acetate, although evidence for the acetate kinase enzyme was lacking.
7 Methanogenesis from methylamines may also be possible, especially at 296 m and 405 m.
8 Methane oxidation using methane monooxygenases and methanol dehydrogenases does not
9 occur in either bacterial or archaeal communities. **It should be noted, however, that the**
10 **ANME-2D archaea are likely to use the methanogenesis pathway in the reverse for oxidizing**
11 **methane anaerobically to carbon dioxide (Haroon et al., 2013). The produced carbon dioxide**
12 **may be fixed by the same archaea and turned in to acetate, which may serve as carbon**
13 **substrate and electron donor and acceptor for a large variety of microorganisms in the**
14 **groundwater.**

15 **Carbohydrate metabolism.** Glycolysis/gluconeogenesis is one of the most common
16 carbohydrate-metabolizing pathways predicted for both the archaeal and bacterial
17 communities (Figure 9). Pyruvate from glycolysis is oxidized to acetyl-CoA by both archaea
18 and bacteria and used in the TCA cycle. The TCA cycle provides for example raw material
19 for many amino acids, such as lysine and glutamate. The butanoate and propanoate
20 metabolisms were also common in the bacterial communities, indicating fermentative
21 metabolism and capability of fatty acid oxidation.

22 **Amino acid metabolism.** Non-essential amino acids, such as alanine, aspartate and glutamate
23 are produced from ammonia and pyruvate or oxaloacetate especially in the archaeal
24 populations. In the archaeal population proline appears to be produced from glutamate.
25 Despite the low use of sulphate as energy source in the microbial communities sulphate and
26 other sulphur compounds are taken up for the production of the amino acids cysteine and
27 methionine by both the archaeal and the bacterial communities. A higher predicted relative
28 abundance of genes involved in aromatic amino acid synthesis (phenylalanine, tyrosine,
29 tryptophane) was seen in the archaeal than in the bacterial communities. Both the archaeal and
30 the bacterial communities synthesise branched chained amino acids (isoleucine, leucine and
31 valine), but only the bacteria degrade them. Especially proteobacteria have been shown to be
32 able to use the branched chained amino acids (isoleucine, leucine and valine) and short

1 chained fatty acids (acetate, butyrate, propionate) as sole energy and carbon source (Kazakov
2 et al., 2009). The branched chained amino acids function as raw material in the biosynthesis
3 of branched chained fatty acids, which regulate the membrane fluidity of the bacterial cell. In
4 salt stress conditions, the proportion of branch-chained fatty acids in the membranes
5 decreases.

6 **Membrane transport.** According to the predicted metagenomes, the microbial cells transport
7 sulphate into the cell, but do not take up nitrate. Nitrogen is taken up as glutamate but not as
8 urea. Iron is taken up by an Fe(III) transport system and an iron complex transport system in
9 the bacterial communities, but generally only by the iron complex transport system in
10 archaea. However, Fe(III) transport system may also exist in the archaeal communities at 405
11 m to 423 m depth, where also some manganese/iron transport systems could be found.
12 Molybdate and phosphate is transported into the cell by molybdate and phosphate ATPases,
13 respectively. Nickel is taken up mainly by a nickel/peptide transport system but also to some
14 extent by a cobalt/nickel transport system. Zink is taken up to some extent by a zink transport
15 system, but transport systems for manganese, manganese/iron, manganese/zink/iron, or
16 iron/zink/copper are negligent. Ammonia is taken up by an Amt transport system.

17

18 **5 Conclusions**

19 The wide diversity of microbial groups in the deep Fennoscandian groundwater at the
20 Olkiluoto site revealed that the majority of the microbial community present belong to only a
21 few microbial taxa while the greatest part of the microbial diversity is represented by low
22 abundance and rare microbiome taxa. The core community was present in all tested samples
23 from different depths, but the relative abundance of the different taxa varied in the different
24 samples. **Nevertheless, the proportion of OTUs found in only a small propostion (e.g. 20%) of**
25 **the samples far surpassed the number of OTUs included in the core communities.**
26 Fermentation or oxidation of fatty acids was a common carbon cycling and energy harvesting
27 metabolic pathways in the bacterial communities whereas the archaea may either produce or
28 consume methane. Glycolysis/gluconeogenesis was predicted to be common in both the
29 archaeal and bacterial communities. In addition both the bacterial and archaeal communities
30 were estimated to contain several different common carbon fixation pathways, such as the
31 Calvin cycle and the reductive TCA and the Wood-Ljungdahl pathway.

32

1 **Acknowledgements**

2 The Illumina sequencing data were made possible by the Deep Carbon Observatory's Census
3 of Deep Life supported by the Alfred P. Sloan Foundation. Illumina sequencing was
4 performed at the Marine Biological Laboratory (Woods Hole, MA, USA) and we are grateful
5 for the assistance of Mitch Sogin, Susan Huse, Joseph Vineis, Andrew Voorhis, Sharon Grim,
6 and Hilary Morrison at MBL and Rick Colwell, OSU. MB was supported by the Academy of
7 Finland (project 261220).

8

1 **References**

- 2 Bano, N., Ruffin, S., Ransom, B. and Hollibaugh, J.T.: Phylogenetic composition of Arctic
3 ocean archaeal assemblages and comparison with Antarctic assemblages. *Appl. Environ.*
4 *Microbiol.* 70, 781-789, 2004.
- 5 Barns, S.M., Fundyga, R.E., Jeffries, M.W. and Pace, N.R.: Remarkable archaeal diversity
6 detected in Yellowstone National Park hot spring environment. *Proc. Natl. Acad. Sci. USA*
7 91, 1609-1613, 1994.
- 8 Berg, I.A., Kockelkorn, D., Ramos-Vera, W.H., Say, R.F., Zarzycki, J., Hügler, M., Alber,
9 B.E. and Fuchs, G.: Autotrophic carbon fixation in archaea. *Nat. Rev. Microbiol.* 8, 447-
10 460, 2010.
- 11 Blankenberg, D., Von Kuster, G., Coraor, N., Ananda, G., Lazarus, R., Mangan, M.,
12 Nekrutenko, A. and Taylor, J.: Galaxy: a web-based genome analysis tool for
13 experimentalists. *Curr. Protoc. Mol. Biol.*, 19, 19.10.1-21, 2010.
- 14 Bomberg, M., Nyssönen, M., Nousiainen, A., Hultman, J., Paulin, L., Auvinen, P. and
15 Itävaara, M.: Evaluation of molecular techniques in characterization of deep terrestrial
16 biosphere. *Open J Ecol* 4, 468-487, 2014.
- 17 Bomberg, M., Nyssönen, M., Pitkänen, P., Lehtinen, A. and Itävaara, M.: Active microbial
18 communities inhabit sulphate-methane interphase in deep bedrock fracture fluids in
19 Olkiluoto, Finland. *Biomed. Res. Int.*, Article ID 979530, 2015.
20 <http://dx.doi.org/10.1155/2015/979530>.
- 21 Caporaso, J. G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F. D., Costello, E. K.,
22 Fierer, N., Gonzalez Pena, A., Goodrich, J. K., Gordon, J. I., Huttley, G. A., Kelley, S. T.,
23 Knights, D., Koenig, J. E., Ley, R. E., Lozupone, C. A., McDonald, D., Muegge, B. D.,
24 Pirrung, M., Reeder, J., Sevinsky, J. R., Turnbaugh, P. J., Walters, W. A., Widmann, J.,
25 Yatsunencko, T., Zaneveld, J. and Knight, R. QIIME allows analysis of high-throughput
26 community sequencing data. *Nature Methods*, 7, 335-336, 2010.
- 27 Cleary, D. F., Becking, L. E., Polónia, A. R., Freitas, R. M. and Gomes, N. C.: Composition
28 and predicted functional ecology of mussel-associated bacteria in Indonesian marine lakes.
29 *Antonie van Leeuwenhoek*, 107, 821-834, 2015.
- 30 DeSantis, T. Z., Hugenholtz, P., Larsen, N., Rojas, M., Brodie, E. L., Keller, K., Huber, T.,
31 Dalevi, D., Hu, P. and Andersen, G. L. Greengenes, a Chimera-Checked 16S rRNA Gene
32 Database and Workbench Compatible with ARB. *Appl. Environ. Microbiol.*, 72, 5069-72,
33 2006.
- 34 Dong, Y., Kumar, C.G., Chia, N., Kim, P.-J., Miller, P.A., Price, N.D., Cann, I.K.O., Flynn,
35 T.M., Sanford, R.A., Krapac, I.G., Locke, R.A.II, Hong, P.-Y., Tamaki, H., Liu, W.-T.,
36 Mackie, R.I., Hernandez, A.G., Wright, C.L., Mikel, M.A., Walker, J.L., Sivaguru, M.,
37 Fried, G., Yannarell, A.C., Fouke, B.W. Halomonas sul- fidaeris-dominated microbial
38 community inhabits a 1.8 km-deep subsurface Cambrian Sandstone reservoir. *Environ.*
39 *Microbiol.* 16, 1695-1708, 2014.
- 40 Eren, A.M., Maignien, L., Sul, W.J., Murphy, L.G., Grim, S.L., Morrison, H.G. and Sogin,
41 M.L. Oligotyping: differentiating between closely related microbial taxa using 16S rRNA
42 gene data. *Meth. Ecol. Evol.*, 4, 1111-1119, 2013.
- 43 Fuchs, G.: Alternative pathways of autotrophic CO₂ fixation. In: *Autotrophic Bacteria*, ed.

- 1 HG Schlegel, B Bowien, pp. 365–82. Berlin: Springer, 1989.
- 2 Giardine, B., Riemer, C., Hardison, R.C., Burhans, R., Elnitski, L., Shah, P., Zhang, Y.,
3 Blankenberg, D., Albert, I., Taylor, J., Miller, W., Kent, W.J., Nekrutenko, A.: Galaxy: a
4 platform for interactive large-scale genome analysis. *Genome Res.*, 15, 1451-5, 2005.
- 5 Goecks, J., Nekrutenko, A., Taylor, J. and The Galaxy Team: Galaxy: a comprehensive
6 approach for supporting accessible, reproducible, and transparent computational research in
7 the life sciences. *Genome Biol.* 11, R86, 2010.
- 8 Hallbeck, L. and Pedersen, K.: Culture-dependent comparison of microbial diversity in deep
9 granitic groundwater from two sites considered for a Swedish final repository of spent
10 nuclear fuel. *FEMS Microbiol. Ecol.*, 81, 66-77, 2012.
- 11 Hammer, Ø., Harper, D.A.T. and Ryan, P.D.: PAST: Paleontological statistics software
12 package for education and data analysis. *Palaeontologia Electronica* 4: 9pp. [http://palaeo-](http://palaeo-electronica.org/2001_1/past/issue1_01.htm)
13 [electronica.org/2001_1/past/issue1_01.htm](http://palaeo-electronica.org/2001_1/past/issue1_01.htm), 2001.
- 14 Haroon, M.F., Hu, S., Shi, Y., Imelfort, M., Keller, J., Hugenholtz, P., Yuan, Z. and Tyson,
15 G.W.: Anaerobic oxidation of methane coupled to nitrate reduction in a novel archaeal
16 lineage. *Nature*, 500, 567-570, 2013.
- 17 Hoehler, T.M. and Jørgensen, B.B.: Microbial life under extreme energy limitation. *Nat Rev*
18 *Microbiol* 11, 83-94, 2013.
- 19 Hügler, M. and Sievert, S. M.: Beyond the Calvin cycle: autotrophic carbon fixation in the
20 ocean. *Mar. Sci.*, 3, 2011.
- 21 Huse, S.M., Dethlefsen, L., Huber, J.A., Welch, D.M., Relman, D.A. and Sogin, M.L.
22 Exploring microbial diversity and taxonomy using SSU rRNA hypervariable tag sequencing.
23 *PLoS Genet*, 4, p.e1000255, 2008.
- 24 Kärki, A. and Paulamäki, S.: Petrology of Olkiluoto. Posiva 2006-02, 2006.
- 25 Kazakov, A.E., Rodionov, D.A., Alm, E., Arkin, A.P., Dubchak, I. and Gelfand, M.S.:
26 Comparative genomics of regulation of fatty acid and branched-chain amino acid utilization
27 in proteobacteria. *J. Bacteriol.*, 191, 52-64, 2009.
- 28 Kutvonen, H.: Nitrogen-cycling bacteria in groundwater of the low and medium active nuclear
29 waste repository in Olkiluoto, Finland (In Finnish) M.Sc. Thesis, Faculty of Agriculture and
30 Forestry, University of Helsinki, Finland, 2015.
- 31 Langille, M. G. I., Zaneveld, J., Caporaso, J. G., McDonald, D., Knights, D., Reyes, J. A.,
32 Clemente, J. C., Burkepille, D. E., Vega Thurder, R. L., Knight, R., Beiko, R. G. and
33 Huttenhower, C.: Predictive functional profiling of microbial communities using 16S rRNA
34 marker gene sequences. *Nature Biotechnol.*, 31, 814-821, 2013.
- 35 Magnabosco, C., Tekere, M., Lau, M. C., Linage, B., Kuloyo, O., Erasmus, M., Cason, E.,
36 van Heerden, E., Borgonie, G. and Onstott, T. L. Comparisons of the composition and
37 biogeographic distribution of the bacterial communities occupying South African thermal
38 springs with those inhabiting deep subsurface fracture water. *Front. Microbiol*, 5, 10-3389,
39 2014.
- 40 Magnabosco, C., Ryan, K., Lau, M.C., Kuloyo, O., Lollar, B.S., Kieft, T.L., van Heerden, E.
41 and Onstott, T.C. A metagenomic window into carbon metabolism at 3 km depth in
42 Precambrian continental crust. *ISME J.*, 10, 730-741, 2016.

- 1 Martin, W., Baross, J., Kelley, D. and Russell, M.J.: Hydrothermal vents and the origin of
2 life. *Nat. Rev. Microbiol.* 6, 805–14, 2008.
- 3 **McMurdie, P.J.; Holmes, S.: Shiny-phyloseq: Web Application for Interactive Microbiome**
4 **Analysis with Provenance Tracking. *Bioinformatics* 31, 282–283, 2015.**
- 5 Miettinen H., Bomberg, M., Nyysönen, M., Salavirta, H. Sohlberg, E. Vikman, M. and
6 Itävaara, M.: The Diversity of Microbial Communities in Olkiluoto Bedrock Groundwaters
7 2009–2013. Olkiluoto, Finland, Posiva Oy. Posiva Working Report 2015-12, 1-160, 2015.
8 www.posiva.fi
- 9 Muyzer, G., de Waal, E.C., and Uitterlinden, A.G.: Profiling of complex microbial
10 populations by denaturing gradient gel electrophoresis analysis of polymerase chain
11 reaction-amplified genes coding for 16S rRNA. *Appl. Environ. Microbiol.* 59, 695–700,
12 1993.
- 13 Nyysönen, M., Bomberg, M., Kapanen, A., Nousiainen, A., Pitkänen, P. and Itävaara, M.:
14 Methanogenic and sulphate-reducing microbial communities in deep groundwater of
15 crystalline rock fractures in Olkiluoto, Finland. *Geomicrobiol. J.*, 29, 863–878, 2012.
- 16 Nyysönen, M., Hultman, J., Ahonen, L., Kukkonen, I.T., Paulin, L., Laine, P., Itävaara, M.
17 and Auvinen, P.: Taxonomically and functionally diverse microbial communities in deep
18 crystalline rocks of the Fennoscandian shield. *ISME J.*, 8, 126-138, 2014.
- 19 **Oksanen, J.; Blanchet, F.G.; Friendly, M.; Kindt, R.; Legendre, P.; McGlinn, D.; Minchin,**
20 **P.R.; O’Hara, R.B.; Simpson, G.L.; Solymos, P.; Stevens, M.H.H.; Szoecs, E.; Wagner, H.:**
21 **vegan: Community Ecology Package. R package version 2.4-0. 2016. [http://CRAN.R-](http://CRAN.R-project.org/package=vegan)**
22 **[project.org/package=vegan](http://CRAN.R-project.org/package=vegan).**
- 23 Osburn, M.R., LaRowe, D.E., Momper, L.M. and Amend, J.P. Chemolithotrophy in the
24 continental deep subsurface: Sanford Underground Research Facility (SURF), USA. *Front.*
25 *Microbiol.*, 5, 610, 2014.
- 26 Pedersen, K., Arlinger, J., Eriksson, S., Hallbeck, A., Hallbeck, L. and Johansson, J.:
27 Numbers, biomass and cultivable diversity of microbial populations relate to depth and
28 borehole-specific conditions in groundwater from depths of 4–450 m in Olkiluoto, Finland.
29 *ISME J.*, 2, 760-775, 2008.
- 30 Pedersen, K., Bengtsson, A., Edlund, J. and Eriksson, L.: Sulphate-controlled diversity of
31 subterranean microbial communities over depth in deep groundwater with opposing
32 gradients of sulphate and methane. *Geomicrobiol. J.*, 31, 617-631, 2014.
- 33 Pedersen, K.: Metabolic activity of subterranean microbial communities in deep granitic
34 groundwater supplemented with methane and H₂. *ISME J.*, 7, 839-849, 2013.
- 35 Pedersen, K.: Subterranean microbial populations metabolize hydrogen and acetate under in
36 situ conditions in granitic groundwater at 450 m depth in the Äspö Hard Rock Laboratory,
37 Sweden. *FEMS Microbiol. Ecol.*, 81, 217-229, 2012.
- 38 Posiva. Olkiluoto site description 2011. POSIVA 2011-02, 2013.
- 39 Purkamo, L., Bomberg, M., Nyysönen, M., Kukkonen, I., Ahonen, L., Kietäväinen, R.,
40 Itävaara, M.: Retrieval and analysis of authentic microbial communities from packer-
41 isolated deep crystalline bedrock fractures: evaluation of the method and time of sampling.
42 *FEMS Microbiol. Ecol.*, 85, 324-337, 2013.

- 1 Purkamo, L., Bomberg, M., Nyysönen, M., Kukkonen, I., Ahonen, L. and Itävaara, M.:
2 Heterotrophic communities supplied by ancient organic carbon predominate in deep
3 Fennoscandian bedrock fluids. *Microb. Ecol.*, 69, 319-332, 2015.
- 4 **R Core Team (2013). R: A language and environment for statistical computing. R Foundation
5 for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0, <http://www.R-project.org/>.**
- 6 Rajala, P., Bomberg, M., Kietäväinen, R., Kukkonen, I., Ahonen, L., Nyysönen, M.,
7 Itävaara, M.: Deep subsurface microbes rapidly reactivate in the presence of C-1
8 compounds. *Microorganisms*, 3, 17-33, 2015.
- 9 Schloss, P.D., Westcott, S.L., Ryabin, T., Hall, J.R., Hartmann, M., Hollister, E.B.,
10 Lesniewski, R.A., Oakley, B.B., Parks, D.H., Robinson, C.J., Sahl, J.W., Stres, B.,
11 Thallinger, G.G., Van Horn, D.J., Weber, C.F.: Introducing MOTHUR: open-source,
12 platform-independent, community-supported software for describing and comparing
13 microbial communities. *Appl. Environ. Microbiol.*, 75, 7537–41, 2009.
- 14 **Sedighi, M., Bennett, D., Masum, S.A., Thomas, H.R., Johansson, E.: Analysis of temperature
15 data at the Olkiluoto. Posiva Working Report 2013-58, 1-78, 2013**
- 16 Shade, A. and Handelsman, J.: Beyond the Venn diagram: the hunt for a core microbiome.
17 *Environ. Microbiol.*, 14, 4-12, 2012.
- 18 Sogin, M.L., Morrison, H.G, Huber, J.A, Welch, D.M., Huse, S.M., Neal, P.R., Arrieta, J.M.
19 and Herndl, G.J.: Microbial diversity in the deep sea and the underexplored “rare
20 biosphere”. *Proc. Natl. Acad. Sci. USA*, 103, 12115-12120, 2006.
- 21 Sohlberg, E., Bomberg, M., Miettinen, H., Nyysönen, M., Salavirta, H., Vikman, M. and
22 Itävaara, M.: Revealing the unexplored fungal communities in deep groundwater of
23 crystalline bedrock fracture zones in Olkiluoto, Finland. *Front. Microbiol.*, 6, 573, 2015.
- 24 Staley, C., Gould, T.J., Wang, P., Phillips, J., Cotner, J.B., Sadowsky, M.J. Core functional
25 traits of bacterial communities in the Upper Mississippi River show limited variation in
26 response to land cover. *Front. Microbiol.* 5, 414, 2014.
- 27 Tsitko, I., Lusa, M., Lehto, J., Parviainen, L., Ikonen, A.T.K, Lahdenperä, A.-M. and
28 Bomberg, M. The variation of microbial communities in a depth profile of an acidic,
29 nutrient-poor boreal bog in southwestern Finland. *Open J. Ecol.*, 4, 832-859, 2014.
- 30 Weisburg, W.G., Barns, S.M., Pelletier, D.A. and Lane, D.J. 16S ribosomal DNA
31 amplification for phylogenetic study. *J. Bacteriol.*, 173, 697-703, 1991.
- 32 Wu, X., Holmfeldt, K., Hubalek, V., Lundin, D., Åström, M., Bertilsson, S. and Dopson, M.
33 Microbial metagenomes from three aquifers in the Fennoscandian shield terrestrial deep
34 biosphere reveal metabolic partitioning among populations. *ISME J.*, on line.
- 35

- 1 Table 1. Geochemical and microbiological measurements from 12 different water conductive fractures in the bedrock of Olkiluoto, Finland.
- 2 The different drillholes are presented at the top of the table. The data is compiled from Posiva (2013) and Miettinen et al. (2015)

Drillhole	OL-KR13	OL-KR6	OL-KR3	OL-KR23	OL-KR5	OL-KR49	OL-KR9	OL-KR9	OL-KR2	OL-KR1	OL-KR44	OL-KR29
Sampling date	3/11/2010	18/5/2010	29/8/2011	15/12/2009	16/10/2012	14/12/2009	31/10/2011	29/8/2011	27/1/2010	26/1/2010	15/1/2013	18/5/2010
Depth (m)	296	328	340	347	405	415	423	510	559	572	693	798
Alkalinity												
mEq/L	2.19	0.37	0.47	0.05	0.27	0.16	0.18	0.13	0.29	0.23	0.49	0.13
Ec mS/m	897	1832	1047	2190	2240	2670	2300	2960	4110	3770	6690	7820
pH	7.9	7.9	7.9	7.5	7.9	8.1	7.7	8.1	8.6	7.8	7.5	7.3
NPOC mg L ⁻¹	10	0	12	5.1	19	3	5.1	6.6	11	5	110	10
DIC mg L ⁻¹	27	4.1	4.1	3.9	0	3	3	0	3.75	3.75	6.5	81
HCO ₃ mg L ⁻¹	134	22.6	25	17.1	16	9.8	11.6	7.3	17.7	14	30	424
N _{tot} mg L ⁻¹	0.71	0	1.1	0.42	1.2	0.16	0.38	0.66	1.1	0.41	10	3.1
NH ₄ ⁺ mg L ⁻¹	0.07	0.03	0.03	0	0	0	0.05	0	0.02	0.04	0.08	0.08
S _{tot} mg L ⁻¹	31	130	12	1.7	1.7	0	4.8	0	0	0	4	0
SO ₄ ²⁻ mg L ⁻¹	79.5	379	32	2.9	3	1.4	13.7	0.9	0.5	0.5	9.6	2
S ₂ ⁻ mg L ⁻¹	5.1		0.38	0.62	2	0.02	0.36	0	0.02	0.13	0.02	0.02
Fe _{tot} mg L ⁻¹	0.0042	0.0037	0.022	0.062	0.2	0.71	0.036	0.02	0	0.49	1.2	560
Fe(II) mg L ⁻¹	0	0	0.02	0.08	0.21	0.53	0.06	0.02	0.02	0.04	1.2	0.46
TDS mg L ⁻¹	4994	10670	5656	12710	12880	15900	13430	18580	25500	23260	37410	53210
K mg L ⁻¹	8.2	9.3	7.6	8.3	18	27	12	17	19	20	24	27
Mg mg L ⁻¹	35	77	17	55	68	19	32	41	18	52	33	136
Ca mg L ⁻¹	460	1100	290	2100	1750	2700	2260	2930	4600	3700	7680	10000
Cl mg L ⁻¹	2920	6230	3400	7930	7950	9940	8220	11500	15700	14600	22800	33500
Na mg L ⁻¹	1320	2800	1850	2530	2990	3110	2790	3970	4980	4720	6570	9150
TNC ml ⁻¹	4.2 × 10 ⁵	1.0 × 10 ⁵	2.4 × 10 ⁵	2.5 × 10 ⁵	2.1 × 10 ⁵	1.5 × 10 ⁴	na	2.9 × 10 ⁴	5.9 × 10 ⁴	8.7 × 10 ⁴	5.5 × 10 ⁴	2.3 × 10 ⁴
16S qPCR ml ⁻¹												
bacteria	7.0 × 10 ⁵	9.5 × 10 ³	2.0 × 10 ⁴	3.6 × 10 ⁵	4.9 × 10 ⁴	1.3 × 10 ⁴	7.2 × 10 ⁴	1.5 × 10 ⁵	1.4 × 10 ⁵	1.9 × 10 ⁴	3.2 × 10 ⁴	1.5 × 10 ⁴
archaea	5.8 × 10 ³	2.0 × 10 ⁴	9.9 × 10 ³	6.3 × 10 ⁴	6.2 × 10 ³	1.5 × 10 ²	4.4 × 10 ⁴	5.2 × 10 ²	7.5 × 10 ²	3.0 × 10 ³	2.6 × 10 ¹	2.8 × 10 ²

1 Table 2a. The total number of sequence reads, observed and estimated (Chao1, ACE) number of OTUs, number of singleton and doubleton
2 OTUs, and Shannon diversity index per sample of the bacterial 16S rRNA gene data set. The analysis results are presented for both the total
3 number of sequence reads per sample as well as for data normalized according tot he sample with the lowest number of sequence reads, i.e.
4 140,000 random sequences per sample.


Bacteria	Number of sequence reads	All sequences						Normalized to 140,000 sequences					
		Observed OTUs	Chao1	ACE	Singletons	Doubletons	Shannon	Observed OTUs	Chao1	ACE	Singletons	Doubletons	Shannon
Sample													
OLKR13/296m	786,346	79,527	87,188	91,360	18,025	21,203	13	37,045	74,288	84,530	22,445	6,762	13
OLKR3/318m	345,433	52,381	53,238	54,961	5,789	19,557	14	39,309	57,793	64,021	19,287	10,061	13
OLKR6/328m	188,812	29,411	35,018	37,269	9,209	7,561	13	26,442	34,964	37,626	10,420	6,369	13
OLKR23/347m	485,154	33,257	37,175	38,895	8,000	8,166	11	20,494	34,268	37,305	10,641	4,109	11
OLKR49/415m	184,052	38,275	49,758	53,525	14,799	9,535	13	34,117	48,804	52,938	15,372	8,043	13
OLKR9/423m	175,295	36,412	44,452	47,571	12,357	9,494	14	33,596	44,496	48,161	13,489	8,345	14
OLKR5/435m	141,886	40,445	70,520	78,340	22,166	8,167	14	40,145	70,288	78,232	22,086	8,090	14
OLKR9/510m	241,312	41,545	51,348	54,535	14,251	10,357	13	33,208	49,115	53,631	15,592	7,640	13
OLKR2/559m	257,789	45,456	72,269	78,325	22,550	9,481	13	32,600	62,318	69,573	19,071	6,118	12
OLKR1/572m	210,659	29,804	35,362	37,491	9,197	7,607	12	25,703	34,934	37,682	10,650	6,142	12
OLKR44/750m	303,058	31,410	31,589	32,188	2,005	11,200	12	25,937	33,448	36,295	10,346	7,124	12
OLKR29/798m	221,524	37,989	45,126	48,042	11,991	10,071	13	31,911	44,957	48,533	14,078	7,594	13

5

1 Table 2b. The total number of sequence reads, observed and estimated (Chao1, ACE) number of OTUs, number of singleton and doubleton
2 OTUs, and Shannon diversity index per sample of the archaeal 16S rRNA gene data set. The analysis results are presented for both the total
3 number of sequence reads per sample as well as for data normalized according to the sample with the lowest number of sequence reads, i.e.
4 17,000 random sequences per sample.


Sample	Number of sequence reads	All sequences						Normalized to 17,000 sequences					
		Observed OTUs	Chao1	ACE	Singletons	Doubletons	Shannon	Observed OTUs	Chao1	ACE	Singletons	Doubletons	Shannon
OLKR13/296m	507,373	27,111	29,516	30,699	5,835	7,076	10	3,957	13,380	15,062	2,867	435	10
OLKR3/318m	271,699	25,491	32,299	34,231	9,205	6,221	11	4,955	15,044	17,238	3,546	622	10
OLKR6/328m	446,380	21,597	22,930	23,781	3,861	5,588	10	3,776	11,705	14,020	2,748	475	9
OLKR23/347m	395,339	20,800	22,403	23,214	4,083	5,199	10	3,919	11,855	13,323	2,755	477	9
OLKR49/415m	210,545	22,600	23,372	24,004	2,975	5,733	12	7,023	17,088	19,874	4,738	1,114	12
OLKR9/423m	697,360	22,014	22,527	23,082	2,381	5,520	9	3,180	9,617	10,586	2,224	383	9
OLKR5/435m	769,026	21,127	22,235	23,078	3,515	5,574	9	2,596	10,114	10,078	1,852	227	9
OLKR9/510m	169,142	12,709	12,782	12,960	713	3,488	11	4,879	11,205	13,215	3,148	782	11
OLKR2/559m	100,101	15,359	24,950	27,026	7,840	3,203	11	5,119	14,497	16,488	3,548	670	11
OLKR1/572m	1,213,360	28,884	33,207	34,832	7,846	7,118	9	2,273	9,233	9,923	1,631	190	9
OLKR44/750m	17,716	6,436	8,748	9,750	2,890	1,805	12	6,325	8,743	9,804	2,921	1,763	12
OLKR29/798m	98,770	15,641	16,720	17,483	3,158	4,617	12	6,951	14,655	17,184	4,483	1,303	12

5




1
 2 Figure 1. Map of Olkiluoto. The boreholes used in this study are marked with a turquoise
 3 triangle and the attached black line depicts the direction of the borehole. (with courtesy of
 4 Pöyry Oy, Nov 17th, 2015 by Eemeli Hurmerinta)


5




6
 7 Figure 2. The concentration of A) microbial cells mL⁻¹ determined by epifluorescence
 8 microscopy and the estimated concentration of B) bacterial and C) archaeal 16S rRNA gene
 9 copies mL⁻¹ groundwater determined by qPCR in water conductive fractures situated at
 10 different depths in the Olkiluoto bedrock.




1
 2 Figure 3. The core A) archaeal and B) bacterial community OTUs detected from at least 80%
 3 of the samples with heatmaps on the abundance of the C) archaeal and D) bacterial core
 4 community profiles. In A) and B) the OTUs are stacked in the columns according to the
 5 number of sequence reads, with the most abundant OTUs at the bottom of the columns. The
 6 OTU segments of the columns are colored according to the family to which they belong. Each
 7 Order is presented as a separate column for each sample.



8
 9
 10 Figure 4. The number of shared A) archaeal and B) bacterial OTUs in the different samples.
 11 The number of shared OTUs is shown on the Y-axis and the proportion of samples on the X-
 12 axis.




1
 2 Figure 5. A UPGMA cladogram clustering the samples based on the A) archaeal and B)
 3 bacterial OTU profile according to the Bray Curtis distance model. Red colour indicates low
 4 abundance and yellow colour indicates high abundance. C) and D) show the corresponding
 5 physicochemical parameters as shown in Table 1, with the lowest values in green, medium
 6 values in yellow and high values in red.



1
2
3 Figure 6. Principal coordinates analysis (PCoA) on the whole OTU profiles of the different
4 samples based on Bray Curtis distance model. The analyses for archaea are shown in A) C)
5 and E, and the bacteria in B), D) and F). In A) and B) the points indicate water type, where
6 the circle is for brackish sulphate-rich water, the square is for brackish chloride-rich water and
7 the diamond is for saline water. The colouring of the points are according to concentration of
8 total solids as indicated in the upper left corner of figure A. In C) and D) the different sizes of
9 the points describe the library size and the positions are the same as in A) and B). E) and F)
10 display the proportion of variance for 12 dimensions, of which 1 and 2 were used for plotting
11 the PCoA.


1



2


3 Figure 7. The nearest sequenced taxon index (NSTI) values calculated by PICRUSt for the
4 bacterial (blue) and archaeal (red) communities. The NSTI value describes the sum of
5 phylogenetic distances of each OTU to its nearest relative with a sequenced reference
6 genome, and measures substitutions per site in the 16S rRNA gene and the weighted the
7 frequency of the each OTU in a sample dataset. A higher NSTI value indicates greater
8 distance to the closest sequenced relatives of the OTUs in each sample.

A.



1


B.



2

3 Figure 8. The relative abundance of predicted genes of the most abundant pathways identified
4 in the A) bacterial and B) archaeal populations in the PICRUSt analysis. The pathways are
5 presented according to KEGG. The samples are ordered according to depth, with OL-
6 KR13/296m as innermost and OL-KR29/798m as the outermost sample.


7




1
 2 **Figure 9.** The microbial carbon metabolism pathway according to KEGG. The predicted
 3 genes combined from all samples were plotted on the map. Green arrows indicate enzymes
 4 predicted only in the archaeal communities, red arrows indicate genes predicted only in the
 5 bacterial communities, black arrows show enzymes predicted in both the archaeal and
 6 bacterial communities.

7
 8
 9

1 Supplementary figures




2
3 Figure S1. The rarefaction curves of observed bacterial (left pane) and archaeal (right pane)
4 OTUs in each sample generated on sequence data normalized to 140,000 reads for bacteria
5 and 17,000 reads for archaea.




7
8 Figure S2. Bacterial oxidative phosphorylation according to KEGG. The predicted genes from
9 the bacterial communities belonging to the oxidative phosphorylation are shown in pink.

10




1



2

3 Figure S3. Predicted genes shown in red of the bacterial A) fatty acid degradation and B) fatty
 4 acid biosynthesis pathways, combined from all samples.




1
2 Figure S4. The predicted genes of enzymes included in the microbial nitrogen metabolism
3 according to KEGG. Enzymes predicted from the bacterial communities are shown in red,
4 archaeal communities in green and enzymes predicted from both archaeal and bacterial
5 communities in green/red. Enzymes not predicted from either community are shown in blue
6 or white.

7

Mineral and organic ion transporters




Metallic cation, iron-siderophore and vitamin B12 transporters



8

1 Figure S5. The genes of ABC transporters predicted from the bacterial (pink), archaeal
 2 (green) or both (pink/green) communities. Genes not predicted in any of the communities are
 3 shown in blue.

4



5

6 Figure S6. The predicted genes of enzymes included in the microbial sulphur metabolism
 7 according to KEGG. Enzymes predicted from the bacterial communities are shown in red,
 8 archaeal communities in green and enzymes predicted from both archaeal and bacterial
 9 communities in green/red. Enzymes not predicted from either community are shown in blue
 10 or white.

11

12

1 **Author responses to Reviewer 1**

2 Reviewer 1

3

4 The manuscript entitled, 'Estimation of microbial metabolism and co-occurrence patterns in
5 fracture groundwaters of deep crystalline bedrock at Olkiluoto, Finland', analyzes the
6 community composition and function of microbial communities in the Finnish subsurface
7 (296-798 m depth). A bacterial and archaeal V6 16S rDNA dataset from 12 different
8 boreholes is the focus of the authors's analysis and discussion. Metadata such as the pH, the
9 concentration of anions and cations, and relative abundance of bacteria and archaea are also
10 reported. From this data, the authors use techniques such as NMDS, network analysis, and
11 PICRUST to analyze the data. I do not believe that the methods outlined in Section 2.7
12 Statistical analyses were applied and analyzed appropriately.

13

14 **Author response; Thank you for your comments. They have all been answered systematically**
15 **below. However, due to all the changes done to the manuscript we would like to propose a**
16 **new name for the paper; 'Microbial communities and their predicted metabolic characteristics**
17 **in deep fracture groundwaters of the crystalline bedrock at Olkiluoto, Finland'**

18

19 Major Comments:

20

21 My first concern is about the taxonomic data matrix from which the statistical analyses were
22 made. From what I can tell, these analyses were performed on counts of taxonomic labels that
23 are illustrated in Figure 3. Based on the legend of Figure 3, it appears that a general
24 taxonomic cut-off was not applied as there are genera such as 'Microbacterium' and then
25 broad categories of phyla such as 'Actinobacteria_Other'. Since Figure 7 illustrates that the
26 V6 sequences did not have good matches to sequences in the reference database the
27 'phylaX_Other' counts are inflated relative to other genera. This imbalance in taxonomic
28 labels is particularly concerning as it makes the interpretation of the statistical analyses
29 downstream difficult. For example, two sites may have similar relative abundances of
30 'phylaX_Other' but the community composition of taxa within the broad category may vary

1 dramatically. Therefore, I would recommend that OTUs or a broader taxonomic classification
2 be used as the taxonomic data matrix in statistical analyses.

3

4 **Author response; Agreed. The NMDS plot has been changed to PCoA plots calculated based**
5 **on the OTU tables (Figure 6).**

6

7 Another concern is in the co-occurrence network analysis. The cut-offs are very relaxed
8 ($R > 0.7$, $p < 0.01$). This leads to an extremely large network with many spurious correlations.
9 As presented, I don't believe network is not informative.

10

11 **Author response; The network was omitted.**

12

13 In the NMDS plots, I think that there should be an indication of how the environmental
14 variables were placed on the NMDS plot or how statements of the type 'No clear
15 environmental factor was identified to drive the communities' (page 9, line 20-25) were
16 made. As written, one is unable to interpret or reproduce their analyses.

17

18 **Author response; The NMDS plot was recalculated using only the OTU tables. The chemistry**
19 **was moved to the OTU heatmap and clustering figure (Fig 5) and added only as absolute**
20 **values in a heatmap below the OTU heatmaps to show the chemistry in connection to how the**
21 **samples were grouped (Figure 5) the description has been clarified.**

22

23 Minor Comments:

24

25

26 Abstract

27

28 - I am confused as to what page 1, lines 15-17 mean

1

2 Author response; The text was clarified.

3

4 Introduction

5

6 - Page 2, line 17: did the previous efforts provide information on the relative abundance of
7 archaea to bacteria? This is a nice point of your MS would be good to know if there are any
8 estimates of bacteria and archaea proportions in the introduction (or if there are any general
9 trends about the proportion of bacteria and archaea)

10

11 Author response; the proportion of archaea vs. bacteria in Olkiluoto deep groundwater has not
12 been published before. Usually the bacteria greatly outnumber the archaea, but as seen in the
13 present data, in specific samples the archaea are pretty strong. I have added a reference from
14 Outokumpu (Finland), where the archaea at most made up only 1% of the total microbial
15 community.

16

17 - Page 3, lines 4-12 it is unclear what environments the citations are from

18 Author response; the text has been clarified.

19

20 Methods

21

22 - Page 5, line 9: what are the negative DNA isolation controls?

23 Author response; these were reagent controls, which means that we submitted plastics and
24 reagents to the same extraction protocol as we did with the actual samples in order to show
25 that the level of contaminating DNA originating from the extraction kits. The text was
26 rewritten.

27

28 Section 2.7

1 - It is unclear what the input data for the statistical analyses were (are they taxa, OTUs, etc).
2 Based on Figure 3, I wonder if the statistical tests were performed with taxonomic labels. I
3 think that these analyses would be better served on the OTUs the authors generated as OTUs
4 are a type of reference free data. Using OTUs would be especially useful as the majority of
5 taxa (based on Figure 3) are labeled as 'phylaX_Other'.

6

7 Author response; we used the taxonomic tables. However, we recalculated the PCoA and
8 UPGMA clustering using the OTU tables as presented in Figures 5 and 6.

9

10 - Why was the Chord distance used for NMDS and the Bray-Curtis for UPGMA clustering? Is
11 there a reason why the same similarity metric used in both?

12

13 Author response; Good point. Bray-Curtis is now used for both analyses.

14

15 Results

16

17 - Due to the difficulty it may be difficult to report temperature of the groundwater. However,
18 it may be useful (for the readers) to report temperature if any information is available.

19

20 Author response; it is known that the temperature of the groundwater is between 10°C and
21 16°C at these depths, increasing linearly with depth. This has been added with a reference to
22 where the data has been reported before. The temperature specifically for our samples was not
23 possible to obtain. The temperature information has been added to the M&M 2.1 Background.

24

25 Discussion

26

27 - Page 12, lines 21-25: I am unsure what the authors are trying to say

28 Author response; we are just trying to say that using the HiSeq we obtained a load more data

1 from which we could identify a lot more taxa.

2

3 - Page 13, lines 4-18: as presented, I am unsure how these interpretations were taken from the
4 NMDS plots. I think that the discussion on this topic needs to be improved

5 Author response; the discussion was changed. However, now when we have used the OTU
6 tables the effect of the environmental factors seem to fade away and we do not see any clear
7 environmental factor that would drive the communities any more.

8

9 - Page 14, lines 2-4: See my concerns about the network in major comments.

10 Author response; the network analysis was omitted and the text corrected accordingly.

11

12 - Page 14, lines 15-16: I am unsure what the logic behind the statement ‘Thus, on higher
13 taxonomical level common traits for specific groups of microorganisms may be revealed’.
14 The writing in the paragraph could be improved to clarify this section.

15 Author response; The text was changed.

16

17 - Page 15, lines 13-20: I found this section difficult to follow

18 Author response; The text was modified.

19

20 - Page 15, lines 24-25: when referring to PICRUST functions, the authors should use
21 ‘predicted’ or similar indication to highlight it is based on the PICRUST results and not other
22 data (such as rates etc).

23 Author response; This has been corrected.

24

25 - The authors refer to methane oxidation several times but do not mention the possibility for
26 anaerobic methane oxidation using a reversed methanogenesis pathway.

27

1 Author response; Actually, on P12L15-16, we say that the AOM may be a common pathway
2 at depths 328 - 423 m. We have included more discussion about this. We see the prediction in
3 Fig 9, but the arrows only go one way...

4

5 Tables and Figures

6

7 - Table 2ab, columns labeled 'Singles' and 'Doubles' to 'Singletons' and 'Doubletons' as
8 caption

9 Author response; Corrected.

10 - Table 2b appears twice

11 Author response; Corrected.

12

13 - Figure 3: What are minor groups (e.g. are they known members of the phyla that are colored
14 as XXX_Other)? I feel like this figure is misleading about taxonomic composition. Based on
15 the legend it appears that a general taxonomic cut-off was not applied as there are genera such
16 as 'Microbacterium' and then broad categories of phyla such as 'Actinobacteria_Other'. Since
17 Figure 7 illustrates that the V6 sequences did not have good matches to sequences in the
18 reference database the 'phylaX_Other' counts are inflated relative to other genera. This
19 imbalance in taxonomic labels may also cause issues in the other statistical analyses
20 downstream.

21

22 Author response; The statistical analyses have now been performed on the OTU tables. Figure
23 3 has been changed to cover only core community OTUs, i.e. OTUs found in at least 80% of
24 the samples. In addition, a new figure 4 has been added showing how the number of shared
25 OTUs drop dramatically when more samples need to share OTUs for thOTUs to be
26 considered core.

27

28

29 - For Figure 9, it would be easier to read if the connections not observed were faded out to

1 emphasize which connections (pathways) were observed

2 Author response; The Figure was edited to fade out the routs that were not seen.

3

4 - I did not see the supplementary tables referred to in the author's response (page 51 line 31)

5 Author response; the tables were uploaded upon manuscript submission, so something must
6 have gone wrong after that. I can see the tables on my submission page.

7