



Zurich, 8th of February 2016

Manuscript bg-2015-347: Time since death and decay rate constants of Norway spruce and European larch deadwood in subalpine forests determined using dendrochronology and radiocarbon dating

Dear Sir or Madam,

Dear Prof. Schöngart,

Enclosed please find the manuscript with the title 'Time since death and decay rate constants of Norway spruce and European larch deadwood in subalpine forests determined using dendrochronology and radiocarbon dating'. As far as possible, we tried to consider all advices of the reviewer.

The work has not been published previously and is not under consideration for publication elsewhere. The publication is approved by all authors and, if accepted, it will not be published elsewhere in the same form.

I thank you for your efforts.

Yours sincerely

Prof. Markus Egli
University of Zurich
Department of Geography

Response to the reviewers' and readers' comments (manuscript bg-2015-347)

We have received the comments of two reviewers (Reviewer 1 and 2) and two readers (T. Kahl and V.-A. Angers). As far as possible, we tried to consider all the suggestions. In response, we made several changes according to the suggestions and criticisms:

- corrections in the manuscript
- additional calculations (as suggest by T. Kahl) using the stage-based matrix model of Kruys et al. (2002) and corresponding addition of the data obtained
- extended discussion and better implementation of the problem of CWD age overestimation (including the fall rate of snags)
- Tables 3 and 4 are now deleted
- Figure 3 is now redrawn

Our point-by-point response to the reviewers' comments is given below. In most points we agree with the reviewers but not for all.

In red: comments of the reviewers

In black: our response

Anonymous referee #1

1) Major concerns:

I mainly suggest delivering more information about the sampling protocol, the sampling sites and the wood composition

Referee 1' main criticism regards the sampling protocol, wood composition and the comparison of coarse woody debris (CWD) decay mechanisms with organic matter decomposition in soils.

We now give more details about the sampling procedure, added the cellulose, lignin and density values as requested by reviewer 1 and revised the discussion part about soil organic matter.

The comparison with organic matter decomposition in soil is not helpful...

The corresponding paragraph is now revised (see also comment below).

2) Details:

p. 14799 l.1: Use "amount", not size.

We really mean size: "amount" is not the correct term as you suggest. It is not trivial to cut and transport to the lab sections from tree trunks (having large dimensions), especially in the early stage of decomposition when the wood is still close of being intact - also considering difficult field conditions such as steep mountain slopes that are not accessible with transport facilities.

p. 14799 l.3: use "rarely" investigated, not poorly.

Ok, we now use 'rarely'.

p. 14799 I.14: are the ages similar or varying, or similarly varying?

This is now corrected.

p. 14799 I. 19: Review the sentence

the sentence is now corrected.

p. 14801 I. 8-12: cite the original references

Publications considering multiple parameters (multiple-exponential model) are:

Wider R.K. and Lang G.E.: A critique of the analytical methods used in examining decomposition data obtained from litter bags, *Ecology*, 63, 1636-1642, doi: 10.2307/1940104, 1982.

Means, J. E., Cromack, K., and MacMillan, P. C.: Comparison of decomposition models using wood density of Douglas-fir logs, *Can. J. Forest R.*, 15, 1092-1098, doi: 10.1139/x85-178, 1985.

Minderman, G.: Addition, decomposition and accumulation of organic matter in forests, *J. Ecol.*, 56, 355-362, doi: 10.2307/2258238, 1968.

Publications considering the lag-time are for example:

Harmon, M. E., Franklin, J. F., Swanson, F. J., Sollins, P., Gregory, S. V., Lattin, J. D, Anderson, N. H., Cline, S. P., Aumen, N. G., Sedell, J. R., Lienkaemper, G. W., Cromack, K., and Cummins, K. W.: Ecology of coarse woody debris in temperate ecosystems, *Adv. Ecol. Res.*, 15, 133–302, doi: 10.1016/S0065-2504(08)60121-X, 1986.

p. 14801 I. 17: Please give a short definition of CWD

A short definition is now given. For a better understanding, this definition is already given in the second sentence of the 'Introduction'.

... and: Yes the studies that were cited before consider CWD.

p. 14802 I. 15-16: would suggest not to relate your results to soil organic matter, especially without giving any references for your statement.

This part is now deleted

p. 14803: The sampling protocol should be complemented with information about the size of your sampling plots and the number of CWD samples taken. Interestingly, larch was mainly sampled at the South-facing sites while spruce was mainly sampled at the North-facing sites.

More information about the sampling protocol is given. We hope that it is clearer now.

p. 14804 I. 15: The methods to determine cellulose and lignin content are described here although the results have been published in another paper. I would suggest adding the results in a table...

The lignin and cellulose values are now added in Tables 5 and 6.

In the methods section: twice the same solution (5% NaOH), is this correct?

A 5% NaOH solution was indeed applied two times (methods). Each time, the NaOH solution was discarded.

p. 14806 l. 1: It would be helpful to add the density values,

The density values are now added in Tables 5 and 6.

p. 14807: The ages of the CWD cluster in two age groups of about 30-50 years and 205-220 years, considering the first row of average age in Tab. 6. How can this be explained, is it a result of the wide range of calibrated ages?

Radiocarbon dating gives for such ages (about 200 years) not hyper-precise results (due to 'plateaus' in the calibration procedure). This is why we added the range with the highest probability (Table 6) to restrict the rather broad ranges. We finally used these values to 'overcome' this difficulty. We added a paragraph about this issue. It is also implemented in the discussion, conclusion and abstract.

p. 14811-12: The paragraph on soil organic matter is unnecessary as it is now....

This paragraph is now revised (large part is omitted) and a less strict relation to soil organic matter is given. Furthermore, the suggested literature is now considered.

Reader T. Kahl

I was a bit surprised by the low decomposition rates that you found for the observed tree species. Could it be that a sampling bias is part of the explanation for these low decomposition rates. In Kruys et al. 2002 (Fig. 2) it is nicely shown that: "Snap-shot sampling at time t means that the proportion of slow-decaying trees will be overestimated".

Kruys' et al. (2002) model is now implemented (in Tables 5 and 6) and discussed (see our interactive response "bgd-12-C8571-2015"). The results obtained using this model only marginally differ from the single-negative-exponential model. We consequently have to assume that there was no significant sampling bias.

Reader V.-A. Angers

... the discussion would be improved if the influence of species-specific snag fall rates were acknowledged ...

The discussion about fall rates of snags is now included. We have to assume that European larch and Norway spruce do not show great differences regarding the fall rate of snags (see our interactive response "bgd-12-C9144-2016").

Referee #2 (J. Schöngart)

1) Major concerns

Tree-ring chronologies from living trees: At each site (n=8) samples were collected from two radii of 5-

6 trees (P. 14803, L. 6-8). Later in the paragraph the authors state a total sampling size of 83 wood cores (29 from larch and 54 from spruce). The authors should better describe how many trees (cores) from each species were sampled at each site.

The table below shows the number of living trees that were sampled at each site. From each tree two cores were taken. In three cases, one of the two cores taken from the same tree were too damaged to be measured (one at N02, one at S06 and one at S07). Therefore, the number of sampled cores and that of measured cores is not always the same. Furthermore, two outliers were excluded from the master chronology, namely one measured core at S07 and one at S08.

Table 1. Number of sampled living trees.

Site	Tree species	n. of sampled trees	n. of sampled cores	n. of measured cores	Outliers
N01	Norway spruce	5	10	10	
N02	Norway spruce	6	12	11	
N03	Norway spruce	5	10	10	
N04	European larch	4	8	8	
S06	Norway spruce	6	12	11	
S07	European larch	5	10	9	1
S08	Norway spruce	6	12	12	1
S09	European larch	6	12	12	

This is now mentioned accordingly in the manuscript text.

The authors state at P. 14806 (L. 18-24) that larch seems to have a more sensitive growth than spruce and the positive and negative pointer years are not synchronous between both species. This might be also result of differences in leaf phenology (spruce is an evergreen species and larch a deciduous species) or an unequal distribution of sampled individuals among the different sites with varying climate conditions and exposition. Even if this is not the main focus of this paper it should be better explained.

An explanation is now given in the text (cf. chapter 3.1).

Indicate the correlation coefficient between both master chronologies. In figure 3 the sample size should be indicated as number of trees not by the number of cores. As I understood the two cores have been cross-dated to a single tree curve which was used to produce the master chronology.

This is now done. The inter-correlation coefficient between the two master chronologies is 0.274 (Pearson correlation).

CWD dating: P. 14803 (L. 21/22) that a total number of dated deadwood... However, counting the numbers of dated larch trees in these tables indicates a higher sample size of a total of 23 trees....

Yes indeed, there was a mistake here. It is now corrected.

Table 5 and Appendix A: I suggest to present all data in one table. It would be also interesting to show if the two dating techniques (cross-dating and radiocarbon dating) come up to the same result.

We think that there is misunderstanding (?). In Table 5, ALL data (decay class 1 – 3) is presented. In Appendix A, only the details of the dated CWD in decay classes 1 – 3 using ^{14}C is presented. We furthermore do not compare the dendro cross-dating with the ^{14}C dating technique (this methodological aspect already has been done by Krüger et al., 2014). Consequently, we think that it is necessary to keep the data presented in Appendix A. It is an addition to Table 5 (and it cannot be unified with Table 5). We slightly changed the caption of Table 5 and hope that it is now less misleading.

Temperature varies along the altitudinal gradient and also rainfall increases about 60% from the lowest altitude to the highest altitude as indicated in table 1. ... how is this correlated with the decay rates?

The correlation analysis using in addition temperature and precipitation is now done. We performed the Kruskal-Wallis test (as it was already done for the other factors: elevation, exposition, species and decay class; now with temperature and precipitation) to assess the influence of these factors on the k -values. Temperature and precipitation obviously did not significantly affect the decay constant k (MAT: p -value = 0.7082; MAP: p -value = 0.4835).

Radiocarbon dating: Due to the Suess effect samples dated of before 1950 AD have widely calibrated age ranges. For me it is not clear how the authors estimated the age range with the highest probability. The high variation of atmospheric radiocarbon due to the high amount of fossil burning during the industrial revolution results in up to five possible ages for one radiocarbon age in the period between 1640 and 1950, which makes it rather difficult to date the dead wood samples.

We now give more information about the dating procedure and discuss our results more critically. It is indeed an important point and we now even address this issue in the abstract.

Radiocarbon dating of CWD having a post-bomb age gives similar results compared to dendrochronological analyses (as shown by Krüger et al., 2014). One major problem in determining the age using ^{14}C is the partially large age range after calibration (due to plateaus) for samples having a pre-bomb age. We showed the 2- σ range in Table 6 (now Table 4), but used the range with the highest probability (that varied, depending on the sample, from 50 to 82.2%; former Table 6; for comparison, commonly the age 1- σ range, i.e. 68%, are reported in literature! ... this already gives rise to a narrower range) for CWD dated to the time period before the bomb peak. This procedure introduces consequently an uncertainty. These methodological constraints have to be kept in mind – a solution of the problem is not available.

Using OxCal and IntCal 13, we obtained the following results for the example that you mention (this was, however, the worst example we had):

14-C age: 170 yr

error: 31 yr

Calibrated age (and probabilities): 1658 – 1700 cal AD (17.4%), 1720 – 1819 cal AD (50%), 1832 – 1880 cal AD (8.6%), 1915 – 1950 (19.5%). In total 95.5% probability.

2) Minor concerns

P. 14802, L. 1-2: The authors should indicate the wood chemical differences between both species.

The wood chemical differences are small (see p. 14801, L. 28). We now present the values for the most evident difference.

P. 14804, L. 16/17: Please indicate which part of the dead wood was sampled to determine α -cellulose. Was this the outermost part of the sample (sapwood)?

Yes. This is now added.

P. 14806, L. 14/15: Please indicate the range of GLK (means and standard deviation) of trees from the same species also considering different sites

We now calculated the GLK values for each single plot.

Table 2. GLK values and standard deviation for each plot.

Site	Tree species	Mean GLK (standard deviation)
N01	Norway spruce	0.58 (0.13)
N02	Norway spruce	0.63 (0.08)
N03	Norway spruce	0.62 (0.08)
N04	European larch	0.69 (0.09)
S06	Norway spruce	0.68 (0.11)
S07	European larch	0.66 (0.10)
S08	Norway spruce	0.68 (0.10)
S09	European larch	0.70 (0.10)

P. 14807, L. 3: Please indicate how many outliers were excluded from the chronology.

To maximise the common signal, one outlier per each species was excluded from the relative master chronology. These two measurements gave a poor correlation with the others, probably due to an elevated number of missing rings.

Table 3: this table contains few data it could be dropped

The table is now skipped and the values are given in chapter 2.6.

Table 4 contains information on the two master chronologies developed by living trees. As this information is already indicated in the text, this table could be dropped to reduce the amounts of tables in the manuscript.

Table 4 is now deleted.

The data of inter-series correlation could also be shown as additional graph in figure 3 for segments of constant periods (25 years for instance). It also would be interesting to calculate the expressed population signal (EPS) for those segments for the two master chronologies according to Wigley et al. (1984), in order to quantify the degree to which samples represent the hypothetical noise-free chronology (EPS-values should achieve more than 0.85 which is a commonly applied quality threshold according to Wigley et al., 1984).

The EPS values are now calculated and displayed in Figure 3.

Figure 1 shows the eight sample sites indicated as “N” and “S”. Please indicate what does “N” and “s” stand for (north-facing and south-facing sites).

This is now done in the figure caption.

We gratefully acknowledge the constructive suggestions of the referees and readers to improve our manuscript.

References:

Krüger, I., Muhr, J., Hartl-Meier, C., Schulz, C., and Borke, W.: Age determination of coarse woody debris with radiocarbon analysis and dendrochronological cross-dating, *Eur. J. Forest Res.*, 133, 931–939, doi: 10.1007/s10342-014-0810-x, 2014.

1 **Time since death and decay rate constants of Norway spruce and European**
2 **larch deadwood in subalpine forests determined using dendrochronology and**
3 **radiocarbon dating**

4

5 M. Petrillo^{1,2}, P. Cherubini², G. Fravolini⁴, M. Marchetti⁴, J. Ascher^{5,6}, M. Schärer¹, H.-A. Synal³, D.
6 Bertoldi⁷, F. Camin⁷, R. Larcher⁷, and M. Egli^{1*}

7

8 ¹Department of Geography, University of Zurich, 8057 Zurich, Switzerland

9 ²WSL Swiss Federal Institute for Forest, Snow and Landscape Research, 8903 Birmensdorf,
10 Switzerland

11 ³Laboratory of Ion Beam Physics, ETH Zurich, 8093 Zurich, Switzerland

12 ⁴Department of Bioscience and Territory, University of Molise, 86090 Pesche, Italy

13 ⁵Department of Agrifood and Environmental Science, University of Florence, 50144 Florence, Italy

14 ⁶Institute of Microbiology, University of Innsbruck, 6020 Innsbruck, Austria

15 ⁷Fondazione Edmund Mach, 38010 San Michele all'Adige, Italy

16

17 *Corresponding author. Tel.: +41 44 635 51 14; fax: +41 44 6356848.

18 E-mail address: markus.egli@geo.uzh.ch (M. Egli).

19

20

21

22

23

24

25 Abstract

26 Due to the large size (e.g. sections of tree trunks) and highly heterogeneous spatial distribution of
27 deadwood, the time scales involved in the coarse woody debris (CWD) decay of *Picea abies* (L.)
28 Karst. and *Larix decidua* Mill. in Alpine forests are largely unknown. We investigated the CWD
29 decay dynamics in an Alpine valley in Italy using the chronosequence approach and the five-decay
30 class system that is based on a macromorphological assessment. For the decay classes 1 to 3, most
31 of the dendrochronological samples were cross-dated to assess the time that had elapsed since tree
32 death, but for decay classes 4 and 5 (poorly preserved tree rings) radiocarbon dating was used. In
33 addition, density, cellulose and lignin data were measured for the dated CWD. The decay rate
34 constants for spruce and larch were estimated on the basis of the density loss using a single negative
35 exponential model, a regression approach and the stage-based matrix model. In the decay classes 1
36 to 3, the ages of the CWD were similar and varied between 1 and 54 years for spruce and 3 and 40
37 years for larch with no significant differences between the classes; classes 1 – 3 are therefore not
38 indicative for deadwood age. This seems to be due to a time lag between the death of a standing tree
39 and its contact with the soil. We found distinct tree species-specific differences in decay classes 4
40 and 5, with larch CWD reaching an average age of 210 years in class 5 and spruce only 77 years.
41 The mean CWD rate constants were estimated to be in the range 0.018 to 0.022 y⁻¹ for spruce and to
42 about 0.012 y⁻¹ for larch. Snapshot sampling (chronosequences) may overestimate the age and mean
43 residence time of CWD. No sampling bias was, however, detectable using the stage-based matrix
44 model. Cellulose and lignin time trends could be derived on the basis of the ages of the CWD. The
45 half-lives for cellulose were 21 y for spruce and 50 y for larch. The half-life of lignin is
46 considerably higher and may be more than 100 years in larch CWD. Consequently, the decay of
47 *Picea abies* and *Larix decidua* is very low. Several uncertainties, however, remain: ¹⁴C dating of
48 CWD from decay classes 4 and 5 and having a pre-bomb age is often difficult (large age range due

Markus Egli 16.10.2015 07:16

Gelösch: i.e.

Markus Egli 23.1.2016 17:10

Gelösch: have been poorly rarely investigated and

Markus Egli 24.1.2016 17:48

Gelösch: commonly employed for forest surveys,

Markus Egli 23.1.2016 19:59

Gelösch: and visual

Markus Egli 23.1.2016 17:11

Gelösch: and some others not having enough tree rings,

Marta Petrillo 13.10.2015 16:04

Gelösch: similar

Markus Egli 16.10.2015 07:19

Gelösch: ly

Markus Egli 16.10.2015 07:19

Gelösch: ying

Markus Egli 23.1.2016 17:15

Gelösch: .

Markus Egli 23.1.2016 17:15

Gelösch: however,

Markus Egli 23.1.2016 17:16

Gelösch: 2

Markus Egli 23.1.2016 17:16

Gelösch: 18

Markus Egli 23.1.2016 17:16

Gelösch: 0.005

Markus Egli 23.1.2016 17:18

Gelösch: to

Markus Egli 16.10.2015 07:27

Gelösch: half-lives (using a multiple-exponential model)

66 [to methodological constraints\) and fall rates of both, European larch and Norway spruce are](#)
67 [missing.](#)

68

69 Keywords: coarse woody debris, decay constants, Norway spruce, European larch, subalpine
70 forests, cellulose, lignin

71

72

73 **1 Introduction**

74 The quantity and residence time of deadwood [or coarse woody debris](#) in Alpine forests are crucial
75 in assessing the carbon cycle to ensure sustainable management of forests. [Coarse woody debris](#)
76 [\(CWD\) is defined as large-sized deadwood pieces, such as stems of dead trees lying on the forest](#)
77 [floor, standing dead trees and stumps, big branches and wood boles in all stages of decomposition.](#)

78 Deadwood plays an important role in maintaining biodiversity in forest ecosystems (Müller and
79 Bütler, 2010) as well as storing carbon (Di Cosmo et al., 2013), and contributing to nutrient cycle
80 processes (Palviainen et al., 2010). The amount of deadwood varies greatly from managed to
81 natural forests. In managed European Alpine forests, for example, the average stock of deadwood is
82 estimated to be about 26 m³/ha, while in old growth Alpine coniferous forests it can be up to 150 –
83 190 m³/ha (Barbati et al., 2014). Residence time for deadwood (e.g. Krüger et al., 2014) — from the
84 moment the tree reaches the forest floor until it loses 95 % of the mass — can range from decades
85 to several hundred years, depending on intrinsic and external factors. These factors include the
86 dimensions of the log, the wood chemistry and the site conditions, in particular the mean annual
87 temperature and soil moisture.

88 Until now, various different sampling designs have been used to determine the time since death to
89 estimate the decay rate of deadwood. Long-term studies can provide reliable results (Müller-Using
90 and Bartsch, 2009), but the slow decay dynamics of wood usually require a decadal observation

Markus Egli 23.1.2016 17:19

Gelöscht: -

... (1)

93 period. Bond-Lamberty and Gower (2008) used the ratio of deadwood mass input into the pool of
94 initial deadwood to estimate its decay rate based on a 7-years observation period. Such time
95 sequences (chronosequence) offer ideal scenarios to study deadwood dynamics. If windthrow, fire
96 regeneration and harvest events are known, the starting point in the timeline of the decay process
97 can be specified. However, the exact year of such events is often uncertain which means precisely
98 dating a tree's death is critical. Dendrochronology can be a helpful tool to determine the year of
99 death, and the technique has been used in several studies to determine the time elapsed since tree-
100 death (Campbell and Laroque, 2007, Lombardi et al., 2008, 2013). Other researchers have used
101 radiocarbon dating to date the last recognizable ring of deadwood. For example, Kueppers et al.
102 (2004) estimated the turnover time of lodgepole pine along a subalpine elevation gradient and
103 Krüger et al. (2014) compared tree-ring cross-dating and radiocarbon dating, demonstrating that the
104 two techniques produce comparable results. The decay rate can be estimated by relating the time-
105 since-death to the density loss or mass loss of deadwood during a given time period (e.g. Busse,
106 1994; Melin et al., 2009). The decay rate is commonly expressed through a decay constant k , which
107 indicates the density loss or mass loss per year. This constant is derived from a decay model
108 (Harmon, 1986), which can be most simply expressed by the equation

$$x_t = x_0 e^{-kt} \quad (1)$$

110 (single-negative-exponential model), where x_t is the density or mass of deadwood at a given time,
111 and x_0 is the initial density or mass (Jenny et al., 1949; Olson, 1963). Other decay models have also
112 been developed that take wood decomposition into account (reviewed by Mackensen et al., 2003).
113 Several authors (Minderman, 1968; Wider and Lang, 1982; Means et al., 1985) consider the
114 different wood components, e.g. bark, sapwood, heartwood and chemical compounds, and combine
115 them in multiple-exponential equations. Other authors (e.g., Harmon et al., 1986) consider the time
116 elapsed from the death of a standing tree to the moment when it falls and comes in contact with the
117 forest floor (lag-time models). In several environments, e.g. on dry mountain slopes, the time lag
118 between death and contact with the forest floor can last for almost the entire decay process

Markus Egli 16.10.2015 07:39

Gelöscht: Some

Marta Petrillo 13.10.2015 16:08

Gelöscht: models

121 (Kueppers et al., 2004). A few models take not only the losses due to heterotrophic respiration and
122 leaching into account, but also losses due to fragmentation (Mackensen et al., 2003).

123 One of the most important components of deadwood is coarse woody debris (CWD). Because the
124 spatial distribution of CWD is highly heterogeneous, only little quantitative data about its long-term
125 decay dynamics are available for European Alpine forests. Decay models in Europe have, therefore,
126 rarely been parameterised using empirically derived decay constants. In the field, the different
127 stages of CWD decomposition are often described by so-called decay classes (as defined by Hunter,
128 1990) through a visual assessment of the wood status (Lombardi et al. 2013). In a previous study,
129 Petrillo et al. (2015) demonstrated that the Hunter classification is particularly suitable for
130 describing changes in the physical-chemical characteristics of European larch (*Larix decidua* Mill.)
131 and Norway spruce (*Picea abies* (L.) Karst.) deadwood in alpine environments. The physical-
132 chemical properties of deadwood changed distinctly during decay and correlated well with the 5
133 decay classes. Furthermore, no substantial differences between spruce and larch decay patterns were
134 found, although the wood chemistry of the living trees differed slightly between these two species
135 [\(significant differences were found in the cellulose content, with 45.1% for spruce and 39.4% for](#)
136 [larch; these differences were, however, negligible already in decay class 1; Petrillo et al., 2015\).](#)

137 European larch and spruce are widespread in the Alps. Although C-stocks in soils are substantial
138 (e.g. Johnston et al., 2004), CWD is a non-negligible C reservoir in subalpine forests (Sandström et
139 al., 2007). Consequently, it is thus very important to know which time scales are involved in CWD
140 decay. Jebrane et al. (2014) showed that Scots pine is more decay resistant than European larch,
141 which suggests that the decay rate of pine is lower. Some species of larch are, however, considered
142 economically valuable due to their hard, heavy and decay-resistant wood (Parker, 1993), which
143 implies that residence time of larch CWD should be longer.

144 The aim of our work was to find out; i) which time scales are involved in CWD decay of *P. abies*
145 and *L. decidua* in the Alps and ii) how these time scales correlate with the five-decay class system.

146 We hypothesised that the CWD decay of these coniferous trees is relatively slow (due to, e.g. the
147 nutrient availability for macro and micro-organisms being unfavourable),

Markus Egli 16.10.2015 08:10
Gelöscht: and comparable to soil organic matter
compounds that are not easily degradable

148

150 **2 Materials and methods**

151

152 **2.1 Site description**

153 The study area is located in the north-eastern Italian Alps, in Val di Sole and Val di Rabbi (Fig. 1;
154 Table 1). The climate of the valleys ranges from temperate to alpine (above the timberline), the
155 mean annual temperature from 8.2 °C at the valley floor to about 0 °C at 2400 m a.s.l., and the
156 mean annual precipitation from approximately 800 to 1300 mm (Sboarina and Cescatti, 2004). The
157 geological substrate is paragneiss debris in all sites. The soil units are Cambisols, Umbrisols and
158 Podzols (WRB: IUSS working group, 2014). The soil properties at each site could be taken from a
159 previous study (Egli et al., 2006). The timberline is close to 2000 – 2200 m a.s.l., with the forests
160 dominated by Norway spruce and, at the highest altitudes, by European larch.

161

162 **2.2 Sampling protocol**

163 Norway spruce and European larch CWD was sampled at eight sites ranging in altitude from 1200
164 m a.s.l. to 2000 m a.s.l. In spring and summer 2013, wood cores from living trees and cross sections
165 of CWD were taken from all sites. At each site, 5 or 6 living trees were sampled in two directions
166 per each tree at 130 cm height (breast height) using an incremental corer (0.5 cm in diameter;
167 Suunto, Finland). The wood cores were wrapped in paper and transported to the laboratory, where
168 they were air dried, fixed onto a flat wooden support and sanded in order to obtain a smooth surface
169 for tree-ring measurements. Before sampling, each CWD was first classified relative to the decay
170 stage. The classification was done *in situ* using the five-class classification system of Hunter (1990)

173 (Table 2), which is based on visual, geometric and tactile features and considers the
174 presence/absence of twigs and bark, the shape of the log section, and the deadwood structure. [To](#)
175 [sample CWD, a circular area of 50 m radius was explored at each plot. In total, 177 CWD samples](#)
176 [were collected \(46 of them were dated\). At the highest sites, the forests consisted predominantly of](#)
177 [larch trees. In addition, one lower site on the south-facing slope \(S7\) also had a predominantly larch](#)
178 [forest. All other sites are spruce dominated.](#) Samples were taken randomly either using a manual
179 saw or, in more advanced stages of decay, simply by hand. If necessary, they were wrapped up with
180 tape to preserve their structure during transport to the laboratory, where they were air dried and
181 sanded. For CWD in more advanced decay stages (decay class 4 and 5), a 25 cm x 30 cm bag was
182 filled. The samples were then oven-dried at 50 °C, but not sanded. [To establish a master](#)
183 [\(dendro\)chronology](#), 83 wood cores were taken from living trees, 29 from larch and 54 from spruce.
184 [Two cores were taken from each tree. In three cases, one of the two cores taken from the same tree](#)
185 [was too damaged to be measured \(one at N02, one at S06 and one at S07\). Furthermore, two](#)
186 [outliers were excluded from the master chronology, namely one measured core at S07 and one at](#)
187 [S08. CWD was dated using 46](#) cross sections from deadwood (18 from larch and 28 from spruce).

188

189 2.3 Dendrochronological dating

190 At each site, the 10 or 12 wood cores taken from living trees were used to build a reference (master)
191 ring-width chronology for each species. Tree rings were first counted and then measured using the
192 LINTAB tree-ring-width measurement device (RINNTECH e.K., Heidelberg, Germany), coupled
193 together with a stereomicroscope (Leica, Germany). The two ring-width measurements from the
194 same tree were first cross-checked and then incorporated into a single average master chronology
195 for each species and for each site. [To maximise the common signal, one outlier per species was](#)
196 [excluded from the relative master chronology. These two tree-ring measurements exhibited a poor](#)
197 [correlation with the other, probably due to an elevated number of missing rings.](#) The statistical
198 software TSAP-win™ (Time Series Analysis Program, RINNTECH e.K., Heidelberg, Germany)

Marta Petrillo 13.10.2015 16:18

Gelöscht: In total

Markus Egli 16.10.2015 08:30

Gelöscht: summarize

201 was used to calculate the *Gleichläufigkeit*, GLK (Kaennel and Schweingruber, 1995), i.e. the
202 agreement between two ring-width series. The correlations among all the ring-width series of living
203 trees and CWD were statistically assessed using the software COFECHA (Holmes et al., 1986).
204 [EPS \(expressed population signal\) was calculated using the statistic software R.](#)

205 The deadwood cross-sections were measured from the most external ring to the pith, along three or
206 four different radial directions. The individual CWD series (i.e. floating chronologies) were
207 matched to the master chronology of the corresponding species. We visually and statistically
208 checked the deadwood series using the GLK to obtain the highest value with the master chronology
209 and to date the year of death of the tree from which the deadwood originated.

210

211 **[2.4 Radiocarbon dating](#)**

212 [The CWD of the decay classes 4 and 5 was too degraded to be dated through tree-ring analysis as](#)
213 [their wood structure was too altered and the tree rings were no longer visible. In such cases, the](#)
214 [outermost part of the CWD was sampled and ¹⁴C-dated \(Fig. 2-A and 2-B\). We selected a small](#)
215 [fragment of 1 to 2 cm³ in volume from the outermost part assumed to have contained the last- tree](#)
216 [rings produced before the tree died \(Fig. 2-C and 2-D\). This small fragment was gently cleaned](#)
217 [with a brush to remove any non-woody elements, such as particles of soil or vegetation like moss.](#)

218 [The organic samples were cleaned using an acid-alkali-acid \(AAA\) treatment. The samples were](#)
219 [then heated under vacuum in quartz tubes with CuO \(oxygen source\) to remove any absorbed CO₂](#)
220 [in the CuO. The tubes were evacuated, sealed and heated in the oven at 900 °C to obtain CO₂. The](#)
221 [CO₂ of the combusted sample was mixed with H₂ \(1:2.5\) and catalytically reduced over iron powder](#)
222 [at 535 °C to elemental carbon \(graphite\). After reduction, the mixture was pressed into a target so](#)
223 [that carbon ratios could be measured by Accelerator Mass Spectrometry \(AMS\) using the 0.2 MV](#)
224 [radiocarbon dating facility \(MICADAS\) of the Laboratory of Ion Beam Physics at the Swiss](#)
225 [Federal Institute of Technology of Zurich \(ETHZ\).](#)

226 [The calendar ages were obtained using the OxCal 4.2 calibration program \(Bronk Ramsey, 2001,](#)
227 [2009\) based on the IntCal 13 calibration curve and for modern samples the Bomb 13NH1 curve](#)
228 [\(Reimer et al., 2013; Hua et al., 2013\) was used. Several samples \(before 1950 AD\) had a widely](#)
229 [calibrated age range. For these samples, we used the age range with the highest probability of](#)
230 [confining the time elapsed since death very strictly.](#)

231

232 **2.5 Determining the cellulose and lignin**

233 To obtain α -cellulose (Boettger et al., 2007), 10 mg of powdered wood were weighed in Teflon
234 pockets for chemical and thermal treatments. [All wood \(sapwood and heartwood\) was](#)
235 [homogenised prior to chemical analysis. We decided to use this procedure, because it was not](#)
236 [possible to distinguish between sapwood and heartwood for the most decayed stages.](#) Samples were
237 first washed in a 5% NaOH solution at 60 °C for two hours and then for an additional two hours
238 with [fresh 5 % NaOH solution \(the NaOH solution was discarded each time\)](#), before finally being
239 rinsed three times using boiling distilled water (see also Petrillo et al., 2015). The samples were
240 then washed in a 7 % NaClO₂ solution at 60 °C for 30 hours, changing the solution at least every 10
241 hours and then rinsed three times with boiling distilled water. The pockets were dried in the oven at
242 50 °C and the cellulose content was determined as the difference between the initial weight and
243 dried samples. The so-called Klason lignin (lignin insoluble in strong acid; Dence and Lin, 1992)
244 was determined gravimetrically after a sequential extraction in which 0.2 g of each sample was
245 washed three times with 5 ml of distilled water at 80 °C. After each washing, the samples were
246 centrifuged for 10 min at 4500 rpm, dried in the oven at 80 °C and washed three times with 5 ml of
247 ethanol. They were then centrifuged again (10 min. at 4500 rpm) and the supernatant was
248 discarded. After being dried at 60 °C in the oven, 60 mg of each sample were treated with 0.6 ml
249 72 % H₂SO₄ in a warm (30 °C) bath for one hour, and then, after adding 16.8 ml of distilled water,
250 in an autoclave at 120 °C for one hour. Subsequently, the samples were filtered and the filtrate used
251 to determine of the acid-soluble lignin. The insoluble lignin was dried in the oven at 105 °C and

Markus Egli 6.10.2015 16:55

Gelöscht: 4

Markus Egli 1.2.2016 15:45

Gelöscht: a

254 determined as the difference between the dry and initial weight.

255 [In total, the cellulose and lignin content was measured for 177 CWD samples.](#)

256

257 **2.6 Estimating decomposition rate constants on the basis of density loss**

258 In a previous investigation (Petrillo et al., 2015), the density of the deadwood samples was
259 measured. To estimate the decay constants, the average densities in class 1 and in class 5 (the
260 earliest and latest decay stages) were used and the single-negative exponential model of Jenny et al.
261 (1949) applied (see eq. 1). Equation 1 was then solved for the decay constant k according to
262 equation 2:

$$263 \quad k = \frac{-\ln\left(\frac{x_t}{x_0}\right)}{t} \quad (2)$$

264 where x_t is the density of each deadwood sample at a given time (i.e. the estimated time elapsed
265 since death), and x_0 the initial density ([0.45 g cm⁻³ for *Picea abies* L. Karst](#) and [0.59 g cm⁻³ for *Larix*
266 *decidua* L.](#)).

267 [The density of all CWD samples was then compared to the related ages to derive the overall
268 decomposition rates. A similar procedure was applied to cellulose and lignin to derive compound-
269 specific decomposition rates of CWD.](#)

270 [Calculating mean residence time in decay classes from a single time point sample, rather than using
271 longitudinal long-term data, tends to overestimate residence time due to a higher probability of
272 inclusion of slow-decaying trees \(Kruys et al., 2002\). Consequently, snapshot sampling may
273 overestimate the age and mean residence time of CWD. Thus, the decay rate could be
274 underestimated. Calculating the overall CWD decay rates by using density values along a
275 chronosequence risks, therefore, that a certain amount of error is introduced. This bias can be
276 corrected using the proposed approach of Kruys et al. \(2002\). The mean residence time of CWD in
277 a particular decay class is](#)

Markus Egli 6.10.2015 16:55

Gelöscht: -

... [2]

$$E_m = \frac{\sum_{i=1}^N b_{mi}}{N} \quad (3)$$

Unknown
Formatiert: Schriftart:Times, 12 pt

where b_{mi} is the residence time of tree i in a specific decay class m and N = trees present during the time period. The estimator of E_m is:

$$\hat{E}_m = \frac{\sum_{i=1}^{n_m} \frac{b_{mi}}{c} \frac{1}{T}}{\sum_{i=1}^{n_{tot}} \frac{1}{c l_i} \frac{1}{T}} = \frac{n_m}{\sum_{i=1}^{n_{tot}} \frac{1}{l_i}} \quad (4)$$

with c = proportion of the logs existing at time point t , n_m = number of trees from which samples were taken in decay class m , n_{tot} is the total number of sampled trees and l_i is the total residence time of each tree across all decay classes. The expressions cb_{mi}/T and cl_i/T are the probabilities of including units b_{mi} and l_i in the sample, respectively. According to Krueys et al. (2002) it was assumed that all wood samples were taken half-way through their residence time in that class. The parameter l_i was calculated as age_i/x ; age_i is the measured time since death of tree i and

$$x = \frac{r_m}{2} + \sum_{k=1}^{m-1} r_k \quad (5)$$

Unknown
Formatiert: Schriftart:Times, 12 pt

where x is the sum of proportions, r_k , assigned to classes preceding the class of tree i plus 50% of the proportion assigned to tree i 's class, $r_m \cdot \hat{E}_m$ can be calculated iteratively for the different classes. Convergence occurred after 5-10 iterations.

3 Results

3.1 Living chronologies

Two master chronologies for spruce and larch were obtained extending over 164 and 141 years, respectively (Fig. 3). The spruce chronology ranged from 1848 to 2012 AD (Fig. 3) and the larch chronology from 1871 to 2012 AD. The tree-ring widths of the same species correlated well among

305 each other with a high GLK. [When considering each individual plot, the GLK values were highest](#)
306 [at S09 \(larch\), with a mean GLK of 0.70 \(\$\pm\$ 0.1\) and lowest at N01 \(spruce\) with a mean GLK of](#)
307 [0.58 \(\$\pm\$ 0.13\).](#) The series inter-correlation coefficients obtained using COFECHA were 0.535 for
308 spruce and 0.641 for larch (Pearson correlation, all series above 0.3281 were significant; 99%
309 confidence interval). [The EPS values \(Fig. 3B\) were in most cases above \(or close\) to 0.85 \(a](#)
310 [threshold value for noise-free chronology; Wigley et al., 1984\), except for *Picea abies* for the](#)
311 [period of 1870 – 1910 AD.](#) Spruce and larch, however, had quite different growth patterns [with a](#)
312 [Pearson correlation coefficient of 0.274 when comparing the two species-specific master](#)
313 [chronologies.](#) The spruce chronology indicated that the trees grew homogeneously throughout the
314 whole observation period, while larch seemed to be more sensitive to climate with marked high and
315 low growth periods (positive and negative pointer years, Fig. 3). Furthermore, the negative and
316 positive pointer years were not synchronous in the larch and spruce master chronologies. [Even](#)
317 [though more larch trees were sampled at south-facing sites and at higher elevation, we can exclude](#)
318 [a bias due to an unbalanced sample distribution. Within the same homogenous climatic region](#)
319 [similar growth pattern are found \(Carrer and Urbinati, 2006\). The differences between the two](#)
320 [master chronologies are rather influenced by the different phenology of the two species. Larch is a](#)
321 [deciduous tree, with a deep rooting system, while spruce is evergreen and has a shallow root](#)
322 [system. Furthermore, the larch master chronology is cyclically influenced by outbreaks of the larch](#)
323 [bud defoliator *Zeiraphera diniana* Gn. that result in the abrupt occurrence of extremely narrow tree](#)
324 [rings which are not observed for spruce \(Esper, 2007\).](#) The growth pattern of some trees, however,
325 differed considerably from that of the master chronology, possibly due to the specific growth
326 conditions of the individual trees, e.g. if their growth was very suppressed because of competition.
327 Such outliers were excluded from the chronologies.

328

329 3.2 Age of coarse woody debris (CWD)

Markus Egli 27.1.2016 09:36

Gelöscht: The two master chronologies is described in Table 4.

332 Most of the samples of the decay classes 1 – 3 could be dendrochronologically dated, but those of
333 decay classes 4 and 5 had to be radiocarbon dated because of the poorly preserved tree rings (Tables
334 [3](#) and [4](#)). In the first three decay classes, the CWD ages of spruce and larch seem to be in a similar
335 range. The values vary from 1 to 54 years. Interestingly, the average age of CWD does not seem to
336 increase from class 1 to 3. The average age was around 10 – 20 years for all decay classes assuming
337 a relatively fast decay. In decay classes 4 and 5, the average and maximum ages of CWD were
338 usually higher for larch than for spruce. In decay class 4, spruce CWD has an average of about 42
339 years (median 43 years; Fig. 4) and larch CWD an average of 87 years (median 45 years). In decay
340 class 5, the average age of spruce CWD increases to 77 years and the age of larch CWD to 210
341 years. This shows that larch wood, particularly in the decay classes 4 and 5, is much more resistant
342 to rotting than spruce. [Several CWDs had an age of around 200 years. The calibration of
343 radiocarbon dates for such ages is, however, complicated by the so-called plateaus that give rise to a
344 relatively wide range of calibrated ages. To minimise the array of possibilities, we used the age
345 range with the highest probability for confining the time elapsed since death more strictly.](#)

346

347 **3.3 Relations between year since death, decay class and physical-chemical properties of** 348 **deadwood**

349 The physico-chemical data for the CWD ($n = 177$) are given in Petrillo et al. (2015) [and Tables 3](#)
350 [and 4](#) so that the density and the cellulose and lignin contents could be plotted as a function of the
351 decay class and age of the CWD (Fig. 5). Since the relationship between the age of the CWD and
352 physical-chemical characteristics was rather stochastic for the decay classes 1 – 3, they were
353 grouped and their average was used for further analysis. The decrease in density and cellulose
354 concentrations and the simultaneous increase in lignin definitely proceed faster for the spruce CWD
355 than for the larch CWD (Fig. 5). An exponential function best describes the trends in the cellulose
356 and lignin concentrations with time.

Markus Egli 27.1.2016 09:39

Gelöscht: 5

Markus Egli 27.1.2016 09:39

Gelöscht: 6

359 [The stage-based matrix model of Kruys et al. \(2002\)](#) was applied to calculate the k -values (Table 5)
360 as a function of tree species and decay stage (summed decay classes). Using the classical
361 [chronosequence approach](#), the decay rate constants per year (y^{-1}) were, furthermore, calculated for
362 each dated sample [based on the density loss of spruce and larch CWD \(Table 6\)](#). For spruce, we
363 obtained an average value of 0.018 (y^{-1}) and for larch 0.012 (y^{-1} ; cf. Table 6). The k -values were
364 non-normally distributed. Using the Kruskal-Wallis statistical test, we assessed the effects of the
365 factors elevation, exposition, [MAT \(mean annual temperature\)](#), [MAP \(mean annual precipitation\)](#),
366 species and decay class on the k -values. None of [these parameters](#) significantly influenced the decay
367 rate constant. Nonetheless, the range of k -values on south-facing plots seem to be slightly higher
368 than those on the north-facing plots, which suggests the decomposition rates are faster on south-
369 exposed slopes (Fig. 6). In addition, the k -values were estimated by comparing the CWD density
370 with their age and by plotting an exponential regression curve (not shown). This approach resulted
371 in lower k -values: 0.012 y^{-1} for spruce and 0.005 y^{-1} for larch. The mean residence time and half-
372 lives are summarised in Table 7. [The differences in mean residence time and rate constants between](#)
373 [Kruys' model and the more classical approach \(chronosequence\) using equation 1 are small \(Table](#)
374 [6\)](#). Kruys' model gave slightly higher decay constants for Norway spruces (0.022 y^{-1}) and the same
375 [values \(0.018 \$y^{-1}\$ \) for European larch \(variant a\) in Table 6\)](#).

376

377

378 **4 Discussion**

379 Although the five-decay class system is well suited to describe changes in the physical and
380 chemical properties of deadwood (Lombardi et al., 2008), no real differences in the age of the CWD
381 classes 1 – 3 could be found. The CWD in decay class 4 and 5 was, however, clearly older. This
382 implies that the first three decay classes are not clearly related to deadwood age. Similarly,
383 Lombardi et al. (2013) found no relationship between the age of CWD and the chemical properties

Markus Egli 22.1.2016 06:19

Gelösch: Based on the density loss of spruce and larch CWD

Marta Petrillo 13.10.2015 16:25

Gelösch:

Markus Egli 22.1.2016 06:09

Gelösch:)

Markus Egli 28.1.2016 09:45

Gelösch: them

389 of decay classes 1 – 3. The main explanation for this unexpected finding is that there is probably a
390 time lag between the death of a standing tree and its contact with the soil (Kueppers et al., 2004;
391 Zielonka, 2006; Lombardi et al., 2013). Standing dead trees, i.e. snags, can remain upright for
392 several years and decay much more slowly than fallen dead trees (Yatskov et al., 2003). Such an
393 effect overshadows a clear age trend in decay. If the species-specific fall rates were known the
394 decay rates could be better assessed. Unfortunately, the fall rates of snags of the studied tree species
395 are unknown at the investigated sites. To our knowledge, no data about fall rates of snags of either
396 species, *Picea abies* (Norway spruce) and *Larix decidua* Mill. (European larch), are available. In
397 this respect, the data situation in North America is much better. A good overview is given for
398 example in Hilger et al. (2012) and Dixon (2015). According to Hilger et al. (2012), Engelmann
399 spruce (*Picea engelmannii* Parry ex Engelm.) and subalpine larch (*Larix lyallii* Parl.) have similar
400 snag fall rates. Due to morphological, ecological and physiological similarities, we have to assume
401 (but we finally cannot prove it) that Norway spruce and European larch should exhibit a similar
402 reaction to Engelmann spruce and subalpine larch. As a consequence, no particular difference in the
403 fall rate between European larch and Norway spruce is to be expected. Therefore, difference in the
404 decay rates between European larch and Norway spruce are hypothesised not to be due to different
405 fall rates.

406 Angers et al. (2012), however, observed that the wood density in snags in boreal forests already
407 decreases after a few years. Decay rates they calculated are comparable to those in our study. The
408 density loss in standing dead trees could be due to the activity of cerambycid larvae, while the
409 activity of the wood decomposers, mainly fungi, was impeded in snags due to the lack of moisture.
410 The discrepancy between the macromorphology of deadwood (and consequently decay class) and
411 the age of deadwood seems therefore to be related to the individual tree death history. Shortly after
412 tree death, in fact, the wood is rapidly colonized by fungi (Zielonka, 2006). The CWD in classes 4
413 and 5 showed a relation to deadwood age that seems to be species-specific since larch CWD is older
414 than spruce in both classes. With respect to the CWD ages in our study, classes 1 to 3 appear to be a

416 single group, while classes 4 and 5 are different. The oldest sample (larch CWD) was about 244
417 years old – a surprisingly old age for wood lying on the forest floor (i.e. not buried). Spruce CWD
418 in decay classes 4 and 5 seem to be significantly younger than larch CWD. Few empirical
419 assessments of time since the death of a tree have been made in Europe. Krüger et al. (2014) used
420 both dendrochronology and radiocarbon dating to assess the time since death of Norway spruce in
421 Bavarian forests. They estimated a total residence time of 61 to 62 years for this species. Our values
422 are slightly lower. One major problem in determining the age using ^{14}C are the sometimes large age
423 ranges obtained after calibration (due to plateaus) for samples having a pre-bomb age. We used the
424 ranges with the highest probabilities (varying from 50 to 82.2%; Table 4; commonly the age 1- σ
425 range, i.e. 68% is considered) for CWD dated to the time period before the bomb peak.
426 Consequently, this procedure introduces an uncertainty. According to Krüger et al. (2014),
427 radiocarbon analysis and dendrochronological cross-dating revealed a similar year of tree death for
428 samples having a post-bomb age. The results of Krüger et al. (2014) suggest that both methods are
429 suitable for the age determination of CWD. In Atlantic Canada, Campbell and Laroque (2007)
430 found an age of 56 to 84 years (depending on the investigated sites) in the latest decay stage (decay
431 class 5; Black spruce and Balsam fir). Lombardi et al. (2008) estimated stumps of beech and silver
432 fir in decay class 3 to be 55 and 59 years which is close to our findings.

433 The decay rates reflect the determined ages of the CWD; and spruce therefore had a higher decay
434 rate constant than larch. Consequently, decay rates are species specific due to, among others things,
435 the initial differences in the physical-chemical properties of the wood of the living trees and in
436 environmental factors. Larch has, for example, a higher density (cf. Fig. 5) and a lower nutrient
437 content than spruce (Petrillo et al., 2015). Shorohova et al. (2014) also found that decay rates can
438 strongly vary among tree species. The decay rate (i.e. 0.032 y^{-1}) they found for spruce was slightly
439 higher than that in our study (Fig. 6). The variability of the decay rates given in the literature may
440 also arise from using different mathematical models or different methods to determine wood density
441 or the age of the CWD. According to Hale and Pastor (1998), the decay rates of oak and maple logs

442 (in a temperate forest) varied between 0.00 and 0.18 y^{-1} (their dating of the logs, however, based on
443 estimates). The decay rates of tree species in a Mediterranean area (Australia; Brown et al., 1996)
444 varied in the range of 0.05 up to 0.22 y^{-1} , while in a cool-continental climate (Alban and Pastor,
445 1993), decay rates were 0.042 and 0.055 for red and jack pine, respectively, 0.07 for spruce and
446 0.08 y^{-1} for aspen. Fukusawa et al. (2014) estimated decay rates by using the annual input of CWD
447 divided by the CWD accumulation, and obtained a value of 0.036 y^{-1} . With the chronosequence
448 approach, however, the rates were in the order of 0.020 – 0.023 y^{-1} .

449 [The determined decay rates for spruce and larch in our investigation seem to be very low \(Table 6\).](#)
450 [As pointed out by Kruys et al. \(2002\), the chronosequence approach, and thus the snapshot](#)
451 [sampling, may overestimate the CWD age and consequently residence time. Thus, the decay rate](#)
452 [may be underestimated. It seems, however, that this error is not overwhelmingly distinct in our case](#)
453 [or even absent. The approach according to Kruys et al. \(2002\) and variant a\) in Table 6 gave similar](#)
454 [results. The regression approach \(variant b\) in Table 6\) probably slightly underestimated the decay](#)
455 [rates.](#)

456 Using mass losses instead of density losses to estimate the decay rates may result in higher values,
457 because the losses for fragmentation are added to the mineralisation losses (Yin, 1999). This might
458 explain why our decay rate constants were lower than those in some other studies (Rock et al.,
459 2008). Moreover, the decay rates are sensitive, at a regional scale, to climatic conditions such as
460 temperature and precipitation (Shorohova et al., 2014), although the decay rates for a mean annual
461 temperature of 0 to 10 °C are, however, quite similar, and rates below 0.04 y^{-1} are often reported
462 (Mackensen et al., 2003). Soil temperature was found to be the main explanatory variable for
463 differences in the decay rates of standard wood, such as aspen and pine (Risch et al., 2013).
464 Although the data are too limited to draw clear conclusion, some of the differences in the decay
465 rates we observed are likely to be due to environmental factors. On south-facing sites, for instance,
466 we found that the decay rates were slightly, but not significantly, higher than those on north-facing
467 sites (Fig. 6), which is comparable to Shorohova's et al. (2014) observations.

468 [Although our measured \$k\$ values are very low, they fit reasonably well to those of the recent](#)
469 [compilation of Russel et al. \(2015\). For environments having a mean annual temperature of < 10](#)
470 [°C, the decay rate constants are usually < 0.1 \(median value is 0.027 for such sites\). The](#)
471 [compilation of Russel et al. \(2015\), however, only considers two sites having *Picea abies* \(\$k\$ values](#)
472 [= 0.044 and 0.027; Krankina et al., 1999; Næsset, 1999\) and none for larch. Together with our](#)
473 [results, a residence time of about 20 – 90 years for *Picea abies* in subalpine \(boreal\) climates might](#)
474 [be suggested.](#)

475 The concentrations of cellulose and lignin in the CWD are given as a function of time in Figure 5.
476 Due to the faster decomposition of cellulose, lignin is relatively enriched. Lignin, however, also
477 decomposes with time. To unravel the decay behaviour of these compounds, a multiple-exponential
478 model was applied (Means et al., 1985; Mackensen et al., 2003), with the general form:

$$479 \quad x = x_1 e^{-k_1 t} + x_2 e^{-k_2 t} \dots + x_n e^{-k_n t} \quad (6)$$

480 where x_i is the density or mass of deadwood at a given time and $x_{1..n}$ are partitioned parameters. The
481 portioning of cellulose and lignin is solved graphically using their mass per unit volume over time,
482 and fitting them to an exponential regression curve. From this, the half-life of cellulose or lignin in
483 the CWD could be calculated:

$$484 \quad t_{1/2} = \frac{\ln(1/2)}{-k} \quad (7)$$

485 where $t_{1/2}$ is the half-life and k is the decay constant (obtained from the exponential regression
486 curve). Using the k -values in Figure 7, the following half-lives were obtained:

- 487 a) for cellulose: 21 years (spruce) and 50 years (larch)
- 488 b) for lignin: 91 years (spruce) and 481 years (larch)

489 Means et al. (1985) were able to derive k -values for cellulose values of 0.0109 to 0.0117 y^{-1} for
490 Douglas-fir logs (in a cool to temperate climate), although age determination (or estimation) was
491 done differently. This would give rise to half-lives in the range of 59 to 64 years. With k -values in
492 the range of 0.0039 to 0.0045 y^{-1} (Means et al., 1985), the half-life of lignin would be in the range

Markus Egli 20.1.2016 21:46
Gelöscht: 3

Markus Egli 20.1.2016 21:46
Gelöscht: 4

495 of 154 – 178 years. In this specific case, the overall decay rates were between 0.006 and 0.0073 y⁻¹.
496 Although cellulose is relatively easily degradable by (micro)organisms, it may persist astonishingly
497 long in larch trees (several decades). Lignin may have a half-life of more than hundred years. These
498 half-lives may be shorter if the decay is related to mass losses and not to density.

499 In decay classes 4 and 5, the CWD starts to become more and more part of the soil. [The further fate](#)
500 [of CWD compounds strongly depends on their interaction with the mineral soil.](#) The introduced
501 organic matter into soils can be either further degraded or stabilised to a certain extent. The
502 persistence of organic matter in soils is largely due to complex interactions between the organic
503 matter and its environment, such as the interdependence of compound chemistry, reactive mineral
504 surfaces, climate, water availability, soil acidity, soil redox state and the presence of potential
505 degraders in the immediate micro-environment (Schmidt et al., 2011). [Together with physical](#)
506 [protection, organo-mineral interactions are generally thought to be the main mechanism for SOM](#)
507 [stabilisation \(e.g., Nierop et al., 2002; Kleber et al., 2005; Marschner et al., 2008\).](#)

Markus Egli 10.10.2015 22:04

Gelösch: How do such data now compare to organic matter in soils?

Markus Egli 10.10.2015 22:10

Gelösch: (SOM)

Markus Egli 10.10.2015 22:08

Gelösch: Most organic components in soils have a mean residence time of just a few years up to about 50 – 60 years. Fire-derived organic matter may have a residence time of decades to as much as 300 years (Schmidt et al., 2011). The measured mean ages of the CWD in decay class 4 or 5, the CWD residence time and the decay rates of cellulose and lignin are very similar to those of soil organic matter compounds (and even fire-derived SOM). The rate of CWD decay processes does not apparently differ very greatly from the degradation of organic matter in soils.

510 5 Conclusions

511 The first 3 decay classes do not seem to reflect the age of the CWD, but they are relevant for the
512 description of its decay stage. [The time lag between the death of a standing tree and its contact with](#)
513 [the soil overshadows a clear age trend.](#) Taking classes 1 – 3 as one group and relating them to the
514 decay classes 4 and 5, a time trend with increasing decay stage can then be detected. This time trend
515 also closely correlates to the wood density, and the cellulose and lignin content. The oldest CWD
516 age of a larch tree reached the considerable age of 244 years. [We used a chronosequence approach](#)
517 [and applied several calculation techniques to estimate the overall decay rate constants of European](#)
518 [larch and Norway spruce. The stage-based matrix model of Kruijs et al. \(2002\) that corrects for](#)
519 [sampling bias was in good agreement with the often-used single-negative-exponential model. The](#)

Markus Egli 23.1.2016 09:20

Gelösch: these

Markus Egli 23.1.2016 09:39

Gelösch: The ages of the CWD in decay classes 4 and 5 even correspond to relatively 'stable' SOM (e.g. fire-derived organic matter). Consequently, the turn-over time of CWD organic matter is in a similar range to that of SOM.

Markus Egli 23.1.2016 09:22

Formatiert: Hervorhebung

541 [regression approach probably underestimated slightly the decay rates.](#) The decay rate constant for
 542 spruce [seems to be in the range of 0,018 to 0,022 \(y⁻¹\)](#) and for larch [it is about 0.012 \(y⁻¹\)](#). The rates
 543 seemed to be slightly higher on south-facing sites (although this was not statistically significant).
 544 [An effect of the altitude on the decay rates was, however, not discernible.](#) Using the dating
 545 approach (dendrochronology and ¹⁴C-dating), the behaviour of cellulose and lignin as a function of
 546 time could be assessed. Our findings demonstrate that lignin in larch may persist particularly long,
 547 with a mean residence time of > 100 years. [This indicates that turnover rates of CWD organic](#)
 548 [matter are even in a comparable range to that of SOM.](#)
 549 More empirical data is, however, needed to ascertain our findings. [A major issue is that fall rates](#)
 550 [between European Larch and Norway spruce could not be compared. Furthermore, the preparation](#)
 551 [and precise dating of CWD is time-consuming, cost intensive and in some cases also difficult](#)
 552 [\(particularly samples with a pre-bomb age in decay classes 4 and 5\).](#) Since CWD represents an
 553 important forest carbon pool, improving the informative potential of the decay classes (including
 554 the dating of the CWD) would contribute to sustainable forest management and make carbon
 555 accounting easier.

Markus Egli 23.1.2016 09:33
 Gelösch: was
 Markus Egli 23.1.2016 09:33
 Gelösch: 012
 Markus Egli 23.1.2016 09:33
 Gelösch: 018
 Markus Egli 23.1.2016 09:34
 Gelösch: 0.005 to
 Markus Egli 23.1.2016 09:34
 Gelösch: , depending on the approach

Markus Egli 23.1.2016 09:37
 Gelösch: T
 Markus Egli 24.1.2016 18:40
 Gelösch:
 Markus Egli 23.1.2016 19:52
 Gelösch: and

558 Acknowledgements

559 This study is part of the DecAlp DACH project no. 205321L_141186. J. Ascher has been funded by
 560 the Fonds zur Förderung der wissenschaftlichen Forschung (FWF) Austria (Project I989-B16). We
 561 are indebted to Dr. Fabio Angeli of the ‘Ufficio distrettuale forestale di Malé’ and his team of
 562 foresters for their support in the field. We would also like to thank Leonora Di Gesualdo for her
 563 help in the sampling wood cores and Michelle Kovacic for preparing samples for radiocarbon
 564 dating. We are grateful to Silvia Dingwall for the English corrections. [Furthermore, we gratefully](#)

573 | [acknowledge the constructive suggestions of the referees \(one unknown reviewer and J. Schöngart\)](#)
574 | [and two readers \(T. Kahl and V.-A. Angers\) to improve the manuscript.](#)

575

576

577 **References**

578 Alban, D. H. and Pastor, J.: Decomposition of aspen, spruce, and pine boles on two sites in
579 Minnesota, *Can. J. Forest Res.*, 23, 1744-1749, doi: 10.1139/x93-220, 1993.

580 Angers, V. A., Drapeau, P., and Bergeron, Y.: Mineralization rates and factors influencing snag
581 decay in four North American boreal tree species, *Can. J. Forest Res.*, 42, 157–166, doi:
582 10.1139/X11-167, 2012.

583 Barbati, A., Marchetti, M., Chirici, G., and Corona, P.: European forest types and forest Europe
584 SFM indicators: Tools for monitoring progress on forest biodiversity conservation, *Forest Ecol.*
585 *Manag.*, 321, 145–157, doi: 10.1016/j.foresco.2013.07.004, 2014.

586 Bond-Lamberty, B. and Gower, S.T.: Decomposition and fragmentation of coarse woody debris:
587 Re-visiting a boreal black spruce chronosequence, *Ecosystems* 11, 831–840, doi:
588 10.1007/s10021-008-9163-y, 2008.

589 Boettger, T., Haupt, M., Knoller, K., Weise, S. M., Waterhouse, J.S., Rinne, K. T., Loader, N. J.,
590 Sonninen, E., Jungner, H., Masson-Delmotte, V., Stievenard, M., Guillemin, M. T., Pierre, M.,
591 Pazdur, A., Leuenberger, M., Filot, M., Saurer, M., Reynolds, C. E., Helle, G., and Schleser, G.
592 H.: Wood cellulose preparation method and mass spectrometric analysis of delta C-13, delta O-
593 18, and nonexchangeable delta H-2 values in cellulose, sugar and starch: An interlaboratory
594 comparison, *Analytical Chemistry*, 79, 4603–4612, doi: 10.1021/ac0700023, 2007.

595 Bronk Ramsey, C.: Bayesian analysis of radiocarbon dates, *Radiocarbon* 51, 337-360, 2009.

596 Brown, S., Mo, J., McPherson, J. K., and Bell, D. T.: Decomposition of woody debris in Western
597 Australian forests, *Can. J. Forest R.*, 26, 954-966, doi: 10.1139/x26-105, 1996.

598 Campbell, L. J. and Laroque, C. P.: Decay progression and classification in two old-growth forests
599 in Atlantic Canada, *Forest Ecol. Manag.*, 238, 293–301, doi: 10.1016/j.foreco.2006.10.027,
600 2007.

601 [Carrer, M. and Urbinati, C.: Long-term change in the sensitivity of tree-ring growth to climate](#)
602 [forcing in *Larix decidua*, *New Phytol.*, 170, 861 – 872, doi: 10.1111/j.1469-8137.2006.01703.x,](#)
603 [2006.](#)

604 Dence, C. W. and Lin, S. Y.: Introduction. In: Lin, S.Y., Dence, C.W. (eds). *Methods in lignin*
605 *chemistry*, Springer, Heidelberg pp. 1–19, 1992.

606 Di Cosmo, L., Gasparini, P., Paletto, A., and Nocetti, M.: Deadwood basic density values for
607 national-level carbon stock estimates in Italy, *Forest Ecol. Manag.*, 295, 51–58, doi:
608 10.1016/j.foreco.2013.01.010, 2013.

609 [Dixon, G.E.: Essentials FVS: A User’s Guide to the Forest Vegetation Simulator. United States](#)
610 [Department of Agriculture, US Forest Service, Fort Collins CO, 2015.](#)

611 Egli, M., Mirabella, A., Sartori, G., Zanelli, R., and Bischof, S.: Effect of north and south exposure
612 on weathering rates and clay mineral formation in Alpine soils, *Catena* 67, 155–174, doi:
613 10.1016/j.catena.2006.02.010, 2006.

614 [Esper, J., Büntgen U., Frank, D.C., Nievergelt, D., and Liebhold, A.: 1200 years of regular](#)
615 [outbreaks in alpine insects, *Proc. R. Soc. London Ser. B.* 274, 671 – 679, doi:](#)
616 [10.1098/rspb.2006.0191, 2007.](#)

617 Hale, C. M. and Pastor, J.: Nitrogen content, decay rates, and decompositional dynamics of hollow
618 versus solid hardwood logs in hardwood forests of Minnesota, U.S.A., *Can. J. Forest R.*, 28,
619 1276-1285, doi: 10.1139/cjfr-28-9-1276, 1998.

620 Harmon, M. E., Franklin, J. F., Swanson, F. J., Sollins, P., Gregory, S. V., Lattin, J. D., Anderson,
621 N. H., Cline, S. P., Aumen, N. G., Sedell, J. R., Lienkaemper, G. W., Cromack, K., and
622 Cummins, K. W.: Ecology of coarse woody debris in temperate ecosystems, *Adv. Ecol. Res.*, 15,
623 133–302, doi: 10.1016/S0065-2504(08)60121-X, 1986.

- 624 [Hilger, A.B., Shaw, C.-H., Metsaranta, J.M., and Kurz, W.A.: Estimation of snag carbon transfer](#)
625 [rates by ecozone and lead species for forests in Canada, *Ecol. Appl.*, 22, 2078–2090,](#)
626 [doi:10.1890/11-2277.1, 2012.](#)
- 627 Holmes, R. L., Adams, R. K., and Fritts, H. C.: Tree-ring chronologies of Western North America:
628 California, Eastern Oregon and Northern Great Basin, with procedures used in chronology
629 development work, including users manuals for computer programs COFECHA and ARSTAN,
630 Chronology series VI, Laboratory of Tree-Ring Research, University of Arizona, Tucson, 182
631 pp., 1986.
- 632 [Hua, Q., Barbetti, M., and Rakowski, A. Z.: Atmospheric radiocarbon for the period 1950-2010,](#)
633 [Radiocarbon, 55, 2059-2072, doi: 10.2458/azu_js_rc.v55i2.16177, 2013.](#)
- 634 Hunter, M. L.: Wildlife, forests and forestry: principles of managing forests for biological diversity.
635 Prentice Hall, Englewood Cliffs (NJ), USA, 370 pp., 1990.
- 636 [IUSS Working Group WRB: World Reference Base for Soil Resources 2014. International soil](#)
637 [classification system for naming soils and creating legends for soil maps, World Soil Resources](#)
638 [Reports No. 106. FAO, Rome, 2014.](#)
- 639 Jebrane, M., Pockrandt, M., and Terziev, N.: Natural durability of selected larch and Scots pine
640 heartwoods in laboratory and field tests, *Int. Biodeter. Biodegr.*, 91, 88–96, doi:
641 10.1016/j.ibiod.2014.03.018, 2014.
- 642 Jenny, H., Gessel, S. P., and Bingham, F. T.: Comparative study of decomposition of organic matter
643 in temperate and tropical regions, *Soil Sci.*, 68, 419–432, doi: 10.1097/00010694-200606001-
644 00017, 1949.
- 645 Kaennel, M. and Schweingruber, F. H.: Multilingual glossary of dendrochronology. Terms and
646 definitions in English, German, French, Spanish, Italian, Portuguese and Russian. Swiss Federal
647 Institute for Snow, Forest and Landscape Research. Haupt, Bern, Switzerland, 1995.

Markus Egli 27.1.2016 08:49

Gelöscht: IPCC: Good practice guidance for land use, land use change and forestry, Hayama: Institute for Global Environmental Strategies, 2003.

- 651 [Kleber, M., Mikutta, R., Torn, M. S., and Jahn, R.: Poorly crystalline mineral phases protect organic](#)
652 [matter in acid subsoil horizons, Eur. J. Soil Sci., 56, 717-725, doi:10.1111/j.1365-](#)
653 [2389.2005.00706.x, 2005](#)
- 654 [Krankina, O. N., Harmon, M. E., and Griazkin, A. V.: Nutrient stores and dynamics of woody](#)
655 [detritus in a boreal forest: modeling potential implications at the stand level, Can. J. For. Res.,](#)
656 [29, 20-32, doi: 10.1139/x98-162, 1999.](#)
- 657 [Kruys, N., Jonsson, B. G., and Ståhl, G.: A stage-based matrix model for decay-class dynamics of](#)
658 [woody debris, Ecol. Applic., 12, 773-781, 2002.](#)
- 659 Kueppers, L. M., Southon, J., Baer, P., and Harte, J.: Dead wood biomass and turnover time,
660 measured by radiocarbon, along a subalpine elevation gradient, *Oecologia*, 141, 641–651, doi:
661 10.1007/s00442-004-1689-x, 2004.
- 662 Krüger, I., Muhr, J., Hartl-Meier, C., Schulz, C., and Borken, W.: Age determination of coarse
663 woody debris with radiocarbon analysis and dendrochronological cross-dating, *Eur. J. Forest*
664 *Res.*, 133, 931–939, doi: 10.1007/s10342-014-0810-x, 2014.
- 665 Lombardi, F., Cherubini, P., Lasserre, B., Tognetti, R., and Marchetti, M.: Tree rings used to assess
666 time since death of deadwood of different decay classes in beech and silver fir forests in the
667 central Apennines (Molise, Italy), *Can. J. For. Res.*, 38, 821–833, doi: 10.1139/X07-195, 2008.
- 668 Lombardi, F., Cherubini, P., Tognetti, R., Coccozza, C., Lasserre, B., and Marchetti, M.:
669 Investigating biochemical processes to assess deadwood decay of beech and silver fir in
670 Mediterranean mountain forests, *Ann. For. Sci.*, 70, 101–111, doi: 10.1007/s13595-012-0230-3,
671 2013.
- 672 Mackensen, J., Bauhus, J., and Webber, E.: Decomposition rates of coarse woody debris – A review
673 with particular emphasis on Australian tree species, *Aust. J. Bot.*, 51, 27– 37, doi:
674 10.1071/BT02014, 2003.
- 675 [Marschner, B., Brodowski, S., Dreves, A., Gleixner, G., Gude, A., Grootes, P. M., Hamer, U.,](#)
676 [Heim, A., Jandl, G., Ji, R., Kaiser, K., Kalbitz, K., Kramer, C., Leinweber, P., Rethemeyer, J.,](#)

677 [Schäffer, A., Schmidt, M. W. I., Schwark, L., and Wiesenberg, G. L. B.: How relevant is](#)
678 [recalcitrance for the stabilization of organic matter in soils? J. Plant Nutr. Soil Sci. 171, 91-110,](#)
679 [doi: 10.1002/jpln.200700049, 2008.](#)

680 Means, J. E., Cromack, K., [and](#) MacMillan, P. C.: Comparison of decomposition models using
681 wood density of Douglas-fir logs, *Can. J. Forest Res.*, 15, 1092-1098, doi: 10.1139/x85-178, 1985.

682 [Minderman, G.: Addition, decomposition and accumulation of organic matter in forests, J. Ecol.,](#)
683 [56, 355-362, doi: 10.2307/2258238, 1968.](#)

684 Moroni, T. M., Hagemann, U., and Beilman, D. W.: Dead wood is buried and preserved in a
685 Labrador boreal forest, *Ecosystems*, 13, 452–458, doi: 10.1007/s10021-010-9331-8, 2010.

686 Müller, J. and Büttler, R.: A review of habitat thresholds for dead wood: a baseline for management
687 recommendations in European forests, *Eur. J. Forest Res.*, 129, 981–992, doi: 10.1007/s10342-
688 010-0400-5, 2010.

689 Müller-Using, S. and Bartsch, N.: Decay dynamic of coarse and fine woody debris of a beech
690 (*Fagus sylvatica* L.) forest in Central Germany, *Eur. J. Forest Res.*, 128, 287–296, doi:
691 10.1007/s10342-009-0264-8, 2009.

692 [Næsset, E.: Decomposition rate constants of Picea abies logs in southeastern Norway, Can. J. For.](#)
693 [Res. 29:372-381, doi: 10.1139/x99-005, 1999.](#)

694 [Nierop, K. G. J., Jansen, B. and Verstraten, J. A.: Dissolved organic matter, aluminium and iron](#)
695 [interactions: precipitation induced by metal/carbon ratio, pH and competition, Sci. Total](#)
696 [Environ., 300, 201-211, 10.1016/S0048-9697\(02\)00254-1, 2002.](#)

697 Olson, J. S.: Energy storage and the balance of producers and decomposers in ecological systems,
698 *Ecology*, 44, 322–331, doi: 10.2307/1932179, 1963.

699 Parker, W. H.: Larix. Flora of North America Editorial Committee (eds.): Flora of North America
700 North of Mexico, Vol. 2, Oxford University Press, pp. 366-368, 1993.

701 Palviainen, M., Finén, L., Laiho, R., Shorohova, E., Kapitsa, E., and Vanha-Majamaa, I.:
702 Phosphorus and base cation accumulation and release patterns in decomposing Scots pine,

703 Norway spruce and silver birch stumps, *Forest Ecol. Manag.*, 260, 1478–1489, doi:
704 10.1016/j.foreco.2010.07.046, 2010.

705 Petrillo, M., Cherubini, P., Sartori, G., Abiven, S., Ascher, J., Bertoldi, D., Camin, F., Barbero, A.,
706 Larcher, R., and Egli, M.: Decomposition of Norway spruce and European larch coarse woody
707 debris (CWD) in relation to different elevation and exposition in an Alpine setting, *iForest*, doi:
708 10.3832/ifor1591-008, 2015.

709 Reimer, P. J., Bard, E., Bayliss, A., Beck, J. W., Blackwell, P. G., Bronk Ramsey, C., Buck, C. E.,
710 Cheng, H., Edwards, R. L., Friedrich, M., Grootes, P. M., Guilderson, T. P., Hafliðason, H.,
711 Hajdas, I., Hatté, C., Heaton, T. J., Hoffmann, D. L., Hogg, A. G., Hughen, K. A., Kaiser, K. F.,
712 Kromer, B., Manning, S. W., Nui, M., Reimer, R. W., Richards, D. A., Scott, E. M., Southon, J.
713 R., Staff, R. A., Turney, C., and van der Plicht, J.: IntCal13 and Marine13 radiocarbon age
714 calibration curves 0–50,000 years cal BP, *Radiocarbon*, 55, 1869–1887, doi:
715 10.2458/azu_js_rc.55.16947, 2013.

716 Risch, C. A., Jurgensen, F. M., Page-Dumroese, D. S., and Schütz, M.: Initial turnover rates of two
717 standard wood substrates following land-use change in subalpine ecosystems in the Swiss Alps,
718 *Can. J. Forest Res.*, 43, 901–910, doi: 10.1139/cjfr-2013-0109, 2013.

719 Rock, J., Badeck, F., and Harmon, M. E.: Estimating decomposition rate constants for European
720 tree species from literature sources, *Eur. J. Forest Res.*, 127, 301–313, doi: 10.1007/s10342-008-
721 0206-x, 2008.

722 [Russel, M. B., Fraver, S. F., Aakala, T., Gove, J. H., Woodall, C. W., D'Amato, A. W., and Ducey,
723 M. J.: Quantifying carbon stores and decomposition in dead wood: A review, *Forest Ecol.
724 Manag.*, 350, 107–128, doi.org/10.1016/j.foreco.2015.04.033, 2015.](#)

725 Sboarina, C. and Cescatti, A.: Il clima del Trentino – Distribuzione spaziale delle principali variabili
726 climatiche. Report 33, Centro di Ecologia Alpina Monte Bondone, Trento, Italy. 2004.

727 Schmidt, M. W. I., Torn, M. S., Abiven, S., Dittmar, T., Guggenberger, G., Janssens, I. A., Kleber,
728 M., Kögel-Knabner, I., Lehmann, J., Manning, D. A. C., Nannipieri, P., Rasse, D. P., Weiner, S.,

729 and Trumbore, S. E.: Persistence of soil organic matter as an ecosystem property, *Nature*, 478,
730 49–56, doi: 10.1038/nature10386, 2011.

731 Shorohova, E. and Kapitsa, E.: Influence of the substrate and ecosystem attributes on the
732 decomposition rates of coarse woody debris in European boreal forests, *Forest Ecol. Manag.*,
733 315, 173–184, doi: 10.1016/j.foreco.2013.12.025, 2014.

734 [Wider R.K. and Lang G.E.: A critique of the analytical methods used in examining decomposition](#)
735 [data obtained from litter bags, *Ecology*, 63, 1636-1642, doi: 10.2307/1940104, 1982.](#)

736 [Wigley, T. M. L., Briffa, K. R., and Jones, P. D.: On the average value of correlated time series,](#)
737 [with applications in dendroclimatology and hydrometeorology, *J. Climate Appl. Meteor.*, 23,](#)
738 [201-213, \[http://dx.doi.org/10.1175/1520-0450\\(1984\\)023<0201:OTAVOC>2.0.CO;2\]\(http://dx.doi.org/10.1175/1520-0450\(1984\)023<0201:OTAVOC>2.0.CO;2\), 1984.](#)

739 WRB, IUSS Working Group: World Reference Base for Soil Resources (2nd ed.). World Soil
740 Resources Reports No. 103, First update, FAO, Rome, 2007.

741 Yatskov, M., Harmon, M. E., and Krankina, O. N.: A chronosequence of wood decomposition in
742 the boreal forests of Russia, *Can. J. Forest Res.*, 33, 1211–1226, doi: 10.1139/X03-033, 2003.

743 Yin, X.: The decay of forest woody debris: numerical modelling and implications based on some
744 300 data cases from North America, *Oecologia*, 121, 81–98, doi: 10.1007/s004420050909, 1999.

745 Zielonka, T.: When does dead wood turn into a substrate for spruce replacement?, *J. Veg. Sci.*, 17,
746 739–746, doi: 10.1111/j.1654-1103.2006.tb02497.x, 2006.

747 **Figure captions**

748

749 **Fig. 1.** Location of the study area with the major vegetation units and investigation sites. Data
750 source: Museo delle Scienze (Trento), CORINE Landcover (Joint Research Center of the European
751 Union) and scilands GmbH. [The site label N indicates north-facing sites and S south-facing sites.](#)

752

753 **Fig. 2.** Cross section of (A) spruce deadwood in the field (site N03) and (B) larch deadwood (site
754 S07). Examples (C and D) of deadwood fragments classified as decay class 4 dated using
755 radiocarbon (outermost part of the wood piece).

756

757 **Fig. 3.** Master chronologies for spruce and larch to cross-date the deadwood (A) and the expressed
758 [population signal \(EPS\) for segments of constant periods \(B\). A noise-free chronology is achieved](#)
759 [with an EPS > 0.85 \(dashed line; Wigley et al., 1984\)](#)

760

761 **Fig. 4.** Box plots of the larch and spruce deadwood age as a function of decay class.

762

763 **Fig. 5.** Relation between the age of spruce and larch CWD and density (A), cellulose % (B) and
764 lignin % (C). The decay classes 1 – 3 were grouped together due to their similar age (Fig. 4).

765

766 **Fig. 6.** Calculated decay rate constants (k) as a function of tree species and site exposure.

767

768 **Fig. 7.** Empirically determined exponential regression curves (principle of multiple-exponential
769 model) for partitioning the decay behaviour of cellulose (A) and lignin (B).

Markus Egli 7.2.2016 12:02

Gelöscht: (continuous line)

Markus Egli 7.2.2016 12:02

Gelöscht: (dashed line)

Table 1. Characteristics of the study sites.

Plot ID	Elevation (m a.s.l.)	Aspect (°N)	Slope (°)	MAP ^a (mm yr ⁻¹)	Parent material	Dominating tree species	Land use	Soil classification (WRB) (Egli et al., 2006)
<i>North-facing sites</i>								
N01	1180	340	31	950	Paragneiss debris	<i>Picea abies</i>	Natural forest (ecological forestry)	Chromi-Episkeletic Cambisol (Dystric)
N02	1390	0	28	1000	Paragneiss debris	<i>Picea abies</i>	Natural forest (ecological forestry)	Chromi-Episkeletic Cambisol (Dystric)
N03	1620	0	29	1060	Paragneiss debris	<i>Picea abies</i>	Natural forest (ecological forestry)	Chromi-Endoskeletal Cambisol (Dystric)
N04	1930	20	12	1180	Paragneiss debris, moraine material	<i>Larix decidua</i>	Originally used as pasture	Episkeletic Podzol
<i>South-facing sites</i>								
S06	1185	160	31	950	Paragneiss debris	<i>Picea abies</i>	Ex-coppice, natural forest (ecological forestry)	Episkeleti-Endoleptic Cambisol (Chromi- Dystric)
S07	1400	145	33	1000	Paragneiss debris	<i>Larix decidua</i>	Natural forest (ecological forestry)	Dystri-Endoskeletal Cambisol
S08	1660	210	33	1060	Paragneiss debris	<i>Picea abies</i>	Natural forest (ecological forestry)	Skeletal Umbrisol
S09	1995	160	25	1180	Paragneiss debris	<i>Larix decidua</i>	Ex pasture, natural forest	Skeletal Umbrisol

^aMAP = mean annual precipitation (Sboarina and Cescatti, 2004)

Table 2. The five decay-class system of log decomposition (according to Hunter, 1990).

Log features	Decay classes				
	1	2	3	4	5
Bark	Intact	Partially absent	Absent	Absent	Absent
Twigs	Present	Partially absent or absent	Absent	Absent	Absent
Shape of radial section	Round	Round	Round	Oval	Very oval
Colour	Original	Original	Faded in the external part	Reddish brown or faded	Reddish or faded
Texture of wood	Intact	Intact	Soft outer layer, intact inner part	Small pieces, soft	Powdery or fibrous, very soft
Contact with soil	Log elevated on what remains of branches	Log in contact with soil	Log in contact with soil	Log in contact with soil	Log in contact with soil and partially buried

Markus Egli 27.1.2016 08:49
Gelöscht: Table 3. Wood density values for Norway spruce and European larch given by the IPCC (2003) and in previous investigations (Petrillo et al., 2015). - [...](#) [3]

Markus Egli 27.1.2016 08:53
Gelöscht: Table 4. Description of the two master chronologies. The inter-series correlation is obtained with the software COFECHA. - [...](#) [4]

Table 3. [Typical properties](#) and ages of Norway spruce and European larch CWD in class 1 – 3. Ages were obtained mostly from dendrochronological measurements and a few* from ¹⁴C-dating (Detail of ¹⁴C-dating cf. Appendix A).

Plot	Tree species	Decay class	Density (g/cm ³)	Cellulose (%)	Lignin (%)	Year of death	CWD age
N03	Norway spruce	1	0.32	34.8	22.4	2009	4
S08	Norway spruce	1	0.40	31.3	33.2	1992	21
N01	Norway spruce	1	0.45	43.4	22.0	1988	25
N02	Norway spruce	1	0.44	41.4	23.3	1969	44
N03	Norway spruce	2	0.39	41.4	35.9	2006	7
N03	Norway spruce	2	0.44	28.4	25.0	2004	9
N03	Norway spruce	2	0.39	27.8	25.9	2004	9
S08	Norway spruce	2	0.36	38.2	23.1	2003	10
N02	Norway spruce	2	0.11	39.2	14.2	1996	17
N03	Norway spruce	2	0.43	40.8	21.5	1993	20
N01	Norway spruce	2	0.39	28.2	24.3	1970	43
N02	Norway spruce	2	0.67	27.8	25.9	1959	54
N03	Norway spruce	3	0.48	43.6	23.4	2012	1
N03	Norway spruce	3	0.38	36.6	24.2	2005	8
N03	Norway spruce	3	0.39	37.4	10.7	2005	8
N01	Norway spruce	3	0.30	22.3	35.1	1979	34
N02	Norway spruce	3	0.48	33.5	24.6	1970	43
S09	European larch	1	0.60	37.2	21.1	2010	3
N04	European larch	1	0.59	44.2	16.9	1973	40
S07*	European larch	1	0.31	21.2	39.6	2007	6
S07	European larch	2	0.58	20.2	38.0	2010	3
S09	European larch	2	0.53	37.3	31.0	2000	13
S07*	European larch	2	0.30	30.4	40.6	2003	10
S07*	European larch	3	0.27	6.7	63.0	2004	9
S09*	European larch	3	0.60	4.2	40.8	1973	40
S09*	European larch	3	0.33	23.3	58.4	1968	45

Markus Egli 27.1.2016 09:33

Gelöscht: 5

Markus Egli 12.10.2015 08:03

Gelöscht:

Markus Egli 27.1.2016 08:09

Formatierte Tabelle

Table 4. Typical properties and radiocarbon data of the deadwood samples (decay classes 4 and 5) as a function of site and tree species.

UZH number	ETH number	Sample code	Site	Tree species	Decay class	Density (g/cm ³)	Cellulose (%)	Lignin (%)	¹⁴ C-age years	±1σ	δ ¹³ C ‰	±δ ¹³ C ‰	cal AD 2 σ	Average age ¹⁾ years	cal AD ²⁾ years (probability)	Average age ¹⁾ years
UZ 6210	ETH-56612	37A	N4	European larch	4	0.26	34.6	27.9	-2931	29	-23.6	1	1962-1974	45		
UZ 6211	ETH-56613	34A	N4	European larch	4	0.40	1.0	47.2	170	31	-24.4	1	1658-1950	209	1720-1819 (50%)	244
UZ 6213	ETH-56615	50A	S9	European larch	4	0.15	2.3	43.2	-659	31	-26.5	1	1957-2003	33		
UZ 6219	ETH-56853	95A	S7	European larch	4	0.21	16.7	38.4	-860	25	-25.5	1	1957-1998	36		
UZ 6227	ETH-56861	209	S7	European larch	4	0.33	0.0	47.2	-2545	25	-25.3	1	1962-1976	44		
UZ 6228	ETH-56862	214	S7	European larch	4	0.33	16.5	28.7	60	25	-23.9	1	1695-1919	206	1867-1919 (53.3%)	120
UZ 6212	ETH-56614	45B	S9	European larch	5	0.34	25.9	29.4	183	31	-27	1	1650-1950	213	1726-1815 (51.5%)	243
UZ 6224	ETH-56858	202	S9	European larch	5	0.25	0.0	58.1	140	25	-28.9	1	1669-1944	207	1798-1944 (52.6 %)	142
UZ 6264	ETH-60747	33	N4	European larch	5	0.28	0.0	49.4	185	25	-29.4	1	1656-1950	220	1728-1810 (54.7 %)	244
UZ 6214	ETH-56616	69A	S6	Norway spruce	4	0.13	23.3	25.2	-1331	30	-31.3	1	1958-1989	40		
UZ 6215	ETH-56849	72A	S6	Norway spruce	4	0.37	0.0	51.7	-2120	25	-23.6	1	1961-1980	43		
UZ 6216	ETH-56850	84A	S8	Norway spruce	4	0.16	32.4	29.1	-4080	25	-27	1	1966-1967	47		
UZ 6220	ETH-56854	97B	N2	Norway spruce	4	0.26	28.1	36.1	-3720	25	-31.4	1	1967-1968	46		
UZ 6221	ETH-56855	98A	N3	Norway spruce	4	0.26	0.0	67.0	-290	25	-19.7	1	1955-2009	31		
UZ 6226	ETH-56860	206	N3	Norway spruce	4	0.29	28.5	20.5	-2150	25	-23.9	1	1962-1979	43		
UZ 6217	ETH-56851	87A	S8	Norway spruce	5	0.13	1.5	66.8	130	25	-26.1	1	1677-1940	205	1800-1940 (59.5%)	143
UZ 6218	ETH-56852	89A	S8	Norway spruce	5	0.27	17.6	27.7	-615	25	-24.6	1	1956-2004	33		
UZ 6222	ETH-56856	106A	N1	Norway spruce	5	0.19	4.3	68.5	-1665	25	-25.2	1	1959-1984	42		
UZ 6223	ETH-56857	197	N1	Norway spruce	5	0.19	0.0	40.2	-4595	25	-24	1	1963-1965	49		
UZ 6225	ETH-56859	205	N2	Norway spruce	5	0.19	0.0	44.3	10	25	-22.9	1	1699-1916	206	1879-1916 (82.2 %)	116

¹⁾ Calculated as the mean value between the maximum and minimum age (2 σ). For this range of years (2 σ), associated probabilities are summed to 95.4 %

²⁾ For samples dated in the period before the bomb peak and giving a wide calibrated age range, the age is also calculated for the most important calibrated time-range associated to the highest, corresponding probability (in brackets)

Table 5. Mean residence time and decay constants calculated using the stage-based matrix model of Krays et al. (2002)

Decay class	No of samples		Mean residence time		Decay constant (yr ⁻¹)	
	<u>Spruce</u>	<u>Larch</u>	<u>Spruce</u>	<u>Larch</u>	<u>Spruce</u>	<u>Larch</u>
<u>1</u>	<u>4</u>	<u>3</u>	<u>77</u>	<u>80</u>	<u>0.013</u>	<u>0.012</u>
<u>1+2</u>	<u>8</u>	<u>3</u>	<u>36</u>	<u>47</u>	<u>0.027</u>	<u>0.021</u>
<u>1+2+3</u>	<u>4</u>	<u>3</u>	<u>28</u>	<u>67</u>	<u>0.036</u>	<u>0.015</u>
<u>1+2+3+4</u>	<u>6</u>	<u>6</u>	<u>63</u>	<u>116</u>	<u>0.016</u>	<u>0.009</u>
<u>1+2+3+4+5</u>	<u>5</u>	<u>3</u>	<u>63</u>	<u>254</u>	<u>0.016</u>	<u>0.004</u>

Table 6. CWD decay parameters based on a) equation 1, b) the regression approach and c) stage-based matrix model of Kruijs et al. (2002).

	Average decay constant k (yr^{-1})	Residence time* (yr)	Half-life* (yr)
a)			
Norway spruce	0.018	56	39
European larch	0.012	83	58
b)			
Norway spruce	0.012	84	58
European larch	0.005	222	154
c)			
Norway spruce	0.022	45	32
European larch	0.012	83	58

*calculated from the average decay constant

Markus Egli 20.1.2016 21:53

Gelöscht: 7

Markus Egli 20.1.2016 21:51

Gelöscht: CWD decay parameters based on equation 1 (a) and on the regression approach (b)

Markus Egli 20.1.2016 21:51

Gelöscht: -

Decay constant k -

Residence time -

Half-life -

a)

... [5]

Appendix A. Radiocarbon data of the deadwood samples of the decay classes 1 – 3.

UZH number	ETH number	Sample code	Site	Tree species	Decay class	C14 age	$\pm 1\sigma$	$\delta^{13}\text{C}$ ‰	$\pm\delta^{13}\text{C}$ ‰	Cal AD $\pm 1\sigma$	Average age years
UZ-6258	ETH-60741	L_10_c1_1	S7	European Larch	1	-435	25	-25.7	1	2006-2009	6
UZ-6260	ETH-60743	S07_dc2_92	S7	European Larch	2	-590	25	-26.9	1	2002-2004	10
UZ-6261	ETH-60744	S07_dc3_96	S7	European Larch	3	-545	25	-26.4	1	2003-2005	9
UZ-6262	ETH-60745	S09_cl3_46	S9	European Larch	3	-2865	25	-29.3	1	1973-1974	40
UZ-6263	ETH-60746	S09_cl3_48	S9	European Larch	3	-2775	25	-23.8	1	1962-1974	45

¹⁾ Calculated as the mean value between the maximum and minimum age (1σ). For this range of years (1σ), associated probabilities summed to 68.2 %