

**BGD**

12, 14869–14910, 2015

**Microbial  
methanogenesis in  
the sulfate-reducing  
zone**

J. Maltby et al.

# Microbial methanogenesis in the sulfate-reducing zone of surface sediments traversing the Peruvian margin

J. Maltby<sup>1</sup>, S. Sommer<sup>1</sup>, A. W. Dale<sup>1</sup>, and T. Treude<sup>1,a</sup>

<sup>1</sup>GEOMAR Helmholtz Centre for Ocean Research Kiel, Department of Marine Biogeochemistry, Wischhofstr. 1–3, 24148 Kiel, Germany

<sup>a</sup>present address: Department of Earth, Planetary & Space Sciences and Atmospheric & Oceanic Sciences, University of California, Los Angeles (UCLA), CA, USA

Received: 7 August 2015 – Accepted: 15 August 2015 – Published: 9 September 2015

Correspondence to: J. Maltby (jmalby@geomar.de) and T. Treude (ttreude@g.ucla.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Abstract

We studied the concurrence of methanogenesis and sulfate reduction in surface sediments (0–25 cm below sea floor, cmbsf) at six stations (70, 145, 253, 407, 770 and 1024 m) along the Peruvian margin (12° S). This oceanographic region is characterized by high carbon export to the seafloor, creating an extensive oxygen minimum zone (OMZ) on the shelf, both factors that could favor surface methanogenesis. Sediments sampled along the depth transect traversed areas of anoxic and oxic conditions in the bottom-near water. Net methane production (batch incubations) and sulfate reduction (<sup>35</sup>S-sulfate radiotracer incubation) were determined in the upper 0–25 cmbsf of multicorer cores from all stations, while deep hydrogenotrophic methanogenesis (> 30 cmbsf, <sup>14</sup>C-bicarbonate radiotracer incubation) was determined in two gravity cores at selected sites (78 and 407 m). Furthermore, stimulation (methanol addition) and inhibition (molybdate addition) experiments were carried out to investigate the relationship between sulfate reduction and methanogenesis.

Highest rates of methanogenesis and sulfate reduction in the surface sediments, integrated over 0–25 cmbsf, were observed on the shelf (70–253 m, 0.06–0.1 and 0.5–4.7 mmol m<sup>-2</sup> d<sup>-1</sup>, respectively), while lowest rates were discovered at the deepest site (1024 m, 0.03 and 0.2 mmol m<sup>-2</sup> d<sup>-1</sup>, respectively). The addition of methanol resulted in significantly higher surface methanogenesis activity, suggesting that the process was mostly based on non-competitive substrates, i.e., substrates not used by sulfate reducers. In the deeper sediment horizons, where competition was probably relieved due to the decline of sulfate, the usage of competitive substrates was confirmed by the detection of hydrogenotrophic activity in the sulfate-depleted zone at the shallow shelf station (70 m).

Surface methanogenesis appeared to be correlated to the availability of labile organic matter (C / N ratio) and organic carbon degradation (DIC production), both of which support the supply of methanogenic substrates. A negative correlation of methanogenesis rates with dissolved oxygen in the bottom-near water was not obvious, however,

**BGD**

12, 14869–14910, 2015

## Microbial methanogenesis in the sulfate-reducing zone

J. Maltby et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



anoxic conditions within the OMZ might be advantageous for methanogenic organisms at the sediment–water interface.

Our results revealed a high relevance of surface methanogenesis on the shelf, where the ratio between surface to deep (below sulfate penetration) methanogenic activity ranged between 0.13 and  $10^5$ . In addition, methane concentration profiles indicate a partial release of surface methane into the water column as well as a partial consumption of methane by anaerobic methane oxidation (AOM) in the surface sediment. The present study suggests that surface methanogenesis might play a greater role in benthic methane budgeting than previously thought, especially for fueling AOM above the sulfate-methane transition zone.

## 1 Introduction

Microbial methanogenesis represents the terminal step of organic matter degradation in marine sediments (Jørgensen, 2006). The process is entirely restricted to a small group of prokaryotes within the domain of the Archaea (Thauer, 1998). Methanogens produce methane from a narrow spectrum of substrates, primarily carbon dioxide ( $\text{CO}_2$ ) and hydrogen ( $\text{H}_2$ ) (hydrogenotrophic pathway), as well as acetate (acetoclastic pathway) (Zinder, 1993). In addition, methanol or methylated compounds such as methylamine are utilized (methylotrophic pathway) (Oremland and Polcin, 1982; Buckley et al., 2008; Zinder, 1993; King et al., 1983). Substrates for methanogenesis are produced during depolymerization and fermentation of organic macromolecules (e.g., sugars, vitamins, amino acids) to smaller monomeric products (Jørgensen, 2006; Schink and Zeikus, 1982; Neill et al., 1978; Donnelly and Dagley, 1980).

Acetoclastic and hydrogenotrophic methanogenesis are predominantly found in deeper sediment zones below sulfate penetration, owing to the competition with sulfate reducers that outcompete methanogens for  $\text{H}_2$  and acetate due to their higher substrate affinity (Oremland and Polcin, 1982; Jørgensen 2006). Furthermore,  $\text{CO}_2/\text{H}_2$  and acetate are the more abundant substrates in deeper sediments as degradabil-

### Microbial methanogenesis in the sulfate-reducing zone

J. Maltby et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ity of organic matter, and with it the substrate variety and availability, decreases with increasing sediment depth (Jørgensen, 2006).

Methanogens avoid competition with sulfate reducers by the utilization of non-competitive substrates, such as methanol or methylamines (Oremland and Polcin, 1982; King et al., 1983). Facilitated by the usage of such non-competitive substrates, sulfate reduction and methanogenesis were found to co-occur in sulfate-containing salt marsh sediments (Oremland et al., 1982; Buckley et al., 2008; Senior et al., 1982). Concurrent activity of sulfate reduction and methanogenesis in the marine environment has mostly been postulated for organic-rich sediments (Mitterer, 2010; Jørgensen and Parkes, 2010; Treude et al., 2005a, 2009; Hines and Buck, 1982; Crill and Martens, 1986); however, research on magnitude and environmental controls of surface methanogenesis is still sparse (Holmer and Kristensen, 1994; Ferdelman et al., 1997).

In a study from Eckernförde Bay, southwestern Baltic Sea, considerable in- vitro methanogenic activity was observed in samples taken from 5 to 40 cm sediment depth (Treude et al., 2005). Although in- vitro activity was measured in sulfate-free setups, methanogenic activity coincided with zones of in-situ sulfate reduction. The authors concluded a coexistence of the two types of organisms, which could be enabled through either the usage of non-competitive substrates, dormancy of methanogens until phases of sulfate depletion, and/or temporal or spatial heterogeneity in the sediments. Eckernförde Bay sediments feature a high input of organic matter due to a shallow water depth (~ 30 m) and pronounced phytoplankton blooms in spring, summer, and fall (Smetacek, 1985). Furthermore, seasonal hypoxia ( $O_2 < 90 \mu\text{M}$ ) or even anoxia ( $O_2 = 0 \mu\text{M}$ ) occur in the deep layers of the water column caused by stratification and degradation of organic matter (Bange et al., 2011). Oxygen-depleted conditions in the bottom water together with frequent input of fresh organic matter possibly favors methanogenesis in surface sediment by offering reduced conditions and non-competitive substrates. In accordance, methanogenesis activity was observed within

**BGD**

12, 14869–14910, 2015

**Microbial  
methanogenesis in  
the sulfate-reducing  
zone**

J. Maltby et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the sulfate-reducing zone of organic-rich and seasonally hypoxic sediments from the Limfjorden sound, Northern Denmark (Jørgensen and Parkes, 2010; Jørgensen, 1977).

The environmental relevance of surface methanogenesis is hitherto unknown. Its closeness to the sediment–water interface makes it a potential source for methane emissions into the water column, unless the methane is microbially consumed before escaping the sediment (Knittel and Boetius, 2009). Methane escapes the sediment either by diffusion or in the form of gas bubbles, when methane saturation is exceeded (Whiticar, 1978; Wever and Fiedler, 1995; Judd et al., 1997; Dimitrov, 2002). How much of the released methane reaches the atmosphere mainly depends on water depth, as methane is consumed within the water column through microbial oxidation (Reeburgh, 2007; Valentine et al., 2001). Thus, coastal areas have higher methane emission potentials than the open ocean (Bange et al., 1994). Once in the atmosphere, methane acts as a very potent greenhouse gas (IPCC, 2014).

In the present study, we focused on the upwelling region off the Peruvian coast, which is another excellent example of an environment, where both factors that potentially favor surface methanogenesis convene, i.e., high export of organic carbon and low dissolved oxygen concentrations in the bottom water. This upwelling region represents one of the most productive systems in the world oceans, creating one of the most intense oxygen minimum zones (OMZ, Kamykowski and Zentara, 1990; Pennington et al., 2006). Oxygen concentrations in waters impinging on the seafloor are below 20  $\mu\text{M}$  or even reach anoxia. Research on surface methanogenesis in upwelling regions is rare and its potential role in the carbon cycling of the Peruvian OMZ is completely unknown. In a study from the central Chilean upwelling area (87 m water depth, 0.5–6 cmbsf), small production of methane was detected despite high sulfate reduction activity, when offering the non-competitive substrate trimethylamine (Ferdelman et al., 1997). The authors concluded that the prevailing methanogens were competing with sulfate reducers for  $\text{H}_2$  and with acetogens for methylamines, explaining the overall low methanogenesis activity observed (Ferdelman et al., 1997).

**Microbial  
methanogenesis in  
the sulfate-reducing  
zone**

J. Maltby et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



---

**Microbial  
methanogenesis in  
the sulfate-reducing  
zone**

J. Maltby et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Even though the Chilean and Peruvian OMZs are connected, commonly known as OMZ in the eastern South Pacific Ocean (ESP) (Fuenzalida et al., 2009), the core of the ESP-OMZ is centered off Peru with an upper boundary at  $< 100$  m and a vertical distribution to  $> 600$  m vs. a thinner OMZ band off Chile constrained between 100–400 m water depth (Fuenzalida et al., 2009). The anoxic conditions in the water column of the OMZ core together with the high export rates of labile organic carbon to the seafloor (Reimers and Suess, 1983; Dale et al., 2015) provide favorable conditions for methanogenesis activity in surface sediments, thus increasing the potential for benthic methane emissions.

Here, we provide first insights into surface methanogenesis in sediment cores ( $< 30$  cmbsf = centimeters below seafloor) taken along the Peruvian shelf and margin. We hypothesize that methanogenesis coexists with sulfate reduction through the utilization of non-competitive substrates. In addition, we postulate that surface methanogenesis depends on the quantity and quality (= freshness) of organic carbon, and the concentrations of dissolved oxygen in the bottom water. We therefore expect spatial variability of surface methanogenesis along the continental shelf and margin. The observed methanogenic activity will be compared to methane concentrations in the bottom-near water to discuss the potential relevance of surface methanogenesis for methane emissions into the pelagic zone.

## 2 Material and methods

### 2.1 Study site and sediment sampling

Samples were taken during the R.V. *Meteor* cruise M92 between 5 January and 3 February 2013 along a depth transect off the Peruvian coast from the shelf ( $\sim 70$  m) to the continental slope ( $\sim 1000$  m). The transect was located in the central part of the ESP-OMZ (Fuenzalida et al., 2009) at  $12^\circ$  S. Further hydrographic details on the study area can be found elsewhere (Dale et al., 2015).

---

**Microbial  
methanogenesis in  
the sulfate-reducing  
zone**

---

J. Maltby et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Sediment cores for the determination of near-surface methanogenesis were collected at six stations along the depth transect at 70, 145, 253, 407, 770 and 1024 m water depth (Fig. 1), using a multiple corer with a mounted camera (TV-MUC). The MUC held seven cores (length: 60 cm, inner diameter: 10 cm) and covered an area of  $\sim 1 \text{ m}^2$ . If necessary, a second MUC was deployed at the same station, thus sediment cores sometimes originated from different MUC casts. Station numbers were assigned in accordance with Dale et al. (2015). After retrieval, sediment cores were transferred to a  $\sim 9^\circ\text{C}$  cold room and processed the same day.

In addition to the MUC, a gravity corer was deployed at two stations (78 and 407 m) for determining deep methanogenesis. The total core length was 400 and 206 cm, respectively. The gravity corer was equipped with a 260 kg weight and a 5 m steal barrel (diameter: 14 cm). The replaceable core liner (PVC, diameter: 12.5 cm) was housed within the barrel and fixed with a core catcher. After retrieval, sediment cores from the gravity corer were sliced into 1 m sections, capped on both sides, and brought to the cold room ( $4^\circ\text{C}$ ) for further processing. Relevant station details for MUC and gravity cores are summarized in Table 1.

## 2.2 Water column sampling

CTD/Rosette water column casts were conducted at the same station as sediment coring (for details see Table 1). Temperature and oxygen data are taken from Dale et al., 2015.

For the analysis of methane concentrations in the bottom-near water, water was sampled ca. 1.5 m above the seafloor from 10 L Niskin bottles mounted on a rosette water sampler. The collected water was filled bubble-free into 60 mL vials (triplicates), each vial containing 3 pellets of sodium hydroxide ( $\text{NaOH}$ ,  $\sim 0.3 \text{ M vial}^{-1}$ ) to stop microbial activity and force dissolved gas into the headspace. After closing the vials with a butyl rubber stopper and a crimp seal, 10 mL of water was removed with a  $\text{N}_2$ -flushed 10 mL syringe and replaced with  $\text{N}_2$  gas from a second syringe to create a headspace in the

sampling vials. Samples were stored and transported at room temperature until further processing.

In the home laboratory, 100  $\mu\text{L}$  of the headspace volume was injected into a Shimadzu GC-2014 gas chromatograph equipped with a flame ionization detector and a HayeSep-T 100/120 column (length 3 m, diameter: 2 mm). Gases were separated isothermally at 75 °C with helium carrier gas. Methane concentrations were calibrated against methane standards (Scotty gases). The detection limit was 0.1 ppm with a precision of 2 %.

### 2.3 Porewater geochemistry

Porewater sampling for MUC cores has been previously described by Dale et al. (2015). In short, one MUC core per station was subsampled in an argon-filled glove bag, to preserve redox sensitive constituents.

The gravity cores at St. 1 (78 m) and St. 8 (407 m) were subsampled at 10–12 different sediment depths (depending on core length) resulting in depth intervals of 20–33 cm. Before sampling, the plastic core liner was cut open with an electric saw at the specific depths. Porewater was extracted by using anoxic (flushed with argon), wetted rhizons (Rhizosphere Research Products, Seeberg-Elverfeldt et al., 2005).

Sulfate concentrations were determined by ion chromatography (Methrom 761) as described previously by Dale et al. (2015).

For DIC analysis, 1.8 mL of porewater was transferred into a 2 mL glass vial, fixed with 10  $\mu\text{L}$  saturated mercury chloride solution and crimp sealed. Samples were stored at 4 °C until further processing in the home laboratory. DIC concentration was determined as  $\text{CO}_2$  with a multi N/C 2100 analyzer (Analytik Jena) following the manufacturer's instructions. Therefore the sample was acidified with phosphoric acid and the outgassing  $\text{CO}_2$  was measured. The detection limit was 20  $\mu\text{M}$  with a precision of 2–3 %.



## 2.4 Sediment porosity and particulate organic carbon/nitrogen

Methodology and data for porosity, particulate organic carbon (POC) and particulate organic nitrogen (PON) have been previously described by Dale et al. (2015).

In short, wet sediment samples were taken from the porewater MUC core and the gravity cores for determination of porosity from the weight difference of wet and freeze-dried sediment. POC and PON were analyzed with a Carlo–Erba element analyzer (NA 1500). Ratios of POC : PON were calculated by division.

## 2.5 Sediment methane

For sediment methane concentration, one MUC core per station was sliced in 2 cm intervals until 20 cm depth, followed by 5 cm intervals until the end of the core (maximum depth = 48 cm). Gravity cores were subsampled according to the above scheme (see Sect. 2.3). From each sampled sediment layer, 2 cm<sup>3</sup> sediment were transferred into a 15 mL serum glass vial containing 5 mL of NaOH (2.5% *w/w*). The vial was closed with a butyl stopper, crimp sealed and shaken thoroughly to stop microbial activity and to force all methane into the headspace. Vials were stored upside down at room temperature until measurement in the home laboratory.

Sediment methane concentration was determined by injecting 0.1 mL of headspace volume into a Shimadzu GC-2014 gas chromatograph as described under Sect. 2.2.

## 2.6 Net methanogenesis activity in MUC cores

Sediment from MUC cores was used to determine net methanogenesis, which is defined as the sum of total methane production and consumption, including all available methanogenic substrates in the sediment. Net methanogenesis was determined by measuring the linear increase of methane concentration in the headspace of closed incubation vials over time. Therefore, one MUC core per station was sliced in 5 cm intervals, transferring 10 cm<sup>3</sup> of sediment in triplicates into a N<sub>2</sub>-flushed 60 mL serum

---

**Microbial  
methanogenesis in  
the sulfate-reducing  
zone**J. Maltby et al.

---

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

glass vial. The sediment core lengths ranged between 25–48 cm, resulting in maximum 10 depth intervals. Ten ml of anoxic deep water overlying each MUC core was added to the vial and the slurry was mixed under a constant N<sub>2</sub> stream (Hungate, 1950) before sealed with a butyl rubber stopper and crimped. The sediment slurry was repeatedly flushed with N<sub>2</sub> through the stopper to guarantee fully anoxic conditions. The vials were incubated in the dark and at 9 °C, which reflected the average in situ temperature along the depth transect (see Table 1). The first gas chromatographic measurement was done directly after preparation of the vials, by injecting 100 µL of headspace sample into the gas chromatograph. The on-board Hewlett Packard-5890 gas chromatograph was equipped with a flame ionization detector and a HayeSep-T 100/120 column (Length 3 m, diameter: 2 mm). Gases were separated isothermally at 75 °C with helium carrier gas. Methane concentrations were calibrated against methane standards. The detection limit was 1 ppm with a precision of < 5 %. Measurements were done in 2–4 day-intervals over a total incubation time of ~ 2 weeks.

## 2.7 Potential non-competitive and competitive methanogenesis in sediment slurries from MUC cores

Sediment slurry experiments were conducted with sediment from St. 1 (70 m) to examine the interaction between sulfate reduction and methanogenesis, as this station revealed highest microbial activity of sulfate reduction and methanogenesis. On board, the sediment core was sliced in 5 cm intervals. Sediment from the 0–5 cm interval and the 20–25 cm interval was transferred completely into 250 mL glass bottles, which were then closed without headspace with a butyl rubber stopper and screw cap. Until further treatment, sediment was stored at 4 °C on board and later in a 1 °C cold room on shore.

Approximately 6 months after the cruise, sediment slurries from both depth intervals were prepared by mixing 5 mL sediment in a 1 : 1 ratio with artificial, fully marine seawater (Widdel and Bak, 1992) before further manipulations.

In total, three different treatments, each in triplicates, were prepared per depth: (1) sulfate-rich (28 mM), serving as a control, (2) sulfate-rich plus molbydate (22 mM) from

now on referred to as molybdate-treatment, and (3) sulfate-rich plus methanol (10 mM) from now on referred to as methanol-treatment.

Molybdate was used as an enzymatic inhibitor for sulfate reduction (Oremland and Capone, 1988). Methanol is a known non-competitive substrate used by methanogens, but not by sulfate reducers (Oremland and Polcin, 1982), which makes it suitable to examine non-competitive methanogenesis.

The sediment slurries were incubated at 9 °C in the dark for 23 days and headspace concentration of methane was measured repeatedly over time on a gas chromatograph. Therefore, 100 µL of headspace was removed from the gas vials and injected into a Shimadzu gas chromatograph (GC-2014) equipped with a methanizer (inactive), a packed HayeSep-D column and a flame ionization detector. The column temperature was 80 °C and the helium flow was set to 12 mL min<sup>-1</sup>. Methane concentrations were measured against methane standards. The detection limit was 0.1 ppm with a precision of < 5%. Rates were determined from the linear increase of methane concentration over time. Due to differences in the linear increase between the three treatments, rates were determined at two different time points: the first period of incubation includes the starting point (day 0) until day 5, the second period includes day 8 to day 23 (Supplement, Fig. S1).

Student's *t* test (independent, two-tailed,  $\alpha = 0.05$ ) was applied to detect significant differences between the three different treatments.

## 2.8 Gross hydrogenotrophic methanogenesis activity in gravity cores

For the determination of surface to deep methanogenesis activity in gravity cores the radiotracer technique using <sup>14</sup>C-bicarbonate was applied (Jørgensen, 1978). With this method only hydrogenotrophic methanogenesis from CO<sub>2</sub>/H<sub>2</sub> can be determined, which is the expected main pathway in deeper sediment layers.

Sampled sediment depths were according to the above scheme (see Sect. 2.3). Circa 5 cm<sup>3</sup> of sediment was sampled in triplicates into glass syringes and then sealed headspace-free with butyl rubber stoppers. Then, <sup>14</sup>C-bicarbonate-tracer (dis-

**Microbial  
methanogenesis in  
the sulfate-reducing  
zone**

J. Maltby et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)





into 1 cm intervals and transferring each sediment layer into 50 mL plastic centrifuge tubes filled with 20 mL zinc acetate (20% *w/w*). Controls were done in triplicates from different depths. Here, the sediment was first fixed with zinc acetate before adding the tracer. Rates for sulfate reduction were determined using the cold chromium distillation procedure according to Kallmeyer et al. (2004).

The yielded sulfate reduction rates have to be treated with caution, due to long (up to 3 half-life times of  $^{35}\text{S}$ ) and unfrozen storage. Storage of sulfate reduction samples without freezing has recently been shown to result in the re-oxidation of  $^{35}\text{S}$ -sulfides, which results in an underestimation of sulfate reduction rates (Røy et al., 2014). During this reaction, zinc sulfide ( $\text{Zn}^{35}\text{S}$ ) and iron sulfide ( $\text{Fe}^{35}\text{S}$ ) are re-oxidized to sulfate by reactive Fe(III), which originates from the reaction of  $\text{Fe}^{2+}$  with oxygen.  $\text{Fe}^{2+}$  is released during the gradual conversion of FeS to ZnS, which has the lower solubility product. Still, we do trust the relative distribution of activity along depth profiles and consider a potential underestimation of absolute rates.

### 3 Results

#### 3.1 Water column oxygen and methane concentration

Dissolved oxygen in the bottom water was below detection limit from St. 1 (70 m) to St. 8 (407 m), subsequently increasing with water depth to  $53\ \mu\text{M}$  at the deepest site (see Table 1 and Dale et al., 2015). At the shallowest St. 1 (70 m) the water was turbid and smelled of sulfide.

Dissolved methane concentrations in the bottom water were high on the shelf (St. 1–6, 70–253 m) and 10 fold lower at the deeper sites (St. 8–10, 407–1024 m; Table 1). The highest measured methane concentration was detected at St. 6 (253 m,  $\sim 80\ \text{nM}$ ) and lowest concentrations were detected at St. 10 (1024 m,  $\sim 4\ \text{nM}$ ).

**BGD**

12, 14869–14910, 2015

## Microbial methanogenesis in the sulfate-reducing zone

J. Maltby et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 3.2 Sediment core description

A detailed sediment description for the porewater geochemistry cores has been already published in detail by Dale et al. (2015). In short, sediments revealed a grey color with a black surface layer at St. 1 (70 m), a dark olive green color at St. 4–8 (145–407 m), and a green-brown color at St. 9 and 10 (770–1024 m). Sediment texture was soft and fluffy at St. 1–6 (70–253 m), and was less soft at the deeper sites. St. 8 (407 m) revealed a fluffy surface layer followed by a dense clay layer > 2 cmbsf sediment depth. In addition, phosphorite nodules were found at the sediment surface (0–2 cmbsf) of St. 8 (407 m).

Mats of the sulfur oxidizing bacteria *Thioploca* spp. (Gallardo, 1977) were visible at the sediment surface at St. 1–6 (70–253 m), with the densest mat at St. 1 (70 m) continuously decreasing with increasing water depth. Sheaths of *Thioploca* were visible until 20–30 cmbsf at St. 1, 4 and 6 (70–253 m).

Foraminifera could be observed at the sediment surface of St. 8 (407 m), St. 9 (770 m) and St. 10 (1024 m). St. 8 (407 m) showed a thick layer of foraminifera ooze on the sediment surface (0–3 cmbsf) while St. 9 (770 m) and St. 10 (1024 m) showed only scattered foraminifera at the sediment surface (0–5 cmbsf).

Macrofauna (large polychaetes, oligochaetes, ophiuroids) were restricted to the sites below the OMZ at St. 9 (770 m) and St. 10 (1024 m), where deep waters were oxygenated. However, small snails (~ 1 cm) were observed at St. 8 (407 m).

## 3.3 Geochemical parameters in MUC cores

Porewater and solid phase geochemistry of sediments retrieved by the MUC cores are shown in Fig. 2. Surface sediment (0–0.5 cmbsf) POC content increased along the continental shelf from 1.6 wt % at the shallow St. 1 (70 m) to a maximum of 15 wt % at St. 8 (253 m). Surface POC content decreased again with increasing water depth showing the lowest POC content at St. 10 (1024 m, 2 wt %). While POC content showed more or less stable profiles throughout the sediment core at St. 1 (70 m, around 3 wt %),

## Microbial methanogenesis in the sulfate-reducing zone

J. Maltby et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

St. 9 (770 m, around 4 wt%), and St. 10 (1024 m, around 3 wt%), POC content was stable only in the upper ~ 10 cmbsf at St. 4 (150 m, around 10 wt%) and St. 6 (253 m, around 15 wt%), followed by a decrease until the deepest sampled depth (2 and 9 wt%, respectively). At St. 8 (407 m), POC content increased with sediment depth below 3 cmbsf (from 4 to 9 wt%), which consisted of dense clay (see above). In the upper 3 cmbsf, POC decreased from ~ 7 to ~ 4 wt%, which was the sediment layer with a more fluffy appearance.

The sediment surface C/N ratio was lowest at St. 1 (70 m, 6.2) and increased along the continental shelf showing the highest surface C/N ratio at St. 10 (1024 m, 11). St. 8 (407 m) was exceptional, as it showed slightly lower surface C/N ratio (8) as at St. 6 (253 m, 9). St. 8 (407 m) was also the only site showing an increase of 4 units in the upper 0–5 cmbsf, followed by stable ratios around 12 throughout the rest of the core. St. 1 and 4 (70 and 145 m) showed shallower increases in C/N ratio in the upper ~ 2 and 1 cmbsf, respectively, followed by stable ratios around 10 until the bottom of the core. At St. 9 and 10 (770 and 1024 m), C/N ratios ranged around 11 and 12, respectively.

The highest increase in methane concentration was observed at St. 1 (70 m). Here, methane increased linearly from the surface (1  $\mu\text{M}$ ) to the bottom of the core (100  $\mu\text{M}$ ). All other stations showed either no clear trend (St. 4= 145 m) or only slight methane increases with depth. At St. 9 (770 m), even a decrease in methane concentration was observed from the surface to the bottom of core.

Besides St. 1 (70 m), which showed a strong decrease in sulfate ( $\text{SO}_4^{2-}$ ) concentration with depth from about 28 mM at the top to about 9 mM at the bottom of the core (43 cmbsf), all other stations showed  $\text{SO}_4^{2-}$  concentrations > 25 mM throughout the cores. At St. 4, 6 and 9 (145, 253, 770 m),  $\text{SO}_4^{2-}$  showed very slight decrease with depth from about 28 mM at the top to about 25 mM at the bottom of the core. Porewater  $\text{SO}_4^{2-}$  concentrations were stable around 28 mM throughout the core at St. 8 and 10 (407 and 1024 m).

Dissolved inorganic carbon (DIC) concentration increased with depth at St. 1–6 (70–253 m). St. 1 (70 m) showed the steepest increase with depth, showing the lowest DIC concentration at the top (2.3 mM) and the highest at the deepest sampled depth (21.6 mM). At St. 4 (153 m), maximum concentration was reached at ~ 23 cmbsf with 4 mM. St. 6 (253 m) showed maximum concentration at the deepest sampled depth with 9 mM. St. 8 and 9 (407 and 770 m) showed stable DIC concentrations around 2.3 mM throughout the core. No DIC data was available for St. 10 (1024 m).

### 3.4 Net methanogenesis and gross sulfate reduction in MUC cores

Maximum net methanogenesis rates were detected at St. 1 (70 m,  $1.1 \pm 0.5 \text{ nmol cm}^{-3} \text{ d}^{-1}$ , 20–25 cmbsf) and St. 6 (253 m,  $1.3 \pm 0.65 \text{ nmol cm}^{-3} \text{ d}^{-1}$ , 25–30 cmbsf). At all other stations, methanogenesis was mostly below  $0.5 \text{ nmol cm}^{-3} \text{ d}^{-1}$  throughout the cores. St. 8 (407 m) showed methanogenesis activity only in the top 10 cmbsf with the maximum at 5–10 cmbsf ( $0.2 \pm 0.5 \text{ nmol cm}^{-3} \text{ d}^{-1}$ ). At St. 9 and 10 (770 and 1024 m), maximum methanogenesis activity was found in the surface layer (0–5 cmbsf) with  $0.3 \pm 0.4 \text{ nmol cm}^{-3} \text{ d}^{-1}$  and  $0.4 \pm 0.6 \text{ nmol cm}^{-3} \text{ d}^{-1}$ , respectively. St. 10 (1024 m) also showed high average methanogenesis at 10–15 cmbsf ( $1.5 \pm 2.5 \text{ nmol cm}^{-3} \text{ d}^{-1}$ ), which was caused by a single high replicate ( $4.3 \text{ nmol cm}^{-3} \text{ d}^{-1}$ ). In the following, e.g., integration of rates, we will exclude this single high replicate, which will be further elaborated in the discussion.

At all stations beside St. 9 (770 m), sulfate reduction activity was highest in the 0–1 cmbsf horizon, followed by a sharp decrease in activity of 20–90% in the subsequent 1–2 cmbsf horizon. Highest measured rates at 0–1 cmbsf were observed at St. 4 (145 m,  $290 \text{ nmol cm}^{-3} \text{ d}^{-1}$ ), followed by St. 1 (70 m,  $270 \text{ nmol cm}^{-3} \text{ d}^{-1}$ ). Surface (0–1 cmbsf) sulfate reduction activity decreased from St. 4 (145 m) to St. 8 (407 m) with concomitant increase in water depth. St. 9 (770 m) was the only site without a distinct surface sulfate reduction maximum. Here, highest rates were found at 7 cmbsf ( $11.2 \text{ nmol cm}^{-3} \text{ d}^{-1}$ ).



## Microbial methanogenesis in the sulfate-reducing zone

J. Maltby et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



St. 6, 8 and 9 (253, 407, and 770 m) showed a second but smaller maximum of sulfate reduction activity. At St. 6 (253 m), this second maximum was situated at 20.5 cmbsf ( $6.2 \text{ nmol cm}^{-3} \text{ d}^{-1}$ ). St. 8 and 9 (407 and 770 m) showed additional maxima at 4.5 cmbsf ( $3.1 \text{ nmol cm}^{-3} \text{ d}^{-1}$ ) and 2.5 cmbsf ( $1.5 \text{ nmol cm}^{-3} \text{ d}^{-1}$ ), respectively. At St. 9 (770 m), sulfate reduction activity was not detectable at most depths  $> 10$  cmbsf. At St. 10 (1024 m), no sulfate reduction activity was detectable throughout the entire core. At St. 9 and 10 (770 and 1024 m) we cannot exclude that sulfate reduction was present but undetectable due to long, unfrozen storage of the samples (see Sect. 2.7).

Figure 3 shows an overview of integrated methanogenesis and sulfate reduction rates (over the upper 0–25 cm) along the depth transect on the Peruvian margin. Highest integrated surface methanogenesis activity was detected on the shelf (70, 145 and 253 m) with  $0.1 \pm 0.03$ ,  $0.06 \pm 0.02$ , and  $0.07 \pm 0.01 \text{ mmol m}^{-2} \text{ d}^{-1}$ , respectively. St. 8 (407 m) revealed the lowest integrated methanogenesis rate of all sites ( $0.02 \pm 0.00 \text{ mmol m}^{-2} \text{ d}^{-1}$ ). St. 9 (770 m) and St. 10 (1024 m) showed integrated methanogenesis activity around  $0.03 \pm 0.02 \text{ mmol m}^{-2} \text{ d}^{-1}$ , respectively.

Integrated sulfate reduction activity decreased along the continental margin with increasing water depth, revealing the highest activity at the St. 1 (70 m,  $4.7 \text{ mmol m}^{-2} \text{ d}^{-1}$ ) and the lowest activity at St. 9 (770 m,  $0.2 \text{ mmol m}^{-2} \text{ d}^{-1}$ ). Please note again, that integrated sulfate reduction rates are probably underestimated due to long, unfrozen storage of the samples (see Sect. 2.7).

### 3.5 Potential competitive and non-competitive methanogenesis in sediment slurries from MUC cores

Results from the sediment slurry experiments, in which we added either the sulfate reduction inhibitor molybdate, the non-competitive substrate methanol, or no additives (control), are shown in Fig. 4. During the first phase of incubation, all three treatments showed rates within the same order of magnitude. Nevertheless, potential methanogenesis rates were significantly higher ( $p < 0.05$ ) in all treatments in the shallow sediment horizon (0–5 cmbsf) compared to the deep horizon (20–25 cmbsf). In addition, po-

tential methanogenesis was always significantly higher in the molybdate and methanol treatment compared to the control.

During the second phase of the incubation (day 8–23), potential methanogenesis showed a different pattern. Rates in the methanol treatment were 350 and 4 times higher compared to the control and molybdate treatment in the 0–5 cm horizon and the 20–25 cm horizon, respectively ( $p < 0.05$ ). Control and molybdate treatments showed no significant difference ( $p > 0.05$ ) in the shallow and deep horizon.

### 3.6 Geochemical parameters and gross hydrogenotrophic methanogenesis activity in gravity cores

At the shallow St. 1 (78 m), POC concentration slightly decreased with depth, from ~ 4 wt % at the surface to about 2–3 wt % at the bottom of the core (385 cmbsf, Fig. 5). At St. 8 (407 m), POC concentrations were slightly higher with values ranging around 8–9 wt % in the upper 120 cmbsf, and then decreasing with depth. The C/N ratio at St. 1 (78 m) remained around 10 throughout the core, while it showed slightly higher values around 12 throughout the core at St. 8 (407 m).

At St. 1 (78 m), the methane concentration increased with depth from 0.1 mM at the surface to the highest measured concentration at 165 cmbsf (~ 5 mM), followed by a decrease to ~ 2 mM at 198 cmbsf. Methane concentration stayed around 2 mM until the deepest measured depth (385 cmbsf).

Methane concentrations at St. 8 (407 m) ranged from 14 to 17  $\mu\text{M}$  in the upper 120 cmbsf, then increased to a maximum of 36  $\mu\text{M}$  at 180 cmbsf, followed by a decrease to 28  $\mu\text{M}$  at the deepest sampled depth (195 cmbsf).

$\text{SO}_4^{2-}$  concentration at St. 1 (78 m) decreased with depth with the highest concentration (10 mM) at the shallowest measured sediment depth (33 cmbsf) and the lowest concentration at 350 cmbsf (0.16 mM). At St. 8 (407 m),  $\text{SO}_4^{2-}$  concentration decreased slightly from ~ 28 mM at the shallowest measured sediment depth (20 cmbsf) to ~ 24 mM at 145 cmbsf, followed by stable concentrations around 25 mM until the bottom of the core.

DIC concentrations were 5–8 times higher at St. 1 (78 m) compared to St. 8 (407 m) and increased with sediment depth from ~ 21 mM at 33 cmbsf to ~ 39 mM at 385 cmbsf. DIC concentrations at St. 8 (407 m) could only be measured at distinct sediment depths due to limited amounts of porewater but still revealed a slight increase with sediment depth (from ~ 3 to ~ 5 mM).

Hydrogenotrophic methanogenesis at St. 1 (78 m) was present but low below 66 cmbsf until it reached a peak between 300 and 400 cmbsf ( $0.7 \text{ nmol cm}^{-3} \text{ d}^{-1}$ ). In contrast, no hydrogenotrophic methanogenesis activity was detected at St. 8 (407 m).

## 4 Discussion

### 4.1 Concurrent activity of methanogenesis and sulfate reduction in surface sediments

Before we discuss the distribution of methanogenesis in the collected sediment cores, it has to be pointed out that the top soft sediment layer (ca. 0–20 cm) of gravity cores is often disturbed or even lost during the coring procedure. Hence, surface parameters in the gravity cores should not be directly compared to the respective depth layers in MUC cores. According to this likely offset, we will use the term “surface methanogenesis/sediments” when referring to MUC cores and “deep methanogenesis/sediments” when referring to gravity cores.

In the present study, methanogenesis and sulfate reduction concurred in surface sediments along the entire depth transect (70–1024 m) on the Peruvian margin ( $12^\circ \text{ S}$ ). Methanogenesis activity was detected in sediment layers that revealed high porewater sulfate concentrations and sulfate reduction activity (besides St. 10, where sulfate reduction was undetectable). Even though absolute sulfate reduction rates were most likely underestimated, we trust relative distribution pattern in the sediment and along the continental margin.

**BGD**

12, 14869–14910, 2015

## Microbial methanogenesis in the sulfate-reducing zone

J. Maltby et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



---

## Microbial methanogenesis in the sulfate-reducing zone

J. Maltby et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



As the competition between methanogens and sulfate reducers for  $H_2$  and acetate was probably never relieved, the detected surface methanogenesis was most likely based on non-competitive substrates such as methanol or methylated compounds including methylated amines or methylated sulfides (Oremland and Polcin, 1982; Oremland and Taylor, 1978; Kiene et al., 1986). Likewise, in a study off Chile (0–6 cm sediment depth, 87 m water depth), surface methanogenesis was found to be coupled to the non-competitive substrate trimethylamine, and not to  $CO_2/H_2$  or acetate, in sediments where sulfate and sulfate reduction was abundant (Ferdelman et al., 1997).

Non-competitive substrate utilization by methanogens in the present study was further confirmed by a significant increase of potential methanogenesis after the addition of methanol to sediment slurries from St. 1 (70 m) (Fig. 4b). The delayed response of methanogenesis after methanol addition (Supplement, Fig. S1), however, suggests that the present microbial methanogenic community was not primarily feeding on methanol. Potentially other non-competitive substrates like dimethyl sulfides were utilized predominantly. While most methylotrophic methanogens are able to use both methanol and methylated amines, growth on dimethyl sulfide appears to be restricted to only a few methylotrophic species (Oremland et al., 1989). Dimethyl sulfides could have build up during the long storage time (~ 6 months) before experimentation. Even though methylated sulfur compounds (e.g., dimethyl sulfide or methanthiol) are mainly produced by organisms in the marine photic zone (e.g., Andreae and Raemdonck, 1983), it was recently postulated that these compounds may also be generated through nucleophilic attack by sulfide on methyl groups in the sedimentary organic matter (Mitterer, 2010). As sulfate reduction was a predominant process in the sediment, it could have delivered sufficient sulfide to produce methylated sulfur compounds. Consequently, results from the sediment slurry experiments might not reflect the activity of the in situ methanogenic community as we cannot exclude community shifts as a response to the availability of alternative substrates that were produced during the long storage.

The utilization of the competitive substrates  $H_2$  and acetate by the methanogens occurs probably only when sulfate reducers are inhibited. Accordingly, potential methano-

**Microbial  
methanogenesis in  
the sulfate-reducing  
zone**

J. Maltby et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

genesis rates in the molybdate treatment of the sediment slurry experiment were significantly higher in the two studied horizons (0–5 and 20–25 cmbsf) compared to the controls during the first phase of the incubation (day 0–5), indicating the usage of competitive substrate facilitated by the inhibition of sulfate reduction. However, in the second phase (day 8–23) of the incubation, rates were much lower in both the control and molybdate treatment and did not show significant differences in both horizons ( $p > 0.05$ ). In this second phase, methane production might have slowed down due to depletion of electron donors.

Hydrogenotrophic methanogenesis in the gravity core from St. 1 (78 m) showed no activity at depths where porewater sulfate concentrations were  $> 0.7$  mM. Instead activity peaked where porewater sulfate was lowest (0.16 mM at 350 cmbsf), supporting the above conclusions regarding competition within the sulfate zone. The observation that sulfate was never completely depleted in the porewater until the bottom of the gravity core, in combination with an increase of iron (II) in the porewater at depths  $> 200$  cmbsf (data not shown), hint to the presence of a cryptic sulfur cycle that is responsible for deep formation of sulfate (Holmkvist et al., 2011; Treude et al., 2014).

In comparison, surface net methanogenesis activity along the Peruvian margin was similar to activity found off Chile at 87 m water depth ( $0$ – $0.6$  nmol cm $^{-3}$  d $^{-1}$ ) (Ferdelman et al., 1997). The slightly higher rates determined in our study (St. 1 = 70 m;  $0.4$ – $1.7$  nmol cm $^{-3}$  d $^{-1}$ ) could be related to different approaches, as our rates represent the sum of net methanogenesis from all available substrates in the sediment, while rates off Chile were based only on CO $_2$ , acetate, and trimethylamine utilization. Hence, total methanogenesis could have been easily underestimated, if methanogenesis was supplied by other substrates, which is not unlikely, as methylotrophic methanogens, which are able to use methanol or methylated amines, were the dominant type of methanogens in these sediments (Ferdelman et al., 1997). Interestingly, the authors detected a high number of acetogens, implicating that acetogenesis competed for methylamines or other methylated compounds (Ferdelman et al., 1997). A competi-

tion with acetogens for methylated substrates is conceivable for our study, but would require further studies.

## 4.2 Surface vs. deep methanogenesis

Maximum single net surface methanogenesis activities detected in our study (0.3–4.3 nmol cm<sup>-3</sup> d<sup>-1</sup>) were found to be at the very low end of or even one order of magnitude lower than organic-rich, sulfate-depleted sediments (9.8–37 nmol cm<sup>-3</sup> d<sup>-1</sup>, 0–40 cmbsf, Treude et al., 2005a, 10–17 nmol cm<sup>-3</sup> d<sup>-1</sup>, 0–30 cmbsf, Schmaljohann, 1996, 100–300 nmol cm<sup>-3</sup> d<sup>-1</sup>, 0–30 cmbsf, Crill and Martens, 1983, 1986, 100–400 nmol cm<sup>-3</sup> d<sup>-1</sup>, 0–3 cmbsf, Alperin et al., 1992). To estimate the overall relevance of surface methanogenesis within the sulfate zone compared to deep methane production, we estimated the deep methane production in our study and compiled an overview of published deep methane production data from the sulfate-free zone of organic-rich sediments (Table 2). For this comparison, the deep methane production was assumed to equal the flux of methane into the sulfate-methane-transition zone (SMTZ), where it is consumed by anaerobic oxidation of methane (AOM). Within the SMTZ, both sulfate and methane are depleted steeply as a result of AOM, thus dividing the sulfate-reducing zone above from the methanogenic zone below. The SMTZ is the main niche for AOM in marine sediments, acting as an important filter for upwards migrating methane (Knittel and Boetius, 2009). The SMTZ can be found at decimeters to tens of meters below the seafloor, depending on the burial rate of reactive organic matter, the depth of the methane production zone, and the transport velocity of methane and sulfate as well as their consumption rates (Knittel and Boetius, 2009).

In the present study, a SMTZ was only detected in the gravity core taken at St. 1 (78 m; Fig. 5), where it was located between 66 and 99 cmbsf, i.e., below the penetration depth of the MUC cores. We estimated a methane flux (= deep methane production) into the SMTZ (from 99 to 66 cmbsf) according to Iversen and Jørgensen (1993) using a seawater methane-diffusion coefficient from Schulz (2006), which was corrected for porosity resulting in a sediment-diffusion coefficient for

### Microbial methanogenesis in the sulfate-reducing zone

J. Maltby et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



methane of  $D_s = 1.325 \times 10^{-5} \text{ cm}^{-2} \text{ s}^{-1}$  at  $15^\circ\text{C}$ . The resulting deep methane production ( $0.8 \text{ mmol m}^{-2} \text{ d}^{-1}$ ) was slightly higher (ratio of 0.13, surface vs. deep) but still in the same magnitude as the integrated surface methanogenesis at St. 1 (70 m;  $0.1 \text{ mmol m}^{-2} \text{ d}^{-1}$ ). Compared to a different study from the Peruvian OMZ, the ratio between shallow ( $0.07$  to  $0.1 \text{ mmol m}^{-2} \text{ d}^{-1}$ , this study) vs. deep ( $8.9 \times 10^{-8}$  to  $2.2 \times 10^{-7} \text{ mmol m}^{-2} \text{ d}^{-1}$ ; Arning et al., 2012) methanogenesis on the shelf (150–250 m) was  $3.2 \times 10^5$  to  $1.1 \times 10^6$ . Both examples highlight the significance of surface methanogenesis, especially on the Peruvian shelf. On the lower Peruvian slope ( $\sim 3800$  m water depth), deep methanogenesis increased (up to  $0.017 \text{ mmol m}^{-2} \text{ d}^{-1}$ ; Arning et al., 2012). In contrast, surface methanogenesis at the deeper St. 10 (1024 m) was lower ( $0.02 \text{ mmol m}^{-2} \text{ d}^{-1}$ ) compared to the shelf indicating a decreasing relevance of surface methanogenesis along the margin with increasing relevance of deep methanogenesis. The decrease of surface methanogenesis with increasing water depth might be correlated to the decreasing organic carbon content and freshness in the sediment (Fig. 6), which will be further discussed in Sect. 4.4.

In comparison with other organic-rich sediments (Table 2), surface methanogenesis off Peru was in the same order of magnitude as most reported deep methanogenesis (e.g., off Namibia, off Chile, Limfjorden). The only exemption is Eckernförde Bay (Baltic Sea), where surface methanogenesis off Peru was less than 15% of deep methanogenesis at Eckernförde Bay. This site has a water depth of only  $\sim 30$  m with high carbon export, featuring extremely high methanogenesis activity below the SMTZ, causing supersaturation and methane gas ebullition (Whiticar, 2002; Treude et al., 2005a).

### 4.3 Potential consumption and emission of surface methane

Due to its closeness to the sediment–water interface, surface methanogenesis along the Peruvian margin could lead to methane emissions from the sediment into the water column. A short diffusion distance, especially in the top most sediment layers, might facilitate a partial escape of methane from consumption by microbes. As surface

BGD

12, 14869–14910, 2015

## Microbial methanogenesis in the sulfate-reducing zone

J. Maltby et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



methanogenesis decreased with water depth (Fig. 3), the methane emission potential appears to be highest on the shelf. Sediment methane concentrations in the 0–2 sediment horizon of all sites along the margin were always higher than bottom-near water methane concentrations ( $\sim 1.5$  m above seafloor; Table 1, Fig. 2), hinting towards an efflux of methane from the sediment. However, more precise profiling of methane at the sediment–water interface would be necessary to confirm this hypothesis. Still, most of the sediment methane profiles suggest methane consumption close to the seafloor to some extent, which would reduce the amount of emitted methane (Fig. 2). AOM might act as an important methane filter at the sediment surface of the shelf stations, where anoxic conditions dominated, while aerobic oxidation might prevail at the deeper stations below the OMZ (St. 9 and 10). The presence of methane oxidation above the SMTZ of organic-rich sediments has been reported earlier (Treude et al., 2005a, b), and could indeed be fueled by surface methanogenesis. An immediate oxidation of the produced methane would explain why sediment methane profiles did not necessarily correlate with peaks in surface methanogenesis (see, e.g., St. 6, 253 m). The importance of AOM for the reduction of methane emissions from surface methanogenesis remains speculative, as explicit data is missing. On the basis of our findings, however, we suggest to consider surface methanogenesis as a possible driver for AOM above the SMTZ in earlier and future studies.

#### 4.4 Factors controlling methanogenesis along the Peruvian margin

For this discussion we excluded the high integrated methane production observed in one of the replicates at station 10 (1024 m), as we do not think that the detected activity ( $0.23 \text{ mmol m}^{-2} \text{ d}^{-1}$ ) is representative for this deep site, especially as sediment POC content was lowest at station 10 compared to the other stations ( $< 4\%$ , Fig. 2). The outlier might have been caused by additional carbon sources in the sediment, e.g., from fecal pellets or organic carbon released from dead infauna, thus stimulating below-surface microbial activities during our incubations (Ziervogel et al., 2014; Bertics et al., 2013).

**Microbial methanogenesis in the sulfate-reducing zone**

J. Maltby et al.

|  |                              |
|--|------------------------------|
| <a href="#">Title Page</a>               |                              |
| <a href="#">Abstract</a>                 | <a href="#">Introduction</a> |
| <a href="#">Conclusions</a>              | <a href="#">References</a>   |
| <a href="#">Tables</a>                   | <a href="#">Figures</a>      |
| <a href="#">◀</a>                        | <a href="#">▶</a>            |
| <a href="#">◀</a>                        | <a href="#">▶</a>            |
| <a href="#">Back</a>                     | <a href="#">Close</a>        |
| <a href="#">Full Screen / Esc</a>        |                              |
| <a href="#">Printer-friendly Version</a> |                              |
| <a href="#">Interactive Discussion</a>   |                              |





## 4.4.1 Oxygen

Oxygen is an important controlling factor, as methanogenesis is an oxygen- and redox-sensitive process (Oremland, 1988). Some methanogens can tolerate oxygen exposure for several hours before they die, however, no methane will be produced in the presence of oxygen (Zinder, 1993).

Comparing integrated surface methanogenesis (over 0–25 cmbsf) from the shallowest to the deepest station (Fig. 3), highest rates ( $> 0.05 \text{ mmol m}^{-2} \text{ d}^{-1}$ ) were detected on the shelf (St. 1, 4 and 6 = 70, 145, 253 m), where oxygen concentrations were below detection (Fig. 6), providing advantageous conditions for methanogenesis, particularly at the very sediment surface, where normally aerobic respiration dominates (Jørgensen, 2006). Below the OMZ, integrated methanogenesis decreased. Bioturbating macrofauna and megafauna (e.g., mussels, polychaetes, oligochaetes) were observed at these sites (St. 9 and 10, 770 and 1024 m) (Mosch et al., 2012), which could have transported oxygen into deeper sediment layers (Orsi et al., 1996), thus leading to less reduced conditions ( $> -200 \text{ mV}$ ) unsuitable for methanogens (Oremland, 1988). However, integrated methanogenesis was lowest at St. 8 (407 m), which still revealed anoxic bottom water. Thus, oxygen might just be advantageous but not the driving factor for surface methanogenesis.

## 4.4.2 Organic matter

The probably most important factor controlling benthic methanogenesis activity is the POC content of the sediment, as it determines the substrate availability and variety, and can thus relieve the competitive situation between methanogens and sulfate reducers (Holmer and Kristensen, 1994; Treude et al., 2009). Hence, we would expect high methanogenesis at sites with high organic carbon load along the Peruvian margin. Conversely, integrated methanogenesis rates are not correlating with sediment POC content (Fig. 6). While POC content was increasing from St. 1 (70 m) to St. 6 (253 m), followed by a decrease until St. 10 (1024 m), integrated methanogenesis showed rather

**BGD**

12, 14869–14910, 2015

### Microbial methanogenesis in the sulfate-reducing zone

J. Maltby et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



---

## Microbial methanogenesis in the sulfate-reducing zone

J. Maltby et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



a decreasing trend with increasing water depth. This deviation might be caused by another factor, as not only the quantity of organic matter is important for microbial degradation but also its quality, i.e., freshness. Numerous studies have shown that the quality of the organic matter is important for the rate and magnitude of microbial organic matter degradation (Westrich and Berner, 1984; Canfield, 1994; Amon et al., 2001; Middelburg, 1989).

Integrated methanogenesis and C/N ratios (indicating the freshness of organic matter) were negatively correlated along the Peruvian margin (Fig. 6), suggesting that fresh, labile organic matter is advantageous for surface methanogenesis. As methanogens consume mostly short, monomeric substrates, they depend on other microbial groups to break down large organic macromolecules (Zinder, 1993). Hence, labile organic matter offers an important supply of methanogenic substrates.

In agreement with this hypothesis, highest integrated methanogenesis rates were observed at St. 1 (70 m), which revealed the freshest organic matter (lowest C/N, Fig. 6) and the highest POC remineralization rates along the Peruvian margin (Dale et al., 2015). The degradation of organic matter within the water column was probably limited at St. 1 (70 m) due to anoxic conditions and high sedimentation rates (Dale et al., 2015); hence, labile organic matter accumulated at the seafloor, thereby increasing the benthic POC degradation and resulting in high substrate availability and variety for the methanogenic community.

Nevertheless, lowest methanogenesis was measured at St. 8 (407 m), which was neither the site of the highest C/N ratio, lowest POC content (Fig. 6), or the lowest POC mineralization (Dale et al., 2015). In this particular case, methanogenesis was most likely controlled by the sediment properties. Methanogenesis activity was undetectable below 10 cmbsf, which coincided with a very dense and sticky clay layer. The POC profile at St. 8 (407 m) revealed lower concentrations in the upper 5 cmbsf, followed by an increase with depth, suggesting that either the organic matter at this station was resistant to microbial attack (indicated by the increase in C/N) or that microbes were not as frequent/active in the dense clay layer as at the surface. Similarly, sulfate

reduction and microbial nitrogen fixation (Gier et al., 2015) showed very low activity at this site (Fig. 2).

## 5 Conclusions

The present study demonstrated that methanogenesis coincides with sulfate reduction in surface sediments (< 30 cmbsf) along the Peruvian margin. The competition with sulfate reducers was partially relieved due to the high load of organic carbon allowing both groups to show concurrent activity through the utilization of non-competitive substrates by the methanogens.

The significance of surface methanogenesis was high on the shelf, where ratios between surface and deep methanogenesis was around 0.13 (this study) or even as high as  $\sim 10^5$  (compared to Arning et al., 2012), and decreased with increasing water depth. Accordingly, we assume that potential methane emissions into the water column, indicated by a higher methane concentration at the sediment surface compared to the bottom water, should be highest on the shelf, where surface methane production rates were highest. Our results further hint towards a partial consumption of methane before reaching the sediment–water interface, probably by anaerobic oxidation of methane (AOM). In this case, surface methanogenesis might act as important supplier of methane for AOM above the SMTZ, which has been largely overseen before.

We postulate that the dominant factor controlling surface methanogenesis is the availability of (primarily labile) organic matter. The high load of organic carbon and resulting high organic carbon mineralization rates secure the supply for methanogenic substrates, especially on the shelf, which mitigates the competition between sulfate reducers and methanogens. Anoxic conditions in the overlying water might be advantageous for the oxygen-sensitive process of methanogenesis, but does not appear to primarily control benthic rates, as they change within the anoxic zones.

### Microbial methanogenesis in the sulfate-reducing zone

J. Maltby et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Interestingly, organic matter made available by bioturbating infauna (e.g., fecal pellets or dead organisms) could be an important additional factor facilitating methanogenesis in surface sediments. As shown in this study, methanogenesis rates vary strongly in bioturbated sediments below the OMZ, sometimes exceeding all other observed methanogenic rates.

Future studies should seek to (1) identify methanogens and their metabolic capabilities in surface sediments, (2) determine the direct interaction between surface methanogenesis and AOM, (3) evaluate the effect of organic matter hot spots on total benthic surface methanogenesis in organic-rich sediments.

**The Supplement related to this article is available online at doi:10.5194/bgd-12-14869-2015-supplement.**

*Author contributions.* J. Maltby and T. Treude designed the experiments. J. Maltby carried out all methanogenesis experiments, T. Treude conducted sulfate reduction measurements. Pore-water measurements of MUC cores were coordinated by A. W. Dale and S. Sommer. J. Maltby prepared the manuscript with contributions of all co-authors.

*Acknowledgements.* We thank the captain and crew of R.V. *Meteor* for field assistance. We thank A. Petersen, S. Kriwanek and S. Cherednichenko and the shipboard scientific party for field and laboratory assistance. For the geochemical analysis we want to thank B. Domeyer, A. Bleyer, U. Lomnitz, R. Suhrberg, S. Trinkler and V. Thoenissen. Additional thanks goes to G. Schuessler, P. Wefers, and S. Krause of the Treude working group for their laboratory assistance. We further thank the authorities of Peru for the permission to work in their territorial waters. This study is a contribution of the Sonderforschungsbereich 754 "Climate – Biogeochemistry Interactions in the Tropical Ocean" (www.sfb754.de), which is supported by the German Research Foundation. Further support came from the Cluster of Excellence "The Future Ocean" funded by the the German Research Foundation.

The article processing charges for this open-access publication were covered by a Research Centre of the Helmholtz Association.

14896

BGD

12, 14869–14910, 2015

**Microbial  
methanogenesis in  
the sulfate-reducing  
zone**

J. Maltby et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## References

- Alperin, M. J., Blair, N. E., Albert, D. B., Hoehler, T. M., and Martens, C. S.: Factors that control the stable isotopic composition of methane produced in an anoxic marine sediment, *Global Biogeochem. Cy.*, 6, 271–291, 1992.
- 5 Amon, R. M. W., Fitznar, H.-P., and Benner, R.: Linkages among the bioreactivity, chemical composition, and diagenetic state of marine dissolved organic matter, *Limnol. Oceanogr.*, 46, 287–297, 2001.
- Andreae, M. O. and Raemdonck, H.: Dimethyl sulfide in the surface ocean and the marine atmosphere: a global view, *Science*, 221, 744–747, 1983.
- 10 Arning, E. T., Van Berk, W., and Schulz, H. M.: Quantitative geochemical modeling along a transect off Peru: carbon cycling in time and space, and the triggering factors for carbon loss and storage, *Global Biogeochem. Cy.*, 26, 1–18, 2012.
- Bange, H. W., Bartell, U. H., Rapsomanikis, S., and Andreae, M. O.: Methane in the Baltic and North Seas and a reassessment of the marine emissions of methane, *Global Biogeochem. Cy.*, 8, 465–480, available at: <http://doi.wiley.com/10.1029/94GB02181> (last access: 29 June 2015), 1994.
- 15 Bange, H. W., Hansen, H. P., Malien, F., Laß, K., Karstensen, J., Petereit, C., Friedrichs, G., and Dale, A.: Boknis Eck Time Series Station (SW Baltic Sea): Measurements from 1957 to 2010, 1, *LOICZ Inprint*, 16–22, 2011.
- 20 Bertics, V. J., Löscher, C. R., Salonon, I., Dale, A. W., Gier, J., Schmitz, R. A., and Treude, T.: Occurrence of benthic microbial nitrogen fixation coupled to sulfate reduction in the seasonally hypoxic Eckernförde Bay, Baltic Sea, *Biogeosciences*, 10, 1243–1258, doi:10.5194/bg-10-1243-2013, 2013.
- Buckley, D. H., Baumgartner, L. K., and Visscher, P. T.: Vertical distribution of methane metabolism in microbial mats of the Great Sippewissett Salt Marsh, *Environ. Microbiol.*, 10, 967–77, 2008.
- 25 Canfield, D. E.: Factors influencing organic carbon preservation in marine sediments, *Chem. Geol.*, 114, 315–329, 1994.
- Crill, P. and Martens, C.: Spatial and temporal fluctuations of methane production in anoxic coastal marine sediments, *Limnol. Oceanogr.*, 28, 1117–1130, 1983.
- 30 Crill, P. M. and Martens, C. S.: Methane production from bicarbonate and acetate in an anoxic marine sediment, *Geochim. Cosmochim. Ac.*, 50, 2089–2097, 1986.

## Microbial methanogenesis in the sulfate-reducing zone

J. Maltby et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Microbial methanogenesis in the sulfate-reducing zone

J. Maltby et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Dale, A. W., Sommer, S., Lomnitz, U., Montes, I., Treude, T., Liebetrau, V., Gier, J., Hensen, C., Dengler, M., Stolpovsky, K., Bryant, L. D., and Wallmann, K.: Organic carbon production, mineralisation and preservation on the Peruvian margin, *Biogeosciences*, 12, 1537–1559, doi:10.5194/bg-12-1537-2015, 2015.

5 Dimitrov, L.: Mud volcanoes – the most important pathways for degassing deeply buried sediments, *Earth Sci. Revi.*, 59, 49–76, 2002.

Donnelly, M. I. and Dagley, S.: Production of methanol from aromatic acids by *Pseudomonas putida*, *J. Bacteriol.*, 142, 916–924, 1980.

10 Ferdelman, T. G., Lee, C., Pantoja, S., Harder, J., Bebout, B. M., and Fossing, H.: Sulfate reduction and methanogenesis in a Thioploca-dominated sediment off the coast of Chile, *Geochim. Cosmochim. Ac.*, 61, 3065–3079, 1997.

Fuenzalida, R., Schneider, W., Garcés-Vargas, J., Bravo, L., and Lange, C.: Vertical and horizontal extension of the oxygen minimum zone in the eastern South Pacific Ocean, *Deep-Sea Res. Pt. II*, 56, 992–1003, 2009.

15 Gallardo, V. A.: Large benthic microbial communities in sulphide biota under Peru–Chile sub-surface countercurrent, *Nature*, 268, 331–332, 1977.

Gier, J., Sommer, S., Löscher, C. R., Dale, A. W., Schmitz, R. A., and Treude, T.: Nitrogen fixation in sediments along a depth transect through the Peruvian oxygen minimum zone, *Biogeosciences Discuss.*, 12, 14401–14440, doi:10.5194/bgd-12-14401-2015, 2015.

20 Hines, M. E. and Buck, J. D.: Distribution of methanogenic and sulfate-reducing bacteria in near-shore marine sediments, *Appl. Environ. Microb.*, 43, 447–453, 1982.

Holmer, M. and Kristensen, E.: Coexistence of sulfate reduction and methane production in an organic-rich sediment, *Mar. Ecol.-Prog. Ser.*, 107, 177–184, 1994.

25 Holmkvist, L., Ferdelman, T. G., and Jørgensen, B. B.: A cryptic sulfur cycle driven by iron in the methane zone of marine sediment (Aarhus Bay, Denmark), *Geochim. Cosmochim. Ac.*, 75, 3581–3599, 2011.

IPCC: Climate Change 2014: Synthesis Report, contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, edited by: T. core writing Team, Pachauri, R. K., and Meyer, L. A., Geneva, Switzerland, 2014.

30 Iversen, N. and Jørgensen, B. B.: Diffusion coefficients of sulfate and methane in marine sediments: influence of porosity, *Geochim. Cosmochim. Ac.*, 57, 571–578, 1993.

Jørgensen, B. B.: The sulfur cycle of a coastal marine sediment (Limfjorden, Denmark), *Limnol. Oceanogr.*, 22, 814–832, 1977.

---

**Microbial  
methanogenesis in  
the sulfate-reducing  
zone**

---

J. Maltby et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Jørgensen, B. B.: A comparison of methods for the quantification of bacterial sulfate reduction in coastal marine sediments: I. Measurements with radiotracer techniques, *Geomicrobiol. J.*, 1, 11–27, 1978.

Jørgensen, B. B.: Bacteria and marine biogeochemistry, in: *Marine Geochemistry*, edited by: Schulz, H. D. and Zabel, M., Springer-Verlag, Berlin/Heidelberg, 173–207, 2006.

Jørgensen, B. B. and Parkes, R. J.: Role of sulfate reduction and methane production by organic carbon degradation in eutrophic fjord sediments (Limfjorden, Denmark), *Limnol. Oceanogr.*, 55, 1338–1352, 2010.

Judd, A., Davies, G., Wilson, J., Holmes, R., Baron, G., and Bryden, I.: Contributions to atmospheric methane by natural seepages on the UK continental shelf, *Mar. Geol.*, 137, 165–189, 1997.

Kallmeyer, J., Ferdelman, T. G., Weber, A., Fossing, H., and Jørgensen, B. B.: Evaluation of a cold chromium distillation procedure for recovering very small amounts of radiolabeled sulfide related to sulfate reduction measurements, *Limnol. Oceanogr.-Meth.*, 2, 171–180, 2004.

Kamykowski, D. and Zentara, S.: Hypoxia in the world ocean as recorded in the historical data set, *Deep-Sea Res.*, 37, 1861–1874, 1990.

Kiene, R. P., Oremland, R. S., Catena, A., Miller, L. G., and Capone, D. G.: Metabolism of reduced methylated sulfur compounds in anaerobic sediments and by a pure culture of an estuarine methanogen, *Appl. Environ. Microb.*, 52, 1037–1045, 1986.

King, G. M., Klug, M. J., and Lovley, D. R.: Metabolism of acetate, methanol, and methylated amines in intertidal sediments of lowes cove, maine, *Appl. Environ. Microb.*, 45, 1848–1853, 1983.

Knittel, K. and Boetius, A.: Anaerobic oxidation of methane: progress with an unknown process, *Annu. Rev. Microbiol.*, 63, 311–34, 2009.

Middelburg, J. J.: A simple rate model for organic matter decomposition in marine sediments, *Geochim. Cosmochim. Ac.*, 53, 1577–1581, 1989.

Mitterer, R. M.: Methanogenesis and sulfate reduction in marine sediments: A new model, *Earth Planet. Sc. Lett.*, 295, 358–366, 2010.

Neill, a R., Grime, D. W., and Dawson, R. M.: Conversion of choline methyl groups through trimethylamine into methane in the rumen, *Biochem. J.*, 170, 529–535, 1978.

## Microbial methanogenesis in the sulfate-reducing zone

J. Maltby et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Niewöhner, C., Hensen, C., Kasten, S., Zabel, M., and Schulz, H. D.: Deep sulfate reduction completely mediated by anaerobic oxidation in sediments of the upwelling area off Namibia, *Geochim. Cosmochim. Ac.*, 62, 455–464, 1998.

Oremland, R. S.: Biogeochemistry of methanogenic bacteria, in: *Biology of Anaerobic Microorganisms*, edited by: Zehnder, A. J. B., J. Wiley and Sons, New York, 641–705, 1988.

Oremland, R. S. and Capone, D. G.: Use of specific inhibitors in biogeochemistry and microbial ecology, in: *Advances in Microbial Ecology*, edited by: Marshall, K. C., Springer US, Boston, MA, 285–383, 1988.

Oremland, R. S. and Polcin, S.: Methanogenesis and sulfate reduction: competitive and non-competitive substrates in estuarine sediments, *Appl. Environ. Microb.*, 44, 1270–1276, 1982.

Oremland, R. S. and Taylor, B. F.: Sulfate Reduction and Methanogenesis in Marine Sediments, *Geochim. Cosmochim. Ac.*, 42, 209–214, 1978.

Oremland, R. S., Kiene, R. P., Mathrani, I., Whiticar, M. J., and Boone, D. R.: Description of an estuarine methylotrophic methanogen which grows on dimethyl sulfide, *Appl. Environ. Microb.*, 55, 994–1002, 1989.

Oremland, R. S., Marsh, L., and Desmarais, D. J.: Methanogenesis in Big Soda Lake, Nevada: an alkaline, moderately hypersaline desert lake, *Appl. Environ. Microb.*, 43, 462–468, 1982.

Orsi, T. H., Werner, F., Milkert, D., Anderson, A. L., and Bryant, W. R.: Environmental overview of Eckernförde Bay, northern Germany, *Geo-Mar. Lett.*, 16, 140–147, 1996.

Pennington, J. T., Mahoney, K. L., Kuwahara, V. S., Kolber, D. D., Calienes, R., and Chavez, F. P.: Primary production in the eastern tropical Pacific: a review, *Prog. Oceanogr.*, 69, 285–317, 2006.

Reeburgh, W.: Oceanic methane biogeochemistry, *Chem. Rev.*, 107, 486–513, 2007.

Reimers, C. E. and Suess, E.: The partitioning of organic carbon fluxes and sedimentary organic matter decomposition rates in the ocean, *Mar. Chem.*, 13, 141–168, 1983.

Røy, H., Weber, H. S., Tarpgaard, I. H., Ferdelman, T. G., and Jørgensen, B. B.: Determination of dissimilatory sulfate reduction rates in marine sediment via radioactive <sup>35</sup>S tracer, *Limnol. Oceanogr.-Meth.*, 12, 196–211, 2014.

Schink, B. and Zeikus, J. G.: Microbial ecology of pectin decomposition in anoxic lake sediments and in defined laboratory cultures of species prevalent in the lake sediment. The turnover, *J. Gen. Microbiol.*, 128, 393–404, 1982.

Schmaljohann, R.: Methane dynamics in the sediment and water column of Kiel Harbour (Baltic Sea), *Mar. Chem.*, 131, 263–273, 1996.



## Microbial methanogenesis in the sulfate-reducing zone

J. Maltby et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Schulz, H. D.: Quantification of early diagenesis: dissolved constituents in marine pore water, in: *Marine Geochemistry*, edited by: Schulz, H. D. and Zabel, M., Springer Berlin Heidelberg, Berlin Heidelberg, 75–124, 2006.

5 Seeberg-Elverfeldt, J., Schluter, M., Feseker, T., and Kolling, M.: Rhizon sampling of porewaters near the sediment–water interface of aquatic systems, *Limnol. Oceanogr.-Meth.*, 3, 361–371, 2005.

Senior, E., Lindström, E. B., Banat, I. M., and Nedwell, D. B.: Sulfate reduction and methanogenesis in the sediment of a saltmarsh on the East coast of the United Kingdom, *Appl. Environ. Microb.*, 43, 987–996, 1982.

10 Smetacek, V.: The annual cycle of Kiel bight plankton: a long-term analysis, *Estuaries*, 8, 145–157, 1985.

Thauer, R. K.: *Biochemistry of methanogenesis: a tribute to Marjory Stephenson*, Microbiology, 144, 2377–2406, 1998.

15 Treude, T., Krüger, M., Boetius, A., and Jørgensen, B. B.: Environmental control on anaerobic oxidation of methane in the gassy sediments of Eckernförde Bay (German Baltic), *Limnol. Oceanogr.*, 50, 1771–1786, 2005a.

Treude, T., Niggemann, J., Kallmeyer, J., Wintersteller, P., Schubert, C. J., Boetius, A., and Jørgensen, B. B.: Anaerobic oxidation of methane and sulfate reduction along the Chilean continental margin, *Geochim. Cosmochim. Ac.*, 69, 2767–2779, 2005b.

20 Treude, T., Smith, C. R., Wenzhöfer, F., Carney, E., Bernardino, A. F., Hannides, A. K., Krgüer, M., and Boetius, A.: Biogeochemistry of a deep-sea whale fall: sulfate reduction, sulfide efflux and methanogenesis, *Mar. Ecol.-Prog. Ser.*, 382, 1–21, 2009.

25 Treude, T., Krause, S., Maltby, J., Dale, A. W., Coffin, R., and Hamdan, L. J.: Sulfate reduction and methane oxidation activity below the sulfate-methane transition zone in Alaskan Beaufort Sea continental margin sediments: implications for deep sulfur cycling, *Geochim. Cosmochim. Ac.*, 144, 217–237, 2014.

Valentine, D. L., Blanton, D. C., Reeburgh, W. S., and Kastner, M.: Water column methane oxidation adjacent to an area of active hydrate dissociation, Eel River Basin, *Geochim. Cosmochim. Ac.*, 65, 2633–2640, 2001.

30 Westrich, J. T. and Berner, R. A.: The role of sedimentary organic matter in bacterial sulfate reduction: the G model tested, *Limnol. Oceanogr.*, 29, 236–249, 1984.

Wever, T. F. and Fiedler, H. M.: Variability of acoustic turbidity in Eckernförde Bay (southwest Baltic Sea) related to the annual temperature cycle, *Mar. Geol.*, 125, 21–27, 1995.

Whiticar, M. J.: Relationships of Interstitial Gases and Fluids During Early Diagenesis in Some Marine Sediments, Sonderforschungsbereich 95, Wechselwirkung Meer-Meeressboden. Ph.D. thesis, Kiel University, 1978.

Whiticar, M. J.: Diagenetic relationships of methanogenesis, nutrients, acoustic turbidity, pockmarks and freshwater seepages in Eckernförde Bay, Mar. Geol., 182, 29–53, 2002.

Widdel, F. and Bak, F.: Gram-negative mesophilic sulfate-reducing bacteria, in: The Prokaryotes, edited by: Balows, A., Trüper, H. G., Dworkin, M., Harder, W., and Schleifer, K.-H., Springer New York, New York, NY, 3352–3378, 1992.

Ziervogel, K., Joye, S. B., and Arnosti, C.: Microbial enzymatic activity and secondary production in sediments affected by the sedimentation pulse following the Deepwater Horizon oil spill, Deep-Sea Res. Pt. II, 1–8, 2014.

Zinder, S. H.: Physiological ecology of methanogens, in: Methanogenesis, edited by: Ferry, J. G., Chapman and Hall, New York, NY, 128–206, 1993.

**BGD**

12, 14869–14910, 2015

**Microbial  
methanogenesis in  
the sulfate-reducing  
zone**

J. Maltby et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Microbial methanogenesis in the sulfate-reducing zone

J. Maltby et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Table 1.** Stations, instruments, chemical/physical parameters in the bottom-near water, and analyses applied to samples along the depth transect on the Peruvian margin (12° S). For abbreviations see footnote.

| Station No | Instrument | Latitude (S) | Longitude (W) | Water depth (m) | O <sub>2</sub> (μM) | Temp. (°C) | CH <sub>4</sub> (nM) | Type of analysis |
|------------|------------|--------------|---------------|-----------------|---------------------|------------|----------------------|------------------|
| 1          | MUC 13     | 12°13.492    | 77°10.511     | 70              |                     |            |                      | All              |
|            | MUC 38     | 12°13.517    | 77°10.084     | 70              |                     |            |                      | SE               |
|            | GC 8       | 12°14.500    | 77°9.611      | 78              |                     |            |                      | GC-All           |
|            | CTD 9      | 12°13.535    | 77°10.522     | 73              | bdl                 | 14         | 38.6                 | WC               |
| 4          | MUC 10     | 12°18.704    | 77°17.790     | 145             |                     |            |                      | All              |
|            | CTD 14     | 12°18.697    | 77°18.004     | 145             | bdl                 | 13.4       | 24.4                 | WC               |
| 6          | MUC 5      | 12°23.321    | 77°24.176     | 253             |                     |            |                      | Gas +PW          |
|            | MUC 6      | 12°23.322    | 77°24.181     | 253             |                     |            |                      | nMG              |
|            | CTD 6      | 12°24.904    | 77°26.314     | 305             | bdl                 | 12         | 79.6                 | WC               |
| 8          | MUC 23     | 12°27.198    | 77°29.497     | 407             |                     |            |                      | Gas + PW         |
|            | MUC 24     | 12°27.197    | 77°29.497     | 407             |                     |            |                      | nMG              |
|            | GC 3       | 12°27.192    | 77°29.491     | 407             |                     |            |                      | GC-All           |
|            | CTD 37     | 12°29.502    | 77°29.502     | 407             | bdl                 | 10.6       | 7.3                  | WC               |
| 9          | MUC 17     | 12°31.374    | 77°35.183     | 770             |                     |            |                      | Gas + PW         |
|            | MUC 18     | 12°31.373    | 77°35.184     | 770             |                     |            |                      | nMG              |
|            | CTD 27     | 12°31.327    | 77°35.265     | 770             | 19                  | 5.5        | 8.4                  | WC               |
| 10         | MUC 28     | 12°35.377    | 77°40.975     | 1024            |                     |            |                      | Gas + PW         |
|            | MUC 29     | 12°35.377    | 77°40.976     | 1024            |                     |            |                      | nMG              |
|            | CTD 11     | 12°34.863    | 77°38.954     | 1010            | 53                  | 4.4        | 3.9                  | WC               |

MUC = multicorer, GC = gravity corer, CTD = CTD/Rosette, bdl = below detection limit (5 μM), All = methane gas analysis, porewater analysis, net methanogenesis analysis, SE = slurry experiment, GC-All = analysis for gravity cores including methane gas analysis, porewater analysis, hydrogenotrophic methanogenesis analysis, WC = Water column analyses, Gas = methane gas analysis, PW = porewater analysis, nMG = net methanogenesis analysis.

## Microbial methanogenesis in the sulfate-reducing zone

J. Maltby et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)


[Printer-friendly Version](#)

[Interactive Discussion](#)



**Table 2.** Comparison of deep methanogenesis in organic-rich sediments from different regions with surface methanogenesis ( $0.02\text{--}0.1\text{ mmol m}^{-2}\text{ d}^{-1}$ ) determined in the present study. The ratio range was achieved by dividing the lowest surface by the highest deep and the highest surface by the lowest deep methanogenic activity, respectively.

|                                 | Water Depth (m) | Depth of SMTZ (mbsf) | Methane flux into the SMTZ = integrated deep methanogenesis ( $\text{mmol m}^{-2}\text{ d}^{-1}$ ) | Ratio between surface methanogenesis (present study) and deep methanogenesis | Reference                   |
|---------------------------------|-----------------|----------------------|--|--|-----------------------------|
| Namibia (SE Atlantic)           | 1312–2060       | 3–10                 | 0.07–0.15  | 0.13–1.43  | Niewöhner et al. (1998)     |
| Eckernförde Bay (SW Baltic Sea) | 25–28           | 0.5–1.5              | 0.66–1.88  | 0.01–0.15  | Treude et al. (2005a)       |
| Chile (SE Pacific)              | 797–2746        | 3–4                  | 0.068–0.13   | 0.15–1.47  | Treude et al. (2005b)       |
| Limfjorden (North Sea)          | 7–10            | 1–1.5                | 0.076  | 0.03–1.32  | Jørgensen and Parkes (2010) |
| Peru (SE Pacific)               | 150–3819        | 2–50                 | $2.2 \times 10^{-7}$ –0.017  | $1.18\text{--}4.55 \times 10^5$  | Arning et al. (2012)        |
| Peru (SE Pacific)               | 70–1024         | 0.7–1                | 0.8  | 0.03–0.13  | present study               |



**Figure 1.** Location of sampling sites off Peru along the depth transect at 12° S. Source: Schlitzer, R., Ocean Data View, <http://odv.awi.de>, 2014.

## BGD

12, 14869–14910, 2015

### Microbial methanogenesis in the sulfate-reducing zone

J. Maltby et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



Microbial  
methanogenesis in  
the sulfate-reducing  
zone


J. Maltby et al.



**Figure 2.** Profiles of particulate organic carbon (POC), C/N ratio, methane (CH<sub>4</sub>), sulfate (SO<sub>4</sub><sup>2-</sup>), DIC (dissolved inorganic carbon), net methanogenesis (MG) rates and sulfate reduction (SR) rates in the MUC cores along the depth transect. For MG, triplicates (symbols) and mean (solid line) are shown. For SR, duplicates are shown. Data points from the overlying water in the MUC core (OLW) are set to 0 cm. Note deviant scale dimension for MG at St. 10 and for SR at St. 1 and 4.

## Microbial methanogenesis in the sulfate-reducing zone

J. Maltby et al.



**Figure 3.** Integrated methanogenesis and sulfate reduction rates (0–25 cm) along the depth transect. For methanogenesis rates (black bars), average values are shown with standard deviation. Note for St. 10 a mean from two replicates is shown without standard deviation (pattern-filled bar) and the outlier is shown separately (cross). For sulfate reduction rates (blue bars), means from two replicates are shown without standard deviation.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## Microbial methanogenesis in the sulfate-reducing zone

J. Maltby et al.




**Figure 4.** Potential methanogenesis rates in sediment slurry experiments from the two sediment intervals (0–5 cm and 20–25 cm) at St. 1 (70 m). The first phase of the incubation shows rates calculated from day 0 to 5 (a), while the second phase of the incubation summarizes the rates from day 8–23 (b). “Control” is the treatment with sulfate-rich (28 mM) artificial seawater medium, “plus Mb” is the treatment with sulfate-rich artificial seawater medium plus molbdate (Mb, 22 mM), and “plus Meth” is defined as the treatment with sulfate-rich artificial seawater medium plus methanol (Meth, 10 mM). Per treatment, average values are shown with standard deviation. Please note the split-up in the diagram in part B and the different x axis for methanogenesis.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)




## Microbial methanogenesis in the sulfate-reducing zone

J. Maltby et al.




**Figure 5.** Profiles of particulate organic carbon (POC), C/N ratio, methane ( $\text{CH}_4$ ), sulfate ( $\text{SO}_4^{2-}$ ), dissolved inorganic carbon (DIC), and hydrogenotrophic methanogenesis (MG) rates in the gravity cores at two stations within the depth transect. For MG, triplicates (symbols) and mean (solid line) are shown.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

## Microbial methanogenesis in the sulfate-reducing zone

J. Maltby et al.



**Figure 6.** Bottom-near water methane (CH<sub>4</sub>) and oxygen (O<sub>2</sub>) concentrations along the depth transect (above). Surface sediment particulate organic carbon (POC) content and C/N ratio together with integrated methanogenesis (MG) rates (0–25 cmbsf) along the depth transect (below). For MG rates, averages are shown with standard deviation beside St. 10, where a mean from two replicates is shown (see text). Please note the secondary y axis.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)