

Supplement of Biogeosciences Discuss., 12, 15537–15581, 2015
<http://www.biogeosciences-discuss.net/12/15537/2015/>
doi:10.5194/bgd-12-15537-2015-supplement
© Author(s) 2015. CC Attribution 3.0 License.



Supplement of

Predicting biomass of hyperdiverse and structurally complex Central Amazon forests – a virtual approach using extensive field data

D. Magnabosco Marra et al.

Correspondence to: D. Magnabosco Marra (dmarra@bgc-jena.mpg.de)

The copyright of individual parts of the supplement might differ from the CC-BY 3.0 licence.

1 Supplement

2 Tables

3 Table S1. Summary of the allometry data from 101 genera and at least 135 tree species (n = 727) included in this study.

<i>No.</i>	<i>Family</i>	<i>Species</i>	<i>Common name</i>	<i>NT</i>	<i>DBH range</i>	<i>H range</i>	<i>WD</i>	<i>WD lev</i>	<i>SG</i>	<i>AGB range</i>
1	Anacardiaceae	<i>Astronium lecointei</i> Ducke	muiracatiara	1	8.5	11.3	0.752	spp	lat	39.3
2	Anacardiaceae	<i>Thyrsodium guianense</i> Sagot ex Marchand	breu de leite	9	5.6-17.7	4.2-19.5	0.630	spp	mid	8.0-245.1
3	Annonaceae	<i>Annona neoinsignis</i> H.Rainer	envira bobó	10	5.4-24.3	6.9-19.8	0.430	spp	pio	5.4-288.2
4	Annonaceae	<i>Bocageopsis multiflora</i> (Mart.) R.E. Fr.	envira surucucu	2	6.0-21.4	9.0-28.5	0.643	spp	mid	22.2-712.2
5	Annonaceae	<i>Duguetia</i> spp.	envira vermelha	4	6.5-13.9	12-17.5	0.787	gen	lat	23.5-165.1
6	Annonaceae	<i>Duguetia surinamensis</i> R.E. Fr.	envira amarela	2	5.2-5.5	6.0-7.3	0.800	spp	lat	15.2-15.8
7	Annonaceae	<i>Guatteria olivacea</i> R.E. Fr.	envira fofa, envira preta lisa	13	6.8-34.2	11-25.6	0.510	spp	pio	26.6-1690.2
8	Annonaceae	<i>Guatteria</i> spp.	envira branca	3	5.9-16.6	10.8-17.8	0.556	gen	pio	14.4-206.6
9	Annonaceae	ni	envira	1	12.4	18.5	0.630	fam	mid	102.9
10	Annonaceae	<i>Unonopsis stipitata</i> Diels	envira preta cascuda	2	5.5-6.5	8.0-10.8	0.686	spp	mid	12.7-33.9
11	Annonaceae	<i>Xylopia benthamii</i> R.E. Fr.	envira taripucu, embiriba	10	6.6-17.8	5.9-19.2	0.600	spp	mid	20.7-210.2
12	Annonaceae	<i>Xylopia</i> spp.	envira pimenta	4	7.7-12.0	9.8-16.0	0.626	gen	mid	31.1-137.5
13	Apocynaceae	<i>Aspidosperma desmanthum</i> Benth. ex Müll. Arg.	pequiá marfim	1	85.0	33.0	0.610	spp	lat	7509.1
14	Apocynaceae	<i>Couma guianensis</i> Aubl.	sorvinha	1	7.0	8.9	0.560	spp	lat	27.0
15	Apocynaceae	<i>Couma utilis</i> (Mart.) Müll. Arg.	sorva	1	8.5	14.5	0.660	spp	lat	35.0
16	Apocynaceae	<i>Geissospermum argenteum</i> Woodson	acariquara branca	3	5.2-9.8	7.6-13.0	0.760	gen	lat	15.6-66.3
17	Araliaceae	<i>Schefflera morototoni</i> (Aubl.) Maguire, Steyerem. & Frodin	morototó	12	5.8-33.1	9.0-27.0	0.437	spp	pio	9.0-909.3
18	Bignoniaceae	<i>Jacaranda copaia</i> (Aubl.) D. Don	jacaranda, pará pará	1	13.5	17.8	0.348	spp	pio	69.8
19	Boraginaceae	<i>Cordia</i> cf. <i>naidophylla</i>	freijó	2	5.6-6.4	8.6-10.3	0.520	gen	mid	14.2-28.4
20	Burseraceae	<i>Protium decandrum</i> (Aubl.) Marchand	breu branco	5	5.8-9.3	9.9-15.4	0.560	spp	mid	15.3-58.5
21	Burseraceae	<i>Protium hebetatum</i> D.C. Daly	breu vermelho	11	5.2-30.0	4.2-25.0	0.579	gen	mid	7.7-1258.9
22	Burseraceae	<i>Protium</i> sp.	breu	1	10.7	15.0	0.579	gen	mid	74.7
23	Burseraceae	<i>Protium trifoliolatum</i> Engl.	breu peludo	1	6.5	12.0	0.640	spp	mid	20.8
24	Burseraceae	<i>Tetragastris panamensis</i> (Engl.) Kuntze	breu manga	13	5.0-17.5	3.9-19.0	0.742	spp	mid	7.9-159.1
25	Caryocaraceae	<i>Caryocar pallidum</i> A.C. Sm.	piquiarana	2	5.9-8.1	8.8-12.6	0.680	spp	lat	21.4-75.8
26	Chrysobalanaceae	<i>Acioa longipendula</i> (Pilg.) Sothers & Prance	castanha de galinha	1	48.3	29.2	0.940	spp	lat	3994.5
27	Chrysobalanaceae	<i>Licania coriacea</i> Benth.	marí bravo	4	5.8-10.5	10.5-12.3	0.880	spp	lat	19.9-113.7
28	Chrysobalanaceae	<i>Licania heteromorpha</i> Benth.	macucú fofó	1	7.0	14.3	0.740	spp	lat	38.8
29	Chrysobalanaceae	<i>Licania oblongifolia</i> Standl.	macucú chiodor	2	5.0-13.5	8.5-17.6	0.878	spp	lat	19.1-178.6

30	Chrysobalanaceae	<i>Licania</i> spp.	macucú	5	5.2-9.9	7.5-17.8	0.822	gen	lat	16.7-121.3
31	Clusiaceae	<i>Tovomita</i> spp.	sapateiro	2	8.7-37.0	17.2-28.4	0.799	gen	mid	69.0-1624.7
32	Combretaceae	<i>Buchenavia</i> sp.	tanimbuca	1	33.8	17.9	0.719	gen	lat	1801.4
33	Connaraceae	<i>Connarus perrottetii</i> (DC.) Planch.	sacaca brava	28	5.2-8.9	6.2-12.3	0.450	gen	pio	15.0-69.0
34	Dichapetalaceae	<i>Tapura amazonica</i> Poepp.	tapura	2	20-21.2	18.9-21.2	0.711	gen	mid	355.8-599.8
35	Elaeocarpaceae	<i>Sloanea</i> sp.	urucurana	1	5.8	7.6	0.827	gen	lat	21.2
36	Euphorbiaceae	<i>Alchorneopsis</i> sp.	supiarana	1	19.5	23.0	0.390	gen	pio	336.5
37	Euphorbiaceae	<i>Croton draconoides</i> Müll. Arg.	sacaca	6	5.1-7.9	5.7-12.0	0.626	gen	pio	9.0-20.5
38	Euphorbiaceae	<i>Croton matourensis</i> Aubl.	dima	69	5.4-28.7	8.7-26.5	0.620	spp	pio	12.9-1066.6
39	Euphorbiaceae	<i>Hevea guianensis</i> Aubl.	seringa vermelha	1	8.0	11.6	0.575	spp	lat	39.1
40	Euphorbiaceae	<i>Mabea angularis</i> Hollander	taquari branco	4	7.0-11.8	10.0-16.2	0.634	gen	mid	20.6-105.0
41	Euphorbiaceae	<i>Mabea piriri</i> Aubl.	taquari vermelho	21	5.3-16.5	6.4-24.2	0.617	spp	mid	13.5-321.5
42	Euphorbiaceae	<i>Mabea</i> spp.	taquari	5	6.2-9.3	11.8-15.0	0.634	gen	mid	22.6-58.2
43	Euphorbiaceae	<i>Micrandra</i> spp.	seringarana	12	5.1-35.0	7.4-27.5	0.848	gen	lat	15.9-1677.5
44	Euphorbiaceae	<i>Micrandropsis scleroxylon</i> (W.A. Rodrigues) W.A. Rodrigues	piãozinho	11	5.7-32.4	7.2-25.0	0.880	spp	lat	22.9-1235.5
45	Euphorbiaceae	<i>Pogonophora schomburgkiana</i> Miers ex Benth.	amarelinho	1	5.6	10.0	0.833	spp	lat	48.2
46	Fabaceae	<i>Andira micrantha</i> Ducke	sucupira preta	2	6.0-8.1	7.5-12.3	1.000	spp	lat	19.0-45.1
47	Fabaceae	<i>Andira</i> spp.	sucupira	3	5.2-35.0	10.0-30.0	0.730	gen	lat	15.4-2012.2
48	Fabaceae	<i>Bocoa viridiflora</i> (Ducke) R.S. Cowan	muirajiboia preta	1	22.6	20.8	0.835	spp	lat	1165.0
49	Fabaceae	<i>Cedrelinga cateniformis</i> (Ducke) Ducke	cedrorana	1	55.0	34.1	0.480	spp	lat	3754.2
50	Fabaceae	<i>Dipteryx magnifica</i> Ducke	cumarurana	1	7.0	16.0	0.940	gen	lat	43.3
51	Fabaceae	<i>Eperua glabriflora</i> (Ducke) R.S. Cowan	muirapiranga	3	7.5-30.5	9.4-20.5	0.759	spp	lat	23.8-1304.4
52	Fabaceae	<i>Inga</i> cf. <i>pezizifera</i>	ingá vermelha	14	5.5-31.3	7.0-23.0	0.650	spp	pio	14.0-1562.8
53	Fabaceae	<i>Inga</i> spp.	ingá branca	11	5.3-24.9	5.9-22.5	0.614	gen	mid	8.3-485.8
54	Fabaceae	<i>Inga thibaudiana</i> DC.	ingá peluda	5	9.231.9	15-22.7	0.657	spp	mid	62.2-1509.0
55	Fabaceae	<i>Macrobium</i> spp.	ingarana	3	10.3-19.5	14.3-17.0	0.606	gen	mid	78.3-374.9
56	Fabaceae	<i>Ormosia</i> spp.	tento	5	8.0-37.5	12.7-25.4	0.676	gen	mid	30.5-1606.9
57	Fabaceae	<i>Parkia</i> sp.	fava parkia	2	6.5-9.5	9.0-12.5	0.532	gen	lat	10.6-85.0
58	Fabaceae	<i>Pterocarpus rohrii</i> Vahl	mututí	2	29.8-35.5	27.3-34.5	0.550	spp	mid	1240.0-1740.6
59	Fabaceae	<i>Stryphnodendron guianense</i> (Aubl.) Benth.	fava camuzé	14	5.5-32.9	9.5-27.0	0.650	spp	mid	7.5-1120.5
60	Fabaceae	<i>Swartzia ingifolia</i> Ducke	acapú amarelo, ingá ferro	1	40.4	21.6	0.815	spp	lat	1345.3
61	Fabaceae	<i>Swartzia</i> cf. <i>recurva</i>	muirajiboia amarela	2	7.7-12.0	11.7-17.2	0.883	gen	lat	39.7-155.5
62	Fabaceae	<i>Tachigali setifera</i> (Ducke) Zarucchi & Herend.	tachi preto	1	7.0	17.2	0.670	spp	pio	36.5
63	Fabaceae	<i>Zygia racemosa</i> (Ducke) Barneby & J.W. Grimes	angelim pedra	3	6.2-10.7	9.0-15.5	0.748	spp	lat	23.9-82.9
64	Fabaceae	<i>Zygia ramiflora</i> (F. Muell.) Kosterm.	ingá copaiba	2	7.2	8.4-11.3	0.727	gen	mid	16.8-33.0
65	Goupiaceae	<i>Goupia glabra</i> Aubl.	cupiúba	17	5.1-14.2	7.7-17.0	0.730	spp	mid	12.9-216.4
66	Humiriaceae	<i>Endopleura</i> spp.	uchí amarelo	1	9.0	16.8	0.775	gen	lat	72.0
67	Humiriaceae	ni.2	uchí preto	1	14.0	18.5	0.819	fam	lat	281.2

68	Humiriaceae	<i>Sacoglottis ceratocarpa</i> Ducke	uchí coco	1	9.5	12.5	0.807	gen	lat	109.9
69	Humiriaceae	<i>Vantanea</i> sp.	uchirana	1	19.0	17.6	0.857	gen	lat	324.8
70	Hypericaceae	<i>Vismia guianensis</i> (Aubl.) Pers.	lacre branco	15	6.1-23.8	8.6-23.5	0.475	spp	pio	12.0-508.1
71	Lauraceae	<i>Aniba ferrea</i> Kubitzki	louro chumbo	1	29.8	26.6	0.709	gen	lat	1484.5
72	Lauraceae	ni.3	louro amarelo	1	9.2	14.5	0.656	fam	mid	58.1
73	Lauraceae	ni.4	louro fofo	1	5.2	7.8	0.656	fam	mid	21.6
74	Lauraceae	<i>Ocotea</i> spp.	louro preto	4	7.0-36.2	10.0-27.0	0.598	gen	lat	25.2-2162.6
75	Lauraceae	<i>Rhodostemonodaphne</i> spp.	louro peludo	1	6.5	12.2	0.395	gen	mid	35.6
76	Lecythidaceae	<i>Cariniana integrifolia</i> Ducke	tauari	6	5.0-7.5	6.4-10.0	0.490	spp	lat	12.2-29.1
77	Lecythidaceae	<i>Couratari</i> sp.	tauari vermelho	1	5.2	10.5	0.549	gen	lat	19.9
78	Lecythidaceae	<i>Eschweilera atropetiolata</i> S.A. Mori	castanha vermelha	1	20.4	25.5	0.753	spp	lat	715.2
79	Lecythidaceae	<i>Eschweilera collina</i> Eyma	ripeiro branco	2	7.7-15.7	11.2-17.7	0.778	spp	lat	20.5-178.0
80	Lecythidaceae	<i>Eschweilera</i> spp.	matamata	8	5.1-34.3	8.6-28.0	0.810	gen	lat	19.6-2588.0
81	Lecythidaceae	<i>Gustavia cf. elliptica</i>	mucurão	3	7.0-10.2	8.2-13.1	0.648	gen	mid	22.3-93.9
82	Lecythidaceae	<i>Lecythis barnebyi</i> S.A. Mori	castanha jarana	2	7.0-31.0	8.3-24.7	0.821	gen	lat	22.9-1785.8
83	Lecythidaceae	<i>Lecythis graciana</i> S.A. Mori	castanha jarana folha miúda	1	16.5	18.0	0.830	spp	lat	348.6
84	Lecythidaceae	<i>Lecythis</i> sp.	ripeiro vermelho	1	42.4	27.6	0.821	gen	lat	3999.8
85	Lecythidaceae	ni.5		1	5.5	9.5	0.713	fam	lat	18.0
86	Malpighiaceae	<i>Byrsonima duckeana</i> W.R. Anderson	murici vermelho	23	5.6-25.7	9.0-22.1	0.671	gen	pio	21.9-1036.6
87	Malvaceae	<i>Scleronema micranthum</i> (Ducke) Ducke	cardeiro	2	11.5-38.4	13.2-27.6	0.595	spp	mid	75.2-2298.9
88	Malvaceae	<i>Theobroma sylvestre</i> Aubl. ex Mart.	cacaúí	1	5.3	10.4	0.470	spp	mid	20.4
89	Melastomataceae	<i>Bellucia dichotoma</i> Cogn.	goiaba de anta	7	6.3-24.5	9.0-21.8	0.607	gen	pio	30.5-775.6
90	Melastomataceae	<i>Miconia argyrophylla</i> DC.	buxuxu canela de velho	3	5.1-26.0	8.2-22.0	0.637	gen	pio	14.1-426.0
91	Melastomataceae	<i>Miconia minutiflora</i> (Bonpl.) DC.	tinteira, tintarana	4	6.0-13.3	6.3-18.2	0.637	gen	mid	32.5-219.2
92	Melastomataceae	<i>Miconia</i> spp.	buxuxu	24	5.3-13.7	7.8-15.6	0.637	gen	pio	13.9-198.4
93	Melastomataceae	<i>Mouriri</i> sp.	mamãozinho	1	63.2	32.0	0.740	gen	mid	6655.1
94	Meliaceae	<i>Guarea</i> spp.	gitó vermelho	8	5.2-7.3	7.2-11.4	0.652	gen	lat	12.9-30.7
95	Meliaceae	<i>Trichilia</i> sp.	gitó branco	1	5.5	10.7	0.740	gen	lat	17.2
96	Moraceae	<i>Brosimum rubescens</i> Taub.	pau rainha	1	10.5	14.8	0.878	spp	lat	101.9
97	Moraceae	<i>Brosimum</i> spp.	muiratinga	4	6.1-8.5	8.1-11.0	0.666	gen	mid	14.2-57.2
98	Moraceae	<i>Helianthostylis sprucei</i> Baill.	falsa rainha	1	9.8	10.4	0.628	spp	mid	63.1
99	Moraceae	<i>Pseudolmedia</i> sp.	muiratinga	1	13.7	19.0	0.630	gen	mid	182.9
100	Moraceae	<i>Sorocea guillemianiana</i> Gaudich.	jaca brava	2	5.3-5.7	8.0	0.612	gen	mid	11.8-12.2
101	Myristicaceae	<i>Iryanthera juruensis</i> Warb.	ucuúba punã	3	10.1-22.0	12.85-22.0	0.663	spp	mid	65.1-544.5
102	Myristicaceae	<i>Iryanthera lancifolia</i> Ducke	ucuúba vermelha	1	8.5	10.9	0.634	gen	mid	38.1
103	Myristicaceae	<i>Iryanthera</i> sp.	ucuúba	1	32.0	25.0	0.634	gen	mid	1036.6
104	Myristicaceae	<i>Virola michelii</i> Heckel	ucuúba preta	1	8.7	13.0	0.529	spp	mid	58.8
105	Myrtaceae	<i>Eugenia</i> spp.	araçá	3	5.9-6.8	8.4-9.8	0.742	gen	mid	21.3-24.7

106	Myrtaceae	ni.6	araçá bravo	3	8.0-14.1	14.1-17.0	0.788	fam	mid	64.8-230.7
107	Nyctaginaceae	<i>Neea</i> spp.	joão mole	5	5.9-9.3	6.8-9.0	0.631	gen	mid	17.3-38.4
108	Ochnaceae	<i>Ouratea</i> cf. <i>dischophora</i>	uchí de morcego	2	5.3-11.1	12.3-14.8	0.743	gen	mid	35.0-123.0
109	Olaceae	ni.7		1	27.0	24.0	0.692	fam	lat	926.3
110	Peraceae	<i>Pera schomburgkiana</i> (Klotzsch) Müll. Arg.	pera	1	6.2	11.0	0.590	spp	mid	30.4
111	Rubiaceae	<i>Amaioua guianensis</i> Aubl.	muiraximbé	1	6.0	8.3	0.670	spp	mid	19.8
112	Rubiaceae	<i>Faramea</i> spp.	taboquinha	3	6.0-15.0	7.7-17.9	0.618	gen	mid	23.9-134.7
113	Rubiaceae	<i>Psychotria</i> spp.	taboca mata gado	2	6.2-6.9	8.5-10.6	0.564	gen	mid	29.5-39.0
114	Rutaceae	<i>Zanthoxylum</i> sp.	limãozinho	1	6.1	7.0	0.602	gen	mid	13.5
115	Salicaceae	<i>Casearia arborea</i> (Rich.) Urb.	piabinha	4	5.5-9.5	9.4-15.0	0.535	spp	pio	37.9-93.0
116	Salicaceae	<i>Laetia procera</i> (Poepp.) Eichler	periquiteira	86	5.2-15.8	4.9-20.3	0.664	spp	pio	7.8-244.7
117	Sapindaceae	<i>Toulicia guianensis</i> Aubl.	pitomba da mata	3	7.8-9.7	13.0-14.7	0.756	spp	mid	33.5-80.0
118	Sapotaceae	<i>Chrysophyllum</i> cf. <i>prieurii</i>	abiurana sabiá, abiurana roxa	1	38.6	28.8	0.715	gen	lat	2890.4
119	Sapotaceae	<i>Micropholis guyanensis</i> subsp. <i>guyanensis</i> (Baehni) T.D. Penn.	maparajuba, balata brava	1	19.4	23.3	0.663	spp	lat	445.7
120	Sapotaceae	<i>Micropholis</i> sp.	balata rosada, chiclete bravo	1	31.0	27.0	0.676	spp	lat	1535.2
121	Sapotaceae	<i>Pouteria anomala</i> (Pires) T.D. Penn.	abiurana olho de veado	1	45.0	32.7	0.765	spp	lat	3729.8
122	Sapotaceae	<i>Pouteria manaosensis</i> (Aubrév. & Pellegr.) T.D. Penn.	abiurana roxa	1	6.2	8.3	0.640	spp	lat	21.7
123	Sapotaceae	<i>Pouteria oblanceolata</i> Pires	cucutiriba folha peluda	2	8.8-13.7	11.8-21.0	0.790	spp	lat	79.4-234.9
124	Sapotaceae	<i>Pouteria</i> spp.	abiurana	5	7.3-44.0	12.2-24.0	0.801	gen	lat	45.4-2934.2
125	Simaroubaceae	<i>Simarouba amara</i> Aubl.	marupá	1	27.5	21.5	0.404	spp	pio	555.6
126	Siparunaceae	<i>Siparuna</i> cf. <i>guianensis</i>	caápitíú	10	5.4-9.3	4.5-13.0	0.593	gen	mid	15.9-92.2
127	Urticaceae	<i>Cecropia purpurascens</i> C.C. Berg	imbaúba roxa	3	11.8-15.5	14.8-16.5	0.430	spp	pio	63.9-106.4
128	Urticaceae	<i>Cecropia sciadophylla</i> Mart.	imbaúba gigante	33	8.4-37.2	10.9-23.0	0.412	spp	pio	21.6-1543.6
129	Urticaceae	<i>Pourouma</i> sp.	imbaubarana	1	17.3	18.0	0.389	gen	pio	125.9
130	Urticaceae	<i>Pourouma tomentosa</i> Mart. ex Miq.	imbaubarana	4	5.7-29.7	8.5-20.0	0.395	spp	pio	8.5-797.8
131	Verbenaceae	<i>Aegiphila</i> sp.	tabaco bravo	2	18.4-20.0	12.4-20.1	0.555	gen	pio	113.3-253.2
132	Violaceae	<i>Leonia cymosa</i> Mart.	mucurão	1	11.4	11.1	0.673	gen	mid	85.0
133	Violaceae	<i>Rinorea racemosa</i> (Mart.) Kuntze	branquinha, canela de jacamim	9	5.0-10.5	8.0-18.0	0.682	spp	mid	14.6-114.8
134	Violaceae	<i>Rinorea</i> cf. <i>guianensis</i>	falsa cupiúba	5	5.3-23.4	9.4-24.0	0.709	gen	mid	18.1-751.9
135	Vochysiaceae	<i>Erisma bracteosum</i> Ducke	quaruba	3	5.4-7.1	7.0-8.9	0.533	gen	lat	15.2-27.0

4 Species' attributes: *Family* botanic family and *Species* species' name accordingly to the APGIII system (Stevens, 2012); *Common name* species'
5 common name in Central Amazon; *NT* number of trees; *DBH range* range of DBH (cm); *H range* range of total height (m); *WD* wood density (g
6 cm⁻³); *WD level* wood density assignment level (*spp* value at species level, *gen* value at genus level and *fam* value at family level); *SG* species'

7 successional group (*pio* pioneer species, *mid* mid-successional species and *lat* late-successional species); and *AGB range* above-ground biomass
 8 (dry) range (kg).

9

10 Table S2. Attributes used to assign the tree species from this study into successional groups.

<i>SG</i>	Architecture, morphology and anatomy						Ecology and life history		
	<i>BFo</i>	<i>CPo</i>	<i>CSi</i>	<i>CCo</i>	<i>mDBH</i>	<i>mWD</i>	<i>GAs</i>	<i>NRe</i>	<i>DSy</i>
<i>pio</i>	<i>pla, ort</i>	<i>can, eme</i>	<i>med, lar</i>	<i>rou, ell</i>	12.4 (5.1,37.2)	0.573 (0.348,0.671)	yes	BW4, BW7, BW14, BW17, SE14, SGa	<i>ane, zoo</i>
<i>mid</i>	<i>ort</i>	<i>und, can</i>	<i>sma, med</i>	<i>elo, con, irr</i>	9.8 (5.0,35.5)	0.656 (0.395,0.833)	yes	BW17, BW24, BW27, SE23, OG, SGa	<i>zoo, ane</i>
<i>lat</i>	<i>unb</i>	<i>can, eme</i>	<i>med, lar</i>	<i>rou, ell, irr</i>	14.0 (5.0,85.0)	0.735 (0.480,1.000)	no	BW24, BW27, OG	<i>aut, zoo</i>

11 Attributes_[references]: *SG* species' successional group (*pio* pioneer species, *mid* mid-successional species and *lat* late-successional species)_[1]; *BFo*
 12 predominant branch form (*pla* plagiotropic, *ort* orthotropic and *unb* unbranched)_[2,3]; *CPo* general tree crown position of a mature tree (*und*
 13 understory/supressed crown, *can* canopy/co-dominant crown and *eme* emergent crown)_[2,3]; *CSi* relative crown size of a mature tree (*sma* small-
 14 size crown, *med* mediun-size crown and *lar* large-size crown)_[2,3]; *CCo* general crown contour of a mature tree (*rou* round, *ell* elliptic, *elo*
 15 elongated, *con* conic and *irr* irregular)_[2,3,4]; *mDBH* mean DBH (cm) (minimum and maximum) of the trees included in this study_[5]; *mWD* mean
 16 wood density (g cm⁻³) (minimum and maximum) of the tree species included in this study_[5]; *GAs* gap-associated species_[3]; *NRe* natural
 17 regeneration mostly observed-niche_[3]; *DSy* major dispersion vector (*ane* anemochory, *aut* autochory and *zoo* zoochory)_[6]. References: [1]
 18 Amaral et al., 2009; Chambers et al., 2009; Kammesheidt, 2000; Marra et al., 2014; Saldarriaga et al., 1998; Swaine and Whitmore, 1988; [2]
 19 Hallé, 1974; Hallé et al., 1978; allometric data from this study (Table S1); [3] Network of permanent plots including an *OG* old-growth forest
 20 (LMF unpublished data [1996-2012 census] and Silva et al., 2002), a *BW4* four year-old blowdown (Marra et al., 2014), a *BW7* seven year-old
 21 blowdown, a *BW14* 14 year-old blowdown, a *BW17* 17 year-old blowdown, a *BW24* 24 year-old blowdown, a *BW27* 27 year-old blowdown
 22 (LMF unpublished data), a *SE14* 14 year-old slash and burn secondary forest, a *SE23* 23 year-old clear cut secondary forest (Lima et al., 2007;

23 Silva, 2007) and *SGa* small ($< c. 2000 \text{ m}^2$) canopy gaps (LMF unpublished data [1996-2012 census]); [4] Myers, 1982; Trichon, 2001; [5]
 24 Allometric data from this study (Tab. 1 and Tab. S1); [6] Camargo et al., 2008; Ferraz et al., 2004; Ribeiro et al., 1999; Saravy et al., 2003; Silva
 25 Junior and Pereira, 2009; Stefanello et al., 2009, 2010; Terborgh et al., 2008.

26

27 Table S3. Coefficients (low [2.5%] and high [97.5%] confidence interval) of 24 above-ground estimation models fit in this study. See the Table 2
 28 for checking the equations and variance modeling approaches, and the Table 3 for checking models' statistics.

Model series	Model	<i>SG</i>	<i>b1</i>	<i>b2</i>	<i>b3</i>	<i>b4</i>	<i>c1</i>	<i>c2</i>
1	M11	G	1.010 (0.858,1.179)	2.052 (2.012,2.093)			190.5 (180.8,200.4)	
	M12	G	-1.148 (-1.262,-1.031)	2.338 (2.289,2.386)			0.359 (0.341,0.378)	
	M13	G	0.336 (0.299,0.377)	2.339 (2.290,2.387)			0.106 (0.085,0.132)	2.374 (2.285,2.464)
2	M21	pio	0.496 (0.262,0.835)	2.191 (2.018,2.371)			148.8 (141.6,156.7)	
		mid	0.248 (0.082,0.521)	2.499 (2.243,2.795)				
		lat	4.284 (3.632,4.989)	1.714 (1.675,1.754)				
	M22	pio	-0.874 (-1.049,-0.705)	2.203 (2.133,2.274)			0.345 (0.328, 0.365)	
		mid	-1.296 (-1.544,-1.053)	2.400 (2.290, 2.512)				
		lat	-1.335 (-1.535,-1.140)	2.481 (2.402,2.562)				
M23	pio	0.445 (0.375,0.522)	2.202 (2.135,2.271)			0.151 (0.107,0.209)	2.199 (2.061,2.335)	
	mid	0.297 (0.235,0.377)	2.394 (2.289,2.487)			0.234 (0.138,0.372)	2.038 (1.820,2.264)	
	lat	0.269 (0.228,0.314)	2.489 (2.422,2.554)			0.060 (0.037,0.092)	2.574 (2.401,2.756)	
3	M31	G	1.714 (1.502,1.953)	2.036 (2.002,2.069)	1.035 (0.944,1.126)		144.3 (137.1,152.0)	
	M32	G	-0.869 (-0.986,-0.753)	2.375 (2.329,2.420)	0.749 (0.624,0.876)		0.329 (0.313,0.347)	
	M33	G	0.415 (0.373,0.460)	2.401 (2.361,2.439)	0.755 (0.631,0.881)		0.150 (0.119,0.186)	2.189 (2.097,2.282)
4	M41	pio	0.332 (0.181,0.548)	2.577 (2.398,2.767)	1.233 (0.981,1.492)		130.5 (124.1,137.6)	

		mid	0.244 (0.097,0.479)	2.484 (2.238,2.760)	-0.070 (-0.582,0.458)			
		lat	3.293 (2.797,3.832)	1.848 (1.804,1.893)	0.716 (0.591,0.841)			
	M42	pio	-0.552 (-0.724,-0.378)	2.279 (2.211,2.346)	0.875 (0.692,1.055)		0.323 (0.307,0.341)	
		mid	-1.193 (-1.465,-0.913)	2.408 (2.306,2.510)	0.284 (-0.125,0.683)			
		lat	-1.141 (-1.364,-0.919)	2.468 (2.394,2.544)	0.510 (0.185,0.829)			
	M43	pio	0.539 (0.454,0.640)	2.326 (2.251,2.397)	0.880 (0.687,1.079)		0.217 (0.151,0.300)	2.005 (1.868,2.150)
		mid	0.362 (0.268,0.490)	2.395 (2.285,2.494)	0.464 (0.054,0.956)		0.211 (0.205,0.329)	2.082 (1.874,2.304)
		lat	0.323 (0.270,0.386)	2.475 (2.409,2.538)	0.481 (0.222,0.750)		0.059 (0.037,0.090)	2.561 (2.389,2.744)
5	M51	G	0.078 (0.051,0.114)	1.561 (1.492,1.636)	1.347 (1.152,1.527)		164.3 (156.2,173.0)	
	M52	G	-2.098 (-2.290,-1.912)	1.928 (1.847,2.010)	0.735 (0.615,0.856)		0.329 (0.312,0.346)	
	M53	G	0.147 (0.117,0.179)	1.971 (1.884,2.057)	0.650 (0.523,0.781)		0.092 (0.073,0.116)	2.405 (2.310,2.501)
6	M61	pio	0.029 (0.007,0.076)	1.926 (1.729,2.125)	1.252 (0.860,1.632)		134.1 (127.2,141.1)	
		mid	0.144 (0.052,0.316)	2.173 (1.822,2.543)	0.508 (0.036,0.900)			
		lat	0.697 (0.454,1.041)	1.414 (1.343,1.482)	0.896 (0.709,1.077)			
	M62	pio	-1.946 (-2.238,-1.663)	1.820 (1.713,1.930)	0.751 (0.583,0.921)		0.314 (0.298,0.331)	
		mid	-2.101 (-2.405,-1.799)	1.900 (1.740,2.062)	0.762 (0.562,0.958)			
		lat	-2.223 (-2.665,-1.814)	2.099 (1.914,2.271)	0.688 (0.406,1.001)			
	M63	pio	0.170 (0.118,0.234)	1.870 (1.757,1.984)	0.664 (0.471,0.863)		0.140 (0.098,0.192)	2.202 (2.067,2.341)
		mid	0.147 (0.101,0.207)	1.970 (1.796,2.149)	0.655 (0.414,0.883)		0.193 (0.112,0.309)	2.090 (1.866,2.328)
		lat	0.120 (0.083,0.169)	2.141 (2.000,2.288)	0.626 (0.379,0.861)		0.056 (0.035,0.084)	2.563 (2.388,2.742)
7	M71	G	0.143 (0.099,0.192)	1.577 (1.517,1.631)	1.273 (1.136,1.435)	0.955 (0.881,1.028)	118.0 (112.1,124.2)	
	M72	G	-1.733 (-1.921,-1.541)	2.010 (1.934,2.085)	0.643 (0.530,0.756)	0.654 (0.537,0.769)	0.305 (0.290,0.321)	
	M73	G	0.186 (0.149,0.227)	2.096 (2.017,2.170)	0.706 (0.584,0.831)	0.573 (0.449,0.698)	0.143 (0.112,0.181)	2.183 (2.083,2.285)
8	M81	pio	0.062 (0.023,0.135)	2.318 (2.121,2.512)	0.802 (0.485,1.121)	1.010 (0.785,1.239)	108.4 (103.0,114.4)	
		mid	0.074 (0.019,0.175)	2.078 (1.760,2.385)	0.933 (0.409,1.516)	0.658 (0.063,1.294)		

	lat	0.243 (0.168,0.345)	1.479 (1.422,1.539)	1.224 (1.061,1.368)	0.851 (0.754,0.948)		
M82	pio	-1.403 (-1.718,-1.397)	1.981 (1.873,2.092)	0.555 (0.385,0.731)	0.713 (0.534,0.886)	0.300 (0.284,0.317)	
	mid	-2.011 (-2.323,-1.685)	1.909 (1.757,2.065)	0.757 (0.566,0.948)	0.225 (-0.145,0.600)		
	lat	-1.982 (-2.447,-1.450)	2.133 (1.954,2.335)	0.609 (0.259,0.909)	0.362 (0.054,0.676)		
M83	pio	0.254 (0.175,0.344)	2.082 (1.958,2.192)	0.760 (0.570,0.954)	0.479 (0.311,0.683)	0.209 (0.145,0.295)	2.005 (1.860,2.151)
	mid	0.175 (0.113,0.255)	1.971 (1.791,2.137)	0.446 (0.074,0.872)	0.661 (0.445,0.901)	0.178 (0.103,0.286)	2.124 (1.899,2.362)
	lat	0.150 (0.116,0.214)	2.173 (2.033,2.314)	0.364 (0.117,0.617)	0.558 (0.321,0.798)	0.058 (0.035,0.088)	2.538 (2.362,2.730)

29 Model series predictors: 1 (DBH diameter at breast height), 2 (DBH and SG species' successional group), 3 (DBH and WD wood density), 4
30 (DBH, WD and SG), 5 (DBH and *H* tree height), 6 (DBH, *H* and SG), 7 (DBH, *H* and WD) and 8 (DBH, *H*, SG and WD).


31

32 Table S4. *RMSE* root-mean-square error and bias from Chave et al. (2014)'s pantropical above-ground biomass estimation model (DBH diameter
33 at breast height, *H* tree height [estimated from a DBH: *H* relationship], WD wood density and E environmental stress as predictors) over the six
34 different successional scenarios included in this study.

Scenarios	Bias (%)	<i>RSME</i> (kg)
Early-succession	-30.7	73.3
Mid-succession	-29.2	81.7
Late-succession	-27.9	115.0
Small-sized	-30.7	99.5
Mid-sized	-30.2	116.0
Large-sized	-29.6	130.2
Mean	-29.7	102.6

35


36 **Figures**



37

38 Figure S1. Fit of the two best models parameterized in this study (M33 [DBH diameter at
 39 breast height and WD wood density as predictors] and M43 [DBH, WD and SG species'
 40 successional group]) and the Chave et al. (2014)'s pantropical model (DBH, H tree height
 41 [estimated from a DBH: H relationship], WD and E environmental stress). Note that Chave et
 42 al. (2014)'s model underestimates the biomass of the small-sized trees (DBH < 21 cm). This
 43 pattern was also observed at the landscape-level (Fig. S2).


44



45

46 Figure S2. Relationship between the predicted and observed AGB above-ground biomass for
 47 the six forest scenarios included in this study. Here, the predictions were made by using
 48 Chave et al. (2014)'s pantropical model, which has DBH diameter at breast height, H tree
 49 height [estimated from a DBH: H relationship], WD wood density and E environmental stress
 50 as predictors.


51



52

53 Figure S3. DBH- and height-distribution of trees included in Chave et al. (2014)'s pantropical
 54 model (n = 4004) and those from this study (n = 727). Note the great density/probability
 55 difference for small- and large-sized trees between the two datasets.


56



57

58 Figure S4. Relationship between predictors (DBH diameter at breast height and WD wood
 59 density) and AGB above-ground biomass of the trees included in this study (n = 727) and
 60 those included in Chave et al. (2014)'s pantropical model (n = 4004).

61



62

63 Figure S5. Relationship between predictors (DBH diameter at breast height and WD wood
 64 density) and the observed AGB above-ground biomass of the trees from old-growth and
 65 secondary forests used to parameterize the biomass estimation models fit in this study and
 66 Chave et al. (2014)'s pantropical model. Note that this study included 596 trees (82% of the
 67 total) harvested in secondary forests, while Chave et al. (2014) included only 220 (5% of the
 68 total). In this study, the representative amount of trees and species from different successional
 69 stages of the same forest type allowed for the inclusion of a wide range of tree architecture
 70 and thus allometries (i.e. from slender to stout trunks, from suppressed or emergent late-
 71 successional species typical of old-growth forests, up to competing or canopy/emergent
 72 pioneer species typical of forest gaps.

73

74 References

75 Amaral, D. D., Vieira, I. C. G., Almeida, S. S., Salomão, R. P., Silva, A. S. L. and Jardim, M.
 76 A. G.: Checklist of remnant forest fragments of the metropolitan area of Belém and historical
 77 value of the fragments, State of Pará, Brazil, Bol. Mus. Para. Emílio Goeldi, 4(3), 231–289,
 78 2009.

79 Camargo, J., Ferraz, I., Mesquita, M., Santos, B. and Brum, H.: Guia de Propágulos &
 80 Plântulas da Amazônia vol.1, 1st ed., edited by Authors, INPA, Manaus, 2008.

81 Chambers, J. Q., Robertson, A. L., Carneiro, V. M. C., Lima, A. J. N., Smith, M. L., Plourde,
82 L. C. and Higuchi, N.: Hyperspectral remote detection of niche partitioning among canopy
83 trees driven by blowdown gap disturbances in the Central Amazon, *Oecologia*, 160(1), 107–
84 117, 2009.

85 da Silva, R. P.: Alometria, estoque e dinâmica da biomassa de florestas primárias e
86 secundárias na região de Manaus (AM), PhD Thesis, Universidade Federal do Amazonas,
87 Brazil, available at: http://www.inpa.gov.br/arquivos/Tese_Biomassa_Roseana_Silva.pdf (last
88 access on 20.08.2015), 2007.

89 da Silva, R. P., Santos, J. dos, Tribuzy, E. S., Chambers, J. Q., Nakamura, S. and Higuchi, N.:
90 Diameter increment and growth patterns for individual tree growing in Central Amazon,
91 Brazil, *For. Ecol. Manage.*, 166, 295–301, 2002.

92 Ferraz, I., Leal Filho, N., Imakawa, A., Varela, V. and Piña-Rodríguez, C.: Basic
93 characteristics for a preliminary ecological ranking of timber species occurring in a non-
94 flooded forest in Central Amazonia, *Acta Amaz.*, 34(4), 621–633, 2004.

95 Hallé, F.: Architecture of trees in the rain forest of Morobe District, New Guinea, *Biotropica*,
96 6(1), 43–50, 1974.

97 Hallé, F., Oldeman, R. A. A. and Tomlinson, P. B.: *Tropical trees and forests: an architectural*
98 *analysis*, Springer-Verlag, Berlin, German Federal Republic, 1978.

99 Kammesheidt, L.: Some autecological characteristics of early to late successional tree species
100 in Venezuela, *Acta Oecol.*, 21(1), 37–48, 2000.

101 Lima, A. J. N., Teixeira, L. M., Carneiro, V. M. C., Santos, J. dos and Higuchi, N.: Análise da
102 estrutura e do estoque de fitomassa de uma floresta secundária da região de Manaus AM, dez
103 anos após corte raso seguido de fogo, *Acta Amaz.*, 37(1), 49–54, 2007.

104 Marra, D. M., Chambers, J. Q., Higuchi, N., Trumbore, S. E., Ribeiro, G. H. P. M., Santos, J.
105 dos, Negrón-Juárez, R. ., Reu, B. and Wirth, C.: Large-scale wind disturbances promote tree
106 diversity in a central Amazon forest., *PLoS One*, 9(8), e103711, 2014.

107 Myers, B. J.: Guide to the identification of some tropical rainforest species from large-scale
108 colour aerial photographs, *Aust. For.*, 45(1), 28–41, 1982.

109 Negrón-Juárez, R. I., Chambers, J. Q., Marra, D. M., Ribeiro, G. H. P. M., Rifai, S. W.,
110 Higuchi, N. and Roberts, D.: Detection of subpixel treefall gaps with Landsat imagery in
111 Central Amazon forests, *Remote Sens. Environ.*, 115(12), 3322–3328, 2011.

112 Ribeiro, J. E. L. S., Hopkins, M. J. G., Vicentini, A., Sothers, C. A., Costa, M. A. da S., Brito,
113 J. M. de, Souza, M. A. D. de, Martins, L. H. P., Lohmann, L. G., Assunção, P. A. C. L.,
114 Pereira, E. da C., Silva, C. F. da, Mesquita, M. R. and Procópio, L. C.: Flora da Reserva
115 Ducke: Guia de Identificação das Plantas Vasculares de uma Floresta de Terra-firme na
116 Amazônia Central, INPA, Manaus, 1999.

117 Saldarriaga, J. G., West, D. C., Tharp, M. L. and Uhl, C.: Long-Term Chronossequence of
118 Forest Succession in the Upper Rio Negro of Colombia and Venezuela, *J. Ecol.*, 76, 938–958,
119 1998.

120 Saravy, S., Freitas, P., Lage, M., Leite, S., Braga, L. and Souza, M.: Síndrome de dispeção em
121 estratos arbóreos em um fragmento de floresta ombrófila aberta e densa em Alta Floresta MT,
122 *Rev. Prog. C. Agro-Amb.*, 2(1), 1–12, 2003.

123 Silva Junior, M. and Pereira, B.: Mais 100 Árvores do Cerrado -Matas de Galeria - Guia de
124 Campo, 1st ed., Rede de Sementes do Cerrado, Brasília, 2009.

125 Stefanello, D., Fernandes-Bulhão, C. and Martins, S. V.: Seed dispersal syndromes in three
126 riparian vegetation sites (source, middle and estuary) along the Pindaíba river, Mato Grosso
127 state, Brazil, *Rev. Árvore*, 33, 1051–1061, 2009.

128 Stefanello, D., Ivanauskas, N. M., Martins, S. V., Silva, E. and Kunz, S. H.: Syndromes of
129 diaspores dispersal of species of riparian vegetation of the river from Pacas, Querência - MT,
130 *Acta Amaz.*, 40(1), 141–150, 2010.

131 Stevens, P. F.: Angiosperm Phylogeny Website, version 12, available at:
132 <http://www.mobot.org/MOBOT/research/APweb/> (last access on 20.08.2015), 2012.

133 Swaine, M. D. and Whitmore, T. C.: On the definition of ecological species groups in tropical
134 rain forests, *Vegetatio*, 75, 81–86, 1988.

135 Terborgh, J., Nuñez-Iturri, G., Pitman, N. C. a, Valverde, F. H. C., Alvarez, P., Swamy, V.,
136 Pringle, E. G. and Paine, C. E. T.: Tree recruitment in an empty forest, *Ecology*, 89(6), 1757–
137 1768, 2008.

138 Trichon, V.: Crown typology and the identification of rain forest trees on large-scale aerial
139 photographs, *Plant Ecol.*, 153, 301–312, 2001.