

**Structuring of  
gastropod  
communities**

T. Hauffe et al.

This discussion paper is/has been under review for the journal Biogeosciences (BG).  
Please refer to the corresponding final paper in BG if available.

# Gastropod diversification and community structuring processes in ancient Lake Ohrid: a metacommunity speciation perspective

T. Hauffe, C. Albrecht, and T. Wilke

Department of Animal Ecology and Systematics, Justus Liebig University, Giessen, Germany

Received: 31 August 2015 – Accepted: 1 September 2015 – Published: 29 September 2015

Correspondence to: T. Hauffe (torsten.hauffe@gmail.com)

Published by Copernicus Publications on behalf of the European Geosciences Union.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



## Abstract

The Balkan Lake Ohrid is the oldest and most speciose freshwater lacustrine system in Europe. However, it remains unclear whether the diversification of its endemic taxa is mainly driven by neutral processes, environmental factors, or species interactions.

This calls for a holistic perspective involving both evolutionary processes and ecological dynamics. Such a unifying framework – the metacommunity speciation model – considers how community assembly affects diversification and vice versa by assessing the relative contribution of the three main community assembly processes, dispersal limitation, environmental filtering, and species interaction.

The current study therefore used the species-rich model taxon Gastropoda to assess how extant communities in Lake Ohrid are structured by performing process based metacommunity analyses. Specifically, the study aimed at (i) identifying the relative importance of the three community assembly processes and (ii) to test whether the importance of these individual processes changes gradually with lake depth or whether they are distinctively related to eco-zones.

Based on specific simulation steps for each of the three processes, it could be demonstrated that dispersal limitation had the strongest influence on gastropod community structures in Lake Ohrid. However, it was not the exclusive assembly process but acted together with the other two processes – environmental filtering, and species interaction. In fact, the relative importance of the three community assembly processes varied both with lake depth and eco-zones, though the processes were better predicted by the latter.

The study thus corroborated the high importance of dispersal limitation for both maintaining species richness in Lake Ohrid (through its impact on community structure) and generating endemic biodiversity (via its influence on diversification processes). However, according to the metacommunity speciation model, the inferred importance of environmental filtering and biotic interaction also suggests a small but significant influence of ecological speciation. These findings contribute to the main goal of the SCOPSCO

## Structuring of gastropod communities

T. Hauffe et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



initiative – inferring the drivers of biotic evolution – and might provide an integrative perspective on biological and limnological dynamics in ancient Lake Ohrid.

## 1 Introduction

Ancient Lake Ohrid on the Balkan Peninsula is the oldest and most speciose fresh-water lacustrine system in Europe (Albrecht and Wilke, 2008; Neubauer et al., 2015). Recently, an International Continental Scientific Drilling Program (ICDP; Wagner et al., 2014) has been conducted in the lake within the research initiative Scientific Collaboration on Past Speciation Conditions (SCOPSCO). One of the major goals of this collaborative project is to infer the drivers of speciation by linking the geological and biotic evolution of the lake over space and time. In order to reach this goal, geological, limnological, and paleontological data from sediment cores are being complemented with phylogenetic and molecular clock data derived from DNA information of extant endemic species.

Though the sediment and molecular data are still being analyzed, first results of the SCOPSCO campaign indicate that climatic, geological, and/or environmental changes over time may have had little direct effect on speciation and extinction processes in endemic biota (Föller et al., 2015). Instead, such changes likely affected abundances of paleospecies as well as species compositions (Albrecht et al., 2010; Belmecheri et al., 2009; Jovanovska et al., 2015; Reed et al., 2010; Wagner et al., 2014), which, in turn may have affected speciation processes. Similar effects could be shown for horizontal and vertical (bathymetric) environmental gradients. They may not only have shaped the structure of extant Ohrid communities (Hauffe et al., 2011; Lorenschat and Schwalb, 2013), but also indirectly fostered species divergence (Schreiber et al., 2011; Wysocka et al., 2008). Speciation, and extinction processes, in turn, also affect the spatial structure of extant communities (Hauffe et al., 2011).

These results question the direct effect of climatic, geological, and/or environmental factors on diversification and call for a holistic perspective involving evolutionary pro-

## Structuring of gastropod communities

T. Hauffe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



cesses (e.g., speciation) and ecological dynamics (e.g., dispersal limitation, species interaction) (reviewed in Hubert et al., 2015).

Such a mechanistic and unifying eco-evolutionary framework – the metacommunity speciation model – was recently proposed by Hubert et al. (2015). It considers how a metacommunity (i.e., a network of local communities linked by dispersal of multiple interacting species; Leibold et al., 2004) is affected by speciation and vice versa.

In principle, a local community is structured by three non-exclusive consecutive processes (e.g., Leibold et al., 2004; van der Plas et al., 2015): (1) random dispersal of individuals to a patch previously unoccupied by the species, (2) environmental filtering that permits the establishment of species whose ecological niche fits the local environmental conditions, and (3) species interaction including competitive exclusion among resident and colonizing species.

From a theoretical point of view, all of these three processes might affect speciation (Hubert et al., 2015). However, their individual contribution and spatial dynamics in Lake Ohrid remain little understood. The latter is particularly true for the bathymetric structure of Lake Ohrid. The lake has a large depth but also a distinct horizontal eco-zonation (reviewed in Albrecht and Wilke, 2008). This leads to the question whether the individual importance of the three processes mentioned above gradually changes with lake depth or whether there are abrupt changes related to transitions between eco-zones. Conceptually, a differentiation between gradual and abrupt changes in structuring processes among eco-zones is of high importance for understanding how limnological complexity and stability affect assembly and diversification processes.

A suitable model taxon to study structuring processes are the Gastropoda as they represent the largest animal taxon in Lake Ohrid with 74 described species, 56 of which being endemic to the lake and its catchment (Albrecht et al., 2014; Albrecht and Wilke, 2008; Hauffe et al., 2011; Radoman, 1983, 1985). In the current study we therefore focus on the question how extant gastropod communities are assembled by performing process based metacommunity analyses. Our working hypothesis is that, in general, dispersal limitation plays a crucial role in structuring communities.

## Structuring of gastropod communities

T. Hauffe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





Though phylogenetic relationships among endemic gastropod taxa of Lake Ohrid are increasingly well characterized, a complete picture is still lacking. For critical taxa we here used the taxonomic assignments of Bodon et al. (2001), Hubendick and Radoman (1959), and Radoman (1983). Taxa were classified in five hierarchical levels: species, 5 genus, species flock, family, and superfamily. Note that we included the level “species flock” for monophyletic groups of endemic species that may comprise more than one genus (Albrecht et al., 2008; Föller et al., 2015; Wilke et al., 2007). Species flocks likely evolved intralacustrine and are common in Lake Ohrid.

## 2.2 Simulation of individual community assembly processes

10 Van der Plas et al. (2015) suggested a simulation method of stepwise community assembly (STEPCAM). It prunes the regional species pool to the size of an observed community by sequentially removing species according to specific simulation processes for dispersal limitation, environmental filtering, and biotic interaction.

STEPCAM simulates dispersal limitation by randomly removing species with a probability inversely related to their relative frequency in the regional species pool. Dispersal 15 limited species are therefore less likely present in the simulated community (van der Plas et al., 2015).

Environmental filtering and species interaction depend on the ecology of the species (i.e., its niche). These ecological requirements are assumed to be reflected by morphological 20 features because the latter are often under selection. Environmental filtering therefore predicts that species of a community show similar morphological traits because environmental conditions only allow the establishment of species that are adapted to the specific niche (Webb et al., 2002). Environmental filtering is simulated by pruning species with trait values most distant to the trait mean of the simulated 25 community.

Community composition driven by species interaction assumes that species with similar niches do not co-exist because they share the same resources, predators,

## Structuring of gastropod communities

T. Hauffe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



or pathogens (Webb et al., 2002). Therefore, this process is simulated by removing species from the regional species pool showing most similar traits.

We here extended this original STEPCAM approach, which only considers continuous species traits, now to allow a mixture of continuous and categorical species traits based on pairwise distance matrices between species. This generalization also permits the use of phylogenetic or taxonomic distances, which are typically strongly correlated with morphological divergence among species.

The number of species removed in each of the three simulation steps can be used as a direct measure for the relative importance of the three structuring processes (see specific aim 1).

Following van der Plas et al. (2015), we here utilized Approximate Bayesian Computation within a Sequential Monte Carlo framework (ABC-SMC) for inferring the number of species removals needed per simulation step, which results in a simulated community that approximates the observed community. The similarity between simulated and observed communities ( $Fit_{total}$ ; van der Plas et al., 2015) is the sum of absolute differences between four indices (i.e., richness, evenness, divergence, and mean distance to the community centroid) based on traits (Villéger et al., 2008) or relatedness (Helmus et al., 2007).

However, calculating  $Fit_{total}$  may be problematic because different indices of community structure are bounded between 0 and 1, whereas others lack an upper limit. Van der Plas et al. (2015) standardized the diversity indices by dividing them by the standard deviation (SD) of the respective index, calculated for all observed communities. However, for large differences in species richness or few sampled communities, this approach can be biased because some indices are not independent of species richness (Villéger et al., 2008) and the SD is less precisely estimated due to the small sample size. Hence, we estimated all SDs using an initial simulation of 500 communities based on observed values for species richness.

Our distance-based community simulations built on the STEPCAM package v1.0 (Janzen and van der Plas, 2014) for the R v3.1.2 statistical environment (R Core Team,

**BGD**

12, 16081–16103, 2015

## Structuring of gastropod communities

T. Hauffe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2015). However, the algorithm was improved for runtime and a fully reproducible analysis including the taxonomic distance and the community matrix is available in the Supplement.

We calculated pairwise taxonomic distances between the 64 collected gastropod species using the taxonomic classification with five hierarchical levels. This method has been described by Clarke and Warwick (1998) and is implemented in the R package `vegan` v2.2-1 (Oksanen et al., 2015). The STEPCAM target for the assembly simulation for each of our 264 sampled communities was a final acceptance rate of 1 in 10 000.

### 2.3 Testing for gradual vs. distinct changes among eco-zones

For testing whether the importance of individual structuring processes changes gradually with lake depth or whether they are distinctively related to eco-zones (see specific aim 2), reliable information for the spatial distribution of these zones is necessary.

Previous assessments of the vertical distribution of eco-zones in Lake Ohrid were based on analyses of habitat types, limnological features, and/or typical species assemblages (Albrecht and Wilke, 2008; Hauffe et al., 2011). In contrast, only a rough approximation of the horizontal dimension exists so far (Stanković, 1960).

Therefore, we first performed an automatic detection of eco-zones using a spatially constrained clustering of ecological networks (Miele et al., 2014). The analysis, implemented in the R package `geoclust` v0.2.0 (Miele et al., 2014), is based on the similarity of gastropod communities and their geographical adjacency in horizontal ( $X$  and  $Y$ ) and vertical ( $Z$ ) dimensions. Composition similarity was calculated using Bray–Curtis distances and the `vegan` v2.2-1 package. Similarities were then arc–sin transformed to ensure normality, as required for ecological networks. The three-dimensional adjacency network was based on (i) vertical (bathymetric) connectivity, and (ii) the cost-distance between collecting points along isobaths. The latter approach had to be used because intralacustrine dispersal of benthic organisms is not well approximated by straight line distances (Heino et al., 2015a). We explored the robustness of our analysis by varying the number of connected neighbors from 15 to 264 with an increment of 10, ensuring

## Structuring of gastropod communities

T. Hauffe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





a fully linked spatial network. For the vertical scale, a connectivity threshold of 8.2 m was determined by a multivariate community correlogram with 9999 permutations in vegan, showing that similarity between gastropod communities decreases significantly if their bathymetric distance exceeds this threshold.

5 In a final step, we used a Bayesian generalized linear model (BGLM; R package MCMCglmm v2.21; Hadfield, 2010) to test whether our multivariate response, i.e., the relative importance of the three community assembly processes, was better predicted by either lake depth or delineated eco-zones (see specific aim 2). As decision criterion between the two competing BGLMs, we utilized the deviance information criterion (DIC), a Bayesian measure of model fit (Spiegelhalter et al., 2002). In both BGLMs, we included species richness and its interaction with lake depth or eco-zones as additional predictor because process-importance may co-vary with species richness. Prior to model fitting, normality of the [0, 1] scaled multivariate response was improved by an additive planar transformation, implemented in the compositions v1.4-1 package (van den Boogaart et al., 2014) for R.

### 3 Results

#### 3.1 Relative importance of the three community assembly processes

Using our ABC-SMC-based STEPCAM approach, we obtained a posterior distribution for the relative contribution of the three community assembly processes (dispersal limitation, environmental filtering, and biotic interaction) in shaping the lake's gastropod compositions. Averaging the posterior distribution over all 264 gastropod communities, dispersal limitation achieved the highest relative importance with 78.7% (95% bootstrapped confidence interval (CI): 75.5–81.7), followed by environmental filtering (mean 12.2%; CI: 9.9–14.6), and species interaction (mean 9.1%; CI: 7.9–10.5).

**BGD**

12, 16081–16103, 2015

## Structuring of gastropod communities

T. Hauffe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 3.2 Testing for gradual vs. distinct changes among eco-zones

Prior to testing whether the importance of the three individual processes changes gradually with lake depth or whether they are related to distinct eco-zones, we performed an automatic detection of eco-zones using a spatially constrained clustering of ecological networks. Applying a range of one to ten groups and different numbers of neighbors in the spatial network, our delineation of eco-zones consistently revealed that seven groups of gastropod communities showed the highest likelihood. These communities relate to seven eco-zones (southeastern (SE) upper littoral 1–3, non-SE upper littoral, lower littoral, upper sublittoral, lower sublittoral). The spatial distribution (horizontal and vertical) of these eco-zones based on 65 neighbors is shown in Fig. 1 (zone terminology roughly follows Stanković, 1960). Clusters of communities across the different parameter settings were highly consistent, indicating a robust delineation of eco-zones.

Finally, we used two BGLMs to test whether the relative importance of the three community assembly processes was better predicted by either lake depth or delineated eco-zones. In general, both assumptions were supported. However, according to the rule of thumb of Bayesian model selection (Bolker, 2008), the difference in DIC was greater than 10 and thus decisively favored eco-zones over lake depth (DIC values of –740 vs. –662, respectively). For a visual presentation of the eco-zone dependent relative importance of dispersal limitation, environmental filtering, and biotic interaction see Fig. 2.

## 4 Discussion

In the current study, we addressed the question of how extant gastropod communities are assembled by performing a process-based metacommunity analysis. More specifically, we attempted to identify the relative importance of three previously suggested community assembly processes (dispersal limitation, environmental filtering, and species interaction) and tested whether their importance changes gradually with

### Structuring of gastropod communities

T. Hauffe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Structuring of gastropod communities

T. Hauffe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



lake depth or is distinctively related to eco-zones. Given that geographical isolation has been stressed as important driver for speciation in Lake Ohrid (reviewed in Albrecht and Wilke, 2008) and adhering to the metacommunity speciation model, our working hypothesis was that dispersal limitation plays a crucial role in structuring communities. We, indeed, could show that dispersal limitation had the strongest influence on gastropod community structures in Lake Ohrid. However, it was not the exclusive assembly process but acted together with the other two processes – environmental filtering, and species interaction (Fig. 2). In fact, the relative importance of the three community assembly processes varied both with lake depth and eco-zones (Fig. 2), though the processes were better predicted by delineated eco-zones.

In the following, these principal findings are interpreted in the context of how ecosystem features influence community assembly. We also discuss the implications of our three community processes for patterns and processes of speciation in endemic gastropods of Lake Ohrid. As structuring processes appear to be strongly related to eco-zones, we first address the result of our automatic eco-zones identification.

### 4.1 The eco-zones of Lake Ohrid

Previously, six horizontal and four vertical eco-zones have been suggested for Lake Ohrid (Stanković, 1960; Albrecht and Wilke, 2008). However, as they were largely based on an a priori differentiation into horizontal and vertical features, we here automatically delineated eco-zones using a three dimensional approach (i.e., involving the three spatial axes  $X$ ,  $Y$ , and  $Z$ ).

The zonation inferred (Fig. 1) resembles, in large parts, the one previously proposed. However, it appears to be slightly more complex in terms of horizontal and vertical structuring. Moreover, the zones show a considerable degree of overlapping, reflecting the patchy nature of some habitats in the lake (see Albrecht and Wilke, 2008). The strong vertical structure largely corresponds to vertically arranged biotopes in Lake Ohrid, such as the “*Chara* belt” in the lower littoral and the “shell zone” in the upper sublittoral (Albrecht and Wilke, 2008). Note that a potential fifth vertical zone, the pro-

fundal, could not be identified in this study as the only species of gastropod previously reported from there has not been found in years; see Hauffe et al., 2011).

Interestingly, we do not see any significant horizontal structuring within and among the deeper vertical zones. However, within the upper littoral (i.e., a zone typically extending from 0 to 3 m water depth), a strong horizontal structure is evident. One eco-zone (“non-SE upper littoral zone”) comprises all shallow water gastropod communities from the northeastern, northern, western, and southwestern parts of the lake. Three additional eco-zones can be found in the southeastern upper littoral (“SE upper littoral zones 1–3”). This area, comprising major spring fields within the lake, has long been known for its high degree of point-endemism (Albrecht et al., 2006; Albrecht and Wilke, 2008; also see Hauffe et al., 2011 Fig. 3a for compositional similarities among these communities).

The emergent pattern of partly overlapping horizontal/vertical eco-zones, indeed, confirms that the gastropods of Lake Ohrid adhere to the metacommunity concept.

## 4.2 Relative importance of community assembly processes

Our results suggest that gastropod communities of Lake Ohrid are substantially structured by the dispersal limitation process (i.e., restricted specimen exchange; mean importance across all communities ~ 80 %; see Fig. 2). Environmental filtering and biotic interaction were of lower importance (12 and 8 %, respectively). However, the individual importance of these three processes depends on lake depth and even more on individual eco-zones (Fig. 2). The importance of dispersal limitation is, for example, relatively low in the SE upper littoral 1 (i.e., the shallowest and most species rich zone in Lake Ohrid) and in the lower sublittoral (i.e., a deep and relatively species poor zone in Lake Ohrid). The opposite is true for environmental filtering and, to a lesser extent, for species interaction (see Fig. 2). This interesting relationship shows that the relative importance of the three community structuring processes does not depend on species richness or lake depth per se. The latter might also explain why eco-zones (i.e., three spatial axes) have a higher explanatory power than lake depth (only one spatial axis).

## Structuring of gastropod communities

T. Hauffe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Structuring of gastropod communities

T. Hauffe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



5 Interestingly, the clearly demonstrated high importance of dispersal limitation for both maintaining species richness in Lake Ohrid (through its impact on community structure) and generating endemic biodiversity (via its influence on diversification processes) stands in contrast to some other studies on Lake Ohrid biota. In fact, previous investigations on gastropod (Schneider et al., 2014), ostracod (Lorenschat et al., 2014), diatom (Reed et al., 2010), and fungal communities (Čomić et al., 2010) suggested a strong environmental control (= environmental filtering) of species composition. However, this is not surprising as other processes were not considered in these studies.

10 In studies that did consider more than one process, the picture was more complex. This concerns gastropod communities in Lake Ohrid (Hauffe et al., 2011) and in other ancient lakes such as Lake Tanganyika (Meyer et al., 2011), but also communities in freshwater systems in general (Heino et al., 2015b).

15 As the importance of assembly processes often depends on spatial scale (Meyer et al., 2011) and resolution (Münkemüller et al., 2014), the sensitivity of the STEPCAM approach to spatial data aggregation should generally be considered. However, as the spatial resolution in our study is relatively high and as dispersal rates of freshwater gastropods are generally low (Kappes and Haase, 2012; Michel et al., 2007), we here refrained from data aggregation.

### 4.3 Metacommunity implications for gastropod speciation

20 If we accept that species divergence and community assembly can be unified in an eco-evolutionary framework because both are based on similar processes (Hubert et al., 2015), then the here inferred importance of processes allows for a fresh look on the diversification mode of Lake Ohrid's endemic taxa. Research on species divergence, in general, and in ancient lakes in particular, has traditionally focused on geographic speciation (Cristescu et al., 2010; Schluter, 2001; also see Albrecht and Wilke, 2008 for a review on geographic speciation modes in Lake Ohrid). Adaptive diversification along environmental gradients, however, highlights the additional role of ecology in driving species divergence in ancient lakes (Schön and Martens, 2004).

## Structuring of gastropod communities

T. Hauffe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Our results support this perspective of non-exclusive geographic and ecological speciation in Lake Ohrid. Though the inferred high importance of dispersal limitations in structuring gastropod communities implies a dominant role of geographic speciation in the lake (also see Albrecht et al., 2006; Kilikowska et al., 2013; Trajanovski et al., 2010; Wysocka et al., 2014), the inferred importance of environmental filtering and species interaction suggests a small but significant influence of ecological speciation.

Interestingly, the relative contribution of geographical speciation (i.e., via dispersal limitation) and ecological speciation (i.e., via environmental filtering and biotic interaction) varies among eco-zones. However, as the total number of eco-zones inferred is relatively low, detailed studies that aim at linking the underlying biotic and abiotic drivers of community processes with modes of speciation are difficult to conduct.

Nonetheless, the fact that ecological speciation appears to be more important both in the SE upper littoral 1 (i.e., a shallow and species-rich zone) and in the lower sublittoral (i.e., a deep and species-poor zone) potentially implies a bimodal contribution of ecological speciation in generating biodiversity over time.

#### 4.4 Methodological implications and outlook

For many questions concerning processes in eco-evolutionary biology, such as dispersal or selection dependent rates of species diversification (Pigot et al., 2010; Rosindell et al., 2015), no analytical solutions are (yet) available. However, the outcome of processes can be simulated by utilizing a wide range of parameter values representing them. The set of parameters resulting in a simulated pattern that resembles the observed one best is then assumed to approximate the processes in question. Our newly developed distance-based STEPCAM approach is a relatively simple model that could be easily extended by making, for instance, dispersal limitation spatially or temporally explicit, but at the risk of overparameterization.

Moreover, we suggest further exploration of the STEPCAM approach using different taxa in Lake Ohrid, for instance the species rich-group of diatoms. A combined study of extant and paleo-communities derived from multiple sediment cores would allow

inferring the relative importance of dispersal limitations, environmental filtering, and species interaction over space and time.

## 5 Conclusions

Our study builds on a prediction of the metacommunity speciation model: if geographic speciation is the prevailing mode of species divergence, as suggested by many evolutionary studies, then dispersal limitation should also structure communities. The results of our analysis corroborate this hypothesis for gastropods in Lake Ohrid. However, the inferred importance of environmental filtering and biotic interaction in community assembly also suggests a small but significant influence of ecological speciation on diversification dynamics; an aspect which previously could not be quantified. Moreover, we show that limnological characteristics such as lake depth and eco-zones do influence the relative importance of assembly processes, whereas former studies could only show that these limnological features give rise to distinct species compositions. These findings contribute to the main goal of the SCOPSCO initiative – inferring the drivers of biotic evolution – and provide an integrative perspective on biological and geological dynamics in ancient Lake Ohrid.

Moreover, our novel distance-based STEPCAM complements the growing eco-evolutionary toolbox that aims at disentangles patterns and processes.

**The Supplement related to this article is available online at doi:10.5194/bgd-12-16081-2015-supplement.**

*Author contributions.* T. Hauffe conceived the study. C. Albrecht and T. Hauffe collected and identified specimens. T. Hauffe performed the analyses. The manuscript was written by T. Hauffe and T. Wilke with contributions from all co-authors. All authors gave final approval for publication.

### Structuring of gastropod communities

T. Hauffe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



*Acknowledgements.* We are grateful to our colleagues at the Hydrobiological Institute Ohrid and the Albanian authorities for their valuable support. Sampling was conducted in accordance with national and provincial regulations. D. Georgiev (Ohrid) supported our field work with his vast local expertise. Fons van der Plas and Thijs Janzen are acknowledged for their enormous help and discussion of the STEPCAM approach. This work was supported by DFG grants to TW (WI 1902/13) and CA (AL 1076/9).

## References

- Albrecht, C. and Wilke, T.: Ancient Lake Ohrid: biodiversity and evolution, *Hydrobiologia*, 615, 103–140, doi:10.1007/s10750-008-9558-y, 2008.
- Albrecht, C., Trajanovski, S., Kuhn, K., Streit, B., and Wilke, T.: Rapid evolution of an ancient lake species flock: freshwater limpets (Gastropoda: *Ancylidae*) in the Balkan Lake Ohrid, *Org. Divers. Evol.*, 6, 294–307, doi:10.1016/j.ode.2005.12.003, 2006.
- Albrecht, C., Wolff, C., Glöer, P., and Wilke, T.: Concurrent evolution of ancient sister lakes and sister species: the freshwater gastropod genus *Radix* in lakes Ohrid and Prespa, *Hydrobiologia*, 615, 157–167, doi:10.1007/s10750-008-9555-1, 2008.
- Albrecht, C., Vogel, H., Hauffe, T., and Wilke, T.: Sediment core fossils in ancient Lake Ohrid: testing for faunal change since the Last Interglacial, *Biogeosciences*, 7, 3435–3446, doi:10.5194/bg-7-3435-2010, 2010.
- Albrecht, C., Föller, K., Clewing, C., Hauffe, T., and Wilke, T.: Invaders vs. endemics: alien gastropod species in ancient Lake Ohrid, *Hydrobiologia*, 739, 163–174, doi:10.1007/s10750-013-1724-1, 2014.
- Belmecheri, S., Namiotko, T., Robert, C., von Grafenstein, U., and Danielopol, D. L.: Climate controlled ostracod preservation in Lake Ohrid (Albania, Macedonia), *Palaeogeogr. Palaeoclimatol.*, 277, 236–245, doi:10.1016/j.palaeo.2009.04.013, 2009.
- Bodon, M., Manganelli, G., and Giusti, F.: A survey of the European valvatiform hydrobiid genera with special reference to *Hauffenia Pollonera*, 1898 (Gastropoda: *Hydrobiidae*), *Malacologia*, 43, 103–215, 2001.
- Bolker, B. M.: *Ecological Models and Data in R*, Princeton University Press, Princeton, NJ, 2008.

## Structuring of gastropod communities

T. Hauffe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





## Structuring of gastropod communities

T. Hauffe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Clarke, K. R. and Warwick, R. M.: A taxonomic distinctness index and its statistical properties, *J. Appl. Ecol.*, 35, 523–531, doi:10.1046/j.1365-2664.1998.3540523.x, 1998.
- Čomić, L., Rankovic, B., Novevska, V., and Ostojic, A.: Diversity and dynamics of the fungal community in Lake Ohrid, *Aquat. Biol.*, 9, 169–176, doi:10.3354/ab00248, 2010.
- 5 Cristescu, M. E., Adamowicz, S. J., Vaillant, J. J., and Haffner, D. G.: Ancient lakes revisited: from the ecology to the genetics of speciation, *Mol. Ecol.*, 19, 4837–4851, doi:10.1111/j.1365-294X.2010.04832.x, 2010.
- Föller, K., Stelbrink, B., Hauffe, T., Albrecht, C., and Wilke, T.: Constant diversification rates of endemic gastropods in ancient Lake Ohrid: ecosystem resilience likely buffers environmental fluctuations, *Biogeosciences Discuss.*, 12, 14271–14302, doi:10.5194/bgd-12-14271-2015, 2015.
- 10 Hadfield, J. D.: MCMC methods for multi-response generalized linear mixed models: the MCMCglmm R package, *J. Stat. Softw.*, 33, 1–22, 2010.
- Hauffe, T., Albrecht, C., Schreiber, K., Birkhofer, K., Trajanovski, S., and Wilke, T.: Spatially explicit analysis of gastropod biodiversity in ancient Lake Ohrid, *Biogeosciences*, 8, 175–188, doi:10.5194/bg-8-175-2011, 2011.
- 15 Heino, J., Melo, A. S., Siqueira, T., Soininen, J., Valanko, S., and Bini, L. M.: Metacommu- nity organisation, spatial extent and dispersal in aquatic systems: patterns, processes and prospects, *Freshwater Biol.*, 60, 845–869, doi:10.1111/fwb.12533, 2015a.
- 20 Heino, J., Soininen, J., Alahuhta, J., Lappalainen, J., and Virtanen, R.: A comparative analysis of metacomunity types in the freshwater realm, *Ecol. Evol.*, 5, 1525–1537, doi:10.1002/ece3.1460, 2015b.
- Helmus, M. R., Bland, T. J., Williams, C. K., and Ives, A. R.: Phylogenetic measures of biodi- versity, *Am. Nat.*, 169, E68–E83, doi:10.1086/511334, 2007.
- 25 Hubendick, B. and Radoman, P.: Studies on the *Gyraulus* species of Lake Ochrid, *Morphol. Ark. Zool.*, 12, 223–243, 1959.
- Hubert, N., Calcagno, V., Etienne, R. S., and Mouquet, N.: Metacomunity specia- tion models and their implications for diversification theory, *Ecol. Lett.*, 18, 864–881, doi:10.1111/ele.12458, 2015.
- 30 Janzen, T. and van der Plas, F.: STEPCAM: ABC-SMC inference of the STEPCAM model, available at: <http://CRAN.R-project.org/package=STEPCAM>, last access: 16 June 2015.
- Jovanovska, E., Cvetkoska, A., Hauffe, T., Levkov, Z., Wagner, B., Sulpizio, R., Francke, A., Albrecht, C., and Wilke, T.: Differential resilience of ancient sister lakes Ohrid and Prespa to

## Structuring of gastropod communities

T. Hauffe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



environmental disturbances during the Late Pleistocene, Biogeosciences Discuss., in press, 2015.

Kappes, H. and Haase, P.: Slow, but steady: dispersal of freshwater molluscs, *Aquat. Sci.*, 74, 1–14, doi:10.1007/s00027-011-0187-6, 2012.

5 Kilikowska, A., Wysocka, A., Burzynski, A., Kostoski, G., Rychlińska, J., and Sell, J.: Patterns of genetic differentiation and population history of endemic isopods (Asellidae) from ancient Lake Ohrid: combining allozyme and mtDNA data, *Cent. Eur. J. Biol.*, 8, 854–875, doi:10.2478/s11535-013-0204-y, 2013.

10 Leibold, M. A., Holyoak, M., Mouquet, N., Amarasekare, P., Chase, J. M., Hoopes, M. F., Holt, R. D., Shurin, J. B., Law, R., Tilman, D., Loreau, M., and Gonzalez, A.: The meta-community concept: a framework for multi-scale community ecology, *Ecol. Lett.*, 7, 601–613, doi:10.1111/j.1461-0248.2004.00608.x, 2004.

Lorenschat, J. and Schwalb, A.: Autecology of the extant ostracod fauna of Lake Ohrid and adjacent waters – a key to paleoenvironmental reconstruction, *Belg. J. Zool.*, 143, 42–68, 2013.

15 Lorenschat, J., Pérez, L., Correa-Metrio, A., Brenner, M., von Bramann, U., and Schwalb, A.: Diversity and spatial distribution of extant freshwater ostracodes (Crustacea) in ancient Lake Ohrid (Macedonia/Albania), *Diversity*, 6, 524–550, doi:10.3390/d6030524, 2014.

20 Meyer, J. R., Michel, E., McIntyre, P. B., Huntington, B. E., Long, D. J., and Lara, G.: Scale-dependent processes of community assembly in an African rift lake, *Freshwater Biol.*, 56, 2082–2093, doi:10.1111/j.1365-2427.2011.02639.x, 2011.

Michel, E., McIntyre, P. B., and Chan, J.: A snail's space sets a snail's pace: movement rates of *Lavigeria* gastropods in Lake Tanganyika, East Africa, *J. Mollus. Stud.*, 73, 195–198, doi:10.1093/mollus/eym013, 2007.

25 Miele, V., Picard, F., and Dray, S.: Spatially constrained clustering of ecological networks, *Methods Ecol. Evol.*, 5, 771–779, doi:10.1111/2041-210X.12208, 2014.

Münkemüller, T., Gallien, L., Lavergne, S., Renaud, J., Roquet, C., Abdulhak, S., Dullinger, S., Garraud, L., Guisan, A., Lenoir, J., Svenning, J.-C., Van Es, J., Vittoz, P., Willner, W., Wohlgemuth, T., Zimmermann, N. E., and Thuiller, W.: Scale decisions can reverse conclusions on community assembly processes, *Global Ecol. Biogeogr.*, 23, 620–632, doi:10.1111/geb.12137, 2014.

## Structuring of gastropod communities

T. Hauffe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Neubauer, T. A., Harzhauser, M., Georgopoulou, E., Kroh, A., and Mandic, O.: Tectonics, climate, and the rise and demise of continental aquatic species richness hotspots, *P. Natl. Acad. Sci. USA*, 112, 11478–11483, doi:10.1073/pnas.1503992112, 2015.
- Oksanen, J., Blanchet, F. G., Kindt, R., Legendre, P., Minchin, P. R., O'Hara, R. B., Simpson, G. L., Solymos, P., Stevens, M. H. H., and Wagner, H.: *vegan: Community Ecology Package*, available at: <http://CRAN.R-project.org/package=vegan>, last access: 12 May 2015.
- Pigot, A. L., Phillimore, A. B., Owens, I. P. F., and Orme, C. D. L.: The shape and temporal dynamics of phylogenetic trees arising from geographic speciation, *Syst. Biol.*, 59, 1–14, doi:10.1093/sysbio/syq058, 2010.
- R Core Team: *R: A language and environment for statistical computing*, Vienna, Austria, available at: <http://www.R-project.org/>, last access: 9 March 2015.
- Radoman, P.: *Hydrobioidea a Superfamily of Prosobranchia (Gastropoda)*, I Systematics, Serbian Academy of Sciences and Arts, Belgrade, 1983.
- Radoman, P.: *Hydrobioidea a Superfamily of Prosobranchia (Gastropoda)*, II Origin, Zoogeography, Evolution in the Balkans and Asia Minor, Serbian Academy of Sciences and Arts, Belgrade, 1985.
- Reed, J. M., Cvetkoska, A., Levkov, Z., Vogel, H., and Wagner, B.: The last glacial-interglacial cycle in Lake Ohrid (Macedonia/Albania): testing diatom response to climate, *Biogeosciences*, 7, 3083–3094, doi:10.5194/bg-7-3083-2010, 2010.
- Rosindell, J., Harmon, L. J., and Etienne, R. S.: Unifying ecology and macroevolution with individual-based theory, *Ecol. Lett.*, 18, 472–482, doi:10.1111/ele.12430, 2015.
- Schluter, D.: Ecology and the origin of species, *Trends Ecol. Evol.*, 16, 372–380, doi:10.1016/S0169-5347(01)02198-X, 2001.
- Schneider, S. C., Cara, M., Eriksen, T. E., Budzacoska Goreska, B., Imeri, A., Kupe, L., Lokoska, T., Patceva, S., Trajanovska, S., Trajanovski, S., Talevska, M., and Veljanoska Sarafiloska, E.: Eutrophication impacts littoral biota in Lake Ohrid while water phosphorus concentrations are low, *Limnologia*, 44, 90–97, doi:10.1016/j.limno.2013.09.002, 2014.
- Schön, I. and Martens, K.: Adaptive, pre-adaptive and non-adaptive components of radiations in ancient lakes: a review, *Org. Divers. Evol.*, 4, 137–156, doi:10.1016/j.ode.2004.03.001, 2004.
- Schreiber, K., Hauffe, T., Albrecht, C., and Wilke, T.: The role of barriers and gradients in differentiation processes of pyrgulinid microgastropods of Lake Ohrid, *Hydrobiologia*, 682, 61–73, doi:10.1007/s10750-011-0864-4, 2011.

## Structuring of gastropod communities

T. Hauffe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Spiegelhalter, D. J., Best, N. G., Carlin, B. P., and Van Der Linde, A.: Bayesian measures of model complexity and fit, *J. Roy. Stat. Soc. B*, 64, 583–639, doi:10.1111/1467-9868.00353, 2002.

Stanković, S.: The Balkan Lake Ohrid and its living world, *Monographiae biologicae*, Vol. IX., edited by: Junk, W., Bodenheimer, F. S., and Weisbach, W. W., Uitgeverij Dr. W. Junk, Den Haag, 358 pp., 1960.

Trajanovski, S., Albrecht, C., Schreiber, K., Schultheiß, R., Stadler, T., Benke, M., and Wilke, T.: Testing the spatial and temporal framework of speciation in an ancient lake species flock: the leech genus *Dina* (Hirudinea: Erpobdellidae) in Lake Ohrid, *Biogeosciences*, 7, 3387–3402, doi:10.5194/bg-7-3387-2010, 2010.

van den Boogaart, K. G., Tolosana, R., and Bren, M.: compositions: compositional data analysis, available at: <http://CRAN.R-project.org/package=compositions>, last access: 16 June 2015.

van der Plas, F., Janzen, T., Ordonez, A., Fokkema, W., Reinders, J., Etienne, R. S., and Olf, H.: A new modeling approach estimates the relative importance of different community assembly processes, *Ecology*, 96, 1502–1515, doi:10.1890/14-0454.1, 2015.

Villéger, S., Mason, N. W. H., and Mouillot, D.: New multidimensional functional diversity indices for a multifaceted framework in functional ecology, *Ecology*, 89, 2290–2301, doi:10.1890/07-1206.1, 2008.

Wagner, B., Wilke, T., Krastel, S., Zanchetta, G., Sulpizio, R., Reicherter, K., Leng, M. J., Grazhdani, A., Trajanovski, S., Francke, A., Lindhorst, K., Levkov, Z., Cvetkoska, A., Reed, J. M., Zhang, X., Lacey, J. H., Wonik, T., Baumgarten, H., and Vogel, H.: The SCOPSCO drilling project recovers more than 1.2 million years of history from Lake Ohrid, *Sci. Drill.*, 17, 19–29, doi:10.5194/sd-17-19-2014, 2014.

Webb, C. O., Ackerly, D. D., McPeck, M. A., and Donoghue, M. J.: Phylogenies and community ecology, *Annu. Rev. Ecol. Syst.*, 33, 475–505, doi:10.1146/annurev.ecolsys.33.010802.150448, 2002.

Wilke, T., Albrecht, C., Anistratenko, V. V., Sahin, S. K., and Yıldırım, M. Z.: Testing biogeographical hypotheses in space and time: faunal relationships of the putative ancient Lake Egirdir in Asia Minor, *J. Biogeogr.*, 34, 1807–1821, doi:10.1111/j.1365-2699.2007.01727.x, 2007.

Wysocka, A., Kostoski, G., Kilikowska, A., Wróbel, B., and Sell, J.: The *Proasellus* (Crustacea, Isopoda) species group, endemic to the Balkan Lake Ohrid: a case of ecological diversification?, *Fund. Appl. Limnol.*, 172, 301–313, doi:10.1127/1863-9135/2008/0172-0301, 2008.

Wysocka, A., Grabowski, M., Sworobowicz, L., Mamos, T., Burzyński, A., and Sell, J.: Origin of the Lake Ohrid gammarid species flock: ancient local phylogenetic lineage diversification, *J. Biogeogr.*, 41, 1758–1768, doi:10.1111/jbi.12335, 2014.

Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper

**BGD**

12, 16081–16103, 2015

## Structuring of gastropod communities

T. Hauffe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



Structuring of  
gastropod  
communities

T. Hauffe et al.



**Figure 1.** Sampling sites and eco-zonation of 264 gastropod communities of ancient Lake Ohrid and its feeder springs. Eco-zones were detected by network analysis weighting links among sampling sites by their geographical proximity and compositional similarity. **(a)** Vertical (bathymetrical) extension of the eco-zones. Due to the spatial density, plots of horizontal distribution of clusters were divided into **(b)** upper littoral, and **(c)** lower littoral and sublittoral.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



Structuring of  
gastropod  
communities

T. Hauffe et al.



**Figure 2.** Box plots showing the eco-zone dependent relative importance of (a) dispersal limitation, (b) environmental filtering, and (c) biotic interaction in structuring gastropod communities. Colors correspond to Fig. 1. These differences in process-importance due to eco-zones are statistically supported by a multivariate Bayesian generalized linear model.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

