

1 **Scaling impacts on environmental controls and spatial heterogeneity of soil organic carbon**
2 **stocks**

3

4 U. Mishra^{1*} and W.J. Riley²

5

6 ¹Environmental Science Division, Argonne National Laboratory, 9700 South Cass Avenue, 240-
7 6143, Argonne, IL 60439, USA.

8

9 ²Earth Sciences Division, Lawrence Berkeley National Laboratory, One Cyclotron Road, 50-
10 4037, Berkeley, California 94720, USA.

11

12

13 *Corresponding author, tel: 630-252-1108, email:umishra@anl.gov

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

1 **Abstract**

2 The spatial heterogeneity of land surfaces affects energy, moisture, and greenhouse gas
3 exchanges with the atmosphere. However, representing heterogeneity of terrestrial hydrological
4 and biogeochemical processes in earth system models (ESMs) remains a critical scientific
5 challenge. We report the impact of spatial scaling on environmental controls, spatial structure,
6 and statistical properties of soil organic carbon (SOC) stocks across the U.S. state of Alaska. We
7 used soil profile observations and environmental factors such as topography, climate, land cover
8 types, and surficial geology to predict the SOC stocks at a 50-m spatial scale. These spatially
9 heterogeneous estimates provide a dataset with reasonable fidelity to the observations at a
10 sufficiently high resolution to examine the environmental controls on the spatial structure of
11 SOC stocks. We upscaled both the predicted SOC stocks and environmental variables from finer
12 to coarser spatial scales ($s = 100$ m, 200 m, 500 m, 1 km, 2 km, 5 km, 10 km) and generated
13 various statistical properties of SOC stock estimates. We found different environmental factors
14 to be statistically significant predictors at different spatial scales. Only elevation, temperature,
15 potential evapotranspiration, and scrub land cover types were significant predictors at all scales.
16 The strengths of control (the median value of geographically weighted regression coefficients) of
17 these four environmental variables on SOC stocks decreased with increasing scale and were
18 accurately represented using mathematical functions ($R^2=0.83-0.97$). The spatial structure of
19 SOC stocks across Alaska changed with spatial scale. Although the variance (sill) and
20 unstructured variability (nugget) of the calculated variograms of SOC stocks decreased
21 exponentially with scale, the correlation length (range) remained relatively constant across scale.
22 The variance of predicted SOC stocks decreased with spatial scale over the range of 50 m to
23 ~500 m, and remained constant beyond this scale. The fitted exponential function accounted for

1 98% of variability in the variance of SOC stocks. We found moderately-accurate linear
2 relationships between mean and higher-order moments of predicted SOC stocks ($R^2 \sim 0.55-0.63$).
3 Current ESMs operate at coarse spatial scales (50 - 100 km), and are therefore unable to
4 represent environmental controllers and spatial heterogeneity of high-latitude SOC stocks
5 consistent with observations. We conclude that improved understanding of the scaling behavior
6 of environmental controls and statistical properties of SOC stocks could improve ESM land
7 model benchmarking and perhaps allow representation of spatial heterogeneity of
8 biogeochemistry at scales finer than those currently resolved by ESMs.

9

10 **1 Introduction**

11

12 Soil organic carbon (SOC) stocks have large spatial heterogeneity globally (Todd-Brown
13 et al., 2013) and in the northern circumpolar permafrost region (Ping, 2013; Hugelius et al.,
14 2014). Achieving an accurate representation of the spatial heterogeneity of existing SOC stocks
15 in earth system models (ESMs) is a prerequisite for predicting future carbon-climate feedbacks
16 (Todd-Brown et al., 2013), given that uncertainty resulting from the distribution of SOC stocks
17 accounts for a large proportion of the overall uncertainty (Burke et al., 2012) in predictions of
18 future greenhouse gas concentrations and associated climate changes. Reliable estimates of
19 regional SOC stocks and their spatial heterogeneity are also essential to gaining a better
20 understanding of existing environmental controls of SOC stocks and their vulnerability to
21 changing climate (Johnson et al., 2011).

22 At present, large differences exist between observation-based and ESM-based SOC stock
23 estimates at high latitudes (Mishra et al., 2013). Several factors likely account for these

1 differences (Koven et al. 2013; Todd-Brown et al. 2013); the relevant factors for the current
2 analyses are that the natural spatial variability of some environmental controls observed to affect
3 SOC stocks are not represented in the models. One potential approach to account for spatial
4 heterogeneity in ESMs is to incorporate the impacts of environmental controls on SOC stocks at
5 a spatial scale consistent with the observations.

6 | Among environmental variables, climate, topography, organisms, parent material, [time](#),
7 | and spatial position (Jenny, 1941; McBratney et al., 2003) have widely been used to predict the
8 | spatial variability of soil properties from plot to regional scales (Thompson and Kolka; 2005;
9 | Vasques et al., 2012; Minasny et al., 2013; Mishra and Riley, 2014). At high latitudes where
10 | solar radiation and evaporation are strongly related to the geometry of the land surface
11 | (McKenzie et al., 2000), topography has been reported to play an important role in determining
12 | the level of SOC at a specific location (Jenny 1980; Birkeland, 1984; Hobbie et al., 2000). Hill
13 | slope processes, such as erosion and deposition, are related to terrain attributes, such as the soil
14 | wetness index (SWI) and sediment transport index (STI). Both the SWI and STI are strong
15 | predictors of SOC stocks and have been used extensively in digital mapping of SOC stocks at
16 | various scales (e.g., Zhu et al., 2001; Thompson et al., 2006; Adhikari et al., 2013).

17 Despite several global and regional studies on SOC inventories, quantitative data on the
18 | relationship between SOC stocks and the environmental controlling factors do not exist at
19 | multiple scales (Lal, 2004; [Torn et al., 2009](#); Chaplot et al., 2010). In spatial prediction of SOC
20 | concentration and stocks, different primary and secondary topographic attributes are calculated
21 | from a digital elevation model (DEM) and used, along with other soil-forming factors such as
22 | climate, land cover type, and parent material, to predict SOC stocks across different scales
23 | (Hancock et al., 2010; Vasques et al., 2010; Kumar et al., 2012). Usually, fine-scale

1 environmental data are preferred over coarser spatial scales as they are more representative of
2 natural landscapes. However, it has also been reported that the highest scale DEMs do not
3 always produce highest accuracy in predicting soil properties and that knowledge of the
4 appropriate DEM scale for a particular landscape is important (Smith et al., 2006; Roecker and
5 Thompson, 2010). Calculated values of topographic indices of a study area differed depending
6 on the spatial scale of the DEM used (Wilson and Gallant, 2000; Thompson et al., 2001; Kienzle,
7 2004). In a recent study, Vasques et al. (2012) showed that the quality (R^2) of predicted carbon
8 stocks consistently decreased with increasing grid size of the environmental data used by the
9 models.

10 Information collected at one scale can be used to infer properties at either smaller or
11 larger scales (Beven 1995), such as when using point observations to estimate regional SOC
12 stocks, or taking areal averages of SOC stocks from ESMs and disaggregating them. The essence
13 of successful scaling is to infer the key patterns from information collected at one scale and use
14 those patterns to make inferences at another scale that maintain consistency in the desired
15 metrics across the scales (Western et al., 2002). We believe that the statistical spatial structure of
16 SOC stocks may be useful in (1) downscaling and inferring fine-scale variability from coarse-
17 scale ESM predictions; (2) model benchmarking; and (3) developing reduced order model
18 formulations relating biogeochemical processes across scales. By model benchmarking, we mean
19 using observation based scaling relationships to test the land biogeochemical representations in
20 earth system models.

21 A large literature exists on spatial scaling of soil moisture, which is a dominant control on
22 SOC dynamics and stocks. Several studies have modeled the spatial variability of soil moisture
23 patterns from relatively smaller to larger scales, and have attempted to characterize the spatial

1 structure based on system properties (Rodriguez-Iturbe et al., 1995; Peters-Lidard et al., 2001;
2 Western et al., 2002; Riley and Shen 2014; Pau et al. 2014). For some systems, soil moisture
3 spatial variance follows a power law decay as a function of spatial area (Manfreda et al., 2007),
4 although in other systems there are clear scale breaks in this relationship (e.g., Das and Mohanty,
5 2008; Joshi and Mohanty, 2010; Pau et al., 2014). Gebremichael et al. (2009) reported that, in a
6 watershed located in the Southern Great Plains of the USA, soil moisture showed scale-
7 invariance, and if the scaling parameters could be estimated from large-scale soil moisture fields,
8 it might be feasible to transform spatial soil moisture fields between scales. Despite the progress
9 made in modeling the scaling behavior of soil moisture, to our knowledge no study exists that
10 has examined the statistical structure of the scaling behavior of SOC stocks.

11 Scaling approaches might also improve prediction of the spatial structure of SOC stocks.
12 The variogram, which relates the variance of SOC stock differences between two points to the
13 distance between the two points (Webster and Oliver, 2007), is one approach to representing the
14 spatial structure of SOC stocks (Western et al., 2002). The main structural parameters of a
15 variogram are the nugget, sill, and range (Western and Blöschl, 1999; Webster and Oliver,
16 2007). A nugget shows the unexplained portion of the variance and the sill is the level at which a
17 variogram flattens and therefore represents the total variance of the variable. The nugget-to-sill
18 ratio can be used to quantify the percentage of overall variance found at a distance smaller than
19 the smallest lag interval, and to classify the spatial dependence of soil properties (Cambardella et
20 al., 1994; Mora-Vallejo et al., 2008).

21 Current ESMs operate at spatial scales larger than 50 km and use a nested subgrid
22 hierarchy approach to represent the land surface heterogeneity (Lawrence et al., 2012). ESM
23 spatial scales are becoming increasingly finer to more accurately represent the localized

1 features of earth systems that affect energy, water, and greenhouse gas fluxes. For example,
2 next generation ESMs such as those being developed by the U.S. Department of Energy
3 (Accelerated Climate Modeling for Energy; ACME) will have a spatial scale for the land
4 model of ~10 km (Bader et al., 2014). In this study, we successively increased the spatial
5 scale (from $s = 50$ m to 10 km) of environmental variables and used observations and
6 geospatial approaches to predict SOC stocks. Throughout this paper, we refer to the ‘scale’
7 (s) as either the distance across which properties are assumed to be homogeneous or the
8 square root of the pixel area satisfying that criteria, and note that the terms ‘scale’ and
9 ‘resolution’ are often interchangeable in this context. The idea was to model the change in
10 predicted heterogeneity of SOC stocks resulting from changing scale of the environmental
11 data. In this study, by the term “scaling” we mean the transfer of information about
12 environmental controls and statistical properties of SOC stocks from one scale to another.
13 We also investigated how environmental controls on SOC stocks and the spatial structure of
14 SOC stocks (the correlation length (range), total variance (sill), and unstructured variability
15 (nugget)) changed as a result of change in spatial scale of environmental variables.

16 **2 Study approach**

17 **2.1 Soil organic carbon observations and environmental variables**

18 We used 556 georeferenced soil profile observations by combining data from two
19 recently published studies of soil organic carbon stocks across Alaska (472 soil profiles from
20 Mishra and Riley, 2012 and 84 additional samples from a recently published database of
21 Michaelson et al., 2013). This database combined the U.S. Department of Agriculture’s National
22 Resource Conservation Service’s pedons with the soil pedon observations collected from the
23 University of Alaska Fairbanks under its northern [latitudes](#) soils ~~research~~ program ([Michaelson](#)

1 | [et al., 2013](#)). This database included measured representative soil profiles from Alaska and
2 | covered all soil types at the soil suborder level (18 suborders) and all 27 major land resource
3 | areas of Alaska. The observed SOC stocks showed a wide range (0.35–296 kg m⁻²) due to
4 | varying sampling depths (0.05–4.5 m depth), various soil types with differing carbon content
5 | levels (Inceptisols to Gelisols), and the large environmental heterogeneity present in the study
6 | area. [About 50% of the observed samples were from the 0.05 to 1 m depth interval, 41% of the](#)
7 | [samples were from the 1- 2 m depth interval, and the remaining samples \(9%\) were from the 2-](#)
8 | [4.5 m depth interval.](#) A large number of observations had SOC stocks ≤ 50 kg m⁻², and about
9 | 49 samples had SOC stocks greater than 100 kg m⁻². A majority of pedons with high SOC stocks
10 | were located in the Arctic region of Alaska, which are areas with soils impacted by
11 | cryoturbation, and the majority of pedons with lower SOC stocks were located in interboreal and
12 | coastal rainforests. The SOC stock data has a unimodal (kurtosis, $K = 9.4$) positively skewed
13 | distribution (coefficient of skewness, $S = 2$), and is unevenly distributed throughout Alaska (Fig.
14 | 1). The fine-scale (spatial scale $s = 50$ m) spatially heterogeneous SOC stock estimates have
15 | uncertainty, as described by the prediction errors in Mishra and Riley (2012). However, that
16 | study indicated that the SOC stock estimates have reasonable fidelity with the observations
17 | withheld from the estimation procedure, and therefore we believe are sufficiently accurate to be
18 | used in the scaling analyses described here. As more data become available, the approaches
19 | described here to evaluate spatial scaling controls can be [readily](#) re-applied.

20 | We collected environmental variables of the study area from various sources. The
21 | topographic attributes were derived from a digital elevation model (DEM) with a 50-m spatial
22 | scale ($s = 50$ m) obtained from the U.S. Geological Survey (USGS) database (Gesch et al.,
23 | 2009). From the DEM, 13 topographic attributes were calculated that have been reported to be

1 useful predictors of the SOC stocks across different environmental conditions using the Spatial
2 Analyst function of ArcGIS (ArcGIS version 10.2, Environmental Systems Research Institute,
3 Inc., Redlands, CA, USA).

4 Climate data for long-term (1961–1990) average annual temperature, precipitation,
5 potential evapotranspiration (PET), and summer shortwave radiation were obtained from the
6 Scenario Network for Alaska and Arctic Planning (SNAP, 2014) database, with a spatial scale of
7 $s = 2$ km.

8 Land cover data of 60 m spatial scale ($s = 60$ m) was extracted for Alaska from the
9 National Land Cover Database (NLCD) database (Homer et al., 2007). In this study, to reduce
10 the number of categorical variables with potentially redundant information, we reclassified the
11 19 NLCD land cover types found in Alaska into 9 major categories. Thus, developed open space,
12 low intensity, medium intensity, and high intensity were classified as a developed category;
13 deciduous, evergreen, and mixed forests were classified as a forest category; dwarf scrub and
14 shrub scrub were classified as a scrub category; shrub, sedge, and moss were classified as a
15 herbaceous category; pasture and cultivated lands were classified as a cultivated category; and
16 woody and herbaceous wetlands were classified as a wetland category. In the reclassified map,
17 the category with the largest land area was the scrub category (43%), followed by forest (25%),
18 barren (8.5%), herbaceous (7%), and wetlands (7%). The remaining surface area (9.5%) was
19 under open water, perennial ice, barren land, and moss vegetation. Indicator variables for the
20 presence or absence of 7 land cover types (except open water and perennial ice) were created and
21 used in the model selection process.

22 The surficial geology data of Alaska was obtained from a USGS database (Karlstrom,
23 1964) in which the entire state of Alaska was classified into 27 different types of surficial

1 geology. The category with the largest land area was mountain alluvium and colluvium (22.5%),
2 followed by coarse rubble (19%), coastal alluvium (8.5%), glacial moraine (7%), and
3 undifferentiated mosaics (6%). The remaining land area (37%) was placed in the remaining
4 22 surficial geology types.

5

6 **2.2 Spatial scaling and environmental controls**

7 We investigated changes in environmental controls on SOC stocks at spatial scales of
8 100 m, 200 m, 500 m, 1 km, 2 km, 5 km, and 10 km. For this purpose, the DEM of a 50-m
9 spatial scale was successively aggregated to 100 m, 200 m, 500 m, 1 km, 2 km, 5 km, and 10 km
10 using the bilinear convolution algorithm of ArcGIS 10.2 (Fig. 2). Topographic attributes were
11 calculated at each spatial scale and included in predicting SOC stocks at specific scales. The land
12 cover types and surficial geology data were aggregated at similar scales using the nearest-
13 neighbors algorithm of ArcGIS 10.2 and were included in predicting SOC stocks at various
14 spatial scales.

15 Median values of the geographically weighted regression coefficients (β) of log-
16 transformed (natural log) SOC stock and environmental variables (that were significant at all
17 spatial scales) were calculated for each spatial scale. Log transformations of SOC stocks were
18 applied to meet the linear regression assumptions of normality and constant variance of errors.
19 The change in the strength of environmental controls (β 's) on SOC stocks resulting from a
20 change in spatial scale was investigated by plotting the median values against the spatial scale at
21 which they were calculated. Several mathematical functions were used to represent the statistical
22 nature of scaling of the environmental controls on SOC stocks.

23

1 **2.3 Selection of environmental predictors, prediction of SOC stocks, and variogram**
2 **modeling**

3 We used best subset regression in SAS 9.3 (SAS Institute, 2011) to generate all possible
4 combinations of environmental variables to predict the SOC stocks across Alaska. We used
5 Mallows' C_p criterion to select three candidate linear models for each spatial scale for which C_p
6 values were close to the number of predictors (Kutner et al., 2004; p.358), from which we
7 subsequently selected one linear model for each spatial scale with uncorrelated and statistically
8 significant environmental predictors. Presence of multicollinearity among selected
9 environmental predictors was tested by calculating the variance influence factors (VIFs) for each
10 of the selected variables. The VIFs for all the variables included in models selected at different
11 spatial resolutions were < 5. High levels of multicollinearity (VIF > 10) can falsely inflate the
12 least squares estimates (Kutner et al., 2004; O'Brien R., 2007; Gomez et al., 2013). The selected
13 environmental predictors were then used in a geographically weighted regression approach
14 (Mishra et al., 2012; 2014) to predict the whole profile SOC stocks across Alaska at different
15 spatial scales using:

$$C(u_i, v_i) = \beta_o(u_i, v_i) + \sum_{k=1}^p \beta_k(u_i, v_i)X_k + \varepsilon(u_i, v_i)$$

16 where C is the SOC stock at a certain location, u_i, v_i are the geographical coordinates, X_k are the
17 environmental predictors, p is the number of independent variables, β_o and β_k are the
18 geographically weighted regression coefficients, and ε is the error term. We used adaptive kernel
19 bandwidth in this study, given that the sample density varied over the study area. The optimal
20 bandwidth was determined by minimizing the corrected Akaike Information Criterion (AICc) as
21 described in Fortheringham et al. (2002).

1 We calculated several statistical parameters, including mean (μ), variance (σ^2), skewness
2 | (S), and kurtosis (K) from the predicted SOC stocks at each specific spatial scale. The predicted
3 | variances were plotted against spatial scale and a best-fit mathematical function was determined.
4 Relationships between the μ and other statistical parameters were investigated by plotting the μ
5 SOC stocks calculated at each spatial scale across Alaska versus its other statistical parameters
6 calculated at the same scale.

7 To study the change in spatial structure of SOC stocks due to spatial scaling, we
8 | calculated the SOC stock variograms (Webster and Oliver, 2007) at different spatial scales:

$$\gamma(h) = \frac{1}{2n} \sum_{i=1}^n \{Z(X_i) - Z(X_i + h)\}^2$$

9 where $Z(X_i)$ and $Z(X_i + h)$ are the measured SOC stocks at X_i and $X_i + h$, respectively; h is the lag;
10 n is the number of paired comparisons at that lag; and $\gamma(h)$ is the semivariance. By varying h in
11 discrete steps, we obtain an ordered set of semivariances (Webster and Oliver, 1992). Suitable
12 mathematical functions were fitted to the calculated semivariance values using weighted least
13 squares (Webster and Oliver, 2007). The best-fit model parameters for each spatial scale are
14 provided in Table 2. We examined the change in correlation length (range), total variance of
15 SOC stocks (sill), and the unstructured variability (nugget) resulting from the change in the
16 spatial scales of environmental variables. A variable is considered to have a strong spatial
17 dependence if the nugget-to-sill ratio is less than 0.25; a moderate spatial dependence if the ratio
18 is between 0.25 and 0.75; and otherwise a weak spatial dependence (Cambardella et al., 1994;
19 Mora-Vallejo et al., 2008). Range, which is also referred as correlation length, is the distance at
20 which the sill is reached and represents the maximum extent of the spatial correlation between

1 predicted SOC stocks. The change in the spatial structure of variability of SOC stocks was
2 evaluated using the nugget-to-sill ratio and the correlation length.

3

4 **3 Results**

5 **3.1 Inferred environmental controls on SOC stocks change with scale**

6 The set of significant environmental predictors and their individual strengths varied with
7 spatial scale. Statistically significant environmental predictors at different spatial scales and the
8 predicted variance of SOC stocks are summarized in Table 1. From the 13 topographic attributes,
9 5 attributes—elevation (meters), slope (degree), aspect (degree from north), soil wetness index
10 (SWI), and sediment transport index (STI)—were significant predictors of SOC stocks at various
11 spatial scales. Among land cover types, barren, scrub, herbaceous, and cultivated land covers
12 were significant predictors. Among different climatic factors, surface air temperature and
13 potential evapotranspiration were significant predictors. Among surficial geology types, the
14 glacial moraine, fluvial, undifferentiated mosaic, coastal delta, and volcanic mountain types were
15 significant predictors at various spatial scales. Among all of the environmental variables
16 investigated in this study, only elevation, temperature, potential evapotranspiration, and scrub
17 land cover types were significant predictors of SOC stocks at all spatial scales. In general, at
18 finer spatial scales, controls of topographic attributes were more prominent, whereas land cover
19 and surficial geology types were more significant predictors at coarser scales (≥ 500 -m spatial
20 scale). Fewer environmental variables controlled the variability of SOC stocks at finer spatial
21 scales, whereas the number of predictors increased at larger spatial scales.

22 Our results showed that the strength of the control (median geographically weighted
23 regression coefficient across Alaska) of elevation on SOC stocks decreased by 31% as the spatial

1 scale increased from 50 m to 1 km. Beyond this scale, we found no change in the control of
2 elevation on predicted SOC stocks (Fig. 3a). The control of temperature on SOC stocks
3 decreased with spatial scale between 50 m and 500 m and became constant at larger scales (Fig.
4 3b). The controls of elevation and temperature on SOC stocks across spatial scales can be
5 accurately modeled by using exponential functions with $R^2=0.83$ and 0.94 , respectively. The
6 control of potential evapotranspiration decreased by 36% as the spatial scale increased from
7 50 m to 500 m, and became constant beyond 500 m (Fig. 3c). The control of potential
8 evapotranspiration on SOC stocks across spatial scales can be modeled by using an exponential
9 decay function ($R^2=0.97$). The control of scrub vegetation (shrubs less than 5 meters tall) on
10 SOC stocks decreased linearly (from 0.3-0.13) with scale over the range 50 m to 10 km (Fig. 3d).
11 At finer scales, scrub vegetation showed higher control on SOC stocks, which decreased by
12 about 57% as the scale increased. Despite the observed decrease in control of scrub vegetation on
13 SOC stocks due to scaling, the relationship between scrub vegetation and SOC stocks remained
14 positive. The control of scrub vegetation on SOC stocks can be modeled using a linear function
15 ($R^2=0.85$).

16

17 **3.2 Scaling impacts on the spatial structure of SOC stocks**

18 The variogram summarizes the statistical structure of spatial variability and is known as a
19 summarizing function of the investigated property. For the predicted SOC stocks, the variograms
20 at finer-scales showed significant amount of unexplained variance (nugget) (Table 2) due to
21 measurement errors and/or microscale spatial heterogeneity not captured across the study area.
22 Both the nugget and sill decreased as the scale increased from finer to coarser spatial scales. The
23 nugget and sill decreased by 75% and 64% respectively as s increased from 50 m to 500 m and

1 remained constant beyond 500 m scale (Fig. 4a). The relationship between these variogram
2 parameters and scale can be accurately described using exponential functions with R^2 of 0.98.
3 The correlation length of SOC stocks remained relatively constant up to about 1100 -1400 km
4 across spatial scales (Fig. 4b). The nugget-to-sill ratio of predicted SOC stocks at 50 m and 100
5 m spatial scales showed moderate spatial dependency (i.e. a nugget-to-sill ratio $>25\%$;
6 Cambardella et al., 1994). However, strong, and relatively similar, spatial dependency (a nugget-
7 to-sill ratio $\leq 25\%$) was predicted for spatial scales between $s = 200$ m and $s = 10$ km (18–21%;
8 Table 2). These results suggest that the predicted SOC stocks show different spatial structure
9 below $s = 100$ m as compared to $s \geq 200$ -m.

10

11 **3.3 Statistical behavior of spatial scaling of SOC stocks**

12 We found different statistical properties in predicted SOC stocks as the spatial scale of
13 environmental variables increased. The variance of predicted SOC stocks decreased by 45% as
14 the scale increased from 50 m to 500 m and remained constant beyond 500 m (Fig. 5). An
15 exponential function based on spatial scale accounted for 98% of variability in the variance of
16 predicted SOC stocks across Alaska. These results suggest that the spatial heterogeneity of SOC
17 stocks decreases exponentially with scale until about 500 m. Beyond this scale, no change in the
18 spatial heterogeneity of SOC stocks was predicted. The higher-order moments (skewness S and
19 kurtosis K) of the predicted SOC stocks decreased as the spatial scale increased (Fig. 6a). S
20 decreased from 1.4 to 0.8 as the scale increased from 50 m to 200 m; and fluctuated between
21 1.25 and 0.74 as the scale increased further. K decreased from 5.7 to 3.6 as the scale increased
22 from 50 m to 200 m and fluctuated between 5.1 and 3.68 beyond a scale of 200 m.

1 | We evaluated the relationships of mean SOC stocks (μ) generated at each spatial scale
2 | with σ^2 , S , and K . Within the investigated range of μ SOC stocks (45.7 to 50.4 kg m⁻²), we found
3 | moderately-accurate linear relationships between μ and σ^2 , S , and K of predicted SOC stocks,
4 | with R² values of 0.59, 0.63, and 0.55, respectively (Figs. 6 b-d). Significant negative slopes of
5 | these relationships indicate that as μ increases the higher-order moments of SOC stock decrease,
6 | and therefore the μ of SOC stocks can be used to predict SOC spatial heterogeneity.
7 |

8 | **4 Discussions**

9 | Soil organic carbon lies at the interface between the atmosphere and pedosphere and can
10 | be altered at time scales relevant to climate change by anthropogenic and climatic factors. As a
11 | result, the spatial heterogeneity of SOC stocks may affect the magnitude of greenhouse gas
12 | fluxes from the land surface and the terrestrial carbon cycle.

13 | Several studies have demonstrated the impact of DEM scale on calculated topographic
14 | attributes (Thompson et al., 2001; McMaster, 2002; Smith et al., 2006; Roecker and Thompson,
15 | 2010). However, very little attention has been paid to the impact of scaling on environmental
16 | controls and the predicted spatial heterogeneity of soil properties. We found that apparent
17 | significant environmental controllers of high-latitude SOC stocks changed with scale. Our
18 | findings are consistent with the results of Vasques et al. (2012), who showed inconsistent
19 | controls of various tropical-system environmental factors in predicting SOC stocks as the scale
20 | of environmental variables increased from 30 m to 1,920 m. Due to a lack of other scaling
21 | studies examining the change in environmental controllers of SOC stocks, we compared our
22 | results with findings from two studies that reported environmental controllers of SOC stocks at
23 | two different scales. Johnson et al. (2011) reported that at the plot scale (~ 1 m²), surface air

1 temperature, topographic attributes, and soil texture were primary controllers of SOC stocks
2 across Alaska. Similarly, Martin et al. (2011) predicted SOC stocks at 12 km scale over France,
3 and reported land use and clay content to be important drivers of SOC stock spatial variability.
4 Our results at 50 m and 10 km scales are partially consistent with these findings, showing
5 significant controls of temperature, land cover types, and topographic attributes on SOC stocks.
6 Our results showed that the strength of controls of environmental factors that were significant
7 decreased as spatial scale increased. The largest decrease was found in the control of scrub land
8 cover type, and the smallest decrease was found in the control of temperature. The rates of these
9 changes can be modeled using simple mathematical functions.

10 The scaling properties of soil moisture have been widely studied (Western and Boschl,
11 1999; Isham et al., 2005; Famiglietti et al., 2008; Li and Rodell, 2013). Some studies reported
12 that the variance of soil moisture follows a power law relationship with surface area (Rodriguez-
13 Iturbe et al. 1995; Manfreda et al., 2007), while others have shown non-simple scaling (Joshi and
14 Mohanty, 2010; Riley and Shen 2014; Pau et al. 2014). Beven (1995) pointed out that it is
15 unlikely that any general scaling theory can be developed because of the dependence of
16 hydrological systems on geological perturbations, and advocated for a disaggregation approach
17 to develop scale-dependent models to represent hydrological heterogeneity. These studies
18 suggested that further work focusing on the factors influencing soil moisture (topography,
19 | vegetation, soil properties, and rainfall) is expected to increase understanding of the mechanisms
20 | affecting the scaling properties of soil moisture.

21 Keeping these inferences in mind and that soil moisture is an important control on SOC
22 dynamics, in this study we used the soil-forming factors that have been documented to affect
23 SOC stocks to study the spatial scaling behavior of SOC stocks. However, in contrast to soil

1 moisture, our results show that the variance of SOC stocks follows an exponential decay with
2 spatial scale up to about 500 m, and then remains constant thereafter. We interpreted
3 environmental controllers on SOC stocks in an equilibrium sense; i.e., over hundreds to
4 thousands of years these controllers are expected to be related to observed SOC stocks. However,
5 transient responses are not expected to follow the same scaling properties, given likely changes
6 in thermokarst formation and other hydrological and geomorphological changes at high latitudes.

7 The spatial structure of SOC stocks results from the spatial structure of soil-forming
8 factors (Jenny, 1941; McBratney et al., 2003). Besides the state factors described in the
9 Introduction, SOC stocks of Alaska and other arctic regions are also impacted by cryopedogenic
10 features (pingos, frost boils, hummocks) and thermal surface erosion resulting from the freezing
11 and thawing of the polygonal land surface (Tarnocai and Bockheim, 2011; Ping et al., 2008; Ping
12 2013). In this study, the spatial organization of predicted SOC stocks increased with spatial scale
13 as indicated by moderately lower nugget-to-sill ratios for spatial scales >200 m. However, the
14 correlation length remained relatively constant in the upscaled SOC stocks across Alaska.

15 One potential approach to represent the spatial heterogeneity of soil properties in ESMs is
16 to relate their statistical properties to the mean state. As an analogy, several soil moisture studies
17 investigated the relationships between soil moisture mean and higher-order statistics, such as σ^2 ,
18 S , and K (Famiglietti et al., 1999; Ryu and Flamiglietti, 2005; Famiglietti et al., 2008; Li and
19 Rodell, 2013). The results from these studies suggest that the mean of soil moisture is often
20 related to its σ^2 , S , and K , although with different functional forms depending on various system
21 properties. Our results indicate a moderate but statistically significant linear relationship between
22 the mean and higher-order moments of SOC stocks (i.e., σ^2 , S , and K decrease linearly with the
23 mean SOC stock within the range of mean SOC stocks investigated in this study (45.5 – 50.4 kg

1 m⁻²). The strength of the linear relationship was highest between μ and S ($R^2=0.63$, $P<0.01$), and
2 lowest between μ and K ($R^2=0.55$, $P<0.05$). However, available observations are insufficient to
3 determine whether the same relationship holds with dynamically changing mean SOC stocks,
4 highlighting an area of further investigation.

5 Current land models integrated in ESMs, such as CLM4.5 (Lawrence et al., 2012; Koven
6 et al., 2013; Tang et al., 2013), use a nested subgrid hierarchy approach to represent the land
7 surface heterogeneity. In this approach the grid cells ($\sim 1^\circ \times 1^\circ$) are divided into nonspatially
8 explicit land units (such as vegetated, lakes, urban, glacier, and crops), columns (with variability
9 in hydrological, snow, and crop management), and plant functional types (accounting for
10 variations in broad categories of plants and bare ground). We have shown here that this type of
11 representation does not characterize the environmental controls and scaling properties of SOC
12 stocks; rectifying this problem would require substantial restructuring of the model's subgrid
13 hierarchy. One potential application of the relationships we developed in this study could be to
14 apply them with coarse-resolution ESM results in order to generate fine-scale spatial
15 heterogeneity parameters of SOC stocks that are more representative of the natural landscape.
16 Currently the land models of most ESMs have a spatial scale of ≥ 50 km. Therefore our results
17 will be directly relevant to next generation ESMs which will operate at high resolution (~ 10 km)
18 (Bader et al., 2014).

19 Median values of regression coefficients of different environmental controls might
20 change if a global or continental scale study would have been conducted. Other environmental
21 factors which are not significant predictors of Alaskan SOC stocks might become significant in
22 different study domains. Further work is required to investigate these factors in other systems
23 and at larger spatial extents.

1 Soil texture has also been reported to be related to SOC stocks. To investigate the use of
2 soil texture in our scaling study, we collected the soil texture data currently used in CLM 4.5
3 (Bonan et al., 2002) that was available at ~8 km spatial resolution across Alaska. However, this
4 data has approximately one soil texture value in each Alaska ecoregion implying that the data
5 must have been derived from a source with a much coarser spatial resolution (International
6 Geosphere-Biosphere Programme soil dataset that had 4931 soil mapping units globally; CLM
7 4.5 Technical notes). Because of this limitation, we were unable to include soil texture in the
8 current scaling study. However, if soil texture information becomes available at finer spatial
9 scales, it could be readily incorporated in future studies using the methods described here.
10

11 **5 Summary and conclusions**

12 Understanding the spatial scaling structure of SOC stocks and their relationships with
13 environmental factors is important for prediction of carbon-climate feedbacks under a changing
14 climate. Here, we estimated the spatial scaling properties of environmental factors and statistical
15 properties of SOC stocks. We conclude that environmental controllers of SOC stocks change
16 with scale, and the strength of environmental controls can be accurately modeled using simple
17 mathematical functions. The variance of predicted SOC stocks decreases exponentially with
18 scale up to about 500 m, and then remains constant thereafter. Our results showed that the mean
19 predicted SOC content is linearly related with its variance, skewness, and kurtosis. If the
20 objective of a study is to represent the spatial heterogeneity of SOC stocks resulting from
21 environmental factors, the prediction spatial scale should be finer than ~500 m. Similarly, the
22 choice of environmental predictors should be based on the intended final spatial scale of
23 predicted SOC stocks. Therefore, the current subgrid structure of many ESM-scale land models

1 does not allow for characterization of the environmental controls and scaling properties affecting
2 SOC stocks. The findings of this study may help to develop scale-appropriate techniques for
3 ESM-scale land models and ultimately reduce the existing uncertainties in carbon-climate
4 feedback predictions.

5
6 *Acknowledgments*

7
8 This research was supported by the Director, Office of Science, Office of Biological and
9 Environmental Research of the US Department of Energy under Argonne National Laboratory
10 contract No. DE-AC02-06CH11357, and the Regional and Global Climate Modeling (RGCM)
11 Program. Contributions of W.J. Riley were supported under Lawrence Berkeley National
12 Laboratory Contract No. DE-AC02-05CH11231 as part of the RGCM program and the Next-
13 Generation Ecosystem Experiment Arctic project. Thanks to G. Michaelson and C.L. Ping for
14 providing access to some of the SOC profile data.

15
16
17
18
19
20
21
22
23

24 **References**

25 Adhikari, K., Bou Kheir, R., Greve, M.B., Bøcher, P.K., Malone, B.P., Minasny, B., McBratney,
26 A.B., and Greve, M.H.: High-Resolution 3-D Mapping of Soil Texture in Denmark, *Soil Sci.*
27 *Soc. Am. J.*, 77, 860-876, 2013.

28
29 [Bonan, G.B., Levis, S., Kergoat, L., and Oleson, K.W.: Landscapes as patches of plant functional](#)
30 [types: An integrating concept for climate and ecosystem models, *Global Biogeochemical*](#)
31 [*Cycles*, 16, 5.1-5.23, 2002.](#)
32

1 Bader, D., Collins, W., Jacob, R., Jones, P., Rasch, P., Taylor, M., Thornton, P., and Williams,
2 D.: Accelerated Climate Modeling for Energy (ACME) Project Strategy and Initial
3 Implementation Plan; 2014. (Available online at <http://climatemodeling.science.energy.gov>).
4

5 Beven, K.: Linking parameters across scales: Subgrid parameterizations and scale-dependent
6 hydrological models, *Hydrol. Process.*, 9, 507–25, 1995.
7

8 Birkeland, P.W.: Soils and geomorphology, Oxford University Press, New York, 1984.
9

10 Burke, E.J., Hartley, I.P., and Jones, C.D.: Uncertainties in the global temperature change caused
11 by carbon release from permafrost thawing, *The Cryosphere*, 6, 1063–1076, 2012.
12

13 Cambardella, C.A., Moorman, T.B., Novak, J.M., Parkin, T.B., Karlen, D.L., Turco, R.F., and
14 Konopka, A.E.: Field-scale variability of soil properties in Central Iowa soils, *Soil Sci. Soc.
15 Am. J.*, 58, 1501–1511, 1994.
16

17 Chaplot, V., Bounthong, B., and Valentin, C.: Soil organic carbon stocks in Laos: Spatial
18 variations and controlling factors, *Glob. Change Biol.*, 16, 1380–1393, 2010.
19

20 Das, N.N. and Mohanty, B. P.: Temporal dynamics of PSR-based soil moisture across spatial
21 scales in an agricultural landscape during SMEX02: A wavelet approach, *Remote Sens.
22 Environ.*, 112, 522–534, 2008.
23

24 Famiglietti, J.S., Devereaux, J.A., Laymon, C.A., Tsegaye, T., Houser, P.R., Jackson, T.J.,
25 Graham, S.T., Rodell, M., and van Oevelen, P.J.: Ground-based investigation of soil moisture
26 variability within remote sensing footprints during the Southern Great Plains 1997 (SGP97)
27 Hydrology Experiment, *Water Resour. Res.*, 35, 1839–1851, 1999.
28

29 Famiglietti, J.S., Ryu, D.R., Berg, A.A., Rodell, M., and Jackson, T.J.: Field observations of soil
30 moisture variability across scales, *Water Resour. Res.*, 44, W01423,
31 doi:10.1029/2006wr005804, 2008.
32

33 Fotheringham, A.S., Brunson, C., and Charlton, M.E.: Geographically Weighted Regression:
34 The Analysis of Spatially Varying Relationships, John Wiley & Sons, UK, 2002.
35

36 Gebremichael, M., Rigon, R., Bertoldi, G., and Over, T.M.: 2009. On the scaling characteristics
37 of observed and simulated spatial soil moisture fields, *Nonlinear Proc. Geoph.*, 16, 141–150,
38 2009.
39

40 Gesch, D., Evans, G., Mauck, J. Hutchinson, J. and Carswell Jr., W.J.: The National Map—
41 Elevation: U.S. Geological Survey Fact Sheet 2009-3053, 4, 2009.
42

43 [Gomez, C., Le Bissonnais, Y., Annabi, M., Bahri, H., and Raclot, D.: Laboratory Vis-NIR](#)
44 [spectroscopy as an alternative method for estimating the soil aggregate stability indexes of](#)
45 [Mediterranean soils, *Geoderma*, 209–210, 86–97, 2013.](#)
46

- 1 Hancock, G.R., Murphy, D., and Evans, K.G.: Hillslope and catchment scale soil organic carbon
2 concentration: An assessment of the role of geomorphology and soil erosion in an
3 undisturbed environment, *Geoderma*, 155, 36–45, 2010.
4
- 5 Hobbie, S.E., Schimel, J.P. Trumbore, S.E. and Randerson, J.R.: Controls over carbon storage
6 and turnover in high-latitude soils, *Glob. Change Biol.*, 6, 196–210, 2000.
7
- 8 Homer, C., Dewitz, J., Fry, J., Coan, M., Hossain, N., Larson, C., Herold, N., McKerrow, A.,
9 VanDriel, J.N., and Wickham, J.: Completion of the 2001 National Land Cover Database for
10 the Conterminous United States, *Photogramm. Eng. Remote Sens.*, 73, 337-341, 2007.
11
- 12 Hugelius G.H., Strauss, J., Zubrzycki, S., Kuhry, P., Harden, J.W., Schuur, E.A.G., Ping, C-L.,
13 Schirrmeister, L., Michaelson, G., Koven, C.D., O'Donnell, J., Elberling, B., Mishra, U.,
14 Camill, P., Yu, Z., and Palmtag, J.: Revised estimate confirms large circumpolar stocks of
15 permafrost carbon but reveals wide uncertainty ranges and remaining data gaps,
16 *Biogeosciences Discussions*, 11, 4771-4822, 2014.
17
- 18 Isham, V., Cox, D.R., Rodriguez-Iturbe, I., Porporato, A., and Manfreda. S.: Representation of
19 space-time variability of soil moisture, *P. Roy. Soc. Lond. A Mat.*, 461, 4035–4055, 2005.
20
- 21 Jenny, H.: *Factors of soil formation: A system of quantitative pedology*, McGraw Hill, New
22 York, 1941.
23
- 24 Jenny, H.: *The soil resource: Origin and behavior*, Springer, New York, 1980.
25
- 26 Johnson, K.D., Harden, J.W., McGuire, A.D., Bliss, N.B., Bockheim, J.G., Clark, M., Nettleton-
27 Hollingsworth, T., Jorgenson, M.T., Kane, E.S., Mack, M., O'Donnell, J., Ping, C.L., Schuur,
28 E.A.G., Turetsky, M.R., and Valentine, D.W.: Soil carbon distribution in Alaska in relation
29 to soil-forming factors, *Geoderma*, 167–168, 71–84, 2011.
- 30 Joshi, C. and Mohanty, B. P.: Physical controls of near-surface soil moisture across varying
31 spatial scales in an agricultural landscape during SMEX02, *Water Resour. Res.*, 46, W12503,
32 doi:10.1029/2010WR009152, 2010.
- 33 Karlstrom, T.N.V.: *Surficial geology of Alaska*. U.S. Geological Survey, Washington, D.C.
34 USGS Misc. Geol. Invest. Map I-357, 2 sheets, scale 1:1,584,000, 1964.
35
- 36 Kienzle, S.: The effect of DEM raster resolution on first order, second order, and compound
37 terrain derivatives, *Transactions in GIS*, 8, 83–111, 2004.
38
- 39 Koven, C.D., Riley, W.J., Subin, Z.M., Tang, J.Y., Torn, M.S., Collins, W.D., Bonan, G.B.,
40 Lawrence, D.M., and Swenson, S.C.: The effect of vertically resolved soil biogeochemistry
41 and alternate soil C and N models on C dynamics of CLM4, *Biogeosciences*, 10, 7109–7131,
42 2013.
43

- 1 Kutner, M.H., Nachtsheim, C.J., and Neter, J.: Applied linear regression models, McGraw-Hill,
2 New York, 2004.
3
- 4 Kumar, S., Lal, R., and Liu, D.: A geographically weighted regression kriging approach for
5 mapping soil organic carbon stock, *Geoderma*, 189-90, 627-634, 2012.
6
- 7 Lal, R.: Soil carbon sequestration to mitigate climate change, *Geoderma*, 123, 1–22, 2004.
8
- 9 Lawrence, P.J., Feddema, J.J., Bonan, G.B., Meehl, G.A., O’Neill, B.C., Oleson, K.W., Levis, S.,
10 Lawrence, D.M., Kluzek, E., Lindsay, K., and Thornton, P.E.: Simulating the
11 biogeochemical and biogeophysical impacts of transient land cover change and wood harvest
12 in the Community Climate System Model (CCSM4) from 1850 to 2100, *J. Climate*, 25,
13 3071–3095, 2012.
14
- 15 Li, B., and Rodell, M.: Spatial variability and its scale dependency of observed and modeled soil
16 moisture over different climate regions, *Hydrol. Earth Syst. Sci.*, 17, 1177–1188, 2013.
17
- 18 Manfreda, S., McCabe, M.F., Fiorentino, M., Rodriguez-Iturbe, I., and Wood, E.F.: Scaling
19 characteristics of spatial patterns of soil moisture from distributed modeling, *Adv. Water*
20 *Resour.*, 30, 2145–2150, 2007.
21
- 22 Martin, M.P., Wattenbach, M., Smith, P., Meersmans, J., Jolivet, C., Boulonne, L., and
23 Arrouays, D.: Spatial distribution of soil organic carbon stocks in France, *Biogeosciences*, 8,
24 1053–1065, 2011.
- 25 McBratney, A.B., Mendonça-Santos, M.L., and Minasny, B.: On digital soil mapping,
26 *Geoderma*, 117, 3–52, 2003.
27
- 28 McKenzie, N.J., Gessler, P.E., Ryan, P.J., and O’Connell, D.A.: The role of terrain analysis in
29 soil mapping, in: *Terrain analysis: Principles and applications*, edited by: Wilson, J.P., and
30 Gallant, J.C., John Wiley & Sons, New York, 245–265, 2000.
31
- 32 McMaster, K.J.: Effects of digital elevation model resolution on derived stream network
33 positions, *Water Resour. Res.*, 38, 1–9, 2002.
34
- 35 Michaelson, G.J., Ping, C.L., and Clark, M.: Soil pedon carbon and nitrogen data for Alaska: An
36 analysis and update, *Open J. Soil Sci.*, doi:10.4236/ojss.2013, 2013.
37
- 38 Minasny, B., McBratney, A.B., Malone, B.P., and Wheeler, I.: Digital mapping of soil carbon,
39 *Adv. Agron.*, 118, 1–47, 2013.
- 40 Mishra, U., and Riley, W.J.: Alaskan soil carbon stocks: Spatial variability and dependence on
41 environmental factors, *Biogeosciences*, 9, 3637–3645, 2012.
42
- 43 Mishra, U., Jastrow, J.D., Matamala, R., Hugelius, G., Koven, C.D., Harden, J.W., Ping, C.L.,
44 Michaelson, G.J., Fan, Z., Miller, R.M., McGuire, A.D., Tarnocai, C., Kuhry, P., Riley,
45 W.J., Schaefer, K., Schuur, E.A.G., Jorgenson, M.T., and Hinzman, L.D.: Empirical

1 estimates to reduce modeling uncertainties of soil organic carbon in permafrost regions: A
2 review of recent progress and remaining challenges, *Environ. Res. Lett.*, 8, 035020,
3 doi:10.1088/1748-9326/8/3/035020, 2013.
4

5 Mishra, U., and Riley, W.J.: Active-layer thickness across Alaska: Comparing observation-
6 based estimates with CMIP5 earth system model predictions, *Soil Sci. Soc. Am. J.*, 78, 894–
7 902, 2014.
8

9 Mora-Vallejo, A., Classens, L., Stoorvogel, J., and Heuvelink, G.B.M.: Small scale digital soil
10 mapping in southeastern Kenya, *Catena*, 76, 44–53, 2008.
11

12 [O'Brien, R.: A caution regarding rules of thumb for variance inflation factors, *Quality and*](#)
13 [*Quantity*, 41, 673–690, 2007.](#)
14

15 Peters-Lidard, C.D., Pan, F., and Wood, E.F.: A re-examination of modeled and measured soil
16 moisture spatial variability and its implications for land surface modeling, *Adv. Water*
17 *Resour.*, 24, 1069–83, 2001.
18

19 Pau, G.S.H., Bisht, G., and Riley, W.J.: A reduced-order modeling approach to represent
20 subgrid-scale hydrological dynamics for land-surface simulations: application in a polygonal
21 tundra landscape, *Geosci. Model Dev.*, 7, 2091–2105, 2014.
22

23 Ping, C.L., Michaelson, G.J., Kimble, J.M., Romanovsky, V.E., Shur, Y.L., Swanson, D.K., and
24 Walker, D.A.: Cryogenesis and soil formation along a bioclimate gradient in Arctic North
25 America, *J. Geophys. Res.*, 113, G03S12, 2008.
26

27 Ping, C.L.: Gelisols: Part I. Cryogenesis and state factors of formation, *Soil Horizons*, 54, 1–5,
28 2013.
29

30 Riley, W.J., and Shen, C.: Characterizing coarse-resolution watershed soil moisture
31 heterogeneity using fine-scale simulations, *Hydrol. Earth Syst. Sci.*, 18, 2463–2483, 2014.
32

33 Roecker, S.M., and Thompson, J.A.: Scale effects on terrain attribute calculation and their use as
34 environmental covariates for digital soil mapping, in *Digital Soil Mapping, Progress in Soil*
35 *Science*, edited by: Boettinger, J.L., Howell, D.W., Moore, A.C., Hartemink, A.E., and
36 Kienast-Brown, S., Springer, Berlin, 55–66, 2010.
37

38 Rodriguez-Iturbe, I., Vogel, G.K., Rigon, R., Entekhabi, D. Castelli, F., and Rinaldo, A.: On the
39 spatial organization of soil moisture fields, *Geophys. Res. Lett.*, 22, 2757–60, 1995.
40

41 Ryu, D., and Famiglietti, J.S.: Characterization of footprint-scale surface soil moisture variability
42 using Gaussian and beta distribution functions during the Southern Great Plains 1997
43 (SGP97) hydrology experiment, *Water Resour. Res.*, 41, W12433,
44 doi:10.1029/2004wr003835, 2005.
45

46 SAS Institute: Base SAS 9.3 Procedures Guide: SAS Institute, Inc. Cary, NC, 2011.

1
2 Smith, M.P., Zhu, A., Burt, J.E. and Stiles, C.: The effects of DEM resolution and neighborhood
3 size on digital soil survey, *Geoderma*, 137, 58–69, 2006.
4
5 Scenarios Network for Alaska and Arctic Planning: <http://www.snap.uaf.edu>, access: 6 March,
6 2014, 2014.
7
8 Tang, J.Y., Riley, W.J., Koven, C.D., and Subin, Z.M.: CLM4-BeTR, a generic biogeochemical
9 transport and reaction module for CLM4: model development, evaluation, and application,
10 *Geosci. Model Dev.*, 6, 127-140, 2013.
11
12 Tarnocai, C., and Bockheim, J.G.: 2011. Cryosolic soils of Canada: Genesis, distribution, and
13 classification, *Can. J. Soil Sci.*, 91,749–762, 2011.
14
15 Thompson, J.A., Bell, J.C., and Butler, C.A.: 2001. Digital elevation model resolution: Effects
16 on terrain attribute calculation and quantitative soil–landscape modeling, *Geoderma*, 100,
17 67–89, 2001.
18
19 Thompson, J.A., and Kolka, R.K.: Soil carbon storage estimation in a forested watershed using
20 quantitative soil landscape modeling, *Soil Sci. Soc. Am. J.*, 69, 1086–1093, 2005.
21
22 Thompson, J.A., Pena-Yewtukhiw, E.M., and Grove, J.H.: Soil–landscape modeling across a
23 physiographic region: Topographic patterns and model transportability, *Geoderma*, 133, 57–
24 70, 2006.
25
26 Todd-Brown, K.E.O., Randerson, J.T., Post, W.M., Hoffman, F.M., Tarnocai, C., Schuur,
27 E.A.G., and Allison, S.D.: Causes of variation in soil carbon predictions from CMIP5 Earth
28 system models and comparison with observations, *Biogeosciences*, 10, 1717–1736, 2013.
29
30 [Torn, M.S., Swanston, C.W., Castanha, C., and Trumbore, S.E.: Storage and turnover of organic](#)
31 [matter in soil, In: Senesi, N., Xing, B., and Huang, P.M. \(eds.\) *Biophysico-chemical*](#)
32 [*Processes Involving Natural Nonliving Organic Matter in Environmental Systems*, pp. 219-](#)
33 [272, John Wiley & Sons, New Jersey, USA, 2009.](#)
34 Vasques, G.M., Grunwald, S. Comerford, N.B., and Sickman, J.O.: Regional modelling of soil
35 carbon at multiple depths within a subtropical watershed, *Geoderma*, 156, 326–336, 2010.
36
37 Vasques, G.M., Grunwald, S. and Myers , D.B.: Influence of the spatial extent and resolution of
38 input data on soil carbon models in Florida, USA, *J. Geophys. Res.*, 117, G04004,
39 doi:10.1029/2012JG001982, 2012.
40
41 Webster, R., and Oliver, M.A.: Sample adequately to estimate variograms of soil properties, *J.*
42 *Soil Sci.*, 43,117–192, 1992.
43
44 Webster, R., and Oliver, M.A.: *Geostatistics for environmental scientists*, John Wiley & Sons,
45 UK, 2007.
46

1 Western, A.W., and Boschl, G.: On the spatial scaling of soil moisture, *J. Hydrol.*, 217, 203–224,
2 1999.
3
4 Western, A.W., Grayson, R.B., and Bloschl, G.: Scaling of soil moisture: A hydrologic
5 perspective, *Annu. Rev. Earth Planet. Sci.*, 30,149–180, 2002.
6
7 Wilson, J.P., and Gallant, J.C.: Digital terrain analysis, In: Wilson, J.P., and Gallant, J.C. (eds.)
8 *Terrain Analysis*, pp. 1–27, John Wiley & Sons, New York, 2000.
9
10 Zhu, A.X., Hudsonb, B., Burta, J., Lubichc, K., and Simonsond, D: Soil Mapping Using GIS,
11 Expert Knowledge, and Fuzzy Logic, *Soil Sci. Soc. Am. J.*, 65, 1463-1472, 2001.

Table 1: Environmental predictors and predicted variance of soil organic carbon stocks across different spatial scales. Black dots represent statistically significant ($P < 0.05$) predictors.

<i>Environmental Variables</i>	<i>Spatial Scale</i>							
	<i>50 m</i>	<i>100 m</i>	<i>200 m</i>	<i>500 m</i>	<i>1 km</i>	<i>2 km</i>	<i>5 km</i>	<i>10 km</i>
Elevation	•	•	•	•	•	•	•	•
Slope								•
Aspect	•							
SWI	•			•			•	
STI				•				
Temperature	•	•	•	•	•	•	•	•
PET	•	•	•	•	•	•	•	•

Barren	•	•	•				•	•
Scrub	•	•	•	•	•	•	•	•
Herb						•		
Pasture						•	•	
Glacial Moraine				•	•		•	•
Fluvial				•	•	•	•	•
Undifferentiated Mosaic							•	
Coastal delta						•		•
Volcanic mountain						•	•	
Predicted variance	476.5	354	304	263	261	257	256.4	258

Table 2: Modeled variogram parameters of predicted soil organic carbon stocks at various spatial scales.

<i>Spatial scale</i> (m)	<i>Nugget</i> (Ln kg m ⁻²) ²	<i>Sill</i> (Ln kg m ⁻²) ²	<i>Nugget/Sill</i> (%)	<i>Range</i> (km)
50.00	0.12	0.45	27	1,484
100.00	0.07	0.26	27	1,094
200.00	0.04	0.20	19.5	1,126
500.00	0.03	0.16	18	1,124

1,000.00	0.03	0.15	19	1,078
2,000.00	0.04	0.17	21	1,170
5,000.00	0.03	0.16	20.5	1,174
10,000.00	0.03	0.16	19	1,129

1 **Figure Legends**

2 Figure 1. Spatial distribution of observation sites, histogram, and summary statistics of observed
3 SOC stocks across Alaska.

4 Figure 2. The spatial scaling approach used in this study. The black dot represents the location of
5 an SOC sampling site, and the rectangles denote different pixel sizes of environmental variables
6 (the rectangles do not represent exact scale).


7 Figure 3. Control of environmental factors on SOC stocks as a function of spatial scale. Each dot
8 is a median geographically weighted regression coefficient (β) for a particular spatial scale. The
9 fitted functions are: (a) Elevation $\beta = -0.0009 + 0.0003 \cdot (1 - \exp(-0.0089 \cdot s))$, adjusted $R^2 = 0.83$,

1 $P < 0.004$, (b) Temperature $\beta = -0.11 + 0.037 \cdot (1 - \exp(-0.012 \cdot s))$, adjusted $R^2 = 0.94$, $P < 0.001$,
2 (c) Potential evapotranspiration $\beta = 0.066 + 0.187 \cdot \exp(-0.035 \cdot s)$, adjusted $R^2 = 0.97$,
3 $P < 0.001$, (d) Scrub $\beta = 0.395 - 0.0696 \cdot s$, adjusted $R^2 = 0.85$, $P < 0.001$.

4 Figure 4. Change of variogram parameters with scale (a), and calculated variograms of log-
5 transformed predicted SOC stocks (b) at spatial scales of 50 m, 100 m, 200 m, 500 m, 1 km,
6 2 km, 5 km, and 10 km. Fitted functions are: Sill = $0.163 + 0.763 \cdot \exp(-0.0197 \cdot s)$, adjusted $R^2 =$
7 0.98 , $P < 0.001$; Nugget = $0.032 + 0.201 \cdot \exp(-0.016 \cdot s)$, adjusted $R^2 = 0.98$, $P < 0.001$.

8 Figure 5. Variance of soil organic carbon stocks as a function of spatial scale. Each dot is
9 predicted variance of SOC stock at each spatial scale across Alaska. The fitted function is: SOC
10 variance = $260.61 + 422.42 \cdot \exp(-0.0318 \cdot s)$, adjusted $R^2 = 0.98$, $P < 0.001$.

11 Figure 6. Higher-order moments of predicted SOC stocks as a function of spatial scale (a), and
12 relationships between predicted mean SOC stocks and its (b) variance, (c) skewness, and (d)
13 kurtosis within the range of predicted mean values of SOC stocks.



1
2
3
4
5
6

Figure 1:





Figure 2:



1




Figure 3:





Figure 4:




Figure 5:




Figure 6: