Dear Dr. Niemann,

We would like to thank for your and other reviewers' valuable comments and constructive suggestions. In our response below, we replied to your comments in detail point by point and explained how we have modified the manuscript for publication in Biogeosciences. Comments and suggestions are shown in black and our responses are shown in blue.

Comments to the Author:

Dear Dr. Chuang,

thank you very much for handing in a revised version of your MS. The content has improved substantially and I only have a few technical details open that I would like you to address. Additionally, one of the reviewers provided some general thoughts.

Helge Niemann

Page Line 12/13: mark inserted sentence with commas: "It is surprising that at many sites, particularly within Groups 1 and 2 in the inner zone of both lagoons (1CEL, 2CEL, 3CEL and 1CH), high concentrations....

Page 10, line 20 ff: modify accordingly: There are several possible explanations for these observations: Firstly.... Secondly.... Thirdly...

Page 14, L31: ions in formula are wrong/incomplete. Sulphate is SO4 (4 in lower case and smaller than the font size than O) 2- (2- in upper case with lower font size than O). Check also chlorinity

Response:

These technical corrections have been revised in the manuscript.

Further thoughts by a reviewer:

1. Why is a lagoon/estuary that is receiving excess sulfate emitting more methane that other mangrove systems? The addition of sulfate would enhance the oxidative capacity of the system.

Response:

Indeed this is a good question and precisely why our study is interesting and important as it is not expected that a system with high sulfate will produce so much methane. As we discuss in the paper our data suggest that the reasons for that are (1) high organic matter and particularly non-competitive substrates and (2) The shallow depth of methane production in many sites and the high concentrations result in bubble formation which escape the sediment area with high oxidative capacity and hence a relatively small fraction of the bubbles (methane gas) gets dissolved and can oxidize. We discuss these issues in the paper.

2. What is the concentration of the freshwaters where the anhydrite was dissolved? To increase the pore water concentration in these coastal sediments by mM amounts requires a significant contribution of sulfate. I would guess the groundwater would contain 5-10 mM sulfate. Is that reasonable?

Response:

Young et al., (2008) have collected groundwater samples in Celestún Lagoon and reported the water chemistry in detail. Some groundwater samples has sulfate concentrations as high as 32 mM.

3. It remains unclear to me why AOM would be inactive even if there is a lot of competition for sulfate. In other systems with high OM and high methane, the competition is eliminated and both systems enabled by spatially separating the organoclastic and AOM zones. Both processes are possible.

Response:

We are not claiming that AOM is inactive just that in our model this process is not capable of reducing methane concentrations to a degree that we see large depletions in methane. In other words it is highly possible that the dissolved organic matter pool which was not measured in this study contains high abundance of competitive and non-competitive substrates to sustain high methane concentrations resulting in a relatively negligible role of AOM in mangrove dominated systems in comparison to other systems. Again the gas formation and short residence time in the sediment due to shallow formation depth, low gas dissolution and fast release all contribute to lower relative impact of AOM.

Other studies such as Lee et al. (2008) also detected the co-existence of porewater sulfate and methane in dwarf red mangrove habitats (Twin Cays, Belize) (Fig. R1) and in that setting like at our study site it was not able to be spatially separated the organoclastic and AOM zones.



Fig. R1: Porewater profiles of sulfate and methane reported in dwarf red mangrove habitats (Twin Cays, Belize) (modified from Lee et al., 2008). Dotted lines indicate base of homogenous surface soil layer at 10 cm. The filled and empty symbols represent two different sampling months.

4. I am still suspect that mangrove systems would have such a high relative abundance of noncompetitive substrates in comparison to salt marshes and places with high vegetative detritus. Fascinating.

Responses:

Indeed this is fascinating and not expected but this is consistent with our data and seen in other studies as well. There are only a few studies for the relative abundance or concentrations of non-competitive substrates in the organic carbon pool reported in the literatures for mangrove systems, salt marshes and places with high vegetative detritus. But where studied the possibility of utilizing noncompetitive substrates for methanogenesis in sulfate containing mangrove sediments have been shown (e.g., Mohanraju et al. (1997), Lyimo et al. (2000) and Purvaja and Ramesh (2001)).

Technical Corrections:

Page 8, line 26: replace 'chlorine' with 'chloride'

Page 9, line 32: put a space between 'sulfate' and 'and'

Response:

These technical corrections have been revised in the manuscript.

I understand that you cannot fully address these concerns, but I'd like to encourage you to address them in the text and provide at least an "educated guess".

We try to emphasize the above responses throughout the manuscript. It is a very interesting system and more studies should be carried out in such systems in the future. Maybe some microbial genetics as well.

With the best wishes, Helge Niemann

References:

Lee, R., Porubsky, W., Feller, I., McKee, K., and Joye, S.: Porewater biogeochemistry and soil metabolism in dwarf red mangrove habitats (Twin Cays, Belize), Biogeochemistry, 87, 181-198, 2008.

Lyimo, T. J., Pol, A., den Camp, H. J. M., Harhangi, H. R., and Vogels, G. D.: Methanosarcina semesiae sp nov., a dimethylsulfide-utilizing methanogen from mangrove sediment, Int. J. Syst. Evol. Micr., 50, 171–178, 2000.

Mohanraju, R., Rajagopal, B. S., and Daniels, L.: Isolation and characterization of a methanogenic bacterium from mangrove sediments, J. Mar. Biotechnol., 5, 147–152, 1997.

Purvaja, R. and Ramesh, R.: Natural and anthropogenic methane emission from coastal wetlands of South India, Environ. Manage., 27, 547–557, 2001.

Young, M. B., Gonneea, M. E., Fong, D. A., Moore, W. S., Herrera-Silveira, J., and Paytan, A.: Characterizing sources of groundwater to a tropical coastal lagoon in a karstic area using radium isotopes and water chemistry, Mar. Chem., 109, 377–394, 2008.

1 2 3	Methane and Sulfate Dynamics in Sediments from Mangrove-dominated Tropical Coastal Lagoons, Yucatán, Mexico
3 4	P-C Chuang ¹ M B Young ² A W Dale ³ L C Miller ² J A Herrera-Silveira ⁴ A
5	Paytan ^{1,5}
6	
7	[1] Department of Earth and Planetary Sciences, University of California Santa Cruz, 1156
8	High St., Santa Cruz, CA 95064, United States
9	[2] U.S. Geological Survey, 345 Middlefield Rd, MS 434, Menlo Park, CA 94025, United
0	States
1	[3] GEOMAR Helmholtz Centre for Ocean Research Kiel, Wischhofstr. 1-3, 24148 Kiel,
2	Germany
3	[4] CINVESTAV-IPN, Unidad Mérida, A.P. 73 CORDEMEX, Mérida, Yucatán, Mexico
4	[5] Institute of Marine Sciences, University of California Santa Cruz, 1156 High St., Santa
5	Cruz, CA 95064, United States

- 16
- 17 Correspondence to: P.-C. Chuang (pchuan2@ucsc.edu; pcchuang2@gmail.com)

- 1 -

1 Abstract

2 Porewater profiles in sediment cores from mangrove-dominated coastal lagoons (Celestún and

- 3 Chelem) on the Yucatán Peninsula, Mexico, reveal the widespread coexistence of dissolved
- 4 methane and sulfate. This observation is interesting as it is not expected that a system with high
- 5 sulfate content will produce much methane and if some methane is produced it typically is
- 6 oxidized through anaerobic oxidation. To explain the observations we used a numerical
- 7 transport-reaction model. The model suggests that methane in the upper sediments is produced
- 8 in the sulfate reduction zone at rates ranging between 0.012 and 31 mmol $m^{-2} d^{-1}$, concurrent
- 9 with sulfate reduction rates between 1.1 and 24 mmol SO_4^{2-} m⁻² d⁻¹. These processes are
- 10 supported by high organic matter content in the sediment and particularly non-competative
- substrates. Indeed sediment slurry incubation experiments show that non-competitive substrates such as trimethylamine (TMA) and methanol can be utilized for microbial methanogenesis at the study sites. The model also indicates that a significant fraction of methane is transported to the sulfate reduction zone from deeper zones within the sedimentary
- 15 column by rising bubbles and gas dissolution. Combined the shallow depths of methane
- 16 production and the fast rising methane gas bubbles reduce the likelihood for oxidation
- 17 <u>allowing</u> a large fraction of the methane formed in the sediments escapes to the overlying water

Deleted: A

Deleted: The model also indicates that a significant fraction of methane is transported to the sulfate reduction zone from deeper zones within the sedimentary column, by rising bubbles and gas dissolution.

Deleted: S

Deleted: Our results suggest that

Deleted: By combining field measurements with transport-reaction modeling, we are able to demonstrate that sediments in coastal lagoons within mangrove ecosystems are characterized by shallow methane production and accumulation depths, likely due to non-competitive substrate utilization in near-surface sediments and extensive bubble transport and dissolution; this may favor high methane emission rates.

20 1 Introduction

column.

18

19

21 Wetlands are the largest natural source of methane (CH₄) to the atmosphere, accounting for 22 between 20-25% of the global atmospheric methane budget (Fung et al., 1991; Whalen, 2005). 23 Methane produced in wetlands is primarily biogenic, originating from microbial activity in 24 anaerobic sediments and soil. Since sulfate-reducing bacteria outcompete methanogens for 25 common substrates (Oremland and Polcin, 1982), freshwater wetlands typically have much 26 higher methane fluxes to the atmosphere than brackish to fully marine wetlands (Bartlett et al., 27 1987; Bartlett et al., 1985; Segarra et al., 2013). Marine and estuarine sediments are generally 28 characterized by comparatively low rates of methanogenesis with a methane production and 29 accumulation zone located deeper within the sediment pile below the sulfate reduction zone 30 (Holmer and Kristensen, 1994; Martens and Val Klump, 1984; Poulton et al., 2005; Segarra et 31 al., 2013). In these marine or estuarine systems methane that diffuses upwards towards the 32 sediment surface can be oxidized both anaerobically (AOM) and aerobically within the 33 sediments and in the water column, reducing emissions to the atmosphere (Whalen, 2005).

34 Despite brackish to marine salinities, methane fluxes comparable to those measured in

- 2 -

Deleted: arising

1 freshwater wetlands have been reported for coastal mangrove-dominated lagoon systems in 2 several places around the world, including Florida (Barber et al., 1988), Puerto Rico (Sotomayor et al., 1994), India (Biswas et al., 2004; Biswas et al., 2007; Purvaja and Ramesh, 3 2000; Purvaja and Ramesh, 2001; Ramesh et al., 1997; Ramesh et al., 2007; Verma et al., 4 5 1999), Tanzania (Kristensen et al., 2008), Thailand (Lekphet et al., 2005), China (Alongi et al., 6 2005), Andaman Islands (Linto et al., 2014) and Australia (Call et al., 2015). The anaerobic 7 and organic-rich sediments found in these systems provide a suitable environment for 8 methanogenesis, yet the extensive supply of sulfate from seawater should favor sulfate 9 reducers over methanogens in the shallow sections of the sediments (Kristensen et al., 2008; 10 Lee et al., 2008). There are, however, several possible ways for coastal mangrove lagoons to 11 sustain relatively high methane fluxes despite high sulfate concentrations. For example, if the microbial activity of sulfate reducers is high and sulfate replenishment from the overlying 12 13 water is slow, sulfate may become depleted in the upper centimeters of the sediment, thus 14 allowing methanogens to occur close to the sediment surface. Additionally, methanogens can 15 co-exist with sulfate reducers when non-competitive substrates (those used only by 16 methanogens and not by sulfate reducers) are available. Moreover, in some systems methane 17 may migrate from deeper in the sediment to shallower depth and to the water column. 18 Typically, a large percentage of the methane produced in sediments is oxidized prior to 19 reaching the atmosphere, and in shallow-water systems, the oxidation takes place primarily in the sediments and not in the water column (Martens and Valklump, 1980; Mitsch and 20 Gosselink, 2000; Weston et al., 2011; Segarra et al., 2013, 2015). However, accumulation and 21 22 transport of methane in gas bubbles reduces the exposure time of methane to oxidants such as 23 oxygen and sulfate, allowing a large fraction of gas to escape the sediment (Barnes et al., 2006; Martens and Valklump, 1980). 24

25 The objective of this study was to examine porewater methane distributions within the 26 sediments of two mangrove-dominated coastal lagoons in Mexico and relate them to sulfate 27 concentrations of the sediments. We aim to gain a better understanding of the factors 28 controlling the methane flux from coastal mangrove-dominated lagoon sediments. To this end, 29 we applied a numerical transport-reaction model based on Wallmann et al. (2006) and Chuang 30 et al. (2013) to simulate porewater methane and sulfate concentration profiles. We also 31 performed sediment slurry incubation experiments to test the effect of competitive and 32 non-competitive substrates on methanogenesis in the lagoon sediments. The results provide 33 quantitative data on methane dynamics in coastal mangrove-dominated lagoon systems and 34 highlight their importance as methane sources to the atmosphere.

- 3 -

35

Deleted: these

1 2 Study sites

2 Fieldwork was conducted in two mangrove-dominated coastal lagoons located on the western

3 Yucatán Peninsula, Mexico (Figure 1). The typical climatological pattern for this area consists

4 of a dry season (March-May), a rainy season (June-October) during which the majority of the

5 annual rainfall (>500mm) occurs, and the "nortes" season (November-February), which is

6 characterized by moderate rainfall (20-60mm) and intermittent high wind speeds greater than

7 80 km hr⁻¹ (Herrera-Silveira, 1994).

8 Celestún Lagoon (20°52'N, 90°22'W) is long, narrow, and relatively shallow (average depth =

9 1.2 m). The inner and middle sections of the lagoon always have lower salinities than the section near the mouth due to year-round discharge of brackish groundwater from multiple 10 11 submarine springs (Young et al., 2008). Salinity within the lagoon fluctuates seasonally, with 12 salinity in the inner zone ranging from 8.9 to 18.2 during the course of this study, grading out 13 to marine salinities at the mouth of the lagoon (Young et al., 2008). The lagoon is surrounded 14 by 22.3 km² of a well-developed mangrove forest, and has experienced relatively little 15 disturbance from human development and/or pollution such as wastewater discharge (Herrera-Silveira et al., 1998). Sediments in Celestún consist primarily of autochthonous 16 17 carbonate ooze.

Chelem Lagoon (21°15'N, 89°45W) (average depth = 0.7 m), in contrast, receives very little 18 19 groundwater input and the surrounding area has been heavily impacted by urban development. 20 Salinity in Chelem ranges from brackish to hypersaline (24.8 - 40.3 during the study period), 21 and vegetation surrounding the lagoon consists of scrub mangrove forest (Young et al., 2008). 22 The construction of Yucalpeten Harbor in 1969 (Valdes and Real, 1998) increased the 23 circulation and resulted in sandy marine sediments entering the lagoon. Sediments in Chelem 24 deposited since 1969 consist of a heavily bioturbated mix of sands and autochthonous carbonate ooze, with a large number of shells of living and dead burrowing organisms (Valdes 25 and Real, 1998). In the following text, CEL and CH denote cores collected from Celestún 26 27 Lagoon and Chelem Lagoon, respectively.

28

29 3 Sampling and analytical methods

30 3.1 Porewater solutes

Sediment cores were collected along lengthwise transects in both lagoons during the three
 different seasons; April 2000 (dry season), December 2000 (nortes season), and October 2001

33 (late rainy season). Duplicate samples (1 1CH Oct01 and 1 2CH Oct01) were collected at

- 4 -

Formatted: Superscript

1 station 1CH in Chelem lagoon. Sediments were sampled using hand-held acrylic push cores (7 2 cm inner diameter) either 30 or 60 cm in length. The push cores had holes drilled along the side 3 at 2 cm intervals, which were sealed with electrical tape prior to sampling. Subsamples for porewater methane analysis were collected in the field immediately after core collection from 4 5 the holes along the sides of the push cores, using plastic 3 mL syringes with the needle attachment end removed. The sediment plugs from the syringes were immediately extruded 6 7 into 20 mL glass Wheaton bottles and sealed with blue butyl stoppers and aluminum crimp 8 caps. 3 mL of degassed Milli-Q water and 0.3 mL of saturated mercuric chloride (HgCl₂) 9 solution were added to create a slurry and halt all biological activity within the sample.

After subsampling, the cores were capped, the holes were resealed, and the cores were transported back to the lab for sectioning and porewater extraction. The cores were extruded and sliced into 2.5 cm depth intervals in an anaerobic glove bag under an N_2 atmosphere and transferred into centrifuge tubes for porewater extraction. Core length was measured immediately after collection and just prior to extrusion in order to correct for compaction during transport. Average compaction was 6% of the total core length, and never exceeded 20%.

Porewater for sulfate (SO₄²⁻) and chloride (Cl⁻) analyses was extracted by centrifuging all the 17 sediment from each depth interval and filtering the porewater through sterile 0.20 µm syringe 18 19 filters. Samples were kept frozen in 20 mL acid-cleaned glass scintillation vials until analysis. 20 Porewater sulfate and chloride concentrations were measured by ion chromatography using a 21 Dionex DX-500 IC equipped with an Ionpac AS9-HC column (4mm) and AG9-HC (4mm) 22 guard column. The samples were diluted 5-fold with Milli-Q water prior to analysis in order to 23 bring the sulfate and chloride within the appropriate analytical range for the ion 24 chromatograph.

25 Methane concentrations for all samples were measured on an SRI 310 Gas Chromatograph 26 (GC) equipped with a flame ionization detector and an Alltech Haysep S 100/120 column (6' x 27 1/8" x 0.085"). Helium was used as the carrier gas at a flow rate of 15 ml/min and the column 28 and detector temperatures were maintained at 50 °C and 150 °C, respectively. Peak integration 29 was performed using Peak Simple NT software. Methane gas standards were prepared by 30 diluting 100% methane in helium, and five standards bracketing the range of sample 31 concentrations were measured at the beginning, middle, and end of each set of analyses. 32 Average standard error of repeat injections of standards throughout a sample run (between 2 to 33 6 hours of continuous analysis) was 1.8% (n=152). Porewater methane concentration in the 34 sediment core subsamples was determined after vigorously shaking of the sealed serum bottles 35 containing the sediment slurries to ensure complete mixing, followed by at least 3 minutes of - 5 -

standing equilibration time to ensure that the porewater methane was fully equilibrated with the
 headspace in the serum bottles. A small volume of headspace (0.25-0.5 μL) was drawn out of
 each serum bottle using a gas-tight syringe, and analyzed for methane concentration on the SRI
 GC. The total volume of porewater in each sample was calculated using the difference
 between the total wet weight of the sediment minus the dry weight of the sediment, correcting
 for the added water and HgCl₂ solution.

7

8 3.2 Sediment slurry incubation experiments

9 Sediment slurry incubations were performed in order to examine changes in methane production over different time intervals and at different substrate concentrations (Table 1). 10 11 Incubations consisted of three competitive substrates (H₂, acetate, formate), two 12 non-competitive substrates (methanol, trimethylamine (TMA)), and four types of controls. The 13 controls (preparation methods are described below) consisted of an un-amended sediment 14 control under anaerobic conditions, an un-amended aerobic control (partial oxygen headspace), 15 a killed control in which the sediment was autoclaved to kill all living organisms in the 16 sediment, and a chemical control in which biological methanogenesis was inhibited through the 17 addition of 2-bromoethanesulfonic acid (BES) to a final concentration of 40 mM within the 18 slurry. Triplicate bottles were prepared for each condition (controls and substrate additions), and methane headspace concentrations were measured at 3-4 time intervals over the course of 19 20 29 days.

21 All the sediment slurries were prepared semi-anaerobically by homogenizing the sediment in a 22 blender with an artificial seawater mixture in a 1:1 ratio under continuous flow of nitrogen gas. 23 Large pieces of leaves, twigs, and shells were removed from the sediment prior to 24 homogenization. 70 mL glass Wheaton bottles were flushed with nitrogen gas for 1 minute 25 prior to the addition of the sediment slurry. 30 mL of slurry was then added to each bottle under 26 continuous nitrogen flow, and the bottles were sealed using blue butyl rubber stoppers and 27 aluminum crimp seals. Substrate additions were made by injecting the substrate solution into the bottle immediately after sealing the bottles, except for the H₂ gas treatment and the aerobic 28 29 control. For the addition of H₂, the entire headspace of the bottles was flushed with 100% H₂ gas. After each headspace sampling the H₂ gas removed by microbial activity in the sediment 30 31 was replaced by inserting a gas tight syringe filled with 100% H₂ gas into the bottles, and allowing the gas to be drawn into the bottles until equilibrium pressure was reached. The 32 33 aerobic controls were prepared like the anaerobic un-amended controls, except that 8 mL (20% 34 of the total headspace) of 100% O_2 was added to the bottles immediately after they were sealed. In order to ensure that the sediment slurries remained aerobic, 100% O2 was added to the 35 - 6 -

- 1 bottles throughout the incubation period. The sediment slurries were kept at room temperate
- 2 (22°C) and agitated continuously on a shaker table throughout the course of the incubations.
- 3 Headspace samples (0.25 mL) were extracted from the bottles at each time interval using a

4 gas-tight syringe. Methane concentrations were measured on an HP 5730A GC equipped with a

- 5 flame ionization detector. GC calibration and creation of standard curves were based on
- 6 successive dilutions of 100% methane. Analytical error was approximately 5% for methane
- 7 concentrations below 10 ppm-v (446 nM), and less than 3% for methane concentrations above
- 8 10 ppm-v as determined by repeat analyses of standards and samples.
- 9
- 10 4 Results

11 4.1 Porewater concentrations of dissolved species

12 Representative porewater methane profiles were plotted alongside sulfate profiles in Fig. 2 and Fig. A1. Profiles were assigned to one of four profile-types based on the relation between 13 14 methane and sulfate distributions down core (see below). Considerable spatial and temporal 15 variability in porewater chemistry was observed with no systematic seasonal differences in 16 concentration trends. For example, porewater methane concentrations varied by up to three 17 orders of magnitude in both lagoons, even between sites in close proximity to each other (i.e. 1CEL and 2CEL, Oct01; 1CH and 2CH, Dec00), and at the same station sampled during 18 19 different seasons (i.e. 2CEL Dec00, Oct01; 1CH Apr00 and Oct01). No consistent differences 20 were evident between the stations at the sides of the lagoons and those located in the center of 21 the lagoons, or between stations located in the inner zone of the lagoons and those located near 22 the mouth. For instance, methane above calculated saturated concentrations (1.1 and 1.3 mM) 23 was observed in cores 1CEL Jul02 (the inner zone of Celestún lagoon) and 14CEL Dec00 24 (near the mouth of Celestún lagoon). This is particularly interesting because the water column 25 at the mouth of the lagoon has much higher salinities than the water of the inner zone (Young 26 et al., 2008). The variability (both spatial and temporal) in the porewater methane 27 concentrations and in the spatial and temporal distribution of profile types suggest a very dynamic system where both concentration and distribution patterns in the porewater vary 28 29 constantly (spatially throughout the lagoons and temporally at distinct sites). Such variability is 30 indicative of rapid methane production and efflux rates.

Porewater sulfate concentrations ranged from 0.21 to 35.3 mM in Celestún lagoon and from 4.13 to 33.5 mM in Chelem lagoon and showed different trends (Fig. 2; Fig. A1). In many of the cores a negative relation between methane and sulfate was observed. Specifically, higher sulfate was associated with lower methane in cores located near the mouth of the lagoons

- 7 -

- 1 (16CEL_Jul02, 16CEL_Oct01, 14CEL_Oct01, 14CEL_Jul02 and 5CH_Apr00) and lower
- 2 sulfate with high methane in the inner zone of the lagoons (e.g. cores 1CEL_Jul02,

3 1CEL_Dec00, 3CEL_Jul02, 3CEL_Apr00, 1_1CH_Oct01, and 1_2 CH_Oct01).

4 The relationship between porewater salinity (represented by chloride concentration), methane

- 5 and sulfate concentrations was spatially and temporally variable (Fig. 3). Generally, higher
- 6 sulfate concentrations were associated with higher chloride in cores located near the mouth of
- 7 the lagoons and lower sulfate with lower chloride in the inner zone of the lagoons (Fig. 3A).
- 8 Despite these general trends there were no clear consistent relationships between methane and
- 9 chloride (Fig. 3B) and sulfate and methane (Fig. 3C) when the data was considered collectively.

10 The lack of consistent trends suggests multiple processes impacting the distribution of methane

11 and sulfate. These include physical processes, such as mixing and dilution by seawater or

12 groundwater, and biological processes such as sulfate reduction, methanogenesis and methane 13 oxidation. Brackish groundwater enters <u>Celestún lagoon through at least 30 subsurface</u> 14 discharge points (Young et al., 2008), and the chloride profiles suggest that some of this 15 groundwater may seep through the sediments, resulting in localized decline in porewater 16 salinities.

To account for mixing with seawater or freshwater and to extract information on the <u>biological</u> and chemical processes controlling the distribution of porewater solutes, the observed sulfate depletion ($[SO_4^{2-}_{dep}]_{OBS}$) relative to seawater was calculated as the difference between the expected sulfate concentration contributed from seawater (based on porewater chloride concentration) and the measured sulfate concentration:

22

23
$$[SO_4^{2-}]_{OBS} = \frac{[SO_4^{2-}]_{(SW)}}{[Cl^{-}]_{(SW)}} \times [Cl^{-}]_{(measured)} - [SO_4^{2-}]_{(measured)}$$
(1)

where 0.05171 is taken as the $\frac{[SO_4^2^-](SW)}{[CI^-](SW)}$ ratio (Pilson, 1998). Positive values indicate that sulfate has been removed from the porewater, most likely through sulfate reduction while negative values indicate an external source of sulfate not associated with chloride that is other than seawater, in this case the groundwater (see discussion below).

Based on the observed trends in sulfate depletion, when considered together with methane, four
different porewater trends can be described, referred to as Groups 1 through 4 here (Fig. 2, Fig.

31 A1). The majority of profiles fell into Group-1 (10 cores); these profiles showed positive

32 sulfate depletion profiles (e.g. sulfate consumption or loss) with methane profiles mirroring the

33 sulfate concentration profiles (methane production or input). The peaks for methane and sulfate

Deleted: the

Deleted: n

1 depletion occurred at the same depth as the lowest measured sulfate concentrations. In Group-2 2 (7 cores), sulfate depletion also showed positive values (sulfate consumption) but not throughout the core. In some cores sulfate depletion was close to zero at shallow depths and 3 then increased with depth and in other cores positive sulfate depletion values appeared at the 4 5 surface of the sediments and then decreased to almost zero at deeper depths. Methane 6 concentrations for this group showed no clear relation to the sulfate profiles. In Group-3 (3 7 cores), sulfate depletion showed negative values (e.g. sulfate addition). The values became 8 more negative toward the deeper sediment starting from zero right at the surface suggesting 9 that sulfate was being added from the bottom of the sediment section. In Group-4 (4 cores), 10 there was almost no sulfate depletion (sulfate concentrations similar to seawater) from the 11 surface to the deeper depths and methane concentrations were low (< 0.25 mM) increasing at 12 depth, indicating a deeper source of methane.

13

14 4.2 Sediment slurry incubation experiments

15 All the sediment slurries with added substrates showed an increase in headspace methane 16 concentration that was significantly greater than those observed with either the un-amended aerobic and anaerobic controls or the treated controls (Figure 4). The greatest increases in 17 18 headspace methane concentration were seen with additions of the two noncompetitive 19 substrates, TMA and methanol. The H₂ treatment showed the next highest methane production 20 rate, followed by formate and acetate. Of the four control conditions, the un-amended, 21 anaerobic treatment had the highest overall increase in headspace methane concentration. The aerobic treatment had an initial higher increase in headspace methane concentration than the 22 23 un-amended, anaerobic treatment, although there was no detectable change in the headspace 24 methane concentration in the aerobic treatment between 150 and 700 hours. Both the 25 autoclaved and BES treatments did not show any changes in headspace methane concentration 26 greater than the instrumental detection limits. The maximum methane production rates for each 27 treatment are listed in Table 1.

28

29 5 Discussion

30 5.1 Co-existence of methane and sulfate in sediments

Seawater transport into the sediment by diffusion and bioirrigation due to the activity of burrowing animals has clear effects on porewater solutes. These processes are a source of seawater sulfate_and mask sulfate loss by microbial reduction. Although, as indicated above, considerable variability in porewater profile distribution trends was observed, and different

-9-

1 profile types were found throughout the lagoons, certain trends were more common at distinct 2 locations. Specifically, sites characterized by sulfate addition from input of seawater into the 3 sediment (cores in Group-4) were found primarily near the mouth of both lagoons where low 4 methane was associated with near-zero sulfate depletion. Negative sulfate depletion (Group-3), 5 on the other hand, which indicates the presence of porewater that is enriched in sulfate relative to chloride, was seen primarily in the middle zone of Celestún Lagoon where groundwater 6 7 springs rich in sulfate due to anhydrite dissolution are present, as reported by Perry et al. (2009; 8 2002). Positive sulfate depletion profiles co-occurring with methane (Groups 1 and 2) were 9 seen throughout the lagoons but mostly at sites in the inner zone of both lagoons, suggesting 10 significant sulfate reduction at rates higher than the replenishment from sulfate rich 11 groundwater or from the overlying seawater and a source of methane to the shallow sections of 12 the sediment.

13 It is surprising that at many sites, particularly within Groups 1 and 2 in the inner zone of both 14 lagoons (1CEL, 2CEL, 3CEL and 1CH), high concentrations of methane and sulfate 15 co-occurred at the same depth in the sediment. Co-existence of methanogenesis and sulfate 16 reduction is not normally observed because sulfate reduction is more energetically favorable 17 than methanogenesis, and sulfate reducers should outcompete methanogens for common 18 substrates such as hydrogen and acetate (Oremland and Polcin, 1982; Jørgensen and Kasten, 19 2006). Moreover, anaerobic oxidation of methane (AOM) coupled with sulfate reduction at the 20 base of the sulfate reducing zone should further deplete methane (Capone and Kiene, 1988; 21 Valentine and Reeburgh, 2000). There are several possible explanations for these observations; 22 Firstly, the high methane concentrations measured in the sulfate rich porewater may be 23 supplied by a rapid non-diffusive mechanism from below the sulfate reduction zone (like rising 24 gas bubbles), limiting the exposure time to AOM. Secondly, methane may be produced in-situ 25 at these depths supported by a high abundance of competitive substrates in the sulfate reduction 26 zone hence sustaining both methanogenesis and sulfate reduction (Holmer and Kristensen, 27 1994). Thirdly, methanogens may instead be able to thrive on various non-competitive 28 substrates (Oremland and Polcin, 1982; Wellsbury and Parkes, 2000; Lee et al., 2008; Taketani 29 et al., 2010). Indeed, use of non-competitive substrates by methanogens, including methanol, 30 trimethylamines and dimethylsulfide, has been reported for mangrove sediments, coastal 31 lagoons and continental shelf sediments (Ferdelman et al., 1997; Lyimo et al., 2000; Mohanraju 32 et al., 1997; Purvaja and Ramesh, 2001; Torres-Alvarado et al., 2013; Maltby et al., 2016). Our 33 slurry incubation experiments demonstrated that the methanogenic community at Celestún is 34 capable of using a wide range of substrates, including H₂, acetate, formate, methanol, and 35 trimethylamine (Fig. 4). Both methanol and trimethylamine are not utilized by sulfate reducers,

- 10 -

Deleted: chlorine

Deleted:

1 which could allow methanogens to thrive in the sulfate reduction zone (Fig. 4). The use of 2 non-competitive substrates by the methanogenic community has important implications for methane fluxes to the atmosphere as it allows for methane production at shallow depths in the 3 sediment and reduces the potential for complete oxidation of methane. Although processes and 4 5 trends similar to those described above have been reported for other mangrove sediments (e.g., 6 Lee et al., 2008; Purvaja and Ramesh, 2001), the co-occurrence of sulfate and methane and 7 related biogeochemical reactions in these reports remain qualitative in nature. In the following 8 section, we use a transport-reaction model to better quantify the processes controlling methane 9 fluxes from the sediments in these mangrove-dominated tropical coastal lagoons.

10

11 5.2 Model set-up and application to mangrove-dominated coastal lagoon sediments

12 In order to understand methane production and consumption and how these processes relate to 13 sulfate dynamics in the lagoon sediments, we used two different approaches to simulate 14 methane and sulfate porewater profiles.

15 In the first approach, a transport-reaction model was applied to profiles of Group-1 where 16 methane and sulfate co-occur with no indication of groundwater sulfate input and where sulfate 17 reduction surpasses sulfate addition from seawater (Fig. 2; Fig. A1). Data in Group-1 have 18 positive net sulfate depletion rates indicative of sulfate reduction. The sulfate depletion is seen 19 within the zone where methane concentrations are high. In these cores the net sulfate depletion 20 rates can be used to derive the minimum methanogenesis rates (see model details in the 21 Appendix). Reactions considered in this first approach include organic matter degradation via 22 heterotrophic sulfate reduction, methane production via methanogenesis and methane addition 23 from gas bubble dissolution (Haeckel et al., 2004; Chuang et al., 2013).

A second approach (detailed in the Appendix) was used for simulating the profiles for Group-2,

Group-3 and Group-4 which show no positive net sulfate depletion rates when integrated over the core length. These sites are affected by groundwater input or by considerable irrigation and input of seawater. Here, the link between sulfate and methane reactions is less clear and hard to

28 quantify directly.

The following equation was solved to quantify the rates of reaction and transport of dissolved
methane and sulfate in the upper 20 cm of the sediments in both approaches (Berner, 1980;
Boudreau, 1997):

32

33 $\Phi \cdot \frac{\partial C}{\partial t} = \frac{\partial \left(\Phi \cdot Ds \cdot \frac{\partial C}{\partial x}\right)}{\partial x} - \frac{\partial \left(\Phi \cdot v \cdot C\right)}{\partial x} + \Phi \cdot$ 34 R_c

- 11 -

(2)

2 where x is sediment depth, t is time, Φ is porosity, Ds is the solute-specific diffusion 3 coefficient in the sediment, C is the concentration of methane or sulfate in the porewater, v is the burial velocity of porewater and R_C is the sum of reactions affecting C (Table A1). Solutes 4 were simulated in moles L⁻¹ of porewater (M). Details of all the reaction terms and parameters 5 6 and how they were derived for each of the two approaches are given in the Appendix. The 7 model assumes steady state conditions to constrain methanogenesis rates at each site. 8 Considering the observed variability in porewater distributions non-steady state simulations 9 would be desirable, yet this would require continuous monitoring of porewater sulfate, methane and chloride concentrations to evaluate temporal changes in sulfate depletion at each site. This 10 11 time-series data is unavailable, hence the modeled 'instantaneous' rates bear uncertainties that 12 currently cannot be quantified accurately.

13 Model derived sulfate depletion and sulfate and methane concentrations are shown in Fig. 2 14 and Fig. A1. Modeled porewater data for Group-1 (the most common trend) show that methane 15 generated from organic matter degradation within the upper sediments is a more important 16 methane source than methane diffusing from below and gas bubble dissolution, as further seen 17 in the results of 1CEL_Jul02 and the sensitivity analysis from 2CEL_Jul02 (Fig. 5A). In 1CEL Jul02, for example, gas dissolution of methane transported from deeper sediments is not 18 19 necessary at all to achieve a good model fit to the data, and *in-situ* methanogenesis alone can 20 reproduce methane concentrations similar to the measured data even though methane 21 concentrations are oversaturated (> 1.1 mM (in situ solubility)) (Fig. 2). In contrast, the modeled methane profile for 2CEL Jul02 (black dashed line) arguably does require the 22 23 inclusion of methane from gas dissolution (R_{MB}) (Fig. 5A). In Fig. 5A, the gray dashed and solid lines represent only gas dissolution in the methane reaction terms (no methanogenesis 24 25 within the modeled 20 cm column) using different gas dissolution constants (k_{MB} values are 0.2 yr⁻¹ and 0.5 yr⁻¹ respectively). The model results shown as the gray dashed line simulate the 26 27 methane concentrations below 10 cm depth, whereas those shown by the gray solid line 28 reproduce methane concentrations in the upper 5 cm, but neither reproduces the data 29 throughout the whole core. Comparing results considering methanogenesis and gas dissolution 30 (black solid line) and methanogenesis only (black dashed line), it is clear that both 31 methanogenesis and some gas dissolution are needed for reproducing the methane distribution observed in core 2CEL Jul02. This illustrates the complexity of controlling processes and the 32 dynamic nature and resulting temporal variability in methane fluxes at this and the other sites 33 34 in the lagoons.

35

1

- 12 -

1 5.3 Model derived depth-integrated turnover rates and fluxes

- 2 Table 2 lists the calculated depth-integrated turnover rates and fluxes for the individual cores.
- 3 For profiles in Group-1, methane sources include methanogenesis within the upper 20 cm
- 4 and/or methane transported from deeper sections (>20 cm) via bubble transport and dissolution.
- 5 Methane can be supported fully by methanogenesis without gas bubble dissolution within the
- 6 modeled upper 20 cm in cores 1CEL Dec00, 1CEL Jul02, 1 1CH Oct01 and 1 2CH Oct01.
- 7 Gas bubble transport from deeper sediments and its dissolution contributes more methane than
- 8 methanogenesis in cores 1CEL_Apr00, 1CEL_Oct01, 2CEL_Dec00 and 3CEL_Jul02.
- 9 Methane sinks include <u>fluxes into</u> the water column or methane diffusion into deeper sediments
- 10 (>20 cm) and oxidation. Our model shows that the major sink for methane, however, is <u>efflux</u>
- 11 to the water column accounting for over 90% of methane produced within the upper 20 cm (e.g.
- 12 1CEL Apr00, 1CEL Oct01, 3CEL Apr00 and 3CEL Jul02). Model derived methane fluxes
- 13 to the water column are listed in Table 2 ($F_{methane (top)}$) and range from 0.012-20 mmol CH₄ m⁻²
- 14 d^{-1} . These are similar to or up to two orders of magnitude larger than fluxes reported for other
- mangrove lagoon systems in Florida (0.02 mmol CH₄ m⁻² d⁻¹, Barber et al., 1988; Harriss et al., 16 1988), Australia (0.03-0.52 mmol CH₄ m⁻² d⁻¹, Kreuzwieser et al., 2003), and India (5.4-20.3
- 16 1988), Australia (0.03-0.52 mmol CH₄ m⁻² d⁻¹, Kreuzwieser et al., 2003), and India (5.4-20.3 17 mmol CH₄ m⁻² d⁻¹, Purvaja and Ramesh, 2001). Since all methane depth profile types were
- in million official in a straticity and realistic 2001). Since an include deput prome types were
- 18 observed throughout the year with no differences in spatial and temporal distribution (seasons
- 19 and sampling locations), our results support the idea that methane fluxes in coastal mangrove
- 20 lagoon systems respond very dynamically to environmental stimuli.
- Sulfate sinks include heterotrophic sulfate reduction and AOM, although the model suggests that AOM plays a minor role compared to heterotrophic sulfate reduction. Sulfate reduction ranges from 1.1 to 24 mmol SO_4^{2-} m⁻² d⁻¹ and is the major sink for both sulfate and organic
- carbon in most cores. Sulfate reduction accounts for 2.2 to 48 mmol C $m^{-2} d^{-1}$ of total anaerobic carbon respiration, which is in the same range of values listed in Kristensen et al.
- 26 (2008) for most mangrove sediments.

27 Mangrove forests are known to be highly productive systems with the capacity to release high 28 concentrations of dissolved organic matter (DOM) to surrounding sediments and porewaters 29 (Kristensen et al., 2008). Tree litter and subsurface root growth provide further significant 30 inputs of organic carbon to mangrove sediments which are unique for this type of system. The rate of organic matter mineralization (R_{POC} ; Eq. A6) derived from sulfate depletion ranges from 31 3.2 mmol C m⁻² d⁻¹ to 110 mmol C m⁻² d⁻¹. Although our modeling approach for determining 32 33 degradation rates is not without uncertainty, it is more accurate than rates derived from 34 down-core trends in organic matter content because of temporal variability in accumulation

- 13 -

Deleted: dissolution and

Deleted: emissions

Deleted: emission

1 rates in this area (Gonneea et al., 2004). Particulate organic matter will also contain a high 2 amount of refractory carbon that is not easy to quantify and separate from the bulk pool. The 3 derived degradation rates likely represent the more labile particulate components and labile 4 DOM that was not considered (or measured) in this study. The high calculated organic carbon 5 oxidation rates derived here are thus not unexpected since mangrove systems in general (e.g. Dittmar et al., 2006; Dittmar and Lara, 2001; Lee 1995; Odum and Heald, 1975) and the 6 7 lagoons in Yucatan in particular are dominated by high concentrations of DOM, a large fraction 8 of which is likely to be labile (Young et al., 2005).

9 Depth-integrated methane production or consumption rates (R_{CH_4}) and net sulfate inputs 10 $(R_{SO_4^{2-}})$ calculated from Eq. (A9) and (A10) for cores in Group-2, Group-3 and Group-4 are listed in Table 2. The methane and sulfate net production/consumption rates ranged from 11 -0.060 to 11 mmol CH₄ m⁻² d⁻¹ and -69 to 21 mmol SO₄²⁻ m⁻² d⁻¹ (Negative values indicate net 12 sulfate or methane consumptions while positive values indicate production or addition from 13 14 external sources). Although sulfate depletion values for cores in Group-2 are positive (e.g. net 15 sulfate reduction), sulfate concentrations at some depths of the porewater are relatively high, suggesting continuous sulfate input from deeper within the sediments or from seawater. Cores 16 17 in Group-3 and Group-4 show negative or zero sulfate depletion that likely results from high 18 rates of sulfate addition from groundwater (Group-3) or seawater (Group-4), thus prohibiting 19 accurate calculation of sulfate reduction and methanogenesis rates. Although, in theory, H₂S 20 oxidation is a possible source for the excess sulfate, we believe that sulfate-rich groundwater 21 input is a more likely source due to correlation between excess sulfate and excess Sr which has 22 been previously described for groundwater in this region (Young et al., 2008). Perry et al. 23 (2002) identified dissolution of evaporites within the freshwater lens at some Yucatán sites as a probable source of excess sulfate in groundwater using the sulfate-to-chloride ratio 24 $(100 \times \frac{[SO_4^{-1}]}{[C]^{-1}}$ (SO). Ratios higher than seawater (average seawater is 10.3) are expected where 25 26 gypsum/anhydrite dissolution occurs (Perry et al. 2002). Another indicator is the Sr/Cl ratio, 27 which is invariably higher in the Yucatan groundwater than in seawater and indicates 28 dissolution of celestite (from evaporite) and/or aragonite (Perry et al., 2002). The region east 29 and south of Lake Chichancanab, referred to as the Evaporite Region by Perry et al. (2002), is 30 characterized by distinctive topography and high sulfate groundwater concentrations (Perry et 31 al., 2002). The groundwater from the Lake Chichancanab area flows northward into the 32 Celestún Estuary which can be recognized by the progressive decrease in the ratio $\frac{|C|^2}{|C|^2}$ groundwater in water from southeast to northwest (Perry et al., 2009). Some groundwater 33

 $[SO_4^{2-}]$

- 14 -

Deleted: ₄/Cl)

Deleted: [
Deleted: SO ₄	
Deleted: Cl]	
Deleted:]	
Deleted: [
Deleted: SO ₄	
Deleted: Cl]	
Deleted:]	

1 samples with sulfate concentrations as high as 32 mM were reported in Young et al. (2008) and

2 the Sr and sulfur trends for Celestún lagoon (Young et al., 2008) are consistent with our

3 interpretation that gypsum/anhydrite dissolution in groundwater is the source of excess sulfate
4 in the porewater of Group-3 in Celestún lagoon. Due to the impact of groundwater, our sulfate

5 reduction and methanogenesis rates estimated using the model are minimum rates and 6 independent rates of groundwater discharge into each core are needed for obtaining more 7 realistic estimates in these sites.

8 In addition to depth-integrated rates, Table 3 also includes maximum methanogenesis/methane 9 production (Max- R_M) and sulfate reduction/consumption (Max- R_{SR}) rates solved by Eq. 2 in the 10 model. Interestingly, the maximum methane production rates estimated from TMA, methanol 11 and H₂ additions to sediments in the slurry incubations (Table 1) are similar to model derived 12 Max- R_M at station 16CEL (Table 3), which is the site from which sediments were collected for 13 the slurry incubations. The rates in the TMA, methanol and H₂ treatments from the slurry 14 incubations (Table 1) and in some of our stations are higher than methane production rates 15 from previously reported coastal freshwater and brackish wetland sediments that were measured using radiolabeled acetate and bicarbonate in slurries (Segarra et al., 2013). 16

17 Modeled Max- R_M in some cores were 1-2 orders of magnitude higher than rates derived from 18 the sediment slurry incubations (e.g., cores 1CEL Jul02, 1 1CH Oct01, 2CEL Oct01 and 19 14CEL Dec00). Although heterotrophic sulfate reduction generally dominates organic matter 20 degradation, Max- R_M values are even higher than the maximum sulfate reduction rates in some 21 cores (1 1CH Oct01, 1 2CH Oct01 and 1CH Dec00). Both the methanogenesis rates 22 measured in the sediment slurry incubations and the modeled maximum methanogenesis rates 23 in this study area were much higher than those reported for some mangrove systems (e.g., 24 Thailand, Kristensen et al., 2000; Malaysia, Alongi et al., 2004; Australia, Kristensen and 25 Alongi, 2006) but similar to other sites in India (Ramesh et al., 2007).

26 AOM is expected to play an important role in oxidizing methane in tropical porewaters with 27 abundant methane and sulfate (Biswas et al., 2007). However, our model results and sensitivity 28 analyses indicate that AOM is insufficient to prevent methane escape to the bottom water, 29 probably because of the abundant organic matter available for sulfate reducers to use instead of methane. In our sensitivity tests (using core 1CEL Oct01 as an example), if AOM is allowed to 30 31 be responsible for sulfate and methane consumption (no heterotrophic sulfate reduction and 32 methanogenesis; $R_{SD} = R_{AOM}$) then methane concentrations would decrease to negative values 33 (gray solid lines in Fig. 5B), which is inconsistent with observations. Although based on our data it is not possible to accurately quantify methane oxidation by calculating the relative 34 35 proportion of sulfate loss due to heterotrophic sulfate reduction and/or AOM, our model results - 15 -

Deleted: has

Deleted: T

1 suggest AOM plays a minor role in this setting. It is also possible to have high rates of methane 2 production and also AOM in the sediments but this is not captured as methane loss because 3 there is more production than depletion. In addition the formation of gas bubbles and short 4 residence time in the sediment due to shallow formation depth, low gas dissolution and fast 5 release all contribute to lower the relative impact of AOM. Other studies such as Lee et al. 6 (2008) also detected the co-existence of porewater sulfate and methane in dwarf red mangrove 7 habitats (Twin Cays, Belize) and in that setting like at our study site it was not able to be 8 spatially separated the methanogenesis and AOM zones. Future investigations on the role of 9 AOM in these dynamic mangrove-dominated tropical coastal lagoons are needed (e.g., Thalasso et al, 1997; Raghoebarsing et al., 2006; Lee et al., 2008; Kristensen et al., 2008; Beal 10 11 et al., 2009; Silvan et al., 2011; Segarra et al., 2013).

12

13 6 Conclusions

14 The variable trends observed in sediment porewater chemistry from mangrove-dominated 15 tropical coastal lagoons, in Yucatán, Mexico indicate a very dynamic system spatially and 16 temporally throughout the year. This can be explained by multiple controlling parameters 17 including physical processes such as mixing and dilution with seawater or groundwater, gas 18 bubble rise and dissolution and microbial processes which operate at different rates during 19 different times at all sites. Although our modeling suggests that organic carbon degradation 20 rates are dominated by heterotrophic sulfate reduction in these cores, methanogenesis both in 21 shallow and deeper sediments is also prevalent. The co-occurrence of methane and sulfate 22 reduction (documented by sulfate depletion) in shallow sediments in this system is explained 23 by high methane production rates supported by some combination of non-competitive 24 substrates and ample dissolved and labile organic matter in the shallow sediments as well as 25 the contributions of methane from deeper sediment through gas rise and dissolution. Model 26 results demonstrate that the largest sink for methane in these sediments is efflux to the water 27 column. Build-up of methane at shallow depths may reduce the fraction of methane that is 28 oxidized prior to entering the water column, thereby increasing the flux at the sediment-water 29 interface. This shallow methane pool may also encourage methane flux through bubble release, 30 which can result in a larger fraction of the methane reaching the atmosphere without being lost 31 to oxidation. Specifically, the ability of the microbial community in these sediments to use non-competitive substrates may allow for methane production in the upper sections of the 32 sediment, potentially contributing to the higher than expected atmospheric methane flux 33 34 measured from mangrove-dominated tropical coastal lagoons.

- 16 -

1

2 Appendix: Modeling procedure used in the evaluation of porewater observations from

3 sediments in mangrove-dominated tropical coastal lagoons, Yucatán, Mexico

4 Details of the modeling procedure and parameters used are described here. The following 5 reactions are considered in the model:

6 7

8

9

11

13

Heterotrophic sulfate reduction (R_{SR}) : $2CH_2O + SO_4^{2-} \rightarrow 2HCO_3^{-} + H_2S$ (R1)

10 Methanogenesis (R_M) : 2CH₂O \rightarrow CO₂+CH₄ (R2)

12 Gas bubble dissolution
$$(R_{MB})$$
: CH₄(g) \rightarrow CH₄(aq) (R3)

The net reaction terms (R_C in Eq. 2) are given in Table A1, boundary conditions are listed in Table A2, best-fit model parameters are given in Table A3 and model derived concentration profiles are shown in Fig. 2 and Fig. A1.

(A1)

17 In Eq. (2), sediment porosity decreases with depth due to steady-state compaction:

18
19
$$\Phi = \Phi_f + (\Phi_0 - \Phi_f) \cdot e^{-px}$$

20

where Φ_f is the porosity below the depth of compaction (0.78 for Celestún and 0.83 for Chelem), Φ_0 is porosity at the sediment surface (0.90 for Celestún and 0.89 for Chelem) and *p* (1/15 cm⁻¹) is the depth attenuation coefficient. These parameters were determined from the measured porosity data at each site or at a nearby site (Eagle, 2002).

Under the assumption of steady state compaction, the burial of porewater was calculated as inBerner (1980):

$$28 \qquad v = \frac{\Phi_f \cdot w_f}{\Phi} \tag{A2}$$

29

27

where w_f is the sedimentation rate of compacted sediments calculated from excess ²¹⁰Pb data (0.25 cm yr⁻¹ for Celestún and 0.35 cm yr⁻¹ for Chelem; Gonneea et al. (2004)). Sediment burial results in the downward movement of both sediment particles and porewater relative to the sediment water interface.

- 17 -

1 The sediment diffusion coefficient of each solute (Ds) was calculated according to Archie's law

2 considering the effect of tortuosity on diffusion (Boudreau, 1997):

$$\begin{array}{ll}
4 & D_s = \Phi^2 \cdot D_M \\
5 &
\end{array} \tag{A3}$$

6 where D_M is the molecular diffusion coefficient at the in situ temperature, salinity and pressure 7 (Table A1) calculated according to Boudreau (1997). We used the same tortuosity coefficient 8 (Φ^2 corresponding to m = 3 in Archie's law) as reported by Wallmann et al. (2006) for 9 fine-grained sediments.

Since net sulfate consumption is observed in Group-1 profiles (Fig. 2; Fig. A1), we used the following calculations to obtain net sulfate depletion rates (R_{SD} ; mmol SO₄²⁻ cm⁻³ yr⁻¹). R_{SD} is proportional to the difference between modeled ($C(SO_4^{2-}_{dep}))$) and measured concentrations ($C(SO_4^{2-}_{dep})_{OBS}$):

3

15
$$R_{SD} = k_{SD} \cdot \left(C \left(SO_4^{2-}{}_{dep} \right)_{OBS} - C \left(SO_4^{2-}{}_{dep} \right) \right)$$
(A4)

16

17 The corresponding kinetic constant is set to be high $(k_{SD} \ge 100 \text{ yr}^{-1})$ to ensure that simulated 18 concentrations are very close to measured values. R_{SD} implicitly includes R_{SR} as well as 19 anaerobic oxidation of methane (R_{AOM}) :

20 21

$$CH_4 + SO_4^{2-} \rightarrow HCO_3^{-} + HS^{-} + H_2O$$
(R4)

22

23 The numerical modeling procedure outlined in Wallmann et al. (2006) is used as a basis to 24 simulate the rate of sedimentary organic carbon degradation (R_{POC}) by sulfate reduction and 25 methanogenesis. Since the measured organic matter content in both lagoons showed evidence 26 for a change in depositional pattern over time (Gonneea et al., 2004 and Eagle, 2002), these 27 measurements cannot be used for reliable organic matter degradation calculations. Hence, R_{SR} 28 (Eq. A5 below) was first calculated and then used to estimate R_{POC} (Eq. A6) and subsequently 29 to derive R_M (Eq. A7). Here, we assume the three reactions (R1, R2 and R4) co-occur in the 30 sulfate reduction zone such that the net reaction for methanogenesis and AOM (reactions 31 R2+R4) is equal to carbon respiration by heterotrophic sulfate reduction (reaction R1). In other 32 words, $R_{SD} = 0.5 R_{POC}$.

33 To approximate the fraction of R_{POC} due to R_M and R_{SR} , a Michaelis-Menten kinetic limitation

34 term is applied to Eq. (A5-A7) (Wallmann et al., 2006):

2
3
$$R_{SR} = R_{SD} = 0.5 \cdot R_{POC} \cdot f_{SO_4^2}$$
 (A5)

5
$$R_{POC} = \frac{R_{SD}}{0.5 \cdot f_{C} \cdot f_{SO_{4}^{2^{-}}}}$$
 (A6)

4

1

7
$$R_M = 0.5 \cdot R_{POC} \cdot (1 - f_{SO_4^{2-}})$$
 (A7)

8

9 where $f_{SO_4^{2-}} = \frac{c_{SO_4^{2-}}}{c_{SO_4^{2-}} + K_{SR}}$ is the Michaelis–Menten rate-limiting term for sulfate reduction.

10 At sites where methanogenesis was insufficient to simulate the measured methane data, 11 methane was added as an external source by dissolution of gas bubbles (Chuang et al., 2013). 12 Gas bubbles were observed in the field. The rate of dissolution of the gas bubbles (R3) rising 13 through the sediment ($CH_{4(g)} \rightarrow CH_4$) was also considered as (Haeckel et al., 2004):

14 15

16
$$R_{MB} = k_{MB} \cdot \left(L_{MB} - C_{CH_4}\right) \qquad \text{if } CH_4 \le L_{MB}$$
(A8)

17

where L_{MB} is the in situ methane gas solubility concentration calculated using the algorithm of (Duan et al., 1992a; Duan et al., 1992b) and the site-specific salinity, temperature and pressure. R_{MB} depends on the first-order rate constant k_{MB} , which is a fitting parameter that lumps together gas dissolution in addition to diffusion of dissolved gas in the bubble tubes and walls.

Since sulfate depletion profile trends in Group-2, Group-3 and Group-4 show evidence of groundwater or seawater input with no positive depth integrated net sulfate depletion rates, the second approach for determining net methane and sulfate reaction rates for porewater data in these three groups is summarized as:

26

27
$$R_{CH_4} = k_{CH_4} \cdot (C_{CH_4} - C_{CH_4})$$
 (A9)

29
$$R_{SO_4^{2-}} = k_{SO_4^{2-}} \cdot (C_{SO_4^{2-}OBS} - C_{SO_4^{2-}})$$
 (A10)

30

31 Net methane and sulfate reaction rates are set to be proportional to the difference between 32 modeled (C_{CH_4} and $C_{SO_4^{2-}}$) and measured concentrations (C_{CH_4} and $C_{SO_4^{2-}}$ or $C_{SO_4^{2-}}$). The

- 19 -

1 corresponding kinetic constants k_{CH_4} and $k_{SO_4^{2-}}$ are listed in Table A3.

2 Methane fluxes at the boundaries were calculated using the model as follows:

3

$$dL_{av}(\mathbf{x})$$

$$4 \quad F_{CH_4}(x) = \Phi(x) \cdot \left(v(x) \cdot C_{CH_4} - D_S \cdot \frac{uC_{CH_4}(x)}{dx} \right)$$
(A11)

5

16

6 where x = 20 cm is the bottom of the simulated core and x = 0 cm is the sediment-water 7 interface.

8 Fixed concentrations were imposed for all solutes at the upper and lower boundaries to values 9 measured at or near the sediment-water interface and at 20 cm. The method-of-lines was used 10 to transfer the set of finite difference equations of the spatial derivatives of the coupled partial 11 differential equations to the ordinary differential equation solver (NDSolve) in MATHEMATICA v. 7.0, using a grid spacing which increased from ca. 0.015 cm at the 12 13 sediment surface to 0.38 cm at depth. Since most of the porewater profiles were fitted directly, 14 only a few years of simulation time (5 yrs) was needed to achieve steady state. Mass balance was typically better than 99.9 %. 15

17 Acknowledgements

18 We thank the staff of the DUMAC Celestún station and the students of CINVESTAV for 19 assistance with laboratory space, lodging, and field work. We thank Tom Lorenson and Ron 20 Oremland of the Menlo Park, CA USGS for facility use and analyses for slurry 21 incubations. This work was funded by Consejo Nacional de Ciencia y Tecnología Ref: 4147-P 22 T9608, 32356T, and CONABIO Ref: B019 to J H-S, NSF INT 009214214 to AP, a Stanford 23 Graduate Fellowship and Lieberman Fellowship to MY and a fellowship of the Postdoctoral 24 Research Abroad Program, sponsored by the National Science Council, Taiwan to PCC (now: 25 MOST (the Ministry of Science and Technology)). We thank John Pohlman and an anonymous reviewer for their thoughtful comments and the associate editor (Helge Niemann) for handling 26 27 the manuscript.

28

29 References

30 Alongi, D. M., Sasekumar, A., Chong, V. C., Pfitzner, J., Trott, L. A., Tirendi, F., Dixon, P., and

31 Brunskill, G. J.: Sediment accumulation and organic material flux in a managed mangrove

32 ecosystem: estimates of land-ocean-atmosphere exchange in peninsular Malaysia, Marine

- 1 Geology, 208, 383-402, 2004.
- 2 Alongi, D. M., Pfitzner, J., Trott, L. A., Tirendi, F., Dixon, P., and Klumpp, D. W.: Rapid
- 3 sediment accumulation and microbial mineralization in forests of the mangrove Kandelia
- 4 candel in the Jiulongjiang Estuary, China, Estuar. Coast. Shelf S., 63, 605–618, 2005.
- Barber, T. R., Burke, R. A., and Sackett, W. M.: Difusive flux of methane from warm wetlands,
 Global Biogeochem. Cy., 2, 411–425, 1988.
- 7 Barnes, J., Ramesh, R., Purvaja, R., Nirmal Rajkumar, A., Senthil Kumar, B., Krithika, K.,
- 8 Ravichandran, K., Uher, G., and Upstill-Goddard, R.: Tidal dynamics and rainfall control N2O
- 9 and CH4 emissions from a pristine mangrove creek, Geophys. Res. Lett., 33, L15405,
 10 doi:10.1029/2006GL026829, 2006.
- ,
- Bartlett, K. B., Harriss, R. C., and Sebacher, D. I.: Methane flux from coastal salt marshes, J.
 Geophys. Res.-Atmos., 90, 5710–5720, 1985.
- Bartlett, K. B., Bartlett, D. S., Harriss, R. C., and Sebacher, D. I.: Methane emissions along a
 salt marsh salinity gradient, Biogeochemistry, 4, 183–202, 1987.
- Beal, E. J., House, C. H., and Orphan, V. J.: Manganese- and Iron-Dependent Marine Methane
 Oxidation, Science, 325, 184-187, 2009.
- Berner, R. A.: Early Diagenesis. A Theoretical Approach, Princeton University Press, Princeton,
 NJ, USA, 241 pp., 1980.
- 19 Biswas, H., Mukhopadhyay, S. K., De, T. K., Sen, S., and Jana, T. K.: Biogenic controls on the
- 20 air-water carbon dioxide exchange in the Sundarban mangrove environment, northeast coast of
- 21 Bay of Bengal, India, Limnol. Oceanogr., 49, 95–101, 2004.
- 22 Biswas, H., Mukhopadhyay, S. K., Sen, S., and Jana, T. K.: Spatial and temporal patterns of
- methane dynamics in the tropical mangrove dominated estuary, NE coast of Bay of Bengal,
 India, J. Marine Syst., 68, 55–64, 2007.
- Boudreau, B. P.: Diagenetic Models and Their Implementation: Modelling Transport and
 Reactions in Aquatic Sediments, Springer-Verlag, Berlin, 414 pp., 1997.
- 27 Call, M., Maher, D. T., Santos, I. R., Ruiz-Halpern, S., Mangion, P., Sanders, C. J., Erler, D. V.,
- 28 Oakes, J. M., Rosentreter, J., Murray, R., and Eyre, B. D.: Spatial and temporal variability of
- carbon dioxide and methane fluxes over semi-diurnal and spring-neap-spring timescales in a
 mangrove creek, Geochim. Cosmochim. Ac., 150, 211–225, 2015.
- 31 Capone, D. G. and Kiene, R. P.: Comparison of microbial dynamics in marine and freshwater
- 32 sediments: contrasts in anaerobic carbon catabolism1, Limnol. Oceanogr., 33, 725–749, 1988.
 - 21 -

- 1 Chuang, P.-C., Dale, A. W., Wallmann, K., Haeckel, M., Yang, T. F., Chen, N.-C., Chen, H.-C.,
- 2 Chen, H.-W., Lin, S., Sun, C.-H., You, C.-F., Horng, C.-S., Wang, Y., and Chung, S.-H.:
- 3 Relating sulfate and methane dynamics to geology: accretionary prism o_shore SW Taiwan,
- 4 Geochem. Geophy. Geosy., 14, 2523–2545, 2013.
- 5 Dittmar, T. and Lara, R. J.: Driving forces behind nutrient and organic matter dynamics in a
 6 mangrove tidal creek in North Brazil, Estuar. Coast. Shelf S., 52, 249–259, 2001.
- Dittmar, T., Hertkorn, N., Kattner, G., and Lara, R. J.: Mangroves, a major source of dissolved
 organic carbon to the oceans, Global Biogeochem. Cy., 20, GB1012,
 doi:10.1029/2005GB002570, 2006.
- Duan, Z., Møller, N., Greenberg, J., and Weare, J. H.: The prediction of methane solubility in
 natural waters to high ionic strength from 0 to 250 _C and from 0 to 1600 bar, Geochim.
 Cosmochim. Ac., 56, 1451–1460, 1992a.
- Duan, Z. H., Møller, N., and Weare, J. H.: An equation of state for the CH4-CO2-H2O
 system.1. pure systems from 0 _C to 1000 _C And 0 to 8000 bar, Geochim. Cosmochim. Ac.,
 56, 2605–2617, 1992b.
- Eagle, M.: Tracing organic matter sources in mangrove estuaries utilizing δ¹³C, δ¹⁵N and C/N
 Ratios. Master Thesis, Stanford University, Stanford, CA, USA, 2002.
- Ferdelman, T. G., Lee, C., Pantoja, S., Harder, J., Bebout, B. M., and Fossing, H.: Sulfate
 reduction and methanogenesis in a Thioploca-dominated sediment off the coast of Chile,
 Geochimica et Cosmochimica Acta, 61, 3065-3079, 1997.
- 21 Fung, I., John, J., Lerner, J., Matthews, E., Prather, M., Steele, L. P., and Fraser, P. J.:
- Threedimensional model synthesis of the global methane cycle, J. Geophys. Res.-Atmos., 96,
 13033–13065, 1991.
- 24 Gonneea, M. E., Paytan, A., and Herrera-Silveira, J. A.: Tracing organic matter sources and
- carbon burial in mangrove sediments over the past 160 years, Estuar. Coast. Shelf S., 61,
 211–227, 2004.
- Haeckel, M., Suess, E., Wallmann, K., and Rickert, D.: Rising methane gas bubbles form
 massive hydrate layers at the seafloor, Geochim. Cosmochim. Ac., 68, 4335–4345, 2004.
- Harriss, R. C., Sebacher, D. I., Bartlett, K. B., Bartlett, D. S., and Crill, P. M.: Sources of
 atmospheric methane in the south Florida environment, Global Biogeochemical Cycles, 2,
 231-243, 1988.
- 32 Herrera-Silveira, J.A.: Nutrients from uderground discharges in a coastal lagoon (Celestún,
 - 22 -

- Yucatán, México), Verhandlungen des Internationalen Verein Limnologie, 25, 1398–1403,
 1994.
- 3 Herrera-Silveira, J. A., Ramìrez R, J., and Zaldivar R, A.: Overview and characterization of the
- 4 hydrology and primary producer communities of selected coastal lagoons of Yucatán, México,
 5 Aquatic Ecosystem Health and Management, 1, 353–372, 1998.
- Holmer, M. and Kristensen, E.: Coexistence of sulfate reduction and methane production in an
 organic-rich sediment, Mar. Ecol.-Prog. Ser., 107, 177–184, 1994.
- 8 Jørgensen, B. B. and Kasten, S.: Sulfur Cycling and Methane Oxidation. In: Marine
- 9 Geochemistry, Schulz, H. D. and Zabel, M. (Eds.), Springer Berlin Heidelberg, Berlin,
 10 Heidelberg, 2006.
- Kreuzwieser, J., Buchholz, J., and Rennenberg, H.: Emission of Methane and Nitrous Oxide by
 Australian Mangrove Ecosystems, Plant Biology, 5, 423-431, 2003.
- 13 Kristensen, E., Andersen, F. Ã., Holmboe, N., Holmer, M., and Thongtham, N.: Carbon and
- nitrogen mineralization in sediments of the Bangrong mangrove area, Phuket, Thailand,
 Aquatic Microbial Ecology, 22, 199-213, 2000.
- 16 Kristensen, E. and Alongi, D. M.: Control by fiddler crabs (Uca vocans) and plant roots
- 17 (Avicennia marina) on carbon, iron, and sulfur biogeochemistry in mangrove sediment, Limnol.
 18 Oceanogr., 51, 1557-1571, 2006.
- Kristensen, E., Bouillon, S., Dittmar, T., and Marchand, C.: Organic carbon dynamics in
 mangrove ecosystems: a review, Aquat. Bot., 89, 201–219, 2008.
- 21 Lee, R., Porubsky, W., Feller, I., McKee, K., and Joye, S.: Porewater biogeochemistry and soil
- metabolism in dwarf red mangrove habitats (Twin Cays, Belize), Biogeochemistry, 87,
 181-198, 2008.
- Lee, S. Y.: Mangrove outwelling a review, Hydrobiologia, 295, 203–212, 1995.

- Lekphet, S., Nitisoravut, S., and Adsavakulchai, S.: Estimating methane emissions from
 mangrove area in Ranong Province, Thailand, Songklanakarin, J. Sci. Technol., 27, 153–163,
 2005.
- 28 Linto, N., Barnes, J., Ramachandran, R., Divia, J., Ramachandran, P., and Upstill-Goddard, R.
- C.: Carbon dioxide and methane emissions from mangrove-associated waters of the Andaman
 islands, Bay of Bengal, Estuar. Coast., 37, 381–398, 2014.
- 31 Lyimo, T. J., Pol, A., den Camp, H. J. M., Harhangi, H. R., and Vogels, G. D.: Methanosarcina
 - semesiae sp nov., a dimethylsulfide-utilizing methanogen from mangrove sediment, Int. J. Syst.
 - 23 -

- 1 Evol. Micr., 50, 171–178, 2000.
- 2 Maltby, J., Sommer, S., Dale, A. W., and Treude, T.: Microbial methanogenesis in the
- 3 sulfate-reducing zone of surface sediments traversing the Peruvian margin, Biogeosciences, 13,
- 4 283-299, 2016.
- Martens, C. S. and Klump, V. J.: Biogeochemical cycling in an organic-rich coastal marine
 basin.1. Methane sediment-water exchange processes, Geochim. Cosmochim. Ac., 44, 471–490,
- 7 1980.
- 8 Martens, C. S. and Klump, V. J.: Biogeochemical cycling in an organic-rich coastal marine 9 basin 4. An organic carbon budget for sediments dominated by sulfate reduction and
- 10 methanogenesis, Geochim. Cosmochim. Ac., 48, 1987–2004, 1984.
- 11 Mitsch, W.J., Gosselink, J.G.: Wetlands. Wiley, New York, p. 936, 2000.
- Mohanraju, R., Rajagopal, B. S., and Daniels, L.: Isolation and characterization of a
 methanogenic bacterium from mangrove sediments, J. Mar. Biotechnol., 5, 147–152, 1997.
- Odum, W. E. and Heald, E. J.: The detritus-based food web of an estuarine mangrove community, in: Estuarine Research, edited by: Cronin, L. E., New York, Academic Press, Inc.,
- 16 265–286, 1975.
- Oremland, R. S. and Polcin, S.: Methanogenesis and sulfate reduction competitive and non
 competitive substrates in estuarine sediments, Appl. Environ. Microb., 44, 1270–1276, 1982.
- Perry, E., Velazquez-Oliman, G., and Marin, L.: The hydrogeochemistry of the karst aquifer
 system of the northern Yucatan Peninsula, Mexico, Int. Geol. Rev., 44, 191–221, 2002.
- 21 Perry, E., Paytan, A., Pedersen, B., and Velazquez-Oliman, G.: Groundwater geochemistry of
- the Yucatan Peninsula, Mexico: constraints on stratigraphy and hydrogeology, J. Hydrol., 367,
 27–40, 2009.
- Pilson, M. E. Q.: An Introduction to the Chemistry of the Sea, Prentice Hall, Upper Saddle
 River, New Jersey, 431 pp., 1998.
- Poulton, Simon, W., Canfield, and Donald, E.: Development of a sequential extraction
 procedure for iron: implications for iron partitioning in continentally derived particulates.
 Elsevier B.V., 2005.
- Purvaja, R. and Ramesh, R.: Human impacts on methane emission from mangrove ecosystems
 in India, Reg. Environ. Change, 1, 86–97, 2000.
- 31 Purvaja, R. and Ramesh, R.: Natural and anthropogenic methane emission from coastal
- 32 wetlands of South India, Environ. Manage., 27, 547–557, 2001.

- 24 -

- 1 Raghoebarsing, A. A., Pol, A., van de Pas-Schoonen, K. T., Smolders, A. J. P., Ettwig, K. F.,
- 2 Rijpstra, W. I. C., Schouten, S., Damste, J. S. S., Op den Camp, H. J. M., Jetten, M. S. M., and
- 3 Strous, M.: A microbial consortium couples anaerobic methane oxidation to denitrification,
- 4 Nature, 440, 918-921, 2006.
- 5 Ramesh, R., Purvaja, G. R., Parashar, D. C., Gupta, P. K., and Mitra, A. P.: Anthropogenic
- forcing on methane e_ux from polluted wetlands (Adyar River) of Madras City, India, Ambio,
 26, 369–374, 1997.
- 8 Ramesh, R., Purvaja, R., Neetha, V., Divia, J., Barnes, J., and Upstill-Goddard, R.: CO2 and
- 9 CH4 emissions from Indian mangroves and its surrounding waters, in: Greenhouse gas and
- 10 carbon balances in mangrove coastal ecosystems, Proceedings, edited by: Tateda, Y., Gendai
- 11 Tosho, Kanagawa, Japan, 153–164, 2007.
- 12 Segarra, K. E. A., Comerford, C., Slaughter, J., and Joye, S. B.: Impact of electron acceptor
- availability on the anaerobic oxidation of methane in coastal freshwater and brackish wetland
 sediments, Geochimica et Cosmochimica Acta, 115, 15-30, 2013.
- 15 Segarra, K. E. A., Schubotz, F., Samarkin, V., Yoshinaga, M. Y., Hinrichs, K. U., and Joye, S.
- 16 B.: High rates of anaerobic methane oxidation in freshwater wetlands reduce potential
- 17 atmospheric methane emissions, Nat Commun, 6, 2015.
- 18 Sivan, O., Adler, M., Pearson, A., Gelman, F., Bar-Or, I., John, S. G., and Eckert, W.:
- Geochemical evidence for iron-mediated anaerobic oxidation of methane, Limnol. Oceanogr.,
 56, 1536-1544, 2011.
- Sotomayor, D., Corredor, J. E., and Morell, J. M.: Methane flux from mangrove sediments
 along the southwestern coast of Puerto-Rico, Estuaries, 17, 140–147, 1994.
- 23 Taketani, R., Yoshiura, C., Dias, A., Andreote, F., and Tsai, S.: Diversity and identification of
- methanogenic archaea and sulphate-reducing bacteria in sediments from a pristine tropical
 mangrove, Antonie van Leeuwenhoek, 97, 401-411, 2010.
- Thalasso, F., Vallecillo, A., García-Encina, P., and Fdz-Polanco, F.: The use of methane as a
 sole carbon source for wastewater denitrification, Water Research, 31, 55-60, 1997.
- 28 Torres-Alvarado, M., José Fernández, F., Ramírez Vives, F., and Varona-Cordero, F.: Dynamics
- of the methanogenic archaea in tropical estuarine sediments, Archaea, 13, 582646,
 doi:10.1155/2013/582646, 2013.
- 31 Valdes, D. S. and Real, E.: Variations and relationships of salinity, nutrients and suspended
- 32 solids in Chelem coastal lagoon at Yucatan, Mexico, Indian J. Mar. Sci., 27, 149–156, 1998.

- 25 -

- Valentine, D. L. and Reeburgh, W. S.: New perspectives on anaerobic methane oxidation,
 Environ. Microbiol., 2, 477–484, 2000.
- 3 Verma, A., Subramanian, V., and Ramesh, R.: Day-time variation in methane emission from
- 4 two tropical urban wetlands in Chennai, Tamil Nadu, India, Curr. Sci. India, 76, 1020–1022,
 5 1999.
- 6 Wallmann, K., Aloisi, G., Haeckel, M., Obzhirov, A., Pavlova, G., and Tishchenko, P.: Kinetics
- of organic matter degradation, microbial methane generation, and gas hydrate formation in
 anoxic marine sediments, Geochim. Cosmochim. Ac., 70, 3905–3927, 2006.
- 9 Wellsbury, P. and Parkes, R. J.: Deep biosphere: source of methane for oceanic hydrate, in:
- 10 Natural gas hydrate in oceanic and permafrost environments, edited by: Max, M. D., Springer,
- 11 New York, 91–104, 2000.
- 12 Weston, N., Vile, M., Neubauer, S., and Velinsky, D.: Accelerated microbial organic matter
- mineralization following salt-water intrusion into tidal freshwater marsh soils, Biogeochemistry,
 102, 135-151, 2011.
- Whalen, S. C.: Biogeochemistry of methane exchange between natural wetlands and the
 atmosphere, Environ. Eng. Sci., 22, 73–94, 2005.
- 17 Young, M. B., Gonneea, M. E., Herrerasilveira, J. A., and Paytan, A.: Export of Dissolved and
- 18 Particulate Carbon and Nitrogen From a Mangrove-Dominated Lagoon, Yucatán Peninsula,
- 19 Mexico, Int. J. Ecol. Environ. Sci., 31, 189–202, 2005.
- 20 Young, M. B., Gonneea, M. E., Fong, D. A., Moore, W. S., Herrera-Silveira, J., and Paytan, A.:
- Characterizing sources of groundwater to a tropical coastal lagoon in a karstic area using
 radium isotopes and water chemistry, Mar. Chem., 109, 377–394, 2008.
- 23

- 26 -

Table 1: Experimental conditions and sampling time intervals for methane headspace concentration analyses of sediment slurry incubations.

1 Table	1: Experimental co	onditions and sampling	time intervals for	methane headspace	e
2 concer	ntration analyses of	of sediment slurry incub	ations.	-	
3	-				
	Treatment	Initial concentration of treatment	Experiment length (days)	Number of measurements	Methane production rate (nmol CH ₄ cm ⁻³ slurry d ⁻¹)
Controls	No amendment (anaerobic)	N ₂ headspace	29	3	1.3×10^{-4} to 2.0×10^{-3}
	Autoclaved	N ₂ headspace	29	3	0 to 2.6×10^{-3}
	Aerobic- O ₂ gas	$16\% O_2$ headspace (0.36 mM)	29	3	5.7×10^{-4} to 3.5×10^{-3}
	BES	40 mM	29	3	$0 \text{ to } 1.3 \times 10^{-4}$
Competitive		100% headspace (1.8	••	2	5 41/ 10-3
substrates	H_2 gas	mM)	29	3	5.4×10^{-5} to 6.2
	Acetate	10 mM	29	3	6.8×10^{-4} to 9.2×10^{-2}
	Formate	10 mM	29	3	6.9×10^{-4} to 1.6×10^{-1}
Noncompetitive					
substrates	Methanol	10 mM	29	4	2.0×10^{-2} to 19
	TMA	10 mM	29	4	5.4×10^{-4} to 40

- 27 -

- 1 Table 2. Model-derived depth-integrated turnover rates (mmol $m^{-2} d^{-1}$), dissolved methane fluxes to the water column (mmol $m^{-2} d^{-1}$) and contributions of
- 2 methanogenesis to net methane production (%) and heterotrophic sulfate reduction to POC degradation (%). CEL and CH represent cores collected from

3 Celestún Lagoon and Chelem Lagoon.

	Length of model	$R_{aa} = R_{aa}$	<i>R</i>	R	<i>R</i>	F	F	$R_{\rm ex}/(R_{\rm ex}+R_{\rm exc})$	$2R_{m}/R_{max}$	R ₅₀ ²⁻	Reu
Group 1	column (cm)	$K_{SD} - K_{SR}$	κ_M	RPOC	<i>𝔅𝔅𝔅𝔅𝔅𝔅𝔅𝔅𝔅𝔅𝔅𝔅𝔅𝔅𝔅𝔅𝔅𝔅𝔅</i>	T methane (top)	T methane (bottom)	$K_{M'}(K_{M} + K_{MB})$	2KSR/KPOC	504	MCH ₄
1CEL Apr00	20	37	0.13	77	0.41	0.59	0.06	25%	07%		
1CEL_Aproo	20	2.7	1.5	7.4	0.41	0.57	0.60	100%	50%		
1CEL_Det01	20	6.2	0.12	12	0 32	0.94	-0.00	279/	0.80/		
1CEL_0001	20	3.6	8.0	23	0.32	6.0	-0.04	2770	310/		
2CEL_Juio2	20	1.1	0.05	23	0.76	0.54	-1.98	5 8%	96%		
2CEL_Decou	20	1.1	0.05	2.5	0.70	0.34	-0.27	5.870	90%		
2CEL_Jui02	20	11	0.08	22	0.05	0.11	-0.02	63%	99%		
SCEL_API00	20	1.5	0.29	5.2	0.24	0.08	0.13	33%	8270		
3CEL_JUI02	20	7.1	0.25	15	2.2	3.0	0.63	10%	9/%		
I_ICH_OctoI	13.75	24	31	110	0	11	-19	100%	44%		
I_2CH_Oct01	20	3.0	26	58	0	20	-5.6	100%	10%		
Group-2											
1CH_Dec00	20					0.52	-7.2			4.5	7.8
1CH_Apr00	20					~0	~0			-3.2	~0
2CH_Dec00	20					~0	~0			6.9	~0
5CH_Apr00	20					0.012	~0			21	0.013
2CEL_Oct01	20					11	-0.01			3.9	11
14CEL_Jul02	20					0.27	~0			3.7	0.27
16CEL_Dec00	20					-0.047	0.013			-1.8	-0.060
Group-3											
5CEL_Apr00	10					0.014	-0.01			-69	0.028
14CEL Dec00	20					3.4	-0.13			10	3.6
14CEL Oct01	20					0.088	-0.01			2.9	0.10
Group-4											
16CEL Jul02	20					0.096	0.02			6.1	0.072
16CEL Oct01	20					~0	~0			0.83	~0
7CH Oct01	20					0.13	~0			2.6	0.14
8CH Dec00	20					~0	~0			0.85	0.012

 $\frac{8 \text{ CH} \text{ Decould}}{R_{SD}} = \frac{20}{2} - \frac{20}{2}$

- 28 -

- Table 3: Maximum model-derived rates of methanogenesis and sulfate reduction for cores in
- Group-1 and maximum model-derived rates of methane production and sulfate consumption for cores in Group-2, Group-3 and Group-4. CEL and CH represent cores collected from

2 3 4 Celestún Lagoon and Chelem Lagoon.

1

	$Max-R_M$ (nmol CH ₄ cm ⁻³ d ⁻¹)	$\frac{\text{Max}-R_{SR}}{(\text{nmol SO}_4^{-2} \text{ cm}^{-3} \text{ d}^{-1})}$
Group-1		
1CEL_Apr00	9.0	304
1CEL_Dec00	116	559
1CEL_Oct01	7.1	740
1CEL_Jul02	564	1425
2CEL_Dec00	4.9	587
2CEL_Jul02	7.4	1323
3CEL_Apr00	20	405
3CEL_Jul02	26	1227
1_1CH_Oct01	2199	1802
1_2CH_Oct01	1959	1476
Group-2		
1CH_Dec00	2531	407
1CH_Apr00	1.6	2687
2CH_Dec00	1.1	2835
5CH_Apr00	2.1	8378
2CEL_Oct01	504	715
14CEL_Jul02	19	394
16CEL_Dec00	2.7	330
Group-4		
5CEL_Apr00	63	5212
14CEL_Dec00	1517	1756
14CEL_Oct01	23	1007
Group-5		
16CEL_Jul02	10	186
16CEL_Oct01	0.08	599
7CH_Oct01	4.1	940
8CH_Dec00	0.57	230

Table A1: Rate expressions applied in the differential equations $(R_C \text{ in Eq. } (2))$

Variable	Rates	Applied cores
SO_4^{2-}	$-R_{SR}$	Group-1
CH_4	$+R_M + R_{MB}$	Group-1
$SO_4^{2-}_{dep}$	$+R_{SD}$	Group-1
SO_4^{2-}	$+R_{SO_{4}^{2-}}$	Group-2, Group-3 and Group-4
CH ₄	$+R_{CH_4}$	Group-2, Group-3 and Group-4

- 30 -

rubie 112. Boundary conditions used in the model.
ruble 112. Doundary conditions used in the model.

	SO_4^{2-}	CH_4	SO4 ²⁻ dep	SO4 ²⁻	CH ₄	SO4 ²⁻ dep	Un
	(top)	(top)	(top)	(bottom)	(bottom)	(bottom)	0.1
Group-1	-	0	4.0	0.5	0.5	5 53 4	
1CEL_Apr00	5	0	4.8	8.5	0.5	5.534	mN
1CEL_Dec00	15	0.16	-2.2	5	0.56	2	mN
1CEL_Oct01	15	0	-2.3	7.5	0.295	4.6	mN
1CEL_Jul02	15	0.1	2.5	/.8	0.35	5.368	mN
2CEL_Jul02	18	0.02	10	18.5	0.035	-2	mN
3CEL_Apr00	6.5	0.25	6.7	3.5	0.825	5.766	mN
3CEL_Jul02	13.8	0.31	12.4	0.5	1.3	3.5	mN
I_ICH_Oct01	15.1	0	12.4	13.2	0.0295	12.03	mN
1_2CH_Oct01	12	0.01	16	10	1	14.641	m
2CEL_Dec00	21	0.01	-0.4431	/.0	0.25	4.0	mix
Group-2	11.5	0.102		92	0.522		m۱
1CH_Decou	32	0.005		12.5	0.006		mN
2CH Dec00	19.9	0.0015		7 96	0.0019		mN
5CH_Apr00	31.7	0.0031		29.1	0.0145		mN
2CEL Oct01	5.0	0.511		7.88	0.734		mN
14CEL Jul02	18.3	0.085		31.5	0.02		mN
16CEL Dec00	8.8	0.038		8.81	0.025		mN
Group-3							
5CEL_Apr00	17	0.047		11.6	0.0275		mN
14CEL_Dec00	20.5	2.1		34.9	0		mN
14CEL_Oct01	20	0.01		33	0.012		mN
Group-4							
16CEL_Jul02	21	0		25.65	0.070		mN
16CEL_Oct01	23	0.00139		25.8	0.0015		mN
7CH_Oct01	20.5	0.00477		19.1	0.01		mN
8CH Dec00	18.6	0		19	0.013		mN

- 31 -

1	Table A2. In	magad and	boot fit	model	noromotors	in aaah	ooro
1	Table AS. III	iposeu anu	UCSI-III	mouer	parameters	in caci	COLE

	T	S	Р	D _{m(SO4²⁻)}	D _{m(CH₄)}	D _{m(SO4} ²⁻ dep)	L_{MB}	k _{MB}	k _{SD}	k _{CH4}	k _{S04} 2-
	(°C)	(-)	(bar)	(cm ² yr ⁻¹)	(cm ² yr ⁻¹)	(cm ² yr ⁻¹)	(mM)	(yr ⁻¹)	(yr ⁻¹)	(yr ⁻¹)	(yr ⁻¹)
Group-1											
1CEL_Apr00	27.3	17.6	1.06	354	598	354	1.2	1	500		
1CEL_Dec00	22.2	16.4	1.06	367	523	367	1.3	0	400		
1CEL_Oct01	31.2	13.9	1.1	382	659	382	1.4	0.6	500		
1CEL_Jul02	30	21.1	1.01	374	640	374	1.1	0	500		
2CEL_Dec00	22	17.7	1.06	315	520	315	1.3	1.6	500		
2CEL_Jul02	28.7	20.8	1.01	364	619	364	1.1	0.1	500		
3CEL_Apr00	28.6	20.2	1.07	363	618	363	1.2	0.9	400		
3CEL_Jul02	30.4	18.2	1.01	377	646	377	1.1	50	500		
1_1CH_Oct01	29.8	32.1	1.01	372	636	372	1.1	0	500		
1_2CH_Oct01	29.8	32.1	1.01	372	636	372	1.1	0	500		
Group-2											
1CH_Dec00	25.2	24.8	1.05	318	556					1000	300
1CH_Apr00	26.3	39.4	1.09	347	583					1000	500
2CH_Dec00	23.9	27.5	1.08	329	547					4000	2000
5CH_Apr00	29.6	38	1.04	382	659					1000	1000
2CEL_Oct01	31.2	14.3	1.1	382	659					1000	500
14CEL_Jul02	31.5	27.4	1.01	385	663					1000	300
16CEL_Dec00	22.6	31.2	1.02	319	529					1000	300
Group-3											
5CEL_Apr00	26.5	21.1	1.06	348	586					1100	3000
14CEL_Dec00	23.5	31.1	1.06	326	541					1000	300
14CEL_Oct01	31.1	13.9	1.07	382	657					1000	500
Group-4											
16CEL_Jul02	30.3	30.5	1.01	376	644					600	300
16CEL_Oct01	29.7	28.2	1.06	319	529					1000	300
7CH_Oct01	29.6	31.3	1.01	371	633					500	300
8CH_Dec00	24.4	31.3	1.05	333	555					500	300

- 32 -



1

2 Figure 1: Maps of (A) the Yucatán Peninsula with lagoon locations, (B) Celestún Lagoon and

- 33 -

3 (C) Chelem Lagoon showing the sampling stations (circles) of sediment cores.



2 Figure 2: Depth profiles of modeled (lines), measured (circles) and calculated (triangles) 3 concentration of dissolved methane (dashed lines; open circles), sulfate (solid lines; solid 4 circles) in the upper panel and sulfate depletion (solid lines; solid triangles), zero sulfate 5 depletion (dashed lines) and chloride concentration (gray circles) in the lower panel for each 6 profile type (Groups 1-4, see text). One selected profile per group is shown here for illustration 7 and the other profiles for each group (9 cores for Group-1, 6 cores for Group-2, 2 cores for Group-3 and 3 cores for Group-4) are presented in the Appendix (Fig. A1). CEL and CH 8 9 represent cores collected from Celestún Lagoon and Chelem Lagoon.

10 11





4 Figure 3: Relationship between (A) $[Cl^-]$ and $[SO_4^{2-}]$, (B) $[Cl^-]$ and $[CH_4]$ and (C) $[CH_4]$ and

5 $[SO_4^{2^-}]$ in porewater samples.

- 35 -



2

1

Figure 4: (A) Headspace methane concentrations in sediment slurry incubations. (B) Expansion
of (A), showing results for acetate, formate, and controls. (C) Expansion of (A), showing
results for controls only. Error bars represent one standard deviation for triplicate sample
bottles.

- 36 -



2 Figure 5: Model sensitivity analysis of methane concentrations for cores in Group-1 to 3 the different processes controlling methane concentrations in porewaters. Black dashed 4 lines denote the standard simulation results: CH₄ production rate = $+ R_{MB} + R_M$. R_M is

4 lines denote the standard simulation results: CH₄ production rate = $+ R_{MB} + R_M$. R_M is 5 methanogenesis, R_{MB} is methane bubble dissolution, R_{AOM} is anaerobic oxidation of

- 37 -

6 methane and R_{SD} is net sulfate depletion.

7

Group-1



6 cores collected from Celestún Lagoon and Chelem Lagoon.



Figure A1: Continued.

- 39 -

Group-3





Group-4



- 41 -