

Supplement of Biogeosciences Discuss., 12, 18315–18344, 2015
<http://www.biogeosciences-discuss.net/12/18315/2015/>
doi:10.5194/bgd-12-18315-2015-supplement
© Author(s) 2015. CC Attribution 3.0 License.



Supplement of

Dead zone or oasis in the open ocean? Zooplankton distribution and migration in low-oxygen medowater eddies

H. Hauss et al.

Correspondence to: H. Hauss (hhauss@geomar.de)

The copyright of individual parts of the supplement might differ from the CC-BY 3.0 licence.

Supplementary information




Figure S1: Daily cycles of moored ADCP target strength (dB) during transit of ACME in 2010 (data from Jan 1 to Mar 14). Data are averaged for days with a mean O₂ concentration (at the depth of the moored O₂ sensor, approximately 50 m) below 20 μmol kg⁻¹ (row A, n=5), 20-50 μmol kg⁻¹ (row B, n=7), and 100-250 μmol kg⁻¹ (row C, n=54). Left: Depth-resolved target strength contour plot; right: target strength at the respective depth levels indicated in the left plots

Table S1. Multinet-based abundance (ind m⁻³) for the five stations sampled during M105.

Day/Night	Station	Depth range	copepods										other crustaceans					gelatinous							other			protists			detritus			
			cop_calanoid_other	cop_eucalanid	cop_nauplii	cop_oithonid	cop_cyclopoid_harpacticoid_other	cop_macrosetella	cop_oncaeaid	crust_euphausiid	crust_amphipod	crust_ostracod	crust_decapod	gel_filt_salp_like	gel_cam_ctenophore	gel_cam_medusa	gel_cam_chaetognath	gel_cam_siphonophore	gel_filt_app	gel_other	moll	fish	egg	polychaeta	pro_sarco_radiolaria_like	pro_sarco_foram_like	det_feces_like	det_darkparticle	det_aggregate	det_fiber				
Day	Eddy Core	0-85	557.2	5.1	1.7	58.0	6.7	2.5	82.7	1.8	0.6	5.7	0.7	3.3	0.0	0.0	0.0	0.0	0.0	32.1	1.1	1.4	0.8	2.2	8.7	0.14	37.2	1.2	3.9	45.0	0.0	10.6	306.0	23.9
	Eddy Core	85-120	156.4	24.5	3.0	12.6	7.2	16.5	107.8	0.4	0.4	54.7	2.7	0.0	0.0	0.0	6.1	1.9	0.0	2.2	8.7	0.33	29.4	2.5	0.8	13.0	0.0	8.3	271.6	20.1				
Day	Eddy Core	120-200	49.3	13.5	1.0	4.7	6.3	1.0	16.0	0.1	0.1	11.5	0.1	0.1	0.0	3.7	0.0	0.0	0.6	8.9	0.04	11.8	0.4	0.2	6.6	0.0	3.5	84.0	5.9					
	Eddy Core	200-300	43.4	4.0	1.6	4.9	0.5	0.6	28.6	0.1	0.0	6.8	0.0	0.0	0.0	0.7	0.5	0.0	0.5	2.2	0.00	8.2	0.2	0.2	4.8	0.0	4.6	52.0	12.1					
Night	Eddy Core	0-100	512.5	25.5	3.0	46.6	10.8	0.9	74.8	3.4	1.1	6.9	0.5	3.4	0.0	31.4	3.4	2.3	0.9	19.0	0.23	38.3	1.6	2.8	30.8	0.0	6.7	129.2	59.7					
	Eddy Core	100-200	33.1	3.0	1.0	8.5	1.2	0.4	25.4	0.5	0.3	7.0	0.1	0.0	0.0	2.1	0.0	0.0	0.0	0.1	0.00	0.1	0.00	0.3	0.8	1.1	0.0	0.8	17.0	4.0				
Day	Eddy Core	200-300	15.5	4.7	0.6	1.7	0.6	0.7	7.5	0.3	0.0	3.3	0.1	0.0	0.0	1.3	0.1	0.0	0.0	0.3	0.00	3.6	0.2	0.0	0.9	0.0	1.1	8.7	2.2					
	Eddy Core	300-600	22.4	3.9	1.8	2.7	0.2	0.6	16.4	0.0	0.0	1.8	0.0	0.0	0.0	1.2	0.0	0.0	0.1	0.0	0.02	4.8	0.3	0.0	0.6	0.0	0.8	17.6	2.6					
Night	Eddy Core	600-800	9.6	3.2	0.9	0.5	0.5	0.0	10.9	0.0	0.0	1.0	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.1	0.00	3.2	0.6	0.0	0.1	0.0	0.3	8.6	1.8					
	Outside Eddy	600-800	9.6	3.2	0.9	0.5	0.5	0.0	10.9	0.0	0.0	1.0	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.1	0.00	3.2	0.6	0.0	0.1	0.0	0.3	8.6	1.8					

Table S2. Multinet-based integrated abundance (ind m⁻², upper 600 m) for the five stations sampled during M105.

Day/Night	Station	Depth range	copepods					other crustaceans				gelatinous					other		protists		detritus								
Day	Eddy Core 0-600m	67382	3180	767	6771	1433	976	19940	194	76	4693	158	335	0	0	3393	210	118	290	3369	41	7204	280	408	5822	0	2402	50884	4916
Night	Eddy Core 0-600m	89537	6940	553	8521	1024	1427	19457	162	9	4563	115	0	94	0	5768	482	752	453	10954	14	8075	2374	890	12614	597	8279	71515	8851
Day	Eddy Mar 0-600m	33563	2986	1377	2339	401	220	13754	178	52	3530	256	109	3	18	2421	581	4740	518	680	38	4857	230	29	1240	94	987	25923	3350
Day	Outside Ek 0-600m	48624	3876	773	4599	824	207	14823	90	102	2611	220	235	0	0	2669	148	1822	271	975	38	4086	396	1928	1431	47	1507	34073	1738
Night	Outside Ek 0-600m	62826	4496	1007	6478	1320	381	15707	431	141	2261	83	344	0	0	3827	353	232	117	1957	29	6169	298	357	3445	0	1098	20784	7360