

This discussion paper is/has been under review for the journal Biogeosciences (BG).  
Please refer to the corresponding final paper in BG if available.

# Changing seasonality of the Baltic Sea

M. Kahru<sup>1</sup>, R. Elmgren<sup>2</sup>, and O. P. Savchuk<sup>3</sup>

<sup>1</sup>Scripps Institution of Oceanography, University of California San Diego, La Jolla, California

<sup>2</sup>Department of Ecology, Environment and Plant Sciences, Stockholm University, Stockholm, Sweden

<sup>3</sup>Baltic Nest Institute, Stockholm University, Stockholm, Sweden

Received: 30 October 2015 – Accepted: 9 November 2015 – Published: 25 November 2015

Correspondence to: M. Kahru (mkahru@ucsd.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

BGD

12, 18855–18882, 2015

## Changing seasonality of the Baltic Sea

M. Kahru et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Abstract

Changes in the phenology of physical and ecological variables associated with climate change are likely to have significant effect on many aspects of the Baltic ecosystems. We apply a set of phenological indicators to multiple environmental variables measured by satellite sensors for 17–35 years to detect possible changes in the seasonality in the Baltic Sea environment. We detect significant temporal changes such as earlier start of the summer season and prolongation of the productive season in multiple variables ranging from basic physical drivers to ecological status indicators. While increasing trends in the absolute values of variables like sea-surface temperature (SST), diffuse attenuation of light (Ked490) and satellite-detected chlorophyll concentration (CHL) are detectable, the corresponding changes in their seasonal cycles are more dramatic. For example, the cumulative sum of  $30\,000\text{ W m}^{-2}$  of surface incoming shortwave irradiance (SIS) was reached 23 days earlier in 2014 compared to the beginning of the time series in 1983. The period of the year with SST of at least  $17^\circ\text{C}$  has almost doubled (from 29 days in 1982 to 56 days in 2014), the period with Ked490 over  $0.4\text{ m}^{-1}$  has increased from about 60 days in 1998 to 240 days in 2013, i.e. quadrupled. The period with satellite-detected CHL of at least  $3\text{ mg m}^{-3}$  has doubled from approximately 110 days in 1998 to 220 days in 2013. While the timing of both the phytoplankton spring and summer blooms have advanced, the annual CHL maximum that in the 1980s corresponded to the spring diatom bloom in May has now switched to the summer cyanobacteria bloom in July.

## 1 Introduction

Estuarine areas world-wide are experiencing rapid changes due to anthropogenic pressures of both local and global nature (Cloern et al., 2015). Ecosystems in the brackish Baltic Sea have been under anthropogenic stress for many decades (Elmgren, 1989, 2001). Both direct effects, such as input of nutrients and various pollutants, and indirect

BGD

12, 18855–18882, 2015

## Changing seasonality of the Baltic Sea

M. Kahru et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



effects through climate change are important. Time series of environmental variables show variability at multiple scales but separating the effects of natural climate variability from the effects of anthropogenic climate change is difficult. We apply a set of phenological indicators to multiple environmental variables, mostly from satellite sensors, to detect changes in the environment. It appears that phenological indicators are very sensitive in detecting environmental change. We show significant changes in the seasonality of both the physical drivers and in ecological indicators of the Baltic Sea. Increasing trends are detectable in the absolute values of sea-surface temperature and water turbidity, but changes in their seasonal cycles are more clear-cut. The seasonality of a number of abiotic and biotic variables for which data are available has changed drastically during the last few decades. These changes are likely to have major effects on many aspects of Baltic ecosystems.

## 2 Methods and datasets

*Datasets.* The datasets used in this analysis and their sources are shown in Table 1. Surface incoming shortwave irradiance (SIS) data were produced from geostationary Meteosat satellites (Mueller et al., 2009; Müller et al., 2015) using a climate version of the Heliosat algorithm (Cano et al., 1986; Beyer et al., 1996) and obtained from the Satellite Application Facility on Climate Monitoring (CM SAF, [http://www.cmsaf.eu/EN/Products/AvailableProducts/Dataset/Dataset\\_node.html](http://www.cmsaf.eu/EN/Products/AvailableProducts/Dataset/Dataset_node.html)).

Sea-surface temperature (SST) data produced by NOAA and NASA were obtained from <http://www.nodc.noaa.gov/SatelliteData/pathfinder4km> (Casey et al., 2010).

Near-surface wind data were assembled from two sources. Data for 1987–2011 are version 3.5a from the Cross-Calibrated Multi-Platform (CCMP) Ocean Surface Wind Components (Atlas et al., 2008), available from <ftp://podaac-ftp.jpl.nasa.gov/allData/ccmp/L3.5a/>. Data for 2013–2015 are from ASCAT, produced by the KNMI Scatterometer Team and available from the EUMETSAT Ocean and Sea Ice SAF (<http://www.osi-saf.org/>).

## Changing seasonality of the Baltic Sea

M. Kahru et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Changing seasonality of the Baltic Sea

M. Kahru et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Satellite-detected coefficient of attenuation of diffuse downwelling light at 490 nm (Ked490) and near-surface chlorophyll concentration (CHL) are produced by the ESA Ocean Colour Climate Change Initiative project (Lavender et al., 2015) using satellite data archives of NASA's SeaWiFS and MODIS-Aqua sensors and ESA's MERIS sensor. Version 2.0 datasets were downloaded from ftp://oc-cci-data:ELaiWai8ae@oceancolour.org/. The Ked490 algorithm uses the Lee et al. (2005) semianalytic method. The CHL algorithm is based on an empirical ratio of remote sensing reflectance (O'Reilly et al., 1998).

Fraction of Cyanobacteria Accumulations (FCA, Kahru et al., 2007; Kahru and Elmgren, 2014) is a form of presenting the frequency of cyanobacteria accumulations that is normalized to the number of unobstructed satellite views of the sea surface. This normalization is needed as clouds often cover the sea surface and make it impossible to detect accumulations.

*Phenological and cumulative indicators.* For the analysis of changes in the annual cycle we use the following phenological indicators (Table 2):

1. Day of year when a threshold value is first reached (DF="day first"),
2. Day of year when a threshold value is last reached (DL="day last"),
3. Duration between the first and last reaching of a threshold value (DD="day duration"),
4. Day of year when the annual maximum occurs (DM="day maximum"),
5. Count of days above a threshold value (DC="day count").

These indicators can be applied to different satellite-derived or in situ variables. Satellite versions of these indicators were spatially averaged over the area of interest, e.g. the whole Baltic Sea or parts of it. The nomenclature of indicators uses two characters showing the type of indicator (e.g. DF, DL, etc.), followed by the variable type (e.g. SST for sea-surface temperature), and followed by the threshold value (e.g. 16 for 16 °C of



in timing, no significant changes in the absolute daily values are evident in the time series.

### 3.2 Sea-surface temperature

As global time series of sea-surface temperature (SST) are available for several decades (Casey et al., 2010), SST is a good variable for phenological analysis. While SST itself is an important driver that directly affects the rates of biological processes, its indirect effects are probably more important, through its influence on the near-surface stratification that suppresses vertical mixing in the water column and affects many ecological processes, such as the flux of nutrients to the surface layer or the accumulation of cyanobacteria near the water surface. As cyanobacteria growth in temperate environments is enhanced by higher temperatures and by near-surface stratification, SST is an important environmental driver for these blooms (Paerl and Huisman, 2008). Cyanobacteria growth typically accelerates when SST exceeds 12 °C and peaks above 17 °C (Hense et al., 2013). We therefore estimated the start of the cyanobacterial growth season as the first day when the spatially averaged SST reached 12 °C (DFSST12), the end of the growth season as the last day with SST of at least 12 °C (DLSST12) and the duration of the growth season as the number of days between those dates (DDSST12= DLSST12–DFSST12). Similarly, the “peak growth season” is parameterized by the respective indicators (DFSST17, DLSST17, DDSST17) for a SST of 17 °C. Significant ( $F$  test,  $p = 0.05$ ) trends towards an earlier start and a later end of the growth season were detected for all those timing indicators (Fig. 4).

It appears that while the springtime warming of the sea surface has become significantly earlier (i.e. the slope of DFSST vs. time is negative), particularly for colder temperatures, the time in summer when the higher temperatures are reached has not changed and the trend is not significant ( $F$  test,  $p = 0.05$ ) for any SST above 16 °C (Fig. 5a). In contrast, the slope for DLSST is positive, indicating delay of the fall cooling, with the highest slopes for SSTs of 15 to 17 °C (Fig. 5a). As a result of the earlier warming and later cooling, the period with SST above a certain threshold (DDSST) has

## Changing seasonality of the Baltic Sea

M. Kahru et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Changing seasonality of the Baltic Sea

M. Kahru et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



changed quite dramatically, particularly for SST values of  $\sim 15\text{--}16^\circ\text{C}$  (Fig. 5b). The rate of change for DDSST16 has been almost a day per year ( $0.98\text{ day year}^{-1}$ ). As a result, the duration of the period with a mean SST of at least  $16^\circ\text{C}$  has increased from about 40 days in 1982 to about 72 days in 2014, i.e. by 32 days (Fig. 5b). Similarly, the duration of the period with SST of at least  $17^\circ\text{C}$  (DDSST17) has increased from 29 days in 1982 to about 56 days in 2014, i.e. almost doubled. These are drastic changes in the phenology of an important driver of ecosystem processes. In contrast, the timing and the duration of the period with the highest SST ( $\sim 22^\circ\text{C}$ ) has not changed significantly. It is important to note that while a significant increase in the mean summertime SST is evident in the Baltic Sea (Belkin, 2009), this increase is rather small compared to the annual range of SST, from about  $0^\circ\text{C}$  in the winter to about  $22^\circ\text{C}$  in the summer. The phenology of the changes gives a more detailed and ecologically highly relevant picture that complements general increase in the mean SST.

### 3.3 Wind

Wind speed and direction are important variables that affect the biology of the Baltic surface waters. In a previous study (Kahru et al., 2007) we found a correlation between the location of the major cyanobacteria accumulations either in the southern or northern half of the Baltic and the strength of wind in the north-easterly direction. We therefore examined the mean annual cycle in the strength of winds in the north-eastern direction (NE-ward winds) and the possible changes to it. The annual cycle of the NE-ward winds has a minimum in April and a subsequent increase after that (Fig. 6a). We examined if the onset of the summer increase in NE-ward winds has changed. While the last 6–7 years (2008 to 2015) show a steady shift towards earlier start of the summer wind increase (Fig. 6b), the large interannual variability precludes establishing a long-term trend and the changes in the starting time of the summer increase in NE-ward winds are not significant.

### 3.4 Attenuation of light, *Ked490*

The coefficient of downwelling light at 490 nm (*Ked490*) is a commonly used indicator of water transparency (e.g., Kratzer et al., 2003; Lee et al., 2005) that can be estimated from a satellite sensor. It reflects both light absorption (e.g. by CDOM and phytoplankton pigments) and scattering of light by particles. The timing of the increase in *Ked490* has changed significantly during the time period with reliable satellite data (1997 to present). The summer period with high *Ked490* corresponding to low water transparency has become earlier and persisted longer into the fall (Fig. 7). Significantly, these changes have been more pronounced for the higher values of *Ked490* than for the lower values of *Ked490*. For example, using the regression lines in Fig. 7, we can estimate that from 1998 to 2013 the period with mean *Ked490* over  $0.1 \text{ m}^{-1}$  has become longer by 35 days whereas over the same period the duration of the mean *Ked490* over  $0.4 \text{ m}^{-1}$  has increased from about 60 days in 1998 to 240 days in 2013, i.e. has become longer by 180 days. However, the last 1–2 years seem to show a reversal of the trend.

### 3.5 Near-surface chlorophyll *a*

The satellite-derived surface concentration of chlorophyll *a* (CHL) is an important measure of the amount of phytoplankton, but is known to have large errors in the Baltic Sea (Darecki and Stramski, 2004; Attila et al., 2013). This is due both to problems in the atmospheric correction procedures and in the separation of phytoplankton pigments from the colored dissolved organic matter (CDOM). The standard CHL algorithm essentially measures total absorption of blue light by a mixture of phytoplankton pigments, non-algal particles and CDOM. The separation of these components is difficult in case of large and variable concentrations of CDOM that are typical of the Baltic Sea. While the satellite-derived CHL overestimates true near-surface CHL in the Baltic Sea due to the high concentration of CDOM (Darecki and Stramski, 2004), the phenology of CHL is still a meaningful indicator of the timing of phytoplankton blooms, as the early

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



increase in turbidity is well correlated with phytoplankton spring bloom and the summer maximum in turbidity is correlated with the cyanobacteria blooms in the central Baltic Sea.

The time series of CHL in the central Baltic Sea (Fig. 8) shows the well-known annual cycle with maxima corresponding to the summer cyanobacteria and the spring diatom blooms. A statistically significant ( $F$  test,  $p = 0.05$ ) increasing trend can be detected in the 1998–2013 time series of mean CHL in central Baltic Sea (Fig. 8) with a slope of  $0.067 \text{ mgm}^{-3} \text{ year}^{-1}$ . However, it is not clear how much of the increase is truly due to an increase in phytoplankton vs. increases in CDOM and non-algal particles. As phytoplankton is expected to have a strong annual cycle in the open Baltic Sea, it is, however, likely that the distinct spring increase in CHL is due to the phytoplankton spring bloom. The timing of the annual CHL maximum (DMCHL) averaged over central Baltic (Fig. 1b) has advanced from 1998 to 2013 at a mean annual rate of  $-1.66 \text{ day year}^{-1}$ . It is remarkable that while in the 1980s (e.g. 1985–1989; Kahru et al., 1991) the spring bloom concentration of chlorophyll  $a$  was approximately double of that of the late summer cyanobacteria bloom (dashed line in Fig. 9a), currently CHL during the summer cyanobacteria bloom is significantly higher (solid line in Fig. 9a). A similar change in chlorophyll values around 1990, with a decrease in spring and an increase in summer, has been reported from the western Gulf of Finland (Raateoja et al., 2005). Another significant change is the earlier occurrence of the summer CHL maximum. While in 1985–1989 it occurred in late summer (August–September, year days 213–270), the maximum now is on 7 July (year day 188). The current annual maximum now occurs at the time that used to be summer minimum in 1985–1989.

Both an earlier start and a later cessation of high CHL values are detected (Fig. 9b). The duration of the annual time period with CHL at least  $3 \text{ mgm}^{-3}$  (DDCHL3) has increased from approximately 110 days in 1998 to approximately 220 days in 2013, i.e. doubled. This increase in the duration of CHL and the associated high absorption of light by various substances is likely to have important ecological consequences. As with Ked490, the last 1–2 years deviate from the general trend.

## Changing seasonality of the Baltic Sea

M. Kahru et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



### 3.6 Frequency of cyanobacteria surface accumulations

Cyanobacteria are a major component of the Baltic phytoplankton community with some unique features. Their ability to fix nitrogen makes them important in driving the nitrogen cycle and stimulating summer primary production (e.g. Larsson et al., 2001; Karlson et al., 2015). The propensity of the co-dominant genus *Nodularia* to form dense surface accumulations makes it feasible to map its distribution using satellite sensors, including some not specifically designed for ocean color applications (Kahru and Elmgren, 2014). This makes it possible to create a time-series that is much longer than time series for ocean color variables like CHL. A 35 year time-series of the frequency of cyanobacteria surface accumulations (FCA) shows a trend towards earlier occurrence as measured by the center of timing of the accumulations (Fig. 10). The estimates of the start and end of the accumulation period are less reliable when based on the older, less sensitive AVHRR sensor and are therefore given only for the second half of the study period, based on the more sensitive and reliable ocean color sensor data. The start and end of the accumulation period seems closely related to the center of timing. Years when the accumulations are late (e.g. 2004), the total period of accumulations also tends to be shorter.

## 4 Discussion

We have applied a uniform set of phenological indicators to a number of variables ranging from physical drivers of the environment, such as incoming shortwave energy, sea-surface temperature and winds, to ecological (Ked490, CHL) and biological (cyanobacteria) components of the Baltic Sea ecosystem. Satellite-derived variables have the advantage of providing extended areal coverage instead of a point sample, with a regular and frequent sampling that is required for estimating phenological indicators. While satellite measurements using the visible and infrared spectrum are limited

BGD

12, 18855–18882, 2015

## Changing seasonality of the Baltic Sea

M. Kahru et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



by cloudy periods, the effective sampling frequency is still much higher than with ship-board monitoring.

While significant trends can be detected in the absolute values of a number of variables, e.g. the increasing trends of SST and CHL, the phenological indicators often show more marked change and clearer trends, even when trends in the mean values are questionable or nonsignificant. For example, the time series of incoming shortwave irradiance (SIS,  $W m^{-2}$ ) are extremely regular with no apparent trends in the absolute daily values but the timing of SIS shows subtle but significant changes in seasonality, with slightly less energy input in the winter and more during spring and summer. As a result of this change in timing, the cumulative sum of  $30\,000 W m^{-2}$  was reached about 23 days earlier in 2013 than at the beginning of the time series in 1983. Over the same period, the center of timing of the summer cyanobacterial bloom (Fig. 10) has become about 17 days earlier, a similarity that is likely to be more than a coincidence. A probable explanation for the delayed cumulative energy input in winter is increased cloudiness.

The mean annual SST has increased during the last decades, in agreement with the overall warming trend. However, changes in the timing of the annual SST cycle are more informative. In contrast to the spring and early summer warming of the sea surface that has advanced significantly in time, the timing of the summer temperature maximum has not changed. As a result of the earlier warming and later cooling, the period with SST above  $16^{\circ}C$  has increased at a mean rate of  $0.98 day year^{-1}$ . In contrast, the timing and the duration of the period with the highest SST ( $\sim 22^{\circ}C$ ) has not changed.

Dramatic changes have occurred recently in the annual cycle of the coefficient of light attenuation (Ked490), an indicator of water transparency. The duration of the period with elevated but intermediate Ked490 has somewhat increased but the duration of the period with high mean Ked490 (over  $0.4 m^{-1}$ ) has lengthened greatly, i.e. by approximately 180 days from 1998 to 2013. Similarly, the duration of the period with high CHL (operationally defined when CHL is at least  $3 mg m^{-3}$ ) has increased as the spring

**BGD**

12, 18855–18882, 2015

## Changing seasonality of the Baltic Sea

M. Kahru et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Changing seasonality of the Baltic Sea

M. Kahru et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



increase has advanced at a mean rate of  $-1.6 \text{ day year}^{-1}$  and the cessation of the high CHL period in the fall bloom has been delayed at the mean rate of  $1.9 \text{ day year}^{-1}$ . These changes in the timing of high Ked490 and CHL are clearly correlated with the changes towards earlier warming and later cooling of SST, but the rate of change is faster. Since the satellite-derived time series of Ked490 and CHL are shorter and less reliable than the time series of SST, their estimated mean rates of change are also less accurate. However, it seems that the response of the biological system to climate change and eutrophication has been amplified compared to the changes in the timing of the physical environment.

The changes in the timing of Ked490 and CHL are related and reflect the increased turbidity and decreased water transparency in the Baltic Sea. An obvious consequence of the increased Ked490 is that less light reaches depths below the surface. While this analysis was done for the central Baltic with small areas of benthic photosynthesis, we can assume that benthic communities in the coastal areas must also be experiencing significantly reduction in light due to the decreased water transparency. We can hypothesize that the increased turbidity and decreased water transparency are related to increased phytoplankton concentrations and increased bacterial production. Likely effects on the rest of the ecosystem including commercially important fisheries should be further evaluated.

The phenological indicators that we applied to a number of satellite-detected variables show significant climate-related changes in the Baltic Sea ecosystems. It appears that the biological response to climate warming is amplified, compared to the rate of change in the physical forcing (SIS and SST). As satellite-derived CHL algorithms in the Baltic Sea are inaccurate and noisy (e.g. Darecki and Stramski, 2004) – primarily due to the confounding influence of high concentrations of CDOM – trends in CHL absolute values are difficult to interpret. For example, an apparently increasing trend in CHL may be influenced by increasing concentrations of colored dissolved organic matter (CDOM) or nonalgal particles. However, the start of the annual increase in CHL is much more likely to reflect the increase in phytoplankton due to the known

phytoplankton spring bloom and the dependence of photosynthesis on the annual cycle of light and stratification. Elevated concentrations of CDOM and nonalgal particles are perhaps more likely to extend the high CHL period in the fall.

Cyanobacterial blooms are a worldwide phenomenon associated with eutrophication of lakes, reservoirs and estuaries, toxic contamination of drinking water, and other undesirable effects. They are a major environmental problem in the North American Great Lakes (Stumpf et al., 2012), in lakes in China (Paerl et al., 2011) and in the Baltic Sea (Funkey et al., 2014; Kahru and Elmgren, 2014). As cyanobacterial growth in temperate zones is enhanced by higher temperatures and near-surface stratification, these blooms are expected to become more frequent as a result of climate change (Pearl and Huisman, 2008, 2009; Wiedner et al., 2007). Cyanobacterial growth typically accelerates above 12 °C and peaks above 17 °C (e.g. Hense et al., 2013). We can therefore assume that the earlier start of the warm-up (measured as DFSST12) and later start of cool-down (measured as DLSST12) indicate improved conditions for cyanobacteria in the Baltic Sea. By this measure, the duration of both the growth season and the “enhanced growth season” (DDSST17) have become significantly longer. The duration of the period with a mean SST of at least 16 °C has increased, approximately, from 40 days in 1982 to 72 days in 2014, i.e. by 32 days (Fig. 5b), and the period with SST of at least 17 °C (DDSST17) from 29 days in 1982 to 56 days in 2014, i.e. almost doubled. Indeed, in accordance with these trends in SST, we see indications that cyanobacteria are becoming more dominant in the phytoplankton community. While in the 1980s the annual CHL maximum corresponded to the diatom spring bloom (in May in the central Baltic), it has now switched to the cyanobacteria bloom in July which has also advanced from its earlier mean timing in August.

## 5 Conclusions

Phenological indicators are sensitive in detecting environmental changes that are often hardly detectable using absolute values of the respective environmental variables.

## Changing seasonality of the Baltic Sea

M. Kahru et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Moreover, these indicators, e.g. the duration of the growth season, may have special ecological significance. Using these phenological indicators we show significant and sometimes drastic changes in the seasonality of the Baltic Sea. These changes are evident in multiple of variables from purely physical to ecological to biological. For several ecologically important variables (Ked490, CHL) the length of the annual period of high values has increased by a factor of two or more during the last couple of decades. The analyses reported above are based on satellite data, meaning that phenological analyses of this type can be made for most areas of the globe, not only for data-rich areas like the Baltic Sea.

*Acknowledgements.* Financial support was provided by the Swedish Research Council Formas and Stockholm University's Baltic Ecosystem Adaptive Management Program. The authors thank Uwe Pfeifroth, Jörg Trentmann, and Christine Träger-Chatterjee for help in accessing SIS data; CM SAF, NASA Ocean Color Processing Group and ESA OC-CCI group for satellite data.

## References

- Atlas, R., Ardizzone, J., and Hoffman, R. N.: Application of satellite surface wind data to ocean wind analysis, Proc. SPIE, 7087, 70870, doi:10.1117/12.795371, 2008.
- Attila, J., Koponen, S., Kallio, K., Lindfors, A., Kaitala, S., and Ylöstalo, P.: MERIS Case II water processor comparison on coastal sites of the northern Baltic Sea, Remote Sens. Environ., 128, 138–149, 2013.
- Belkin, I. M.: Rapid warming of large marine ecosystems, Prog. Oceanogr., 81, 207–213, 2009.
- Beyer, H. G., Costanzo, C., and Heinemann, D.: Modifications of the Heliosat procedure for irradiance estimates from satellite images, Sol. Energy, 56, 207–212, 1996.
- Cano, D., Monget, J. M., Albuissou, M., Guillard, H., Regas, N., and Wald, L.: A method for the determination of the global solar-radiation from meteorological satellite data, Sol. Energy, 37, 31–39, 1986.
- Casey, K. S., Brandon, T. B., Cornillon, P., and Evans, R.: The past, present and future of the AVHRR Pathfinder SST program, in: Oceanography from Space: Revisited, edited by: Barale, V., Gower, J.F.R., Alberotanza, L., New York, 323–341, doi:10.1111/gcb.13059, Springer, 2010.

## Changing seasonality of the Baltic Sea

M. Kahru et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Changing seasonality of the Baltic Sea

M. Kahru et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Cloern, J. E., Abreu, P. C., Carstensen, J., Chauvaud, L., Elmgren, R., Grall, J., Greening, H., Johansson, J. O. R., Kahru, M., Sherwood, E. T., Xie, J., and Yin, K.: Human activities and climate variability drive fast-paced change across the world's estuarine-coastal ecosystems, *Glob. Change Biol.*, 1–17, doi:10.1111/gbc.13059, 2015.

Darecki, M. and Stramski, D.: An evaluation of MODIS and SeaWiFS bio-optical algorithms in the Baltic Sea, *Remote Sens. Environ.*, 89, 326–350, 2004.

Elmgren, R.: Man's impact on the ecosystem of the Baltic Sea: energy flows today and at the turn of the century, *Ambio*, 18, 326–332, 1989.

Elmgren, R.: Understanding human impact on the Baltic ecosystem: changing views in recent decades, *Ambio*, 30, 222–231, 2001.

Funkey, C. P., Conley, D. J., Reuss, N. S., Humborg, C., Jilbert, T., and Slomp, C. P.: Hypoxia sustains cyanobacteria blooms in the Baltic Sea, *Environ. Sci. Technol.*, 48, 2598–2602, 2014.

Hense, I., Markus Meier, H. E., and Sonntag, S.: Projected climate change impact on Baltic Sea cyanobacteria, *Clim. Chang.*, doi:10.1007/s10584-013-0702-y, 2013.

Kahru, M. and Elmgren, R.: Multidecadal time series of satellite-detected accumulations of cyanobacteria in the Baltic Sea, *Biogeosciences*, 11, 3619–3633, doi:10.5194/bg-11-3619-2014, 2014.

Kahru, M., Kaasik, E., and Leebein, A.: Annual cycle of particle size fractions and phytoplankton biomass in the northern Baltic proper, *Mar. Ecol.-Prog. Ser.*, 69, 117–124, 1991.

Kahru, M., Savchuk, O. P., and Elmgren, R.: Satellite measurements of cyanobacterial bloom frequency in the Baltic Sea: interannual and spatial variability, *Mar. Ecol.-Prog. Ser.*, 343, 15–23, 2007.

Kahru, M., Brotas, V., Manzano-Sarabia, M., and Mitchell, B. G.: Are phytoplankton blooms occurring earlier in the Arctic?, *Glob. Change Biol.*, 115, 1733–1739, doi:10.1111/j.1365-2486.2010.02312.x, 2010.

Karlson, A. M. L., Duberg, J., Motwani, N. H., Hogfors, H., Ploug, H., Svedén, J. B., Garbaras, A., Sundelin, B., Hajdu, S., Larsson, U., Elmgren, R., and Gorokhova, E.: Nitrogen fixation by cyanobacteria stimulates production in Baltic food-webs, *Ambio*, 44, 413–426, 2015.

Kratzer, S., Hakansson, B., and Sahlin, C.: Assessing Secchi and photic zone depth in the Baltic Sea from satellite data, *Ambio*, 32, 577–585, 2003.

## Changing seasonality of the Baltic Sea

M. Kahru et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Larsson, U., Hajdu, S., Walve, J., and Elmgren, R.: Baltic Sea nitrogen fixation estimated from the summer increase in upper mixed layer total nitrogen, *Limnol. Oceanogr.*, 46, 811–820, 2001.

Lavender, S., Jackson, T., and Sathyendranath, S.: The Ocean Colour Climate Change Initiative, *Ocean Chall.*, 21, 29–31, 2015.

Lee, Z., Du, K., and Arnone, R.: A model for the diffuse attenuation coefficient of downwelling irradiance, *J. Geophys. Res.*, 110, C02016, 1–10, 2005.

Mueller, R., Matsoukas, C., Gratzki, A., Behr, H., and Hollmann, R.: The CM-SAF operational scheme for the satellite based retrieval of solar surface irradiance - A LUT based eigenvector hybrid approach, *Remote Sens. Environ.*, 113, 1012–1024, 2009.

Müller, R., Pfeifroth, U., Träger-Chatterjee, C., Cremer, R., Trentmann, J., and Hollmann, R.: Surface Solar Radiation Data Set - Heliosat (SARAH) - 1 edn, Satellite Application Facility on Climate Monitoring, doi:10.5676/EUM\_SAF\_CM/SARAH/V001, 2015.

O'Reilly, J., Maritorena, S., Mitchell, B. G., Siegel, D., Carder, K. L., Garver, S., Kahru, M., and McClain, C.: Ocean color chlorophyll algorithms for SeaWiFS, *J. Geophys. Res.*, 103, 24937–24953, 1998.

Paerl, H. W. and Huisman, J.: Climate - Blooms like it hot, *Science*, 320, 5872, 57–58, 2008.

Paerl, H. W. and Huisman, J.: Climate change: a catalyst for global expansion of harmful cyanobacterial blooms, *Environ. Microbiol.*, 1, 27–37, 2009.

Paerl, H. W., Xu, H., McCarthy, M. J., Zhu, G., Quin, B., Yi, L., and Gardner, W. S.: Controlling harmful cyanobacterial blooms in a hyper-eutrophic lake (Lake Taihu, China): the need for a dual nutrient (N and P) management strategy, *Water Res.*, 45, 5, 1973–1983, 2011.

Raateoja, M., Seppälä, J., Kuosa, H., and Myrberg, K.: Recent changes in trophic state of the Baltic Sea along SW coast of Finland, *Ambio*, 34, 188–191, 2005.

Stumpf, R. P., Wynne, T. T., Baker, D. B., and Fahnenstiel, G. L.: Interannual variability of cyanobacterial blooms in Lake Erie, *PLoS One*, 8, e42444, doi:10.1371/journal.pone.0042444, 2012.

Wiedner, C., Rucker, J., Brüggemann, R., and Nixdorf, B. S.: Climate change affects timing and size of populations of an invasive cyanobacterium in temperate regions, *Oecologia*, 152, 473–484, 2007.

## Changing seasonality of the Baltic Sea

M. Kahru et al.

**Table 1.** Satellite-derived datasets used here to produce phenological indicators.

Variable	Acronym	Units	Sample sources
Surface Incoming Shortwave Irradiance	SIS	$\text{W m}^{-2}$	EUMETSAT, NOAA, NASA doi:10.5676/EUM SAF_CM/SARAH/V001
Sea-surface temperature	SST	$^{\circ}\text{C}$	NOAA, NASA, GHRSSST http://www.nodc.noaa.gov/SatelliteData/pathfinder4km/ http://ghrsst.jpl.nasa.gov
Near-surface wind	WSP, directional wind	$\text{m s}^{-1}$	NASA https://podaac.jpl.nasa.gov/ EUMETSAT Ocean and Sea Ice SAF, http://www.osi-saf.org/
Coefficient of diffuse light attenuation	KED	$\text{m}^{-1}$	ESA http://www.esa-oceancolour-cci.org/
Near-surface chlorophyll <i>a</i> concentration	CHL	$\text{mg m}^{-3}$	ESA http://www.esa-oceancolour-cci.org/
Fraction of Cyanobacteria Accumulations	FCA	unitless	Kahru and Elmgren (2014)

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Changing seasonality of the Baltic Sea

M. Kahru et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




**Table 2.** Summary of proposed climate indicators.

Explanation	Type	Example variable groups
First day of reaching a threshold	DF	DFSST, DFCHL, DFKED
Last day of reaching a threshold	DL	DLSST, DLCHL, DLKED
Duration of the period between DF and DL	DD	DFSST, DDCHL, DDKED
Count of days over the threshold	DC	DCSST, DCCHL, DCKED
Day of reaching the annual maximum	DM	DMSST, DMCHL, DMKED
First day of reaching a cumulative threshold	DFCUM	DFCUMSIS, DFCUMSST
Cumulative count of days above threshold	DCCUM	DCCUMSIS, DCCUMSST


## Changing seasonality of the Baltic Sea

M. Kahru et al.



**Figure 1.** Maps of the study areas. **(a)** Baltic Sea area excluding shallow coastal areas and the Bothnian Bay (Kahru et al., 2007). **(b)** Central Baltic Sea including the following sub-basins: Northern Baltic Proper, Western and Eastern Gotland Basin and the Southeastern Baltic Proper. The shaded areas are used to calculate averages for, respectively, Baltic and central Baltic Sea.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)



**Figure 2.** Time series of the daily surface incoming shortwave irradiance (SIS,  $\text{W m}^{-2}$ ) derived from geostationary Meteosat sensors, averaged over the shaded area of the Baltic Sea shown in Fig. 1a.

## BGD

12, 18855–18882, 2015

### Changing seasonality of the Baltic Sea

M. Kahru et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)


[Printer-friendly Version](#)

[Interactive Discussion](#)



Changing seasonality  
of the Baltic Sea

M. Kahru et al.



**Figure 3.** Changing seasonality in cumulative surface incoming shortwave irradiance (SIS,  $\text{W m}^{-2}$ ) averaged over the area of the Baltic Sea indicated in Fig. 1a. **(a)** DFCUMSIS, i.e. day of year when the annual sum of daily mean SIS reaches the following thresholds: 200, 500, 1000, 2000, 3000, 5000, 10 000, 20 000 and 30 000  $\text{W m}^{-2}$ . For each threshold the circles show the day of the year and the line shows the respective linear regression. **(b)** Slopes of linear regressions in panel **(a)**.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I ◀

▶ I

◀

▶

Back


Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Figure 4.** Changes in satellite-detected SST phenology in the Baltic Sea (area indicated in Fig. 1a). The symbols and regression lines are (left to right): first day when 12°C is reached (DFSST12, open circles), first day when 17°C is reached (DFSST17, filled circles), last day when 17°C is reached (DLSST17, filled squares), last day when 12°C is reached (DLSST12, open squares).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## Changing seasonality of the Baltic Sea

M. Kahru et al.



**Figure 5.** (a) Rate of change ( $\text{day year}^{-1}$ ) in the day of year when a certain SST level is first reached (DFSST, filled circles) and when a certain SST level is last reached in a season (DLSST, open squares) for the Baltic Sea (area indicated in Fig. 1a). (b) Increase in the duration of a period with SST above a certain level (DDSST) from 1982 to 2014.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Changing seasonality of the Baltic Sea

M. Kahru et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)


[Back](#)

[Close](#)


[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



**Figure 6.** (a) Annual cycle in the mean monthly wind speed in the NE-direction in the Baltic Sea (area indicated in Fig. 1a). (b) Day of the year when the 5-day mean wind in the NE direction exceeds 2 m s<sup>-1</sup> for the first time after the April minimum.



**Figure 7.** Temporal changes in the start and end of  $Ked_{490} = 0.1 \text{ m}^{-1}$  **(a)** and  $Ked_{490} = 0.4 \text{ m}^{-1}$  **(b)** for the central Baltic Sea (area indicated in Fig. 1b). The circles show the first day of the year (DFKED) and the squares show the last day of the year (DLKED) with the respective value of  $Ked_{490}$ . The regression lines have been estimated for the time period of 1998–2013.


[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


# BGD

12, 18855–18882, 2015

## Changing seasonality of the Baltic Sea

M. Kahru et al.




**Figure 8.** Time series of the 5-day mean CHL ( $\text{mg m}^{-3}$ ) in the central Baltic Sea (area indicated in Fig. 1b) derived from the ESA-CCI processing of SeaWiFS, MERIS and Aqua-MODIS satellite data (Lavender et al., 2015).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Changing seasonality  
of the Baltic Sea

M. Kahru et al.



**Figure 9.** (a) Mean annual cycle of CHL in central Baltic Sea (area indicated in Fig. 1b) for 1997–2013 (solid line) compared with CHL measured in situ in 1985–1989 (circles, dashed line, from Kahru et al., 1991). (b) Temporal changes in the start and end of the “high chlorophyll season” ( $\text{CHL} \geq 3 \text{ mg m}^{-3}$ ) in the Baltic Sea. The connected markers and the respective regression lines are (from left to right): day when  $\text{CHL} = 3.0 \text{ mg m}^{-3}$  is reached for the first time (DFCHL3, circles) and day when  $\text{CHL} = 3.0 \text{ mg m}^{-3}$  is reached for the last time (DLCHL3, squares) during the season.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## Changing seasonality of the Baltic Sea

M. Kahru et al.



**Figure 10.** Temporal change in the timing of cyanobacteria accumulations for the Baltic Sea. The circles connected with a solid line show the “center of timing” (modified after Kahru and Elmgren, 2014, by adding 2014 data). The red line is the estimated linear regression for the Baltic (area in Fig. 1a). The gray dotted lines show the mean start and end of the accumulations for the whole Baltic Sea. For reference, 1 July is year day 182 (year day 183 in a leap year), year day 200 is 19 July (18 July on a leap year).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

