

1 Global warming potential and greenhouse gas intensity in rice agriculture driven by
2 high yields and nitrogen use efficiency: A 5-year field study

3

4 Xiaoxu Zhang^a, Xin Xu^a, Yinglie Liu^a, Jinyang Wang^{a,b}, Zhengqin Xiong^{a,*}

5 ^a Jiangsu Key Laboratory of Low Carbon Agriculture and GHGs Mitigation, College of
6 Resources and Environmental Sciences, Nanjing Agricultural University, Nanjing, 210095,
7 China

8 ^b State Key Laboratory of Soil and Sustainable Agriculture, Institute of Soil Science, Chinese
9 Academy of Sciences, Nanjing 210008, China

10

11 *Corresponding author (Z. Xiong):

12 Tel: +86-25-84395148,

13 Fax: +86-25-84395210,

14 E-mail: zqxiong@njau.edu.cn

15 **Abstract:** Our understanding of how global warming potential (GWP) and greenhouse gas
16 intensity (GHGI) is affected by management practices aimed at food security with respect to
17 rice agriculture remains limited. In the present study, a 5-year field experiment was conducted
18 in China to evaluate the effects of integrated soil-crop system management (ISSM) mainly
19 consisting of different nitrogen (N) fertilization rates and split, manure, Zn and Na₂SiO₃
20 fertilization and planting density on GWP and GHGI after accounting for carbon dioxide (CO₂)
21 equivalent emissions from all sources including methane (CH₄) and nitrous oxide (N₂O)
22 emissions, agrochemical inputs and farm operations and sinks (i.e., soil organic carbon
23 sequestration). For the improvement of rice yield and agronomic nitrogen use efficiency (NUE),
24 four ISSM scenarios consisting of different chemical N rates relative to the local farmers'
25 practice (FP) rate were carried out, namely, ISSM-N1 (25% reduction), ISSM-N2 (10%
26 reduction), ISSM-N3 (FP rate) and ISSM-N4 (25% increase). The results showed that
27 compared with the FP, the four ISSM scenarios significantly increased the rice yields by 10, 16,
28 28 and 41% and the agronomic NUE by 75, 67, 35 and 40%, respectively. In addition,
29 compared with the FP, the ISSM-N1 and ISSM-N2 scenarios significantly reduced the GHGI
30 by 14 and 18%, respectively, despite similar GWPs. The ISSM-N3 and ISSM-N4 scenarios
31 remarkably increased the GWP and GHGI by an average of 69 and 39%, respectively. In
32 conclusion, the ISSM strategies are promising for both food security and environmental
33 protection, and the ISSM scenario of ISSM-N2 is the optimal strategy to realize high yields
34 and high NUE together with low environmental impacts for this agricultural rice field.

35

36 **1 Introduction**

37 Rapid population growth and economic development place a growing pressure on increasing
38 food production (Barrett, 2010). An increase in global crop production of 100% would be
39 necessary to sustain the projected demand for human food and livestock feed in 2050 (Tilman
40 et al., 2011). Rice is the staple food for nearly 50% of the world's people, mainly in Asia
41 (Frolking et al., 2002). According to FAO (2010), approximately 600 million people in
42 Asia-Pacific region are suffering from hunger and malnutrition. With the region's population
43 projected to increase by another billion by mid-century, new approaches to increase food
44 production are needed (Chen et al., 2014). With a limited agricultural land area, the intensive
45 agricultural regions of China are facing serious environmental problems due to large inputs of
46 chemical fertilizer and low nitrogen use efficiency (NUE) (Ju et al., 2009; Makino, 2011).
47 Thus, integrated soil-crop system management (ISSM), which redesigns the whole production
48 system based on the local environment and draws on appropriate fertilizer compounds and
49 application ratios, crop densities and advanced water management regimes, has been
50 advocated and developed to simultaneously increase crop productivity and NUE with low
51 carbon dioxide (CO₂) equivalent emissions per unit product in China (Chen et al., 2014). The
52 key points of the ISSM are to integrate soil and nutrient management with high-yielding
53 cultivation systems, to integrate the utilization of various nutrient sources and match nutrient
54 supply to crop requirements, and to take all soil quality improvement measures into
55 consideration (Zhang et al., 2011).

56 Carbon dioxide, methane (CH₄) and nitrous oxide (N₂O) are the most important
57 greenhouse gases (GHGs) that contribute to global warming (IPCC, 2013). The concept of
58 global warming potential (GWP) has been applied to agricultural lands by taking in to
59 account of the radiative properties of all GHG emissions associated with agricultural
60 production and soil organic carbon (SOC) sequestration, expressed as CO₂ eq. ha⁻¹ yr⁻¹
61 (Robertson and Grace, 2004; Mosier et al., 2006). Although agriculture releases significant
62 amounts of CH₄ and N₂O into the atmosphere, the net emission of CO₂ equivalents from
63 farming activities can be partly offset by changing agricultural management to increase the
64 soil organic matter content and/or decrease the emissions of CH₄ and N₂O (Mosier et al., 2006;
65 Smith et al., 2008). If global agricultural techniques are improved, the mitigation potential of

66 agriculture (excluding fossil fuel offsets from biomass) is estimated to be approximately 5.5–
67 6.0 Pg CO₂ eq. yr⁻¹ by 2030 (Smith et al., 2008). However, the release of CO₂ during the
68 manufacturing and application of N fertilizer to crops and from fuel used in machines for
69 farm operations can counteract these mitigation efforts (West and Marland, 2002). Therefore,
70 when determining the GWP of agroecosystems, there is a need to account for all sources of
71 GHG emissions, including the emissions associated with agrochemical inputs (Ei) and farm
72 operations (Eo) and sinks, e.g. soil organic carbon (SOC) sequestration (Sainju et al., 2014).

73 Information on the effects of ISSM scenarios on GWP and greenhouse gas intensity
74 (GHGI) of agricultural systems is limited in China (Ma et al., 2013; Liu et al., 2015). The
75 annual rotation of summer rice-upland crop is a dominant cropping system in China. Previous
76 studies were mainly focused on the initial influences of ISSM practices on CH₄ and N₂O
77 emissions, but did not account for the contributions of CO₂ emissions from Ei and Eo (Ma et
78 al., 2013; Zhang et al., 2014). In this study, we evaluated GWP and GHGI of rice-wheat crop
79 rotation managed under several scenarios of ISSM by taking CO₂ equivalents emissions from
80 all sources and sinks into account for 5 years. We hypothesized that the ISSM strategies
81 would reduce the overall GWP and GHGI compared with local farmers' practices (FP). The
82 specific objectives of this study were to (i) evaluate the effects of different ISSM scenarios on
83 GWP and GHGI; (ii) determine the main sources of GWP and GHGI in a rice-wheat cropping
84 system; and (iii) elucidate the overall performance for each ISSM scenario for different
85 targets to increase grain yields and NUE and reduce GWP and GHGI.

86 **2 Materials and Methods**

87 2.1 Experimental site

88 A 5-year field experiment was conducted at the Changshu agro-ecological experimental
89 station (31°32'93"N, 120°41'88"E) in Jiangsu Province, China. This is a typical, intensively
90 managed agricultural area where the cropping regime is dominated by a flooded rice (*Oryza*
91 *sativa* L.)-drained wheat (*Triticum aestivum* L.) rotation system. The site is characterized by a
92 subtropical humid monsoon climate with a mean annual air temperature of 15.6, 15.2 and
93 15.8 °C and precipitation of 878, 1163 and 984 mm for three years, respectively. The soil of
94 the field is classified as an *Anthrosol* with a sandy loam texture of 6% sand (1–0.05 mm), 80%
95 silt (0.05–0.001 mm), and 14% clay (<0.001 mm), which developed from lacustrine sediment.

96 The major properties of the soil at 0–20 cm can be described as follows: bulk density, 1.11 g
97 cm⁻³; pH, 7.35; organic matter content, 35.0 g kg⁻¹; and total N, 2.1 g kg⁻¹. The daily mean air
98 temperatures and precipitation during the study period from June 15, 2011, to June 15, 2014,
99 are given in the supplementary resource 1.

100 2.2 Experimental design and management

101 A completely randomized block design was established in 2009 with four replicates of six
102 treatments, including no nitrogen (NN) and FP as controls, and four ISSM scenarios at
103 different chemical N fertilizer application rates relative to the local FP rate (300 kg N ha⁻¹),
104 namely ISSM-N1 (25% reduction), ISSM-N2 (10% reduction), ISSM-N3 (FP rate) and
105 ISSM-N4 (25% increase). The designed ISSM scenarios (only for rice but not for wheat)
106 included a redesigned split N fertilizer application, a balanced fertilizer application that
107 included sodium silicate, zinc sulphate, rapeseed cake (C/N=8) providing an additional 112.5
108 kg N ha⁻¹, and additional phosphorus and potassium, and different transplanting densities,
109 used as the main techniques for improving rice yield and agronomic NUE. The agronomic
110 NUE was calculated as the difference in grain yield between the plots that received N
111 application and the NN plot, divided by the total N rate which included chemical N fertilizer
112 and rapeseed cake in the ISSM-N3 and ISSM-N4 scenarios. The details of the fertilizer
113 applications, irrigation, and field management practices of the six different treatments are
114 presented in Table 1. Further information was described previously (Zhang et al., 2014). Each
115 plot was 6 m × 7 m in size with an independent drainage/irrigation system.

116 One midseason drainage (about one week) and final drainage before harvest were used
117 during the rice-growing season, whereas the plots only received precipitation during the
118 wheat-growing season. The N fertilizer was split into a 6:2:0:2 or 5:1:2:2 ratio of basal
119 fertilizer and topdressings for the rice crop and a 6:1:3 ratio for the wheat crop. All of
120 phosphorous (P), silicon (Si), zinc (Zn) were applied as basal fertilizers for both crops and
121 rapeseed cake manure was applied for rice crop. Potassium (K) was added as a split (1:1)
122 application to the rice crop and all as basal fertilizer for the wheat crop. The basal fertilization
123 occurred at the time of rice transplanting and wheat seeding. The topdressing was applied at
124 the tillering, elongation and panicle stages of the rice crop and at the seedling establishment
125 and elongation stages of the wheat crop. Aboveground biomass including crop grains and

126 straws were removed out of the fields for all the treatments.

127 2.3 Gas sampling and measurements

128 We measured the CH₄ and N₂O emissions from each plot of the field experiment over five
129 annual cycles from the 2009 rice-growing season to the 2014 wheat-growing season. The
130 initial 2-yr measurements during the 2009–2011 rice-wheat rotational systems were described
131 in our previous study (Ma et al., 2013). Emissions were measured manually using the
132 static-opaque chamber method. Each replicate plot was equipped with a chamber with a size
133 of 50 cm × 50 cm × 50 cm or 50 cm × 50 cm × 110 cm, depending on the crop growth and
134 plant height. The chamber was placed on a fixed PVC frame in each plot and wrapped with a
135 layer of sponge and aluminum foil to minimize air temperature changes inside the chamber
136 during the period of sampling. Gas samples were collected from 9:00 to 11:00 am using an
137 airtight syringe with a 20-ml volume at intervals of 10 min (0, 10, 20 and 30 min after
138 chamber closure). The fluxes were measured once a week and more frequently after fertilizer
139 application or a change in soil moisture.

140 The gas samples were analyzed for CH₄ and N₂O concentrations using a gas
141 chromatograph (Agilent 7890A, Shanghai, China) equipped with two detectors. Methane was
142 detected using a hydrogen flame ionization detector (FID), and N₂O was detected using an
143 electron capture detector (ECD). Argon-methane (5%) and N₂ were used as the carrier gas at a
144 flow rate of 40 ml min⁻¹ for N₂O and CH₄ analysis, respectively. The temperatures for the
145 column and ECD detector were maintained at 40 °C and 300 °C, respectively. The oven and
146 FID were operated at 50 °C and 300 °C, respectively. The CH₄ and N₂O fluxes rate were
147 calculated using a linear increase in the two gas concentrations over time described by Jia et
148 al. (2012).

149 2.4 Topsoil organic carbon sequestration measurements

150 To measure the organic carbon content of the topsoil as described by Zhang et al. (2014), soil
151 samples were collected after the wheat harvest in 2009 and 2014 from all experimental plots
152 at a plowing depth of 0–20 cm. The soil organic carbon sequestration rates (SOCSR) were
153 calculated as follows (Liu et al., 2015):

$$154 \text{SOCSR (t C ha}^{-1} \text{ yr}^{-1}) = (\text{SOC}_t - \text{SOC}_0) / T \times \gamma \times (1 - \delta_{2\text{mm}}/100) \times 20 \times 10^{-1} \text{ (1)}$$

155 In Eq. (1), SOC_t (g C kg⁻¹) and SOC₀ (g C kg⁻¹) are the SOC contents measured in the

156 soils sampled after the wheat was harvested in 2014 and 2009, respectively. T refers to the
157 experimental period (yr). γ and $\delta_{2\text{mm}}$ are the average bulk density and the gravel content (>2
158 mm) of the topsoil (0–20 cm), respectively.

159 2.5 GWP and GHGI measurements

160 To better understand the overall GHG impact of the rice-wheat crop rotation managed
161 under different ISSM scenarios, the GWP and GHGI were calculated as the following
162 equations (Myhre et al., 2013):

$$163 \quad \text{GWP (kg CO}_2 \text{ eq. ha}^{-1} \text{ yr}^{-1}) = 28 \times \text{CH}_4 + 265 \times \text{N}_2\text{O} + \text{Ei} + \text{Eo} - 44/12 \times \text{SOCSR} \quad (2)$$

$$164 \quad \text{GHGI (kg CO}_2 \text{ eq. kg}^{-1} \text{ grain yield yr}^{-1}) = \text{GWP}/\text{grain yield} \quad (3)$$

165 In Eq. (2), Ei (kg CO₂ eq. ha⁻¹ yr⁻¹), Eo (kg CO₂ eq. ha⁻¹ yr⁻¹) and SOCSR (kg C ha⁻¹
166 yr⁻¹) represent CO₂ equivalent emissions from the agrochemical inputs, farm operations and
167 soil organic carbon sequestration rate, respectively. The global warming potential of 1 kg CH₄
168 and 1 kg N₂O are 28 and 265 kg CO₂ equivalents respectively (without inclusion of
169 climate-carbon feedbacks), based on 100-yr time scale (Myhre et al., 2013). 12 and 44 refers
170 to molecular weights of C and CO₂, respectively. The grain yield is expressed as the air-dried
171 grain yield.

172 Therefore, the GWP of the cropland ecosystem equals the total CO₂ equivalent emissions
173 minus the SOC change per unit land area. In addition to CH₄ and N₂O emissions, we
174 considered CO₂ equivalent emissions associated with the use of agrochemical inputs (Ei),
175 such as the manufacture and transportation of the N, P and K fertilizers (Snyder et al., 2009),
176 and farm operations (Eo), such as the water used for irrigation (Zhang et al., 2013) and diesel
177 fuel (Huang et al., 2013a). The CO₂ equivalent emissions of N fertilizer were calculated as the
178 mean value of the C emissions of 1.3 kg C equivalent kg⁻¹ (Lal, 2004). Similarly, the CO₂
179 equivalent for irrigation was calculated from the total amount of water used during the
180 rice-growing season; the coefficient for the C cost was 5.16 (kg C eq. cm⁻¹ ha⁻¹) originated
181 from the value of 257.8 kg C eq. ha⁻¹ for a 50 cm of irrigation provided by Lal (2004). The
182 CO₂ equivalents of other Ei (P and K fertilization, manure, herbicide, pesticide and fungicide
183 applications) and Eo (tillage, planting, harvest, and farm machinery production) were
184 recorded and also estimated by coefficients provided by Lal (2004) since no specific
185 coefficients were available. We collected the data specific to China's fertilizer manufacture

186 and consumption, and obtained the C emission coefficients to be 0.07 and 0.1 kg C eq. kg⁻¹ of
187 active ingredient for Si and Zn fertilizer, respectively. The C emission factor for these farm
188 operations depends on diesel used as fuel or electricity. Chemical fertilizers were hand
189 spraying broadcasted for each fertilization event. Detailed information of each Ei and Eo
190 component for rice and wheat crop season was presented in Supplementary resource 2.

191 2.6 Statistical analysis

192 Repeated-measures multivariate analysis of variance (MANOVA) and linear relationships
193 were determined using JMP 7.0, ver. 7.0 (SAS Institute, USA, 2007). The F-test was applied
194 to determine whether there were significant differences among practices, years and their
195 interaction at $P < 0.05$. One-way analysis of variance was conducted to determine the
196 emissions of CH₄ and N₂O, and the grain yield among the different treatments. Tukey's HSD
197 test was used to determine whether significant differences occurred between the treatments at
198 a level of $P < 0.05$. Normal distribution and variance uniformity were checked and all data
199 were consistent with the variance uniformity ($P > 0.05$) within each group. The results are
200 presented as the means and standard deviation (mean \pm SD, n = 4).

201 3 Results

202 3.1 Crop production and agronomic NUE

203 During the three cropping rotations from 2011 to 2014, the rice and wheat yields varied
204 significantly among the treatments (Table 2). The grain yields ranged from 5.83 to 12.11 t
205 ha⁻¹ for rice and 1.75 to 6.14 t ha⁻¹ for wheat. On average over the three cycles, the annual
206 rice yield of the FP was significantly lower than that of the ISSM scenarios of ISSM-N1,
207 ISSM-N2, ISSM-N3 and ISSM-N4. Compared with the FP, rice grain yields increased by 10%
208 and 16% for the ISSM-N1 and ISSM-N2 scenarios, respectively, i.e., with the lower N input,
209 by 28% for the ISSM-N3 scenario with the same N input and by 41% for the ISSM-N4
210 scenario with the highest N input. However, we did not observe any significant increases in
211 the wheat-grain yields compared with the FP except for the ISSM-N4 scenario. Statistical
212 analysis indicated that rice and wheat yields from the three years were not significantly
213 influenced by the interaction of cultivation patterns and cropping year (Table 3).

214 The agronomic NUE for the rice and wheat of the fertilized plots ranged from 9.2 to 16.1
215 and 19.5 to 24.7 kg grain kg N⁻¹, respectively (Fig. 1). The higher NUE in the wheat season

216 was mainly due to the relatively lower N fertilizer (40%) rates used for wheat compared with
217 that for rice. As expected, the rice agronomic NUE significantly increased by 75, 67, 35 and
218 40% for the ISSM-N1, ISSM-N2, ISSM-N3 and ISSM-N4 scenarios, respectively, compared
219 with the FP (Fig. 1). For the wheat crop, the agronomic NUE increased by 12 and 14% in the
220 ISSM-N1 and ISSM-N2 scenarios, respectively, and slightly decreased in the ISSM-N3 and
221 ISSM-N4 scenarios compared with the FP, mainly because the current ISSM strategy was
222 only designed for rice and not wheat production.

223 3.2 CH₄ and N₂O emissions

224 All plots showed similar CH₄ emission patterns, being a source in the rice season and
225 negligible in the wheat season (Fig. 2). During the three annual rice-wheat rotations from
226 2011 to 2014, the CH₄ fluxes ranged from -3.89 to 99.67 mg C m⁻² h⁻¹. The seasonal CH₄
227 emissions varied significantly among the treatments during the rice-growing season (Table 3,
228 Fig. 2). No significant difference was found between the FP, ISSM-N1 and ISSM-N2 plots.
229 Temporal variation was significant during the three cycles (Table 3, *P* < 0.001). Averaged
230 across years, the CH₄ emission was greater in the ISSM-N3 and ISSM-N4 plots than in the
231 NN, FP, ISSM-N1 and ISSM-N2 plots (Table 2, *P* < 0.05). However, compared with the NN
232 plots, the FP, ISSM-N1 and ISSM-N2 plots with inorganic fertilizer application resulted in
233 increased CH₄ emission rates of 59.9, 41.9 and 43.0%, respectively, averaged over the
234 rice-growing seasons. The CH₄ emission rates were further enhanced by 198.5% in the
235 ISSM-N3 plots and by 246.7% in the ISSM-N4 plots.

236 The annual N₂O fluxes varied from -33.1 to 647.5 μg N₂O-N m⁻² h⁻¹, with most N₂O
237 emissions occurring during the wheat-growing season after fertilization events, and several
238 small emission peaks during the rice-growing season (Fig. 3). With respect to the N
239 application effect, the annual cumulative N₂O emissions for all four ISSM scenarios were
240 significantly higher than that in NN (*P* < 0.05). Relative to the FP plot, the ISSM-N1 and
241 ISSM-N2 scenarios decreased the annual N₂O emissions by an average of 41% and 22%,
242 respectively (Table 2). The ISSM-N4 scenario significantly increased the cumulative N₂O
243 emissions by 46% (*P* < 0.05) because this system received highest inorganic N fertilizer (25%
244 higher than that in FP) and additional N via manure application compared to the FP practice,
245 although there was no significant difference between the ISSM-N3 and FP plots.

246 3.3 Annual GWP and GHGI

247 Based on the perspective of the carbon footprint, we included the GHG emissions associated
248 with all of the inputs (Ei and Eo), and SOC sequestration was expressed as kg CO₂ eq. ha⁻¹
249 yr⁻¹. The CO₂ equivalent emissions associated with Ei and Eo are presented in Table 4. The
250 CO₂ equivalents rates from N fertilizer dominated not only the chemical input section (67–75%
251 of Ei) but also the total CO₂ equivalents from agricultural management (45–50% of the sum
252 of the Ei and Eo). And irrigation was the second largest source of CO₂ equivalents associated
253 with agricultural management after N fertilizer (19–31% of the sum of the Ei and Eo). The
254 GWP ranged from 8425 to 22711 kg CO₂ eq. ha⁻¹ yr⁻¹ for the NN and the ISSM-N4 plots,
255 respectively (Table 5). Although fertilized treatments increased the annual CH₄ and N₂O
256 emissions in comparison with the NN plot, it also increased the SOC sequestration in these
257 cropping systems. Of the main field GHGs that were directly emitted, CH₄ accounted for 59–
258 78% of the GWP in all plots. An increase in the annual SOC content led to a significant
259 decrease in the GWP (contributed 5–9% decrease of the GWP except in the NN plot). The
260 CO₂ equivalents from agricultural management practices, emissions associated with Ei
261 (2493–4300 CO₂ eq. ha⁻¹ yr⁻¹) were higher than those associated with Eo (1296–1708 CO₂ eq.
262 ha⁻¹ yr⁻¹) in the fertilized plots. There was no significant difference in the annual GWP
263 observed between the FP, ISSM-N1 and ISSM-N2 plots (Table 5). Across the three years,
264 ISSM-N1 and ISSM-N2 slightly reduced the GWP by 12 and 10%, respectively; however,
265 ISSM-N3 and ISSM-N4 significantly increased the GWP by an average of 55 and 84%,
266 respectively, in comparison with the FP.

267 The GHGI was used to express the relationship between GWP and grain yield. The
268 GHGIs (kg CO₂ eq. t⁻¹ grain) in this study ranged from 712 to 1245 kg CO₂ eq. t⁻¹ grain
269 (Table 5). The significant difference in the GHGI of grain was found between the FP and the
270 ISSM strategies. Compared with the FP, ISSM-N1 and ISSM-N2 significantly reduced the
271 GHGI by 14 and 18%, respectively, mainly due to the increased grain yield and SOC
272 sequestration as well as reduced GHG emissions for the ISSM strategies of reasonable N
273 fertilizer management and suitable planting density. Although N fertilizer or
274 organic/inorganic combination fertilizer application reduced the SOC losses caused by crop
275 cultivation and increased the grain yields, the GHGIs were generally higher for the ISSM-N3

276 and ISSM-N4 scenarios than the ISSM-N1 and ISSM-N2 scenarios due to further increases in
277 CH₄ and N₂O emissions.

278 **4 Discussion**

279 4.1 Grain yield and agronomic NUE as affected by ISSM strategies

280 Grain yields are directly related to fertilizer management. The MANOVA results indicated
281 that the rice and wheat grain yields were significantly affected by the cultivation strategies
282 (Table 3, $P < 0.001$), which is in agreement with previous results (Chen et al., 2011; Zhang et
283 al., 2011). Compared with the FP, rice yields increased significantly by all four ISSM
284 scenarios (Table 2). However, the wheat grain yield decreased significantly when the N
285 fertilizer rate was reduced by 25% (N1 scenario). It has been reported in previous studies that
286 ISSM strategies can effectively improve the rice grain yield (Ma et al., 2013; Liu et al., 2015).
287 First, the adjusted transplanting density for the ISSM-N1, ISSM-N2, and ISSM-N3 scenarios
288 would produce a positive effect on rice yield by influencing rice colony structure, which
289 agreed with Wu et al. (2005). Second, split application of N fertilizer to match crop demand in
290 the ISSM-N1, ISSM-N2, ISSM-N3 and ISSM-N4 scenarios would significantly increase
291 agronomic NUE and rice yield which had been reported previously by Liu et al. (2009). In the
292 present study, ISSM-N1 and ISSM-N2 significantly increased annual rice production by 10
293 and 16%, respectively, in comparison with the FP (Table 2). This finding is consistent with the
294 results of Peng et al. (2006), who reported that a 30% reduction in the total N rate during the
295 early vegetative stage did not reduce the yield but slightly increased it when combined with
296 the modified farmers' fertilizer practice. Third, integrated management of three
297 macronutrients: N, P and K as well as the two micronutrients: Si and Zn were considered as
298 essential for sustainable high crop yields. Additional Si and Zn fertilizers for the ISSM-N3
299 and ISSM-N4 scenarios would support better seedling establishment and reduce both biotic
300 and abiotic stress, thus produced higher yields (Wang et al., 2005; Slaton et al., 2005;
301 Kabata-Pendias and Mukherjee, 2007; Hossain et al., 2008). As expected, when the total N
302 rate was at the FP rate and/or increased by 25%, in combination with other ISSM strategies
303 (e.g. rapeseed cake manure, additional P and K, applying Si and Zn fertilizer), the rice yield in
304 these ISSM-N3 and ISSM-N4 plots increased substantially by 28 and 41%, respectively.

305 Based on a long-term fertilizer experiment, Shang et al. (2011) reported that organic fertilizer
306 incorporation significantly increased the early rice grain yield. This may have resulted from
307 the organic fertilizer applied in combination with adequate nutrients contributing to alleviate
308 potential yield limiting factors of rice.

309 It has been suggested that N losses vary depending on the timing, rate, and method of N
310 application, as well as the source of N fertilizer (Zhu, 1997). In addition to high rates of N and
311 improper timing of N application, rapid N losses (via ammonia volatilization, denitrification,
312 surface runoff, and leaching) are important factors that cause low agronomic NUE of irrigated
313 rice in China (Peng et al., 2006). Compared with the FP plot, the rice agronomic NUE was
314 significantly increased by 75, 67, 35 and 40% under the ISSM-N1, ISSM-N2, ISSM-N3 and
315 ISSM-N4 scenarios, respectively (Fig. 1). The higher rice agronomic NUE in our study over
316 the experimental period could be due to the greatly reduced N losses by leaching and
317 volatilization as well as the improvement of N bioavailability in the rice crop season (Zhao et
318 al., 2015). Organic/inorganic combination fertilizer application also increases uptake by crops
319 compared with the traditional farmers' practice (Peng et al., 2006). These findings suggest
320 that the ISSM strategy is an effective method for improving grain yield and agronomic NUE
321 for future sustainable rice agriculture in China.

322 4.2 CH₄ and N₂O emissions as affected by ISSM strategies

323 During the three years, the annual cumulative CH₄ emissions, on average, varied from 133 to
324 469 kg C ha⁻¹yr⁻¹ (Table 2), and these values fell within the range of 4.1 to 1015.6 kg CH₄
325 ha⁻¹ observed previously in a rice field (Huang et al., 2004). Methane emissions were highest
326 during rice season, but only during the flooding period. Mainly because CH₄ was produced in
327 the anaerobic zones of submerged soils by methanogens and is oxidized into CO₂ by
328 methanotrophs in the aerobic zones of wetland soils and in upland soils (Le Mer and Roger,
329 2001). The MANOVA results indicated that obvious effects of cultivation patterns and
330 years on CH₄ emissions were found during the rice-wheat rotations (Table 3, *P* < 0.001). The
331 CH₄ emissions were not significantly affected by the cycles but affected by crop season
332 (Table 5, Fig. 2). In this study, no significant difference in CH₄ emission was observed
333 between the FP, ISSM-N1 and ISSM-N2 plots. However, compared with the FP plot, the
334 ISSM-N3 and ISSM-N4 scenarios emitted 87 and 118% more CH₄, respectively (Table 5),

335 which is probably due to the incorporation of the organic rapeseed cake manure. Previous
336 reports support the observations that CH₄ emissions were significantly increased with the
337 application of organic amendments (Ma et al., 2009; Thangarajan et al., 2013; Zou et al.,
338 2005). Apparently, additional application of Si and Zn fertilizers had no significant effect on
339 CH₄ and N₂O fluxes, which was consistent with the result of Xie et al. (2015). Moreover, rice
340 growth was found to be significantly increased under the ISSM-N3 and ISSM-N4 scenarios.
341 In this case, the organic matter inputs such as root litter and rhizodeposits in the ISSM-N3 and
342 ISSM-N4 scenarios were probably also higher than in the other plots, and thus soil C input,
343 which served as an additional source of substrates for the methanogens in the rice paddies,
344 likely contributed to the increase in CH₄ emissions (Ma et al., 2009). Finally, because the rice
345 plants acted as the main pathway for CH₄ transports from the soil to the atmosphere, the
346 higher biomass may have facilitated more CH₄ emissions (Yan et al., 2005).

347 Denitrification and nitrification are the main processes that produce N₂O in the soil (Paul
348 et al., 1993). The N₂O emission patterns varied during the rice and wheat growing seasons
349 which were partially associated with the anaerobic conditions prevailing in a rice paddy.
350 Changes in the soil water content strongly influenced the soil N₂O emissions and resulted in
351 negligible N₂O emissions when the rice field was flooded (Fig. 3), which is consistent with
352 previous reports (Akiyama et al., 2005; Murdiyarso et al., 2010). When the soil water content
353 was below saturation, N₂O emissions increase with soil moisture; however, N₂O emissions
354 gradually decreased with the soil saturation condition (Rudaz et al., 1999). A relatively high
355 N₂O peak was observed in the first two weeks of the wheat-growing season (Fig. 3), possibly
356 because soil changes from flooded to drained conditions may have enhanced N₂O release
357 (Deng et al., 2012). Alternation of drainage and flooding may induce large amounts of N₂O
358 emissions, particularly in fertilized systems; this has commonly been shown in earlier studies
359 (Wang et al., 2012; Xiong et al., 2007; Zou et al., 2005). The seasonal and annual rates of N₂O
360 emissions were significantly affected by the cultivation practices and years (Table 3).
361 Compared with the FP plot, the ISSM-N2 scenario significantly decreased the seasonal N₂O
362 emissions in this study, which may have resulted from a reduction in the N fertilizer rate
363 (Table 1, Table 2). The total N₂O emissions decreased by 7–38% and 26–42% in the rice and
364 wheat seasons, respectively, when the conventional N management (300 kg N ha⁻¹ for rice

365 and 180 kg N ha⁻¹ per crop for wheat) changed to optimum N management (225–270 kg N
366 ha⁻¹ for rice and 135–162 kg N ha⁻¹ per crop for wheat). It is likely that more N₂O was
367 emitted (Mosier et al., 2006) as a result of the additional N made available to the soil
368 microbes through N fertilizer application, which also probably contributed increased CH₄
369 emissions (Banger et al., 2013). Strategies that can reduce N fertilization rates without
370 influencing crop yields can inevitably lower GHG emissions (Mosier et al., 2006). Nitrogen
371 leaching and volatilization are the important components of reactive N releases but not
372 included in the current GHG budget.

373 4.3 GWP and GHGI as affected by ISSM strategies

374 The GWP in our study (10871–22711 kg CO₂ eq. ha⁻¹) with the ISSM strategies was higher
375 than that in a double-cropping cereal rotation (1346–4684 kg CO₂ eq. ha⁻¹) and a rice-wheat
376 annual rotation (290–4580 kg CO₂ eq. ha⁻¹) reported by Huang et al. (2013b) and Yang et al.
377 (2015), respectively. Dominant CH₄ emissions as well as additional CO₂ emissions due to the
378 use of machinery/equipment for irrigation and farm operations under the ISSM strategies may
379 increase the GWP more than in other cropping systems (emit more CO₂ equivalent emissions
380 of 2439–5694 kg CO₂ eq. ha⁻¹ for agricultural management practices in the present study).
381 However, the current GWP was comparable to that of a double-rice cropping system (13407–
382 26066 kg CO₂ eq. ha⁻¹) (Shang et al., 2011). The GHGIs, which ranged from 0.71 to 1.25 kg
383 CO₂ eq. kg⁻¹ grain in this study, were slightly higher than previous estimates of 0.24–0.74 kg
384 CO₂ eq. kg⁻¹ grain from rice paddies with midseason drainage and organic manure
385 incorporation (Qin et al., 2010; Li et al., 2006) but were lower than the DNDC model
386 estimates for continuous waterlogged paddies (3.22 kg CO₂ eq. kg⁻¹ grain) (Li et al., 2006).
387 Differences in GWP or GHGI were found in the cultivation patterns over the three rice-wheat
388 rotations (Table 5). The ISSM-N1 and ISSM-N2 scenarios with optimized ISSM strategies led
389 to a lower GWP than the FP by a certain extent, but there were not significant differences
390 among the FP, ISSM-N1 and ISSM-N2 plots (Table 5). Compared with the FP, the ISSM-N1
391 and ISSM-N2 scenarios dramatically reduced the GHGI, which was mainly due to higher
392 yields. In spite of the similar GWP compared with the FP plot, the lowest GHGI (0.71 kg CO₂
393 eq. kg⁻¹ grain) was obtained under the ISSM-N2 scenario. This finding is consistent with the
394 suggestion made by Burney et al. (2010), i.e., that the net effect of higher yields offsets

395 emissions. It is well known that CH₄ emissions dominate the GWP in rice paddies (Ma et al.,
396 2013; Shang et al., 2011). In comparison to the GWP (12371 kg CO₂ eq. ha⁻¹yr⁻¹) and GHGI
397 (0.87 kg CO₂ eq. kg⁻¹ grain) of the FP, the ISSM-N3 and ISSM-N4 scenarios increased both
398 the GWP and GHGI, mainly because these scenarios notably increased the CH₄ emissions
399 compared with the FP, which resulted in relatively higher GWP (Table 5).

400 Agricultural management practices that change one type of GWP source/sink may also
401 impact other sources/sinks and therefore change the GWP and GHGI (Mosier et al., 2006;
402 Shang et al., 2011). Although the N-fertilizer plots, especially those with the incorporation of
403 organic fertilizer, increased the annual CH₄ and N₂O emissions, they increased the SOC
404 sequestration in this cropping system, which is agreement with previous reports (Huang and
405 Sun, 2006). This was mainly due to the enhanced incorporation of rapeseed cake and crop
406 residue associated with higher crop productivity (Ma et al., 2013). In the present study, the
407 ISSM-N2 scenario with ISSM strategies decreased the CH₄ and N₂O emissions as well as the
408 energy consumption related to irrigation and the manufacture and transport of N fertilizer
409 (depending on coal combustion), ultimately leading to a decrease in the GWP relative to the
410 FP plot. Moreover, despite the lower N fertilizer input, the grain yield did not decline and the
411 GHGI of the ISSM-N2 scenario was thus lower than of the FP plot, indicating less
412 consumption of CO₂ equivalents per unit of grain produced. We demonstrate that high yield
413 and agronomic NUE, together with low GWP, are not conflicting goals by optimizing ISSM
414 strategies.

415 4.4 Main components of GWP and GHGI and implementation significance for the ISSM 416 strategies

417 Determining the main components of the GWP and GHGI in specific cropping systems is
418 very important for mitigating GHG emissions in the future because the benefits of C
419 sequestration would be negated by CH₄ and N₂O emissions and the CO₂ equivalents released
420 with the use of high N fertilizer application rates (Schlesinger, 2010). In the current study, the
421 five main components of the CO₂ equivalents for the GWP were ranked in decreasing order of
422 importance as follows: CH₄ emissions > agrochemical inputs of N fertilizer > farm operations
423 related to irrigation > SOC sequestration > N₂O emissions (Table 5). In each crop, CH₄ and
424 irrigation were important for rice, but less important for wheat, in which N₂O losses were

425 expected to have a higher weight (Supplementary resource 2). Methane emissions, the most
426 important component of GWP in this typical rice-wheat rotation system, could be further
427 mitigated by some other strategies, such as reasonable irrigation (Zou et al., 2005; Wang et al.,
428 2012).

429 Although N fertilizer application increased SOC sequestration when it was applied
430 with rapeseed cake manure, this benefit was consistently overshadowed, on a CO₂ equivalent
431 basis, by the increases in CH₄ and N₂O emissions (Table 5). Similar results have been
432 reported, i.e., GHG emissions substantially offset SOC increases (Six et al., 2004). It is
433 possible that the realization of reducing the GWP and GHGI in China should focus on
434 increasing the SOC and simultaneously decreasing the CO₂ equivalents from CH₄ emissions
435 and N fertilizer inputs. Several studies reported possible methods for these types of mitigation
436 strategies, such as optimizing the chemical fertilizer application amount and rate (Ju et al.,
437 2011), the amount of water used for irrigation (Gao et al., 2015), and the timing and rate of N
438 using the in-season N management approach, as well as improving the N fertilizer
439 manufacturing technologies (Zhang et al., 2013), and using nitrification inhibitors or
440 polymer-coated controlled-release fertilizers (Hu et al., 2013).

441 China is a rapidly developing country that faces the dual challenge of substantially
442 increasing grain yields at the same time as reducing the very substantial environmental
443 impacts of intensive agriculture (Chen et al., 2011). We used the ISSM strategies to develop a
444 rice production system that achieved mean yields of 10.63 t ha⁻¹ (an increment of almost 24%)
445 and an agronomic NUE of 13.20 kg grain kg N⁻¹ (an increment of 43%) in long-term field
446 experiments compared with current farmers' practices. The ISSM redesigned the whole
447 production system only for the rice crop based on the local environment and drawing on
448 appropriate fertilizer varieties and application ratios, crop densities and an advanced water
449 regime management. If the ISSM strategies were also developed for the rotated wheat crop,
450 the overall performance of the whole rice-wheat system would be much improved, with
451 further increases in yield and reductions in the GWP and GHGI. We conclude that the ISSM
452 strategies are promising, particularly the ISSM-N2 scenario, which is the most favorable to
453 realize higher yields with lower environmental impact. The proposed ISSM strategies can
454 provide substantial benefits to intensive agricultural systems and can be applied feasibly using

455 current technologies.

456 **5 Conclusions**

457 Reasonable agricultural management practices are the key to reducing GHG emissions from
458 agricultural ecosystems. This study provided an insight into the complete GHG emission
459 accounting of the GWP and GHGI affected by different ISSM scenarios. After a five-year
460 field experiment, we found that the CH₄ emissions, production of N fertilizer, irrigation, SOC
461 sequestration and N₂O fluxes were the main components of the GWP in a typical rice-wheat
462 rotation system. In contrast with the FP, ISSM-N1 and ISSM-N2 significantly reduced the
463 GHGI, though they resulted in similar GWPs, and ISSM-N3 and ISSM-N4 remarkably
464 increased the GWP and GHGI. By adopting the ISSM-N2 strategy, the conventional N
465 application rate was reduced by 10% while the rice yield was significantly increased by 16%,
466 the NUE was improved by 67% and the GHGI was lowered. ISSM scenarios could be
467 adopted for both food security and environmental protection with specific targets. We propose
468 that the ISSM-N2 scenario is the most appropriate management strategy (10% reduction of N
469 input, no rapeseed manure and higher plant density) for realizing higher yields and NUE,
470 together with some potential to reduce GHGI by integrated soil-crop management. For
471 simultaneously mitigating GHG emissions, further research on integrated soil-crop system
472 managements is required particularly for mitigating CH₄ emissions in sustainable rice
473 agriculture.

474 **Acknowledgments** We sincerely appreciate two anonymous reviewers for their critical and
475 valuable comments to help improve this manuscript. This work was jointly supported by the
476 National Science Foundation of China (41171238, 41471192), the Special Fund for
477 Agro-Scientific Research in the Public Interest (201503106) and the Ministry of Science and
478 Technology (2013BAD11B01).

479 **References**

- 480 Akiyama, H., Yagi, K. and Yan, X. Y.: Direct N₂O emissions from rice paddy fields: summary of
481 available data, *Global Biogeochem. Cy.*, 19, GB002378, doi:10.1029/2004GB002378, 2005.
- 482 Banger, K., Tian, H. and Lu, C.: Do nitrogen fertilizers stimulate or inhibit methane emissions from
483 rice fields?, *Glob. Change Biol.*, 18, 3259-3267, 2013.
- 484 Barrett, C. B.: Measuring food insecurity, *Science*, 327 (5967), 825-828, 2010.
- 485 Burney, J. A., Davis, S. J. and Lobell, D. B.: Greenhouse gas mitigation by agricultural intensification,
486 *Proc. Natl. Acad. Sci. U.S.A.*, 107, 12052-12057, 2010.
- 487 Chen, X., Cui, Z., Vitousek, P. M., Cassman, K. G., Matson, P. A., Bai, J., Meng, Q., Hou, P., Yue, S.
488 and Römheld, V.: Integrated soil-crop system management for food security, *P. Natl. Acad. Sci.*
489 *USA.*, 108, 6399-6404, 2011.
- 490 Chen, X., Cui, Z., Fan, M., Vitousek, P., Zhao, M., Ma, W., Wang, Z., Zhang, W., Yan, X. and Yang, J.:
491 Producing more grain with lower environmental costs, *Nature*, 514, 486-489, 2014.
- 492 Deng, J., Zhou, Z., Zheng, X., Liu, C., Yao, Z., Xie, B., Cui, F., Han, S. and Zhu, J.: Annual emissions
493 of nitrous oxide and nitric oxide from rice-wheat rotation and vegetable fields: a case study in
494 the Tai-Lake region, China, *Plant Soil*, <http://dx.doi.org/10.1007/s11104-012-1223-6>, 2012.
- 495 Food and Agriculture Organization Statistical Data (FAOSTAT), Available online at:
496 <http://www.fao.org/docrep/003/x6905e/x6905e04.htm>, Rome, Italy, 2010.
- 497 Frolking, S., Qiu, J., Boles, S., Xiao, X., Liu, J., Zhuang, Y., Li, C. and Qin, X.: Combining remote
498 sensing and ground census data to develop new maps of the distribution of rice agriculture in
499 China, *Global Biogeochem. Cy.*, doi:16, 10.1029/2001GB001425, 2002.
- 500 Gao, B., Ju, X., Meng, Q., Cui, Z., Christie, P., Chen, X. and Zhang, F.: The impact of alternative
501 cropping systems on global warming potential, grain yield and groundwater use, *Agr. Ecosyst.*
502 *Environ.*, 203, 46-54, 2015.
- 503 Hossain, M. A., Jahiruddin, M., Islam, M. R., and Mian, M. H.: The requirement of zinc for
504 improvement of crop yield and mineral nutrition in the maize-mungbean-rice system. *Plant*
505 *and Soil*, 306, 13-22, 2008.
- 506 Hu, X., Su, F., Ju, X., Gao, B., Oenema, O., Christie, P., Huang, B., Jiang, R. and Zhang, F.:
507 Greenhouse gas emissions from a wheat-maize double cropping system with different nitrogen
508 fertilization regimes, *Environ. Pollut.*, 176, 198-207, 2013.
- 509 Huang, J., Chen, Y., Sui, P. and Gao, W.: Estimation of net greenhouse gas balance using crop-and
510 soil-based approaches: Two case studies, *Sci. Total Environ.*, 456, 299-306, 2013a.
- 511 Huang, T., Gao, B., Christie, P. and Ju, X.: Net global warming potential and greenhouse gas intensity
512 in a double-cropping cereal rotation as affected by nitrogen and straw management,
513 *Biogeosciences*, 10, 7897-7911, 2013b.
- 514 Huang, Y. and Sun, W.: Changes in topsoil organic carbon of croplands in mainland China over the last
515 two decades, *Chinese Sci Bull.*, 51, 1785-1803, 2006.
- 516 Huang, Y., Zhang, W., Zheng, X. H., Li, J. and Yu, Y. Q.: Modeling methane emission from rice
517 paddies with various agricultural practices, *J. Geophys. Res. Atmos.*, 109, D08,
518 doi:10.1029/2003JD004401, 2004.
- 519 IPCC.: *Climate Change 2013: The Physical Science Basis: working group I contribution to the Fifth*
520 *Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge University
521 Press, Stockholm, 2013.
- 522 Jia, J. X., Ma, Y. C. and Xiong, Z. Q.: Net ecosystem carbon budget, net global warming potential and

- 523 greenhouse gas intensity in intensive vegetable ecosystems in China, *Agr. Ecosyst. Environ.*,
524 150, 27-37, 2012.
- 525 Ju, X., Xing, G., Chen, X., Zhang, S., Zhang, L., Liu, X., Cui, Z., Yin, B., Christie, P. and Zhu, Z.:
526 Reducing environmental risk by improving N management in intensive Chinese agricultural
527 systems, *P. Natl. Acad. Sci. USA*, 106, 3041-3046, 2009.
- 528 Ju, X., Lu, X., Gao, Z., Chen, X., Su, F., Kogge, M., Römheld, V., Christie, P. and Zhang, F.: Processes
529 and factors controlling N₂O production in an intensively managed low carbon calcareous soil
530 under sub-humid monsoon conditions, *Environ. Pollut.*, 159, 1007-1016, 2011.
- 531 Kabata-Pendias, A. and Mukherjee, A. B.: Trace elements from soil to human. Springer Science &
532 Business Media, 2007.
- 533 Lal, R.: Carbon emission from farm operations, *Environ. Int.*, 30, 981-990, 2004.
- 534 Le Mer, J. and Roger, P.: Production, oxidation, emission and consumption of methane by soils: a
535 review, *European Journal of Soil Biology*, 37, 25-50, 2001.
- 536 Li, C. S., Salas, W., DeAngelo, B. and Rose, S.: Assessing alternatives for mitigating net greenhouse
537 gas emissions and increasing yields from rice production in China over the next twenty years,
538 *J. Environ. Qual.*, 35, 1554-1565, 2006.
- 539 Liu, L., Xue Y., Sun, X., Wang, Z. and Yang, J.: Effects of water management methods on grain yield
540 and fertilizer-nitrogen use efficiency in rice, *Chin J Rice Sci.*, 23, 282-288, 2009.
- 541 Liu, Y. L., Zhou, Z., Zhang, X., Xu, X., Chen, H. and Xiong, Z.: Net global warming potential and
542 greenhouse gas intensity from the double rice system with integrated soil-crop system
543 management: A three-year field study, *Atmos. Environ.*, 116, 92-101, 2015.
- 544 Ma, J., Ma, E., Xu, H., Yagi, K. and Cai, Z.: Wheat straw management affects CH₄ and N₂O emissions
545 from rice fields, *Soil Biol. Biochem.*, 41, 1022-1028, 2009.
- 546 Ma, Y., Kong, X., Yang, B., Zhang, X., Yan, X., Yang, J. and Xiong, Z.: Net global warming potential
547 and greenhouse gas intensity of annual rice-wheat rotations with integrated soil-crop system
548 management, *Agr. Ecosyst. Environ.*, 164, 209-219, 2013.
- 549 Makino, A.: Photosynthesis, grain yield, and nitrogen utilization in rice and wheat, *Plant physiol.*, 155,
550 125-129, 2011.
- 551 Mosier, A. R., Halvorson, A. D., Reule, C. A. and Liu, X. J.: Net global warming potential and
552 greenhouse gas intensity in irrigated cropping systems in northeastern Colorado, *J. Environ.*
553 *Qual.*, 35, 1584-1598, 2006.
- 554 Murdiyarso, D., Hergoualc'h, K. and Verchot, L. V.: Opportunities for reducing greenhouse gas
555 emissions in tropical peatlands, *P. Natl. Acad. Sci. USA*, 107, 19655-19660, 2010.
- 556 Myhre, G., Shindell, D., Bréon, F.M., Collins, W., Fuglestedt, J., Huang, J., Koch, D., Lamarque, J.F.,
557 Lee, D., Mendoza, B., Nakajima, T., Robock, A., Stephens, G., Takemura, T. and Zhang, H.:
558 Anthropogenic and natural radiative forcing. In: Stocker, T.F., Qin, D., Plattner, G.K., Tignor,
559 M., Allen, S.K., Boschung, J., Nauels, A., Xia, Y., Bex, V., Midgley, P.M. (Eds.), *Climate*
560 *Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth*
561 *Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University*
562 *Press, Cambridge, UK, New York, NY, USA, 659-740, 2013.*
- 563 Paul, J.W., Beauchamp, E.G. and Zhang, X.: Nitrous and nitric oxide emissions during nitrification and
564 denitrification from manure-amended soil in the laboratory, *Can. J. Soil. Sci.*, 73, 539-553,
565 1993.
- 566 Peng, S., Buresh, R. J., Huang, J., Yang, J., Zou, Y., Zhong, X., Wang, G. and Zhang, F.: Strategies for

567 overcoming low agronomic nitrogen use efficiency in irrigated rice systems in China, *Field*
568 *Crop. Res.*, 96, 37-47, 2006.

569 Qin, Y. M., Liu, S.W., Guo, Y. Q., Liu, Q. and Zou, J. W.: Methane and nitrous oxide emissions from
570 organic and conventional rice cropping systems in Southeast China, *Biol. Fertil. Soils.*, 46,
571 825-834, 2010.

572 Robertson, G. P. and Grace, P. R.: Greenhouse gas fluxes in tropical and temperate agriculture: The
573 need for a full-cost accounting of global warming potentials, *Environment Development and*
574 *Sustainability*, 6, 51-63, 2004.

575 Rudaz, A. O., Wälti, E., Kyburz, G., Lehmann, P. and Fuhrer, J.: Temporal variation in N₂O and N₂
576 fluxes from a permanent pasture in Switzerland in relation to management, soil water content
577 and soil temperature, *Agr. Ecosyst. Environ.*, 73, 83-91, 1999.

578 Sainju, U. M., Stevens, W. B., Caesar-TonThat, T., Liebig, M. A. and Wang, J.: Net global warming
579 potential and greenhouse gas intensity influenced by irrigation, tillage, crop rotation, and
580 nitrogen fertilization, *J. Environ. Qual.*, 43, 777-788, 2014.

581 Schlesinger, W. H.: On fertilizer-induced soil carbon sequestration in China's croplands, *Glob. Change*
582 *Biol.*, 16, 849-850, 2010.

583 Shang, Q., Yang, X., Gao, C., Wu, P., Liu, J., Xu, Y., Shen, Q., Zou, J. and Guo, S.: Net annual global
584 warming potential and greenhouse gas intensity in Chinese double rice-cropping systems: a
585 3-year field measurement in long-term fertilizer experiments, *Glob. Change Biol.*, 17,
586 2196-2210, 2011.

587 Six, J., Ogle, S. M., Conant, R. T., Mosier, A. R. and Paustian, K.: The potential to mitigate global
588 warming with no-tillage management is only realized when practised in the long term, *Glob.*
589 *Change Biol.*, 10, 155-160, 2004.

590 Slaton, N. A., Norman, R. J. and Wilson, C. E.: Effect of zinc source and application time on zinc
591 uptake and grain yield of flood-irrigated rice, *Agron. J.*, 97, 272-278, 2005.

592 Smith, P., Martino, D., Cai, Z., Gwary, D., Janzen, H., Kumar, P., McCarl, B., Ogle, S., O'Mara, F. and
593 Rice, C.: Greenhouse gas mitigation in agriculture, *Philos. T. R. Soc. B.*, 363, 789-813, 2008.

594 Snyder, C., Bruulsema, T., Jensen, T. and Fixen, P.: Review of greenhouse gas emissions from crop
595 production systems and fertilizer management effects, *Agr. Ecosyst. Environ.*, 133, 247-266,
596 2009.

597 Thangarajan, R., Bolan, N. S., Tian, G., Naidu, R. and Kunhikrishnan, A.: Role of organic amendment
598 application on greenhouse gas emission from soil, *Sci. Total Environ.*, 465, 72-96, 2013.

599 Tilman, D., Balzer, C., Hill, J., Befort, B. L. and Affiliations, A.: Global food demand and the
600 sustainable intensification of agriculture, *P. Natl. Acad. Sci. USA*, 108, 20260-20264, 2011.

601 Wang, J., Zhang, X., Xiong, Z., Khalil, M.A.K., Zhao, X., Xie, Y. and Xing, G.: Methane emissions
602 from a rice agroecosystem in South China: Effects of water regime, straw incorporation and
603 nitrogen fertilizer, *Nutr. Cycl. Agroecosyt.*, 93, 103-112, 2012.

604 Wang, L.J., Nie, Q., Li, M., Zhang, F.S., Zhuang, J.Q., Yang, W.S., Li, T.J. and Wang, Y.H.:
605 Biosilicified structures for cooling plant leaves: a mechanism of highly efficient midinfrared
606 thermal emission, *Appl. Phys. Lett.*, 87, 194105, doi:10.1063/1.2126115, 2005.

607 West, T. O. and Marland, G.: A synthesis of carbon sequestration, carbon emissions, and net carbon
608 flux in agriculture: comparing tillage practices in the United States, *Agr. Ecosyst. Environ.*, 91,
609 217-232, 2002.

610 Wu, C., Ye, D., Lin, H., Ni, R., Lai L. and Lin, H.: Effects of transplanting density on rice yield and its

611 quality, Chinese Agricultural Science Bulletin, 21, 190-205, 2005.

612 Xie, Y., Zhang, J., Jiang, H., Yang, J., Deng, S., Li, X., Guo, J., Li, L., Liu, X. and Zhou, G.: Effects of
613 different fertilization practices on greenhouse gas emissions from paddy soil, Journal of
614 Agro-Environment Science, 3, 578-584, 2015.

615 Xiong, Z. Q., Xing, G. X. and Zhu, Z. L.: Nitrous oxide and methane emissions as affected by water,
616 soil and nitrogen, Pedosphere, 17, 146-155, 2007.

617 Yan, X. Y., Yagi, K., Akiyama, H. and Akimoto, H.: Statistical analysis of the major variables
618 controlling methane emission from rice fields, Glob. Change Biol., 11, 1131-1141, 2005.

619 Yang, B., Xiong, Z., Wang, J., Xu, X., Huang, Q. and Shen, Q.: Mitigating net global warming
620 potential and greenhouse gas intensities by substituting chemical nitrogen fertilizers with
621 organic fertilization strategies in rice-wheat annual rotation systems in China: A 3-year field
622 experiment, Ecol. Eng., 81, 289-297, 2015.

623 Zhang, F., Cui, Z., Fan, M., Zhang, W., Chen, X. and Jiang, R.: Integrated soil-crop system
624 management: reducing environmental risk while increasing crop productivity and improving
625 nutrient use efficiency in China, J. Environ Qual., 40, 1051-1057, 2011.

626 Zhang, W., Dou, Z., He, P., Ju, X., Powlson, D., Chadwick, D., Norse, D., Lu, Y., Zhang, Y. and Wu, L.:
627 New technologies reduce greenhouse gas emissions from nitrogenous fertilizer in China, P.
628 Natl. Acad. Sci. USA, 110, 8375-8380, 2013.

629 Zhang, X., Fan, C., Ma, Y., Liu, Y., Li, L., Zhou, Q. and Xiong, Z.: Two approaches for net ecosystem
630 carbon budgets and soil carbon sequestration in a rice-wheat rotation system in China, Nutr.
631 Cycl. Agroecosys., 100, 301-313, 2014.

632 Zhao, M., Tian, Y., Ma, Y., Zhang, M., Yao, Y., Xiong, Z., Yin, B. and Zhu, Z.: Mitigating gaseous
633 nitrogen emissions intensity from a Chinese rice cropping system through an improved
634 management practice aimed to close the yield gap, Agr. Ecosyst. Environ., 203, 36-45, 2015.

635 Zhu, Z.: Fate and management of fertilizer nitrogen in agro-ecosystems. Nitrogen in Soils of China,
636 Springer Netherlands, 239-279, 1997.

637 Zou, J. W., Huang, Y., Jiang, J. Y., Zheng, X. H. and Sass, R.L.: A 3-year field measurement of methane
638 and nitrous oxide emissions from rice paddies in China: effects of water regime, crop residue,
639 and fertilizer application, Global Biogeochem. Cy., 19, GB002401,
640 doi:10.1029/2004GB002401, 2005.

Table 1

The establishment of different treatments for the annual rice-wheat rotations during the 2011–2014 cycle.

| Scenario | NN ^a | FP | ISSM-N1 | ISSM-N2 | ISSM-N3 | ISSM-N4 |
|--|----------------------|----------------|----------------|----------------|--------------------|--------------------|
| Rice-growing season | | | | | | |
| Chemical fertilizer application rate (N:P ₂ O ₅ :K ₂ O:Na ₂ SiO ₃ :ZnSO ₄ , kg ha ⁻¹) | 0:90:120:0:0 | 300:90:120:0:0 | 225:90:120:0:0 | 270:90:120:0:0 | 300:108:144:225:15 | 375:126:180:225:15 |
| Split N application ratio | | 6:2:0:2 | 5:1:2:2 | 5:1:2:2 | 5:1:2:2 | 5:1:2:2 |
| Rapeseed cake manure (t ha ⁻¹) | 0 | 0 | 0 | 0 | 2.25 ^c | 2.25 |
| Water regime | F-D-F-M ^b | F-D-F-M | F-D-F-M | F-D-F-M | F-D-F-M | F-D-F-M |
| Planting density (cm) | 20×20 | 20×20 | 20×15 | 20×15 | 20×15 | 20×20 |
| Wheat-growing season | | | | | | |
| Chemical fertilizer application rate (N:P ₂ O ₅ :K ₂ O, kg ha ⁻¹) | 0:90:180 | 180:90:180 | 135:90:180 | 162:90:180 | 180:108:216 | 225:126:270 |
| Split N application ratio | | 6:1:3 | 6:1:3 | 6:1:3 | 6:1:3 | 6:1:3 |
| Seed sowing density (kg ha ⁻¹) | 180 | 180 | 180 | 180 | 180 | 180 |

^aNN, no N application; FP, farmers' practice; The four integrated soil-crop system management (ISSM) practices at different nitrogen application rates relative to the FP rate of 300 kg N ha⁻¹ for the rice crop and 180 kg N ha⁻¹ for the wheat crop, namely, ISSM-N1 (25% reduction), ISSM-N2 (10% reduction), ISSM-N3 (FP rate) and ISSM-N4 (25% increase). Urea, calcium biphosphate and potassium chloride were used as N, P and K fertilizer respectively.

^bF-D-F-M, flooding-midseason drainage-re-flooding-moist irrigation.

^c112.5 kg N ha⁻¹ in the form of rapeseed cake was applied as basal fertilizer and included in the total N rate for calculating agronomic NUE.

Table 2Seasonal CH₄ and N₂O emissions, and yields during rice and wheat cropping seasons in the three cycles of 2011–2014.

| Treatment | Rice season | | | Wheat season | | |
|--------------------------------|---|--|--------------------------------|---|--|--------------------------------|
| | CH ₄ (kg C ha ⁻¹) | N ₂ O (kg N ha ⁻¹) | Yield (t ha ⁻¹) | CH ₄ (kg C ha ⁻¹) | N ₂ O (kg N ha ⁻¹) | Yield (t ha ⁻¹) |
| 2011 | | | | | | |
| NN | 153±10.8c | 0.03±0.05c | 5.85±0.08f | - 0.48±0.63a | 0.45±0.09d | 1.74±0.18d |
| FP | 266±25.3b | 0.11±0.08c | 8.38±0.35e | - 0.48±1.86a | 1.43±0.19b | 5.67±0.20b |
| ISSM-N1 | 212±30.3bc | 0.08±0.03c | 9.27±0.26d | 0.78±0.97a | 0.65±0.11cd | 5.05±0.16c |
| ISSM-N2 | 220±32.5bc | 0.17±0.11bc | 9.79±0.44c | 2.25±2.07a | 0.80±0.06c | 5.71±0.18b |
| ISSM-N3 | 518±58.9a | 0.38±0.15ab | 10.81±0.26b | 0.04±3.23a | 1.40±0.10b | 5.31±0.26bc |
| ISSM-N4 | 561±50.9a | 0.37±0.07a | 11.76±0.24a | - 0.09±1.40a | 1.93±0.09a | 6.15±0.15a |
| 2012 | | | | | | |
| NN | 149±25.8d | 0.13±0.10c | 5.80±0.22f | - 4.32±7.29a | 0.65±0.09d | 1.73±0.11c |
| FP | 239±34.5c | 0.33±0.11bc | 8.72±0.62e | 4.85±10.30a | 2.13±0.43ab | 5.64±0.34ab |
| ISSM-N1 | 226±30.4cd | 0.27±0.07bc | 9.43±0.34d | 1.46±6.38a | 1.39±0.14c | 4.94±0.38b |
| ISSM-N2 | 228±32.6cd | 0.38±0.29bc | 9.99±0.50c | - 1.02±0.84a | 1.77±0.38bc | 5.78±0.59ab |
| ISSM-N3 | 431±26.8b | 0.52±0.16ab | 10.92±0.61b | 2.45±8.35a | 2.19±0.24ab | 5.39±0.39ab |
| ISSM-N4 | 536±58.7a | 0.78±0.13a | 12.24±0.60a | 5.91±6.18a | 2.61±0.42a | 6.10±0.49a |
| 2013 | | | | | | |
| NN | 101±39.2b | 0.16±0.09b | 5.84±0.15f | - 1.45±1.34a | 0.35±0.06c | 1.80±0.03c |
| FP | 141±25.2b | 0.43±0.39ab | 8.67±0.26e | - 3.70±1.76a | 0.80±0.20ab | 5.70±0.30ab |
| ISSM-N1 | 135±15.7b | 0.19±0.16ab | 9.66±0.29d | - 1.00±1.61a | 0.49±0.16bc | 5.15±0.20b |
| ISSM-N2 | 129±32.2b | 0.26±0.13ab | 10.15±0.07c | - 0.79±1.60a | 0.69±0.24abc | 5.80±0.18ab |
| ISSM-N3 | 256±45.6a | 0.59±0.42ab | 11.14±0.10b | - 0.62±1.14a | 0.71±0.10ab | 5.51±0.33ab |
| ISSM-N4 | 304±22.3a | 0.74±0.40a | 12.34±0.16a | 0.55±1.68a | 1.02±0.11a | 6.19±0.63a |
| Average 2011–2013 ^a | | | | | | |
| NN ^b | 135±19.6d | 0.11±0.05c | 5.83±0.04f | - 2.08±1.89a | 0.48±0.07d | 1.75±0.04d |
| FP ^b | 215±19.9c | 0.29±0.13bc | 8.59±0.25e | 0.22±3.96a | 1.45±0.24b | 5.67±0.16b |
| ISSM-N1 ^b | 191±19.2c | 0.18±0.06c | 9.45±0.18d | 0.42±2.77a | 0.84±0.08c | 5.04±0.08c |
| ISSM-N2 ^b | 192±11.6c | 0.27±0.12bc | 9.98±0.25c | 0.15±0.58a | 1.08±0.12c | 5.76±0.22ab |
| ISSM-N3 ^b | 402±23.8b | 0.50±0.16ab | 10.95±0.13b | 0.63±3.51a | 1.43±0.05b | 5.40±0.16bc |
| ISSM-N4 ^b | 467±39.2a | 0.68±0.15a | 12.11±0.28a | 2.12±2.57a | 1.85±0.16a | 6.14±0.35a |

^aMean ± SD, different lower case letters within the same column for each item indicate significant differences at $P < 0.05$ according to Tukey's multiple range test.

^bSee Table 1 for treatment codes.

Table 3

Repeated-measures analysis of variance (MANOVA) for the effects of cultivation patterns (P) and cropping year (Y) on mean CH₄ and N₂O emissions, and mean rice and wheat grain yields in the 2011–2014 cycle.

| Crop season | Source | <i>df</i> | CH ₄ (kg C ha ⁻¹) | N ₂ O (kg N ha ⁻¹) | Yield (t ha ⁻¹) |
|-------------|------------------|-----------|---|--|--------------------------------|
| Rice | Between subjects | | | | |
| | P | 5 | 35.3*** | 3.71*** | 123*** |
| | Within subjects | | | | |
| | Y | 2 | 20.7*** | 0.88** | 1.15** |
| | P×Y | 10 | 6.73*** | 0.15 | 0.37 |
| Wheat | Between subjects | | | | |
| | P | 5 | 0.26 | 14.8*** | 76.3*** |
| | Within subjects | | | | |
| | Y | 2 | 0.55* | 15.1*** | 0.08 |
| | P×Y | 10 | 0.83 | 4.39*** | 0.05 |
| Rice-Wheat | Between subjects | | | | |
| | P | 5 | 37.2*** | 24.2*** | 153*** |
| | Within subjects | | | | |
| | Y | 2 | 20.5*** | 5.83*** | 0.70* |
| | P×Y | 10 | 6.50*** | 1.11 | 0.17 |

df – degrees of freedom, * $P < 0.05$, ** $P < 0.01$, and *** $P < 0.001$ represent significant at the 0.05, 0.01 and 0.001 probability level, respectively.

Table 4

Agricultural management practices for chemical inputs and farm operations and contributions to carbon dioxide equivalents (kg CO₂ eq. ha⁻¹yr⁻¹) in the annual rice-wheat rotations from 2011 to 2014 (chemical inputs and farm operations used in each year were similar except for irrigation water).

| Treatment | Chemical inputs (kg ha ⁻¹) ^a | | | | | | | | Farm operations (kg ha ⁻¹) ^c | | | | | | | |
|-----------------|---|-----|-----|-----|----|-----------|-------------|-----------|---|------|------|--------------------|-----------------------|-------------|--------------|---------------------------|
| | N | P | K | Si | Zn | Herbicide | Insecticide | Fungicide | Irrigation (cm) ^b | | | Tillage and raking | Crop planting (event) | Farm manure | Crop harvest | Farm machinery production |
| | | | | | | | | | 2011 | 2012 | 2013 | | | | | |
| NN ^d | 0 | 180 | 300 | 0 | 0 | 2 | 18 | 4 | 75 | 80 | 80 | 37 | 2 | 0 | 11 | 135 |
| FP | 480 | 180 | 300 | 0 | 0 | 2 | 20 | 4.4 | 75 | 80 | 80 | 37 | 2 | 0 | 11 | 147 |
| ISSM-N1 | 360 | 180 | 300 | 0 | 0 | 2 | 20 | 4.4 | 50 | 65 | 55 | 37 | 2 | 0 | 11 | 139 |
| ISSM-N2 | 432 | 180 | 300 | 0 | 0 | 2 | 20 | 4.4 | 50 | 65 | 55 | 37 | 2 | 0 | 11 | 177 |
| ISSM-N3 | 480 | 216 | 360 | 225 | 15 | 2 | 27 | 6 | 50 | 65 | 55 | 37 | 2 | 2250 | 11 | 177 |
| ISSM-N4 | 600 | 252 | 450 | 225 | 15 | 2 | 41 | 9 | 50 | 65 | 55 | 37 | 2 | 2250 | 11 | 275 |
| | Chemical inputs (Ei) (kg CO ₂ eq. ha ⁻¹) | | | | | | | | Farm operations (Eo) (kg CO ₂ eq. ha ⁻¹) | | | | | | | |
| NN | 0 | 132 | 165 | 0 | 0 | 46 | 338 | 53 | 1419 | 1514 | 1514 | 127 | 23 | 0 | 37 | 36 |
| FP | 2288 | 132 | 165 | 0 | 0 | 46 | 375 | 59 | 1419 | 1514 | 1514 | 127 | 23 | 0 | 37 | 39 |
| ISSM-N1 | 1716 | 132 | 165 | 0 | 0 | 46 | 375 | 59 | 946 | 1230 | 1041 | 127 | 23 | 0 | 37 | 37 |
| ISSM-N2 | 2059 | 132 | 165 | 0 | 0 | 46 | 375 | 59 | 946 | 1230 | 1041 | 127 | 23 | 0 | 37 | 47 |
| ISSM-N3 | 2288 | 158 | 198 | 58 | 6 | 46 | 506 | 79 | 946 | 1230 | 1041 | 127 | 23 | 62 | 37 | 47 |
| ISSM-N4 | 2860 | 185 | 248 | 58 | 6 | 46 | 768 | 129 | 946 | 1230 | 1041 | 127 | 23 | 62 | 37 | 73 |

^aThe carbon emission coefficients were 1.3,0.2,0.15, 6.3, 5.1 and 3.9 C cost (kg C eq. kg⁻¹ active ingredient) per applied nitrogen fertilizer, phosphorus, potassium, herbicide, insecticide and fungicide, respectively, as referred to in Lal (2004). We collected data specific to China's fertilizer manufacture and consumption, and then estimated carbon emissions coefficients were 0.07 and 0.1 C cost (kg C eq. kg⁻¹ active ingredient) per applied Si and Zn fertilizer, respectively.

^bThe carbon emission coefficient for irrigation was 5.16 C cost (kg C eq. cm⁻¹ ha⁻¹) as referred to in Lal (2004).

^cThe carbon emission coefficients were 0.94, 3.2, 0.0075, 0.94 and 0.0725 C cost (kg C eq. kg⁻¹ active ingredient) for tillage and raking, crop planting, per farm manure application, harvesting, spraying and threshing, respectively, as referred to in Lal (2004).

^dSee Table 1 for treatment codes.

Table 5

Mean global warming potential (GWP) and greenhouse gas intensity (GHGI) over the three rice season, wheat season and annual cycles of the 2011rice season–2014wheat season.

| Treatment | CH ₄ | N ₂ O | Ei | Eo | SOCSR | GWP ^a | Grain yield | GHGI ^b |
|---------------------|--|------------------|------|------|------------|------------------|-------------------------------------|--|
| | kg CO ₂ eq. ha ⁻¹ yr ⁻¹ | | | | | | t ha ⁻¹ yr ⁻¹ | kg CO ₂ eq. t ⁻¹ grain |
| Rice season | | | | | | | | |
| NN | 5026±733d | 44±20c | 424 | 1601 | -396±164c | 7492±706d | 5.83±0.04f | 1285±123b |
| FP | 8035±742c | 121±53bc | 1859 | 1603 | 585±198ab | 11032±555c | 8.59±0.25e | 1285±68b |
| ISSM-N1 | 7132±716c | 75±24c | 1502 | 1191 | 246±218b | 9654±800c | 9.45±0.18d | 1021±81c |
| ISSM-N2 | 7186±434c | 112±49bc | 1716 | 1198 | 355±97ab | 9858±484c | 9.98±0.25c | 989±67c |
| ISSM-N3 | 15005±888b | 208±66ab | 2037 | 1260 | 691±252a | 17818±786b | 10.95±0.13b | 1626±54a |
| ISSM-N4 | 17427±1463a | 284±60a | 2626 | 1280 | 773±174a | 20844±1452a | 12.11±0.28a | 1720±108a |
| Wheat season | | | | | | | | |
| NN | -78±71a | 201±28d | 310 | 104 | -396±164c | 934±214b | 1.75±0.04d | 533±125a |
| FP | 8±148a | 605±99b | 1206 | 105 | 585±198ab | 1339±129b | 5.67±0.16b | 236±21b |
| ISSM-N1 | 16±103a | 351±32c | 991 | 105 | 246±218b | 1217±342b | 5.04±0.08c | 241±68b |
| ISSM-N2 | 6±22a | 451±49c | 1120 | 108 | 355±97ab | 1329±109b | 5.76±0.22ab | 231±26b |
| ISSM-N3 | 23±131a | 598±20b | 1302 | 108 | 691±252a | 1340±290b | 5.40±0.16bc | 247±48b |
| ISSM-N4 | 79±96a | 772±66a | 1674 | 114 | 773±174a | 1867±175a | 6.14±0.35a | 305±33b |
| Rice-wheat rotation | | | | | | | | |
| NN ^d | 4948±704d ^c | 246±26d | 734 | 1705 | -792±327c | 8425±711d | 7.58±0.04d | 1111±94b |
| FP | 8043±858c | 725±49b | 3065 | 1708 | 1170±396ab | 12371±583c | 14.26±0.36c | 868±29c |
| ISSM-N1 | 7141±709c | 426±55c | 2493 | 1296 | 491±435b | 10871±990c | 14.50±0.14c | 750±68d |
| ISSM-N2 | 7192±424c | 563±86c | 2836 | 1306 | 709±193ab | 11187±552c | 15.74±0.44b | 712±52d |
| ISSM-N3 | 15028±833b | 806±77b | 3339 | 1368 | 1383±503a | 19158±761b | 16.36±0.18b | 1171±37ab |
| ISSM-N4 | 17506±1396a | 1056±58a | 4300 | 1394 | 1545±348a | 22711±1438a | 18.26±0.46a | 1245±93a |

^aGWP (kg CO₂ eq. ha⁻¹yr⁻¹) = 28 × CH₄ + 265 × N₂O + Ei + Eo - 44/12 × SOCSR, Ei (agrochemical inputs), Eo (farm operations), SOCSR (SOC sequestration rate) is divided by 2 to roughly estimate the GWP from rice and wheat season, respectively. All other items were actually measured for each season.

^bGHGI (kg CO₂ eq. t⁻¹ grain) = GWP/grain yields

^cDifferent lower case letters within the same column for each item indicate significant differences at *P*<0.05 based on Tukey's multiple range tests.

^dSee Table 1 for treatment codes.

Fig 1 Rice and wheat agronomic nitrogen use efficiency (NUE) in 2011–2014 in Changshu, China.

Different letters indicate a significant difference between treatments ($p < 0.05$). See Table 1 for treatment codes.

Fig 2 Seasonal variation of methane (CH_4) fluxes from the rice-wheat rotation cropping systems from 2011 to 2014. The black and gray part in figure separates different grain growth periods. See Table 1 for treatment codes. The solid arrows indicate fertilization.

Fig 3 Seasonal variation of nitrous oxide (N_2O) fluxes from rice-wheat rotation cropping systems in three annual cycles over the period 2011–2014. The black and gray part in the figure separates different growth periods. See Table 1 for treatment codes. The solid arrows indicate fertilization.




Fig. 1




Fig. 2




Fig. 3