

Dear Dr. Tina Treude: Thank you very much for your great support! We have corrected our manuscript according to your valuable comments. Please see the point-to-point answers and tracking manuscript. Thank you very much!

**Associate Editor Decision: Publish subject to technical corrections (28 Apr 2016) by Tina Treude**

**Comments to the Author:**

**Dear Zhang et al.**

**it is with pleasure to inform you that your manuscript is now acceptable for publication in Biogeoscience.**

**I have just two technical comments.**

**Number one, I found the second sentence in the abstract (Line 17-23) very long and difficult to read. It could help to either break it up or insert commas or semicolons.**

A: Thank you very much. We have revised those sentences by reorganize them according to your valuable comments. Abstract lines 17-26.

**Number two, we were just notified by our editors in chief that supplementary material is not permanently archived at BG. I therefore would like to encourage you to incorporate your Suppl. Material 1 and 2 into the main manuscript, especially if it is relevant to understand your study.**

A: Thank you for your comment. We have corrected Suppl. Material 1 and 2 to Fig. 1 and Table 2, respectively. Revised accordingly the whole manuscript.

**With kind regards**

**Tina Treude**

**BG Editor**

Thank you very much once again for your great support!

Sincerely yours,

Zhengqin (on behalf of all authors)

~~~~~  
Prof. Zhengqin Xiong, PhD  
College of Resources and Environmental Sciences  
Nanjing Agricultural University  
Weigang #1, Nanjing, 210095 PRC  
zqxiong@njau.edu.cn  
86-13605188915 (cell)  
86-25-84395148 (O)  
ORCID <https://orcid.org/0000-0003-4743-7325>  
~~~~~

1 Global warming potential and greenhouse gas intensity in rice agriculture driven by  
2 high yields and nitrogen use efficiency

3

4 Xiaoxu Zhang<sup>a</sup>, Xin Xu<sup>a</sup>, Yinglie Liu<sup>a</sup>, Jinyang Wang<sup>a,b</sup>, Zhengqin Xiong<sup>a,\*</sup>

5 <sup>a</sup> Jiangsu Key Laboratory of Low Carbon Agriculture and GHGs Mitigation, College of  
6 Resources and Environmental Sciences, Nanjing Agricultural University, Nanjing, 210095,  
7 China

8 <sup>b</sup> State Key Laboratory of Soil and Sustainable Agriculture, Institute of Soil Science, Chinese  
9 Academy of Sciences, Nanjing 210008, China

10

11 \*Corresponding author (Z. Xiong):

12 Tel: +86-25-84395148,

13 Fax: +86-25-84395210,

14 E-mail: zqxiong@njau.edu.cn

15 **Abstract:** Our understanding of how global warming potential (GWP) and greenhouse gas  
16 intensity (GHGI) is affected by management practices aimed at food security with respect to  
17 rice agriculture remains limited. In the present study, a field experiment was conducted in  
18 China to evaluate the effects of integrated soil-crop system management (ISSM) ~~mainly~~  
19 ~~consisting of different nitrogen (N) fertilization rates and split, manure, Zn and Na<sub>2</sub>SiO<sub>3</sub>~~  
20 ~~fertilization and planting density~~ on GWP and GHGI after accounting for carbon dioxide (CO<sub>2</sub>)  
21 equivalent emissions from all sources, including methane (CH<sub>4</sub>) and nitrous oxide (N<sub>2</sub>O)  
22 emissions, agrochemical inputs and farm operations and sinks (i.e., soil organic carbon  
23 sequestration). The ISSM was mainly consisted of different nitrogen (N) fertilization rates  
24 and split, manure, Zn and Na<sub>2</sub>SiO<sub>3</sub> fertilization and planting density For the improvement of  
25 rice yield and agronomic nitrogen use efficiency (NUE), ~~four~~ four ISSM scenarios consisting of  
26 different chemical N rates relative to the local farmers' practice (FP) rate were carried out,  
27 namely, ISSM-N1 (25% reduction), ISSM-N2 (10% reduction), ISSM-N3 (FP rate) and  
28 ISSM-N4 (25% increase). The results showed that compared with the FP, the four ISSM  
29 scenarios significantly increased the rice yields by 10, 16, 28 and 41% and the agronomic NUE  
30 by 75, 67, 35 and 40%, respectively. In addition, compared with the FP, the ISSM-N1 and  
31 ISSM-N2 scenarios significantly reduced the GHGI by 14 and 18%, respectively, despite  
32 similar GWPs. The ISSM-N3 and ISSM-N4 scenarios remarkably increased the GWP and  
33 GHGI by an average of 69 and 39%, respectively. In conclusion, the ISSM strategies are  
34 promising for both food security and environmental protection, and the ISSM scenario of  
35 ISSM-N2 is the optimal strategy to realize high yields and high NUE together with low  
36 environmental impacts for this agricultural rice field.

37

## 38 **1 Introduction**

39 Rapid population growth and economic development place a growing pressure on increasing  
40 food production (Barrett, 2010). An increase in global crop production of 100% would be  
41 necessary to sustain the projected demand for human food and livestock feed in 2050 (Tilman  
42 et al., 2011). Rice is the staple food for nearly 50% of the world's people, mainly in Asia  
43 (Frolking et al., 2002). According to FAO (2010), approximately 600 million people in  
44 Asia-Pacific region are suffering from hunger and malnutrition. With the region's population  
45 projected to increase by another billion by mid-century, new approaches to increase food  
46 production are needed (Chen et al., 2014). With a limited agricultural land area, the intensive  
47 agricultural regions of China are facing serious environmental problems due to large inputs of  
48 chemical fertilizer and low nitrogen use efficiency (NUE) (Ju et al., 2009; Makino, 2011).  
49 Thus, integrated soil-crop system management (ISSM), which redesigns the whole production  
50 system based on the local environment and draws on appropriate fertilizer compounds and  
51 application ratios, crop densities and advanced water management regimes, has been  
52 advocated and developed to simultaneously increase crop productivity and NUE with low  
53 carbon dioxide (CO<sub>2</sub>) equivalent emissions per unit product in China (Chen et al., 2014). The  
54 key points of the ISSM are to integrate soil and nutrient management with high-yielding  
55 cultivation systems, to integrate the utilization of various nutrient sources and match nutrient  
56 supply to crop requirements, and to take all soil quality improvement measures into  
57 consideration (Zhang et al., 2011).

58 Carbon dioxide, methane (CH<sub>4</sub>) and nitrous oxide (N<sub>2</sub>O) are the most important  
59 greenhouse gases (GHGs) that contribute to global warming (IPCC, 2013). The concept of  
60 global warming potential (GWP) has been applied to agricultural lands by taking into account  
61 of the radiative properties of all GHG emissions associated with agricultural production and  
62 soil organic carbon (SOC) sequestration, expressed as CO<sub>2</sub> eq. ha<sup>-1</sup> yr<sup>-1</sup> (Robertson and Grace,  
63 2004; Mosier et al., 2006). Although agriculture releases significant amounts of CH<sub>4</sub> and N<sub>2</sub>O  
64 into the atmosphere, the net emission of CO<sub>2</sub> equivalents from farming activities can be partly  
65 offset by changing agricultural management to increase the soil organic matter content and/or  
66 decrease the emissions of CH<sub>4</sub> and N<sub>2</sub>O (Mosier et al., 2006; Smith et al., 2008). If global  
67 agricultural techniques are improved, the mitigation potential of agriculture (excluding fossil

68 fuel offsets from biomass) is estimated to be approximately 5.5–6.0 Pg CO<sub>2</sub> eq. yr<sup>-1</sup> by 2030  
69 (Smith et al., 2008). However, the release of CO<sub>2</sub> during the manufacturing and application of  
70 N fertilizer to crops and from fuel used in machines for farm operations can counteract these  
71 mitigation efforts (West and Marland, 2002). Therefore, when determining the GWP of  
72 agroecosystems, there is a need to account for all sources of GHG emissions, including the  
73 emissions associated with agrochemical inputs (E<sub>i</sub>) and farm operations (E<sub>o</sub>) and sinks, e.g.  
74 soil organic carbon (SOC) sequestration (Sainju et al., 2014).

75 Information on the effects of ISSM scenarios on GWP and greenhouse gas intensity  
76 (GHGI) of agricultural systems is limited in China (Ma et al., 2013; Liu et al., 2015). The  
77 annual rotation of summer rice-upland crop is a dominant cropping system in China. Previous  
78 studies were mainly focused on the initial influences of ISSM practices on CH<sub>4</sub> and N<sub>2</sub>O  
79 emissions, but did not account for the contributions of CO<sub>2</sub> emissions from E<sub>i</sub> and E<sub>o</sub> (Ma et  
80 al., 2013; Zhang et al., 2014). In this study, we evaluated GWP and GHGI of rice-wheat crop  
81 rotation managed under several scenarios of ISSM by taking CO<sub>2</sub> equivalents emissions from  
82 all sources and sinks into account for 3 years. We hypothesized that the ISSM strategies  
83 would reduce the overall GWP and GHGI compared with local farmers' practices (FP). The  
84 specific objectives of this study were to (i) evaluate the effects of different ISSM scenarios on  
85 GWP and GHGI; (ii) determine the main sources of GWP and GHGI in a rice-wheat cropping  
86 system; and (iii) elucidate the overall performance for each ISSM scenario for different  
87 targets to increase grain yields and NUE and reduce GWP and GHGI.

## 88 **2 Materials and Methods**

### 89 2.1 Experimental site

90 A field experiment was conducted at the Changshu agro-ecological experimental station  
91 (31°32'93"N, 120°41'88"E) in Jiangsu Province, China. This is a typical, intensively managed  
92 agricultural area where the cropping regime is dominated by a flooded rice (*Oryza sativa*  
93 L.)-drained wheat (*Triticum aestivum* L.) rotation system. The site is characterized by a  
94 subtropical humid monsoon climate with a mean annual air temperature of 15.6, 15.2 and  
95 15.8 °C and precipitation of 878, 1163 and 984 mm for three years, respectively. The soil of  
96 the field is classified as an *Anthrosol* with a sandy loam texture of 6% sand (1–0.05 mm), 80%  
97 silt (0.05–0.001 mm), and 14% clay (<0.001 mm), which developed from lacustrine sediment.

98 The major properties of the soil at 0–20 cm can be described as follows: bulk density, 1.11 g  
99  $\text{cm}^{-3}$ ; pH, 7.35; organic matter content, 35.0  $\text{g kg}^{-1}$ ; and total N, 2.1  $\text{g kg}^{-1}$ . The daily mean air  
100 temperatures and precipitation during the study period from June 15, 2011, to June 15, 2014,  
101 are given in the [Fig. supplementary resource 1](#).

## 102 2.2 Experimental design and management

103 A completely randomized block design was established in 2009 with four replicates of six  
104 treatments, including no nitrogen (NN) and FP as controls, and four ISSM scenarios at  
105 different chemical N fertilizer application rates relative to the local FP rate (300  $\text{kg N ha}^{-1}$ ),  
106 namely ISSM-N1 (25% reduction), ISSM-N2 (10% reduction), ISSM-N3 (FP rate) and  
107 ISSM-N4 (25% increase). The designed ISSM scenarios (only for rice but not for wheat)  
108 included a redesigned split N fertilizer application, a balanced fertilizer application that  
109 included sodium silicate, zinc sulphate, rapeseed cake (C/N=8) providing an additional 112.5  
110  $\text{kg N ha}^{-1}$ , and additional phosphorus and potassium, and different transplanting densities,  
111 used as the main techniques for improving rice yield and agronomic NUE. The agronomic  
112 NUE was calculated as the difference in grain yield between the plots that received N  
113 application and the NN plot, divided by the total N rate which included chemical N fertilizer  
114 and N supplied from rapeseed cake in the ISSM-N3 and ISSM-N4 scenarios. The details of  
115 the fertilizer applications, irrigation, and field management practices of the six different  
116 treatments are presented in Table 1. Further information was described previously (Zhang et  
117 al., 2014). Each plot was 6 m  $\times$  7 m in size with an independent drainage/irrigation system.

118 One midseason drainage (about one week) and final drainage before harvest were used  
119 during the rice-growing season, whereas the plots only received precipitation during the  
120 wheat-growing season. The N fertilizer was split into a 6:2:0:2 or 5:1:2:2 ratio of basal  
121 fertilizer and topdressings for the rice crop and a 6:1:3 ratio for the wheat crop. All of  
122 phosphorous (P), silicon (Si), zinc (Zn) were applied as basal fertilizers for both crops and  
123 rapeseed cake manure was applied for the rice crop. Potassium (K) was added as a split (1:1)  
124 application to the rice crop and all as basal fertilizer for the wheat crop. The basal fertilization  
125 occurred at the time of rice transplanting and wheat seeding. The topdressing was applied at  
126 the tillering, elongation and panicle stages of the rice crop and at the seedling establishment  
127 and elongation stages of the wheat crop. Aboveground biomass including crop grains and

128 straws were removed out of the fields for all the treatments.

### 129 2.3 Gas sampling and measurements

130 We measured the CH<sub>4</sub> and N<sub>2</sub>O emissions from each plot of the field experiment over five  
131 annual cycles from the 2009 rice-growing season to the 2014 wheat-growing season. The  
132 initial 2-yr measurements during the 2009–2011 rice-wheat rotational systems were described  
133 in our previous study (Ma et al., 2013). Emissions were measured manually using the  
134 static-opaque chamber method. Each replicate plot was equipped with a chamber with a size  
135 of 50 cm × 50 cm × 50 cm or 50 cm × 50 cm × 110 cm, depending on the crop growth and  
136 plant height. The chamber was placed on a fixed PVC frame in each plot and wrapped with a  
137 layer of sponge and aluminum foil to minimize air temperature changes inside the chamber  
138 during the period of sampling. Gas samples were collected from 9:00 to 11:00 am using an  
139 airtight syringe with a 20-ml volume at intervals of 10 min (0, 10, 20 and 30 min after  
140 chamber closure). The fluxes were measured once a week and more frequently after fertilizer  
141 application or a change in soil moisture.

142 The gas samples were analyzed for CH<sub>4</sub> and N<sub>2</sub>O concentrations using a gas  
143 chromatograph (Agilent 7890A, Shanghai, China) equipped with two detectors. Methane was  
144 detected using a hydrogen flame ionization detector (FID), and N<sub>2</sub>O was detected using an  
145 electron capture detector (ECD). Argon-methane (5%) and N<sub>2</sub> were used as the carrier gas at a  
146 flow rate of 40 ml min<sup>-1</sup> for N<sub>2</sub>O and CH<sub>4</sub> analysis, respectively. The temperatures for the  
147 column and ECD detector were maintained at 40 °C and 300 °C, respectively. The oven and  
148 FID were operated at 50 °C and 300 °C, respectively. The CH<sub>4</sub> and N<sub>2</sub>O fluxes were  
149 calculated using a linear increase in the two gas concentrations over time as described by Jia  
150 et al. (2012).

### 151 2.4 Topsoil organic carbon sequestration measurements

152 To measure the organic carbon content of the topsoil as described by Zhang et al. (2014), soil  
153 samples were collected after the wheat harvest in 2009 and 2014 from all experimental plots  
154 at a plowing depth of 0–20 cm. The soil organic carbon sequestration rates (SOCSR) were  
155 calculated as follows (Liu et al., 2015):

$$156 \text{ SOCSR (t C ha}^{-1} \text{ yr}^{-1}) = (\text{SOC}_t - \text{SOC}_0) / T \times \gamma \times (1 - \delta_{2\text{mm}}/100) \times 20 \times 10^{-1} \text{ (1)}$$

157 In Eq. (1), SOC<sub>t</sub> (g C kg<sup>-1</sup>) and SOC<sub>0</sub> (g C kg<sup>-1</sup>) are the SOC contents measured in the

158 soils sampled after the wheat was harvested in 2014 and 2009, respectively. T refers to the  
159 experimental period (yr).  $\gamma$  and  $\delta_{2\text{mm}}$  are the average bulk density and the gravel content (>2  
160 mm) of the topsoil (0–20 cm), respectively.

## 161 2.5 GWP and GHGI measurements

162 To better understand the overall GHG impact of the rice-wheat crop rotation managed  
163 under different ISSM scenarios, the GWP and GHGI were calculated as the following  
164 equations (Myhre et al., 2013):

$$165 \quad \text{GWP (kg CO}_2 \text{ eq. ha}^{-1} \text{ yr}^{-1}) = 28 \times \text{CH}_4 + 265 \times \text{N}_2\text{O} + \text{Ei} + \text{Eo} - 44/12 \times \text{SOCSR} \quad (2)$$

$$166 \quad \text{GHGI (kg CO}_2 \text{ eq. kg}^{-1} \text{ grain yield yr}^{-1}) = \text{GWP}/\text{grain yield} \quad (3)$$

167 In Eq. (2), Ei (kg CO<sub>2</sub> eq. ha<sup>-1</sup> yr<sup>-1</sup>), Eo (kg CO<sub>2</sub> eq. ha<sup>-1</sup> yr<sup>-1</sup>) and SOCSR (kg C ha<sup>-1</sup>  
168 yr<sup>-1</sup>) represent CO<sub>2</sub> equivalent emissions from the agrochemical inputs, farm operations and  
169 soil organic carbon sequestration rate, respectively. The global warming potential of 1 kg CH<sub>4</sub>  
170 and 1 kg N<sub>2</sub>O are 28 and 265 kg CO<sub>2</sub> equivalents respectively (without inclusion of  
171 climate-carbon feedbacks), based on 100-yr time scale (Myhre et al., 2013). 12 and 44 refers  
172 to molecular weights of C and CO<sub>2</sub>, respectively. The grain yield is expressed as the air-dried  
173 grain yield.

174 Therefore, the GWP of the cropland ecosystem equals the total CO<sub>2</sub> equivalent emissions  
175 minus the SOC change per unit land area. In addition to CH<sub>4</sub> and N<sub>2</sub>O emissions, we  
176 considered CO<sub>2</sub> equivalent emissions associated with the use of agrochemical inputs (Ei),  
177 such as the manufacture and transportation of the N, P, and K fertilizers (Snyder et al., 2009),  
178 and farm operations (Eo), such as the water used for irrigation (Zhang et al., 2013) and diesel  
179 fuel (Huang et al., 2013a). The CO<sub>2</sub> equivalent emissions of N fertilizer were calculated as the  
180 mean value of the C emissions of 1.3 kg C equivalent kg<sup>-1</sup> N (Lal, 2004). Similarly, the CO<sub>2</sub>  
181 equivalent for irrigation was calculated from the total amount of water used during the  
182 rice-growing season; the coefficient for the C cost was 5.16 (kg C eq. cm<sup>-1</sup> ha<sup>-1</sup>) originated  
183 from the value of 257.8 kg C eq. ha<sup>-1</sup> for a 50 cm of irrigation provided by Lal (2004). The  
184 CO<sub>2</sub> equivalents of other Ei (P and K fertilization, manure, herbicide, pesticide and fungicide  
185 applications) and Eo (tillage, planting, harvest, and threshing) were recorded and also  
186 estimated by coefficients provided by Lal (2004) since no specific coefficients were available  
187 for local conditions. We collected the data specific to China's fertilizer manufacture and

188 consumption, and obtained the C emission coefficients to be 0.07 and 0.1 kg C eq. kg<sup>-1</sup> of  
189 active ingredient for Si and Zn fertilizer, respectively. The C emission factor for these farm  
190 operations depends on diesel used as fuel or electricity. Chemical fertilizer was hand  
191 broadcasted for each fertilization event. Detailed information of each Ei and Eo component  
192 for rice and wheat crop seasons are presented in [TableSupplementary resource 2](#).

## 193 2.6 Statistical analysis

194 Repeated-measures multivariate analysis of variance (MANOVA) and linear relationships  
195 were determined using JMP 7.0, ver. 7.0 (SAS Institute, USA, 2007). The F-test was applied  
196 to determine whether there were significant differences among practices, years and their  
197 interaction at  $P < 0.05$ . One-way analysis of variance was conducted to compare the  
198 cumulative fluxes of CH<sub>4</sub> and N<sub>2</sub>O, and grain yield among the different treatments. Tukey's  
199 HSD test was used to determine whether significant differences occurred between the  
200 treatments at a level of  $P < 0.05$ . Normal distribution and variance uniformity were checked  
201 and all data were consistent with the variance uniformity ( $P > 0.05$ ) within each group. The  
202 results are presented as the means and standard deviation (mean  $\pm$  SD, n = 4).

## 203 3 Results

### 204 3.1 Crop production and agronomic NUE

205 During the three cropping rotations from 2011 to 2014, the rice and wheat yields varied  
206 significantly among the treatments (Table [32](#)). The grain yields ranged from 5.83 to 12.11 t  
207 ha<sup>-1</sup> for rice and 1.75 to 6.14 t ha<sup>-1</sup> for wheat. On average over the three cycles, the annual  
208 rice yield of the FP was significantly lower than that of the ISSM scenarios of ISSM-N1,  
209 ISSM-N2, ISSM-N3 and ISSM-N4. Compared with the FP, rice grain yields increased by 10%  
210 and 16% for the ISSM-N1 and ISSM-N2 scenarios, respectively, i.e., with the lower N input,  
211 by 28% for the ISSM-N3 scenario with the same N input and by 41% for the ISSM-N4  
212 scenario with the highest N input. However, we did not observe any significant increases in  
213 the wheat-grain yields compared with the FP except for the ISSM-N4 scenario. Statistical  
214 analysis indicated that rice and wheat yields from the three years were not significantly  
215 influenced by the interaction of cultivation patterns and cropping year (Table [43](#)).

216 The agronomic NUE for the rice and wheat of the fertilized plots ranged from 9.2 to 16.1  
217 and 19.5 to 24.7 kg grain kg N<sup>-1</sup>, respectively (Fig. [24](#)). The higher NUE in the wheat season

218 was mainly due to the relatively lower N fertilizer (40%) rates used for wheat compared with  
219 that for rice. As expected, the rice agronomic NUE significantly increased by 75, 67, 35 and  
220 40% for the ISSM-N1, ISSM-N2, ISSM-N3 and ISSM-N4 scenarios, respectively, compared  
221 with the FP (Fig. 24). For the wheat crop, the agronomic NUE increased by 12 and 14% in the  
222 ISSM-N1 and ISSM-N2 scenarios, respectively, and slightly decreased in the ISSM-N3 and  
223 ISSM-N4 scenarios compared with the FP, mainly because the current ISSM strategy was  
224 only designed for rice and not wheat production.

### 225 3.2 CH<sub>4</sub> and N<sub>2</sub>O emissions

226 All plots showed similar CH<sub>4</sub> emission patterns, being a source in the rice season and  
227 negligible in the wheat season (Fig. 32). During the three annual rice-wheat rotations from  
228 2011 to 2014, the CH<sub>4</sub> fluxes ranged from -3.89 to 99.67 mg C m<sup>-2</sup> h<sup>-1</sup>. The seasonal CH<sub>4</sub>  
229 emissions varied significantly among the treatments during the rice-growing season (Table 43,  
230 Fig. 32). No significant difference was found between the FP, ISSM-N1 and ISSM-N2 plots.  
231 Temporal variation was significant during the three cycles (Table 43, *P* < 0.001). Averaged  
232 across years, the CH<sub>4</sub> emission was greater in the ISSM-N3 and ISSM-N4 plots than in the  
233 NN, FP, ISSM-N1 and ISSM-N2 plots (Table 32, *P* < 0.05). However, compared with the NN  
234 plots, the FP, ISSM-N1 and ISSM-N2 plots with inorganic fertilizer application resulted in  
235 increased CH<sub>4</sub> emission rates of 59.9, 41.9 and 43.0%, respectively, averaged over the  
236 rice-growing seasons. The CH<sub>4</sub> emission rates were further enhanced by 198.5% in the  
237 ISSM-N3 plots and by 246.7% in the ISSM-N4 plots.

238 The annual N<sub>2</sub>O fluxes varied from -33.1 to 647.5 μg N<sub>2</sub>O-N m<sup>-2</sup> h<sup>-1</sup>, with most N<sub>2</sub>O  
239 emissions occurring during the wheat-growing season after fertilization events, and several  
240 small emission peaks during the rice-growing season (Fig. 43). With respect to the N  
241 application effect, the annual cumulative N<sub>2</sub>O emissions for all four ISSM scenarios were  
242 significantly higher than that in NN (*P* < 0.05). Relative to the FP plot, the ISSM-N1 and  
243 ISSM-N2 scenarios decreased the annual N<sub>2</sub>O emissions by an average of 41% and 22%,  
244 respectively (Table 32). The ISSM-N4 scenario significantly increased the cumulative N<sub>2</sub>O  
245 emissions by 46% (*P* < 0.05) because this system received highest inorganic N fertilizer (25%  
246 higher than that in FP) and additional N via manure application compared to the FP practice,  
247 although there was no significant difference between the ISSM-N3 and FP plots.

### 248 3.3 Annual GWP and GHGI

249 Based on the perspective of the carbon footprint, we included the GHG emissions associated  
250 with all of the inputs (Ei and Eo), and SOC sequestration was expressed as kg CO<sub>2</sub> eq. ha<sup>-1</sup>  
251 yr<sup>-1</sup>. The CO<sub>2</sub> equivalent emissions associated with Ei and Eo are presented in Table 54. The  
252 CO<sub>2</sub> equivalents rates from N fertilizer dominated not only the chemical input section (67–75%  
253 of Ei) but also the total CO<sub>2</sub> equivalents from agricultural management (45–50% of the sum  
254 of the Ei and Eo). Irrigation was the second largest source of CO<sub>2</sub> equivalents associated with  
255 agricultural management after N fertilizer (19–31% of the sum of the Ei and Eo). The GWP  
256 ranged from 8425 to 22711 kg CO<sub>2</sub> eq. ha<sup>-1</sup> yr<sup>-1</sup> for the NN and the ISSM-N4 plots,  
257 respectively (Table 65). Although fertilized treatments increased the annual CH<sub>4</sub> and N<sub>2</sub>O  
258 emissions in comparison with the NN plot, it also increased the SOC sequestration in these  
259 cropping systems. Of the main field GHGs that were directly emitted, CH<sub>4</sub> accounted for  
260 59–78% of the GWP in all plots. An increase in the annual SOC content led to a significant  
261 decrease in the GWP (contributed 5–9% decrease of the GWP except in the NN plot). The  
262 CO<sub>2</sub> equivalents from agricultural management practices, emissions associated with Ei  
263 (2493–4300 CO<sub>2</sub> eq. ha<sup>-1</sup> yr<sup>-1</sup>) were higher than those associated with Eo (1296–1708 CO<sub>2</sub> eq.  
264 ha<sup>-1</sup> yr<sup>-1</sup>) in the fertilized plots. There was no significant difference in the annual GWP  
265 observed between the FP, ISSM-N1 and ISSM-N2 plots (Table 65). Across the three years,  
266 ISSM-N1 and ISSM-N2 slightly reduced the GWP by 12 and 10%, respectively; however,  
267 ISSM-N3 and ISSM-N4 significantly increased the GWP by an average of 55 and 84%,  
268 respectively, in comparison with the FP.

269 The GHGI was used to express the relationship between GWP and grain yield. The  
270 GHGIs (kg CO<sub>2</sub> eq. t<sup>-1</sup> grain) in this study ranged from 712 to 1245 kg CO<sub>2</sub> eq. t<sup>-1</sup> grain  
271 (Table 65). The significant difference in the GHGI of grain was found between the FP and the  
272 ISSM strategies. Compared with the FP, ISSM-N1 and ISSM-N2 significantly reduced the  
273 GHGI by 14 and 18%, respectively, mainly due to the increased grain yield and SOC  
274 sequestration as well as reduced GHG emissions for the ISSM strategies of reasonable N  
275 fertilizer management and suitable planting density. Although N fertilizer or  
276 organic/inorganic combination fertilizer application reduced the SOC losses caused by crop  
277 cultivation and increased the grain yields, the GHGIs were generally higher for the ISSM-N3

278 and ISSM-N4 scenarios than the ISSM-N1 and ISSM-N2 scenarios due to further increases in  
279 CH<sub>4</sub> and N<sub>2</sub>O emissions.

## 280 **4 Discussion**

### 281 4.1 Grain yield and agronomic NUE as affected by ISSM strategies

282 Grain yields are directly related to fertilizer management. The MANOVA results indicated  
283 that the rice and wheat grain yields were significantly affected by the cultivation strategies  
284 (Table 43,  $P < 0.001$ ), which is in agreement with previous results (Chen et al., 2011; Zhang  
285 et al., 2011). Compared with the FP, rice yields increased significantly by all four ISSM  
286 scenarios (Table 32). However, the wheat grain yield decreased significantly when the N  
287 fertilizer rate was reduced by 25% (N1 scenario). It has been reported in previous studies that  
288 ISSM strategies can effectively improve the rice grain yield (Ma et al., 2013; Liu et al., 2015).  
289 First, the adjusted transplanting density for the ISSM-N1, ISSM-N2, and ISSM-N3 scenarios  
290 would produce a positive effect on rice yield by influencing rice colony structure, which  
291 agreed with Wu et al. (2005). Second, split application of N fertilizer to match crop demand in  
292 the ISSM-N1, ISSM-N2, ISSM-N3 and ISSM-N4 scenarios would significantly increase  
293 agronomic NUE and rice yield which had been reported previously by Liu et al. (2009). In the  
294 present study, ISSM-N1 and ISSM-N2 significantly increased annual rice production by 10  
295 and 16%, respectively, in comparison with the FP (Table 32). This finding is consistent with  
296 the results of Peng et al. (2006), who reported that a 30% reduction in the total N rate during  
297 the early vegetative stage did not reduce the yield but slightly increased it when combined  
298 with the modified farmers' fertilizer practice. Third, integrated management of three  
299 macronutrients: N, P and K as well as the two micronutrients: Si and Zn were considered as  
300 essential for sustainable high crop yields. Additional Si and Zn fertilizer for the ISSM-N3 and  
301 ISSM-N4 scenarios would support better seedling establishment and reduce both biotic and  
302 abiotic stress, thus produced higher yields (Wang et al., 2005; Slaton et al., 2005;  
303 Kabata-Pendias and Mukherjee, 2007; Hossain et al., 2008). As expected, when the total N  
304 rate was at the FP rate and/or increased by 25%, in combination with other ISSM strategies  
305 (e.g. rapeseed cake manure, additional P and K, applying Si and Zn fertilizer), the rice yield in  
306 these ISSM-N3 and ISSM-N4 plots increased substantially by 28 and 41%, respectively.

307 Based on a long-term fertilizer experiment, Shang et al. (2011) reported that organic fertilizer  
308 incorporation significantly increased the early rice grain yield. This may have resulted from  
309 the organic fertilizer applied in combination with adequate nutrients contributing to alleviate  
310 potential yield limiting factors of rice.

311 It has been suggested that N losses vary depending on the timing, rate, and method of N  
312 application, as well as the source of N fertilizer (Zhu, 1997). In addition to high rates of N and  
313 improper timing of N application, rapid N losses (via ammonia volatilization, denitrification,  
314 surface runoff, and leaching) are important factors that cause low agronomic NUE of irrigated  
315 rice in China (Peng et al., 2006). Compared with the FP plot, the rice agronomic NUE was  
316 significantly increased by 75, 67, 35 and 40% under the ISSM-N1, ISSM-N2, ISSM-N3 and  
317 ISSM-N4 scenarios, respectively (Fig. 24). The higher rice agronomic NUE in our study over  
318 the experimental period could be due to reduced N losses by leaching and volatilization as  
319 well as the improvement of N bioavailability in the rice crop season (Zhao et al., 2015).  
320 Organic/inorganic combination fertilizer application also increases uptake by crops compared  
321 with the traditional farmers' practice (Peng et al., 2006). These findings suggest that the ISSM  
322 strategy is an effective method for improving grain yield and agronomic NUE for future  
323 sustainable rice agriculture in China.

#### 324 4.2 CH<sub>4</sub> and N<sub>2</sub>O emissions as affected by ISSM strategies

325 During the three years, the annual cumulative CH<sub>4</sub> emissions, on average, varied from 133 to  
326 469 kg C ha<sup>-1</sup>yr<sup>-1</sup> (Table 32), and these values fell within the range of 4.1 to 1015.6 kg CH<sub>4</sub>  
327 ha<sup>-1</sup> observed previously in a rice field (Huang et al., 2004). Methane emissions were highest  
328 during rice season, but only during the flooding period. Mainly because CH<sub>4</sub> was produced in  
329 the anaerobic zones of submerged soils by methanogens and is oxidized into CO<sub>2</sub> by  
330 methanotrophs in the aerobic zones of wetland soils and in upland soils (Le Mer and Roger,  
331 2001). The MANOVA results indicated that obvious effects of cultivation patterns and  
332 years on CH<sub>4</sub> emissions were found during the rice-wheat rotations (Table 43,  $P < 0.001$ ). The  
333 CH<sub>4</sub> emissions were not significantly affected by the cycles but affected by crop season  
334 (Table 65, Fig. 32). In this study, no significant difference in CH<sub>4</sub> emission was observed  
335 between the FP, ISSM-N1 and ISSM-N2 plots. However, compared with the FP plot, the  
336 ISSM-N3 and ISSM-N4 scenarios emitted 87 and 118% more CH<sub>4</sub>, respectively (Table 65),

337 which is probably due to the incorporation of the organic rapeseed cake manure. Previous  
338 reports support the observations that CH<sub>4</sub> emissions were significantly increased with the  
339 application of organic amendments (Ma et al., 2009; Thangarajan et al., 2013; Zou et al.,  
340 2005). Apparently, additional application of Si and Zn fertilizers had no significant effect on  
341 CH<sub>4</sub> and N<sub>2</sub>O fluxes, which was consistent with the result of Xie et al. (2015). Moreover, rice  
342 growth was found to be significantly increased under the ISSM-N3 and ISSM-N4 scenarios.  
343 In this case, the organic matter inputs such as root litter and rhizodeposits in the ISSM-N3 and  
344 ISSM-N4 scenarios were probably also higher than in the other plots, and thus soil C input,  
345 which served as an additional source of substrates for the methanogens in the rice paddies,  
346 likely contributed to the increase in CH<sub>4</sub> emissions (Ma et al., 2009). Finally, because the rice  
347 plants acted as the main pathway for CH<sub>4</sub> transports from the soil to the atmosphere, the  
348 higher biomass may have facilitated more CH<sub>4</sub> emissions (Yan et al., 2005).

349 Denitrification and nitrification are the main processes that produce N<sub>2</sub>O in the soil (Paul  
350 et al., 1993). The N<sub>2</sub>O emission patterns varied during the rice and wheat growing seasons  
351 which were partially associated with the anaerobic conditions prevailing in a rice paddy.  
352 Changes in the soil water content strongly influenced the soil N<sub>2</sub>O emissions and resulted in  
353 negligible N<sub>2</sub>O emissions when the rice field was flooded (Fig. 43), which is consistent with  
354 previous reports (Akiyama et al., 2005; Murdiyarso et al., 2010). When the soil water content  
355 was below saturation, N<sub>2</sub>O emissions increase with soil moisture; however, N<sub>2</sub>O emissions  
356 gradually decreased with the soil saturation condition (Rudaz et al., 1999). A relatively high  
357 N<sub>2</sub>O peak was observed in the first two weeks of the wheat-growing season (Fig. 43),  
358 possibly because soil changes from flooded to drained conditions may have enhanced N<sub>2</sub>O  
359 release (Deng et al., 2012). Alternation of drainage and flooding may induce large amounts of  
360 N<sub>2</sub>O emissions, particularly in fertilized systems; this has commonly been shown in earlier  
361 studies (Wang et al., 2012; Xiong et al., 2007; Zou et al., 2005). The seasonal and annual rates  
362 of N<sub>2</sub>O emissions were significantly affected by the cultivation practices and years (Table 43).  
363 Compared with the FP plot, the ISSM-N2 scenario significantly decreased the seasonal N<sub>2</sub>O  
364 emissions in this study, which may have resulted from a reduction in the N fertilizer rate  
365 (Table 1, Table 32). The total N<sub>2</sub>O emissions decreased by 7–38% and 26–42% in the rice and  
366 wheat seasons, respectively, when the conventional N management (300 kg N ha<sup>-1</sup> for rice

367 and 180 kg N ha<sup>-1</sup> per crop for wheat) changed to optimum N management (225–270 kg N  
368 ha<sup>-1</sup> for rice and 135–162 kg N ha<sup>-1</sup> per crop for wheat). It is likely that more N<sub>2</sub>O was  
369 emitted (Mosier et al., 2006) as a result of the additional N made available to the soil  
370 microbes through N fertilizer application, which also probably contributed increased CH<sub>4</sub>  
371 emissions (Banger et al., 2013). Strategies that can reduce N fertilization rates without  
372 influencing crop yields can inevitably lower GHG emissions (Mosier et al., 2006). Nitrogen  
373 leaching and volatilization are the important components of reactive N releases but not  
374 included in the current GHG budget.

#### 375 4.3 GWP and GHGI as affected by ISSM strategies

376 The GWP in our study (10871–22711 kg CO<sub>2</sub> eq. ha<sup>-1</sup>) with the ISSM strategies was higher  
377 than that in a double-cropping cereal rotation (1346–4684 kg CO<sub>2</sub> eq. ha<sup>-1</sup>) and a rice-wheat  
378 annual rotation (290–4580 kg CO<sub>2</sub> eq. ha<sup>-1</sup>) reported by Huang et al. (2013b) and Yang et al.  
379 (2015), respectively. Dominant CH<sub>4</sub> emissions as well as additional CO<sub>2</sub> emissions due to the  
380 use of machinery/equipment for irrigation and farm operations under the ISSM strategies may  
381 increase the GWP more than in other cropping systems (emit more CO<sub>2</sub> equivalent emissions  
382 of 2439–5694 kg CO<sub>2</sub> eq. ha<sup>-1</sup> for agricultural management practices in the present study).  
383 However, the current GWP was comparable to that of a double-rice cropping system  
384 (13407–26066 kg CO<sub>2</sub> eq. ha<sup>-1</sup>) (Shang et al., 2011). The GHGIs, which ranged from 0.71 to  
385 1.25 kg CO<sub>2</sub> eq. kg<sup>-1</sup> grain in this study, were slightly higher than previous estimates of  
386 0.24–0.74 kg CO<sub>2</sub> eq. kg<sup>-1</sup> grain from rice paddies with midseason drainage and organic  
387 manure incorporation (Qin et al., 2010; Li et al., 2006) but were lower than the DNDC model  
388 estimates for continuous waterlogged paddies (3.22 kg CO<sub>2</sub> eq. kg<sup>-1</sup> grain) (Li et al., 2006).  
389 Differences in GWP or GHGI were found in the cultivation patterns over the three rice-wheat  
390 | rotations (Table 65). The ISSM-N1 and ISSM-N2 scenarios with optimized ISSM strategies  
391 | led to a lower GWP than the FP by a certain extent, but there were not significant differences  
392 | among the FP, ISSM-N1 and ISSM-N2 plots (Table 65). Compared with the FP, the ISSM-N1  
393 | and ISSM-N2 scenarios significantly reduced the GHGI, which was mainly due to higher  
394 | yields. In spite of the similar GWP compared with the FP plot, the lowest GHGI (0.71 kg CO<sub>2</sub>  
395 | eq. kg<sup>-1</sup> grain) was obtained under the ISSM-N2 scenario. This finding is consistent with the  
396 | suggestion made by Burney et al. (2010), i.e., that the net effect of higher yields offsets

emissions. It is well known that CH<sub>4</sub> emissions dominate the GWP in rice paddies (Ma et al., 2013; Shang et al., 2011). In comparison to the GWP (12371 kg CO<sub>2</sub> eq. ha<sup>-1</sup>yr<sup>-1</sup>) and GHGI (0.87 kg CO<sub>2</sub> eq. kg<sup>-1</sup> grain) of the FP, the ISSM-N3 and ISSM-N4 scenarios increased both the GWP and GHGI, mainly because these scenarios notably increased the CH<sub>4</sub> emissions compared with the FP, which resulted in relatively higher GWP (Table 65).

Agricultural management practices that change one type of GWP source/sink may also impact other sources/sinks and therefore change the GWP and GHGI (Mosier et al., 2006; Shang et al., 2011). Although the N-fertilizer plots, especially those with the incorporation of organic fertilizer, increased the annual CH<sub>4</sub> and N<sub>2</sub>O emissions, they increased the SOC sequestration in this cropping system, which is agreement with previous reports (Huang and Sun, 2006). This may be due to the incorporation of rapeseed cake and enhanced below-ground crop residue associated with higher crop productivity (Ma et al., 2013). In the present study, the ISSM-N2 scenario with ISSM strategies decreased the CH<sub>4</sub> and N<sub>2</sub>O emissions as well as the energy consumption related to irrigation and the manufacture and transport of N fertilizer (depending on coal combustion), ultimately leading to a decrease in the GWP relative to the FP plot. Moreover, despite the lower N fertilizer input, the grain yield did not decline and the GHGI of the ISSM-N2 scenario was thus lower than of the FP plot, indicating less consumption of CO<sub>2</sub> equivalents per unit of grain produced. We demonstrate that high yield and agronomic NUE, together with low GWP, are not conflicting goals by optimizing ISSM strategies.

#### 4.4 Main components of GWP and GHGI and implementation significance for the ISSM strategies

Determining the main components of the GWP and GHGI in specific cropping systems is very important for mitigating GHG emissions in the future, because the benefits of C sequestration would be negated by CH<sub>4</sub> and N<sub>2</sub>O emissions and the CO<sub>2</sub> equivalents released with the use of high N fertilizer application rates (Schlesinger, 2010). In the current study, the five main components of the CO<sub>2</sub> equivalents for the GWP were ranked in decreasing order of importance as follows: CH<sub>4</sub> emissions > agrochemical inputs of N fertilizer > farm operations related to irrigation > SOC sequestration > N<sub>2</sub>O emissions (Table 65). Of the two crops, CH<sub>4</sub> and irrigation were important for rice, but less important for wheat, in which N<sub>2</sub>O losses were

427 | expected to have a higher weight ([Supplementary resource Table 2](#)). Methane emissions, the  
428 | most important component of GWP in this typical rice-wheat rotation system, could be further  
429 | mitigated by some other strategies, such as reasonable irrigation (Zou et al., 2005; Wang et al.,  
430 | 2012).

431 |         Although N fertilizer application increased SOC sequestration when it was applied  
432 | with rapeseed cake manure, this benefit was consistently overshadowed, on a CO<sub>2</sub> equivalent  
433 | basis, by the increases in CH<sub>4</sub> and N<sub>2</sub>O emissions (Table [65](#)). Similar results have been  
434 | reported, i.e., GHG emissions substantially offset SOC increases (Six et al., 2004). It is  
435 | possible that the realization of reducing the GWP and GHGI in China should focus on  
436 | increasing the SOC and simultaneously decreasing the CO<sub>2</sub> equivalents from CH<sub>4</sub> emissions  
437 | and N fertilizer inputs. Several studies reported possible methods for these types of mitigation  
438 | strategies, such as optimizing the chemical fertilizer application amount and rate (Ju et al.,  
439 | 2011), the amount of water used for irrigation (Gao et al., 2015), and the timing and rate of N  
440 | using the in-season N management approach, as well as improving the N fertilizer  
441 | manufacturing technologies (Zhang et al., 2013), and using nitrification inhibitors or  
442 | polymer-coated controlled-release fertilizers (Hu et al., 2013).

443 |         China is a rapidly developing country that faces the dual challenge of substantially  
444 | increasing grain yields and at the same time reducing the substantial environmental impact of  
445 | intensive agriculture (Chen et al., 2011). We used the ISSM strategies to develop a rice  
446 | production system that achieved mean yields of 10.63 t ha<sup>-1</sup> (an increment of almost 24%)  
447 | and an agronomic NUE of 13.20 kg grain kg N<sup>-1</sup> (an increment of 43%) in long-term field  
448 | experiments compared with current farmers' practices. The ISSM redesigned the whole  
449 | production system only for the rice crop based on the local environment and drawing on  
450 | appropriate fertilizer varieties and application ratios, crop densities and an advanced water  
451 | regime management. If the ISSM strategies were also developed for the rotated wheat crop,  
452 | the overall performance of the whole rice-wheat system would be much improved, with  
453 | further increases in yield and reductions in the GWP and GHGI. We conclude that the ISSM  
454 | strategies are promising, particularly the ISSM-N2 scenario, which is the most favorable to  
455 | realize higher yields with lower environmental impact. The proposed ISSM strategies can  
456 | provide substantial benefits to intensive agricultural systems and can be applied feasibly using

457 current technologies.

## 458 **5 Conclusions**

459 Reasonable agricultural management practices are the key to reducing GHG emissions from  
460 agricultural ecosystems. This study provided an insight into the complete GHG emission  
461 accounting of the GWP and GHGI affected by different ISSM scenarios. After a three-year  
462 field experiment, we found that the CH<sub>4</sub> emissions, production of N fertilizer, irrigation, SOC  
463 sequestration and N<sub>2</sub>O fluxes were the main components of the GWP in a typical rice-wheat  
464 rotation system. In contrast with the FP, ISSM-N1 and ISSM-N2 significantly reduced the  
465 GHGI, though they resulted in similar GWPs, and ISSM-N3 and ISSM-N4 remarkably  
466 increased the GWP and GHGI. By adopting the ISSM-N2 strategy, the conventional N  
467 application rate was reduced by 10% while the rice yield was significantly increased by 16%,  
468 the NUE was improved by 67% and the GHGI was lowered by 23%. ISSM scenarios could be  
469 adopted for both food security and environmental protection with specific targets. We propose  
470 that the ISSM-N2 scenario is the most appropriate management strategy (10% reduction of N  
471 input, no rapeseed manure and higher plant density) for realizing higher yields and NUE,  
472 together with some potential to reduce GHGI by integrated soil-crop management. For  
473 simultaneously mitigating GHG emissions, further research on integrated soil-crop system  
474 managements is required particularly for mitigating CH<sub>4</sub> emissions in sustainable rice  
475 agriculture.

476 **Acknowledgments** We sincerely appreciate two anonymous reviewers for their critical and  
477 valuable comments to help improve this manuscript. This work was jointly supported by the  
478 National Science Foundation of China (41171238, 41471192), the Special Fund for  
479 Agro-Scientific Research in the Public Interest (201503106) and the Ministry of Science and  
480 Technology (2013BAD11B01).

481 **References**

- 482 Akiyama, H., Yagi, K. and Yan, X. Y.: Direct N<sub>2</sub>O emissions from rice paddy fields: summary of  
 483 available data, *Global Biogeochem. Cy.*, 19, GB002378, doi:10.1029/2004GB002378, 2005.
- 484 Banger, K., Tian, H. and Lu, C.: Do nitrogen fertilizers stimulate or inhibit methane emissions from  
 485 rice fields?, *Glob. Change Biol.*, 18, 3259-3267, 2013.
- 486 Barrett, C. B.: Measuring food insecurity, *Science*, 327 (5967), 825-828, 2010.
- 487 Burney, J. A., Davis, S. J. and Lobell, D. B.: Greenhouse gas mitigation by agricultural intensification,  
 488 *Proc. Natl. Acad. Sci. U.S.A.*, 107, 12052-12057, 2010.
- 489 Chen, X., Cui, Z., Vitousek, P. M., Cassman, K. G., Matson, P. A., Bai, J., Meng, Q., Hou, P., Yue, S.  
 490 and Römheld, V.: Integrated soil-crop system management for food security, *P. Natl. Acad. Sci.*  
 491 *USA.*, 108, 6399-6404, 2011.
- 492 Chen, X., Cui, Z., Fan, M., Vitousek, P., Zhao, M., Ma, W., Wang, Z., Zhang, W., Yan, X. and Yang, J.:  
 493 Producing more grain with lower environmental costs, *Nature*, 514, 486-489, 2014.
- 494 Deng, J., Zhou, Z., Zheng, X., Liu, C., Yao, Z., Xie, B., Cui, F., Han, S. and Zhu, J.: Annual emissions  
 495 of nitrous oxide and nitric oxide from rice-wheat rotation and vegetable fields: a case study in  
 496 the Tai-Lake region, China, *Plant Soil*, <http://dx.doi.org/10.1007/s11104-012-1223-6>, 2012.
- 497 Food and Agriculture Organization Statistical Data (FAOSTAT), Available online at:  
 498 <http://www.fao.org/docrep/003/x6905e/x6905e04.htm>, Rome, Italy, 2010.
- 499 Frolking, S., Qiu, J., Boles, S., Xiao, X., Liu, J., Zhuang, Y., Li, C. and Qin, X.: Combining remote  
 500 sensing and ground census data to develop new maps of the distribution of rice agriculture in  
 501 China, *Global Biogeochem. Cy.*, doi:16, 10.1029/2001GB001425, 2002.
- 502 Gao, B., Ju, X., Meng, Q., Cui, Z., Christie, P., Chen, X. and Zhang, F.: The impact of alternative  
 503 cropping systems on global warming potential, grain yield and groundwater use, *Agr. Ecosyst.*  
 504 *Environ.*, 203, 46-54, 2015.
- 505 Hossain, M. A., Jahiruddin, M., Islam, M. R., and Mian, M. H.: The requirement of zinc for  
 506 improvement of crop yield and mineral nutrition in the maize-mungbean-rice system. *Plant*  
 507 *and soil*, 306, 13-22, 2008.
- 508 Hu, X., Su, F., Ju, X., Gao, B., Oenema, O., Christie, P., Huang, B., Jiang, R. and Zhang, F.:  
 509 Greenhouse gas emissions from a wheat-maize double cropping system with different nitrogen  
 510 fertilization regimes, *Environ. Pollut.*, 176, 198-207, 2013.
- 511 Huang, J., Chen, Y., Sui, P. and Gao, W.: Estimation of net greenhouse gas balance using crop-and  
 512 soil-based approaches: Two case studies, *Sci. Total Environ.*, 456, 299-306, 2013a.
- 513 Huang, T., Gao, B., Christie, P. and Ju, X.: Net global warming potential and greenhouse gas intensity  
 514 in a double-cropping cereal rotation as affected by nitrogen and straw management,  
 515 *Biogeosciences*, 10, 7897-7911, 2013b.
- 516 Huang, Y. and Sun, W.: Changes in topsoil organic carbon of croplands in mainland China over the last  
 517 two decades, *Chinese Sci Bull.*, 51, 1785-1803, 2006.
- 518 Huang, Y., Zhang, W., Zheng, X. H., Li, J. and Yu, Y. Q.: Modeling methane emission from rice  
 519 paddies with various agricultural practices, *J. Geophys. Res. Atmos.*, 109, D08,  
 520 doi:10.1029/2003JD004401, 2004.
- 521 IPCC.: *Climate Change 2013: The Physical Science Basis: working group I contribution to the Fifth*  
 522 *Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge University  
 523 *Press*, Stockholm, 2013.
- 524 Jia, J. X., Ma, Y. C. and Xiong, Z. Q.: Net ecosystem carbon budget, net global warming potential and

525 greenhouse gas intensity in intensive vegetable ecosystems in China, *Agr. Ecosyst.*  
526 *Environ.*, 150, 27-37, 2012.

527 Ju, X., Xing, G., Chen, X., Zhang, S., Zhang, L., Liu, X., Cui, Z., Yin, B., Christie, P. and Zhu, Z.:  
528 Reducing environmental risk by improving N management in intensive Chinese agricultural  
529 systems, *P. Natl. Acad. Sci. USA*, 106, 3041-3046, 2009.

530 Ju, X., Lu, X., Gao, Z., Chen, X., Su, F., Kogge, M., Römheld, V., Christie, P. and Zhang, F.: Processes  
531 and factors controlling N<sub>2</sub>O production in an intensively managed low carbon calcareous soil  
532 under sub-humid monsoon conditions, *Environ. Pollut.*, 159, 1007-1016, 2011.

533 Kabata-Pendias, A. and Mukherjee, A. B.: Trace elements from soil to human. Springer Science &  
534 Business Media, 2007.

535 Lal, R.: Carbon emission from farm operations, *Environ. Int.*, 30, 981-990, 2004.

536 Le Mer, J. and Roger, P.: Production, oxidation, emission and consumption of methane by soils: a  
537 review, *European Journal of Soil Biology.*, 37, 25-50, 2001.

538 Li, C. S., Salas, W., DeAngelo, B. and Rose, S.: Assessing alternatives for mitigating net greenhouse  
539 gas emissions and increasing yields from rice production in China over the next twenty years,  
540 *J. Environ. Qual.*, 35, 1554-1565, 2006.

541 Liu, L., Xue Y., Sun, X., Wang, Z. and Yang, J.: Effects of water management methods on grain yield  
542 and fertilizer-nitrogen use efficiency in rice, *Chin J Rice Sci.*, 23, 282-288, 2009.

543 Liu, Y. L., Zhou, Z., Zhang, X., Xu, X., Chen, H. and Xiong, Z.: Net global warming potential and  
544 greenhouse gas intensity from the double rice system with integrated soil-crop system  
545 management: A three-year field study, *Atmos. Environ.*, 116, 92-101, 2015.

546 Ma, J., Ma, E., Xu, H., Yagi, K. and Cai, Z.: Wheat straw management affects CH<sub>4</sub> and N<sub>2</sub>O emissions  
547 from rice fields, *Soil Biol. Biochem.*, 41, 1022-1028, 2009.

548 Ma, Y., Kong, X., Yang, B., Zhang, X., Yan, X., Yang, J. and Xiong, Z.: Net global warming potential  
549 and greenhouse gas intensity of annual rice-wheat rotations with integrated soil-crop system  
550 management, *Agr. Ecosyst. Environ.*, 164, 209-219, 2013.

551 Makino, A.: Photosynthesis, grain yield, and nitrogen utilization in rice and wheat, *Plant physiol.*, 155,  
552 125-129, 2011.

553 Mosier, A. R., Halvorson, A. D., Reule, C. A. and Liu, X. J.: Net global warming potential and  
554 greenhouse gas intensity in irrigated cropping systems in northeastern Colorado, *J. Environ.*  
555 *Qual.*, 35, 1584-1598, 2006.

556 Murdiyarso, D., Hergoualc'h, K. and Verchot, L. V.: Opportunities for reducing greenhouse gas  
557 emissions in tropical peatlands, *P. Natl. Acad. Sci. USA*, 107, 19655-19660, 2010.

558 Myhre, G., Shindell, D., Bréon, F.M., Collins, W., Fuglestedt, J., Huang, J., Koch, D., Lamarque, J.F.,  
559 Lee, D., Mendoza, B., Nakajima, T., Robock, A., Stephens, G., Takemura, T. and Zhang, H.:  
560 Anthropogenic and natural radiative forcing. In: Stocker, T.F., Qin, D., Plattner, G.K., Tignor,  
561 M., Allen, S.K., Boschung, J., Nauels, A., Xia, Y., Bex, V., Midgley, P.M. (Eds.), *Climate*  
562 *Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth*  
563 *Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University*  
564 *Press, Cambridge, UK, New York, NY, USA, 659-740, 2013.*

565 Paul, J.W., Beauchamp, E.G. and Zhang, X.: Nitrous and nitric oxide emissions during nitrification and  
566 denitrification from manure-amended soil in the laboratory, *Can. J. Soil. Sci.*, 73, 539-553,  
567 1993.

568 Peng, S., Buresh, R. J., Huang, J., Yang, J., Zou, Y., Zhong, X., Wang, G. and Zhang, F.: Strategies for

569 overcoming low agronomic nitrogen use efficiency in irrigated rice systems in China, *Field*  
570 *Crop. Res.*, 96, 37-47, 2006.

571 Qin, Y. M., Liu, S.W., Guo, Y. Q., Liu, Q. and Zou, J. W.: Methane and nitrous oxide emissions from  
572 organic and conventional rice cropping systems in Southeast China, *Biol. Fertil. Soils.*, 46,  
573 825-834, 2010.

574 Robertson, G. P. and Grace, P. R.: Greenhouse gas fluxes in tropical and temperate agriculture: The  
575 need for a full-cost accounting of global warming potentials, *Environment Development and*  
576 *Sustainability*, 6, 51-63, 2004.

577 Rudaz, A. O., Wälti, E., Kyburz, G., Lehmann, P. and Fuhrer, J.: Temporal variation in N<sub>2</sub>O and N<sub>2</sub>  
578 fluxes from a permanent pasture in Switzerland in relation to management, soil water content  
579 and soil temperature, *Agr. Ecosyst. Environ.*, 73, 83-91, 1999.

580 Sainju, U. M., Stevens, W. B., Caesar-TonThat, T., Liebig, M. A. and Wang, J.: Net global warming  
581 potential and greenhouse gas intensity influenced by irrigation, tillage, crop rotation, and  
582 nitrogen fertilization, *J. Environ. Qual.*, 43, 777-788, 2014.

583 Schlesinger, W. H.: On fertilizer-induced soil carbon sequestration in China's croplands, *Glob. Change*  
584 *Biol.*, 16, 849-850, 2010.

585 Shang, Q., Yang, X., Gao, C., Wu, P., Liu, J., Xu, Y., Shen, Q., Zou, J. and Guo, S.: Net annual global  
586 warming potential and greenhouse gas intensity in Chinese double rice-cropping systems: a  
587 3-year field measurement in long-term fertilizer experiments, *Glob. Change Biol.*, 17,  
588 2196-2210, 2011.

589 Six, J., Ogle, S. M., Conant, R. T., Mosier, A. R. and Paustian, K.: The potential to mitigate global  
590 warming with no-tillage management is only realized when practised in the long term, *Glob.*  
591 *Change Biol.*, 10, 155-160, 2004.

592 Slaton, N. A., Norman, R. J. and Wilson, C. E.: Effect of zinc source and application time on zinc  
593 uptake and grain yield of flood-irrigated rice, *Agron. J.*, 97, 272-278, 2005.

594 Smith, P., Martino, D., Cai, Z., Gwary, D., Janzen, H., Kumar, P., McCarl, B., Ogle, S., O'Mara, F. and  
595 Rice, C.: Greenhouse gas mitigation in agriculture, *Philos. T. R. Soc. B.*, 363, 789-813, 2008.

596 Snyder, C., Bruulsema, T., Jensen, T. and Fixen, P.: Review of greenhouse gas emissions from crop  
597 production systems and fertilizer management effects, *Agr. Ecosyst. Environ.*, 133, 247-266,  
598 2009.

599 Thangarajan, R., Bolan, N. S., Tian, G., Naidu, R. and Kunhikrishnan, A.: Role of organic amendment  
600 application on greenhouse gas emission from soil, *Sci. Total Environ.*, 465, 72-96, 2013.

601 Tilman, D., Balzer, C., Hill, J., Befort, B. L. and Affiliations, A.: Global food demand and the  
602 sustainable intensification of agriculture, *P. Natl. Acad. Sci. USA*, 108, 20260-20264, 2011.

603 Wang, J., Zhang, X., Xiong, Z., Khalil, M.A.K., Zhao, X., Xie, Y. and Xing, G.: Methane emissions  
604 from a rice agroecosystem in South China: Effects of water regime, straw incorporation and  
605 nitrogen fertilizer, *Nutr. Cycl. Agroecosyt.*, 93, 103-112, 2012.

606 Wang, L.J., Nie, Q., Li, M., Zhang, F.S., Zhuang, J.Q., Yang, W.S., Li, T.J. and Wang, Y.H.:  
607 Biosilicified structures for cooling plant leaves: a mechanism of highly efficient midinfrared  
608 thermal emission, *Appl. Phys. Lett.*, 87, 194105, doi:10.1063/1.2126115, 2005.

609 West, T. O. and Marland, G.: A synthesis of carbon sequestration, carbon emissions, and net carbon  
610 flux in agriculture: comparing tillage practices in the United States, *Agr. Ecosyst. Environ.*, 91,  
611 217-232, 2002.

612 Wu, C., Ye, D., Lin, H., Ni, R., Lai L. and Lin, H.: Effects of transplanting density on rice yield and its

613 quality, Chinese Agricultural Science Bulletin, 21, 190-205, 2005.

614 Xie, Y., Zhang, J., Jiang, H., Yang, J., Deng, S., Li, X., Guo, J., Li, L., Liu, X. and Zhou, G.: Effects of  
615 different fertilization practices on greenhouse gas emissions from paddy soil, Journal of  
616 Agro-Environment Science, 3, 578-584, 2015.

617 Xiong, Z. Q., Xing, G. X. and Zhu, Z. L.: Nitrous oxide and methane emissions as affected by water,  
618 soil and nitrogen, Pedosphere, 17, 146-155, 2007.

619 Yan, X. Y., Yagi, K., Akiyama, H. and Akimoto, H.: Statistical analysis of the major variables  
620 controlling methane emission from rice fields, Glob. Change Biol., 11, 1131-1141, 2005.

621 Yang, B., Xiong, Z., Wang, J., Xu, X., Huang, Q. and Shen, Q.: Mitigating net global warming  
622 potential and greenhouse gas intensities by substituting chemical nitrogen fertilizers with  
623 organic fertilization strategies in rice-wheat annual rotation systems in China: A 3-year field  
624 experiment, Ecol. Eng., 81, 289-297, 2015.

625 Zhang, F., Cui, Z., Fan, M., Zhang, W., Chen, X. and Jiang, R.: Integrated soil-crop system  
626 management: reducing environmental risk while increasing crop productivity and improving  
627 nutrient use efficiency in China, J. Environ Qual., 40, 1051-1057, 2011.

628 Zhang, W., Dou, Z., He, P., Ju, X., Powlson, D., Chadwick, D., Norse, D., Lu, Y., Zhang, Y. and Wu, L.:  
629 New technologies reduce greenhouse gas emissions from nitrogenous fertilizer in China, P.  
630 Natl. Acad. Sci. USA, 110, 8375-8380, 2013.

631 Zhang, X., Fan, C., Ma, Y., Liu, Y., Li, L., Zhou, Q. and Xiong, Z.: Two approaches for net ecosystem  
632 carbon budgets and soil carbon sequestration in a rice-wheat rotation system in China, Nutr.  
633 Cycl. Agroecosys., 100, 301-313, 2014.

634 Zhao, M., Tian, Y., Ma, Y., Zhang, M., Yao, Y., Xiong, Z., Yin, B. and Zhu, Z.: Mitigating gaseous  
635 nitrogen emissions intensity from a Chinese rice cropping system through an improved  
636 management practice aimed to close the yield gap, Agr. Ecosyst. Environ., 203, 36-45, 2015.

637 Zhu, Z.: Fate and management of fertilizer nitrogen in agro-ecosystems. Nitrogen in Soils of China,  
638 Springer Netherlands, 239-279, 1997.

639 Zou, J. W., Huang, Y., Jiang, J. Y., Zheng, X. H. and Sass, R.L.: A 3-year field measurement of methane  
640 and nitrous oxide emissions from rice paddies in China: effects of water regime, crop residue,  
641 and fertilizer application, Global Biogeochem. Cy., 19, GB002401,  
642 doi:10.1029/2004GB002401, 2005.

**Table 1**

The establishment of different treatments for the annual rice-wheat rotations during the 2011–2014 cycle.

Scenario	NN <sup>a</sup>	FP	ISSM-N1	ISSM-N2	ISSM-N3	ISSM-N4
Rice-growing season						
Chemical fertilizer application rate (N:P <sub>2</sub> O <sub>5</sub> :K <sub>2</sub> O:Na <sub>2</sub> SiO <sub>3</sub> :ZnSO <sub>4</sub> , kg ha <sup>-1</sup> )	0:90:120:0:0	300:90:120:0:0	225:90:120:0:0	270:90:120:0:0	300:108:144:225:15	375:126:180:225:15
Split N application ratio		6:2:0:2	5:1:2:2	5:1:2:2	5:1:2:2	5:1:2:2
Rapeseed cake manure (t ha <sup>-1</sup> )	0	0	0	0	2.25 <sup>c</sup>	2.25
Water regime	F-D-F-M <sup>b</sup>	F-D-F-M	F-D-F-M	F-D-F-M	F-D-F-M	F-D-F-M
Planting density (cm)	20×20	20×20	20×15	20×15	20×15	20×20
Wheat-growing season						
Chemical fertilizer application rate (N:P <sub>2</sub> O <sub>5</sub> :K <sub>2</sub> O, kg ha <sup>-1</sup> )	0:90:180	180:90:180	135:90:180	162:90:180	180:108:216	225:126:270
Split N application ratio		6:1:3	6:1:3	6:1:3	6:1:3	6:1:3
Seed sowing density (kg ha <sup>-1</sup> )	180	180	180	180	180	180

<sup>a</sup>NN, no N application; FP, farmers' practice; The four integrated soil-crop system management (ISSM) practices at different nitrogen application rates relative to the FP rate of 300 kg N ha<sup>-1</sup> for the rice crop and 180 kg N ha<sup>-1</sup> for the wheat crop, namely, ISSM-N1 (25% reduction), ISSM-N2 (10% reduction), ISSM-N3 (FP rate) and ISSM-N4 (25% increase). Urea, calcium biphosphate and potassium chloride were used as N, P and K fertilizer respectively.

<sup>b</sup>F-D-F-M, flooding-midseason drainage-re-flooding-moist irrigation.

<sup>c</sup>112.5 kg N ha<sup>-1</sup> in the form of rapeseed cake was applied as basal fertilizer and included in the total N rate for calculating agronomic NUE.

**Table 2**

Agricultural management practices for chemical inputs and farm operations in the rice and wheat cropping seasons.

<u>Treatment</u>	<u>Rice season</u>													
	<u>Chemical inputs (kg ha<sup>-1</sup>)<sup>a</sup></u>								<u>Farm operations<sup>b</sup></u>					
	<u>N</u>	<u>P</u>	<u>K</u>	<u>Si</u>	<u>Zn</u>	<u>Herbicide</u>	<u>Insecticide</u>	<u>Fungicide</u>	<u>Irrigation</u>	<u>Tillage</u>	<u>Crop planting</u>	<u>Farm manure</u>	<u>Crop harvest</u>	<u>Threshing<sup>c</sup></u>
								<u>(cm)</u>	<u>(kg diesel ha<sup>-1</sup>)</u>	<u>(event)</u>	<u>(kg ha<sup>-1</sup>)</u>	<u>(kg diesel ha<sup>-1</sup>)</u>	<u>(kw•h ha<sup>-1</sup>)</u>	
<u>NN</u>	<u>0</u>	<u>90</u>	<u>120</u>	<u>0</u>	<u>0</u>	<u>1</u>	<u>13</u>	<u>2</u>	<u>78</u>	<u>20</u>	<u>1</u>	<u>0</u>	<u>6</u>	<u>74</u>
<u>FP</u>	<u>300</u>	<u>90</u>	<u>120</u>	<u>0</u>	<u>0</u>	<u>1</u>	<u>13</u>	<u>2.4</u>	<u>78</u>	<u>20</u>	<u>1</u>	<u>0</u>	<u>6</u>	<u>80</u>
<u>ISSM-N1</u>	<u>225</u>	<u>90</u>	<u>120</u>	<u>0</u>	<u>0</u>	<u>1</u>	<u>13</u>	<u>2.4</u>	<u>57</u>	<u>20</u>	<u>1</u>	<u>0</u>	<u>6</u>	<u>74</u>
<u>ISSM-N2</u>	<u>270</u>	<u>90</u>	<u>120</u>	<u>0</u>	<u>0</u>	<u>1</u>	<u>13</u>	<u>2.4</u>	<u>57</u>	<u>20</u>	<u>1</u>	<u>0</u>	<u>6</u>	<u>100</u>
<u>ISSM-N3</u>	<u>300</u>	<u>108</u>	<u>144</u>	<u>225</u>	<u>15</u>	<u>1</u>	<u>17</u>	<u>3.5</u>	<u>57</u>	<u>20</u>	<u>1</u>	<u>2250</u>	<u>6</u>	<u>100</u>
<u>ISSM-N4</u>	<u>375</u>	<u>126</u>	<u>180</u>	<u>225</u>	<u>15</u>	<u>1</u>	<u>26</u>	<u>5</u>	<u>57</u>	<u>20</u>	<u>1</u>	<u>2250</u>	<u>6</u>	<u>175</u>
									<u>Wheat season</u>					
<u>NN</u>	<u>0</u>	<u>90</u>	<u>180</u>	<u>0</u>	<u>0</u>	<u>1</u>	<u>5</u>	<u>2</u>	<u>0</u>	<u>17</u>	<u>1</u>	<u>0</u>	<u>5</u>	<u>61</u>
<u>FP</u>	<u>180</u>	<u>90</u>	<u>180</u>	<u>0</u>	<u>0</u>	<u>1</u>	<u>7</u>	<u>2</u>	<u>0</u>	<u>17</u>	<u>1</u>	<u>0</u>	<u>5</u>	<u>67</u>
<u>ISSM-N1</u>	<u>135</u>	<u>90</u>	<u>180</u>	<u>0</u>	<u>0</u>	<u>1</u>	<u>7</u>	<u>2</u>	<u>0</u>	<u>17</u>	<u>1</u>	<u>0</u>	<u>5</u>	<u>65</u>
<u>ISSM-N2</u>	<u>162</u>	<u>90</u>	<u>180</u>	<u>0</u>	<u>0</u>	<u>1</u>	<u>7</u>	<u>2</u>	<u>0</u>	<u>17</u>	<u>1</u>	<u>0</u>	<u>5</u>	<u>77</u>
<u>ISSM-N3</u>	<u>180</u>	<u>108</u>	<u>216</u>	<u>0</u>	<u>0</u>	<u>1</u>	<u>10</u>	<u>2.5</u>	<u>0</u>	<u>17</u>	<u>1</u>	<u>0</u>	<u>5</u>	<u>77</u>
<u>ISSM-N4</u>	<u>225</u>	<u>126</u>	<u>270</u>	<u>0</u>	<u>0</u>	<u>1</u>	<u>15</u>	<u>4</u>	<u>0</u>	<u>17</u>	<u>1</u>	<u>0</u>	<u>5</u>	<u>100</u>

<sup>a</sup>There was no machinery used for fertilizer application.

<sup>b</sup>Tillage and crop harvest, crop planting, and threshing were calculated by diesel fuel (kg ha<sup>-1</sup>), event and electricity (kw•h ha<sup>-1</sup>), respectively.

<sup>c</sup>Electricity energy is calculated according to the power and working hours. The power of thresher is 15 kilowatt in this experiment.

**Table 32**

Seasonal CH<sub>4</sub> and N<sub>2</sub>O emissions, and yields during rice and wheat cropping seasons in the three cycles of 2011–2014.

Treatment	Rice season			Wheat season		
	CH <sub>4</sub> (kg C ha <sup>-1</sup> )	N <sub>2</sub> O (kg N ha <sup>-1</sup> )	Yield (t ha <sup>-1</sup> )	CH <sub>4</sub> (kg C ha <sup>-1</sup> )	N <sub>2</sub> O (kg N ha <sup>-1</sup> )	Yield (t ha <sup>-1</sup> )
2011						
NN	153±10.8c	0.03±0.05c	5.85±0.08f	-0.48±0.63a	0.45±0.09d	1.74±0.18d
FP	266±25.3b	0.11±0.08c	8.38±0.35e	-0.48±1.86a	1.43±0.19b	5.67±0.20b
ISSM-N1	212±30.3bc	0.08±0.03c	9.27±0.26d	0.78±0.97a	0.65±0.11cd	5.05±0.16c
ISSM-N2	220±32.5bc	0.17±0.11bc	9.79±0.44c	2.25±2.07a	0.80±0.06c	5.71±0.18b
ISSM-N3	518±58.9a	0.38±0.15ab	10.81±0.26b	0.04±3.23a	1.40±0.10b	5.31±0.26bc
ISSM-N4	561±50.9a	0.37±0.07a	11.76±0.24a	-0.09±1.40a	1.93±0.09a	6.15±0.15a
2012						
NN	149±25.8d	0.13±0.10c	5.80±0.22f	-4.32±7.29a	0.65±0.09d	1.73±0.11c
FP	239±34.5c	0.33±0.11bc	8.72±0.62e	4.85±10.30a	2.13±0.43ab	5.64±0.34ab
ISSM-N1	226±30.4cd	0.27±0.07bc	9.43±0.34d	1.46±6.38a	1.39±0.14c	4.94±0.38b
ISSM-N2	228±32.6cd	0.38±0.29bc	9.99±0.50c	-1.02±0.84a	1.77±0.38bc	5.78±0.59ab
ISSM-N3	431±26.8b	0.52±0.16ab	10.92±0.61b	2.45±8.35a	2.19±0.24ab	5.39±0.39ab
ISSM-N4	536±58.7a	0.78±0.13a	12.24±0.60a	5.91±6.18a	2.61±0.42a	6.10±0.49a
2013						
NN	101±39.2b	0.16±0.09b	5.84±0.15f	-1.45±1.34a	0.35±0.06c	1.80±0.03c
FP	141±25.2b	0.43±0.39ab	8.67±0.26e	-3.70±1.76a	0.80±0.20ab	5.70±0.30ab
ISSM-N1	135±15.7b	0.19±0.16ab	9.66±0.29d	-1.00±1.61a	0.49±0.16bc	5.15±0.20b
ISSM-N2	129±32.2b	0.26±0.13ab	10.15±0.07c	-0.79±1.60a	0.69±0.24abc	5.80±0.18ab
ISSM-N3	256±45.6a	0.59±0.42ab	11.14±0.10b	-0.62±1.14a	0.71±0.10ab	5.51±0.33ab
ISSM-N4	304±22.3a	0.74±0.40a	12.34±0.16a	0.55±1.68a	1.02±0.11a	6.19±0.63a
Average 2011–2013 <sup>a</sup>						
NN <sup>b</sup>	135±19.6d	0.11±0.05c	5.83±0.04f	-2.08±1.89a	0.48±0.07d	1.75±0.04d
FP <sup>b</sup>	215±19.9c	0.29±0.13bc	8.59±0.25e	0.22±3.96a	1.45±0.24b	5.67±0.16b
ISSM-N1 <sup>b</sup>	191±19.2c	0.18±0.06c	9.45±0.18d	0.42±2.77a	0.84±0.08c	5.04±0.08c
ISSM-N2 <sup>b</sup>	192±11.6c	0.27±0.12bc	9.98±0.25c	0.15±0.58a	1.08±0.12c	5.76±0.22ab
ISSM-N3 <sup>b</sup>	402±23.8b	0.50±0.16ab	10.95±0.13b	0.63±3.51a	1.43±0.05b	5.40±0.16bc
ISSM-N4 <sup>b</sup>	467±39.2a	0.68±0.15a	12.11±0.28a	2.12±2.57a	1.85±0.16a	6.14±0.35a

<sup>a</sup>Mean ± SD, different lower case letters within the same column for each item indicate significant differences at  $P < 0.05$  according to Tukey's multiple range test.

<sup>b</sup>See Table 1 for treatment codes.

**Table 43**

Repeated-measures analysis of variance (MANOVA) for the effects of cultivation patterns (P) and cropping year (Y) on mean CH<sub>4</sub> and N<sub>2</sub>O emissions, and mean rice and wheat grain yields in the 2011–2014 cycle.

Crop season	Source	<i>df</i>	CH <sub>4</sub> (kg C ha <sup>-1</sup> )	N <sub>2</sub> O (kg N ha <sup>-1</sup> )	Yield (t ha <sup>-1</sup> )
Rice	Between subjects				
	P	5	35.3***	3.71***	123***
	Within subjects				
	Y	2	20.7***	0.88**	1.15**
	P×Y	10	6.73***	0.15	0.37
Wheat	Between subjects				
	P	5	0.26	14.8***	76.3***
	Within subjects				
	Y	2	0.55*	15.1***	0.08
	P×Y	10	0.83	4.39***	0.05
Rice-Wheat	Between subjects				
	P	5	37.2***	24.2***	153***
	Within subjects				
	Y	2	20.5***	5.83***	0.70*
	P×Y	10	6.50***	1.11	0.17

*df* – degrees of freedom, \*  $P < 0.05$ , \*\*  $P < 0.01$ , and \*\*\* $P < 0.001$  represent significant at the 0.05, 0.01 and 0.001 probability level, respectively.

**Table 54**

Agricultural management practices for chemical inputs and farm operations and contributions to carbon dioxide equivalents (kg CO<sub>2</sub> eq. ha<sup>-1</sup>yr<sup>-1</sup>) in the annual rice-wheat rotations from 2011 to 2014 (chemical inputs and farm operations used in each year were similar except for irrigation water).

Treatment	Chemical inputs (kg ha <sup>-1</sup> ) <sup>a</sup>								Farm operations <sup>c</sup>							
	N	P	K	Si	Zn	Herbicide	Insecticide	Fungicide	Irrigation (cm) <sup>b</sup>			Tillage and raking (kg diesel ha <sup>-1</sup> )	Crop planting (event)	Farm manure (kg ha <sup>-1</sup> )	Crop harvest (kg diesel ha <sup>-1</sup> )	Threshing (kw•h ha <sup>-1</sup> )
									2011	2012	2013					
NN <sup>d</sup>	0	180	300	0	0	2	18	4	75	80	80	37	2	0	11	135
FP	480	180	300	0	0	2	20	4.4	75	80	80	37	2	0	11	147
ISSM-N1	360	180	300	0	0	2	20	4.4	50	65	55	37	2	0	11	139
ISSM-N2	432	180	300	0	0	2	20	4.4	50	65	55	37	2	0	11	177
ISSM-N3	480	216	360	225	15	2	27	6	50	65	55	37	2	2250	11	177
ISSM-N4	600	252	450	225	15	2	41	9	50	65	55	37	2	2250	11	275
	Chemical inputs (E <sub>i</sub> ) (kg CO <sub>2</sub> eq. ha <sup>-1</sup> )								Farm operations (E <sub>o</sub> ) (kg CO <sub>2</sub> eq. ha <sup>-1</sup> )							
NN	0	132	165	0	0	46	338	53	1419	1514	1514	127	23	0	37	36
FP	2288	132	165	0	0	46	375	59	1419	1514	1514	127	23	0	37	39
ISSM-N1	1716	132	165	0	0	46	375	59	946	1230	1041	127	23	0	37	37
ISSM-N2	2059	132	165	0	0	46	375	59	946	1230	1041	127	23	0	37	47
ISSM-N3	2288	158	198	58	6	46	506	79	946	1230	1041	127	23	62	37	47
ISSM-N4	2860	185	248	58	6	46	768	129	946	1230	1041	127	23	62	37	73

<sup>a</sup>The carbon emission coefficients were 1.3, 0.2, 0.15, 6.3, 5.1 and 3.9 C cost (kg C eq. kg<sup>-1</sup> active ingredient) per applied nitrogen fertilizer, phosphorus, potassium, herbicide, insecticide and fungicide, respectively, as referred to in Lal (2004). We collected data specific to China's fertilizer manufacture and consumption, and then estimated carbon emissions coefficients were 0.07 and 0.1 C cost (kg C eq. kg<sup>-1</sup> active ingredient) per applied Si and Zn fertilizer, respectively.

<sup>b</sup>The carbon emission coefficient for irrigation was 5.16 C cost (kg C eq. cm<sup>-1</sup> ha<sup>-1</sup>) as referred to in Lal (2004).

<sup>c</sup>The carbon emission coefficients were 0.94 C cost (kg C eq. kg<sup>-1</sup> diesel) for tillage, raking and harvesting, 3.2 C cost (kg C eq. event<sup>-1</sup>) for crop planting, 0.0075 C cost (kg C eq. kg<sup>-1</sup>) for farm manure application and 0.0725 C cost (kg C eq. (kw•h)<sup>-1</sup>) for threshing, as referred to in Lal (2004).

<sup>d</sup>See Table 1 for treatment codes.

**Table 65**

Mean global warming potential (GWP) and greenhouse gas intensity (GHGI) over the three rice season, wheat season and annual cycles of the 2011rice season–2014wheat season.

Treatment	CH <sub>4</sub>	N <sub>2</sub> O	Ei	Eo	SOCSR	GWP <sup>a</sup>	Grain yield	GHGI <sup>b</sup>
	kg CO <sub>2</sub> eq. ha <sup>-1</sup> yr <sup>-1</sup>					t ha <sup>-1</sup> yr <sup>-1</sup>		kg CO <sub>2</sub> eq. t <sup>-1</sup> grain
Rice season								
NN	5026±733d	44±20c	424	1601	-396±164c	7492±706d	5.83±0.04f	1285±123b
FP	8035±742c	121±53bc	1859	1603	585±198ab	11032±555c	8.59±0.25e	1285±68b
ISSM-N1	7132±716c	75±24c	1502	1191	246±218b	9654±800c	9.45±0.18d	1021±81c
ISSM-N2	7186±434c	112±49bc	1716	1198	355±97ab	9858±484c	9.98±0.25c	989±67c
ISSM-N3	15005±888b	208±66ab	2037	1260	691±252a	17818±786b	10.95±0.13b	1626±54a
ISSM-N4	17427±1463a	284±60a	2626	1280	773±174a	20844±1452a	12.11±0.28a	1720±108a
Wheat season								
NN	-78±71a	201±28d	310	104	-396±164c	934±214b	1.75±0.04d	533±125a
FP	8±148a	605±99b	1206	105	585±198ab	1339±129b	5.67±0.16b	236±21b
ISSM-N1	16±103a	351±32c	991	105	246±218b	1217±342b	5.04±0.08c	241±68b
ISSM-N2	6±22a	451±49c	1120	108	355±97ab	1329±109b	5.76±0.22ab	231±26b
ISSM-N3	23±131a	598±20b	1302	108	691±252a	1340±290b	5.40±0.16bc	247±48b
ISSM-N4	79±96a	772±66a	1674	114	773±174a	1867±175a	6.14±0.35a	305±33b
Rice-wheat rotation								
NN <sup>d</sup>	4948±704d <sup>c</sup>	246±26d	734	1705	-792±327c	8425±711d	7.58±0.04d	1111±94b
FP	8043±858c	725±49b	3065	1708	1170±396ab	12371±583c	14.26±0.36c	868±29c
ISSM-N1	7141±709c	426±55c	2493	1296	491±435b	10871±990c	14.50±0.14c	750±68d
ISSM-N2	7192±424c	563±86c	2836	1306	709±193ab	11187±552c	15.74±0.44b	712±52d
ISSM-N3	15028±833b	806±77b	3339	1368	1383±503a	19158±761b	16.36±0.18b	1171±37ab
ISSM-N4	17506±1396a	1056±58a	4300	1394	1545±348a	22711±1438a	18.26±0.46a	1245±93a

<sup>a</sup>GWP (kg CO<sub>2</sub> eq. ha<sup>-1</sup>yr<sup>-1</sup>) = 28 × CH<sub>4</sub> + 265 × N<sub>2</sub>O + Ei + Eo - 44/12 × SOCSR, Ei (agrochemical inputs), Eo (farm operations), SOCSR (SOC sequestration rate) is divided by 2 to roughly estimate the GWP from rice and wheat season, respectively. All other items were actually measured for each season.

<sup>b</sup>GHGI (kg CO<sub>2</sub> eq. t<sup>-1</sup> grain) = GWP/grain yields

<sup>c</sup>Different lower case letters within the same column for each item indicate significant differences at *P*<0.05 based on Tukey's multiple range tests.

<sup>d</sup>See Table 1 for treatment codes.

[Fig 1 Daily mean air temperature and precipitation during the rice-wheat rotation in 2011–2014 in Changshu, China.](#)

**Fig 24** Rice and wheat agronomic nitrogen use efficiency (NUE) in 2011–2014 in Changshu, China.

Different letters indicate a significant difference between treatments ( $p < 0.05$ ). See Table 1 for treatment codes.

**Fig 32** Seasonal variation of methane ( $\text{CH}_4$ ) fluxes from the rice-wheat rotation cropping systems from 2011 to 2014. The black and gray part in figure separates different grain growth periods. See Table 1 for treatment codes. The solid arrows indicate fertilization.

**Fig 43** Seasonal variation of nitrous oxide ( $\text{N}_2\text{O}$ ) fluxes from rice-wheat rotation cropping systems in three annual cycles over the period 2011–2014. The black and gray part in the figure separates different growth periods. See Table 1 for treatment codes. The solid arrows indicate fertilization.