

Author response

bg-2015-599

(referee comments are printed in *italic*, author responses are printed in blue)

Answers to Comments of Referee #1

Comment: *Both the title and the abstract focus on carbon budget. Readers, however, would be more interested in the total GHG budget actually. Thus I suggest authors describe more about the components and characteristics of the GHG budget of the system in the abstract and text and reflect it in the title as well.*

Answer: We are well aware that the quantification of the full GHG budget of the pasture system is an important final target (application) of our research. However, we think that from a scientific point of view the quantification of the pasture carbon budget is important and complex enough to be studied in an individual paper on its own right. Thus, as clearly declared in the title and in the abstract, the focus of this paper is the carbon budget of a pasture including the discussion of the different components (fluxes) contributing to the carbon budget of the same system but with different boundaries. The additional presentation of the GHG budget at the end of the manuscript is done only for context reasons and to compare the magnitude and typical uncertainty of the carbon budget (NECB) to the other GHG fluxes.

A consistent evaluation of the GHG budget would need a detailed assessment of the N₂O exchange which is beyond the scope of this paper and will be presented elsewhere.

We did not add more about the GHG budget but revised the text and omitted the phrasing 'GHG budget'.

Comment: *Fluxes related the grazing were monitored for 99 days only, the way that the results were extended to a year was not clearly described. Was it a linear exploration?*

Answer: While the carbon budget (NECB) for quantifying the soil C sequestration of the study field is quantified for the entire year, some animal related budget components are first determined for the cow herd and thus need then an appropriate time attribution to the study field (here 99 days for the year 2013).

We made some rephrasing in the text to clarify this issue and we added a Figure (Fig. 1) with the pasture days to Sect. 2.1 where the different durations are marked in different colors. We also added the attributed time used in Eq. (4) to Table S2 to make that even more clear.

Comment: *Descriptions on the determination of the uncertainties of NECB and other fluxes are over simplified.*

Answer: As suggested by referee #2 we included a more detailed description of uncertainty calculations in the Supplementary and also added some details in Sect. 2.4.

Answers to Referee Comments of Susanna Rutledge (Referee #2)

Main concerns

- 1) *For a paper that claims to discuss the uncertainties in the NECB ('flux uncertainties' is even in the title), the uncertainties are really not very well discussed in the paper. I assume the calculations have been done correctly, but they need to be described in much more detail to allow their reproduction by fellow scientists. In several sections (e.g. in P20078, L27/28; Section 2.4.1, L24; Section 2.4.3) short descriptions of final uncertainty estimates of the components of the NECB have been provided, but almost all these sections need to provide more information and clarification. An additional section in the Supplementary Materials would be most useful so that the main text remains uncluttered.*

As suggested, we added a more systematic and detailed description of the uncertainty calculation/estimation of the various carbon budget components in the Supplement. The following uncertainty estimations are now addressed in the Supplement:

- **Animal intake:** description of uncertainty estimation of dry matter intake, carbon content of forage and concentrates
- **Milk carbon content:** description of uncertainty estimation for $F_{C\text{-products}}$
- **CO₂ exchange:** more details to the uncertainty estimation of $F_{C\text{-CO}_2}$
- **Fertilization:** description of uncertainty estimation of $F_{C\text{-fertil}}$

For example:

- *I realise that for feed intake (Section 2.4.2) the uncertainties may be very hard to determine. Was any attempt made to estimate the uncertainty in $E_{DM\text{-intake}}$ (which would feed into $F_{C\text{-grazing}}$)?*

Section S1.1 in the Supplement now includes a detailed description of the uncertainty estimation for $E_{DM\text{-intake}}$ including the propagation for the error of $F_{C\text{-grazing}}$.

Also, do I read correctly that it is implied that the uncertainty for the amount of supplement feed provided was assumed to be zero?

We recalculated the uncertainty value for the supplement feed ($F_{C\text{-feed,off}}$) originally stated as 6% to a total of 16%. This value was derived as a combination from the estimated uncertainties of the weighing of the fresh matter intake for each cow (15%), the DM content analysis (4%) and the C content analysis (1%). A detailed description of this procedure was added to Sect. S1.1 in the Supplement.

- *It would be helpful if the uncertainties in DM amount, DM content and C content were spelled out explicitly (e.g. P20082, L2-3).*

Uncertainty values were added at the end of Sect. 2.4.2 and the calculation/determination was described in detail in Sect. S1.1 in the Supplement.

- *Section 2.4.3 uncertainties in excreta need more explanation. Uncertainties in which budget terms contributed to the uncertainties in $F_{C\text{-excreta}}$ and how were uncertainties combined?*

Originally, we quantified the excreta term as the residual of the animal C budget (which makes the quantification of an independent uncertainty very difficult). But a reconsideration based on the referee comment led us to the conclusion that it is preferable to estimate the excreta term (and its uncertainty) separately based on the measured feed digestibility. A detailed description of the modified calculation was added to Sect. 2.4.3. In this way the closure of the animal C budget can serve as a really independent (and illustrative) consistency test. The modified calculations led to minor changes in the NECB budget components. The previously almost identical NECB results differ now somewhat more but still within the uncertainty range.

Figure 4 and Tables S1-S2 were adjusted accordingly.

- 2) *It would be helpful if the actual contribution and uncertainties of the components of the GHG balance would be provided in the supplementary material (in addition to Figure 5 in the main text). It would appear that nowhere in the paper the contributions from CO₂, CH₄ and N₂O are actually summed to one total GHG budget. It is unclear to me why the authors haven't done this. This would also allow the GHG budget to be compared with that found in other studies.*

A table with the numbers of the pasture GHG fluxes and their uncertainty ranges was added in the supplementary material (Table S3). In order to better connect the GHG and the carbon fluxes, we changed the bar diagram with the GHG fluxes (new Fig. 6) from vertical to horizontal bars and now display both NECB results.

Yet we want to point out that the focus of the paper is on the carbon budget and not on the GHG budget, which is only presented for context reasons (see also comment to referee #1). For clarification of this focus for the readers, the titles of Sect. 2.5 and 3.4 were changed to 'Comparison to other pasture greenhouse gas fluxes' and 'Comparison to other greenhouse gas fluxes of the dairy cow pasture', respectively.

Other comments

Methods

The budget calculations considered only the 99 days of the year that the cows were grazing the study site (P20075, L10). However, later on the authors state that the NECB was determined for a full calendar year (P20075, L25). These statements are confusing because they seem to contradict each other. From reading a further explanation on P20078 I assume the statement in P20075, L10 only applies to cow-related C fluxes and not all budget components. If this is correct then the statement in P20075, L10 needs re-phrasing to make this clear.

Figure 1 indicating the pasture and off-pasture time over the entire year was added to the manuscript. The last sentence of Sect. 2.1, which seemed to mislead the reader (see also referee #1 commented on this issue), was revised. The last two paragraphs of Sect. 2.2 were also revised and the attributed times used to calculate the NECB components have been added to Table S2.

Section 2.4.1 about live weight increase: I couldn't follow these calculations. If cows weighted on average 640 kg (Section 2.1, L25), then a 6% increase would equal about 38.4 kg per cow over the grazing season of 99 days. Per day, per cow, this is 0.38 kg and not 0.2 kg. Did I miss something?

We added the duration (209 d = grazing season) which was used to calculate the live weight increase at the beginning of the second paragraph in Sect. 2.4.1.

I would also add here the full calculations about the implications of LW increase presented currently in P20084-L26 onwards (which requires the C content of meat which is currently missing from Section 2.4.1) so that it is dealt with in one place.

We moved the respective paragraph from the results section to Sect. 2.4.1 as suggested by the reviewer.

Results and discussion

P20090, L3-6. Can the authors give a possible explanation for this difference in NECB between your findings and these other studies/study sites?

Such an explanation would be very speculative in our opinion. We abstained from a detailed interpretation of the NECB results because, as stated in Sect. 3.4, we think that several years of measurements are necessary for budget results that would allow a meaningful comparison with other studies/sites.

P20091, L4-6. I agree that the simultaneous application of both methods is useful as a consistency check, and am impressed at the level of agreement of the two methods. However, as I understand it the two methods were not entirely independent because the estimation of $F_{C-grazing}$ (needed for Method II) was not based on actual measurements of pasture biomass removed, but instead derived indirectly from milk production (which was also used in Method I). This may be worth mentioning.

We added a corresponding statement at the end of Sect. 3.3. In addition, as mentioned above, we implemented a less constricted (more independent) estimation of the excretion flux, which led to a larger (yet not significant) difference between the two NECB results.

P20091. The (size and contributions from individual gases of the) GHG budget should be discussed in more details and the findings compared to other studies. I realise that the GHG budget may not have been the main focus of the paper, but if the authors choose to present the results regarding the GHG balance, they need to link them better to the existing literature. I feel it would also be worth adding a few words about the GHG balance to the abstract.

For this issue, see the answer given under Main Concerns 2 (above) and the answer to the first comment of Referee #1.

Minor comments

P20080-16. I assume the units of Ec-milk are gC head⁻¹ d⁻¹? Add 'per day' to L16.

'Per cow and day' was added in the first paragraph of Sect. 2.4.1.

Section 2.4.2 L15. Is EDM-intake in kg CM d⁻¹ head⁻¹? Add 'per head'

Added were needed.

P20083, L6. What is EKL?

EKL is the Swiss Federal Commission for Air Hygiene (Eidgenössische Kommission für Lufthygiene EKL) and was used here as a citation. We changed the sentence in Sect. 2.5 and added "... based on the report of the Swiss Federal Commission for Air Hygiene (FCAH, 2004)". The Reference was also changed to English.

Section 2.4.2 L14 Replace 'meat gain' with 'live weight gain' to match wording in Section 2.4.1
All 'meat gain' were changed to live weight gain (see Sect. 2.4 and 2.5)

Section 2.4.2 L7 Conversion factor needs reference.

A reference to IPCC 2006 Guidelines was added in Sect. 2.4.2.

Section 3.1, L13. 'The applied models' – it may be helpful here to refer back to Section 2.4.2
Reference to Sect. 2.4.2 was added in Sect. 3.1.

P20085, L13. This proportion of C excreted in dung was actually not determined by Rutledge et al, 2014 but by Woodward, S.L., Waghorn, G.C., Bryant, M.A., Benton, A., 2012. Can diverse pasture mixtures reduce nitrogen losses? In: Jacobs, J. (Ed.), Proceedings of the 5th Australasian Dairy Science Symposium, Melbourne, pp. 463-464.

Reference was changed as suggested.

P20085 - L25. 'components of higher magnitude' – maybe just say 'larger budget components'
Changed as suggested

P20085, L28. I got $-189 \text{ gC m}^{-2} \text{ yr}^{-1}$ when I add up all exports in Table S2 for NECBtot, not -245 ?

The referee calculated correctly. However, the adjustments of some budget components needed a recalculation of this number anyway. The total export of the new calculation is $-202 \text{ gC m}^{-2} \text{ yr}^{-1}$.

P20087, L22 If total losses in the NECBtot method were indeed -189 , the contribution of $F_{\text{C-CH}_4, \text{cows}}$ to these losses was 9%, not 7%

The number after the adjustment of some budget calculation is now 8% ($-17/-202$).

P20087 last paragraphs of Section 3.2. It may be worth stating that even if small losses of $\sim 10 \text{ gC m}^{-2} \text{ yr}^{-1}$ were added to the calculated NECB's, the conclusion wouldn't change (i.e. the would remain C neutral)

A sentence was added at the end of the last paragraph of Sect. 3.2, stating that even if leaching and erosion would be considered, the conclusions on the budget result do not change.

P20089, L18-19. You may want to add a reference to Kirschbaum MUF, Rutledge S, Kuijper IA, Mudge PL, Puche N, Wall AM, et al. Modelling carbon and water exchange of a grazed pasture in New Zealand constrained by eddy covariance measurements. Science of the Total Environment. 2015; 512–513(0):273-86. They also concluded the risk of underestimating cow respiration losses if grazing events are not captured completely.

Reference was added.

P20090, L26-27. Awkward phrasing. Maybe say "... carbon-neutral budget, both methods resulted in considerable uncertainties, with slightly lower uncertainties when using the NECBtot approach (system...."

Was changed accordingly in Sect. 4.

Determination of the carbon budget of a pasture: effect of system boundaries and flux uncertainties

R. Felber^{1,2}, D. Bretscher¹, A. Mürner³, A. Neftel¹, and C. Ammann¹

¹Agroscope Research Station, Climate and Air Pollution, Zürich, Switzerland

²ETH Zürich, Institute of Agricultural Sciences, Zürich, Switzerland

³Agroscope Research Station, Milk and Meat Production, Posieux, Switzerland

Abstract

Carbon (C) sequestration in the soil is considered as a potential important mechanism to mitigate greenhouse gas (GHG) emissions of the agricultural sector. It can be quantified by the net ecosystem carbon budget (NECB) describing the change of soil C as the sum of all relevant import and export fluxes. NECB was investigated here in detail for an intensively grazed dairy pasture in Switzerland. Two budget approaches with different system boundaries were applied: NECB_{tot} for system boundaries including the grazing cows and NECB_{past} for system boundaries excluding the cows. CO₂ and CH₄ exchange induced by soil/vegetation processes as well as direct emissions by the animals were derived from eddy covariance measurements. Other C fluxes were either measured (milk yield, concentrate feeding) or derived based on animal performance data (intake, excreta). For the investigated year, both approaches resulted in a small **near-neutral C budget: NECB_{tot} -27 ± 62 g C m⁻² yr⁻¹ and NECB_{past} 23 ± 76 g C m⁻² yr⁻¹**. The considerable uncertainties, depending on the approach, were mainly due to errors in the CO₂ exchange or in the animal related fluxes. The **comparison of the NECB results with the annual exchange of other GHG** revealed CH₄ emissions from the cows to be the major contributor **in terms of CO₂-equivalents**, but with much lower uncertainty compared to NECB. Although only one year of data limit the representativeness of the carbon budget results, they demonstrated the important contribution of the non-CO₂ fluxes depending on the chosen system boundaries and the effect of their propagated uncertainty in an exemplary way. The simultaneous application and comparison of both NECB approaches provides a useful

Gelöscht: non-significant C loss: NECB_{tot} -13 ± 61 g C m⁻² yr⁻¹ and NECB_{past} -17 ± 81

Gelöscht: associated GHG budget

32 consistency check for the carbon budget determination and can help to identify and eliminate
33 systematic errors.

34 **1 Introduction**

35 The agricultural sector is the third major contributor of anthropogenic induced greenhouse gas
36 (GHG) emissions and accounts for 14% of global GHG emissions (IPCC, 2014). Depending on
37 the country and the agricultural production system, agriculture can account for more than 50%
38 of total national GHG emissions (UNFCCC, 2014). Whereas agricultural activities mainly lead
39 to emissions of CH₄ and N₂O, agricultural land potentially can be either a source or a sink for
40 atmospheric CO₂ (Tubiello et al., 2015) by changing the carbon (C) storage in the soil. Grazing
41 land management, cropland management and restoration of organic soils are considered as the
42 most cost-effective mitigation options for the agriculture sector (IPCC, 2014), and carbon
43 sequestration, i.e., the increase of soil organic carbon (SOC), in grassland is seen as the key
44 issue (Soussana et al., 2010).

45 To fully account for the GHG effect of an agricultural system, the exchange of all relevant
46 GHGs needs to be determined. Whereas N₂O and CH₄ emissions can be directly measured, the
47 carbon source or sink of an agricultural ecosystem is more difficult to quantify. Changes in
48 SOC can be measured from repeated soil sampling over longer time periods (several years) but
49 are difficult to detect for shorter-term assessments because of the generally large background
50 and high spatial variability (Smith, 2004). For shorter (e.g., annual) timescales the net
51 ecosystem carbon balance (NECB) approach can be used (Chapin et al., 2006). It determines
52 the carbon storage change as the net budget of all C containing import and export fluxes to/from
53 the ecosystem. In natural ecosystems the NECB is mainly determined by the net CO₂ exchange
54 with the atmosphere including uptake by photosynthesis and release by plant and soil
55 respiration. In managed agricultural grasslands additional non-CO₂ carbon imports (e.g.,
56 through manure application) and exports (e.g., through biomass removal) in liquid, solid, or
57 gaseous form are important contributions for the determination of NECB. The NECB of a
58 grazed pasture is also strongly influenced by the C cycling in the animals.

59 While the experimental determination of ecosystem CO₂ exchange and its problems and
60 uncertainties has been investigated in many publications, only few studies have experimentally
61 assessed the NECB of pasture ecosystems and its quality up to now (e.g., Soussana et al., 2007;
62 Mudge et al., 2011; Rutledge et al., 2015). The GHG exchange of agricultural ecosystems is

63 generally determined and described as flux per surface area, whereas the emission of CH₄ and
64 N₂O of livestock production is often measured or calculated per animal, based on mass or
65 energy budgets as used in the IPCC approaches (IPCC, 2006) followed by up-scaling to national
66 or global GHG emission inventories.

67 Felber et al. (2015, 2016) showed how CH₄ and CO₂ fluxes over a pasture with grazing dairy
68 cows can be determined using the eddy covariance (EC) technique. Here we combine and
69 complement those measurements with the non-gaseous C fluxes to determine the annual NECB
70 of the dairy pasture. Two budget approaches with different system boundaries are applied and
71 their advantages and practical limitations (necessary input data and quality) are discussed. To
72 link the NECB and its uncertainty to the full GHG budget of the pasture system, it is compared
73 to the emissions of CH₄ and N₂O in terms of CO₂-equivalents.

74 2 Material and methods

75 2.1 Study site

76 The study site is the same as described in Felber et al. (2015, 2016). The experiment was
77 conducted in 2013 on a pasture field of 3.6 ha at the Agroscope research farm near Posieux on
78 the western Swiss plateau (46°46'04", N 7°06'28" E) at an altitude of 642 m above sea level
79 with normal annual rain amount of 1075 mm and temperature of 8.9 °C (MeteoSchweiz, 2014).

80 The pasture vegetation consists of a grass-clover mixture (mainly *Lolium perenne* and *Trifolium*
81 *repens*). It was last renovated in August 2007 and has since then been used as pasture for various
82 livestock (dairy, beef cattle, calves). On average the pasture was fertilized with 120 kg nitrogen
83 (N) per year in addition to the livestock excreta. The soil is classified as stagnic Anthrosol with
84 a loam texture and a C content of the upper soil layer (0 to 20 cm) of 29 g kg⁻¹.

85 During the grazing season (9 April–4 November 2013) a herd of 20 Holstein and Red Holstein
86 x Simmental crossbred dairy cows with a mean live weight of 640 ± 70 (SD) kg was managed
87 in a rotational grazing system during day and night. Twice per day the cows left the pasture for
88 milking in the barn ([see Fig. 1](#)) where they were also offered concentrate supplement according
89 to their milk production level. Cow positions were recorded by GPS devices to determine
90 pasture presence time on 30 min basis. The pasture was divided into six paddocks of equal size
91 and were grazed for one to three days depending on herbage height. Grazing was interrupted [in](#)
92 [some cases](#) due to unfavorable environmental conditions (risk of frost, too high temperatures,

Gelöscht: for single days

94 or too wet soil conditions). The fodder provided by the 3.6 ha study field was not sufficient for
95 continuous grazing of the herd during the entire season. Therefore, additional pasture was
96 needed for certain periods. However, the budget calculations applied here only consider carbon
97 fluxes related to the specific study pasture.

Gelöscht: During these times the cows were feed in the barn.

Gelöscht: the continuous feeding

Gelöscht: the time periods (99 days in total) when the cows grazed on the

99 2.2 Carbon budget concept

100 In agricultural ecosystems the change of the SOC stock over time represents a sink or source of
101 atmospheric CO₂. The effect of changes in living plant biomass can often be neglected (due to
102 the lack of woody biomass accumulation) when looking at full years including a complete
103 vegetation season or longer periods. With the NECB approach, the SOC stock change is
104 determined by closing the carbon mass budget of the ecosystem:

$$106 \frac{\Delta \text{SOC}}{\Delta t \cdot A} \approx \text{NECB} \equiv \sum_x F_{C-x} \quad (1)$$

107 where A is the surface area under consideration and F_{C-x} are all relevant carbon mass exchange
108 fluxes through the ecosystem boundaries by various pathways x (in gaseous, liquid, or solid
109 form). Here we follow the ecological sign convention, in which positive flux and NECB values
110 indicate a C uptake by the system and negative values a C loss from the system (Chapin et al.,
111 2006). In the present study we determined the NECB for a full calendar year. This is a common
112 procedure in temperate and boreal regions of the northern hemisphere with start/end in the
113 winter season to avoid effects of carbon storage in living plant biomass and of uncertainties in
114 the attribution of management related fluxes.

116 For dairy pasture systems, the choice of system boundaries for the determination of the NECB
117 is not as obvious as for other ecosystems, because of the (temporal) presence of the grazing
118 animals. Two approaches with different boundaries were chosen here to estimate the change of
119 SOC stock expressed as NECB (Fig. 2). In these budget calculations, we neglect C loss due to
120 leaching and erosion because they could not be measured in this experiment, and are assumed
121 to be very small compared to the major fluxes.

Gelöscht: 1

122 The first approach (Fig. 2a) deduces the carbon budget from all relevant C fluxes of the *total*
123 system including the grazing animals (NECB_{tot}) similar as applied by Soussana et al. (2007)
124 and Rutledge et al. (2015). In this approach animal respiration and products count as C exports,

Gelöscht: 1

131 beside other C losses from the pasture. Since the cows had to leave the pasture twice a day for
 132 milking in the barn, this system also comprises cow fluxes during these off-pasture phases.
 133 $NECB_{tot}$ is determined as:

$$134 \quad NECB_{tot} = F_{C-CO_2,tot} + F_{C-CH_4,soil} + F_{C-CH_4,cows} + F_{C-fertil} + F_{C-products} +$$

$$135 \quad + F_{C-feed,off} + F_{C-resp,off} + F_{C-excreta,off} \quad (2)$$

137 where $F_{C-CO_2,tot}$ is the net CO_2 exchange of the total grazing system including cow respiration
 138 (during their presence on the pasture), $F_{C-CH_4,soil}$ is the CH_4 uptake or loss from the soil
 139 including deposited dung on the pasture and $F_{C-CH_4,cows}$ is the CH_4 emission from enteric
 140 fermentation, $F_{C-fertil}$ is the imported C in organic fertilizers, and $F_{C-products}$ is the C exported in
 141 animal products milk and meat (live weight gain). It has to be noted, that the C stock change in
 142 animal live weight is treated here as an export flux and thus it is not part of the resulting net
 143 ecosystem budget. For the time share the cows spent off-pasture, the intake of supplementary
 144 feed ($F_{C-feed,off}$) as well as the loss by animal respiration ($F_{C-resp,off}$) and excreta ($F_{C-excreta,off}$) are
 145 considered.

147 The system boundaries of the second approach ($NECB_{past}$, Fig. 2b) comprise only the pasture
 148 (soil and vegetation); the cows are outside the system but contribute to the budget by exporting
 149 forage and importing excreta. This approach has been applied e.g. by Skinner (2008). $NECB_{past}$
 150 is determined as:

$$151 \quad NECB_{past} = F_{C-CO_2,past} + F_{C-CH_4,soil} + F_{C-fertil} + F_{C-grazing} + F_{C-excreta,past} \quad (3)$$

153 where $F_{C-CO_2,past}$ is the net CO_2 exchange of the pasture without cow respiration, $F_{C-grazing}$ is
 154 grass biomass C removed by grazing, and $F_{C-excreta,past}$ is the C import by excreta on the pasture.
 155 The individual flux terms contributing to the budgets in Eqs. (2) and (3) act for different time
 156 periods; fluxes related to the pasture field act for the full year (i.e., $F_{C-CO_2,tot}$, $F_{C-CO_2,past}$,
 157 $F_{C-CH_4,soil}$, $F_{C-fertil}$), while the cow related fluxes act only for the time periods associated with
 158 grazing on the investigated pasture (including the adjacent milking time) and were calculated
 159 as the attributed temporal fraction. In the study year the cows grazed for a total of 99 days on
 160 the investigated pasture (hereafter referred to as 'total grazing days', see Fig. 1) applying to
 161

Gelöscht: 1

Gelöscht: entire

164 $F_{C-CH_4,cows}$, $F_{C-grazing}$, $F_{C-products}$ and $F_{C-feed,off}$ (see Table S2 in the Supplement). Even on these
 165 grazing days, the cows had to leave the pasture and go to the barn twice a day for milking. The
 166 average time for one milking event (including the time for moving between pasture and barn,
 167 indicated by the GPS position) was 3.1 h. Thus the effective time spent on the investigated
 168 pasture was reduced to 73.1 days (hereafter referred to as 'effective pasture time'), applying to
 169 $F_{C-excreta,past}$. The complementary 'off-pasture time' of 25.9 days applies to $F_{C-resp,off}$ and
 170 $F_{C-excreta,off}$.

171 Annual animal related C fluxes were aggregated from average daily animal exchange rates E_{C-x}
 172 (in units of g C head⁻¹ d⁻¹) over the mean number of animals ($n_{cow} = 19.7$) and allocated to the
 173 total pasture area ($A = 36000$ m²):

$$175 F_{C-x} = E_{C-x} \cdot \frac{n_{cow}}{A} \cdot T_x \quad (4)$$

176 where T_x is the accountable time period for the flux F_{C-x} as described above. The sign may
 177 change between F_{C-x} and E_{C-x} depending on the examined system boundaries. The uncertainty
 178 of the NECB was calculated by Gaussian error propagation of the individual uncertainties of
 179 the fluxes contributing to the budget. A detailed description of the individual error
 180 determination can be found in the Supplement, if not specified in the main text.

183 2.3 Determination of area related fluxes

184 2.3.1 CO₂ fluxes

185 Net CO₂ exchange of the pasture was determined as net ecosystem exchange (NEE) using the
 186 EC technique as described in Felber et al. (2016). NEE was determined under the
 187 micrometeorological sign convention (negative for downward/uptake, positive for
 188 upward/loss), thus F_{C-CO_2} used here has the opposite sign of NEE. Annual F_{C-CO_2} was calculated
 189 either from gap filled flux data including cases with cow respiration ($F_{C-CO_2,tot}$) or only from
 190 data without cow respiration contribution ($F_{C-CO_2,past}$). The selection of $F_{C-CO_2,past}$ data was
 191 achieved using GPS cow position information and the flux footprint distribution. The
 192 uncertainties of the annual CO₂ fluxes were determined from combined random and systematic
 193 uncertainties. Random uncertainty was estimated from varying the input data before gap filling

Gelöscht: and $F_{C-products}$

Gelöscht: $F_{C-grazing}$ and

Gelöscht: , was determined by the sum of all 30 min intervals during which the cows were on the pasture for the entire interval (indicated by the GPS positions) plus one-half of the intervals which were attributed to moving between pasture and barn

Gelöscht: The mean time for one milking event (including the time for moving between pasture and barn) was 3.1 h, thus the total time spent outside of the pasture was 25.9 days (hereafter referred to as 'off-pasture time') applying to $F_{C-feed,off}$.

Gelöscht: two NEEs

207 (adding random noise or additional gaps) and systematic uncertainty was estimated from
208 varying the applied selection threshold for low turbulence conditions (u^* filtering). The
209 difference between the $F_{C-CO_2,tot}$ and $F_{C-CO_2,past}$ corresponds to the area related cow respiration
210 flux, which could be converted to an average cow respiration $E_{C-resp} = 4.6 \text{ kg C head}^{-1} \text{ d}^{-1}$ (for
211 details see Felber et al., 2016). They estimated different uncertainties for cow respiration, here
212 we use the rather conservative uncertainty of $\pm 1.6 \text{ kg C head}^{-1} \text{ d}^{-1}$.

213

214 **2.3.2 CH₄ fluxes**

215 CH₄ emissions of the pasture soil and surface ($F_{C-CH_4,soil}$) were determined from EC data
216 without direct cow influence (for details see Felber et al., 2015). Flux intervals were selected
217 based on GPS data of cow positions. Small generally positive fluxes in a typical range of 0 to
218 $15 \text{ nmol m}^{-2} \text{ s}^{-1}$ were found. Even though some temporal variations in median diurnal and
219 seasonal cycles were observed, a constant soil/surface CH₄ emission over the year of $4 \pm 3 \text{ nmol}$
220 $\text{m}^{-2} \text{ s}^{-1}$ is assumed for the budget calculation. This value integrates emissions induced from
221 cow excreta and CH₄ sources and sinks of the soil. The uncertainty of the pasture CH₄ fluxes
222 was estimated from the uncertainty range of $\pm 50\%$ covering the temporal variation of weekly
223 medians.

224 Felber et al. (2015) also determined in-situ animal CH₄ emissions from EC data. Cow CH₄
225 fluxes were corrected by the weights of individual cow position contributions to convert area
226 integrated data into emissions per animal. The average animal CH₄ emission amounted to 423
227 $\pm 24 \text{ g CH}_4 \text{ head}^{-1} \text{ d}^{-1}$. This seasonal average animal exchange rate was converted to a carbon
228 exchange and back to a corresponding area related flux $F_{C-CH_4,cows}$ using Eq. (4) for the
229 timespan of total grazing days.

230

231 **2.3.3 Fertilizer application**

232 In the study year, two fertilizer applications took place: Before the beginning of the grazing
233 season (6 March) cattle slurry was applied by trailing hose at a rate of $43 \text{ m}^3 \text{ ha}^{-1}$. Dry organic
234 matter of the slurry was determined according to VDLUFA (2000) recommendations and the
235 C content of the dry matter of 52% was adopted from previous comparisons with elemental
236 analysis for similar slurry. The uncertainty of the slurry C import was assumed to be 17%
237 (Ammann et al., 2009). Nitrogen applied by the slurry amounted to 70 kg N ha^{-1} . An additional

238 50 kg N ha⁻¹ was applied as urea in June. Due to the C/N ratio of 1/2 in urea, this corresponds
239 to a very small C import.

240 2.4 Determination of animal related fluxes

241 The animal related carbon fluxes can be examined under the aspect of the animal C budget (in
242 units g C head⁻¹ d⁻¹) balancing gain with loss and storage terms:

$$243 E_{C\text{-intake}} = E_{C\text{-resp}} + E_{C\text{-CH}_4,\text{cow}} + E_{C\text{-milk}} + E_{C\text{-meat}} + E_{C\text{-excreta}} \quad (5)$$

244 Ingested C in feed ($E_{C\text{-intake}} = E_{C\text{-grazing}} + E_{C\text{-feed,off}}$) is partitioned into respired CO₂ ($E_{C\text{-resp}}$),
245 loss of CH₄ by enteric fermentation ($E_{C\text{-CH}_4,\text{cow}}$), the C in milk ($E_{C\text{-milk}}$) and live weight gain
246 ($E_{C\text{-meat}}$), and the C in the excreta ($E_{C\text{-excreta}}$). The determination of $E_{C\text{-resp}}$ and $E_{C\text{-CH}_4,\text{cow}}$ was
247 already described in the previous sections. The quantification of the other terms is explained in
248 the following.

249 2.4.1 Products

250 The animal production terms $E_{C\text{-milk}}$ and $E_{C\text{-meat}}$ were estimated from monitored daily milk
251 yield and live weights measured after milking. Milk was sampled individually on one day per
252 week and analyzed for fat, protein and lactose content. Energy-corrected milk yields (ECM)
253 adjusted to a gross energy content of 3.14 MJ kg⁻¹ were calculated from daily milk yields
254 according to Arrigo et al. (1999) using fat, protein and lactose contents. The C content was
255 calculated using an energy to C content ratio of 21 ± 1.9 g C MJ⁻¹ (for details see Sect. S1.2).
256 Using data from the entire grazing period an average milk C output per cow and day ($E_{C\text{-milk}}$)
257 was derived with an uncertainty of 9%.

258 The live weight (LW) of the dairy cows slightly increased by around 6% over the entire grazing
259 season of 209 days corresponding to an average daily increase of 0.2 kg LW head⁻¹ d⁻¹.
260 Applying the value of 0.14 kg C (kg fresh meat)⁻¹ (Avila, 2006) the C incorporated into meat
261 results in 0.025 kg C head⁻¹ d⁻¹, which is less than 2% of milk C yield and thus negligible here.
262 Even for beef cattle, $E_{C\text{-meat}}$ is generally small (Allard et al., 2007) and thus sometimes
263 neglected in carbon budget calculations (e.g., Soussana et al., 2007).

264 $F_{C\text{-products}}$ was calculated from $E_{C\text{-milk}}$ by Eq. (4) using the number of total grazing days.

265

Gelöscht: meat

Gelöscht: and assuming linear relationship for these components, when no measurements were available

Gelöscht: ECM was adjusted to a gross energy content of 3.14 MJ kg⁻¹ (Arrigo et al., 1999) and t

Gelöscht: (determined in previous experiments by Münger, 1997).

Gelöscht: only

Gelöscht: (on average 0.2 kg LW d⁻¹).

[1] verschoben

Gelöscht: The live weight gain of the cows was around 0.2 kg d⁻¹ (around 6% increase over the grazing season).

Gelöscht: This corresponds to a C accumulation in meat of <0.05 kg C head⁻¹ d⁻¹. Thus for dairy cows $E_{C\text{-meat}}$ was assumed to be negligible compared to $E_{C\text{-milk}}$ (Soussana et al., 2007).¶

Gelöscht: The uncertainty of $F_{C\text{-products}}$ was estimated from the combination of uncertainties of the ECM and the ratio between milk gross energy and C content. The latter effect was dominating and led to a total uncertainty of 10% for $F_{C\text{-products}}$.

289 **2.4.2 Feed intake**

290 The dry matter (DM) feed of the cows was estimated using two different approaches: i) by the
291 Tier 2 model given in the IPCC Guidelines (IPCC, 2006) and ii) based on the Swiss feeding
292 recommendations and nutrition tables for ruminants (Arrigo et al., 1999). The former approach
293 estimates gross energy intake of the cows from net energy requirements for maintenance,
294 activity (grazing), and production (milk yield). The gross energy intake is then converted to
295 DM intake using the default factor of 18.45 MJ (kg DM)⁻¹ (IPCC, 2006). The second model
296 uses the following equations (Eq. 6a for primiparous and Eq. 6b for multiparous cows):

297
298 $E_{\text{DM-intake}} = 0.33 \cdot \text{ECM} + 0.29 \cdot \text{lacW} - 0.0047 \cdot \text{lacW}^2 + 6.0$ (6a)

299 $E_{\text{DM-intake}} = 0.33 \cdot \text{ECM} + 0.17 \cdot \text{lacW} - 0.0025 \cdot \text{lacW}^2 + 8.8$ (6b)

300

301 where ECM is in kg head⁻¹ d⁻¹ and lacW is the actual lactation week of the cow. Additional
302 intake corrections were applied for deviations from standard live weight (600 kg and 650 kg
303 LW for Eqs. 6a/b, respectively) and standard annual milk production (6500 kg and 7500 kg
304 respectively). Estimated $E_{\text{DM-intake}}$ was i) 18.8 kg DM head⁻¹ d⁻¹ and ii) 18.5 kg DM head⁻¹ d⁻¹.

305 We used 18.5 ± 2.7 kg DM head⁻¹ d⁻¹ for the further calculations because this value is based on
306 the actual production state of the cows in contrast to the value from approach i), which is based
307 on the IPCC standard parameterization.

308 Besides the grazing on the pasture, the cows were offered a minor amount of supplement
309 feeding (concentrates) depending on individual milk production level of each cow. Daily
310 concentrate intake was recorded for each cow, on average it amounted to 1.3 ± 0.2 kg DM
311 head⁻¹ d⁻¹ over the grazing period.

312 Carbon (and N) content of pasture forage and concentrates were measured by dry combustion
313 (VDLUFA, 2000) of weekly sampled pasture forage and from periodically analyzed
314 concentrate samples ($n = 6$ over the grazing period). A carbon content of 433 ± 9 g C (kg DM)⁻¹
315 was measured for pasture forage and 430 ± 9 g C (kg DM)⁻¹ for the concentrates. With these
316 information the total average daily carbon intake ($E_{\text{C-intake}}$) per cow was derived. $F_{\text{C-feed,off}}$ was

317 calculated from the daily concentrate intake alone. $F_{\text{C-grazing}}$ was calculated for the total grazing
318 days from the difference between $E_{\text{C-intake}}$ and $E_{\text{C-feed,off}}$ with an uncertainty of ±16% (see Table
319 S2).

Gelöscht: 1.5

Gelöscht: 426

Gelöscht: 429

Gelöscht: The uncertainty (6%) of $F_{\text{C-feed,off}}$ was derived from the combined uncertainties of the DM content and the C content determination.

Gelöscht: 15

327 2.4.3 Excreta

328 Excreta output could not be measured directly in this study, and it is generally difficult to
329 measure for grazing animals. But the ratio of $E_{C\text{-excreta}}$ relative to the animal intake was
330 estimated from the analysis of the feed digestibility. For this purpose, 50 grass samples taken
331 during the grazing season were analyzed by Tilley and Terry (1963). This resulted in an average
332 feed organic matter digestibility of 0.72 with an uncertainty range of ± 0.07 . Because the carbon
333 content in the excreted dung (c. 50% of organic matter, see e.g., Pettygrove et al., 2010) is
334 higher than in the feed (43% of organic matter acc. to sample analysis) the effective carbon
335 digestibility reduces to 0.68. Accordingly $E_{C\text{-excreta}}$ was estimated as 32 ± 8 % of the animal
336 carbon intake. $F_{C\text{-excreta,past}}$ and $F_{C\text{-excreta,off}}$ were calculated from $E_{C\text{-excreta}}$ for the effective
337 pasture time and the off-pasture time, respectively, using Eq. (4).

338

339 2.5 Comparison to other pasture greenhouse gas fluxes

340 For a quantitative comparison of the NECB to the other relevant GHG fluxes of the pasture
341 system, the CH_4 and N_2O emissions were converted to CO_2 equivalents based on their of global
342 warming potential (GWP). Here we used the 100 year GWPs; 25 $\text{CO}_2\text{-eq.}$ for CH_4 and 298
343 $\text{CO}_2\text{-eq.}$ for N_2O (Solomon et al., 2007). The system boundaries were the same as for the
344 determination of the NECB_{tot} , i.e., the effects of the investigated pasture including the animals
345 during pasture days are taken into account. Correspondingly, area related fluxes are accounted
346 for the entire year, while cow related fluxes are accounted for the total pasture days (time spent
347 on the pasture plus the adjacent milking periods).

348 The average CH_4 emissions of the soil and the cow emissions were derived by EC
349 measurements as mentioned in Sect. 2.3.2 and allocated to the respective time periods.

350 Emissions of N_2O in terms of N mass were estimated according to:

351

$$352 F_{\text{N-N}_2\text{O}} = (F_{\text{N-fertil}} + F_{\text{N-resid}} + F_{\text{N-dep}}) \cdot f_1 + F_{\text{N-excreta}} \cdot f_2 \quad (8)$$

353

354 where $F_{\text{N-fertil}}$, $F_{\text{N-resid}}$ and $F_{\text{N-dep}}$ are the N inputs by fertilizers, plant residues, and atmospheric
355 deposition, and $f_1 = 0.01$ and $f_2 = 0.02$ are the default N_2O emission factors due to the respective
356 N inputs according to the IPCC guidelines (IPCC, 2006). $F_{\text{N-fertil}}$ was determined from

Gelöscht: was not measured in this study. But $E_{C\text{-excreta}}$ was estimated by closing the average cow C budget (Eq. 5). The uncertainty was estimated to 46% (resulting from the combination of uncertainties of the other budget terms but limited by plausibility considerations).

Gelöscht: pasture Greenhouse gas budget

Gelöscht: consideration of the full GHG budget of the pasture system, the NECB needs to be quantitatively related to CH_4 and N_2O emissions in terms

366 management records and the analysis of the applied slurry (see Sect. 2.3.3) and amounted to
367 120 kg N ha⁻¹ in total for the study year. The amount of N deposited from the atmosphere was
368 estimated to 25 kg N ha⁻¹ yr⁻¹ based on [the report of the Swiss Federal Commission for Air](#)
369 [Hygiene](#) (FCAH, 2014).

370 The other two terms in Eq. (8), were estimated with the help of the animal N balance, which
371 can be formulated in a similar way as the animal carbon balance in Eq. (5) but without gaseous
372 pathways:

$$374 E_{N\text{-intake}} = E_{N\text{-milk}} + E_{N\text{-meat}} + E_{N\text{-excreta}} \quad (9)$$

375
376 $E_{N\text{-intake}}$ is the uptake of N in the feed and the average value was quantified based on the average
377 N content of pasture forage (28 g N (kg DM)⁻¹) and concentrates (17 g N (kg DM)⁻¹). The
378 intake of the cow is portioned into N in milk ($E_{N\text{-milk}}$), [live weight gain](#) ($E_{N\text{-meat}}$), and excreta
379 ($E_{N\text{-excreta}}$). Average milk N output ($E_{N\text{-milk}}$) was determined from the mean ECM yield (22.7
380 kg head⁻¹ d⁻¹) and associated measured protein contents ranging from 2.8 to 4.5% and a protein-
381 to-N conversion factor of 6.38 (IPCC, 2006). Nitrogen accumulation in meat due to weight gain
382 (see e.g., Estermann et al., 2002) was very small and thus assumed negligible (like for C, see
383 Sect. 2.4.1). $E_{N\text{-excreta}}$ was estimated by closing the N balance (Eq. 9) and was used to calculate
384 $F_{N\text{-excreta}}$ in analogy to Eq. (4) for the effective pasture time resulting in a value of 152 kg N
385 ha⁻¹ yr⁻¹.

386 Nitrogen input from plant residues $F_{N\text{-resid}} = 51$ kg N ha⁻¹ yr⁻¹ was estimated as 25% of the
387 livestock N intake during the grazing period based on Walther et al. (1994) and AGRIDEA
388 (2007).

389 3 Results and discussion

390 3.1 Carbon budget of the dairy cows

391 Animal C budget considerations serve to estimate, constrain or validate animal related C fluxes
392 that contribute to the pasture system NECB. Results derived for the mean daily C budget for
393 the cows used in this study are shown in Fig. 3 together with the N budget (detailed numbers
394 can be found in Table S1). The values represent averages over all cows in the herd and over the
395 entire grazing season. The average cow needed a daily feed intake of 18.5 kg DM corresponding

Gelöscht: meat

Gelöscht: 2

398 to 8.0 kg C. The determination of the feed intake was a very important factor for the assessment
399 of the cow budget. Because in-situ determination of forage intake during grazing is challenging
400 (Undi et al., 2008), the total feed intake was calculated based on the net energy requirements of
401 the animals, which in turn were based on the actual animal performance (milk yield, live
402 weight). The applied models (Sect. 2.4.2) showed only a small difference of 0.3 kg DM head⁻¹
403 d⁻¹. Gibb et al. (2007) reported intake values for grazing dairy cows between 25 and 30 g DM
404 (kg LW)⁻¹. For the live weight of the cows in this study, this would result in intake rates of 16
405 and 18 kg DM head⁻¹ d⁻¹, which is within the estimated uncertainty range (± 2.7 kg DM head⁻¹
406 d⁻¹) of our result.

407 Of the total C intake the largest share (57%) was emitted as CO₂ and a much smaller part (4%)
408 as CH₄. A considerable amount (19%) of the C intake was processed into the milk and 32%
409 was released as excreta. The animal carbon budget shows an imbalance of 12% (see Table S1),
410 which reflects the overall budget uncertainty. Most of C was lost by respiration, which also has
411 the largest uncertainty. The value was determined from EC measurements and was found to be
412 at the upper range of animal respiration rates for dairy cows reported in the literature (see Felber
413 et al., 2016 and references therein). In contrast to the carbon budget, the largest part of the N
414 intake (75%) was excreted in urine and dung.

415 The relative share of excreta C loss is very similar to the 34% share in terms of DM reported
416 by Woodward et al. (2012) for dairy cows. The resulting imbalance of the animal budget,
417 although within the range of uncertainties, may indicate that the estimated C loss due to
418 respiration tends to be overestimated. Indeed the value of 4.6 kg C head⁻¹ d⁻¹ lies in the upper
419 range of measurements with comparable cows (see Felber et al., 2016). However, Soussana et
420 al. (2010) investigating cow C budgets for cut forage, which was fed off-pasture, found that
421 56% to 59% of intake C was respired as CO₂.

423 3.2 Carbon budget of the pasture system

424 Carbon budget components and balance results for the two different NECB approaches (system
425 boundaries) used in this study are shown in Fig. 4 (detailed numbers are listed in Table S2 in
426 the Supplement). While for NECB_{tot} a small negative and for NECB_{past} a small positive value
427 was determined, both results are attributed a considerable uncertainty range and are thus not
428 significantly different from zero nor from each other. NECB_{past} with the larger uncertainty also
429 resulted from larger budget components (fluxes). A total C import of 429 g C m⁻² yr⁻¹ to the

Gelöscht: From

Gelöscht: The residual C was released as excreta (20%).

[1] nach oben: The live weight gain of the cows was around 0.2 kg d⁻¹ (around 6% increase over the grazing season). Applying the value of 0.14 kg C (kg fresh meat)⁻¹ (Avila, 2006) the C incorporated into meat results in 0.025 kg C d⁻¹, which is less than 2% of milk C yield and thus negligible here. Even for beef cattle, E_{C-meat} is generally small (Allard et al., 2007), and thus sometimes neglected in carbon budget calculations (e.g., Soussana et al., 2007). ¶

Gelöscht: The amount of C and N in the excreta was estimated by closing the animal budget, because direct measurements were not available and generally difficult for a grazing system. The amount of C in the excreta (mainly in the dung) is strongly related to the digestibility of the forage. In-vitro digestibility measurements of the forage showed that around 71% of the feed was digested (data not shown). This number has to be considered as lower limit because it does not account for the digestibility of the concentrate. Thus the 20% of C in the excreta can be considered as a reasonable estimate, although it was determined as a (small) difference from other large animal budget terms. Yet the relative share of excreta loss is considerably lower than the 34% share in terms of DM reported by Rutledge et al. (2012) for dairy cows. This discrepancy

Gelöscht: may be

Gelöscht: present C cow budgets in g C m⁻² yr⁻¹ for cut forage that is fed off-pasture. They also found that 56% to 59% of intake C is

Gelöscht: 3

Gelöscht: Very similar, slightly negative values were determined for NECB_{tot} and NECB_{past}. Yet both values

Gelöscht: of higher magnitude

Gelöscht: 389

464 pasture (soil/vegetation ecosystem) was balanced by a total C loss of $-406 \text{ g C m}^{-2} \text{ yr}^{-1}$. For
 465 the NECB_{tot} approach, total import ($176 \text{ g C m}^{-2} \text{ yr}^{-1}$) and export ($-202 \text{ g C m}^{-2} \text{ yr}^{-1}$) were less
 466 than half as large (it has to be noted that in this consideration the annual net CO_2 exchange is
 467 used, not the gross exchange). This difference is due to the predominantly 'internal' processing
 468 of the biomass in the NECB_{tot} system. Accordingly, the largest budget term in the NECB_{tot}
 469 approach was the milk export ($F_{\text{C-products}} = -82 \text{ g C m}^{-2} \text{ yr}^{-1}$), while the largest term in the
 470 $\text{NECB}_{\text{past}}$ approach, the biomass export by grazing ($F_{\text{C-grazing}} = -404 \text{ g C m}^{-2} \text{ yr}^{-1}$), was five
 471 times larger. Additionally, combining the C lost as respired CO_2 when the cows were off-
 472 pasture and the net C imported as CO_2 into the system resulted in a zero-sum situation for the
 473 CO_2 exchange in the NECB_{tot} approach, but was the main contributor to the NECB_{tot}
 474 uncertainty. As discussed in detail in Felber et al. (2016), the difference in the net CO_2 exchange
 475 between the two approaches corresponds to the (annually averaged) effect of cow respiration
 476 while on the pasture. Although this annual cow respiration flux ($180 \text{ g C m}^{-2} \text{ yr}^{-1}$) is typically
 477 much lower than the respiration of the pasture soil/vegetation (J r me et al., 2014), it is larger
 478 than many other carbon budget terms and thus very important for the NECB quantification.
 479 The time that the cows spent each day in the barn for milking represents an important
 480 'disturbance' of the NECB_{tot} . The sum of the three specific off-pasture fluxes ($F_{\text{C-feed,off}}$,
 481 $F_{\text{C-resp,off}}$, $F_{\text{C-excreta,off}}$) results in a net off-pasture carbon loss of $-71 \text{ g C m}^{-2} \text{ yr}^{-1}$. The
 482 relatively small C import due to concentrate feeding only partially balanced the loss through
 483 animal respiration and excreta.
 484 While the resulting NECB values for a single year cannot be considered as fully representative
 485 for the site nor for pasture systems in general, they show the contribution of different C fluxes
 486 to the total budget and the effect of their (propagated) uncertainty in an exemplary way. As
 487 shown in Fig. 4, the resulting uncertainty of $\text{NECB}_{\text{past}}$ ($\pm 76 \text{ g C m}^{-2} \text{ yr}^{-1}$) was larger than for
 488 NECB_{tot} ($\pm 62 \text{ g C m}^{-2} \text{ yr}^{-1}$). These uncertainties are comparable to the uncertainty ranges
 489 reported by Rutledge et al. (2015) for annual NECB_{tot} values of a dairy pasture system (± 50 to
 490 $\pm 86 \text{ g C m}^{-2} \text{ yr}^{-1}$). Because in the present study the determination of most non-gaseous C fluxes
 491 typically have relative errors of 10 to 20%, it may be concluded that the larger absolute
 492 uncertainty of $\text{NECB}_{\text{past}}$ compared to NECB_{tot} was due to the larger individual C fluxes in this
 493 approach. This mainly applies to the largest flux $F_{\text{C-grazing}}$ that dominated the $\text{NECB}_{\text{past}}$
 494 uncertainty. The grazing intake was inferred using an empirical model based on measured milk
 495 yield, composition and animal live weight. The model uncertainty is also the main contributor

Gel scht: 245

Gel scht: 57

Gel scht: 3

Gel scht: 81

Gel scht: 61

Gel scht: from the

Gel scht: , because more

503 to the uncertainty of $F_{C\text{-grazing}}$ (see Sect. S1.1). However, direct intake measurements on the
504 pasture are difficult and would probably not yield more accurate results.

Gelöscht: (see Sect. 3.1)

505 The largest uncertainty contribution in the $NECB_{tot}$ approach was due to the CO_2 exchange flux,
506 although the magnitude of this term was not very large. The uncertainty of F_{C-CO_2} was mainly
507 determined by the gaps in the CO_2 flux measurement and although the calculation of $F_{C-CO_2,tot}$
508 is based on a larger flux dataset than $F_{C-CO_2,past}$ (for which all fluxes influenced by cows were
509 removed before gap filling) the former had a larger uncertainty (for details see Felber et al.,
510 2016). The uncertainty of the annual CO_2 exchange has an absolute rather than a relative
511 characteristic because, like the $NECB$, it is itself the result of large compensating fluxes of
512 opposite signs (Ammann et al., 2009; Felber et al., 2016).

513 Another important component in both $NECB$ approaches was the C import by slurry
514 application, which was also shown for other managed grasslands (Ammann et al., 2007;
515 Soussana et al., 2007). Only by specific sampling and analysis of the applied slurry, the relative
516 error could be limited to <20%, because the DM and thus also the C content in slurry can easily
517 vary by a factor of four.

518 Carbon lost as CH_4 from the soil was the lowest flux in both systems accounting for less than
519 1% of total C loss. While this term appears to be negligible, this is not the case for the animal
520 CH_4 emission ($F_{C-CH_4,cows}$) with a contribution of 8% to the total C loss in the $NECB_{tot}$ system.

Gelöscht: 7

521 In any case the CH_4 fluxes play a much more prominent role when compared to other GHG
522 fluxes in terms of global warming potential (cf. Sect. 3.4).

Gelöscht: in the GHG budget

523 Beside the quality and representativeness of the determination of the various C fluxes, also the
524 completeness of the budget with all relevant components is important. In the present study, the
525 loss of C through leaching and erosion were not measured, but assumed to be small compared
526 to the other C fluxes. Carbon loss through leaching in other managed grasslands was found to
527 be in the range of 5 to 11 g C m⁻² yr⁻¹ (Allard et al., 2007; Zeeman et al., 2010; Rutledge et al.,
528 2015). The loss through erosion can be assumed to be again smaller due to the flat topography
529 and the closed vegetation cover in this study. Even if a value for leaching and erosion in the
530 order of 10 g C m⁻² yr⁻¹ would be including in the budget calculation, the result of the budgets
531 would hardly be affected (i.e., the $NECB$ values would remain non-significant).

536 3.3 Applicability of the NECB approaches

537 The applicability of the two different NECB approaches depends on their specific requirements
538 and the corresponding available information for the investigated pasture system. For the
539 NECB_{past} approach the adequate determination of the relatively large CO₂ exchange flux relies
540 on the capability to distinguish between measurement intervals with and without cow influence.
541 In the present study, GPS position information of the cows in combination with a flux footprint
542 model allowed an explicit distinction of fluxes with and without cow contributions and a
543 detailed determination of times when the cows were on- or off-pasture. The separation of CO₂
544 (and CH₄) fluxes was achieved based on the actual stocking density in the flux footprint (for
545 details see Felber et al., 2015). The effect of the chosen threshold for this separation on the
546 resulting annual net CO₂ exchange is illustrated in Fig. 5. Above an average stocking rate of
547 about 3 heads ha⁻¹ in the footprint the cow respiration led to a strong change of the net CO₂
548 exchange, although these cases accounted for only about 5% of all flux data (before gap filling).
549 The required degree of detail of the position information depends on the grazing management,
550 stocking density and division of the pasture around the measurement tower. Felber et al. (2015)
551 showed that information of paddock occupation and the assumption of homogeneously
552 distributed cows within the paddock resulted in comparable results of cow CH₄ emission
553 estimates for the division used in this experiment. For pasture systems with a distinct alternation
554 of grazing and non-grazing phases (e.g., Jérôme et al., 2014) a simple time schedule based flux
555 separation, without further animal position information, may also be sufficient, but needs to be
556 tested. However, for a free-range (continuous grazing) pasture system where the cows are
557 allowed to graze all around the measurement tower at all times, the NECB_{past} approach would
558 not be feasible; pasture/soil CO₂ and CH₄ exchange ($F_{C-CO_2,past}$ and $F_{C-CH_4,soil}$) can only be
559 determined, if sufficient and defined periods without cow influence on the EC flux
560 measurement are available.

561 While the NECB_{past} approach necessitates a proper identification of pasture CO₂ fluxes without
562 cow respiration, it does not rely on off-pasture information. However, the import and export of
563 C in excreta and forage needs to be determined. Thus the NECB_{past} approach may be suitable
564 for systems with known animal performance and/or short intensive grazing phases, for which
565 the grazing export can be well constrained. The NECB_{past} approach is also suitable for grassland
566 systems with mixed management (grazing and harvest), because the harvest export can be
567 treated in the same way as grazing export (Skinner, 2008).

Gelöscht: 4

569 The $NECB_{tot}$ approach is more suitable (or even the only choice) for continuous grazing systems
 570 (e.g., Allard et al., 2007). For beef cattle pastures, the $NECB_{tot}$ approach can even be simplified,
 571 because the off-pasture phases are avoidable. While a separation of the fluxes influenced by
 572 cow respiration is not necessary in this approach, it needs to be assured that cow respiration
 573 contributions are fully represented in $NECB_{tot}$, i.e. that the cows show a temporally
 574 representative presence in the flux footprint (see Felber et al., 2015). Otherwise the annual
 575 $F_{C-CO_2,tot}$ would be affected by a systematic error as also noted by Kirschbaum et al. (2015).
 576 Generally, for any pasture system it is advisable to record as detailed information of non-
 577 gaseous C fluxes, cow positions, and grazing time schedules as possible, because the
 578 simultaneous application of both approaches and their inter-comparison provides the most
 579 defensible results for the C budget. Because the two $NECB$ approaches partly include the same
 580 fluxes (e.g., $F_{C-fertil}$) or are based on the same information (e.g., $F_{C-excreta,past}$ and $F_{C-excreta,off}$)
 581 they cannot be considered as totally independent. However, the dominant contributions and
 582 their uncertainties may be considered as statistically independent.

584 3.4 Comparison to other greenhouse gas fluxes of the dairy cow pasture

585 The $NECB$ results are compared to the effect of other GHG fluxes for the investigated pasture
 586 system in Fig. 6. In terms of CO_2 -equivalents, the CH_4 emissions from the animals contributed
 587 the most to GHG emissions, while the CH_4 emission from soil (including animal excreta) was
 588 10 times lower but not negligible. N_2O emissions contributed about one fourth to the total
 589 emissions. Due to the non-significant effect of the C storage change (near neutral $NECB$), this
 590 grazing system may not be considered as a C sink and thus a mitigation option for GHG
 591 emissions as suggested by other studies (Soussana et al., 2010; Rutledge et al., 2015).
 592 However, for a reliable assessment of the C budget of a pasture, measurements over several
 593 years are crucial. Environmental as well as management factors will have a large influence on
 594 the annual budget and determine whether a system acts as a C sink or a source. For example,
 595 plowing during restoration process of a pasture can lead to a considerable loss of C that was
 596 sequestered over several years, also affecting N_2O emissions (Ammann et al., 2013; Merbold
 597 et al., 2014).

598 In contrast to $NECB$ and CH_4 emissions, which were determined experimentally using the EC
 599 method, N_2O emissions were roughly estimated here based on modelled N cycling of the cows

Gelöscht: Greenhouse gas budget

Gelöscht: result for $NECB_{tot}$ is

Gelöscht: in the GHG budget

Gelöscht: (including cows during pasture time) shown in Fig. 5

Gelöscht: The non-significant loss of C (negative $NECB$) tends to increase the emission effect of the other GHGs. Thus,

Gelöscht: that showed pastures being a C sink

Gelöscht: The considerably large uncertainty of the $NECB$ determined the uncertainty of the GHG budget.

Gelöscht: GHG

Gelöscht: and to evaluate its C sequestration potential

Gelöscht: GHG budget and decide

Gelöscht: for GHGs

Gelöscht: $_{tot}$

616 and applied fertilizers relying on standardized emission factors. A more comprehensive picture,
617 accounting for the specific environmental conditions, could be achieved by the direct
618 determination of N₂O fluxes also using the EC method. Such measurements will be performed
619 in a follow-up project investigating the N cycling of the same pasture (NiceGras: Nitrogen
620 Cycling and Emissions of Grazing Systems).

621 4 Conclusions

622 The C storage change of a grazed pasture system was determined by two NECB approaches
623 with different system boundaries to investigate their data requirements and associated
624 uncertainties. While both approaches yielded similar results indicating a near carbon-neutral
625 budget, both methods resulted in considerable uncertainties, with slightly lower uncertainties,
626 for the NECB_{tot} approach (system boundaries including cows). Whereas the C budget results
627 for the investigated single year cannot be considered as fully representative for the longer term,
628 they demonstrate the contribution of the different C fluxes to the total budget and the effect of
629 their (propagated) uncertainty in an exemplary way. The simultaneous application and
630 comparison of both NECB approaches provides a useful consistency check for the NECB
631 determination and can help to identify and eliminate larger systematic errors. Additionally, the
632 consideration of the cow C budget can be used to quantify and check the consistency of animal
633 fluxes needed in the determination of the NECB.

634 The NECB result was compared to the effect of the other GHG fluxes from the pasture system
635 (CH₄ and N₂O normalized to CO₂-equivalents). While CH₄ emission by the cows played a very
636 minor role in the C budget, it clearly dominates the GHG emissions due to its larger greenhouse
637 warming potential. Due to its relatively low variability the CH₄ emission from enteric
638 fermentation (depending on animal state and performance) has a much lower uncertainty than
639 the NECB of the pasture field, which is the net effect of large fluxes of opposite sign.

640 While the determination of the non-gaseous fluxes in the C budget could mostly be improved
641 by more comprehensive sampling and analyses, the uncertainty due to the CO₂ exchange
642 measurements is to a certain part inevitable for the given site and management regime, because
643 the accuracy of the CO₂ exchange monitoring by EC is limited by the (micro-) meteorological
644 conditions, especially calm nighttime conditions, and by the variability of the animal presence
645 and density in the footprint. However, the uncertainty may be reduced to some degree by better
646 constrained animal C budgets (especially intake and respiration). This may be achieved by

Gelöscht: very

Gelöscht: a considerable uncertainty was estimated with a moderate advantage

Gelöscht: (and N)

Gelöscht: budget

Gelöscht: the

Gelöscht: in

Gelöscht: it

655 prolonged field measurements over several years in combination with C cycling measurements
656 on the individual animals.

657 **Acknowledgements**

658 We gratefully acknowledge the funding from the Swiss National Science Foundation (Grant
659 No. 205321_138300) and the EU-FP7 Project ECLAIRE. We wish to thank Hubert Bollhalder,
660 Roman Gubler, Veronika Wolff, Andreas Rohner, Manuel Schuler, Markus Jocher, Manuela
661 Falk, Lukas Eggerschwiler and Bernard Papaux for support with the sensors and in the field.
662

663 **References**

- 664 AGRIDEA: Pflanzen und Tiere 2008: Wirz Handbuch, AGRIDEA Lindau / Wirz Verlag,
665 Lindau/Basel, Switzerland., 2007.
- 666 Allard, V., Soussana, J.-F., Falcimagne, R., Berbigier, P., Bonnefond, J. M., Ceschia, E.,
667 D'hour, P., Hénault, C., Laville, P., Martin, C. and Pinarès-Patino, C.: The role of grazing
668 management for the net biome productivity and greenhouse gas budget (CO₂, N₂O and CH₄)
669 of semi-natural grassland, *Agric. Ecosyst. Environ.*, 121(1–2), 47–58,
670 doi:10.1016/j.agee.2006.12.004, 2007.
- 671 Ammann, C., Flechard, C. R., Leifeld, J., Neftel, A. and Fuhrer, J.: The carbon budget of newly
672 established temperate grassland depends on management intensity, *Agric. Ecosyst. Environ.*,
673 121(1–2), 5–20, doi:10.1016/j.agee.2006.12.002, 2007.
- 674 Ammann, C., Spirig, C., Leifeld, J. and Neftel, A.: Assessment of the nitrogen and carbon
675 budget of two managed temperate grassland fields, *Agric. Ecosyst. Environ.*, 133(3–4), 150–
676 162, doi:10.1016/j.agee.2009.05.006, 2009.
- 677 Ammann, C., Leifeld, J., Jocher, M., Neftel, A. and Fuhrer, J.: Effect of grassland renovation
678 on the greenhouse gas budget of an intensive forage production system, in *Advances in Animal*
679 *Biosciences*, vol. 4, p. 284, Cambridge University Press, Dublin, Ireland, 23–26 June 2013.,
680 2013.
- 681 Arrigo, Y., Chaubert, C., Daccord, R., Gagnaux, D., Gerber, H., Guidon, D., Jans, F., Kessler,
682 J., Lehmann, E., Morel, I., Münger, A., Rouel, M. and Wyss, U.: Fütterungsempfehlungen und
683 Nährwerttabellen für Wiederkäuer: das grüne Buch, 4th ed., Eidgenössische Forschungsanstalt
684 für Nutztiere, Zollikofen, Switzerland., 1999.
- 685 Avila, R.: The ecosystem models used for dose assessments in SR-Can, Swedish Nuclear Fuel
686 and Waste Management Co., Stockholm, Sweden. [online] Available from:
687 http://www.iaea.org/inis/collection/NCLCollectionStore/_Public/38/021/38021344.pdf
688 (Accessed 3 June 2015), 2006.
- 689 Chapin, F. S., Woodwell, G. M., Randerson, J. T., Rastetter, E. B., Lovett, G. M., Baldocchi,
690 D. D., Clark, D. A., Harmon, M. E., Schimel, D. S., Valentini, R., Wirth, C., Aber, J. D., Cole,

691 J. J., Goulden, M. L., Harden, J. W., Heimann, M., Howarth, R. W., Matson, P. A., McGuire,
692 A. D., Melillo, J. M., Mooney, H. A., Neff, J. C., Houghton, R. A., Pace, M. L., Ryan, M. G.,
693 Running, S. W., Sala, O. E., Schlesinger, W. H. and Schulze, E.-D.: Reconciling Carbon-cycle
694 Concepts, Terminology, and Methods, *Ecosystems*, 9(7), 1041–1050, doi:10.1007/s10021-005-
695 0105-7, 2006.

696 Estermann, B. L., Wettstein, H.-R., Sutter, F. and Kreuzer, M.: Nutrient and energy conversion
697 of grass-fed dairy and suckler beef cattle kept indoors and on high altitude pasture, *Anim. Res.*,
698 50(6), 477–494, doi:10.1051/animres:2001109, 2002.

699 FCAH: Ammoniak-Immissionen und Stickstoffeinträge (Ammonia Immissions and Nitrogen
700 Imports), Federal Commission for Air Hygiene (FCAH), Bern, Switzerland., 2014.

701 Felber, R., Münger, A., Neftel, A. and Ammann, C.: Eddy covariance methane flux
702 measurements over a grazed pasture: effect of cows as moving point sources, *Biogeosciences*,
703 12(12), 3925–3940, doi:10.5194/bg-12-3925-2015, 2015.

704 Felber, R., Neftel, A. and Ammann, C.: Discerning the cows from the pasture: Quantifying and
705 partitioning the NEE of a grazed pasture using animal position data, *Agric. For. Meteorol.*, 216,
706 37–47, doi:10.1016/j.agrformet.2015.09.018, 2016.

707 Gibb, M.: Grassland management with emphasis on grazing behaviour, in *Fresh herbage for*
708 *dairy cattle: the key to a sustainable food chain*, edited by A. Elgersma, J. Dijkstra, and S.
709 Tamminga, pp. 141–157, Springer Publishing, Dordrecht, Netherlands. [online] Available
710 from: <http://library.wur.nl/ojs/index.php/frontis/article/view/1250> (Accessed 22 April 2015),
711 2007.

712 IPCC: 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Vol 4 Ariculture,
713 Forestry and Other Land Use, Intergovernmental Panel on Climate Change, Hayama, Japan.
714 [online] Available from: <http://www.ipcc-nggip.iges.or.jp/public/2006gl/index.htm> (Accessed
715 31 March 2015), 2006.

716 IPCC: Climate Change 2014: Synthesis Report, Contribution of Working Groups I, II and III
717 to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, Core

718 Writing Team, edited by R. K. Pachauri and L. A. Meyer, Intergovernmental Panel on Climate
719 Change, Geneva, Switzerland., 2014.

720 Jérôme, E., Beckers, Y., Bodson, B., Heinesch, B., Moureaux, C. and Aubinet, M.: Impact of
721 grazing on carbon dioxide exchanges in an intensively managed Belgian grassland, *Agric.*
722 *Ecosyst. Environ.*, 194, 7–16, doi:10.1016/j.agee.2014.04.021, 2014.

723 Kirschbaum, M. U. F., Rutledge, S., Kuijper, I. A., Mudge, P. L., Puche, N., Wall, A. M.,
724 Roach, C. G., Schipper, L. A. and Campbell, D. I.: Modelling carbon and water exchange of a
725 grazed pasture in New Zealand constrained by eddy covariance measurements, *Sci. Total*
726 *Environ.*, 512–513, 273–286, doi:10.1016/j.scitotenv.2015.01.045, 2015.

727 Merbold, L., Eugster, W., Stieger, J., Zahniser, M., Nelson, D. and Buchmann, N.: Greenhouse
728 gas budget (CO₂, CH₄ and N₂O) of intensively managed grassland following restoration,
729 *Glob. Change Biol.*, 20(6), 1913–1928, doi:10.1111/gcb.12518, 2014.

730 MeteoSchweiz: GRA_norm8110.pdf, [online] Available from:
731 http://www.meteoswiss.admin.ch/files/kd/climsheet/de/GRA_norm8110.pdf (Accessed 13
732 October 2014), 2014.

733 Mudge, P. L., Wallace, D. F., Rutledge, S., Campbell, D. I., Schipper, L. A. and Hosking, C.
734 L.: Carbon balance of an intensively grazed temperate pasture in two climatically contrasting
735 years, *Agric. Ecosyst. Environ.*, 144(1), 271–280, doi:10.1016/j.agee.2011.09.003, 2011.

736 Pettygrove, G. S., Heinrich, A. L. and Eagle, A. J.: Dairy manure nutrient content and forms,
737 University of California Manure Technical Guide Series for Crop Management Professionals.
738 [online] Available from: www.manuremanagement.ucdavis.edu/files/134369 (Accessed 26
739 April 2016), 2010.

740 Rutledge, S., Mudge, P. L., Campbell, D. I., Woodward, S. L., Goodrich, J. P., Wall, A. M.,
741 Kirschbaum, M. U. F. and Schipper, L. A.: Carbon balance of an intensively grazed temperate
742 dairy pasture over four years, *Agric. Ecosyst. Environ.*, 206, 10–20,
743 doi:10.1016/j.agee.2015.03.011, 2015.

744 Skinner, R. H.: High Biomass Removal Limits Carbon Sequestration Potential of Mature
745 Temperate Pastures, *J. Environ. Qual.*, 37(4), 1319--1326, doi:10.2134/jeq2007.0263, 2008.

746 Smith, P.: How long before a change in soil organic carbon can be detected?, *Glob. Change*
747 *Biol.*, 10(11), 1878--1883, doi:10.1111/j.1365-2486.2004.00854.x, 2004.

748 Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K. B., Tignor, M. and
749 Miller, H. L.: *Climate Change 2007: The Physical Science Basis. Contribution of Working*
750 *Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*,
751 Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA. [online]
752 Available from: https://www.ipcc.ch/publications_and_data/ar4/wg1/en/spm.html (Accessed
753 23 May 2015), 2007.

754 Soussana, J. F., Allard, V., Pilegaard, K., Ambus, P., Amman, C., Campbell, C., Ceschia, E.,
755 Clifton-Brown, J., Czobel, S., Domingues, R., Flechard, C., Fuhrer, J., Hensen, A., Horvath,
756 L., Jones, M., Kasper, G., Martin, C., Nagy, Z., Neftel, A., Raschi, A., Baronti, S., Rees, R. M.,
757 Skiba, U., Stefani, P., Manca, G., Sutton, M., Tuba, Z. and Valentini, R.: Full accounting of the
758 greenhouse gas (CO₂, N₂O, CH₄) budget of nine European grassland sites, *Agric. Ecosyst.*
759 *Environ.*, 121(1–2), 121–134, doi:10.1016/j.agee.2006.12.022, 2007.

760 Soussana, J. F., Tallec, T. and Blanfort, V.: Mitigating the greenhouse gas balance of ruminant
761 production systems through carbon sequestration in grasslands, *Animal*, 4(3), 334--350,
762 doi:10.1017/S1751731109990784, 2010.

763 Tilley, J. M. A. and Terry, R. A.: A two-stage technique for the in vitro digestion of forage
764 crops, *Grass Forage Sci.*, 18(2), 104–111, doi:10.1111/j.1365-2494.1963.tb00335.x, 1963.

765 Tubiello, F. N., Salvatore, M., Ferrara, A. F., House, J., Federici, S., Rossi, S., Biancalani, R.,
766 Condor Golec, R. D., Jacobs, H., Flammini, A., Prosperi, P., Cardenas-Galindo, P.,
767 Schmidhuber, J., Sanz Sanchez, M. J., Srivastava, N. and Smith, P.: The Contribution of
768 Agriculture, Forestry and other Land Use activities to Global Warming, 1990-2012, *Glob.*
769 *Change Biol.*, 21(7), 2655--2660, doi:10.1111/gcb.12865, 2015.

770 Undi, M., Wilson, C., Ominski, K. H. and Wittenberg, M.: Comparison of techniques for
771 estimation of forage dry matter intake by grazing beef cattle, *Can. J. Soil Sci.*, 88, 693--701,
772 doi:10.4141/CJAS08041, 2008.

773 UNFCCC: Synthesis and Assessment Report on the Greenhouse Gas Inventories Submitted in
774 2014, United Nations Framework Convention on Climate Change, Bonn, Germany. [online]
775 Available from: http://www.ccrasa.com/library_1/12491%20-%20UNFCCC%20-%20Synthesis%20and%20assessment%20report%20on%20the%20greenhouse%20gas%20in%20inventories%20submitted%20in%202010.pdf (Accessed 11 June 2015), 2014.

778 VDLUFA: Die Untersuchung von Sekundärrohstoffdüngern, Kultursubstraten und
779 Bodenhilfsstoffen, Verband Deutscher Landwirtschaftlicher Untersuchungs- und
780 Forschungsanstalten, VDLUFA-Verlag, 204 pp., Speyer, Speyer., 2000.


781 Walther, U., Menzi, H., Ryser, J.-P., Flisch, R., Jeangros, B., Maillard, A. and Vuilloud, P. A.:
782 Grundlagen für die Düngung im Acker- und Futterbau, *Agrarforschung*, 1(7), 1--40, 1994.

783 Zeeman, M. J., Hiller, R., Gilgen, A. K., Michna, P., Plüss, P., Buchmann, N. and Eugster, W.:
784 Management and climate impacts on net CO₂ fluxes and carbon budgets of three grasslands
785 along an elevational gradient in Switzerland, *Agric. For. Meteorol.*, 150(4), 519–530,
786 doi:10.1016/j.agrformet.2010.01.011, 2010.

787


788

789 **Figures**



790

791 Figure 1. Duration of grazing on the study field (green bars) and for other pastures (gray)
792 over the day and year. The 'effective pasture time' of 73.1 days (total of green bars) plus the
793 adjacent 'off-pasture time' for milking of 25.9 days (blue bars) resulted in 'total grazing days'
794 of 99 days. White areas mark other times spent in the barn. White and gray bars are not
795 considered in the budget calculation.




796

797 **Figure 2.** Illustration of the two approaches to determine the net ecosystem carbon budget of
 798 a dairy pasture using different system boundaries (dashed red line): **(a)** NECB_{tot} using system
 799 boundaries including the cows; **(b)** NECB_{past} using system boundaries excluding the cows.
 800 Relevant carbon fluxes through the system boundaries are marked in blue (gaseous fluxes:
 801 light blue, liquid/solid fluxes: dark blue).

802


Gelöscht: 1



804
 805 **Figure 3.** Average daily carbon (blue arrows) and nitrogen (green arrows) budget of the studied
 806 dairy cows. The budget was closed by adjusting the amount of excreta loss.

Gelöscht: 2

807




809
 810 **Figure 4.** Components and uncertainties (95% confidence range) of annual carbon budget
 811 determined with (a) the total system and (b) the pasture system approach as illustrated in Fig.
 812 3. NECB was calculated according to Eqs. (2) and (3). Flux direction is defined according to
 813 ecological sign convention: positive values indicate imports to the system, negative values
 814 indicate export (loss) from the system. Filled bars indicate values derived from direct
 815 measurements, hatched bars indicate values that are modelled with measured and modelled
 816 data.

817

Gelöscht: 3

Gelöscht: 2




820

821 **Figure 5:** Effect of CO₂ flux selection based on the observed cow stocking density within the
 822 flux footprint on the annual CO₂ exchange ($F_{C-CO_2} = -NEE$) and number of fluxes used for the
 823 gap filling (bars). The dark blue diamond symbol represents $F_{C-CO_2,tot}$, the light blue triangle
 824 represents $F_{C-CO_2,past}$.

825

Gelöscht: 4



827
 828 **Figure 6:** Comparison of greenhouse gas fluxes of the pasture system including cows during
 829 pasture use to the NECBs for the two system boundaries. The ecological sign convention is
 830 used: negative values indicate a source from the system to the atmosphere. N₂O emissions are
 831 modelled, whereas the other emissions are measurements. Detailed numbers can be found in
 832 [Table S3](#).

Gelöscht: 5

Gelöscht: G

Gelöscht: loss/emission

833 **Supplementary material**

834 **S1 Uncertainty estimation of selected C budget components**

835 **S1.1 Animal intake**

836 The uncertainty of dry matter (DM) intake is dominated by the (systematic) uncertainty of the applied
837 empirical model (Eq. 6a/b) based on animal performance and characteristics (milk yield and
838 composition, live weight, etc.). To estimate this uncertainty we used results of a multi model validation
839 study by Jensen et al. (2015). They present in their Table 4 root mean square prediction errors (RMSPE)
840 for different published DM intake models. We selected the results of four models that use similar input
841 data like our model, i.e., the models by NRC (2001), Volden et al. (2011), Huhtanen et al. (2011) and
842 Gruber et al. (2004). We converted their RMSPE to relative errors and averaged them. Finally the
843 systematic model uncertainty (15%) was estimated as twice the average relative error.

844 Carbon content of pasture forage and concentrates were measured by dry combustion (VDLUFA, 2000)
845 of weekly sampled pasture forage ($n = 34$, but data from samples contaminated with soil were excluded)
846 and from periodically analyzed concentrate samples ($n = 6$ over the grazing period). The uncertainties
847 of the average C content was limited by the C analyzer uncertainty of 2%. For the concentrate intake
848 also the average DM to fresh matter ratio needed to be quantified from oven dried samples ($n = 6$). Its
849 uncertainty (4%) was estimated as 2SE.

850 Depending on individual production state of the cows they were offered concentrate in weighing
851 troughs. Mean daily fresh matter of concentrate ration amounted to $1.5 \text{ kg head}^{-1} \text{ d}^{-1}$. We assume a total
852 uncertainty of portion and weighing trough of 15% (expert guess). The uncertainty of concentrate C
853 intake was calculated by error propagation from the uncertainties of C content analysis (2%), the DM
854 content analysis (4%) and the weighing of the fresh matter intake for each cow (15%) resulting in a total
855 uncertainty of 16% for $F_{C\text{-feed,off}} = 0.6 \pm 0.1 \text{ kg C head}^{-1} \text{ d}^{-1}$.


856 Daily grazing C intake $F_{C\text{-grazing}}$ ($7.5 \pm 1.2 \text{ kg C head}^{-1} \text{ d}^{-1}$) was calculated from the difference between
857 total required C intake ($8.0 \pm 1.2 \text{ kg C head}^{-1} \text{ d}^{-1}$) and the offered C concentrate. The uncertainty of 16%
858 resulted from the error propagation of the uncertainties of total and concentrate C intake.

859

860 **S1.2 Milk carbon content**

861 The uncertainty of the milk yield related carbon flux was clearly dominated by the estimation of the
862 milk carbon content, which was not directly measured in this study. In a previous experiment Münger
863 (1997) determined the relationship between milk C content and milk gross energy content (Fig. S1).
864 Milk samples were collected during a study comparing energy utilization of three different dairy cattle
865 breeds over a whole lactation cycle. Energy content of the milk (estimated) was calculated according to
866 Arrigo et al. (1999) from sample contents of fat, protein and lactose as determined by mid-infrared

867 spectroscopy (Milkoscan, Foss A/B, Hillerød, DK). Carbon content was determined using the total
868 combustion of freeze-dried samples and subsequent gas analysis (CHN-600 Elemental Analyzer, Leco
869 Inc., St. Joseph MI, USA). A relationship of 21 g C MJ⁻¹ was derived from this experiment. The
870 uncertainty was estimated by fitting outer bands to the data comprising 95% of the points (dashed lines
871 in Fig. S1) resulting in a relative uncertainty of 9% (± 1.9 g C MJ⁻¹).
872



873
874 **Fig. S1:** Relationship between measured milk carbon content and milk gross energy content estimated
875 from measured fat, protein and lactose contents according to Arrigo et al. (1999): $y = 20.8 x$, $R^2 = 0.99$.
876 The dashed lines indicate the uncertainty range limits ($y = 23 x$ and $y = 19.2 x$).
877

878 **S1.3 CO₂ exchange**

879 Measured CO₂ exchange of the pasture system needed a gap filling procedure to derive an annual data
880 series without gaps. Felber et al. (2016) used the REddyProcWeb online partitioning and gap filling tool
881 (www.bgc-jena.mpg.de/bgi/index.php/Services/REddyProcWeb) with two different data sets: i) a data
882 set with fluxes that include fluxes with cow contribution to quantify $F_{C-CO_2,tot}$ and ii) the same data set
883 but without fluxes with cow contributions to quantify $F_{C-CO_2,past}$. The total uncertainty of the annual CO₂
884 fluxes (54 g C m⁻² yr⁻¹ and 44 g C m⁻² yr⁻¹, respectively) was determined from combined random and
885 systematic uncertainties. As reported by Felber et al. (2016), the existence of a high fraction of gaps and
886 the uncertainty of the filled data was the dominant error source. Its effect was estimated by a series of
887 simulations, in which additional gaps were introduced by randomly shifting the original gap structure
888 time series before gap filling. The corresponding results are presented in Fig. S2.
889




Fig. S2: Cumulative gap filled CO₂ fluxes (a) $F_{C-CO_2,tot}$ and (b) $F_{C-CO_2,past}$ simulated with additional gaps introduced by randomly shifting the original gap structure time series before gap filling. The colored lines indicate the time series with the original gap structure.

S1.4 Fertilization

The uncertainty of $F_{C-fertil}$ was combined from the uncertainty of the slurry and the urea application in the study year. The uncertainty of slurry application was estimated from previous studies in Switzerland. Ammann et al. (2009) measured C contents in slurry over several years and we adopted their uncertainty value of 17%. Only by specific sampling and analysis of the applied slurry (see Sect. 2.3.3), the relative error could be limited to <20%, because the DM and thus also the C content in slurry can easily vary by a factor of four.

The uncertainty of urea C was assumed to be close to zero because, at one hand, the absolute C amount ($2 \text{ g C m}^{-2} \text{ yr}^{-1}$) was small compared to the amount of C from slurry ($75 \text{ g C m}^{-2} \text{ yr}^{-1}$) and, at the other hand, the C content and the amount of applied urea can be determined very accurately. Thus the uncertainty of $F_{C-fertil}$ corresponds directly to the uncertainty of slurry C.

907 **S2 Budget results with uncertainties**

908 **Table S1:** Components of the average carbon (C) and nitrogen (N) budget of the dairy cows (Eq. 5 and
 909 9) with uncertainties (95% confidence range). The **N** budget was closed by adjusting the amount of
 910 excreta loss.

	Animal C exchange rate		Animal N exchange rate	
	(kg C head ⁻¹ d ⁻¹)	(% of intake)	(g N head ⁻¹ d ⁻¹)	(% of intake)
$E_{C/N}$ -intake	8.0 ± <u>1.2</u>	100	508 ± 137	100
E_{C} -resp	4.6 ± 1.6	57	-	-
E_{C} -CH _{4,cow}	0.3 ± 0.02	4	-	-
$E_{C/N}$ -milk	1.5 ± <u>0.1</u>	19	124 ± 13	24
$E_{C/N}$ -meat	<0.1	<1	<5	<1
$E_{C/N}$ -excreta	<u>2.6 ± 0.8</u>	<u>32</u>	380 ± 138	75
(Im-)balance	<u>-1.0 ± 2.0</u>	<u>12</u>		

Gelöscht: 2.2

Gelöscht: 0.2

Gelöscht: 1.6 ± 0.7

Gelöscht: 20

911

916 **Table S2:** Components and uncertainties (95% confidence range) of annual carbon fluxes ($\text{g C m}^{-2} \text{yr}^{-1}$)
 917 determined for the total system and pasture system approach. NECB was calculated according to Eqs.
 918 (2) and (3). Flux direction is defined according to ecological sign convention: positive values indicate
 919 imports to the system, negative values indicate export (loss) from the system of interest.

	Total system (incl. cows)	Pasture only (excl. cows)	<u>Attributed time</u> <u>used in Eq. (4)</u>
$F_{\text{C-CO}_2,\text{tot}}$	$+68 \pm 54$		<u>full year</u>
$F_{\text{C-CO}_2,\text{past}}$		$+248 \pm 44$	<u>full year</u>
$F_{\text{C-CH}_4,\text{soil}}$	-2 ± 1	-2 ± 1	<u>full year</u>
$F_{\text{C-CH}_4,\text{cows}}^{1)}$	-17 ± 1		<u>99 days</u>
$F_{\text{C-fertil}}^{2)}$	$+77 \pm 13$	$+77 \pm 13$	<u>full year</u>
$F_{\text{C-grazing}}$		-404 ± 65	<u>99 days</u>
$F_{\text{C-excreta,past}}$		$+104 \pm 30$	<u>73.1 days</u>
$F_{\text{C-products}}$	-82 ± 7		<u>99 days</u>
$F_{\text{C-feed,off}}$	$+31 \pm 5$		<u>99 days</u>
$F_{\text{C-resp,off}}$	-65 ± 23		<u>25.9 days</u>
$F_{\text{C-excreta,off}}$	-37 ± 11		<u>25.9 days</u>
NECB	-27 ± 62	$23 \pm 76^{3)}$	<u>full year</u>

Gelöscht: ± 61

Gelöscht: 64 ± 29

Gelöscht: 8

Gelöscht: 2

Gelöscht: 23 ± 10

Gelöscht: 13 ± 61

Gelöscht: -17 ± 81

Formatiert: Hochgestellt

920 ¹⁾ including $F_{\text{C-CH}_4,\text{cows}}$ during pasture and off-pasture times

921 ²⁾ $75 \text{ g C m}^{-2} \text{yr}^{-1}$ as cattle slurry and $2 \text{ g C m}^{-2} \text{yr}^{-1}$ as urea

922 ³⁾ For the uncertainty calculation of $\text{NECB}_{\text{past}}$ it was taken into account that the errors of
 923 $F_{\text{C-grazing}}$ and $F_{\text{C-excreta,past}}$ are highly correlated, because the excretion was calculated as
 924 a fraction of the animal intake (Sect. 2.4.3).
 925

933 **Table S3:** Comparison of components and uncertainties of the pastures greenhouse gas fluxes (g CO₂-
934 eq. m⁻² yr⁻¹) and the carbon sequestration determined for the total system (NECB_{tot}) and the pasture
935 system (NECB_{past}). The ecological sign convention is used: negative values indicate emission from the
936 system to the atmosphere. N₂O emissions are modelled, whereas the other emissions are measurements.

	<u>mean</u>	<u>uncertainty</u>
<u>N₂O</u>	<u>-219</u>	<u>-438/+153</u>
<u>CH₄,cows</u>	<u>-573</u>	<u>±33</u>
<u>CH₄,soil</u>	<u>-50</u>	<u>±38</u>
<u>NECB_{tot}</u>	<u>-98</u>	<u>±226</u>
<u>NECB_{past}</u>	<u>+85</u>	<u>±179</u>

937
938

939 **S3 References**

- 940 Ammann, C., Spirig, C., Leifeld, J. and Neftel, A.: Assessment of the nitrogen and carbon budget of two
941 managed temperate grassland fields, *Agric. Ecosyst. Environ.*, 133(3–4), 150–162,
942 doi:10.1016/j.agee.2009.05.006, 2009.
- 943 Arrigo, Y., Chaubert, C., Daccord, R., Gagnaux, D., Gerber, H., Guidon, D., Jans, F., Kessler, J.,
944 Lehmann, E., Morel, I., Münger, A., Rouel, M. and Wyss, U.: Fütterungsempfehlungen und
945 Nährwerttabellen für Wiederkäuer: das grüne Buch, 4th ed., Eidgenössische Forschungsanstalt für
946 Nutztiere, Zollikofen, Switzerland., 1999.
- 947 Felber, R., Neftel, A. and Ammann, C.: Discerning the cows from the pasture: Quantifying and
948 partitioning the NEE of a grazed pasture using animal position data, *Agric. For. Meteorol.*, 216, 37–47,
949 doi:10.1016/j.agrformet.2015.09.018, 2016.
- 950 Gruber, L., Schwarz, F. J., Erdin, D., Fischer, B., Spiekers, H., Steingass, H., Meyer, U., Chassot, A.,
951 Jilg, T., Omermaier, A. and Gruggenberg, T.: Vorhersage der Futteraufnahme von Milchkühen -
952 Datenbasis von 10 Forschungs-und Universitätsinstituten Deutschlands, in *VDLUFA Schriftenreihe 60*,
953 pp. 484–504, VDLUFA-Kongress, Rosstock., 2004.
- 954 Huhtanen, P., Rinne, M., Mäntysaari, P. and Nousiainen, J.: Integration of the effects of animal and
955 dietary factors on total dry matter intake of dairy cows fed silage-based diets, *animal*, 5(5), 691–702,
956 doi:10.1017/S1751731110002363, 2011.
- 957 Jensen, L. M., Nielsen, N. I., Nadeau, E., Markussen, B. and Nørgaard, P.: Evaluation of five models
958 predicting feed intake by dairy cows fed total mixed rations, *Livest. Sci.*, 176, 91–103,
959 doi:10.1016/j.livsci.2015.03.026, 2015.
- 960 Münger, A.: Energie- und Stickstoffverwertung bei Milchkühen verschiedener Rassen, Diss. ETH Nr.
961 11929, Eidgenössische Technische Hochschule ETH Zürich, Switzerland, Zürich, Switzerland., 1997.
- 962 NRC, (National Research Council): *Nutrient Requirements of Dairy Cattle*, National Academy Press,
963 Washington, D. C., 2001.
- 964 VDLUFA: *Die Untersuchung von Sekundärrohstoffdüngern, Kultursubstraten und Bodenhilfsstoffen*,
965 *Verband Deutscher Landwirtschaftlicher Untersuchungs- und Forschungsanstalten*, VDLUFA-Verlag,
966 204 pp., Speyer, Speyer., 2000.
- 967 Volden, H., Nielsen, N. I., Åkerlind, M., Larsen, M., Havrevoll, Ø. and Rygh, A. J.: Prediction of
968 voluntary feedintake, in *The Nordic Feed Evaluation System*, edited by H. Volden, pp. 113–126,
969 Wageningen Academic Publishers, Wageningen, The Netherlands., 2011.