

A probabilistic assessment of calcium carbonate export and dissolution in the modern ocean

G. Battaglia^{1,2}, M. Steinacher^{1,2}, and F. Joos^{1,2}

¹Climate and Environmental Physics, Physics Institute, University of Bern, Bern, Switzerland

²Oeschger Centre for Climate Change Research, University of Bern, Bern, Switzerland

Correspondence to: G. Battaglia (battaglia@climate.unibe.ch)

Abstract. The marine cycle of calcium carbonate (CaCO_3) is an important element of the carbon cycle and co-governs the distribution of carbon and alkalinity within the ocean. However, CaCO_3 export fluxes and mechanisms governing CaCO_3 dissolution are highly uncertain. We present an observationally-constrained, probabilistic assessment of the global and regional CaCO_3 budgets.

5 Parameters governing pelagic CaCO_3 export fluxes and dissolution rates are sampled using a Monte Carlo scheme to construct a 1000 member ensemble with the Bern3D ocean model. Ensemble results are constrained by comparing simulated and observation-based fields of excess dissolved calcium carbonate (TA^*). The minerals calcite and aragonite are modelled explicitly and ocean–sediment fluxes are considered. For local dissolution rates either a strong or a weak dependency on CaCO_3 saturation is assumed. In addition, there is the option to have saturation-independent dissolution above the saturation horizon. The median (and 68 % confidence interval) of the constrained model ensemble for global biogenic CaCO_3 export is 0.90 (0.72 – 1.05) Gt-C yr^{-1} , that is within the lower half of previously published estimates (0.4 – 1.8 Gt-C yr^{-1}). The spatial pattern of CaCO_3 export is broadly consistent with earlier assessments. Export is large in the Southern Ocean, the tropical
10 Indo–Pacific, the northern Pacific and relatively small in the Atlantic. The constrained results are robust across a range of diapycnal mixing coefficients and, thus, ocean circulation strengths. Modelled ocean circulation and transport time scales for the different setups were further evaluated with CFC11 and radiocarbon observations. Parameters and mechanisms governing dissolution are hardly constrained by either the TA^* data or the current compilation of CaCO_3 flux measurements such
15 that model realisations with and without saturation-dependent dissolution achieve skill. We suggest to apply saturation-independent dissolution rates in Earth System Models to minimise computational costs.

1 Introduction

The cycling of calcium carbonate (CaCO_3) forms an important component of the marine carbon
25 cycle. It co-governs the surface-to-deep gradients of alkalinity and dissolved inorganic carbon (DIC) in the ocean (Volk and Hoffert, 1985), dominates deep ocean alkalinity fluxes, and influences the

surface fields of DIC and alkalinity, thereby providing the background conditions for the uptake of excess anthropogenic carbon from the atmosphere. However, the export and dissolution fluxes of CaCO_3 are highly uncertain. For example, current estimates of CaCO_3 export diverge by a factor of ~ 4 ($0.4\text{--}1.8 \text{ Gt-C yr}^{-1}$, summarised in Berelson et al., 2007).

The CaCO_3 cycle is driven by calcifying organisms such as coccolithophorids, foraminifera or pteropods, which remove calcium (Ca^{2+}), alkalinity, and DIC from the pelagic surface ocean waters to form shells and structures of CaCO_3 . These CaCO_3 particles are eventually exported out of the surface, gravitationally sink through the water column and dissolve at depth or get buried in ocean sediments. The formation and dissolution of CaCO_3 introduces a vertical gradient in alkalinity and DIC. These gradients have been sustained against the counteracting forces of physical mixing and transport, which would otherwise have removed these gradients. This redistribution of alkalinity and carbon by biogenic and physical transport affects the partitioning of carbon between the ocean and atmosphere and a reduction, expected under ongoing ocean acidification, or even a complete stop of CaCO_3 export tend to decrease atmospheric CO_2 by a few ppm, and in the extreme case by up to ~ 50 ppm, on century time scales (Heinze, 2004; Gangstø et al., 2011).

There are different mineral forms of CaCO_3 and solubility is higher for high-magnesium calcite (typically present in fish) and aragonite (typically present in free-swimming pelagic sea snails and sea slugs (pteropods)) than for calcite (typically present in calcifying algae (coccolithophorids)). Thermodynamic considerations suggest that dissolution of CaCO_3 particles occurs only when the product of the calcium and carbonate ion concentrations in the surrounding environment is below the saturation product. The saturation product of all minerals increases with increasing pressure (Mucci, 1983). At depth, respiration of organic matter additionally decreases the concentration of CO_3^{2-} . As a result, the bulk of the water in the deep ocean is undersaturated with respect to CaCO_3 minerals, generally enabling their dissolution, and oversaturated in the upper ocean, thermodynamically hindering their dissolution (Steinacher et al., 2009). Overall, the quantitative understanding of dissolution kinetics is, however, low and published estimates of saturation-dependent dissolution kinetic parameters range over several orders of magnitude (summarised in Sarmiento and Gruber, 2006). CaCO_3 dissolution may, nevertheless, still occur in the upper ocean in suitable, undersaturated microenvironments which would be present for instance in the guts of zooplankton, suspended organic aggregates, or fecal pellets (Bishop et al., 1980; Milliman et al., 1999; Jansen and Wolf-Gladrow, 2001). There are in fact several tracer-based studies reporting CaCO_3 dissolution above the saturation horizon of bulk seawater (Barrett et al., 2014; Feely et al., 2002, 2004; Sabine et al., 2002b; Chung et al., 2003). In a modelling study, Friis et al. (2006), nevertheless, demonstrated that the method which is often employed to derive these upper ocean dissolution rates (Berelson et al., 2007, see Discussion section on TA^* -CFC age method), might not be applicable, because this method neglects physical transport and mixing of alkalinity. It is therefore still debated where and how fast settling CaCO_3 particles are dissolved in the water column. Another complication,

typically neglected in previous studies on open water CaCO_3 dissolution, is arising from ocean–
65 sediment interactions and the influence of associated burial and redissolution fluxes on alkalinity
and carbon concentrations in the open ocean (Archer, 1996).

Here we propose an alternative, probabilistic assessment of the global CaCO_3 budget. We con-
sider explicitly the transport and mixing of alkalinity and account for ocean–sediment interactions
to probabilistically constrain the CaCO_3 cycle with the observation-based distribution of the TA^*
70 tracer. TA^* reflects the imprint of the CaCO_3 cycle on alkalinity (see Sect. 2) and is therefore im-
pacted by CaCO_3 export, water column dissolution, physical transport and mixing of the released
alkalinity and DIC and ocean–sediment fluxes. For the probabilistic flux assessment, a large range
of calcite and aragonite export and dissolution flux parameterisations are employed within our Earth
System model of intermediate complexity (EMIC) – the Bern3D model – in a Monte Carlo setup
75 with 1000 members. The Bern3D model calculates the corresponding modelled steady-state tracer
concentrations of TA^* . The most probable export and dissolution fluxes are then the ones resulting
in modelled TA^* fields that match closely the observation-derived TA^* distribution. The aim is to
assign uncertainty estimates that will be consistent with both the observations and model equations
– a classical data assimilation problem (see Sect. 3.3). It is expensive to perform simulations with
80 interactive sediments (Heinze et al., 1999; Gehlen et al., 2006; Tschumi et al., 2011) due to long spin-
up times required to bring ocean and sediments in equilibrium and we account a posteriori for the
influence of sediment burial and dissolution fluxes on TA^* (see Sect. 3.3). This approach represents
an alternative to the interpretation of concentrations without a physical model or the interpretation of
flux/rate measurements within the water column, which are generally much sparser (~ 156 flux mea-
85 surements in the water column (Wilson et al., 2012) and ~ 56 benthic dissolution flux measurements
(Berelson et al., 2007), globally) and more difficult to obtain. We apply the Wilson et al. (2012) data
compilation of water column fluxes as an additional constraint for comparison. Additional sensitivity
analyses with respect to vertical diffusion (k_{dia} , see Sect. 4) are illustrated. The results are compared
and contrasted to data-based estimates of export and dissolution (as summarised in Berelson et al.,
90 2007). Finally, implications for the parameterisation of CaCO_3 dissolution in Earth System Models
are discussed.

2 Observation-derived TA^*

Total alkalinity data are from the GLODAP carbon climatology (Key et al., 2004) and salinity
(Antonov et al., 2010), temperature (Locarnini et al., 2010), oxygen and phosphate (Garcia et al.,
95 2010a, b) data are from the World Ocean Atlas. These observational data serve to split the alkalinity
signal into its different physical and biogeochemical components such as TA^* , our target variable
in the data assimilation. The gridded data products from GLODAP and the World Ocean Atlas are
derived from samples taken during the previous few decades. We first regridded all required, grid-

ded datasets to the Bern3D model grid (40x41x32 grid boxes) using the area-weighted regridding
 100 method of Ferret before deriving the other properties.

TA*, our target variable in the data assimilation, is a constructed tracer (Feely et al., 2002; Sabine
 et al., 2002a; Chung et al., 2003; Koeve et al., 2014) to exclusively capture the imprint of CaCO₃
 dissolution on alkalinity. TA* is, in this sense, one of three components of measured total alkalinity
 (TA, Eq. 1, mean concentration 2.427 mol m⁻³). TA* can be extracted from TA by accounting for
 105 preformed (TA⁰, Eq. 2) and remineralised alkalinity (TA^r, Eq. 3):

$$TA = TA^0 + TA^r + TA^*. \quad (1)$$

TA⁰ (mean concentration ~ 2.389 mol m⁻³) is the background or preformed concentration, set at
 the ocean surface, and mixed conservatively in the ocean interior. Accordingly, TA^r and TA* are by
 definition zero in the surface ocean. To describe TA⁰ in the ocean interior, one relies on a multi-linear
 110 regression relationship for surface ocean total alkalinity based on surface ocean salinity (*S*) and
 PO (PO = O₂ + $r_{-O_2:PO_4} \cdot PO_4$, $r_{-O_2:PO_4} = 170$, Broecker, 1974) and sometimes surface ocean
 temperature (*T*) as explanatory variables (all conservative variables, see Eq. (2), Gruber et al., 1996;
 Sabine et al., 2002a; Feely et al., 2002).

$$TA^0 = a_0 + a_1 \cdot S + a_2 \cdot T + a_3 \cdot PO \quad (2)$$

The coefficients, a_i , are estimated from observations of TA (Key et al., 2004), *S* (Antonov et al.,
 115 2010), *T* (Locarnini et al., 2010), and PO (Garcia et al., 2010a, b) in the surface ocean. The linear
 regression fit is sometimes further subdivided to include only specific basins (Feely et al., 2002;
 Koeve et al., 2014). Table 1 summarises different regressions, including previously published re-
 gression estimates as well as new estimates by us calculated on the Bern3D model grid (B3D). The
 120 root mean squared deviations (RMSE) between these fits are smaller than 0.00465 mol m⁻³ (i.e.
 smaller than 0.2 %, excluding the Gruber et al. (1996) equation, which relies on an older database
 from 1996). From these different options, we accordingly chose the Global B3D linear regression as
 robust regression for TA⁰.

TA^r (mean concentration -18 mmol m⁻³) is linked stoichiometrically to the apparent oxygen
 125 utilisation (AOU, Garcia et al., 2010b, Eq. 3) and accounts for decreases in TA due to the oxidation
 of organic nitrogen, phosphorous, and sulfur (OM).

$$TA^r = r_{Alk:OM} \cdot r_{NO_3:-O_2} \cdot AOU = 1.26 \cdot 16/170 \cdot AOU \quad (3)$$

We set $r_{Alk:OM}$ to 1.26 (Kanamori and Ikegami, 1982) and $r_{NO_3:O_2}$ to 16/170 (Anderson and
 Sarmiento, 1994) to uniformly, and globally link AOU changes to changes in TA (as in Feely et al.,
 130 2004; Koeve et al., 2014). Wolf-Gladrow et al. (2007) propose an 8 % higher value for $r_{Alk:OM}$ of
 1.36, based on a different sulphur to carbon ratio. In addition, we note that AOU has been suggested
 to overestimate true oxygen utilisation by 20–25 % (Ito et al., 2004; Duteil et al., 2013). Accordingly,
 TA^r might be associated with an uncertainty of ~ 20 %.

The remaining signal, then, is TA^* (mean concentration $\sim 57 \text{ mmol m}^{-3}$), the changes in alkalinity due only to the CaCO_3 cycle (Fig. 1). The global average RMSE of any of the described ways (16 in total) of accounting for TA^+ (25 % lower AOU or different $r_{\text{Alk:OM}}$) and TA^0 (either 2 or 3 explanatory variables, and either global or regional) from our reference choices is $\sim 4 \text{ mmol m}^{-3}$ (average RMSE 3.9, 3.8, 3.1 mmol m^{-3} in the Atlantic, Pacific and Indian Ocean, respectively, i.e. $\sim 7 \%$). It is to note that this approach, inherent to its empirical nature, yields slightly negative TA^* values in some places. TA^* integrates to about 37.5 Pmol C or 75 Pmol Alk-equivalents of which $\sim 41 \%$ come to lie above the calcite saturation horizon (similar to Koeve et al. (2014) who find 44.7 % of the TA^* inventory above the calcite saturation horizon). These estimates are robust across the different sources of uncertainty. TA^* concentrations are expressed in alkalinity equivalents throughout this paper and we do not divide TA^* concentrations by a factor of 2 as done in many observational studies which express TA^* in terms of carbon changes. TA^* inventories are in Pmol C and CaCO_3 fluxes are given in carbon units (Gt-C yr^{-1} , $\text{mmol-C m}^{-3} \text{ yr}^{-1}$, $\text{mmol-C m}^{-2} \text{ yr}^{-1}$).

3 The modelling framework

To constrain the alkalinity fluxes associated with CaCO_3 cycling, we introduce different export and dissolution fluxes (with a total of 15 degrees of freedom) within the biogeochemistry component of the Bern3D dynamic ocean model. The parameters of the new formulations describing the export and dissolution fluxes of CaCO_3 are varied using a Monte Carlo sampling method (McKay et al., 1979; Steinacher et al., 2013) and the resulting model ensemble, including 1000 members, is run to a pre-industrial steady state producing a broad range of solutions (see Sect. 3.2). The alkalinity fluxes associated with the CaCO_3 cycle are then constrained by comparing observation-based and simulated TA^* data using a Bayesian approach following Steinacher et al. (2013); a skill score is assigned to each ensemble member and used as a weight for the computation of median values and probability density functions from the ensemble results. The sediment module is not included in the ensemble due to extensive computational cost (~ 150 vs. $\sim 8 \text{ CPU h}$ for a single member run with and without the sediment, respectively). Instead, we re-run the model configurations which achieved the best skill scores with interactive sediments to account for CaCO_3 burial and sediment redissolution a posteriori (see Sect. 3.3.3).

3.1 The Bern3D Model

The Bern3D model couples a dynamic ocean, sea ice, an energy-moisture balance atmosphere, a marine biogeochemical cycle, a dynamic global vegetation model, and an ocean sediment module (Müller et al., 2006; Tschumi et al., 2011; Ritz et al., 2011; Roth et al., 2014). Here an ocean version with a horizontal resolution of 41 by 40 grid cells and 32 logarithmically-scaled vertical layers is used (see also Roth et al., 2014). The horizontal resolution is the same for the components

atmosphere, ocean, sea ice, and sediments of the Bern3D model. Transport and mixing of tracers in the ocean is based on Edwards et al. (1998) and Müller et al. (2006) as a three-dimensional frictional geostrophic model. The model has an isopycnal diffusion scheme and Gent–McWilliams parameterisation for eddy-induced transport (Griffies, 1998). The NCEP/NCAR monthly wind-stress climatology (Kalnay et al., 1996) is prescribed at the surface. Air–sea gas exchange for CO₂ is implemented according to OCMIP-2 protocols (Najjar et al., 1999; Orr and Najjar, 1999). The global mean air-sea transfer rate is reduced by 19% compared to OCMIP-2 to match observation-based estimates of natural and bomb-produced radiocarbon (Müller et al., 2008). A two-dimensional energy moisture balance model represents the atmosphere (Ritz et al., 2011). The model is spun up to equilibrium under preindustrial conditions, with atmospheric CO₂ set to 278 ppm. The spin-up period is 4000 years without the sediment module and 50 000 years with the sediment module. Remaining model drifts are negligible. We implicitly neglect potential changes in TA* over the industrial period by comparing model results for preindustrial conditions with TA* data reconstructed from recent measurements. Such changes are negligible in simulations with prescribed anthropogenic forcing in the Bern3D model.

The marine biogeochemical module computes the cycling of carbon, alkalinity, phosphate, iron, oxygen, silica, and carbon isotopes. New production of organic material in the model is limited by temperature, light, phosphate, and iron following Doney et al. (2006) as described by Parekh et al. (2008) and Tschumi et al. (2011). One third (33 %) of the new production is exported out of the euphotic zone (defined at 75 m) as particulate organic matter (POM), with the remainder contributing to the dissolved organic matter pool. Within biogeochemically similar regions/provinces a parameter termed rain ratio linearly scales the pattern of POM export to CaCO₃ export (silica-limitation is not considered here). We define eight such regions, each assigned an independent value for the export rain ratio parameter (mol inorganic carbon / mol organic carbon exported, later given in % inorganic to organic carbon exported). These include the Pacific, Atlantic and Indian sections of the Southern Ocean (< 35° S, separated at 240° W, 63° W and 30° E), the tropical (35° S to 30° N) Pacific, Atlantic, and Indian Ocean, and the northern (> 30° N) Atlantic (including the Arctic) and Pacific Ocean. A global parameter (f_{calc}) determines how much of the total CaCO₃ export flux represents calcite and how much represents aragonite ($1 - f_{\text{calc}}$). Abiotic CaCO₃ precipitation is virtually absent in today’s ocean and not considered (reviewed in Sarmiento and Gruber, 2006).

In the model, we implemented TA* as an explicit, idealised tracer (Koeve et al., 2014). It captures the alkalinity equivalents of CaCO₃ dissolution whenever it occurs and mixes this signal, accordingly. TA* values are set to zero throughout the surface ocean.

In sensitivity simulations with the sediment module enabled, the flux of CaCO₃, and of other particles, reaching the seafloor is passed to the sediment module from where a fraction potentially re-dissolves back into the water column. In simulations without the sediment module, the entire flux reaching the ocean floor re-dissolves back into the water column. Simulated TA* concentrations tend

205 to be lower with the sediment module enabled than without sediment module, because a fraction of the CaCO_3 export flux is removed from the ocean and buried in the geosphere. The sediment diagenesis model (Heinze et al., 1999; Gehlen et al., 2006; Tschumi et al., 2011) features the same horizontal resolution as the ocean model and 10 layers resolving the top 10 cm of the seafloor. It dynamically calculates the transport, remineralisation/redissolution and bioturbation of solid material
 210 within the top 10 cm of the seafloor as well as pore-water chemistry and diffusion as described in detail in Tschumi et al. (2011). Solutes diffuse over a boundary layer of 1 cm between the sediment column and the lowermost ocean grid cell. Four solid components (CaCO_3 , opal, POM and clay) and pore water substances (carbon and carbon isotopes, total alkalinity, phosphate, nitrate, oxygen and silicic acid) are modelled. The pore water carbonate ion concentration determines whether, and
 215 at which rate, CaCO_3 dissolves. Aragonite and calcite are not distinguished within the sediment module and CaCO_3 is assumed to be in the form of calcite. Any solid material that is pushed out of the diagenetic zone disappears into the subjacent diagenetically consolidated zone. During the spin-up of the ocean–sediment model, the net loss of alkalinity, and other tracers such as carbon and nutrients, to the sediments is immediately replaced by corresponding riverine inputs which are
 220 distributed uniformly along the coastlines (and then taken to be part of TA^0). The riverine input is diagnosed at the end of the spin-up.

The model features the main water masses and mixing time scales of the ocean, an essential prerequisite to realistically simulate TA^* and other tracers. The simulated and observed distributions of the ventilation tracers $\Delta^{14}\text{C}$ and CFC11 are provided in the supplementary information along with
 225 a Taylor Diagram (Taylor, 2001) of CFC11, $\Delta^{14}\text{C}$, temperature, salinity, DIC, TA, PO_4 , oxygen, and TA^* (Fig. A1 to A3). Globally, the correlation coefficient and standard deviation of the median relative to the standard deviation of the observations ($\sigma_{\text{rel.}}^{\text{obs.}}$) are 0.94 and 0.99 for $\Delta^{14}\text{C}$ and 0.91 and 0.98 for CFC11.

3.2 CaCO_3 dissolution within the water column

230 The dissolution equations for calcite and aragonite are implemented using equations with identical functional forms, but with different parameters for each mineral. In the following, we do not explicitly distinguish the two minerals to ease notation. The dissolution of calcite and aragonite below the euphotic zone is assumed to be a function of the saturation state of the bulk seawater, Ω , and the particle concentration per unit of water volume, $[\text{CaCO}_3]$ (see Gangstø et al. (2011) for a discussion):

$$235 \quad \frac{d[\text{CaCO}_3]}{dt} = -k_{\text{eff}}(\Omega) \times [\text{CaCO}_3]. \quad (4)$$

k_{eff} denotes a first order rate constant for either calcite or aragonite. It is defined as:

$$k_{\text{eff}}(\Omega) = k_0 \times (1 - \Omega)^n \times H(\Omega - 1) + k_{\text{bg}}. \quad (5)$$

k_0 and k_{bg} are rates in units of 1/time. H is the Heaviside function, which is zero for supersaturated and 1 for undersaturated water. The saturation state is defined by the ratio of the product of

240 calcium ion concentration times carbonate ion concentrations to the saturation product, K_{sp} (Mucci, 1983):

$$\Omega = \frac{[\text{Ca}^{2+}][\text{CO}_3^{2-}]}{K_{\text{sp}}}. \quad (6)$$

Ω values larger than one correspond to oversaturated and Ω values smaller than one to undersaturated conditions. Ω and thus k_{eff} are grid cell specific. At super-saturation, the dissolution rate equals the
 245 constant background rate k_{bg} (which can be zero). With increasing under-saturation, the dissolution rate increases towards its maximum value ($k_{\text{bg}} + k_0$). n is a unitless parameter and determines the deviation from linearity of this increase. For simplicity and to avoid the addition of further free parameters, a constant sinking velocity, v , is assumed and identical for both calcite and aragonite particles. The flux profile of CaCO_3 then takes the form

$$250 \quad F_{i,j}(z_k) = F_{i,j}(z_{k-1}) \cdot \exp\left(\frac{-k_{\text{eff}}(\Omega_{i,j,k})}{v} \cdot \Delta z_k\right) \quad (7)$$

where F is the downward flux of either calcite or aragonite particles per unit area evaluated at the bottom of each tracer grid cell at depth z_k . i , j , and k are grid cell indices indicating longitude, latitude, and depth. Δz_k denotes grid cell height. The export flux is set equal to F ($z = 75$ m) at the depth of the euphotic zone. Particles are dissolved instantaneously and sinking is not explicitly resolved in
 255 this formulation, reducing computational costs. v/k_{eff} is in units of m, and, if assumed constant, can be interpreted as the dissolution length scale, i.e. the depth at which the flux has decreased to $1/e$ (to $\sim 37\%$) of the export flux at 75 m. Any particles reaching the sea floor are dissolved completely in the appropriate lowermost box, except when the sediment module is included.

In Fig. 2 we illustrate three dissolution cases to explore the sensitivity of TA^* to dissolution profiles which cover the sampled uncertainty in dissolution rate parameters. Table 2 (middle column, $k_{\text{dia, ref}}$) summarises the resulting TA^* inventories and their relative distribution. The three dissolution rate profiles are selected to represent a case with constant, saturation-independent, dissolution (constant) and two cases where this background dissolution rate is set to zero and CaCO_3 dissolves only below the saturation horizon. In the fast (slow) case, aragonite (chosen to represent 10 % of
 265 the total) and calcite (90 % of total export) dissolves quickly (slowly) below the saturation horizon. Aragonite and calcite dissolve within a few hundred meters below the saturation horizon in the fast case, while most CaCO_3 dissolves on the ocean floor in the slow case.

The choice of the dissolution rate profile has a substantial influence on the simulated TA^* inventory (Fig. 2, Table 2). The global TA^* inventory is 38 Pmol C for the case with a constant, saturation-independent dissolution and a rain ratio of $\sim 7\%$, which is close to the observation-derived inventory of 37 Pmol C. The simulated inventory is 48 and 63 Pmol C for the fast and slow cases (where no
 270 dissolution occurs above the saturation horizon), respectively, and by that substantially higher. TA^* accumulates in the deep ocean when all CaCO_3 is dissolved below the saturation horizon of aragonite and calcite and no dissolution is permitted above. The TA^* inventory and concentrations are

275 sensitive to the choice of the dissolution rate profile, supporting our choice of TA^* as a target variable
to constrain dissolution rates.

3.3 Ensemble simulations and metrics for skill assessment

3.3.1 The Monte Carlo Ensemble

Following Steinacher et al. (2013) and Steinacher and Joos (2016) we run a 1000 member Latin
280 Hypercube ensemble to constrain the export flux out of the surface ocean, and the dissolution of
aragonite and calcite within the water column. Latin Hypercube sampling (McKay et al., 1979) is
a statistical, Monte Carlo method to generate controlled random samples from a multidimensional
distribution (15 dimensions in our case). The defined parameter ranges are divided into equally prob-
able intervals (1000 in our case). Random samples are then generated in each interval. This method
285 ensures that the sampled values are representative of the real variability while minimising the number
of required samples and thus the computational costs. We sample 15 parameters and apply uniform
priors based on literature information. The free parameters are the eight rain-ratio parameters (prior
range: 0 to 18 % for the Pacific and South Indian regions, prior range: 0 to 16 % for the tropical
Indian, prior range: 0 to 7 % for the North and tropical Atlantic, and prior range: 0 to 10 % for the
290 South Atlantic), defining the total amount of $CaCO_3$ export for each of the six regions. The prior
ranges of the rain ratios for the three Atlantic regions are limited to maximal 10%. This selection is
based on results from previous ensemble setups that revealed an overestimation of TA^* for high rain
ratios in the Atlantic domain. Further we include the fraction f_{calc} (1–0.5) defining the split between
aragonite and calcite export, and 3×2 parameters governing the dissolution kinetics for calcite and
295 aragonite, respectively. These are k_0 (0.05–10 day^{-1}) and n (1–4), describing fast and slow dissolu-
tion kinetics as a function of undersaturation, and k_{bg}/v (0–1/2500 m^{-1}), the length scale associated
with a constant, background dissolution rate acting both above and below the saturation horizon. v
is kept constant at 100 $m day^{-1}$.

The observation-based saturation-state of the bulk seawater with respect to aragonite and calcite is
300 prescribed for each model grid cell. It was calculated with the carbonate chemistry package seacarb
(Gattuso et al., 2010) from GLODAP (Key et al., 2004) and World Ocean Atlas 2009 (WOA09) (Lo-
carnini et al., 2010; Antonov et al., 2010). Seacarb calculates carbonate chemistry based on pressure,
temperature, salinity, alkalinity, DIC, silica and phosphate. This reduces computational costs as the
carbonate chemistry package of the online model requires substantial computational time. Also, this
305 avoids mismatches in the modelled and observation-based saturation states, which are also due to
model deficiencies in the cycling of organic matter and physical transport. Mismatches in modelled
and observed saturation states are particularly large in the North Pacific, where the modelled cal-
cite saturation horizon is up to 1.5 km too deep. The calcite saturation horizon is well represented
in the South Pacific, Indian and Atlantic by the model. The results presented in Sect. 4 suggest

310 that estimated CaCO_3 export production fields and dissolution rates are insensitive to the choice of the saturation field, because saturation-dependent and saturation independent parameterizations of dissolution yield similar TA^* fields.

3.3.2 Skill scores

Global skill scores, S_m , are assigned to each member, m , of the Latin-Hypercube ensemble:

$$315 \quad S_m = \exp(-0.5 \cdot \text{MSE}^{\text{rel}}). \quad (8)$$

MSE^{rel} is the relative mean squared error of the simulated TA^* concentrations from member m with respect to observation-derived TA^* :

$$\text{MSE}^{\text{rel}} = \sum_j a_j \times \frac{(\text{TA}_j^{\text{*model}} - \text{TA}_j^{\text{*obs}} - \text{TA}_j^{\text{*sedcorr}})^2}{\sigma^2}. \quad (9)$$

The sum includes all grid cells (indexed j). $\text{TA}_j^{\text{*model}}$ denotes simulated TA^* concentrations for ensemble member m and $\text{TA}_j^{\text{*obs}}$ observation-based TA^* concentrations estimated by using the Global B3D regression (Sect. 2). $\text{TA}_j^{\text{*sedcorr}}$ is a correction term arising from CaCO_3 burial in sediments, further explained below. a_j is the grid cell volume used as weight in the sum. σ^2 represents the combined error of the observation-based TA^* estimates and of the model and sets the scale against which model deviations are evaluated. Model deviations from the observations are considered large or small relative to the magnitude of σ^2 (Schmittner et al., 2009). The total uncertainty in observation-derived, gridded TA^* data is difficult to estimate and includes uncertainties due to extrapolation of limited number of measurements, uncertainties in individual tracer measurements, and in the computation of TA^* from tracer data. The error associated with the procedure to compute TA^* (average RMSE 3.9, 3.8, 3.1 mmol m^{-3} in the Atlantic, Pacific and Indian Ocean, respectively, see Sect. 2) is small compared to the model error (the best run achieves a RMSE of 11, 18, 18 mmol m^{-3} in the Atlantic, Pacific and Indian Ocean, respectively). Following Steinacher et al. (2013) and Schmittner et al. (2009), we estimate σ^2 as the (volume-weighted) variance of the model–data discrepancy for the ensemble member with the lowest MSE (this variance is 275 $(\text{mmol m}^{-3})^2$); this corresponds to MSE^{rel} close to unity for the best-fitting ensemble member.

335 The skill scores S_m of the individual ensemble members are likelihood-type functions corresponding to a Gaussian distribution of the data-model discrepancy ($\text{TA}^{\text{*model}} - \text{TA}^{\text{*obs}} - \text{TA}^{\text{*sedcorr}}$) with zero mean and variance σ^2 . S_m is an indication of the relative performance/credibility of each individual model configuration. Configurations which have relatively small deviations from the data are judged more probable than configurations which differ greatly from the observations.

340 S_m are used as weight to compute probability density functions (PDFs) and related measures such as the median (50th percentile) and the 16th and 84th percentiles defining the one standard deviation confidence interval (1σ) of the ensemble results. PDFs represent weighted and normalised histograms of the variables of interest. The normalisation is such that the integral over a PDF equals

1. A cubic spline interpolation is used to arrive at a continuous PDF from the discrete, normalised
345 histogram. For the computation of median and confidence ranges the histograms are converted to
cumulative distribution functions (CDFs). We interpolate linearly within the discrete CDFs to arrive
at the chosen percentiles (i.e. CDF=(0.16, 0.5, 0.68)).

The above explanations apply for any simulated quantity of interest. In the following we will
present PDFs, median values, and 1σ confidence ranges for aragonite and calcite export and dis-
350 solution as well as for tracer concentrations at individual grid cells or integrated over regions or
the whole ocean. Spatial integrations are done for each ensemble member individually and before
computing the PDFs and associated measures from the full ensemble.

3.3.3 A first order correction for ocean–sediment TA^* fluxes

CaCO_3 burial removes alkalinity from the ocean water column and lowers concentrations and the
355 overall TA^* inventory relative to a run without the sediment module. Riverine input compensates this
loss. This input of alkalinity is added to the surface ocean and by that to the part of the preformed al-
kalinity component (TA^0), leaving TA^* unchanged. CaCO_3 export and dissolution within the water
column and the corresponding fluxes of alkalinity and TA^* remain largely unchanged between runs
with and without sediments.

360 Ideally, the ensemble would be run fully interactively with the sediment module enabled to ac-
count for all important processes within the CaCO_3 cycle. However, this is computationally too
expensive as the sediment module requires a long spin up to achieve equilibrium. A first order cor-
rection term $\text{TA}^{*\text{sedcorr}}$ that accounts for the influence of CaCO_3 burial and dissolution fluxes on
 TA^* is estimated as follows. First, the skill scores are computed as described above with $\text{TA}^{*\text{sedcorr}}$
365 set to zero. Then, the 14 best ensemble members are selected and re-run with the sediment module
enabled. The mean difference in the TA^* fields between the simulations with and without sediments
yields the sediment correction, $\text{TA}^{*\text{sedcorr}}$.

These simulations with sediments yield a mean global burial flux of $0.12 \text{ Gt-C yr}^{-1}$ (see Table 3:
 $0.011 \text{ Gt-C yr}^{-1}$ in the Atlantic, $0.062 \text{ Gt-C yr}^{-1}$ in the Pacific, and $0.042 \text{ Gt-C yr}^{-1}$ in the Indian
370 Ocean). This is within the estimate by Feely et al. (2004) of $0.1\text{--}0.14 \text{ Gt-C yr}^{-1}$.

The sediment burial correction on TA^* is largest in the Pacific and smallest in the Atlantic (Figs. 3
and 4). The global TA^* inventory in runs with the sediment is 6.9 Pmol C lower compared to the
runs without the sediment (see Table 3: 0.9 Pmol C in the Atlantic, 4.6 Pmol C in the Pacific, and
 1.4 Pmol C in the Indian Ocean). This reduction is equal to the inventory of the $\text{TA}^{*\text{sedcorr}}$ correction
375 and ~ 16 and $\sim 20\%$ of the observation-derived TA^* inventory of the Pacific and Indian Ocean,
respectively. TA^* concentrations are affected relatively uniformly below 1 km and differences in
 TA^* between simulations with and without sediment module tend to vanish toward the surface ocean
(Figs. 3 and 4). Correspondingly, the spatial patterns of TA^* are very similar for simulations with
and without sediments; correlation coefficients are > 0.99 . We expect that the correction will tend

to increase the export flux of CaCO_3 in the optimisation to compensate for the loss by burial, but will not strongly affect dissolution parameters as the spatial patterns of TA^* remain similar with or without correction.

A few caveats apply to this first order estimate of $\text{TA}^{\text{sedcorr}}$. First, our approach involves a few, likely minor, technical inconsistencies. Aragonite is not treated explicitly in the sediment module and all CaCO_3 is assumed to be in the form of calcite; this may tend to bias sediment burial high as calcite is less soluble. The saturation state within the sediments is computed interactively with modelled ocean boundary conditions; this may locally lead to inconsistencies as CaCO_3 dissolution within the ocean water column is computed using the prescribed, observation-based saturation state. The alkalinity flux associated with organic matter remineralisation within the sediment is not explicitly distinguished and included in the flux of TA^* from sediments to the ocean; this results in a bias on order of 5 % in the redissolution flux and a negligible influence on the sediment correction.

4 Results

4.1 Observation-derived vs. simulated TA^*

Reconstructed TA^* (Fig. 1, $\text{TA}^{*\text{obs}}$) is – by definition – close to zero at the ocean surface and correspondingly low within the well-ventilated North Atlantic Deep Waters, Antarctic Intermediate and Mode Waters, and within North Pacific Intermediate Waters. TA^* concentrations in the deep ocean are increasing with the age of water masses (see Fig. A2 of $\Delta^{14}\text{C}$ which is a proxy for water mass age) as the dissolution of CaCO_3 continues to add TA^* along the flow path. TA^* concentrations are around 40 to 50 mmol m^{-3} in the deep Atlantic (Antarctic Bottom Water) and in the deep Southern Ocean and increase to 130 mmol m^{-3} in the northern Pacific. The reconstructed basin-mean profiles in the Pacific and Indian basin (Fig. 4) show strong gradients in the upper 1500 m and relatively uniform values below that. Concentrations are generally much lower in the Atlantic than in the Pacific.

The unconstrained model ensemble yields a large range of TA^* concentrations (Fig. 4, grey shading). The optimisation procedure greatly reduces this range in simulated TA^* to a comparably narrow confidence interval (Fig. 4, green shading representing the 68 % confidence interval). For example, basin-averaged concentrations in the deep Pacific (4000 m) range between 11 to 287 mmol m^{-3} in the unconstrained ensemble, while the corresponding 68 % confidence interval in TA^* is 73 to 122 mmol m^{-3} in the constrained ensemble. The selected a priori parameter ranges are therefore wide enough to result in a very broad range of TA^* concentrations; the Bayesian optimisation framework confines this initial range around or close to the observation-based values.

The median field from the constrained model ensemble generally captures the observation-based TA^* pattern (Fig. 5). The correlation coefficient (r) between the two fields are 0.83, 0.88, and 0.87 in the Atlantic, Pacific, and Indian Ocean, respectively, and the RMSE between the two fields are 17.7, 23.8, 19.8 mmol m^{-3} for the respective basins. These deviations correspond, respectively, to 57, 28

415 and 28 % of their mean TA^* concentration in each basin. Large positive deviations are found in the intermediate waters of the Pacific. This is also evident in the observation-derived basin-mean profiles of TA^* , which generally fall within the 68 % confidence interval of the constrained model ensemble (Fig. 4). In the Atlantic, the ensemble median concentrations are, on basin-average, higher than the reconstructed ones. In the Pacific and Indian Oceans, the median clearly overestimates TA^* in 420 the thermocline (the 68 % confidence range does not include the observations there) and somewhat underestimates TA^* in the deep ocean. This is likely linked to known deficiencies in the model's circulation. Intermediate and mode waters (with low TA^* concentrations) are not penetrating far enough towards the equator. As a consequence, mixing of TA^* depleted surface waters is too low and TA^* concentrations are too high in the thermocline. Alternatively, we cannot exclude that dissolution 425 may be overestimated in the thermocline of the Pacific and Indian Ocean. This data-model mismatch could potentially be reduced by introducing more than one background dissolution rate constant (k_{bg}) or a depth-dependent particle sinking velocity. However, this may simply mask deficiencies in the circulation and we do not attempt such a solution. Generally, the correlation between observed and modelled TA^* is remarkably high.

430 4.2 Probabilistic estimates of CaCO_3 and alkalinity fluxes

The estimated global median export flux of CaCO_3 at 75 m – with its 68 % confidence interval – is 0.90 (0.72–1.05) Gt-C yr^{-1} (Table 4 and Fig. 6). Basin-wide, we find CaCO_3 median export fluxes (and 68 % confidence intervals) of 0.12 (0.078–0.17) Gt-C yr^{-1} from the Atlantic, 0.55 (0.39–0.68) Gt-C yr^{-1} from the Pacific, and 0.23 (0.14–0.32) Gt-C yr^{-1} from the Indian Ocean (Table 4 435 and Fig. 6). Regionally, largest export fluxes are simulated in the Southern Ocean sector of the Pacific, in the Pacific equatorial upwelling regions, and in the northwestern Pacific (Fig. 7). In the Indian Ocean export fluxes are highest in its eastern tropical regions and in its section of the Southern Ocean. In the tropical and northern Atlantic export fluxes are generally low, consistent with the low TA^* values in the bulk of NADW and AABW.

440 Median CaCO_3 export production per unit area in the constrained ensemble is considerably lower in the Atlantic sector of the Southern Ocean ($135 \text{ mmol-C m}^{-2} \text{ yr}^{-1}$) as compared to the Pacific ($301 \text{ mmol-C m}^{-2} \text{ yr}^{-1}$) and Indian ($326 \text{ mmol-C m}^{-2} \text{ yr}^{-1}$) sectors (Fig. 7 top). This is attributable to the choice of the regional boundaries for the rain ratio regions and the assumption that the spatial pattern of export within a region is identical to the pattern simulated by the standard 445 version of the model. The standard model yields relatively little zonal variation in the CaCO_3 export fluxes in the Southern Ocean in contrast to the data assimilation with lower than zonally-averaged export in the Atlantic. This reflects the much lower TA^* reconstructed in the deep Atlantic as compared to the deep Pacific and Indian (Fig. 1). A large export in the Atlantic sector of the Southern Ocean tends to yield high simulated TA^* concentrations in the Antarctic Bottom Water that fills the deep 450 Atlantic. The Monte Carlo data assimilation therefore requires low CaCO_3 export in the Atlantic

sector to minimize model-data mismatches in the deep Atlantic. It is difficult to correctly represent water mass formation and circulation in the Southern ocean and our model may be biased. A known bias is that the Atlantic Bottom Water circulation is too sluggish, also evidenced by simulated low radiocarbon signatures (Figs A2, A4). The influence of a potential bias in South Atlantic export on
455 global CaCO_3 export is estimated to be relatively small; assuming the same (median) CaCO_3 export per unit area in the Atlantic sector as estimated for the Indian sector would yield $0.06 \text{ Gt-C yr}^{-1}$ higher export than suggested by the ensemble median. While our Monte Carlo approach is suitable to estimate export fluxes over larger regions, the detailed spatial patterns in CaCO_3 export remain unconstrained.

460 Globally, the deposition flux on the respective deepest cells integrates to $0.16 (0.11\text{--}0.23) \text{ Gt-C yr}^{-1}$, i.e., $\sim 18 \%$ of the export flux (Fig. 7, Table 4). Local deposition depends on the local CaCO_3 export and on how much dissolution is occurring in the water column, which itself depends on the saturation state and on the depth of the water column. Particularly in the North Atlantic, along coastlines and in the Southern Ocean, high fluxes reach the ocean floor. These are dissolved into the
465 water column in the ensemble setup without sediment module. Accordingly, $\sim 82 \%$ of the global median CaCO_3 export dissolves in the water column. More specifically, $\sim 37 \%$ of the CaCO_3 export dissolves in the upper water column above 1500 m , $\sim 44 \%$ below 1500 m depth, with the remaining $\sim 18 \%$ dissolving at the sea floor (Table 4).

Average dissolution profiles for aragonite (red) and calcite (olive) in different ocean regions are
470 displayed in Fig. 8. A peak in aragonite and calcite dissolution is located at or below the depth of the aragonite and calcite saturation horizon, respectively. These dissolution peaks are associated with the saturation-dependent dissolution rate coefficients (Eq. 5). As is the saturation horizon, these peaks are located deep down in the water column of the north and tropical Atlantic region (at $3\text{--}4 \text{ km}$ depth for aragonite), at intermediate depth of the South Atlantic, Indian, and Pacific (1.5 km for aragonite)
475 and at relatively shallow depth of the tropical and north Pacific region ($\sim 700 \text{ m}$ for aragonite). As the calcite saturation horizon is found deeper in the water column than the aragonite saturation horizon, so are these dissolution peaks located deeper down in the water column for calcite than for aragonite.

Our constrained ensemble includes non-zero values for the background dissolution rate. Conse-
480 quently, calcite and aragonite are dissolving throughout the water column, irrespective of the saturation state in the Atlantic, Pacific and Indian Ocean. The percentage of the export flux which dissolves in waters supersaturated with respect to calcite are 72% ($60\text{--}80 \%$), 43% ($30\text{--}53 \%$), 68% ($55\text{--}78 \%$) in the Atlantic, Pacific and Indian Ocean, respectively. We will further investigate in Sect. 4.4 to which extent the finding that a fraction of the CaCO_3 export dissolves above the calcite
485 saturation horizon and our export and dissolution flux estimates are robust.

4.3 Sensitivity of results to ocean–sediment interactions, and circulation

4.3.1 Ocean-sediment interactions

As discussed in Sect. 3.3, CaCO_3 deposition on, burial within, and redissolution from ocean sediments affects TA^* concentrations mainly in the Pacific and Indian Ocean in our model. To assess
490 uncertainties in CaCO_3 fluxes arising from uncertainties associated with the sediment correction, we set the sediment correction ($\text{TA}^{*\text{sedcorr}}$) of the TA^* field to zero to calculate potential skill scores (Eq. 9). This alternative case also illustrates the potential error due to the neglect of ocean–sediment fluxes. Export fluxes of CaCO_3 are 2, 12, 8, and 8 % smaller in the Atlantic, Pacific, Indian and global ocean, respectively while the variance is about the same. This is not surprising as we have
495 already noted (Sect. 3.3) that a certain burial flux tends to decrease the TA^* pattern uniformly, i.e. the simulations with and without the sediment correlate highly in terms of TA^* . The PDF of fluxes is therefore shifted to higher export fluxes, while the preference for dissolution parameters remains the same. The CaCO_3 that dissolves above 1500 m is 4 % higher, 5 % lower, the same, and 4 % lower in the Atlantic, Pacific, Indian and global ocean, respectively, in the setup that neglects sediment fluxes
500 compared to the standard setup.

4.3.2 Implications of different diapycnal diffusivities illustrated for three dissolution rate profiles

The diapycnal diffusivity, k_{dia} of the model is varied to probe uncertainties related to the magnitude of the ocean overturning circulation. k_{dia} is either set to 0.1 (low), 0.2 (standard), or
505 $0.5(\text{high}) \times 10^{-4} \text{ m}^2 \text{ s}^{-1}$. Increasing k_{dia} increases the strength of the overturning circulation, and deep ocean ventilation. Maximum Atlantic Meridional Overturning is 16, 18, and 23 Sv (Sverdrups), Southern Ocean overturning is -16 , -14 , and -15 Sv and maximal deep Pacific overturning is -13 , -14 , and -20 Sv for the low, standard and high k_{dia} simulations, respectively. We compare the simulated (natural) $\Delta^{14}\text{C}$ of DIC and CFC11 distributions to their corresponding observations (Key et al.,
510 2004) to evaluate the physical transport (see Appendix). Both are conservative tracers with a time information and indicative of the ventilation time scales of the thermocline and the deep ocean, respectively. The observation-based (Key et al., 2004) global mean (natural) $\Delta^{14}\text{C}$ of DIC is -151‰ . The reference simulation achieves a mean (natural) $\Delta^{14}\text{C}$ of DIC of -160‰ . Correspondingly, ocean radiocarbon signatures become too low (-176‰) with the low diapycnal mixing rate and
515 too high with the high (-126‰) diapycnal mixing rate (see Fig. A4). Simulated surface-to-deep $\Delta^{14}\text{C}$ gradients are too low (high) relative to the observed gradients for the high (low) diapycnal diffusivity parameter, thereby indicating too fast (slow) surface-to-deep water exchange (Fig. A4). The global observation-based CFC11 inventory is estimated to 575 Mmol. The reference simulation yields 513 Mmol CFC11 as the mean inventory over the modelled period between 1990 and
520 2000. In the low mixing simulation, this inventory is even lower (479 Mmol) and in the high mixing

simulation it is too high (631 Mmol). On a global scale, our reference choice is therefore in better agreement with these physical tracers (see Fig. A4).

We vary k_{dia} for three illustrative dissolution rate profiles introduced in Sect. 3.2 (see Fig. 2 and Table 2). In the low diapycnal diffusivity case, CaCO_3 export is 11 % lower and in the high diapycnal diffusivity case, CaCO_3 export is 30 % higher compared to the standard case. Larger overturning and mixing yields more nutrient input into the euphotic zone and thus more organic matter export. CaCO_3 export is independent of alkalinity in our model and does not depend on the choice of the dissolution rate. The simulated TA^* patterns remain similar and correlation between the patterns is at least 0.93. Basin-average profiles in TA^* vary little in the upper ocean and the deep Pacific ($< 14 \text{ mmol m}^{-3}$) and modestly in the deep Atlantic ($< 44 \text{ mmol m}^{-3}$) and deep Indian ($< 18 \text{ mmol m}^{-3}$) when varying diapycnal diffusivity between 0.1 and $0.5 \times 10^{-4} \text{ m}^2 \text{ s}^{-1}$. Perhaps somewhat surprisingly, the simulated TA^* inventory is relatively weakly affected by the choice of k_{dia} ; the global TA^* inventory varies by less than 7 % across the range of k_{dia} (Table 2). In other words, variations in the magnitude of ocean ventilation do hardly affect the TA^* inventory for a given dissolution rate profile. Higher (lower) export under higher (lower) mixing compensate each other. These changes in TA^* inventory are smaller than the influence of the sediment correction (see Table 3) or the choice of the dissolution profile. In conclusion, simulated TA^* is only weakly affected by uncertainties in the diapycnal mixing coefficient.

The choice of the dissolution rate profile has a substantial influence on the simulated TA^* inventory (Table 2). As mentioned previously, TA^* accumulates in the deep ocean when all CaCO_3 is dissolved below the saturation horizon of aragonite and calcite and no dissolution is permitted above. This raises the question whether surface-to-deep transport is too slow in our model. As mentioned above, the radiocarbon signatures as well as CFC11 concentrations are on average close to observations (Figs. A3 and A2). Increasing ocean ventilation by increasing k_{dia} results in too young radiocarbon signatures and higher than observed CFC11 concentrations. However, it does not substantially reduce the overestimation of the TA^* inventory in those cases.

4.4 How to parameterise CaCO_3 dissolution in an Earth System Model?

An important question is how to formulate the dissolution rate of calcite and aragonite particles in Earth System Models. Should the dissolution rate be a function of the simulated aragonite and calcite saturation state of the surrounding water? Should dissolution above the saturation horizon be permitted? In our ensemble, the degrees of freedom allow for either fast or slow dissolution kinetics as a function of saturation, and the option to also dissolve CaCO_3 above the saturation horizon with a constant dissolution rate. For a more systematic understanding of which parameterisations achieve high skill we return to the sensitivity simulations (Fig. 2).

Interestingly, the different dissolution schemes yield highly correlated patterns of TA^* ; the correlation coefficient between the global fields from the fast and slow dissolution scheme is 0.85, and

between the fields from the fast and constant scheme is 0.88 in simulations with the same CaCO_3 export (Fig. 2). The high correlation between fast and slow dissolution schemes is not surprising as only 35% of the modern ocean's volume is actually undersaturated with respect to calcite. Differences therefore cannot emerge over much of the ocean and skill is therefore distributed across the range of parameter values sampled.

Figure 9 displays the relationship between model skill, background dissolution rate, CaCO_3 export, and TA^* inventories in the model ensemble. To achieve a high skill (green-to-red coloured dots in the upper panels of Fig. 9), an individual ensemble member needs to reproduce the observation-based TA^* inventory (dashed line) within a limited range. We identify an export range within which TA^* can be reproduced skillfully (vertical green range in the upper panels of Fig. 9). The background rate is responsible for this spread. This is illustrated by plotting the value of the background dissolution rate, k_{bg} (lower panels of Fig. 9) as a function of the TA^* inventory and export. Generally, the higher the global CaCO_3 export flux, the higher the background dissolution rate required to achieve a high skill. Likewise, lower export can be distributed skillfully without dissolution in supersaturated waters ($k_{\text{bg}} = 0$). Apparently, there are trade-offs between the magnitude of export and the applied dissolution parameterisation in terms of TA^* , suggesting that export and dissolution parameters can only be constrained simultaneously within limits when using TA^* as the only constraint. In other words, neither the dissolution scheme nor its parameters are well constrained by the observation-based TA^* field. Corresponding to the high correlation, the RMSE between normalised observed and normalised modelled fields vary within a limited range (42 to 58%) for the three dissolution cases. The high spatial correlation in simulated TA^* and uncertainties in CaCO_3 export make it difficult to distinguish different dissolution parameterisations. The magnitude of CaCO_3 exports modulates absolute TA^* concentrations and thus model-data bias and root mean square errors. Given these uncertainties, we cannot objectively determine the preferred dissolution scheme.

4.5 Flux measurements as additional constraint

Global sediment trap data represent another observational constraint of the CaCO_3 cycle. The database of Wilson et al. (2012, see Fig. A5) includes 156 measurements globally and is an update of the Honjo et al. (2008) compilation. For comparison, we constrained the model ensemble using these sediment trap data instead of the TA^* data as target (Figs. A5 and A6). Skill-scores are calculated individually for the Pacific, Atlantic, and Indian Ocean. CaCO_3 export fluxes, dissolution profiles and parameters constrained with the sediment trap data are consistent within uncertainties with those constrained by the TA^* data. The sediment trap data, however, yield wider uncertainty ranges, as illustrated in Fig. A6, and do therefore not permit us to reduce uncertainty ranges any further.

5 Discussion

5.1 Export of CaCO_3

Berelson et al. (2007) summarised current estimates of CaCO_3 export out of the euphotic zone (based on models and data) to $0.4\text{--}1.8 \text{ Gt-C yr}^{-1}$ (spanning factor ~ 4). Our constrained median
595 estimate of ~ 0.90 (ensemble range: $0.72\text{--}1.05$) Gt-C yr^{-1} therefore lies on the lower part of these previously published estimates. At the end, these authors suggest that global CaCO_3 export must be higher than 1.6 Gt-C yr^{-1} . This estimate is based on sediment trap data and other information constraining the flux to the deep ocean ($> 2000 \text{ m}$) to $0.6 \pm 0.4 \text{ Gt-C yr}^{-1}$ and results obtained with the so-called $\text{TA}^*\text{-CFC}$ age method suggesting an upper ocean dissolution of 1 Gt-C yr^{-1} .
600 The $\text{TA}^*\text{-CFC}$ age method is heavily criticised by Friis et al. (2006) and tends to bias estimates systematically towards high values (Friis et al., 2006). This method and its shortcomings are further discussed in the next section on upper ocean dissolution. While our estimated flux to the deep ocean of 0.52 ($0.43\text{--}0.61$) Gt-C yr^{-1} is roughly consistent with the budget of Berelson et al. (2007), their export estimate, and upper ocean dissolution, is in clear conflict with our results and those of other
605 studies (Lee, 2001; Sarmiento and Gruber, 2006; Jin et al., 2006) that apply a range of different methodologies. We attribute this mismatch to deficiencies in the $\text{TA}^*\text{-CFC}$ age method, implying that the export estimate by Berelson et al. (2007) is biased high.

Global scale CaCO_3 export has been previously estimated (see Fig. 10). Jin et al. (2006) diagnosed a global CaCO_3 export of 1.1 Gt-C yr^{-1} by restoring annual-mean potential alkalinity to ob-
610 servations (Key et al., 2004) in the euphotic zone and in the whole water column within the Modular Ocean Model. Sediment burial and sediment-ocean fluxes are included implicitly in this approach. They provide a relatively small uncertainty range of 0.8 to 1.2 Gt-C yr^{-1} which falls within our range. Most of the uncertainty is attributed to uncertainties in the alkalinity data by these authors. Sarmiento and Gruber (2006) estimated CaCO_3 export ($\sim 0.5 \text{ Gt-C yr}^{-1}$) by combining satellite
615 NPP (as a mean of the three algorithms (Behrenfeld and Falkowski, 1997; Carr, 2002; Marra et al., 2003)) with the organic particle export model of Dunne et al. (2005) and the rain-ratio estimate of Sarmiento et al. (2002). Lee (2001) derived net CaCO_3 production ($\sim 0.92 \pm 0.3 \text{ Gt-C yr}^{-1}$) from seasonal potential alkalinity decreases. Both approaches focus on information of the surface ocean without taking advantage of information displayed in the full biogeochemical depth profile. In addition,
620 Ridgwell et al. (2007) used an Ensemble Kalman Filter approach to assimilate total alkalinity and phosphate data into their model and determined a global CaCO_3 export flux of 1.2 Gt-C yr^{-1} . On average, these four studies yield a global mean CaCO_3 export of $0.93 \text{ Gt-C yr}^{-1}$, close to our median estimate of $0.90 \text{ Gt-C yr}^{-1}$.

We find peaks in zonally-averaged export in the North Pacific, the tropical Pacific and Indian and
625 in the Southern Ocean and low export fluxes in the subtropics and the tropical Atlantic (Fig. 11). This is in agreement with the results of Jin et al. (2006) and Lee (2001), with the exception that

Lee (2001) suggest little CaCO_3 export in all tropical regions. A significant CaCO_3 export in the tropics is consistent with deep ocean sediment trap data (Francois et al., 2002; Berelson et al., 2007). As noted by Jin et al. (2006), the low estimates of Lee (2001) for tropical regions is likely related to small signals in seasonal alkalinity in the tropics, hampering their calculations. Concerning the magnitude of the export, our zonally-averaged values are similar to these two studies in the Pacific sector of the Southern Ocean, and smaller in the Atlantic and Indian Southern Ocean sectors as well as in the North Pacific.

Interestingly, Lee (2001) find significant CaCO_3 export in the North Atlantic, in contrast to Jin et al. (2006), Sarmiento and Gruber (2006), and this study. The relatively low North Atlantic export suggested by these latter studies appears to be in conflict with the occurrence of coccolithophorid blooms in this region (Brown and Yoder, 1994). On the other hand, the low TA^* inventory in the Atlantic (Fig. 1) argues for a limited export of CaCO_3 in this basin.

Carter et al. (2014) defined another alkalinity tracer, termed Alk^* , that isolates the portion of the alkalinity signal that varies in response to calcium carbonate cycling and exchanges with terrestrial and sedimentary environments from the portion that varies in response to freshwater and organic matter cycling. These authors compiled a riverine input of alkalinity into the low-latitude Atlantic ($> 40^\circ \text{S}$ and $< 40^\circ \text{N}$) equivalent to $0.057 \text{ Gt-C yr}^{-1}$, which is $\sim 40\%$ of the global continentally derived alkalinity. Their Alk^* tracer in the Atlantic has the lowest open-ocean surface concentrations despite this large riverine source. They conclude that these large riverine inputs must therefore be more than balanced by strong net CaCO_3 formation. For comparison, our estimates of CaCO_3 export between 35°S and 30°N in the Atlantic are $0.06 (0.018\text{--}0.1) \text{ Gt-C yr}^{-1}$. We note, however, that a direct comparison between these two studies remains difficult. The river input does not strictly have to be compensated by the export flux but rather by the burial flux. Also, the river input gets mixed and is subducted and transported southward by North Atlantic Deep Water and interpretation of concentrations without explicit consideration of transport and mixing is always difficult.

Regional patterns of CaCO_3 export (Fig. 10) vary among the different studies. In our approach CaCO_3 export fluxes are scaled by a rain ratio to simulated export of particulate organic carbon within each of the eight considered regions. Thus, the total export for the three Southern Ocean sectors, the Indian Ocean, the tropical and northern Pacific and the tropical and northern Atlantic is constrained by the TA^* data, but not the pattern within each of these regions.

5.2 CaCO_3 dissolution in the upper ocean

The CaCO_3 leaving the surface ocean dissolves within the water column, at the sea floor or gets buried. Berelson et al. (2007) get to global dissolution rates of $\sim 1 \text{ Gt-C yr}^{-1}$ within upper level waters, clearly higher than our TA^* -based estimate of $0.35 (0.26\text{--}0.46) \text{ Gt-C yr}^{-1}$. On a regional level, only two out of ten regional estimates by Berelson et al. (2007) are within our Latin-Hypercube uncertainty ranges; these are the estimates for the tropical Pacific (taken as 5°N to 5°S) and the

low- and mid-latitude Atlantic (40° S to 40° N) (Table 5). High dissolution rates in the range of ~ 0.1 to $0.4 \text{ mmol-C m}^{-3} \text{ yr}^{-1}$ are estimated by Barrett et al. (2014) for a transect in the upper tropical and northern North Atlantic. These values are much larger than our estimates of order $0.01 \text{ mmol-C m}^{-3} \text{ yr}^{-1}$ for the upper tropical and northern Atlantic. These high estimates are based on the measured decrease of suspended CaCO_3 particles with depth multiplied with a CaCO_3 particle settling velocity of 80 m day^{-1} and neglecting any temporal trend in the CaCO_3 particle concentration. These estimates may be affected by uncertainties in the assumed particle settling velocity. CaCO_3 particles settling velocities are reported by Jansen et al. (2002), to vary greatly (0.15 to 3440 m day^{-1}) and to be typically order one m day^{-1} for coccolithophorides and several 100 m day^{-1} for foraminifera and pteropods. Our dissolution rates would be consistent with the measured depth gradient in suspended biogenic CaCO_3 particles for an average settling velocity of a few m day^{-1} .

As mentioned above, we link the differences between the estimates of Berelson et al. (2007) and this study to methodological problems (Friis et al., 2006) associated with the TA^* -CFC age method that very likely introduce a high bias in the results of Berelson et al. (2007).

The TA^* -CFC age method relies on deduced, observation-derived TA^* concentrations and estimates of water mass age, typically derived from measurements of chlorofluorocarbons (CFCs) and their known atmospheric history. TA^* concentrations are plotted against their CFC-age and a line is fitted to this data. The higher the TA^* concentration for a given water mass, the more TA^* must have been added by dissolution to this particular water parcel according to this method. The slope of the relationship between TA^* and age is in this sense the CaCO_3 dissolution rate ($\text{mol volume}^{-1} \text{ time}^{-1}$).

The method has been criticised for its neglect of explicit transport and mixing processes. In particular, Friis et al. (2006) noted that TA^* signals ended up above the saturation horizon in their model run, even though there was explicitly no dissolution allowed to occur there. This is confirmed by our sensitivity simulations with no dissolution above the saturation horizon (Fig. 2). This finding does not depend on the choice of the numerical model as mixing within the ocean must spread the TA^* signal within the ocean and establish a surface-to-deep gradient in TA^* in the upper ocean even when all CaCO_3 dissolves at great depth. The TA^* -CFC age method does not account for such processes and assigns dissolution rates in the waters above the saturation horizon irrespective of whether or not the signal stems from the deep ocean. Our approach to combine TA^* data within an ocean transport model and the approach by Jin et al. (2006) or Lee (2001) avoid this shortcoming.

In addition, we notice, that in most regions the upper ocean TA^* distribution remains remarkably similar within our ensemble, even for very different dissolution rate profiles (see Fig. 2). And in particular, that, for a given export, k_{slow} dissolution rate profiles, with most of the dissolution at or near the ocean bottom, tend to reach highest values of TA^* both in the deep ocean and the thermocline (see also inventories in Table 2). In these cases, most of the TA^* source is added to slowly ventilated waters in the deep where it accumulates over time along deep water flow paths. This high TA^* signal

is being brought to the thermocline and eventually to the surface where TA^* is reset to its preformed value of zero. As a result, a larger TA^* gradient is established across the thermocline for deep compared to shallow dissolution. The TA^* -CFC age method would therefore have a tendency to assign higher dissolution rates in the wrong cases. This further illustrates the difficulty to uniquely relate upper ocean tracer concentrations to either dissolution or mixing processes.

5.3 $CaCO_3$ dissolution in the deep ocean

Turning to the deep ocean, Berelson et al. (2007) suggest based on sediment trap and benthic dissolution data that the particle flux below 2000 m is $0.6 \pm 0.3 \text{ Gt-C yr}^{-1}$, sea floor dissolution for sites $> 2000 \text{ m}$ averages $0.4 \pm 0.3 \text{ Gt-C yr}^{-1}$, and carbonate burial in deep marine sediments is 0.1 Gt-C yr^{-1} . Our estimate of the particle flux below 1500 m is $0.52 (0.43\text{--}0.61) \text{ Gt-C yr}^{-1}$. Thus, the compilation of sediment trap data by Berelson et al. (2007) roughly supports our particle flux at 1500 m. However, the split between sea floor dissolution and open water dissolution in the deep is different. We estimate that most of the deep ocean particle flux dissolves within the water column ($0.40 (0.32\text{--}0.47) \text{ Gt-C yr}^{-1}$ below 1500 m). The steady state burial flux for the runs with interactive sediments is $0.12 \text{ Gt-C yr}^{-1}$, corresponding to the burial of $1.95 \times 10^{13} \text{ mol Alk, yr}^{-1}$, this is comparable in magnitude but smaller than the total alkalinity input by rivers estimated to be $2.3 \times 10^{13} \text{ mol yr}^{-1}$ by Carter et al. (2014) or the burial flux estimated by Dunne et al. (2012) of $0.121 \text{ Gt-C yr}^{-1}$. We note that $CaCO_3$ formation by coral reefs and burial in shallow coastal waters (Milliman, 1993; Vecsei and Berger, 2004) is not considered in our coarse resolution model.

5.4 Parameterisations of $CaCO_3$ within Earth System Models

Uncertainties appear too large to objectively determine well-defined parameter ranges for the dissolution rates of calcite and aragonite (judging from both the TA^* and the flux data compilation). A good agreement between simulated and observed TA^* fields can be achieved irrespective of whether dissolution is assumed to depend on the calcite or aragonite saturation state or whether dissolution rates are assumed to be constant throughout the water column. We recall that the computation of the saturation state by carbonate chemistry routines across all grid cells and for each model time step poses a considerable computational burden. For simplicity and to minimise computational costs, we therefore recommend to describe $CaCO_3$ dissolution by a constant dissolution rate in Earth System Models as long as these uncertainties exist. This yields an exponential particle flux profile when assuming constant settling velocities throughout the water column. This approach is used in previous studies (Archer and Maier-Reimer, 1994; Ridgwell et al., 2007). A shortcoming of the application of an exponential particle flux profile for $CaCO_3$ is that it is not easy to account for the potential influence on dissolution of changes in environmental variables, including a decrease in saturation state as expected under ongoing ocean acidification, or in the quality, form, and size distribution of exported $CaCO_3$ particles.

735 6 Summary and conclusions

Constraining the CaCO_3 cycle comes down to three fundamental questions: how much CaCO_3 is exported from the surface ocean, where is this CaCO_3 dissolved in the water column and how much is buried in sediments. Here, we setup a probabilistic framework to constrain the CaCO_3 budget within the Bern3D EMIC with observationally-based TA^* as robust target variable. The saturation

740 state of water with respect to calcite and aragonite is prescribed using observational estimates to provide realistic boundary conditions for the CaCO_3 dissolution parameterization. In addition to the uncertainty estimates obtained by our Bayesian Latin-Hypercube framework, we also consider uncertainties related to the choice of metrics to define the assimilation target, including flux measurements as alternative target variable, uncertainties in ocean transport, and ocean sediment interactions.

745 We estimate that 0.72–1.05 Gt-C with a best estimate of 0.90 Gt-C are exported out of the surface ocean each year in the form of biogenic CaCO_3 . Of this about 37 % (0.33 (0.2–0.4) Gt-C yr⁻¹) are estimated to dissolve in the open water column above 1500 m, and 44 % (0.40 (0.32–0.47) Gt-C yr⁻¹) in the open water column below, with the reminder (0.16 (0.113–0.23) Gt-C yr⁻¹) deposited on open ocean sediments, globally. Sensitivity simulations with interactive sediments suggest that about 30% of the deposition flux dissolves back into the ocean and 70% gets buried in

750 consolidated sediments.

We find that the higher the export fluxes within the constrained, likely ranges, the more likely dissolution above the saturation horizon is needed to distribute TA^* skillfully in the model. Different kinds of dissolution schemes (with and without dissolution above saturation) achieve realistic

755 TA^* distributions within the export ranges identified. Therefore, background dissolution above the saturation horizon cannot be ruled out from this Latin-Hypercube ensemble evaluated within the Bern3D EMIC. Future progress likely depends on a better observational characterisation of particle concentrations, size distribution, and settling within the water column. Physical transport and mixing reduces concentration gradients such that conceptually different dissolution schemes (e.g.

760 dissolution permitted above saturation or not) cannot be distinguished statistically. This implies that concentrations cannot be used to infer dissolution rates directly. It also implies that dissolution parameterisations remain uncertain. For simplicity and to minimise computational costs, we suggest to use saturation-independent parameterisations of CaCO_3 dissolution within Earth System Models.

Appendix: Model evaluation and additional constraints

765 It is an essential prerequisite for the model to feature the main water masses and mixing time scales of the ocean to realistically simulate biogeochemical tracers such as TA^* . In this Appendix, we graphically document ocean model performance by comparing simulated and observation-based distributions for a range of tracers. Results are for a pre-industrial steady state of the Bern3D ocean model configuration with a horizontal resolution of 40 by 41 grid cells and 32 vertical layers as coupled to

an energy balance and sea ice module without sediment module. The atmospheric history of CFC11 is prescribed according to Bullister (2011). Here we do not account for potential changes in ocean circulation and CFC11 solubility over the industrial period.

Figure A1 provides a Taylor Diagram (Taylor, 2001) of CFC11, $\Delta^{14}\text{C}$, temperature, salinity, DIC, TA, PO_4 , oxygen, and TA^* of the standard model configuration with a constant rain ratio of 7 % and of the median TA^* of the weighted model ensemble. Modelled and observed distributions of the ventilation tracers $\Delta^{14}\text{C}$ and CFC11 are compared along a section through the Atlantic, Southern Ocean and Pacific (Figs. A2 and A3). Simulated and observation-based basin-mean vertical gradients of CFC11 and natural $\Delta^{14}\text{C}$ are compared in Fig. A4 for three simulations where diapycnal mixing is set to a low, the standard, and a high value ($k_{\text{dia}} = (0.1, 0.2, 0.5) \times 10^{-4} \text{ m}^2 \text{ s}^{-1}$). Results bracket observation-based profiles with best agreement between model and observations for the standard setup.

Acknowledgements. Many thanks to an anonymous reviewer and Wolfgang Koeve for their critical and constructive reviews. Also, thanks to Kitack Lee, Xin Jin and Niki Gruber for sharing their data, and to Raphael Roth for his contributions to the modelling framework. This work was supported by the Swiss National Science Foundation and the European Project CARBOCHANGE (264879) which received funding from the European Commission's Seventh Framework Programme (FP7/20072013). We acknowledge the support from the International Space Science Institute (ISSI). This publication is an outcome of the ISSI's Working Group on "Carbon Cycle Data Assimilation: How to consistently assimilate multiple data streams".

References

- 790 Anderson, L. A. and Sarmiento, J. L.: Redfield ratios of remineralization determined by nutrient data analysis ,
Global Biogeochemical Cycles, 8, 65–80, 1994.
- Antonov, J. I., Seidov, D., Boyer, T. P., Locarnini, R. A., Mishonov, A., Garcia, H. E., Baranova, O. K., Zweng,
M. M., and Johnson, D. R.: World Ocean Atlas 2009, Volume 2: Salinity, NOAA Atlas NESDIS 69, U.S.
Government Printing Office, Washington, D.C., 2010.
- 795 Archer, D.: A data-driven model of the global calcite lysocline, Global Biogeochemical Cycles, 10, 511–526,
doi:10.1029/96GB01521, <http://dx.doi.org/10.1029/96GB01521>, 1996.
- Archer, D. and Maier-Reimer, E.: Effect of deep-sea sedimentary calcite preservation on atmospheric CO₂
concentration, Nature, 367, 260–263, doi:10.1038/367260a0, 1994.
- Barrett, P. M., Resing, J. A., Buck, N. J., Feely, R. A., Bullister, J. L., Buck, C. S., and Landing, W. M.: Calcium
800 carbonate dissolution in the upper 1000 m of the eastern North Atlantic, Global Biogeochemical Cycles, 28,
386–397, doi:10.1002/2013GB004619, <http://dx.doi.org/10.1002/2013GB004619>, 2014.
- Behrenfeld, M. J. and Falkowski, P. G.: Photosynthetic rates derived from satellite-based chlorophyll concen-
tration, Limnology and oceanography, 42, 1–20, 1997.
- Berelson, W. M., Balch, W. M., Najjar, R., Feely, R. a., Sabine, C., and Lee, K.: Relating estimates of CaCO₃
805 production, export, and dissolution in the water column to measurements of CaCO₃ rain into sediment traps
and dissolution on the sea floor: A revised global carbonate budget, Global Biogeochemical Cycles, 21,
doi:10.1029/2006GB002803, <http://doi.wiley.com/10.1029/2006GB002803>, 2007.
- Bishop, J. K. B., Collier, R., Kettens, D., and Edmond, J.: The chemistry, biology, and vertical flux of particulate
matter from the upper 1500 m of the Panama Basin, Deep Sea Res. A., pp. 615–616, 1980.
- 810 Broecker, W. S.: “NO”, a conservative water-mass tracer , Earth Planet. Sc. Lett., 23, 100–107, 1974.
- Brown, C. W. and Yoder, J. a.: Coccolithophorid blooms in the global ocean, Journal of Geophysical Research,
99, 7467–7482, 1994.
- Bullister, J. L.: Atmospheric CFC-11, CFC-12, CFC-113, CCl₄ and SF₆ Histories.
http://cdiac.ornl.gov/ftp/oceans/CFC_ATM_Hist/, Carbon Dioxide Information Analysis Center, Oak
815 Ridge National Laboratory, US Department of Energy, Oak Ridge, Tennessee, 2011.
- Carr, M.-E.: Estimation of potential productivity in Eastern Boundary Currents using remote sensing, Deep-Sea
Research Part II: Topical Studies in Oceanography, 49, 59–80, [http://www.scopus.com/inward/record.url?
eid=2-s2.0-0036140387{&}partnerID=40{&}md5=580afff1a0622a900f0bd61548e6abbf](http://www.scopus.com/inward/record.url?eid=2-s2.0-0036140387{&}partnerID=40{&}md5=580afff1a0622a900f0bd61548e6abbf), 2002.
- Carter, B. R., Toggweiler, J. R., Key, R. M., and Sarmiento, J. L.: Processes determining the marine alkalinity
820 and carbonate saturation distributions, Biogeosciences Discussions, 11, 11 139–11 178, doi:10.5194/bgd-11-
11139-2014, <http://www.biogeosciences-discuss.net/11/11139/2014/>, 2014.
- Chung, S.-N., Lee, K., Feely, R. A., Sabine, C. L., Millero, F. J., Wanninkhof, R., Bullister, J. L., Key, R. M.,
and Peng, T.-H.: Calcium carbonate budget in the Atlantic Ocean based on water column inorganic car-
bon chemistry, Global Biogeochemical Cycles, 17, doi:10.1029/2002GB002001, [http://dx.doi.org/10.1029/
2002GB002001](http://dx.doi.org/10.1029/2002GB002001), 2003.
- 825 Doney, S. C., Lindsay, K., Fung, I., and John, J.: Natural Variability in a Stable, 1000-yr Global Coupled
Climate–Carbon Cycle Simulation, J. Climate, 19, 3033–3054, doi:10.1175/JCLI3783.1, 2006.

Dunne, J. P., Armstrong, R. a., Gnanadesikan, A., and Sarmiento, J. L.: Empirical and mechanistic models for the particle export ratio, *Global Biogeochemical Cycles*, 19, 1–16, doi:10.1029/2004GB002390, <http://www.agu.org/pubs/crossref/2005/2004GB002390.shtml>, 2005.

Dunne, J. P., Hales, B., and Toggweiler, J. R.: Global calcite cycling constrained by sediment preservation controls, *Global Biogeochemical Cycles*, 26, doi:10.1029/2010GB003935, <http://dx.doi.org/10.1029/2010GB003935>, gB3023, 2012.

Duteil, O., Koeve, W., Oschlies, A., Bianchi, D., Kriest, I., Galbraith, E. D., and Mearns, R.: A new estimate of ocean oxygen utilization points to a reduced rate of respiration in the ocean interior, *Biogeosciences*, 10, 7723–7738, 2013.

Edwards, N. R., Willmott, A. J., and Killworth, P. D.: On the Role of Topography and Wind Stress on the Stability of the Thermohaline Circulation, *J. Phys. Oceanogr.*, 28, 756–778, 1998.

Feely, R. A., Sabine, C. L., Lee, K., Millero, F. J., Lamb, M. F., Greeley, D., Bullister, J. L., Key, R. M., Peng, T.-H., Kozyr, A., Ono, T., and Wong, C. S.: In situ calcium carbonate dissolution in the Pacific Ocean, *Global Biogeochemical Cycles*, 16, 12–91, doi:10.1029/2002GB001866, <http://dx.doi.org/10.1029/2002GB001866>, 2002.

Feely, R. a., Sabine, C. L., Lee, K., Berelson, W., Kleypas, J., Fabry, V. J., and Millero, F. J.: Impact of anthropogenic CO₂ on the CaCO₃ system in the oceans., *Science (New York, N.Y.)*, 305, 362–6, doi:10.1126/science.1097329, <http://www.ncbi.nlm.nih.gov/pubmed/15256664>, 2004.

Francois, R., Honjo, S., Krishfield, R., and Manganini, S.: Factors controlling the flux of organic carbon to the bathypelagic zone of the ocean, *Global Biogeochemical Cycles*, 16, 34–1–34–20, doi:10.1029/2001GB001722, <http://doi.wiley.com/10.1029/2001GB001722>, 2002.

Friis, K., Najjar, R. G., Follows, M. J., and Dutkiewicz, S.: Possible overestimation of shallow-depth calcium carbonate dissolution in the ocean, *Global Biogeochemical Cycles*, 20, n/a–n/a, doi:10.1029/2006GB002727, <http://doi.wiley.com/10.1029/2006GB002727>, 2006.

Gangstø, R., Joos, F., and Gehlen, M.: Sensitivity of pelagic calcification to ocean acidification, *Biogeosciences*, 8, 433–458, doi:10.5194/bg-8-433-2011, <http://www.biogeosciences.net/8/433/2011/>, 2011.

Garcia, H. E., Locarnini, R. A., Boyer, T. P., Antonov, J. I., Baranova, O. K., Zweng, M. M., and Johnson, D. R.: World Ocean Atlas 2009, Volume 3: Dissolved Oxygen, Apparent Oxygen Utilization, and Oxygen Saturation, NOAA Atlas NESDIS 70, U.S. Government Printing Office, Washington, D.C., 2010a.

Garcia, H. E., Locarnini, R. A., Boyer, T. P., Antonov, J. I., K, Z. M., Zweng, M. M., Baranova, O., Johnson, D. R., Seidov, D., Mishonov, A., and D. R., J.: World Ocean Atlas 2009, Volume 4: Nutrients (phosphate, nitrate, silicate) Utilization, and Oxygen Saturation, NOAA Atlas NESDIS, U.S. Government Printing Office, Washington, D.C., 2010b.

Gattuso, J., Epitalon, J., Lavigne, H., Orr, J., Gentili, B., Hofmann, A., Proye, A., Soetaert, K., and Rae, J.: seacarb: Seawater Carbonate Chemistry, available online at <http://CRAN.R-project.org/package=seacarb>, 2010.

Gehlen, M., Bopp, L., Emprin, N., Aumont, O., Heinze, C., Ragueneau, O., Mer, D., and Copernic, P.: Reconciling surface ocean productivity, export fluxes and sediment composition in a global biogeochemical ocean model, *Biogeosciences*, 3, 521–537, doi:10.5194/bg-3-521-2006, 2006.

Griffies, S. M.: The Gent–McWilliams Skew Flux, *J. Phys. Oceanogr.*, 28, 831–841, doi:10.1175/1520-0485(1998)028<0831:TGMSF>2.0.CO;2, 1998.

Gruber, N., Sarmiento, J. L., and Stocker, T. F.: An improved method for detecting anthropogenic CO₂ in the oceans, *Global Biogeochemical Cycles*, 10, 809–837, doi:10.1029/96GB01608, <http://dx.doi.org/10.1029/96GB01608>, 1996.

Heinze, C.: Simulating oceanic CaCO₃ export production in the greenhouse, *Geophysical Research Letters*, 31, L16 308, doi:10.1029/2004GL020613, <http://doi.wiley.com/10.1029/2004GL020613>, 2004.

Heinze, C., Maier-Reimer, E., Winguth, A. M. E., and Archer, D.: A global oceanic sediment model for long-term climate studies, *Global Biogeochem. Cy.*, 13, 221–250, 1999.

Honjo, S., Manganini, S. J., Krishfield, R. a., and Francois, R.: Particulate organic carbon fluxes to the ocean interior and factors controlling the biological pump: A synthesis of global sediment trap programs since 1983, *Progress in Oceanography*, 76, 217–285, doi:10.1016/j.pocean.2007.11.003, <http://linkinghub.elsevier.com/retrieve/pii/S0079661108000025>, 2008.

Ito, T., Follows, M. J., and Boyle, E. A.: Is AOU a good measure of respiration in the ocean? , *Geophys. Res. Lett.*, 31, 2004.

Jansen, H. and Wolf-Gladrow, D. A.: Carbonate dissolution in copepod guts: A numerical model, *Mar. Ecol. Prog. Ser.*, 221, 199–207, 2001.

Jansen, H., Zeebe, R. E., and Wolf-Gladrow, D. A.: Modeling the dissolution of settling CaCO₃ in the ocean, *Global Biogeochemical Cycles*, 16, 11–1–11–16, doi:10.1029/2000GB001279, <http://dx.doi.org/10.1029/2000GB001279>, 2002.

Jin, X., Gruber, N., Dunne, J. P., Sarmiento, J. L., and Armstrong, R. a.: Diagnosing the contribution of phytoplankton functional groups to the production and export of particulate organic carbon, CaCO₃, and opal from global nutrient and alkalinity distributions, *Global Biogeochemical Cycles*, 20, 1–17, doi:10.1029/2005GB002532, <http://www.agu.org/pubs/crossref/2006/2005GB002532.shtml>, 2006.

Kalnay, E., Kanamitsu, M., Kistler, R., Collins, W., Deaven, D., Gandin, L., Iredell, M., Saha, S., White, G., Woollen, J., Zhu, Y., Chelliah, M., Ebisuzaki, W., Higgins, W., Janowiak, J., Mo, K. C., Ropelewski, C., Wang, J., Leetmaa, A., Reynolds, R., Jenne, R., and Joseph, D.: The NCEP/NCAR 40-year reanalysis project, *B. Am. Meteorol. Soc.*, 77, 437–471, doi:10.1175/1520-0477(1996)077<0437:TNYRP>2.0.CO;2, 1996.

Kanamori, S. and Ikegami, H.: Calcium–alkalinity relationship in the North Pacific , *J. Oceanogr. Soc. Jpn*, 38, 57–62, 1982.

Key, R. M., Kozyr, a., Sabine, C. L., Lee, K., Wanninkhof, R., Bullister, J. L., Feely, R. a., Millero, F. J., Mordy, C., and Peng, T.-H.: A global ocean carbon climatology: Results from Global Data Analysis Project (GLODAP), *Global Biogeochem. Cy.*, 18, GB4031, doi:10.1029/2004GB002247, <http://doi.wiley.com/10.1029/2004GB002247>, 2004.

Koeve, W., Duteil, O., Oschlies, a., Kähler, P., and Segschneider, J.: Methods to evaluate CaCO₃ cycle modules in coupled global biogeochemical ocean models, *Geoscientific Model Development*, 7, 2393–2408, doi:10.5194/gmd-7-2393-2014, <http://www.geosci-model-dev.net/7/2393/2014>, 2014.

Lee, K.: Global net community production estimated from the annual cycle of surface water total dissolved inorganic carbon, *Limnology and oceanography*, 46, 1287–1297, 2001.

- Locarnini, R. A., Mishonov, A., Antonov, J. I., Boyer, T. P., Garcia, H. E., Baranova, O. K., Zweng, M. M., and Johnson, D. R.: World Ocean Atlas 2009, Volume 1: Temperature, Tech. rep., U.S. Government Printing Office, Washington, D.C, 2010.
- Marra, J., Ho, C., and Trees, C. C.: An algorithm for the calculation of primary production from remote sensing data, Lamont-Doherty Earth Obs., Palisades, N.Y, 2003.
- McKay, M. D., Beckman, R. J., and Conover, W. J.: A comparison of three methods for selecting values of input variables in the analysis of output from a computer code, *Technometrics*, 21, 239–245, 1979.
- Milliman, J. D.: Production and accumulation of calcium carbonate in the ocean: Budget of a nonsteady state, *Global Biogeochemical Cycles*, 7, 927–957, doi:10.1029/93GB02524, <http://dx.doi.org/10.1029/93GB02524>, 1993.
- Milliman, J. D., Troy, P. J., Balch, W. M., Adams, A. K., Li, Y. H., and Mackenzie, F. T.: Biologically mediated dissolution of calcium carbonate above the chemical lysocline?, *Deep-Sea Res. Pt. 1*, 46, 1653–1669, doi:10.1016/S0967-0637(99)00034-5, 1999.
- Mucci, A.: The solubility of calcite and aragonite in seawater at various salinities, temperatures, and one atmosphere total pressure, *American Journal of Science*, 1983.
- Müller, S. A., Joos, F., Edwards, N. R., and Stocker, T. F.: Water mass distribution and ventilation time scales in a cost-efficient, three-dimensional ocean model, *J. Climate*, 19, 5479–5499, doi:10.1175/JCLI3911.1, 2006.
- Müller, S. A., Joos, F., Edwards, N. R., and Stocker, T. F.: Modeled natural and excess radiocarbon: Sensitivities to the gas exchange formulation and ocean transport strength, *Global Biogeochem. Cy.*, 22, 14 pp, doi:10.1029/2007GB003065, 2008.
- Najjar, R. G., Orr, J., Sabine, C. L., and Joos, F.: Biotic-HOWTO. Internal OCMIP Report, Tech. rep., LSCE/CEA Saclay, Gif-sur-Yvette, France, 1999.
- Orr, J. and Najjar, R. G.: Abiotic-HOWTO. Internal OCMIP Report, Tech. rep., LSCE/CEA Saclay, Gif-sur-Yvette, France, 1999.
- Parekh, P., Joos, F., and Müller, S. A.: A modeling assessment of the interplay between aeolian iron fluxes and iron-binding ligands in controlling carbon dioxide fluctuations during Antarctic warm events, *Paleoceanography*, 23, PA4202, doi:10.1029/2007PA001531, 2008.
- Ridgwell, A., Hargreaves, J. C., Edwards, N. R., Annan, J. D., Lenton, T. M., Marsh, R., Yool, A., and Watson, A.: Marine geochemical data assimilation in an efficient Earth System Model of global biogeochemical cycling, pp. 87–104, 2007.
- Ritz, S. P., Stocker, T. F., and Severinghaus, J. P.: Noble gases as proxies of mean ocean temperature: sensitivity studies using a climate model of reduced complexity, *Quaternary Sci. Rev.*, 30, 3728–3741, doi:10.1016/j.quascirev.2011.09.021, 2011.
- Roth, R., Ritz, S. P., and Joos, F.: Burial-nutrient feedbacks amplify the sensitivity of atmospheric carbon dioxide to changes in organic matter remineralisation, *Earth System Dynamics*, 5, 321–343, doi:10.5194/esd-5-321-2014, <http://www.earth-syst-dynam.net/5/321/2014>, 2014.
- Sabine, C. L., Feely, R. A., Key, R. M., Bullister, J. L., Millero, F. J., Lee, K., Peng, T.-H., Tilbrook, B., Ono, T., and Wong, C. S.: Distribution of anthropogenic CO₂ in the Pacific Ocean, *Global Biogeochemical Cycles*, 16, 17–30, doi:10.1029/2001GB001639, <http://dx.doi.org/10.1029/2001GB001639>, 2002a.

- 945 Sabine, C. L., Key, R. M., Feely, R. A., and Greeley, D.: Inorganic carbon in the Indian Ocean: distribution and dissolution processes, *Global Biogeochemical Cycles*, 16, 2002b.
- Sarmiento, J. L. and Gruber, N.: *Ocean biogeochemical dynamics*, Princeton University Press, 2006.
- Sarmiento, J. L., Dunne, J., Gnanadesikan, a., Key, R. M., Matsumoto, K., and Slater, R.: A new estimate of the CaCO_3 to organic carbon export ratio, *Global Biogeochemical Cycles*, 16, 54–1–54–12, doi:10.1029/2002GB001919, <http://doi.wiley.com/10.1029/2002GB001919>, 2002.
- 950 Schmittner, A., Urban, N. M., Keller, K., and Matthews, D.: Using tracer observations to reduce the uncertainty of ocean diapycnal mixing and climate-carbon cycle projections, *Global Biogeochemical Cycles*, 23, 1–16, doi:10.1029/2008GB003421, 2009.
- Steinacher, M. and Joos, F.: Transient Earth system responses to cumulative carbon dioxide emissions: linearities, uncertainties, and probabilities in an observation-constrained model ensemble, *Biogeosciences*, 13, 1071–1103, <http://www.biogeosciences.net/13/1071/2016>, 2016.
- 955 Steinacher, M., Joos, F., Bopp, L., Cadule, P., Doney, S. C., Gehlen, M., Schneider, B., and Segschneider, J.: Projected 21st century decrease in marine productivity : a multi-model analysis, pp. 7933–7981, 2009.
- Steinacher, M., Joos, F., and Stocker, T. F.: Allowable carbon emissions lowered by multiple climate targets., *Nature*, 499, 197–201, doi:10.1038/nature12269, <http://www.ncbi.nlm.nih.gov/pubmed/23823728>, 2013.
- 960 Taylor, K. E.: Summarizing multiple aspects of model performance in a single diagram, *Journal of Geophysical Research*, 106, 7183, doi:10.1029/2000JD900719, 2001.
- Tschumi, T., Joos, F., Gehlen, M., and Heinze, C.: Deep ocean ventilation, carbon isotopes, marine sedimentation and the deglacial CO_2 rise, *Clim. Past*, 7, 771–800, doi:10.5194/cp-7-771-2011, 2011.
- 965 Vecsei, A. and Berger, W. H.: Increase of atmospheric CO_2 during deglaciation: Constraints on the coral reef hypothesis from patterns of deposition, *Global Biogeochem. Cy.*, 18, GB1035, doi:10.1029/2003GB002147, 2004.
- Volk, T. and Hoffert, M. I.: Ocean carbon pumps: Analysis of relative strengths and efficiencies in ocean-driven atmospheric CO_2 changes, *The Carbon Cycle and Atmospheric CO_2 : Natural Variations Archean to Present*, pp. 99–110, 1985.
- 970 Wilson, J. D., Barker, S., and Ridgwell, A.: Assessment of the spatial variability in particulate organic matter and mineral sinking fluxes in the ocean interior: Implications for the ballast hypothesis, *Global Biogeochemical Cycles*, 26, doi:10.1029/2012GB004398, <http://dx.doi.org/10.1029/2012GB004398>, 2012.
- Wolf-Gladrow, D. A., Zeebe, R. E., Klaas, C., Körtzinger, A., and Dickson, A. G.: Total alkalinity: The explicit conservative expression and its application to biogeochemical processes, *Marine Chemistry*, 106, 287–300, doi:10.1016/j.marchem.2007.01.006, <http://www.sciencedirect.com/science/article/pii/S0304420307000047>, 2007.
- 975

Table 1. Overview on suggested ways of regression for TA^0 either including two (S , PO) or three (S , PO , T) explanatory variables and either as a global or basin-wide fit. The last column shows the root mean square deviation relative to the Global B3D estimate.

	Equation [ueq kg^{-1}]	Mean [mol m^{-3}]	RMSE from Global B3D [mol m^{-3}]
Gruber et al. (1996):	$\text{TA}^0 = (367.5 + 59.9 \text{ psu}^{-1} \cdot S + 0.074 \text{ kg umol}^{-1} \cdot \text{PO}) \text{ ueq kg}^{-1}$	2.377	0.012
Feely et al. (2002):	$\text{TA}^0 = 148.7 + 61.36 \cdot S + 0.0941 \cdot \text{PO} - 0.582 \cdot T_{\text{pot}}$	2.39	0.0015
Friis et al. (2006):	as Feely et al. (2002)		
Global B3D 2V:	$\text{TA}^0 = (297.51 + 56.399 \text{ psu}^{-1} \cdot S + 0.1259 \text{ kg umol}^{-1} \cdot \text{PO}) \text{ ueq kg}^{-1}$	2.387	0.0022
Global B3D:	$\text{TA}^0 = (345.64 + 56.03 \text{ psu}^{-1} \cdot S + 0.069 \text{ kg umol}^{-1} \cdot \text{PO} - 0.9^\circ \text{C}^{-1} \cdot T) \text{ ueq kg}^{-1}$	2.389	0
Regional B3D:		2.385	0.0047
Atlantic	$\text{TA}^0 = (688.15 + 44.97 \text{ psu}^{-1} \cdot S + 0.129 \text{ kg umol}^{-1} \cdot \text{PO} + 1.34^\circ \text{C}^{-1} \cdot T) \text{ ueq kg}^{-1}$		
Pacific	$\text{TA}^0 = (381.05 + 55.26 \text{ psu}^{-1} \cdot S + 0.049 \text{ kg umol}^{-1} \cdot \text{PO} - 1.20^\circ \text{C}^{-1} \cdot T) \text{ ueq kg}^{-1}$		
Indian	$\text{TA}^0 = (637.70 + 47.38 \text{ psu}^{-1} \cdot S + 0.078 \text{ kg umol}^{-1} \cdot \text{PO} - 0.54^\circ \text{C}^{-1} \cdot T) \text{ ueq kg}^{-1}$		

Gruber et al. (1996) is the original, global regression based on two explanatory variables (based on their 1996 database), Feely et al. (2002) and Friis et al. (2006) are both taken from Sabine et al. (2002b) also including temperature as an additional explanatory variable and fitted to a Pacific subset. Global B3D 2V is a global regression with two explanatory variables (based on WOA09 Locarnini et al., 2010; Antonov et al., 2010; Garcia et al., 2010a, b) on the Bern3D grid ($32 \times 40 \times 41$ grid cells), Global B3D includes three explanatory variables and Regional B3D further distinguishes each basin, separately.

Table 2. Results from sensitivity simulations. CaCO_3 export and TA^* inventories for different physical mixing (diapycnal mixing coefficient, k_{dia}) and CaCO_3 dissolution schemes. For these illustrative simulations, calcite and aragonite particles were assigned equal parameter values and 10 % of export is assumed to be in the form of aragonite. Fast: $k_0 = 10 \text{ day}^{-1}$, $n = 1$, $f_{\text{calc}} = 0.9$, $k_{\text{bg}} = 0$; slow: $k_0 = 0.16 \text{ day}^{-1}$, $n = 2$, $f_{\text{calc}} = 0.9$, $k_{\text{bg}} = 0$; constant: $k_{\text{bg}}/v = 1/2900 \text{ m}^{-1}$.

	$k_{\text{dia, low}}$ $0.1 \times 10^{-4} \text{ m}^2 \text{ s}^{-1}$	$k_{\text{dia, ref}}$ $0.2 \times 10^{-4} \text{ m}^2 \text{ s}^{-1}$	$k_{\text{dia, high}}$ $0.5 \times 10^{-4} \text{ m}^2 \text{ s}^{-1}$
Export [Gt-C yr^{-1}]	0.727	0.818	1.071
TA* Inventory [Pmol C] (fraction of which lies above $\Omega_{\text{calc}} = 1$)			
Fast	47.81 (54 %)	47.87 (52 %)	44.5 (51 %)
Slow	62.49 (45 %)	63.01 (44 %)	58.93 (43 %)
Constant	38.12 (52 %)	38.35 (51 %)	35.44 (50 %)

Table 3. Mean sediment burial fluxes [Gt-C yr^{-1}] and observation-based vs. simulated TA^* inventories [Pmol C]. $\text{TA}^{*\text{sedcorr}}$ are estimated by comparing output from the 14 Bern3D simulations that agree best with observation-based TA^* concentrations and with and without sediment module. The median estimate is from the constrained 1000 member ensemble (without sediment module).

	Atlantic	Pacific	Indian	Global
Burial flux [Gt-C yr^{-1}]	0.011	0.062	0.042	0.117
TA^* Inventory [Pmol C]				
$\text{TA}^{*\text{obs}}$	2.5	27.9	7.0	37.4
$\text{TA}^{*\text{sedcorr}}$	0.9	4.6	1.4	6.9
$\text{TA}^{*\text{obs}+\text{sedcorr}}$	3.4	32.4	8.4	44.1
$\text{TA}^{*\text{median}}$	5.4	31.1	8.7	45.5




Figure 1. The TA^* tracer captures exclusively the influence of CaCO_3 dissolution on alkalinity. An observationally-based estimate of the TA^* distribution [mmol m^{-3}] (based on GLODAP Key et al. (2004) and WOA09 Locarnini et al., 2010; Antonov et al., 2010; Garcia et al., 2010a, b) yielding an inventory of $\sim 37.5 \text{ Pmol C}$ of which $\sim 41\%$ comes to lie above the calcite saturation horizon. The calcite ($\sigma_{\text{calcite}}=1$) and aragonite ($\sigma_{\text{aragonite}}=1$) saturation horizon are shown by the blue lines. Displayed are results for a cross-section through the Atlantic (25° W), across the Southern Ocean (58° S) into the Pacific, and through the Pacific (175° W).

Table 4. Constrained fluxes (median and 68 % confidence interval, c.i.) of biogenic CaCO_3 [Gt-C yr^{-1}].

	Atlantic		Pacific		Indian		Global	
	median	c.i.	median	c.i.	median	c.i.	median	c.i.
Export at 75 m								
70 – 30°N	0.021	[0.007-0.037]	0.081	[0.027-0.134]				
30°N – 35°S	0.061	[0.018-0.103]	0.307	[0.149-0.457]	0.143	[0.053-0.223]		
>35°S	0.041	[0.014-0.07]	0.174	[0.057-0.254]	0.088	[0.03-0.14]		
70°N – 90°S	0.121	[0.078-0.171]	0.549	[0.391-0.697]	0.23	[0.137-0.316]	0.897	[0.72-1.049]
Dissolution in waters shallower than 1500 m								
70 – 30°N	0.006	[0.002-0.011]	0.048	[0.015-0.086]				
30°N – 35°S	0.015	[0.004-0.03]	0.118	[0.049-0.2]	0.043	[0.016-0.08]		
>35°S	0.01	[0.003-0.021]	0.043	[0.013-0.082]	0.02	[0.006-0.042]		
70°N – 90°S	0.033	[0.018-0.054]	0.216	[0.125-0.32]	0.064	[0.032-0.11]	0.328	[0.202-0.444]
Deposition on sediments shallower than 1500 m								
70 – 30°N	0.003	[0.001-0.005]	0.008	[0.002-0.014]				
30°N – 35°S	0.003	[0.001-0.006]	0.008	[0.004-0.013]	0.015	[0.005-0.024]		
>35°S	0	[0.0-0.001]	0.007	[0.002-0.011]	0	[0.0-0.001]		
70°N – 90°S	0.007	[0.004-0.01]	0.023	[0.016-0.029]	0.016	[0.006-0.025]	0.045	[0.034-0.056]
Dissolution in waters deeper or equal to 1500 m								
70 – 30°N	0.008	[0.003-0.013]	0.014	[0.003-0.035]				
30°N – 35°S	0.03	[0.011-0.048]	0.133	[0.068-0.2]	0.053	[0.022-0.087]		
>35°S	0.023	[0.009-0.038]	0.085	[0.031-0.129]	0.043	[0.015-0.07]		
70°N – 90°S	0.06	[0.039-0.083]	0.233	[0.165-0.305]	0.097	[0.061-0.139]	0.395	[0.32-0.469]
Deposition on sediments deeper or equal to 1500 m								
70 – 30°N	0.003	[0.0-0.007]	0.002	[0.001-0.007]				
30°N – 35°S	0.009	[0.001-0.021]	0.031	[0.012-0.062]	0.019	[0.005-0.039]		
>35°S	0.005	[0.001-0.011]	0.02	[0.006-0.041]	0.016	[0.005-0.03]		
70°N – 90°S	0.018	[0.009-0.033]	0.056	[0.028-0.099]	0.035	[0.018-0.065]	0.116	[0.071-0.179]
Deposition on all sediments								
70 – 30°N	0.006	[0.001-0.012]	0.011	[0.003-0.021]				
30°N – 35°S	0.013	[0.002-0.027]	0.04	[0.017-0.073]	0.036	[0.011-0.061]		
>35°S	0.005	[0.001-0.012]	0.028	[0.008-0.051]	0.016	[0.005-0.03]		
70°N – 90°S	0.025	[0.014-0.043]	0.081	[0.049-0.126]	0.052	[0.026-0.087]	0.164	[0.113-0.227]

Table 5. Dissolution rates of biogenic CaCO_3 [Gt-C yr^{-1}] in the upper water column (200–1500 m depth levels) based on the TA^* -CFC age method as summarised in Berelson et al. (2007), and as constrained by TA^* data in our Latin-Hypercube ensemble (including the sediment correction, median and 68 % c.i.). The dissolution estimates by Berelson et al. (2007) were assigned an estimated uncertainty of ~ 50 %.

Location	Berelson et al. (2007)	This study median	c.i.
Atlantic			
>40°N	0.060	0.006	[0.002-0.011]
40°N - 40°S	0.010	0.02	[0.008-0.036]
>40°S	0.040	0.009	[0.003-0.017]
Pacific			
>40°N	0.070	0.039	[0.013-0.067]
40°N - 5°N	0.330	0.063	[0.032-0.099]
5°N - 5°S	0.020	0.035	[0.015-0.058]
5°S - 40°S	0.000	0.039	[0.019-0.066]
>40°S	0.160	0.034	[0.011-0.061]
Indian			
40°N - 40°S	0.160	0.062	[0.026-0.104]
>40°S	0.140	0.021	[0.009-0.038]
Total	1.0	0.346	[0.225-0.461]




Figure 2. Results from sensitivity simulations applying three different illustrative dissolution schemes. Left column: CaCO_3 dissolution rates. Right column: the resulting steady state TA^* for these three contrasting parameterisations of the CaCO_3 dissolution rate. The dissolution rate is set to increase fast (top row), slowly (middle row) with undersaturation of CaCO_3 , or is set constant throughout the water column (bottom row). The standard version of the Bern3D model without sediment module is applied and the fraction of CaCO_3 export in the form of aragonite is set to 10 %. At least 43 % of the dissolution signal is simulated above the calcite saturation horizon irrespective of whether dissolution is allowed to occur above the saturation horizon (bottom row) or not (middle and top row). This points to the importance of physical transport in shaping the distribution of TA^* .




Figure 3. Influence of ocean–sediment fluxes on TA^* . The mean offset in TA^* between simulations with and without sediment module is shown for the 14 ensemble members with the highest skill. Displayed are results for a cross-section through the Atlantic (25° W), across the Southern Ocean (58° S) into the Pacific, and through the Pacific (175° W). Note the colourbar.




Figure 4. Model ensemble vs. observation-based basin-mean TA^* profiles. The grey shading shows the unconstrained prior and the green shading shows the constrained (68 % confidence interval) distribution of the model ensemble. Lines represent the median of the constrained ensemble (green), observation-based TA^* (black dashes), and observation-based TA^* corrected for a sediment burial flux of $0.12 \text{ Gt-C yr}^{-1}$ (black solid). The corresponding Southern Ocean sector is included in the averaging.




Figure 5. Observation-based vs. simulated TA^* . Left column: modelled (median) TA^* , observed TA^* , and their difference in a cross-section through the Atlantic (25° W), Southern Ocean (58° S), and Pacific (175° W). Right column: the same for a cross-section along 95° E in the Indian Ocean. The correlation coefficient, relative standard deviation ($\sigma_{rel}^{obs.}$) and root mean square error (RMSE) between the simulated and observation-based fields are: 0.83, 1.09, and 17.7 mmol m^{-3} in the Atlantic; 0.88, 0.8, and 23.8 mmol m^{-3} in the Pacific; 0.87, 0.86, and 19.8 mmol m^{-3} in the Indian Ocean.




Figure 6. Probability density functions of basin-wide and global CaCO_3 fluxes as constrained by the observation-based TA^* distribution (corrected for a sediment burial flux of $0.12 \text{ Gt-C yr}^{-1}$). Note the different scaling of the x axes.




Figure 7. Top: median CaCO_3 export, globally 0.90 ($0.72\text{--}1.05$) Gt-C yr^{-1} . Bottom: median CaCO_3 fluxes reaching the ocean floor, globally 0.164 ($0.113\text{--}0.227$) Gt-C yr^{-1} . Note the different scaling of the colourbars.




Figure 8. Constrained open-water dissolution rate profiles for aragonite (red) and calcite (olive). Ensemble medians (solid lines) and 68 % confidence intervals (shadings) are for spatial averages across individual regions and including only grid cells where the model water column extends to a depth of 5000 m.




Figure 9. The simulated TA* inventory vs. the simulated CaCO₃ export of each ensemble member, coloured according to model skill (top) and background dissolution rate (bottom) for the Atlantic, Pacific, and Indian Ocean (columns). Each circle represents results from an individual simulation and its colour indicates the regional skill score (top) or the value of the background dissolution coefficient for calcite (bottom). Green shadings show the 68% confidence range in basin-wide TA* inventories and CaCO₃ export of the constrained model ensemble and black-dashed lines indicate the estimated TA* inventories based on the observations including the sediment correction. Runs with low and high background dissolution can skillfully represent the TA* distribution. Runs with a higher CaCO₃ export tend to require a higher background dissolution to achieve a good skill.




Figure 10. Observationally-constrained CaCO_3 export fields as estimated by different studies. Global total export is estimated to 0.5 by Sarmiento and Gruber (2006), to 1.14 by Jin et al. (2006), to 1.1 by Lee (2001), and to $0.90 \text{ Gt-C yr}^{-1}$ by this study.




Figure 11. Zonally-integrated export fluxes of the constrained Bern3D-model ensemble (median and 68 % confidence interval) compared to estimates by Sarmiento and Gruber (2006), Jin et al. (2006) and Lee (2001).




Figure A1. Taylor diagram (Taylor, 2001) for global and basin-wide volume-weighted oceanic tracer distributions as simulated by the Bern3D standard setup with $k_{\text{dia}} = 0.2 \times 10^{-4} \text{ m}^2 \text{ s}^{-1}$. Observation-derived fields are taken from GLODAP (Key et al., 2004) and the World Ocean Atlas 2009 (Locarnini et al., 2010; Antonov et al., 2010; Garcia et al., 2010a, b).




Figure A2. (Top) Distributions of (natural) $\Delta^{14}\text{C}$ of DIC [‰] as simulated with a standard k_{dia} of $0.2 \times 10^{-4} \text{ m}^2 \text{ s}^{-1}$ at steady state and (bottom) as estimated from $\Delta^{14}\text{C}$ measurements (Key et al., 2004). The correlation coefficient, $\sigma_{\text{rel}}^{\text{obs}}$, and RMSEs are: 0.89, 1.46, 23.77 ‰ in the Atlantic; 0.95, 0.97, 16.28 ‰ in the Pacific; 0.9, 0.78, 21.1 ‰ in the Indian Ocean. The section displayed is through the Atlantic (25° W), the Southern Ocean (58° S), and the Pacific (175° W).




Figure A3. (Top) Distributions of CFC11 [nmol m⁻³] as simulated with a standard k_{dia} of $0.2 \times 10^{-4} \text{ m}^2 \text{ s}^{-1}$ averaged over the model years 1990-2000 and (bottom) as observed (Key et al., 2004). The correlation coefficient, $\sigma_{\text{rel.}}^{\text{obs.}}$ and RMSEs are: 0.9, 1.01, 0.45 nmol m⁻³ in the Atlantic; 0.94, 1, 0.29 nmol m⁻³ in the Pacific; 0.83, 0.92, 0.54 nmol m⁻³ in the Indian Ocean. The section displayed is through the Atlantic (25° W), the Southern Ocean (58° S), and the Pacific (175° W).




Figure A4. Basin-wide average profiles for (natural) $\Delta^{14}\text{C}$ of DIC [‰] at steady state (top) and for CFC11 [nmol m^{-3}] averaged over the model years 1990-2000 (bottom). Results are for simulations with a low (green dotted), standard (green solid), and high (green dashes) value of the diapycnal mixing coefficient k_{dia} (0.1 , 0.2 and $0.5 \times 10^{-4} \text{ m}^2 \text{ s}^{-1}$). Observation-based estimates are in black. The corresponding Southern Ocean sector is included in the averaging.




Figure A5. The global sediment trap data collection of Wilson et al. (2012, see their auxiliary material at doi:10.1029/2012GB004398) on the Bern3D grid. The measurements are located at different depths (> 1500 m). If more than one measurement was assigned to the same cell, the mean was chosen.




Figure A6. As Fig. 9 but coloured according to model skill with respect to the sediment trap database evaluated by basin. This target variable constrains export and dissolution to wider ranges (yellow to red colours) as compared to the TA^* target (light green shading). Generally, high skill scores with respect to regional, sediment-corrected TA^* are also associated with high skill scores with respect to regional fluxes. Only few models are in good agreement with the few flux measurements in the Indian Ocean.