



College of Soil and Water Conservation
BEIJING FORESTRY UNIVERSITY
Qinghua East Road 35, HAIDIAN DISTRICT
BEIJING 100083
P. R. CHINA
Tel: +86-(0)10-623388097
Fax: +86-(0)10-62337873

8

9 **Response to the Reviews of MS No.: bg-2014-536**

10

11 **Journal:** *Biogeosciences (BG)*

12 **Title:** *Responses of energy partitioning and surface resistance to drought in a poplar*
13 *plantation in northern China*

14 **Author(s):** *M.C. Kang et al.*

15 **MS No.:** *bg-2014-536*

16 **MS Type:** *Research Article*

17

June 23, 2015

18 **Dear Associate Editor, Prof. &, Dr. Christopher A. Williams,**

19

20 First of all, we thank you very much for your insightful review and comment on our paper. In
21 particular, we really appreciate your constructive suggestions on the poplar plantations effects
22 on the adjacent ecosystems and livelihoods, which have definitely broaden our discussion and
23 improve the quality of our paper. As you and other reviewers pointed out that it is hard to
24 distinct drought for our four year study as irrigation was applied in years when precipitation
25 was less than average, we have changed our title into “Energy partitioning and surface
26 resistance of a poplar plantation in Northern China”. Of course, we presented our result and
27 discussion accordingly. Overall, we have revised the whole paper and answered all the
28 questions and comments raised accordingly.

29

30 All our co-authors have contributed substantially to the revision of the paper and we believe
31 that the revised paper meets the high quality requirement. The reply to each comment and
32 question is presented one by one following this cover letter.

33

34 Should you have any inquiries about this resubmission, please feel free to contact me at any
35 time:

36

Zhiqiang Zhang, Professor & Ph D

37

College of Soil and Water Conservation

38

Beijing Forestry University

39

Qinghua East Road 35#, Haidian District

40

Beijing 100083, P.R. China

41

Tel. 00 86 10 62338097

42

Fax 00 86 10 62337873


43

Email: Zhqzhang@bjfu.edu.cn or Zhqzhang_bjfu@yahoo.com.cn

1
2
3

4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32

Best regards,
Yours sincerely

A handwritten signature in cursive script that reads "Zhigang Zhang". The signature is written in black ink and is positioned below the typed name "Yours sincerely".

On behalf of all authors

See Reply on the next page.

1 **Reply to comments:**

2 Associate Editor Initial Decision: Reconsider after major revisions (12 May 2015) by
3 Christopher A. Williams

4 Comments to the Author:

5 This is a solid study that should be deserving of publication. It presents a valuable dataset and
6 lends new insights not only into the hydrologic status of these particular poplar plantations
7 located in a water-limited, mid-latitude environment but also into fundamental relationships
8 among attributes of the surface water and energy balances. However, in my view, and similar
9 to statements by the anonymous reviewers, the interpretations need to be corrected in a few
10 places, both in terms of relationships among reported variables (e.g. R_s vs. LAI and LE/LE_{eq}
11 vs. R_s), as well as in terms of the sustainability and perceived threats of the plantation activity.

12 **1. Assessment of Sustainability Is Still Not Aligned With This Study's Findings:** The
13 sustainability of the plantation operation is inadequately quantified and this term should be
14 either dropped or more carefully employed. The addition of Section 4.3 does not ameliorate the
15 situation. What might be unsustainable is the use of groundwater, either directly by the trees or
16 also via irrigation, to support the plantation, particularly if the rate of groundwater extraction
17 exceeds the rate of recharge. That has not been assessed here, and while that outcome is
18 plausible or even likely, the methods employed in this study are insufficient to fully substantiate
19 the claim. The main finding is that the poplar plantations evapotranspire all of the water that is
20 supplied via precipitation and irrigation. That is an important finding but does not directly
21 address the question of sustainability. The paper does not clearly document how the
22 establishment and operation of these poplar plantations is a threat to adjacent ecosystems or
23 may compromise the long-term sustainability of livelihoods in the region. I agree that
24 discussion of these points is acceptable and even warranted but I urge the authors to do so within
25 the bounds of what is supported scientifically in this study. For example, I would argue that the
26 following statements could be part of the discussion: In wet years, the plantation itself is in
27 hydrologic balance with the water that arrives as precipitation, with evapotranspiration
28 consuming nearly all of the precipitation. The same is true in dry years, but irrigation increases
29 ET even further by depleting groundwater. Even if the plantations are in hydrologic balance
30 with water delivered as precipitation, their existence and operation could be a threat to adjacent
31 ecosystems and livelihoods if those rely on runoff or groundwater recharge from the areas
32 where the plantation has been sited. In the absence of the plantations it is likely that groundwater

1 recharge would increase, especially given the sandy textured soil that tends to allow rapid
2 infiltration and percolation as well as limit moisture delivery to the atmosphere directly from
3 the soil surface itself. While poplar plantation growth in this water-limited location might be
4 sustained by the modest precipitation in the region, it could still be unsustainable for the broader
5 context of the region's ecosystems and livelihoods. To truly assess this one would need to study
6 (a) the surface water balance at the same site pre-plantation or at an adjacent, similar site but
7 without a plantation, and/or (b) groundwater levels both spatially and temporally. New text
8 appears notes that Zhang et al. 2014 documented water table decline over the last 30 years, so
9 some of this may well already be substantiated. If that's the case, the argumentation needs to be
10 rebuilt to note that connection more clearly.

11 *Reply: We really appreciate your great and constructive suggestions which help us broaden our*
12 *discussion and improve the quality of our paper. Based on your review comments, we have*
13 *dropped the “sustainability” term and taken your suggestions into our discussion (see in*
14 *Revised MS P38, L24-P39, L5). We have revised the text “the sustainability of these plantations*
15 *needs to be evaluated” as “To further understand the acclimation of poplar species to semiarid*
16 *environment and evaluate the potential impacts of these plantations on the broader context of*
17 *the region’s water supply.”(see in Revised MS P22, L6-9), and changed “(3) evaluate the long-*
18 *term sustainability of poplar plantations in a water limiting region in northern China.” into*
19 *“(3) evaluate the long-term potential impact of poplar plantations on the availability of water*
20 *for adjacent ecosystems and livelihoods in water-limited region” (see in Revised MS P24, L21-*
21 *23) ;*

22

23 **2. Cannot State in the Abstract That the Groundwater Table is in Decline:** This is not
24 studied or reported here. You could potentially report that, or certainly discuss it, but it is not a
25 major finding from the present study.

26 *Reply: We have changed the text the text “In general, ... long-term sustainability and livelihoods*
27 *in the region” into “All physiological and bioclimatological metrics indicated that the water*
28 *demands of the poplar plantation were greater than the amount available through precipitation,*
29 *highlighting the poor match of a water-intensive species like poplar for this water limited*
30 *region.” (see in Revised MS P22, L27-29);*

31

1 **3. Cannot Conclude that Fast-growing, Water-intensive Species Like the Poplar**
2 **Plantation Are Poorly Adapted for Water-Limited Regions:** From the data presented, the
3 plantation growth appears healthy and well-adapted even to this dry setting. While irrigation
4 was applied in the drier years, it is not clear how much growth or damage (mortality?) would
5 have occurred in the dry years if irrigation had not been applied. Statements about the plantation
6 being poorly adapted for water limited regions must be removed, for example the last statement
7 of the conclusions.

8 *Reply: Thanks for helping us to clarify and make our statement to the point. We have revised*
9 *the last statement of the conclusions as “Even at mean long-term precipitation, the water*
10 *demand of poplar plantation may consume nearly all of it and leave little for run-off and*
11 *groundwater recharge in this semi-arid region, potentially compromising the region’s*
12 *ecosystems and livelihoods.”(see in Revised MS P39 L25-28);*

13

14 **4. Drought Impacts Are Only Weakly Assessed:** Because of the irrigation it is not really
15 possible to quantify drought response. The irrigation clearly had the potential to mediate the
16 impacts of lower precipitation in 2006 and 2009, thus the true effects of drought cannot be
17 quantified.

18 *Reply: Yes, we could not quantify the true effects of drought on energy partitioning and bulk*
19 *resistance parameters of poplar plantation due to the application of irrigation in dry years. As*
20 *the effects of drought were still notable even under the irrigation; it is, therefore, logical to infer*
21 *that the effects of drought would be much clearer if the irrigation had not been applied in dry*
22 *years. We revised the statements in the abstract “The partitioning of available energy to latent*
23 *heat (LE) flux decreased from 0.62 to 0.53 under meteorological drought” as “The partitioning*
24 *of available energy to latent heat flux (LE) decreased from 0.62 to 0.53 under mediated*
25 *meteorological drought by irrigation applications”(see in Revised MS P22 L15-17);*

26

27 **5. Correlation Between Resistance and LAI at a Seasonal Scale Does Not Indicate**
28 **Influence of Other Factors (P15):** On the contrary, doesn't this evidence primary control by
29 LAI, which varies seasonally, at least when water is less limiting? Larger scatter in the
30 relationship for the dry year(s) does indicate additional control by other factors. The
31 argumentation needs to be corrected and clarified here.

1 *Reply: Based on your suggestion, we have corrected the statement “The strong correlation*
2 *between R_s and LAI in wet years (Fig.6) that R_s in dry years was also influenced by other*
3 *physiological and non-physiological...” into “Compared to the strong correlation between R_s*
4 *and LAI in wet years, the increased scatter in the R_s -LAI relationship during dry years (Fig.6)*
5 *suggests that R_s in dry years was also influenced by other physiological and non-physiological...”*
6 *(see in Revised MS P36, L23-25);*

7

8 **6. Decline in LE/LEeq with Increasing R_s is Required by the Way R_s is Derived from**
9 **Inversion of the Penman-Monteith Equation (P16):** This should not be presented as a finding
10 given that the relationship is essentially implied by the formulation.

11 *Reply: We have revised the text “Similar to Baldocchi (1994), LE/LEeq declined with increasing*
12 *R_s during the growing season (Fig.7),” as “As essentially implied by the Penman-Monteith*
13 *equation, LE/LEeq exponentially related with R_s during the growing season (Fig.7).”(see in*
14 *Revised MS P37, Line 19-22);*

15

16 Considerable grammar, syntax, spelling, and diction errors remain, particularly (but not only)
17 in the text that has been revised or added as part of the revision. The manuscript must be edited
18 to correct these errors before it can be accepted for publication.

19 *Reply: Thanks for your suggestion for ensuring the publication quality. All the native English*
20 *speaking co-authors of this paper have edited grammar, syntax, spelling, and diction errors.*
21 *Revised content can be seen in authors' changes in manuscript.*

22

23 **P2**

24 L10: "latent heat" to "latent heat flux"

25 L11: "sensible heat" to "sensible heat flux"

26 L20: "thread" to "threat"

27 *Reply: Thanks for your careful reading. Corrected (see in Revised MS P22, L16, 17,30).*

28

29 **P3**

1 L10: "pf" to "of"

2 L28: "most of previous and current are only concentrated on" to "most previous and current
3 studies concentrated only on"

4 L30: "is central important for" to "is of central importance for"

5 L27 to 31: recommend edit from "Whereas ..." to "Most prior work has concentrated primarily
6 on the water balance of forest ecosystems, with less emphasis on the relationship of forest
7 ecosystems to their environmental setting. Much can be learned from exploring the partitioning
8 of available energy and ecosystem response to meteorological forcing such as droughts. Not
9 only are they of central importance for understanding water and carbon balances (), but they
10 also help elucidate the degree to which forest water use is in balance with supply from
11 precipitation, and hence the degree to which plantations located in water limited regions are
12 sustainable in the long-term."

13 *Reply: Thanks for your help on improving the quality of the paper. We appreciated and*
14 *corrected each of suggestions (see Revised MS in P23, L10; L35-L7(P24));*

15

16 **P4**

17 L5: "in water limited" to "in this water limited"

18 L22: "shrubs as the understory layer were low at density due to manual removal" to "understory
19 shrubs were kept at low density by manual removal"

20 *Reply: Thanks for your careful reading and review comments. Corrected (see Revised MS in*
21 *P24 L22; P25 L3);*

22

23 **P5**

24 L1-2: "... from [the] southeast (during [the] growing season and [the] northwest [(outside of the
25 growing season)]."

26 *Reply: Corrected(see in Revised MS P25, L12-13);.*

27 I cannot spend the time to provide additional writing edits but further corrections are required
28 before this can be accepted.

1 *Reply: Thanks for your suggestion for ensuring the publication quality. The native English*
2 *speaking co-authors have edited the writing of manuscript.*

3

4 **Changes in manuscript:**

5 **P21**

6 L1-2: The title has been changed to “*Energy partitioning and surface resistance of a poplar*
7 *plantation in northern China*”

8

9 **P22 (Abstract)**

10 L2-32: the abstract has been revised as “Poplar (*Populus sp.*) plantations have been, on one
11 hand, broadly used in northern China for urban greening, combating desertification, as well as
12 for paper and wood production. On the other hand, such plantations have been questioned
13 occasionally for their possible negative impacts on the water availability due to higher water
14 use nature of poplar trees compared with other tree species in water limited dryland regions. To
15 further understand the acclimation of poplar species to semiarid environment and evaluate the
16 potential impacts of these plantations on the broader context of the region’s water supply, we
17 examine the variability of bulk resistance parameters and energy partitioning in a poplar
18 (*Populus euramericana* CV. “74/76”) plantation located in northern China over a four-year
19 period encompassing both dry and wet conditions. The partitioning of available energy to latent
20 heat flux (LE) decreased from 0.62 to 0.53 under mediated meteorological drought by irrigation
21 applications. A concomitant increase in sensible heat flux (H) resulted in the increase of a
22 Bowen ratio from 0.83 to 1.57. Partial correlation analysis indicated that surface resistance (R_s)
23 normalized by leaf area index (LAI) ($R_s:LAI$) increased by 50% under drought conditions and
24 was the dominant factor controlling the Bowen ratio. Furthermore, R_s was the main factor
25 controlling LE during the growing season, even in wet years, as indicated by the decoupling
26 coefficient ($\Omega = 0.45$ and 0.39 in wet and dry years, respectively). R_s was also a major regulator
27 of the LE/LE_{eq} ratio, which decreased from 0.81 in wet years to 0.68 in dry years. All
28 physiological and bioclimatological metrics indicated that the water demands of the poplar
29 plantation were greater than the amount available through precipitation, highlighting the poor
30 match of a water-intensive species like poplar for this water limited region.”

1 **P23 (Introduction)**

2 L9-10: “i.e., more than” to “over”;

3 L10: “the same species” to “poplar species”;

4 L11: “its” to “their”;

5 L16: “may even” to “could”;

6 L19-20: “Thus, poplar plantation ... such as northern China.” to “Thus, poplar plantation may
7 have higher productivity but also higher water use (Zhou et al., 2013) than other tree species.”;

8 L21-26: “However, over the past 50 years, northern China has ... , while the wide spread
9 use ...for these impacts.” to “The intensive land use practices in northern China over the past
10 50 years, supported by irrigation, are thought to have triggered the decline in its water table,
11 land degradation and increases in surface air temperature and severe droughts (Ding et al., 2007;
12 Qiu et al., 2012; Wang et al., 2008; Zhang et al., 2014).”;

13 L26-29: “studying the drought response ... are not sufficient” to “understanding the
14 contribution of current land cover, including the poplar plantations on the regional water
15 resources is essential for the long-term sustainability of ecosystem services and human
16 wellbeing in this region.”

17 L31-9(P12): “Whereas, most of previous and current studies...(Guo et al., 2010; Jamiyansharav
18 et al., 2011; Sun et al., 2010; Takagi et al., 2009; Wu et al., 2007), ...in water limited regions”
19 to “To date, most researches have concentrated primarily on the water balance of forest
20 ecosystems, with less emphasis on the relationship of forest ecosystems to their environmental
21 setting. Much can be learned from exploring the partitioning of available energy and ecosystem
22 response to meteorological forcing such as droughts. Not only are these of central importance
23 for understanding the water and carbon balance (Guo et al., 2010; Jamiyansharav et al., 2011;
24 Sun et al., 2010; Takagi et al., 2009; Wu et al., 2007), but they also help elucidate the degree to
25 which forest water use is in balance with supply from precipitation, and hence the degree to
26 which plantations located in water limited regions are sustainable in the long-term.”;

27

28 **P24 (Materials and Methods)**

1 L10-16: “The goal of ...in northern China.” To “To investigate the variations of energy
2 partitioning and associated evapotranspiration of poplar plantation under different climate
3 conditions and highlight the management strategies for such plantation forests in water limited
4 region, we evaluated energy partitioning at different water availabilities in a ten-year-old poplar
5 (*Populus euramericana* CV. “74/76”) plantation on sandy soil in northern China.”

6 L17-18: “changes in the surface resistance and energy partitioning in the water demanding
7 poplar species” to “increase in the surface resistance and affect energy partitioning via
8 increasing the Bowen ratio.”

9 L21-23: “(3) evaluate the long term sustainability ... in water limiting region” to “(3) evaluate
10 the long-term potential impact of poplar plantation on the availability of water for adjacent
11 ecosystems and livelihoods in water limited region.”;

12 L31: “were 16.2±1.6 m” to “was 16.2±1.6 m (mean±SD)”;

13

14 **P25**

15 L7: “...11.6°C, and maximum ...” to “11.6°C; maximum...”;

16 L15: “The upper two meter” to “The top two meters”; “well drained” to “well-drained”;

17 L16: “with bulk density” to “with a bulk density”;

18 L17-19: “The groundwater has ..., and has declined...per year.” to “The mean groundwater
19 depth over the past nine years (2001–2009) was 16.5±0.2 m, and declined...per year.”;

20 L22-25: “The amount of flood ... through 2009” to “The site was irrigated using pumped
21 groundwater, and the amount of water supplied was estimated from the water meter records at
22 the three adjacent wells on a weekly basis from 2006 through 2009”;

23 L25: “...included tilling, and weeding...” to “...have included tilling and weeding...”;

24 L29: “in June of 2005” to “in June 2005”;

25 L30-31: “...in size. The observation site has a sufficiently wide fetch of...” to “...in size, with
26 a fetch of ...”; delete “water”;

27

28 **P26**

- 1 L3-7: “The CO₂/H₂O sensor ..., the analyzer was calibrated every year.” to “The anemometer
2 head was installed towards a predominant wind direction (southeast), and the IRGA was
3 installed at a slight vertical angle tilted northward (< 20 degree) between the sonic path and
4 anemometer body. The IRGA was calibrated every year.”;
- 5 L13: “by” to “with”;
- 6 L15: “at 21 m height” to “at height of 21 m”; “...with temperature...” to “...with a temperate...”
- 7 L17: “20 m” to “20 m above ground”;
- 8 L18-20: “Soil heat flux was ...were measured with three soil heat... with three
9 thermocouples...” to “Soil heat flux and soil temperatures, respectively, were measured with
10 three soil heat transducers (HFT3, CSI) and three thermocouples...”;
- 11 L25: “... at 10 Hz, and ...” to “...at 10 Hz and ...”;
- 12 L28-29: “The raw ...Processor,” to “The 30-minute mean fluxes were calculated from raw 10
13 Hz data with an EC Processor software,”;
- 14
- 15 **P27**
- 16 L3: “30-min” to “the 30-min”;
- 17 L9-16: “In this study,...were not filled.” to “Data gaps shorter than 2 hours were filled using
18 linear regressions between the flux of interest and net radiation (R_n), gaps between 2 hours and
19 7 days in length were filled using mean diurnal variation (MDV) method (Falge et al., 2001),
20 and gaps longer than 7 days were not filled.”;
- 21 L17-20: “Four year ... (China, 2006).” to “The four year study period was classified into “wet”
22 and “dry” years distinctively. A dry year referred to a year with annual precipitation less than
23 85% of the 20-year average according to the National Standard of People's Republic of China
24 (GB/T 20481-2006) (China, 2006) and “wet” when above it.”;
- 25 L22: “driving forces” to “environmental forcing”;
- 26 L23: “...water fluxes and...” to “...water fluxes, and ...”;
- 27 L26: “The regulations of surface exchange” to “The regulation of surface energy and gas
28 exchange”;
- 29 L28: “less station than those” to “less stationary than”;

1 L29-31: “The midday was ... was usually the strongest.” To “The midday was defined as the
2 period from 10:00 to 15:00 LST when the coupling between vegetation and the atmosphere was
3 the strongest.”

4

5 **P28**

6 L21: “from 10:00 to 15:00 LST” to “10:00-15:00 LST”;

7

8 **P29**

9 L7: “latent heat” to “the latent heat”;

10 L11: “which” to “that”;

11 L21: “approaching to” to “approaching”;

12

13 **P30 (Results)**

14 L3-8: “...of ecosystem. It,...evaporation (Arain et al., 2003).” To “of an ecosystem, indicating
15 whether soil water supply for evapotranspiration of an ecosystem was under limited. An
16 LE/LE_{eq} of < 1 indicates water stress and suppressed evapotranspiration. Conversely, $LE/LE_{eq} >$
17 1.26 indicates unrestricted water supply, and only available energy limits evapotranspiration
18 (Arain et al., 2003).”

19 L13-14: “biophysical variables” to “the biophysical variables”; “among” to “across”;

20 L21-26: “The annual precipitation ...in 2007 and 2008.” To “The annual precipitation rates in
21 the four years of study differed from the long-term (i.e., 1990–2009) average (556 mm yr^{-1}).
22 Thus, years 2006 and 2009 were drier and 2007 and 2008 were wetter than the mean (Table 1).
23 The interannual contrast was exaggerated by the seasonality of rainfall.”;

24 L30: “of growing season” to “during the growing season”;

25

26 **P31**

27 L6: “(57 mm), and...” to “(57 mm) and...”;

- 1 L7: “smallest” to “the smallest”;
- 2 L9: “of 2009” to “in 2009”; “higher-than” to “higher than”;
- 3 L11-12: add “ $\Delta T = 1.3 \text{ }^\circ\text{C}$ ”;
- 4 L13-14: “in June in 2006” to “in June 2006”, “July in 2007...” to “in July 2007...”, “June for
5 2006” to “June 2006”;
- 6 L19: “with” to “to”;
- 7 L27: “growing season” to “the growing season”;
- 8 L30-31: “with a lower value ... ($p < 0.001$)” to “with a lower value in wet years (2.1% in 2007)
9 than in the dry years (4.9% in 2006; $p < 0.001$).”; delete “Additionally...4.9% in 2006.”;
- 10
- 11 **P32**
- 12 L1: “four growing seasons” to “the four growing seasons”;
- 13 L2: “MI” to “MJ”;
- 14 L3: “between 6.0% in 2007 and 6.8% in 2009 and showed...” to “from 6.0% in 2007 to 6.8%
15 in 2009, showing...”;
- 16 L5-27: “LE was the dominant ... under drought stress.” to “Partitioning of R_n into LE and H
17 differed significantly between the wet and dry years (Table 3; $F = 17.599$, $p < 0.001$). The
18 dominant turbulent energy flux during the early growing season was sensible heat flux (H) with
19 or without drought stress except in 2006 when the irrigation was applied (Table 3). Then LE
20 was the dominant driver of energy partitioning during the middle and late growing seasons
21 under drought stress. The average daytime total LE was about 20% greater in wet years (6.77
22 MJ m^{-2}) than in dry years (5.72 MJ m^{-2} , $p < 0.01$). The timing of peak LE was weakly related
23 to drought, peaking in July in 2006, 2008 and 2009, and in August in 2007. The peak value of
24 daytime total LE was 16.61 MJ m^{-2} , 17.01 MJ m^{-2} , 19.72 MJ m^{-2} and 16.27 MJ m^{-2} , in 2006–
25 2009 respectively. The daily evaporative fraction ($\text{LE}/(R_n - G)$) was significantly higher in wet
26 years (60.3% and 64.8% in 2007 and 2008, respectively) (64.8%) than in dry years (57.1% and
27 50.4% in 2006 and 2009, respectively; $p < 0.05$).”;
- 28 L28: “rapid” to “a rapid”;

- 1 L29-30: “(April-June) and end (September-October)” to “(April-June) and the end
2 (September-October)”;
- 3 L31-32: “growing seasons” to “the growing seasons”, “180-250” to “180-250”, “180-290” to
4 “180-290”, “of dry year” to “in the dry years”;
- 5
- 6 **P33**
- 7 L1: “of the wet year” to “in the wet years”;
- 8 L3: “(Table 3), and had much...” to “(Table 3); with much...”;
- 9 L8-10: “190-250” to “190-250”; add “A significantly negative relationship was found between
10 the R_s and LAI during the wet years (Fig.6).”;
- 11 L10: “surface resistance (R_s)” to “ R_s ”;
- 12 L11-12 “in 2008 (54.1 s m⁻¹ leaf area) was lowest among four years (i.e., $p < 0.05$)” to “was
13 lowest during the wettest year (2008, 54.1 s m⁻¹ leaf area; $p < 0.05$)”
- 14 L12-13: “year” to “years”;
- 15 L14: “of 2006” to “in 2006”; “greatly” to “much”;
- 16 L15-16: “in unstressed periods ($p < 0.001$)” to “during unstressed periods ($p < 0.001$, Table 3)”;
17 delete the sentence “In addition,... (Fig.6).”
- 18 L17: “in June, and” to “in June and”;
- 19 L18: “growing season” to “the growing season”;
- 20 L22: add “and”;
- 21 L23: “that of the dry years” to “in the dry years”;
- 22 L24: “that in dry years” to “in the dry years”;
- 23 L27: “0.89, and...” to “0.89 and...”;
- 24
- 25 **P34 (Discussion)**
- 26 L4: add “0.40.”;
- 27 L6: “lower values” to “lower in value”;

- 1 L10-12: “The energy balance ratio...the closure of the energy budget” to “The energy balance
2 ratio (E_{BR}) at the current study”; “fluxes, and...” to “fluxes and”;
- 3 L15: “site-year” to “site-years”;
- 4 L16: “0.34-1.69” to “0.34–1.69”;
- 5 L18: “...the biomes, and based on...” to “...the biomes and based on...”;
- 6 L25: “energy partitioning” to “energy partitioning to sensible and latent heat”;
- 7 L27: “To the extent that canopy” to “Canopy”;
- 8 L28: “...properties, they could...” to “...properties to some extent and could...”;
- 9 L29: “therefore impact” to “thereby impacting”;
- 10
- 11 **P35**
- 12 L1: “a detectable response of $LE/(R_n-G)$ and Bowen ratio” to “detectable responses of $LE/(R_n-$
13 $G)$ and the Bowen ratio”;
- 14 L6: “...water supply, similar...” to “...water supply; a similar...”;
- 15 L8: “the most part” to “most”;
- 16 L9-10: “non-stressed periods in other 3 years, which...” to “in the non-stressed periods of the
17 other 3 years. This variation...”;
- 18 L15: “such as, 0.74...” to “such as 0.74...”;
- 19 L16: “..., 0.89” to “...and 0.89”;
- 20 L19: “a higher” to “higher”;
- 21 L21-22: “low water holding capacity of the sandy soil, and high...” to “the sandy soil’s low
22 water holding capacity and the high...”;
- 23 L25: “energy partitioning” to “the energy partitioning”;
- 24 L28: “of” to “on”;
- 25
- 26 **P36**
- 27 L1: delete “,”

- 1 L2-5: delete the sentences “Wilson et al. (2002b)...parameters (Cho et al., 2012).”
- 2 L6: “similar to R_s ” to “ R_s similarly”;
- 3 L10-13: “The drought...(e.g., Noormets et al., 2008)” to “It has been shown that drought stress
4 during the canopy development affects leaf area and may have lasting effects on canopy gas
5 exchange through the entire growing season, even after the moisture limitation is removed
6 (Noormets et al., 2008)”;
- 7 L14: “researches” to “studies”;
- 8 L15: “of this poplar plantation” to “in the current study”; “Euphrates” to “the Euphrates”;
- 9 L16: “leaf area” to “LAI⁻¹”; “Gansu Poplar” to “the Gansu Poplar”;
- 10 L17-18: “leaf area” to “LAI⁻¹”; “northwest China” to “semiarid regions”;
- 11 L19: “leaf area” to “LAI⁻¹”; delete “in Iceland”;
- 12 L20: “leaf area” to “LAI⁻¹”; “in Canada (Blanken et al., 1997)” to “(Blanken et al., 1997) in
13 mesic temperate regions”;
- 14 L22: “..., and modulated...” to “... and modulated...”;
- 15 L23-25: “The strong correlation...(Fig.6)...” to “Compared to the strong correlation between
16 R_s and LAI in wet years, the increased scatter in the R_s -LAI relationship during dry years
17 (Fig.6)”;
- 18 L27: “this study area” to “the current study”; “mean” to “the mean”;
- 19 L28: “across site-year for forests” to “reported for temperate forests”;
- 20 L29: add “was”;
- 21 L30-31: “likely due to ... in China.” To “as might be expected given the predominant climatic
22 conditions”;
- 23
- 24 **P37**
- 25 L1: “correlated, and ...” to “correlated and ...”;
- 26 L2-3: “dry years” to “dry years (Fig. 10)”;

1 L3-9: “The Bowen ratio and R_s ... with the growing R_s ” to “The water limitation during the dry
2 years manifested in disproportional increase in R_s than the Bowen ratio; this response may
3 serve as an indicator when water reserves are being depleted. At the extremes, the relationship
4 converges, but as water becomes limiting, stomatal closure and increased R_s do not appear to
5 be able to affect the seasonal dynamics of the Bowen ratio.”;

6 L10: “respectively, had ... effects,” to “had ... effects, respectively,”;

7 L12-14: “both of R_s and R_i ...in wet years.” to “the regulation of the Bowen ratio by R_s and R_i
8 seemed stronger in dry than in wet years.”; delete “Finally,”;

9 L19-22: “Similar to ...the growing season” to “As essentially implied by the *Penman-Monteith*
10 equation, LE/LE_{eq} exponentially related to R_s during the growing season”;

11 L25: “1.1-1.4 range” to “1.1–1.4 range typical in temperate deciduous forest”;

12 L26-27: “characterized by ... forest biome...” to “drier than these reference sites...”;

13 L29: “sandy soil and a low ground water...” to “the sandy soil and the low ground water...”;

14 L31-32: “growing season” to “the growing season”;

15

16 **P38**

17 L1: “growing season” to “the growing season”; “which were” to “as”;

18 L3: “Implication ... establishment” to “Implications ... establishments”;

19 L4-11(P27): revise the context “To our knowledge,... is not sustainable.” to “As forestry is a
20 long-term endeavor, with the economic payback decades from stand establishment, the
21 availability of resources for the stand to prosper should come naturally to natural resource
22 managers. Supplementing limiting resources directly (fertilization, irrigation) or indirectly
23 (competition control, site preparation, thinning) is commonplace in commercial forestry, but it
24 has to be sustainable in the broader context of the region's ecosystems and livelihoods. Earlier,
25 we reported that the water needs of poplar plantation exceed the annual precipitation in the
26 region and plant survival during dry years depends on irrigation from groundwater (Zhang et
27 al., 2014). In the current study, energy partitioning to latent and sensible heat and surface
28 resistance was sensitive to climatological drought—even under the irrigation—as indicated by
29 low LE/LE_{eq} (< 1) and low values of the decoupling coefficient (Ω) (Zhu et al., 2014); the dry

1 surface conditions dominated the poplar plantation in both wet and dry years. In wet years, the
2 plantation itself is in hydrologic balance with the water that arrives as precipitation, with
3 evapotranspiration consuming nearly all of the precipitation. The same is true in dry years, but
4 irrigation increases *ET* even further by depleting groundwater. Even if the plantations were in
5 hydrologic balance with water delivered as precipitation, their existence and operation could be
6 a threat to adjacent ecosystems and livelihoods if those rely on runoff or groundwater recharge
7 from the areas where the plantation has been sited. In the absence of the plantations it is likely
8 that groundwater recharge would increase, especially given the sandy textured soil that tends to
9 allow rapid infiltration and percolation as well as limits moisture delivery to the atmosphere
10 directly from the soil surface itself. While poplar plantation growth in this water-limited
11 location might be sustained by the modest precipitation in the region, it could still be
12 unsustainable for the broader context of the region's ecosystems and livelihoods. However,
13 further study to truly assess these effects is needed by comparing the surface water balance and
14 /or spatial and temporal variations of groundwater levels at an adjacent, similar site without a
15 plantation.”

16

17 **P39 (Conclusions)**

18 L15: “growing seasons” to “the growing seasons”;

19 L17: “correspondingly displayed” to “resulted in”;

20 L19: “in dry years was 33% higher than that in wet years” to “was 33% higher in dry than in
21 wet years”;

22 L20: “Accordingly” to “Correspondingly”; “impact” to “effects”;

23 L21: “,” to “.”;

24 L23: delete “overall”;

25 L24-30: “the dry climate ... water limited regions.” to “the permanent limitation of plant water
26 use and surface energy partitioning by water availability. Even at mean long-term precipitation,
27 the water demand of poplar plantation may consume nearly all of it and leave little for run-off
28 and groundwater recharge in this semi-arid region, potentially compromising the region’s
29 ecosystems and livelihoods.”

30

1 **P40**

2 L3: add “(Grant No. 201204102)”;

3 L4: “scholarship” to “financial”;

4 L6-7: add “Dr. Christopher A. Williams (Associate Editor) and”;

5 L18-20: delete reference “Baldocchi, D...1994.”;

6

7 **P41**

8 L1-2: delete reference “Chi, J., ..., 2012.”;

9

10 **P43**

11 L20-21: delete reference “Richardson, B., ...1999.”;

12

13 **P44**

14 L1-3: delete reference “Watt, M. S., ..., 2005.”

15

16 **P47**

17 L10: “statistic” to “statistics”;

18

19 **P51**

20 L13: “brace” to “brackets”;

21

22 **P54**


23 L12: “midday” to “midday (10:00-15:00 LST)”;

24 L14-15: delete “; Midday means ... local standard time.”;

25

26 **P55**

- 1 L2: “midday” to “midday (10:00-15:00 LST)”;
- 2 L4-5: delete “; Midday means ... 15:00 p.m. LST.”;
- 3
- 4 **P58**
- 5 L9: “four growing seasons” to “the four growing seasons”;
- 6
- 7 **P60**
- 8 L4: change Figure 10 as followed:



- 9
- 10
- 11
- 12
- 13
- 14
- 15
- 16
- 17
- 18

1 ~~Responses of e~~**Energy partitioning and surface resistance**
2 ~~to drought in~~**of a poplar plantation in northern China**

3

4 **M. Kang¹, Z. Zhang¹, A. Noormets², X. Fang¹, T. Zha¹, J. Zhou³, G. Sun⁴, S. G.**
5 **McNulty⁴, and J. Chen⁵**

6 [1]{Key Laboratory of Soil and Water Conservation and Desertification Combating, Ministry
7 of Education, College of Soil and Water Conservation, Beijing Forestry University, Beijing
8 100083, PR China}

9 [2]{Department of Forestry and Environmental Resources, North Carolina State University,
10 Raleigh, NC, USA}

11 [3]{Beijing Municipal Station of Agro-environmental Monitoring, Beijing, 100029, PR China}

12 [4]{Eastern Forest Environmental Threat Assessment Center, Southern Research Station,
13 USDA Forest Service, Raleigh, NC, USA}

14 [5]{Landscape Ecology & Ecosystem Science (LEES) Lab, Center for Global Change and
15 Earth Observations (CGCEO), and Department of Geography, Michigan State University, East
16 Lansing, MI 48823, USA}

17 Corresponding to: Zhiqiang Zhang (zhqzhang@bjfu.edu.cn)

18

19

20

21

22

23

24

25

26

27

1 Abstract

2 Poplar (*Populus sp.*) plantations have been, on one hand, broadly used broadly in northern China
3 for urban greening, combating desertification, urban greening, and as well as for paper and
4 wood production in northern China. On the other hand, such plantations have been questioned
5 occasionally for their possible negative impacts on the water availability due to higher water
6 use nature of poplar trees compared with other tree species in water limited dryland regions. To
7 further understand the acclimation of poplar species to semiarid environment and evaluate
8 ~~However, given the high water use by the species and the regional dry climate, the sustainability~~
9 potential impacts of these plantations on the broader context of the region's water supply, needs
10 to be evaluated. Currently, the understanding of the acclimation of the species to the semiarid
11 environment is limited, impeding assessments of their long term success and impact on the
12 environment. In this study we examine the variability of bulk resistance parameters and energy
13 partitioning in a poplar (*Populus euramericana* CV. "74/76") plantation located in northern
14 China over a four-year period encompassing both dry and wet conditions in a poplar (*Populus*
15 *euramericana* CV. "74/76") plantation located in northern China. The partitioning of available
16 energy to latent heat flux (LE) decreased from 0.62 to 0.53 under mediated meteorological
17 drought by irrigation applications. A concomitant increase in sensible heat flux (H) resulted in
18 the increase of a Bowen ratio from 0.83 to 1.57. Partial correlation analysis indicated that
19 surface resistance (R_s) normalized by leaf area index (LAI) (i.e., R_s :LAI) increased by 50%
20 under drought conditions and became was the dominant factor controlling the Bowen ratio.
21 Furthermore, R_s was the major-main factor controlling LE during the growing season, even in
22 wet years, as indicated by the decoupling coefficient ($\Omega = 0.45$ and 0.39 in wet and dry years,
23 respectively). R_s was also a major regulator of and the LE/LE_{eq} ratio, which decreased ranging
24 from 0.81 in wet years to and 0.68 in wet and dry years, respectively. In general, the dry climate
25 dominated the poplar plantation ecosystem regardless of soil water availability suggesting that
26 fast growing and water use intensive species like poplar plantations are poorly suited for the
27 water limited region. All physiological and bioclimatological metrics indicated that the water
28 demands of the poplar plantation were greater than the amount available through precipitation,
29 highlighting the poor match of a water-intensive species like poplar for this water limited region.
30 ~~The required irrigation for sustaining these forests also presents a thread to the adjacent~~
31 ~~ecosystems because of their role in reducing ground water table, and may compromise long-~~
32 ~~term sustainability and livelihoods in the region.~~

1 1 Introduction

2 Poplar (*Populus sp.*) plantations are the most dominant broadleaf forest ecosystems throughout
3 northern and central China, due to their rapid growth rates, high productivity and wide
4 adaptability ([Gielen and Ceulemans, 2001](#); [Wilske et al., 2009](#); [Zhang et al., 2011](#)). Since the
5 late-1970s, with the implementation of the “Three-North Shelterbelt Program” (1978), the
6 “Combating Desertification Project” (1991) and the “Grain for Grain Program” (1999) ([Wilske](#)
7 [et al., 2009](#)), poplar plantations have been playing a vital role in timber production, bioenergy,
8 urban greening, desertification control, and carbon sequestration ([Martín-García et al., 2011](#);
9 [Zhou et al., 2013](#)). By 2007, China had the largest poplar plantation area in the world (~~i.e., more~~
10 ~~than~~over 7.0 million ha, [Fang, 2008](#)). However, indiscriminate use ~~pf of the same~~poplar species
11 beyond ~~its~~their native range and habitats may result in unanticipated consequences. For
12 example, the use of poplars in water limited regions may increase the risk of environmental
13 degradation, soil moisture deficit, hydrologic and vegetation changes ([Gao et al., 2014](#)).

14 Poplars require large quantities of water throughout the growing season, and may
15 experience water limitation even on the mesic sites ([Kim et al., 2008](#); [Stanturf and Oosten,](#)
16 [2014](#)). For example, poplar plantations ~~may even~~could cause the transformation of wetlands
17 into dry land due to the water-pumping effect on groundwater ([Li et al., 2014](#); [Migliavacca et](#)
18 [al., 2009](#)). Thus, poplar plantations, which have higher productivity but also higher water use
19 ([Zhou et al., 2013](#)) than other ~~forests, clearly require large quantities of irrigation in water~~
20 ~~limited areas such as northern China~~tree species.

21 ~~However, The intensive land use practices in northern China~~ over the past 50 years,
22 ~~supported by irrigation, are~~northern China has experienced ~~thought to have triggered the decline~~
23 ~~in its water table, –land degradation the decline of the water tableand, –land degradation, large~~
24 increases in surface air temperature and severe droughts ([Ding et al., 2007](#); [Qiu et al., 2012](#);
25 [Wang et al., 2008](#); [Zhang et al., 2014](#)), ~~while the ,wide spread use of irrigation has been cited~~
26 ~~as one of possible causes for these impacts~~. Therefore, understanding the contribution of current
27 land cover, including the poplar plantations on the regional water resources ~~studying the~~
28 ~~drought response of poplars under water shortage~~ is essential for long-term sustainability of
29 ecosystem services and human wellbeing in this region ~~effective management of water resource~~
30 ~~over this region and avoiding the use of water intensive species in ecological restoration and~~
31 ~~reforestation efforts if the environmental resources are not sufficient~~. ~~Whereas~~To date, most
32 researches have concentrated primarily on the water balance of forest ecosystems, with less

1 emphasis on the relationship of forest ecosystems to their environmental setting. Much can be
2 learned from exploring the partitioning of available energy and ecosystem response to
3 meteorological forcing such as droughts. Not only are these of central importance for
4 understanding the water and carbon balance (Guo et al., 2010; Jamiyansharav et al., 2011; Sun
5 et al., 2010; Takagi et al., 2009; Wu et al., 2007), but they also help elucidate the degree to
6 which forest water use is in balance with supply from precipitation, and hence the degree to
7 which plantations located in water limited regions are sustainable in the long-term and thus
8 understanding the adaption and long term sustainability of plantation establish in water limited
9 regions.

10 To investigate the variations of energy partitioning and associated evapotranspiration of
11 poplar plantation The goal of the current study was to examine how forest water and energy
12 balances vary under different climatic climate conditions and how to best manage the highlight
13 the management strategies for such plantation forests to maximize ecological benefits in water
14 limited region. ~~Therefore,~~ we evaluated ~~drought responses in~~ energy partitioning at different
15 water availabilities in a ten-year-old poplar (*Populus euramericana* CV. “74/76”) plantation on
16 sandy soil in northern China. We hypothesized that drought would trigger significant
17 ~~increasechanges~~ in the surface resistance and affect energy partitioning via increasing the
18 Bowen ratio in the water demanding poplar species. Specifically, the objectives of this study
19 were to: (1) quantify the seasonal and inter-annual variability of energy partitioning and bulk
20 resistance parameters; (2) partition the control of energy partitioning to biological and
21 climatological components; and (3) evaluate the long-term ~~sustainability~~ potential impact of
22 poplar plantations on the availability of water for adjacent ecosystems and livelihoods in a ~~water~~
23 ~~-limiting limited~~ region ~~in northern China~~.

24 **2 Materials and Methods**

25 **2.1. Study site**

26 The study was carried out in a managed poplar (*Populus euramericana* CV. “74/76”)
27 plantation at the Daxing Forest Farm, which is located in the southern suburbs of Beijing, China
28 (116°15'07"E, 39°31'50"N, 30 m a.s.l.). The trees were planted in 1998 with 3 m × 2 m spacing,
29 dead or low-vigor trees were replaced with new saplings in 2001 and 2003. The stand
30 characteristics over the four years of study are provided in Table 1. At the end of 2009, the
31 average height of the trees ~~were was~~ 16.2±1.6 m (mean±SD), and the diameter at breast height

1 (DBH) was 14.1 ± 1.6 cm. The average leaf area index (LAI) of the stand increased over time.
2 During the growing season, ~~shrubs as the understory layer were low at density due to understory~~
3 ~~shrubs were kept at low density by~~ manual removal. Perennial herbs included *Chenopodium*
4 *glaucum* Linn., *Medicago sativa* L., *Melilotus officinalis* (L.) Lam., *Salsola collina* Pall., and
5 *Tribulus terrestris* L.

6 The local climate is classified as sub-humid warm temperate zone, with a mean (1990–
7 2009) annual temperature of 11.6°C , ~~and~~ maximum and minimum temperature are 40.6°C and
8 -27.4°C , respectively. The annual precipitation ranges from 262 mm to 1058 mm (1952–2000),
9 with an average of 556 mm, of which 60%–70% falls from July to September (Daxing Weather
10 Station, $116^\circ 19' 56''$ E, $39^\circ 43' 24''$ N). The annual frost-free period lasts 209 days, and the
11 total sunshine-hour reaches 2772 h per year with $15.5 \text{ MJ m}^{-2} \text{ d}^{-1}$ of incoming solar radiation.
12 The average wind speed is 2.6 m s^{-1} and it mostly comes from ~~the~~ southeast (during ~~the~~ growing
13 season) and ~~the~~ northwest (~~outside of the during non-growing~~ season).

14 The study area is on the alluvial plain of the Yongding River, and is flat with an average
15 slope of $< 5^\circ$. The ~~upper-top~~ two meters of the soil is mostly composed of ~~well-well~~-drained
16 fluvial sand with a bulk density of 1.43 – 1.47 g cm^{-3} , and a pH of 8.25 – 8.39 . The soil porosity
17 is about 40% and capillary porosity is 32%. The ~~mean~~ groundwater ~~depth over the past nine~~
18 ~~years (2001–2009) table h~~ was ~~an annual average of~~ 16.5 ± 0.2 m ~~below the ground in the past~~
19 ~~nine years (2001 to 2009)~~, and ~~has~~ declined at an average rate of 0.6 m per year. The maximum
20 pan evaporation occurs from May through June, exceeding precipitation for the same period.
21 Severe drought during the beginning of the growing season (from April to June) in northern of
22 China is common. The ~~amount of flood irrigationsite~~ was ~~irrigated applied by using pumping~~
23 ~~pumped~~ groundwater, and ~~the amount of water supplied was estimated from~~ ~~back-calculated the~~
24 ~~based on the water meter~~ records ~~of the water meters from~~ the three ~~adjacent~~ wells on a weekly
25 basis from 2006 through 2009. Other management practices ~~have~~ included tilling— ~~and~~
26 weeding since the establishment of the plantations.

27 **2.2. Eddy covariance system**

28 The micrometeorological and eddy flux measurements were conducted at a 32m tower in the
29 center of the study site, which was established in June ~~of~~ 2005. The foot-print of the eddy flux
30 covariance system, was about 1 km x 1 km in size, ~~with~~. ~~The observation site has a~~ ~~sufficiently~~
31 ~~wide~~-fetch of at least 300 m in all directions. Fluxes of CO_2 , ~~water~~-sensible heat and latent heat

1 were calculated based on the eddy-covariance (EC) principles. The sensors included a CO₂/H₂O
2 infrared analyzer (Li-7500; LI-COR, Inc., Lincoln, NE, USA) and a three-dimensional sonic
3 anemometer (CSAT-3; Campbell Scientific, Inc., CSI, UT, USA). The ~~CO₂/H₂O~~
4 ~~sensor~~~~anemometer~~ head was installed towards a predominant wind direction (southeast), and
5 the IRGA was installed at ~~with~~ a slightly vertical angle tilted northward (< 20 degree) ~~and~~
6 ~~downwind of the sonic anemometer in the predominant wind direction;~~between the sonic path
7 and anemometer body. ~~the~~~~The IRGA~~~~analyzer~~ was calibrated every year. The EC sensors were
8 mounted initially at a height of 16 m in 2006. This was increased to about 18 m before the start
9 of the growing season in 2007, and again to 20 m in February 2009 to ensure that the sensors
10 remained well above the tree canopy.

11 Net radiation was measured with net radiometers (Q7.1, REBS, Seattle, WA, USA) and
12 (CNR-1; Kipp and Zonen, Delft, Netherlands) at 26 m above the ground. Photosynthetically
13 active radiation (PAR) was measured ~~by~~with a PAR quantum sensor (LI-190SB; LI-COR, Inc.)
14 mounted at 20 m. The atmospheric pressure was measured by a barometric pressure sensor
15 (CS105, CSI) at height of 21 m~~height~~. Air temperatures and humidity were measured with a
16 temperature and relative humidity probe (HMP45C; Vaisala, Helsinki, Finland) at 5, 10, 15 and
17 20 m above ground. Precipitation was measured with a tipping bucket rain gauge (TE525-L;
18 Texas Electronics, USA) at 22.5m. Soil heat flux and soil temperatures, respectively, were
19 measured with ~~was determined with~~ three soil heat transducers (HFT3, CSI) ~~and,~~ ~~and soil~~
20 ~~temperatures were measured with~~ three thermocouples (TCAV107; CSI) located at depths of 5,
21 10 and 20 cm below the soil surface. Soil water content was measured with TDR sensors
22 (CS616; CSI) buried at 20 and 50cm.

23 With the exception of the rain gauge, all microclimatic data were recorded with a data-
24 logger (CR23X; CSI) at 30 min intervals and the fluctuations in wind speed, sonic temperature
25 and CO₂ and H₂O concentrations were sampled at 10 Hz, and recorded by a CR5000 data-
26 logger (CSI).

27 **2.3. Data processing and QA/QC**

28 The 30-minute mean fluxes were calculated from raw 10 Hz data ~~were processed~~ with an EC
29 Processor software, version 2.3 ([Noormets et al., 2010](#)). The program is designed for
30 reprocessing EC flux data and can calculate half-hour mean eddy covariance fluxes of carbon,
31 water and energy. The wind coordinates were rotated using the planar fit method ([Paw U et al.](#),

1 [2000](#); [Wilczak et al., 2001](#)). Fluxes were corrected for additional sensor heating ([Burba et al.,](#)
2 [2008](#)) and fluctuations in air density ([Webb et al., 1980](#)). The data quality controls included:
3 screening of the 30-min mean eddy covariance fluxes based on instrument quality flag, integral
4 turbulence characteristics ([Foken and Wichura, 1996](#)), flux stationarity, atmospheric stability,
5 and adequate turbulent mixing([Goulden et al., 1996](#)). The threshold of friction velocity (μ_*)
6 below which flux loss occurred was determined from the seasonal binned relationship between
7 nighttime turbulent flux of CO₂ and friction velocity (μ_*) ([Schmid et al., 2003](#)). The threshold
8 was consistent across different seasons, but differed slightly between years: 0.18 m s⁻¹ (2006),
9 0.12 m s⁻¹ (2007), 0.14 m s⁻¹ (2008) and 0.13 m s⁻¹ (2009). Data gaps shorter than 2 hours were
10 filled using linear regressions between the flux of interest and net radiation (R_n), gaps between
11 2 hours and 7 days in length were filled using mean diurnal variation (MDV) method~~In this~~
12 ~~study, the MDV (mean diurnal variation) method ([Falge et al., 2001](#)), was used to fill the data~~
13 ~~gaps, the linear relationship between LE or H and net radiation (R_n) was used to gap fill each~~
14 ~~flux when short period (< 2h) flux data were missing. A ± 7 day moving average was used to fill~~
15 ~~each flux gaps for period between 2 h and 7 days. and gaps~~ Gaps longer than 7 days were not
16 filled.

17 The ~~four~~ year study period was classified into “wet” and “dry” years distinctively. ~~The~~
18 A dry year referred to a year ~~the meteorological drought when yearly~~ with annual precipitation
19 less than ~~75~~85% of the 20-year average according to the National Standard of People's Republic
20 of China (GB/T 20481-2006) ([China, 2006](#)) and “wet” when above it. Years 2007 and 2008
21 were classified as ‘wet’ while 2006 and 2009 were ‘dry’ year, respectively. We focused on the
22 growing season when the ~~driving force~~ environmental forcing (e.g., solar radiation, and
23 temperature) for energy and water fluxes, and the physiological response of vegetation were
24 usually strong. In this study, the strongest forcing days occurred approximately between day
25 100 (mid-April) and day 300 (late October). The daytime was defined as the period between
26 the sunrise and sunset with PAR > 4 $\mu\text{mol m}^{-2}\text{s}^{-1}$. The regulations of surface energy and gas
27 exchange are often different during nocturnal periods ([Mahrt, 1999](#)), with heat fluxes at night
28 typically weaker and markedly less ~~station-stationary~~ than those during the daytime ([Wilson et](#)
29 [al., 2002b](#)). The midday was defined as the period from 10:00 ~~a.m.~~ to 15:00 ~~p.m.~~ at local
30 standard time LST, ~~—~~ when the interaction-coupling between vegetation and the environment
31 atmosphere was ~~usually~~ the strongest.

2.4. Biophysical characteristics

The availability of relative extractable water (REW) content was calculated to analyze the ecosystem response on drought stress. According to [Granier et al. \(2007\)](#), soil water stress was assumed to occur when the REW dropped below the threshold of 0.4. Daily REW is calculated as,

$$REW = \frac{VWC - VWC_{\min}}{VWC_{\max} - VWC_{\min}} \quad (1)$$

where VWC_{\min} and VWC_{\max} are the minimum and maximum soil volumetric water content across the four years, respectively.

The Bowen ratio (β) reflects the influence of microclimate and the hydrological cycle on the energy partitioning and water use of the ecosystem ([Perez et al., 2008](#)). The midday β is calculated as Eq. (2),

$$\beta = \frac{H}{LE} \quad (2)$$

Based on the daytime half-hourly and daytime totals of turbulent energy fluxes, the energy balance ratio (E_{BR}) is calculated as Eq. (3),

$$E_{BR} = \frac{\sum(H + LE)}{\sum(R_n - G - S)} \quad (3)$$

where S is the latent and sensible heat storage in the air-column below the EC system and is calculated as in Eq. (4) ([Dou et al., 2006](#)),

$$S = \int_0^{hc} \rho c_p \frac{\partial T}{\partial t} dz + \int_0^{hc} \frac{\rho c_p}{\gamma} \frac{\partial e}{\partial t} dz \quad (4)$$

where hc is the height of eddy flux system measurement (32 m), T is air temperature in the air-column below hc , and e is water vapor pressure.

During midday periods (~~from 10:00 to~~ 15:00 LST), the *Penman-Monteith* approximation was inverted to calculate the surface resistance (R_s) in Eq. (5) ([Kumagai et al., 2004](#)),

$$R_s = \frac{\rho c_p (\delta_e / LE)}{\gamma} + \left(\frac{\Delta}{\gamma} \beta - 1 \right) R_a \quad (5)$$

where R_s is the surface resistance to water vapor transport ($s \text{ m}^{-1}$), representing four components: bulk stomatal resistance of the canopy, bulk boundary layer resistance of the vegetation, bulk

1 ground resistance, and bulk boundary layer resistance of the ground ([Admiral et al., 2006](#); [Cho](#)
2 [et al., 2012](#); [Perez et al., 2008](#); [Wilson et al., 2002b](#)).

3 R_i is the climatological resistance (s m^{-1}) indicating the atmospheric demand ([Wilson et](#)
4 [al., 2002b](#)) and is calculated as,

$$5 \quad R_i = \frac{\rho c_p \delta_e}{\gamma A} \quad (6)$$

6 where A is the available energy ($R_n - G$); ρ is air density (kg m^{-3}), c_p is the specific heat of the
7 air ($1005 \text{ J kg}^{-1} \text{ K}^{-1}$); δ_e is the atmospheric vapor pressure deficit (Pa); LE is the latent heat flux;
8 Δ is the change of saturation vapor pressure with temperature (Pa K^{-1}); γ is the psychrometric
9 constant ($\approx 67 \text{ Pa K}^{-1}$); β is the Bowen ratio.

10 R_a is the aerodynamic resistance of the air layer between the canopy and the flux
11 measurement height (s m^{-1}), ~~which—that~~ reflects the aerodynamic properties of turbulent
12 transport in the near surface boundary layer ([Holwerda et al., 2012](#); [Zhang et al., 2007](#)). R_a is
13 calculated following [Hossen et al. \(2011\)](#) and [Migliavacca et al. \(2009\)](#),

$$14 \quad R_a = r_{a,m} + r_b = \frac{\mu}{\mu_*^2} + 6.2\mu_*^{-2/3} \quad (7)$$

15 where $r_{a,m}$ is the aerodynamic resistance for momentum transfer, and r_b is the quasi-laminar
16 boundary-layer resistance, μ is the wind speed, and μ_* is the friction velocity.

17 The decoupling coefficient (Ω) explains the degree of coupling between the atmosphere
18 and the vegetation, and describes the relative control of evapotranspiration by surface resistance
19 and net radiation ([Pereira, 2004](#)). The Ω value ranges from 0 to 1, with values approaching zero
20 indicating that LE is highly sensitive to surface resistance and ambient humidity deficit. The Ω
21 value approaching ~~to~~ 1 indicates that LE or evapotranspiration is mostly controlled by net
22 radiation ([Jarvis and McNaughton, 1986](#)),

$$23 \quad \Omega = \frac{\Delta + \gamma}{\Delta + \gamma(1 + \frac{R_s}{R_a})} \quad (8)$$

24 The equilibrium evaporation (LE_{eq}) is the climatologically determined evaporation
25 (atmospheric demand) over an extensive wet surface and is dependent only on R_n and
26 temperature. It is calculated as,

$$27 \quad LE_{eq} = \frac{\Delta(R_n - G)}{\Delta + \gamma} \quad (9)$$

1 The ratio LE/LE_{eq} , which is also known as the Priestley–Taylor α , reflects the control of
2 evaporation by atmospheric and physiological factors, LE/LE_{eq} characterizes the surface
3 dryness of an ecosystem. ~~It, therefore, indicates indicating~~ whether soil water supply for
4 evapotranspiration of an ecosystem ~~is was~~ under ~~limitation limited or not~~. An LE/LE_{eq} of < 1
5 ~~represents indicates an ecosystem under~~ water stress ~~, and, therefore, experiences reductions~~
6 ~~in suppressed~~ evapotranspiration; ~~whereas~~ ~~Conversely,~~ LE/LE_{eq} ~~of~~ > 1.26 indicates ~~an~~
7 ~~ecosystems of~~ unrestricted water supply, and only available energy limits ~~evaporation~~
8 ~~evapotranspiration~~ (Araim et al., 2003). The LE/LE_{eq} is dependent of leaf area index (LAI), soil
9 water content, meteorological conditions (e.g., wind speed, solar radiation, VPD, air
10 stratification stability, convection, advection surface resistance), vegetation types, and altitude
11 ([Guo et al., 2008](#)).

12 2.5. Statistical analysis

13 Repeated measurement ANOVA (SPSS) was used for quantifying the changes of all the
14 biophysical variables, energy fluxes ~~,~~ and bulk parameters ~~among across~~ years. The t test was
15 used to compare the differences of biophysical variables among different studies. The partial
16 correlation analysis was used to distinguish the impacts of each of the three resistance
17 parameters (R_s , R_i and R_a) on the Bowen ratios. All analyses were accessed at $\alpha = 0.05$.

18

19 3 Results

20 3.1 Environmental conditions

21 The annual precipitation rates in the four ~~study~~ years of study differed from the long-term (i.e.,
22 1990–2009) average (556 mm yr^{-1}) ~~(1990–2009)~~. Thus, years 2006 and 2009 were drier and
23 2007 and 2008 were wetter than the mean (Table 1). The interannual contrast was exaggerated
24 by the seasonality of rainfall. Precipitation was 74 mm below this long term mean in 2006 and
25 159 mm in 2009. Whereas rainfall exceeded the 20 year mean by over 100 mm in 2007 and
26 2008. Generally, over 90% precipitation of each year occurred in April–October, but with
27 different timing and magnitude among the years. The study site was irrigated during the dry
28 years of 2006 and 2009 to mitigate drought conditions (Fig.1). Seasonal drought stress ($REW <$
29 0.4) occurred during periods in the late growing season of 2006 and 2009, the spring of 2007
30 and 2009, but not at all in 2008 (Fig. 2a–d). In 2006, precipitation ~~of during the~~ growing season

1 reached 467 mm, of which 51% had occurred by July. The amount of irrigation was 35 mm in
2 April, 21 mm in May and 30 mm in September. The two seasonal drought periods separately
3 were #1_06 (from DOY 164 to 192) and #2_06 (from DOY 231 to 300). The total rainfall in
4 2007 and 2008 was similar, but more evenly distributed throughout the year in 2008. In 2007,
5 drought stress occurred during DOY 110-143 (#1_07) and 151-200 (#2_07). A single rain event
6 in late May (57 mm), and a few large precipitation events ($> 25 \text{ mm d}^{-1}$) in July were recorded.
7 The amount of rainfall in 2009 was the smallest among the four years, during which 195mm of
8 irrigation was applied from March to September. There were several short and scattered
9 droughts across the growing season ~~of~~ in 2009 (Fig.2d). Despite the higher- than normal rainfall
10 in the two wet years, there was no flooding or overland runoff.

11 The growing season T_a in 2008 was significantly lower than that in 2007 and 2009 (dT=
12 1.3 ° C, $p < 0.05$, Fig.2 e-h). The years differed in the spring warm-up and the timing of peak
13 temperature (by up to 35.9 ° C). The maximum air temperature occurred in June ~~in~~ 2006 and
14 2009, and in July ~~in~~ 2007 and 2008. The warmest month was June ~~for~~ 2006 ($27.1 \pm 2.4 \text{ ° C}$).

15 The daytime average VPD of the four growing seasons (Fig.2 e-h) was $1.3 \pm 0.7 \text{ kPa}$. The
16 mean VPD in wet years (i.e., 2007 and 2008) was $1.2 \pm 0.7 \text{ kPa}$, which was significantly lower
17 ($F=6.093$, $p < 0.01$) than that in dry years (i.e., 2006 and 2009, $1.3 \pm 0.8 \text{ kPa}$). The VPD of the
18 growing seasons in 2008 (i.e., $1.1 \pm 0.5 \text{ kPa}$) was lower than those in the other years ($p < 0.05$).
19 Higher T_a and lower precipitation in May 2007 led to higher VPD compared with-to the same
20 period in 2006 and 2008 ($p < 0.001$). Furthermore, the VPD was the highest in June 2009 (i.e.,
21 $2.3 \pm 1.1 \text{ kPa}$, $p < 0.05$) and the lowest in 2008 (i.e., $1.0 \pm 0.5 \text{ kPa}$, $p < 0.01$).

22 **3.2 Seasonal changes in energy partitioning and β**

23 The energy partitioning trends of daytime total net radiation (R_n) into latent, sensible heat fluxes
24 (LE and H), soil heat fluxes (G) and heat storage of canopy (S) for the year 2006-2009 were
25 presented in Fig.3. Among these years, R_n varied with solar radiation ($R > 0.95$, $\alpha = 0.01$ level),
26 reached the maximum in July, and gradually decreased until the late October (in dry years) or
27 November (in wet years). During the growing season, there were no significant difference in
28 average daytime total R_n between wet and dry years. The average of daytime total G during the
29 growing season displayed great seasonal and annual differences among these years ($p < 0.05$),
30 with a lower value in wet years (2.1% in 2007) than ~~that of~~ the dry years (4.9% in 2006; $p <$
31 0.001). ~~Additionally, G only accounted for a small proportion of R_n , which ranged from 2.1%~~

1 ~~in 2007 to 4.9% in 2006.~~ Moreover, the average value of daytime total S among the four growing
2 seasons were 0.46 MJ m^{-2} , 0.49 MJ m^{-2} , 0.51 MJ m^{-2} , 0.54 MJ m^{-2} , respectively. S/R_n varied
3 ~~between from~~ 6.0% in 2007 ~~and to~~ 6.8% in 2009 ~~and, showed showing~~ no differences between
4 the wet and dry years.

5 Partitioning of R_n into LE and H differed significantly between the wet and dry years
6 (Table 3; $F = 17.599$, $p < 0.001$). The dominant turbulent energy flux during the early growing
7 season was sensible heat flux (H) with or without drought stress except in 2006 when the
8 irrigation was applied (Table 3). Then LE was the dominant driver of energy partitioning during
9 the middle and late growing seasons under drought stress. The average daytime total LE in wet
10 years was about 20% greater in wet years (6.77 MJ m^{-2}) than that of in dry years (5.72 MJ m^{-2} ,
11 $p < 0.01$). The timing of peak LE was weakly related to drought, peaking in July in 2006, 2008
12 and 2009, and in August in 2007. ~~LE was the dominant turbulent flux with changes of R_n , and~~
13 ~~started to rapidly increase in mid-April and reached a maximum in July for all 3 years (i.e., in~~
14 ~~2006, 2008 and 2009), except but August for 2007.~~ The peak value of daytime total LE was
15 16.61 MJ m^{-2} , 17.01 MJ m^{-2} , 19.72 MJ m^{-2} and 16.27 MJ m^{-2} , in 2006 ~~to~~ 2009 respectively.
16 The daily evaporative fraction (H became the main consumer of the growing season R_n in
17 October for the dry years and in November for the wet years. Among the four years, $LE/(R_n -$
18 $G)$ was significantly higher in wet years (60.3% and 64.8% in 2007 and 2008, respectively)
19 (64.8%) than those in dry years (57.1% and 50.4% in 2006 and 2009, respectively; $p < 0.05$).
20 2006 (57.1%), 2007 (60.3%) and 2009 (50.4%) ($p < 0.05$). $LE/(R_n - G)$ was much lower in 2009
21 than those in other 3 years ($p < 0.01$). ~~Partitioning of R_n into LE and H differed significantly~~
22 ~~between the wet and dry years ($F = 17.599$, $p < 0.001$) (Table 3). The average daytime total LE~~
23 ~~in wet years was greater (6.77 MJ m^{-2}) than that of dry years (5.72 MJ m^{-2} , $p < 0.01$).~~ The
24 ~~dominant turbulent energy flux during the early growing season was sensible heat flux (H) with~~
25 ~~or without drought stress, except in 2006 when the irrigation were applied (Table 3). Then LE~~
26 ~~was the dominant driver of energy partitioning during the middle and late growing season under~~
27 ~~drought stress.~~

28 The seasonal variation of the midday Bowen ratio (β) displayed a rapid and significant
29 trend across the growing season, especially at the beginning (April–June) and the end
30 (September–October) of the growing season (Fig. 4). The Bowen ratios during the middle of
31 the growing seasons were all smaller than 1, and approximately lasted from DOY 180–250 in
32 the dry year and from DOY 180–290 in the wet years. The average midday β ~~of in the~~ dry

1 years was greater (1.57) than that ~~of in~~ the wet years (0.83; $F=19.176$, $p < 0.001$). The Bowen
2 ratio showed differences in response to drought stress across the four growing seasons (Table
3 3); ~~and had with~~ much higher values (> 1) during the drought periods in 2007 and 2009, but
4 not in 2006. The Bowen ratio was smaller than 1 during drought stressed periods in 2008.

5 3.3 Biophysical controls of energy partitioning

6 The R_s varied widely at the beginning and the end of growing season, but changed steadily
7 within a low range during the middle of growing season by comparison. Moreover, these lower
8 R_s in the dry year lasted a shorter period (DOY 190–250) than in the wet year (Fig. 5a). A
9 significantly negative relationship was found between the R_s and LAI during the wet years
10 (Fig.6). Overall, the seasonal average of ~~surface resistance~~ (R_s) normalized by leaf area index
11 (LAI) (i.e., $R_s:LAI$) was lowest during the wettest year in (2008, -54.1 s m^{-1} leaf area) was
12 lowest among the four years (i.e., $p < 0.05$). The $R_s:LAI$ in the dry years (106.8 s m^{-1} leaf area)
13 was 50% higher than in the wet years (71.2 s m^{-1} leaf area) ($p < 0.001$). The $R_s:LAI$ in the
14 seasonal drought stressed periods ~~of in~~ 2006, 2007 and 2009 were greatly much higher than
15 those in during unstressed periods ($p < 0.001$, Table 3). ~~In addition, a significantly negative~~
16 ~~relationship was found between the R_s and LAI during the wet years (Fig.6).~~

17 The average midday R_i peaked in June, and decreased in July/August before reaching a
18 second peak in October (Fig. 5b). The seasonal average R_i during the growing season ranged
19 from 68.3 s m^{-1} to 77.9 s m^{-1} , with a mean value of 74.4 s m^{-1} , and showed no difference among
20 the four growing seasons ($p > 0.05$). Figure 5c presents the seasonal and annual variations of
21 midday R_a . The average R_a for the four growing seasons was 23.2 ± 8.5 s m^{-1} , ranging from 10.6
22 to 43.5 s m^{-1} , 9.7 to 52.5 s m^{-1} , 6.5 to 43.1 s m^{-1} , and 9.7 to 74.5 s m^{-1} , from 2006 to 2009,
23 respectively. R_a in 2007 was significantly higher than ~~that of in~~ the dry years ($p < 0.01$), while
24 R_a in 2008 was smaller than ~~that in~~ the dry years ($p < 0.001$). However, there were no significant
25 differences between dry and wet years R_a ($p > 0.05$).

26 The seasonal changes of LE/LE_{eq} value varied between 0.4 and 1.0 during most of the
27 growing seasons (Fig. 5d). The average LE/LE_{eq} of the four years were 0.76, 0.73, 0.89, and
28 0.63, respectively. The mean LE/LE_{eq} of the dry years (0.68) was lower than that of wet years
29 (0.81; $p < 0.001$). Specifically, the value of LE/LE_{eq} in drought periods of 2007 and 2009 were
30 much smaller. A significantly exponential relationship existed between the LE/LE_{eq} and R_s
31 during the growing season (Fig.7).

1 The decoupling coefficient (Ω) across the growing season peaked in mid-July in 2008 and
2 in early August in the other years (Fig. 5e). The mean Ω for the four years was 0.41, 0.46, 0.43
3 and 0.39 (Table 3), respectively, and was significantly higher in wet year (0.45) than that in dry
4 year (0.40; $F=9.460$, $p < 0.01$). Compared to the value during unstressed periods, the
5 decoupling coefficient during the seasonal drought periods (#1_06, #2_06; #1_07, #2_07 and
6 #1_09, #2_09, #3_09) was much lower in values.

8 4 Discussion

9 4.1 Energy partitioning and Bowen ratio

10 ~~The energy balance ratio (E_{BR}) is a way of evaluating scalar flux estimates from EC techniques.~~
11 ~~The energy balance ratio (E_{BR}) at the current study~~In this study, ~~the closure of the energy budget~~
12 was 0.88 based on daytime 30-minute fluxes, and > 0.96 based on daytime totals (Table 2). The
13 annual mean E_{BR} at our site was similar to the values of eight ChinaFlux sites, which averaged
14 0.83 and ranged from 0.58 to 1.00 (Li et al., 2005). The energy budget is also consistent with
15 the 50 site-years of flux data from 22 in FLUXNET sites, which had energy closure of 0.34–
16 1.69 (Mean = 0.84, Wilson et al., 2002a). A recent analysis of 173 FLUXNET sites also found
17 an average closure of 0.84 (Stoy et al., 2013), although the authors also detected consistent
18 differences among the biomes, and based on metrics of landscape heterogeneity. In addition to
19 the known reasons for decreasing energy balance closure (Hernandez-Ramirez et al., 2010; Li
20 et al., 2005; Nakai et al., 2006; Stoy et al., 2013), management operations at our site (e.g.,
21 irrigation, tilling and partial felling) may also affect the energy balance. Although the causes of
22 surface energy balance closure continues to be debated (Stoy et al., 2013) and will not be
23 conclusively answered in the current study, the results reported here are similar to other
24 FLUXNET sites.

25 The surface energy partitioning to sensible and latent heat depends on water potential
26 gradient and surface resistance (Arain et al., 2003; Baldocchi et al., 2000; Chen et al., 2009).
27 ~~To the extent that e~~Canopy development (Guo et al., 2010), rainfall dynamics and irrigation
28 (Ozdogan et al., 2010) affect these properties to some extent and, ~~they~~ could directly lead to a
29 change in soil moisture and the evaporation component of LE, ~~therefore impact thereby~~
30 impacting energy partitioning and β (Chen et al., 2009; Ozdogan et al., 2010). However, the
31 impact of precipitation on the Bowen ratio may vary by even at any site (Tang et al., 2014). In

1 our study, ~~a~~-detectable responses of $LE/(R_n-G)$ and the Bowen ratio to drought stress and non-
2 stress periods were observed in response to soil water supply (Table 3) with a 50 mm threshold
3 on average (Fig 8). The variability of energy partitioning during the growing season was highly
4 sensitive to water availability from precipitation and irrigation. On an annual scale, the Bowen
5 ratio appeared linearly related to the total growing season precipitation ($R^2=0.89$, $p < 0.05$).
6 Thus, the Bowen ratio is very responsive to the site water supply; a similar finding was reported
7 in [Grünwald and Bernhofer \(2007\)](#) in a temperate spruce forest.

8 By contrast, β varied from 0.18 to 0.71, with a mean of 0.35 ± 0.15 during ~~the most~~
9 ~~partmost~~ of the growing season in 2008 and in the non-stressed periods ~~in-of-the~~ other 3 years; ~~which~~-
10 ~~This variation~~ was close to 0.42 for deciduous forests ([Wilson et al., 2002b](#)) and 0.55 in
11 a temperate Douglas-fir ([Humphreys et al., 2003](#)), which is also similar to the variations in a
12 ponderosa pine forest in the western United States ([Goldstein et al., 2000](#)) and a deciduous
13 broadleaved forest in the southern United States ([Wilson and Baldocchi, 2000](#)). Seasonal
14 drought stress had a discernible impact on the Bowen ratio of this poplar plantation. However,
15 compared to the reported β values such as; 0.74 in a temperate mixed forest ([Wu et al., 2007](#)),
16 0.81 in a boreal Scots pine forest ([Launiainen, 2010](#)) and; 0.89 in a loblolly pine plantation ([Sun](#)
17 [et al., 2010](#)), the average β in wet years were close to the above values. β was higher in seasonal
18 drought periods and dry years than most temperate coniferous forests (Mean = 1.07, ([Wilson et](#)
19 [al., 2002b](#)), which typically had ~~a~~-higher β values. The high β value in this study reflects the
20 semi-arid conditions, and suggests a low tree water supply which might be resulted from the
21 combination of low rainfall, the sandy soil's low water holding capacity ~~of the sandy soil~~, and
22 the high plant and atmospheric water demand. It has been suggested that the large-scale
23 establishment of poplar plantation in sandy semi-arid regions of northern China could have an
24 adverse impact on the region's groundwater reserves ([Li et al., 2014](#); [Petzold et al., 2011](#)). Our
25 findings corroborate the hypothesis that drought would trigger significant changes in the energy
26 partitioning of water-demanding poplar species in a water-stressed region.

27 4.2 Biophysical control on Bowen ratio

28 The Bowen ratio is dependent ~~of-on~~ the interactions of climatic and biological factors ([Perez et](#)
29 [al., 2008](#); [Wilson and Baldocchi, 2000](#)). R_i quantifies the climatic control on energy partitioning
30 and tends to decrease the Bowen ratio. A higher R_i implies a warm and dry climate in continental
31 regions ([Raupach, 2000](#); [Wilson et al., 2002b](#)). R_s reflects the physiological control on surface

1 energy exchange of an ecosystem (Costa et al., 2010; Launiainen, 2010; Zhou et al., 2010); and
2 generally increases the Bowen ratio. ~~Wilson et al. (2002b) reported that R_s was the dominant~~
3 ~~factor in controlling the variability of the Bowen ratio of forests in temperate regions. A linear~~
4 ~~relation was also found between the Bowen ratio and R_s normalized by aerodynamic (R_a) and~~
5 ~~climatological resistance (R_i) parameters (Cho et al., 2012).~~

6 In this study, ~~similar to R_s~~ similarly varied seasonally with plant phenology, and showed
7 similar seasonal characteristics to other deciduous forests during the course of the growing
8 season (Cabral et al., 2010; Kutsch et al., 2008; Li et al., 2012). As reported by Tchebakova et
9 al. (2002), R_s in seasonal drought stressed periods was much higher than that in non-stressed
10 periods. ~~It has been shown that The~~ drought stress during the canopy development affects leaf
11 area and may have lasting effects on canopy gas exchange through the entire growing season,
12 even after the moisture limitation is removed in 2007 led to lower leaf area and higher canopy
13 resistance (e.g., Noormets et al., 2008), which may explain significant difference in R_s between
14 wet year 2007 and 2008 (Fig.9). Compared with the R_s in other ~~research studies~~, the R_s :LAI
15 in dry years ~~of this poplar plantation in the current study~~ was close to that of the Euphrates
16 Poplar (*Populus euphratica* Oliv.) ($130.2 \text{ s m}^{-1} \text{ leaf area LAI}^{-1}$) and smaller than that of the
17 Gansu Poplar (*Populus gansuensis* Wang et Yang) ($189.4 \text{ s m}^{-1} \text{ LAI}^{-1} \text{ leaf area}$) in northwest
18 China semiarid regions (Chen et al., 2004). In wet years it was similar to that of poplar (58.6 s
19 $\text{m}^{-1} \text{ LAI}^{-1} \text{ leaf area}$) ~~in Iceland~~ (Wilson et al., 2002b); and boreal aspen during the full-leaf period
20 ($51.8 \text{ s m}^{-1} \text{ LAI}^{-1} \text{ leaf area}$) ~~in Canada~~ (Blanken et al., 1997) in mesic temperate regions. R_s is
21 primarily driven by solar radiation, moisture availability and VPD (Fernández et al., 2009; Li
22 et al., 2012); and modulated by leaf area and stomatal resistance, which in turn changes as a
23 function of the above factors (Wilson and Baldocchi, 2000). ~~The~~ Compared to the strong
24 correlation between R_s and LAI in wet years, the increased scatter in the R_s -LAI relationship
25 during dry years (Fig.6) suggests that R_s in dry years was also influenced by other physiological
26 and non-physiological (e.g., soil evaporation, canopy structure and turbulence) factors (Wilson
27 et al., 2002b). The mean R_i in ~~this study~~ at the current study was higher than the mean R_i
28 ~~across site year for forests reported for temperate forests~~ in Wilson et al. (2002b) ($t=5.91$,
29 $df=741$, $p < 0.001$), but was ~ 50% lower than the value reported by Li et al. (2009) in a vineyard
30 in Gansu Province in China ($t= -29.87$, $df=741$, $p < 0.001$), ~~likely due to the warm dry climate~~
31 of the northern region in China as might be expected given the predominant climatic conditions.

1 On the seasonal [scale](#), the Bowen ratio and R_s of this poplar plantation were correlated,
2 and consistent with [Wilson et al. \(2002b\)](#) and [Li et al. \(2009\)](#), but differed in wet and dry years
3 ~~(Fig 10). The water limitation during the dry years manifested in disproportional increase in R_s~~
4 ~~than the Bowen ratio; this response may serve as an indicator when water reserves are being~~
5 ~~depleted. At the extremes, the relationship converges, but as water becomes limiting, stomatal~~
6 ~~closure and increased R_s do not appear to be able to affect the seasonal dynamics of the Bowen~~
7 ~~ratio. The Bowen ratio and R_s were linearly related in wet years ($R^2=0.98$, $p < 0.001$), and~~
8 ~~correlated exponentially in dry years ($R^2=0.93$, $p < 0.001$, Fig.10), during which the sensitivity~~
9 ~~of the Bowen ratio on R_s increased with the growing R_s . The partial correlation analysis~~
10 indicated that R_s and R_i , ~~respectively~~, had strong positive and negative effects, ~~respectively~~, on
11 β in both wet and dry years (Table 4), which could not be detected through correlation analysis
12 (e.g., the impact of R_i and R_a on β). Furthermore, ~~the regulation of the Bowen ratio by both~~
13 ~~controlling roles of R_s and R_i on the Bowen ratio in dry years~~ seemed ~~greater stronger in dry~~
14 than ~~that~~ in wet years. ~~Finally~~, R_a had a significant negative impact on the Bowen ratio in wet
15 years, but not in dry years.

16 The average LE/LE_{eq} in the growing season was 0.74 at our site, which is similar to
17 deciduous forests (0.72) ([Wilson et al., 2002b](#)), but smaller than at a temperate broad-leaved
18 forest (0.82) ([Komatsu, 2005](#)). The average Ω value of 0.42 ± 0.22 (0.39-0.46) was close to the
19 other forests (0.26-0.4, [Wilson and Baldocchi, 2000](#); 0.25-0.43, [Motzer et al., 2005](#)). ~~As~~
20 ~~essentially implied by the Penman-Monteith equation, LE/LE_{eq} exponentially related to R_s~~
21 ~~during the growing season Similar to Baldocchi (1994), LE/LE_{eq} declined with increasing R_s~~
22 ~~during the growing season~~ (Fig.7), which is equivalent to the logarithmic relationship between
23 LE/LE_{eq} and G_s (surface conductance) reported by other studies ([Chen et al., 2009](#); [Hossen et](#)
24 [al., 2011](#); [Zhu et al., 2014](#)). The asymptotic value of LE/LE_{eq} in dry years (0.89) and wet years
25 (0.96) were both lower than the 1.1–1.4 range ~~typical in temperate deciduous forest~~ reported
26 by [Monteith \(1995\)](#), indicating that our study site was ~~characterized by drier surface conditions~~
27 ~~than average for the deciduous forest biome drier than these reference sites~~ during both dry and
28 wet years. The low LE/LE_{eq} values under dry surface conditions of the ecosystem in this study
29 may also be related to the high porosity of ~~the~~ sandy soil and ~~a the~~ low ground water table ([Zhao](#)
30 [et al., 2013](#)). Overall, as indicated by the lower Ω values and the significant correlation
31 coefficients between LE/LE_{eq} and R_s , the R_s was the major factor controlling the LE during ~~the~~
32 growing season, which was consistent with the relations between R_s and the Bowen ratio. In
33 addition, LE was more coupled to the atmosphere during the dry years and seasonal drought

1 periods across the growing season, ~~which were~~ reported in other studies ([Bagayoko et al.,](#)
2 [2007](#); [Bracho et al., 2008](#); [Zha et al., 2013](#)).

3 **4.3 Implications for poplar plantation establishments**

4 As forestry is a long-term endeavor, with the economic payback decades from stand
5 establishment, the availability of resources for the stand to prosper should come naturally to
6 natural resource managers. Supplementing limiting resources directly (fertilization, irrigation)
7 or indirectly (competition control, site preparation, thinning) is commonplace in commercial
8 forestry, but it has to be sustainable in the broader context of the region's ecosystems and
9 livelihoods. Earlier, we reported that the water needs of poplar plantation exceed the annual
10 precipitation in the region and plant survival during dry years depends on irrigation from
11 groundwater (Zhang et al., 2014). To our knowledge, there is no and it is hard to develop a
12 metrics for the sustainability of forest plantation, even though there are a couple of studies
13 defining the sustainability of forest plantation by site and plantation productivity for
14 commercial purpose only (e.g., [Richardson et al., 1999](#); [Watt et al., 2005](#)) other than in a broader
15 sense of the plantation and environment interactions that were our focus in the current paper.
16 Our previous study indicated that annual water use of the plantation was even higher than the
17 annual precipitation (Zhang et al., 2014) and thus the irrigation was applied in dry years by
18 pumping groundwater (Table 1). Such water abstraction for irrigating plantation and agriculture
19 crops have led to the dramatic water table decline in the last 30 years (Zhang et al., 2014). In
20 the current study, energy partitioning to latent and sensible heat and surface resistance was
21 dramatically responsive sensitive to climatological drought—even under the irrigation—and
22 as indicated by low LE/LE_{eq} (< 1) and low values of the decoupling coefficient (Ω) ([Zhu et al.,](#)
23 [2014](#)); the dry surface conditions dominated the poplar plantation ~~no matter~~ in both wet ~~or~~ and
24 dry years. In wet years, the plantation itself is in hydrologic balance with the water that arrives
25 as precipitation, with evapotranspiration consuming nearly all of the precipitation. The same is
26 true in dry years, but irrigation increases ET even further by depleting groundwater. Even if the
27 plantations were in hydrologic balance with water delivered as precipitation, their existence and
28 operation could be a threat to adjacent ecosystems and livelihoods if those rely on runoff or
29 groundwater recharge from the areas where the plantation has been sited. In the absence of the
30 plantations it is likely that groundwater recharge would increase, especially given the sandy
31 textured soil that tends to allow rapid infiltration and percolation as well as limits moisture
32 delivery to the atmosphere directly from the soil surface itself. While poplar plantation growth

1 in this water-limited location might be sustained by the modest precipitation in the region, it
2 could still be unsustainable for the broader context of the region's ecosystems and livelihoods.
3 However, further study to truly access these effects is needed by comparing the surface water
4 balance and /or spatial and temporal variations of groundwater levels at an adjacent, similar site
5 without a plantation, which led to the shortage of water use in poplar plantation. In other words,
6 the poplar plantation would consume much water which comes from precipitation or
7 groundwater to maintain its ecological services, while the required irrigation for sustaining
8 these forests may present a threat to the adjacent ecosystems because of their role in reducing
9 ground water table, and may compromise long term sustainability and livelihoods in the region.
10 Therefore, from the viewpoint of hydrologic balance as well as interactions with atmosphere,
11 growing poplar trees in a water stressed region is not sustainable.

13 **5 Conclusions**

14 The seasonal drought stress affected the dynamics of individual turbulent energy fluxes and the
15 surface resistances in the poplar plantation during the growing seasons. Partitioning of available
16 energy into latent (LE) and sensible heat (H) flux responded to meteorological drought and
17 ~~correspondingly displayed~~ resulted in higher β in dry years (1.57) than that in wet years (0.83).
18 Similar to the response of the Bowen ratio on drought conditions, the LAI normalized surface
19 resistance (R_s :LAI) was 33% higher in dry years ~~was 33% higher~~ than ~~that~~ in wet years.
20 ~~Accordingly~~ Correspondingly, the contrasting ~~impact effects~~ of R_s and R_i on the Bowen ratio
21 were stronger in dry years than in wet years, while the effect of R_a was stronger in wet years, ~~;~~
22 R_s was the major factor in controlling energy partitioning during the growing season, as
23 indicated by the relatively low decoupling coefficient (Ω) values. Furthermore, the ~~overall~~ low
24 LE/LE_{eq} (< 1) of poplar plantations indicated that the permanent limitation of plant water use
25 and surface energy partitioning by water availability. Even at mean long-term precipitation, the
26 water demand of poplar plantation may consume nearly all of it and leave little for run-off and
27 groundwater recharge in this semi-arid region, potentially compromising the region's
28 ecosystems and livelihoods. ~~dry climate dominated in this water limited region, which~~
29 ~~suggested that the fast growing and water intensive species like the poplar plantation are poorly~~
30 ~~adapted for the water limited regions.~~

1 *Acknowledgements.* This study was financially supported by the National Special Research
2 Program for Forestry entitled “Forest Management Affecting the Coupling of Ecosystem
3 Carbon and Water Exchange with Atmosphere” ([Grant No. 201204102](#)). First author also
4 thanks the ~~scholarship-financial~~ support by Beijing Municipality Educational Committee under
5 the graduate student training program. Partially supported by the US-China Carbon Consortium
6 (USCCC) is acknowledged as well. Authors thank [Dr. Christopher A. Williams \(Associate](#)
7 [Editor\)](#) and anonymous reviewers for their insightful comments which helped to improve our
8 original manuscript greatly.

11 **References**

- 12 Admiral, S. W., Lafleur, P. M., and Roulet, N. T.: Controls on latent heat flux and energy partitioning at a peat bog
13 in eastern Canada, *Agr Forest Meteorol*, 140, 308-321, 2006.
- 14 Arain, M. A., Black, T. A., Barr, A. G., Griffis, T. J., Morgenstern, K., and Nesic, Z.: Year-round observations of the
15 energy and water vapour fluxes above a boreal black spruce forest, *Hydrol Process*, 17, 3581-3600, 2003.
- 16 Bagayoko, F., Yonkeu, S., Elbers, J., and de Giesen, N. V.: Energy partitioning over the west African savanna: Multi-
17 year evaporation and surface conductance measurements in Eastern Burkina Faso, *J Hydrol*, 334, 545-559, 2007.
- 18 ~~Baldocchi, D.: A comparative study of mass and energy exchange over a closed C3 (wheat) and an open C4 (corn)-
19 canopy: I. The partitioning of available energy into latent and sensible heat exchange, *Agr Forest Meteorol*, 67,
20 191-220, 1994.~~
- 21 Baldocchi, D., Kelliher, F. M., Black, T. A., and Jarvis, P.: Climate and vegetation controls on boreal zone energy
22 exchange, *Global Change Biol*, 6, 69-83, 2000.
- 23 Blanken, P. D., Black, T. A., Yang, P. C., Neumann, H. H., Nesic, Z., Staebler, R., den Hartog, G., Novak, M. D., and
24 Lee, X.: Energy balance and canopy conductance of a boreal aspen forest: Partitioning overstory and understory
25 components, *Journal of Geophysical Research: Atmospheres*, 102, 28915-28927, 1997.
- 26 Bracho, R., Powell, T. L., Dore, S., Li, J. H., Hinkle, C. R., and Drake, B. G.: Environmental and biological controls on
27 water and energy exchange in Florida scrub oak and pine flatwoods ecosystems, *J Geophys Res -Biogeo*, 113,
28 G02004, doi: 10.1029/2007JG000469, 2008.
- 29 Burba, G. G., McDermitt, D. K., Grelle, A., Anderson, D. J., and Xu, L.: Addressing the influence of instrument
30 surface heat exchange on the measurements of CO₂ flux from open-path gas analyzers, *Global Change Biol*, 14,
31 1854-1876, 2008.
- 32 Cabral, O. M. R., Rocha, H. R., Gash, J. H. C., Ligo, M. A. V., Freitas, H. C., and Tatsch, J. D.: The energy and water
33 balance of a Eucalyptus plantation in southeast Brazil, *J Hydrol*, 388, 208-216, 2010.
- 34 Chen, R., Kang, E., Zhang, Z., Zhao, W., Song, K., Zhang, J., and Lan, Y.: Estimation of tree transpiration and
35 response of tree conductance to meteorological variables in desert-oasis system of Northwest China, *Science in*
36 *China Series D: Earth Sciences*, 47, 9-20, 2004.
- 37 Chen, S. P., Chen, J. Q., Lin, G. H., Zhang, W. L., Miao, H. X., Wei, L., Huang, J. H., and Han, X. G.: Energy balance
38 and partition in Inner Mongolia steppe ecosystems with different land use types, *Agr Forest Meteorol*, 149, 1800-
39 1809, 2009.
- 40 China, S. A. o. t. P. s. R. o.: Classification of meteorological drought. In: National Standard of People's Republic of
41 China GB/T 20481-2006, China Standard Press, Beijing, 2006.

- 1 ~~Cho, J., Oki, T., Yeh, P. J. F., Kim, W., Kanae, S., and Otsuki, K.: On the relationship between the Bowen ratio and~~
2 ~~the near-surface air temperature, *Theor Appl Climatol*, 108, 135-145, 2012.~~
- 3 Costa, M. H., Biajoli, M. C., Sanches, L., Malhado, A. C. M., Hutrya, L. R., da Rocha, H. R., Aguiar, R. G., and de
4 Araujo, A. C.: Atmospheric versus vegetation controls of Amazonian tropical rain forest evapotranspiration: Are
5 the wet and seasonally dry rain forests any different?, *J Geophys Res-Biogeog*, 115, G04021, doi:
6 10.1029/2009JG001179, 2010.
- 7 Ding, Y. H., Ren, G. Y., Zhao, Z. C., Xu, Y., Luo, Y., Li, Q. P., and Zhang, J.: Detection, causes and projection of climate
8 change over China: An overview of recent progress, *Adv Atmos Sci*, 24, 954-971, 2007.
- 9 Dou, J. X., Zhang, Y. P., Yu, G. R., Zhao, S. J., Wang, X., and Song, Q. H.: A preliminary study on the heat storage
10 fluxes of a tropical seasonal rain forest in Xishuangbanna, *Sci China Ser D*, 49, 163-173, 2006.
- 11 Falge, E., Baldocchi, D., Olson, R., Anthoni, P., Aubinet, M., Bernhofer, C., Burba, G., Ceulemans, R., Clement, R.,
12 Dolman, H., Granier, A., Gross, P., Grünwald, T., Hollinger, D., Jensen, N.-O., Katul, G., Keronen, P., Kowalski, A., Lai,
13 C. T., Law, B. E., Meyers, T., Moncrieff, J., Moors, E., Munger, J. W., Pilegaard, K., Rannik, Ü., Rebmann, C., Suyker,
14 A., Tenhunen, J., Tu, K., Verma, S., Vesala, T., Wilson, K., and Wofsy, S.: Gap filling strategies for defensible annual
15 sums of net ecosystem exchange, *Agr Forest Meteorol*, 107, 43-69, 2001.
- 16 Fang, S.: Silviculture of poplar plantation in China: A review, *Yingyong Shengtai Xuebao*, 19, 2308-2316, 2008.
- 17 Fernández, M. E., Gyenge, J., and Schlichter, T.: Water flux and canopy conductance of natural versus planted
18 forests in Patagonia, South America, *Trees*, 23, 415-427, 2009.
- 19 Foken, T. and Wichura, B.: Tools for quality assessment of surface-based flux measurements, *Agr Forest Meteorol*,
20 78, 83-105, 1996.
- 21 Gao, Y., Zhu, X., Yu, G., He, N., Wang, Q., and Tian, J.: Water use efficiency threshold for terrestrial ecosystem
22 carbon sequestration in China under afforestation, *Agr Forest Meteorol*, 195-196, 32-37, 2014.
- 23 Gielen, B. and Ceulemans, R.: The likely impact of rising atmospheric CO₂ on natural and managed *Populus*: a
24 literature review, *Environ Pollut*, 115, 335-358, 2001.
- 25 Goldstein, A. H., Hultman, N. E., Fracheboud, J. M., Bauer, M. R., Panek, J. A., Xu, M., Qi, Y., Guenther, A. B., and
26 Baugh, W.: Effects of climate variability on the carbon dioxide, water, and sensible heat fluxes above a ponderosa
27 pine plantation in the Sierra Nevada (CA), *Agr Forest Meteorol*, 101, 113-129, 2000.
- 28 Goulden, M. L., Munger, J. W., Fan, S.-M., Daube, B. C., and Wofsy, S. C.: Measurements of carbon sequestration
29 by long-term eddy covariance: methods and a critical evaluation of accuracy, *Global Change Biol*, 2, 169-182,
30 1996.
- 31 Grünwald, T. and Bernhofer, C.: A decade of carbon, water and energy flux measurements of an old spruce forest
32 at the Anchor Station Tharandt, *Tellus B*, 59, 387-396, 2007.
- 33 Granier, A., Reichstein, M., Breda, N., Janssens, I. A., Falge, E., Ciais, P., Grünwald, T., Aubinet, M., Berbigier, P.,
34 Bernhofer, C., Buchmann, N., Facini, O., Grassi, G., Heinesch, B., Ilvesniemi, H., Keronen, P., Knohl, A., Kostner, B.,
35 Lagergren, F., Lindroth, A., Longdoz, B., Loustau, D., Mateus, J., Montagnani, L., Nys, C., Moors, E., Papale, D.,
36 Peiffer, M., Pilegaard, K., Pita, G., Pumpanen, J., Rambal, S., Rebmann, C., Rodrigues, A., Seufert, G., Tenhunen, J.,
37 Vesala, I., and Wang, Q.: Evidence for soil water control on carbon and water dynamics in European forests during
38 the extremely dry year: 2003, *Agr Forest Meteorol*, 143, 123-145, 2007.
- 39 Guo, H. Q., Zhao, B., Chen, J. Q., Yan, Y. E., Li, B., and Chen, J. K.: Seasonal Changes of Energy Fluxes in an Estuarine
40 Wetland of Shanghai, China, *Chinese Geogr Sci*, 20, 23-29, 2010.
- 41 Guo, I., Li, Q., Yan, C., Mei, X., and Li, Y.: Effects of small area irrigation on water and heat transport of winter
42 wheat field under drought condition[J], *Transactions of the Chinese Society of Agricultural Engineering*, 24, 20-
43 24, 2008.
- 44 Hernandez-Ramirez, G., Hatfield, J. L., Prueger, J. H., and Sauer, T. J.: Energy balance and turbulent flux partitioning
45 in a corn-soybean rotation in the Midwestern US, *Theor Appl Climatol*, 100, 79-92, 2010.

- 1 Holwerda, F., Bruijnzeel, L. A., Scatena, F. N., Vugts, H. F., and Meesters, A. G. C. A.: Wet canopy evaporation from
2 a Puerto Rican lower montane rain forest: The importance of realistically estimated aerodynamic conductance, *J*
3 *Hydrol*, 414, 1-15, 2012.
- 4 Hossen, M. S., Mano, M., Miyata, A., Baten, M. A., and Hiyama, T.: Surface energy partitioning and
5 evapotranspiration over a double-cropping paddy field in Bangladesh, *Hydrol Process*, 26, 1311-1320, doi:
6 10.1002/hyp.8232, 2011.
- 7 Humphreys, E. R., Black, T. A., Ethier, G. J., Drewitt, G. B., Spittlehouse, D. L., Jork, E. M., Nesic, Z., and Livingston,
8 N. J.: Annual and seasonal variability of sensible and latent heat fluxes above a coastal Douglas-fir forest, *British*
9 *Columbia, Canada, Agr Forest Meteorol*, 115, 109-125, 2003.
- 10 Jamiyansharav, K., Ojima, D., Pielke, R. A., Parton, W., Morgan, J., Beltrán-Przekurat, A., LeCain, D., and Smith, D.:
11 Seasonal and interannual variability in surface energy partitioning and vegetation cover with grazing at shortgrass
12 steppe, *J Arid Environ*, 75, 360-370, 2011.
- 13 Jarvis, P. G. and McNaughton, K. G.: Stomatal Control of Transpiration: Scaling Up from Leaf to Region. In:
14 *Advances in Ecological Research*, MacFadyen, A. and Ford, E. D. (Eds.), Academic Press, London NW1 7BY, UK,
15 1986.
- 16 Kim, H.-S., Oren, R., and Hinckley, T. M.: Actual and potential transpiration and carbon assimilation in an irrigated
17 poplar plantation, *Tree Physiol*, 28, 559-577, 2008.
- 18 Komatsu, H.: Forest categorization according to dry-canopy evaporation rates in the growing season: comparison
19 of the Priestley-Taylor coefficient values from various observation sites, *Hydrol Process*, 19, 3873-3896, 2005.
- 20 Kumagai, T. o., Saitoh, T. M., Sato, Y., Morooka, T., Manfroi, O. J., Kuraji, K., and Suzuki, M.: Transpiration, canopy
21 conductance and the decoupling coefficient of a lowland mixed dipterocarp forest in Sarawak, Borneo: dry spell
22 effects, *J Hydrol*, 287, 237-251, 2004.
- 23 Kutsch, W. L., Hanan, N., Scholes, B., McHugh, I., Kubheka, W., Eckhardt, H., and Williams, C.: Response of carbon
24 fluxes to water relations in a savanna ecosystem in South Africa, *Biogeosciences*, 5, 1797-1808, doi: 10.5194/bg-
25 5-1797-2008, 2008.
- 26 Launiainen, S.: Seasonal and inter-annual variability of energy exchange above a boreal Scots pine forest,
27 *Biogeosciences*, 7, 3921-3940, doi: 10.5194/bg-7-3921-2010, 2010.
- 28 Li, S., Tong, L., Li, F. S., Zhang, L., Zhang, B. Z., and Kang, S. Z.: Variability in energy partitioning and resistance
29 parameters for a vineyard in northwest China, *Agr Water Manage*, 96, 955-962, 2009.
- 30 Li, Y. Z., Qin, H. Y., Xie, Y. H., Wang, W., Chen, X. S., and Zhang, C. M.: Physiological mechanism for the reduction
31 in soil water in poplar (*Populus deltoides*) plantations in Dongting Lake wetlands, *Wetl Ecol Manag*, 22, 25-33,
32 2014.
- 33 Li, Z., Niu, L., Yuan, F., Guan, D., Wang, A., Jin, C., and Wu, J.: Canopy conductance characteristics of poplar in
34 agroforestry system in west Liaoning Province of Northeast China . , *Chinese Journal of Applied Ecology*, 23, 2975-
35 2982, 2012.
- 36 Li, Z., Yu, G., Wen, X., Zhang, L., Ren, C., and Fu, Y.: Energy balance closure at ChinaFLUX sites, *Sci China Ser D*, 48,
37 12, 51-62, 2005.
- 38 Mahrt, L.: Stratified Atmospheric Boundary Layers, *Bound-Lay Meteorol*, 90, 375-396, 1999.
- 39 Martín-García, J., Jactel, H., and Diez, J. J.: Patterns and monitoring of *Sesia apiformis* infestations in poplar
40 plantations at different spatial scales, *Journal of Applied Entomology*, 135, 382-392, 2011.
- 41 Migliavacca, M., Meroni, M., Manca, G., Matteucci, G., Montagnani, L., Grassi, G., Zenone, T., Teobaldelli, M.,
42 Goded, I., Colombo, R., and Seufert, G.: Seasonal and interannual patterns of carbon and water fluxes of a poplar
43 plantation under peculiar eco-climatic conditions, *Agr Forest Meteorol*, 149, 1460-1476, 2009.
- 44 Monteith, J.: A reinterpretation of stomatal responses to humidity, *Plant, Cell & Environment*, 18, 357-364, 1995.
- 45 Motzer, T., Munz, N., Küppers, M., Schmitt, D., and Anhuif, D.: Stomatal conductance, transpiration and sap flow
46 of tropical montane rain forest trees in the southern Ecuadorian Andes, *Tree Physiol*, 25, 1283-1293, 2005.

- 1 Nakai, T., Van Der Molen, M., Gash, J., and Kodama, Y.: Correction of sonic anemometer angle of attack errors,
2 *Agr Forest Meteorol*, 136, 19-30, 2006.
- 3 Noormets, A., McNulty, S. G., DeForest, J. L., Sun, G., Li, Q., and Chen, J.: Drought during canopy development has
4 lasting effect on annual carbon balance in a deciduous temperate forest, *New Phytol*, 179, 818-828, 2008.
- 5 Noormets, A., Zhou, R., Chen, J., and Billesbach, D.: EC_Processor page :
6 <http://www4.ncsu.edu/~anoorme/ECP/index.html>, (last access: 20 September 2014), 2010.
- 7 Ozdogan, M., Rodell, M., Beaudoin, H. K., and Toll, D. L.: Simulating the Effects of Irrigation over the United States
8 in a Land Surface Model Based on Satellite-Derived Agricultural Data, *J Hydrometeorol*, 11, 171-184, 2010.
- 9 Paw U, K. T., Baldocchi, D. D., Meyers, T. P., and Wilson, K. B.: Correction Of Eddy-Covariance Measurements
10 Incorporating Both Advective Effects And Density Fluxes, *Bound-Lay Meteorol*, 97, 487-511, 2000.
- 11 Pereira, A. R.: The Priestley–Taylor parameter and the decoupling factor for estimating reference
12 evapotranspiration, *Agr Forest Meteorol*, 125, 305-313, 2004.
- 13 Perez, P. J., Castellvi, F., and Martínez-Cob, A.: A simple model for estimating the Bowen ratio from climatic factors
14 for determining latent and sensible heat flux, *Agr Forest Meteorol*, 148, 25-37, 2008.
- 15 Petzold, R., Schwarzel, K., and Feger, K. H.: Transpiration of a hybrid poplar plantation in Saxony (Germany) in
16 response to climate and soil conditions, *Eur J Forest Res*, 130, 695-706, 2011.
- 17 Qiu, G., Yin, J., and Geng, S.: Impact of Climate and Land-Use Changes on Water Security for Agriculture in
18 Northern China, *Journal of Integrative Agriculture*, 11, 144-150, 2012.
- 19 Raupach, M.: Equilibrium evaporation and the convective boundary layer, *Bound-Lay Meteorol*, 96, 107-142, 2000.
- 20 ~~Richardson, B., Skinner, M. F., and West, G.: The role of forest productivity in defining the sustainability of
21 plantation forests in New Zealand, *Forest Ecol Manag*, 122, 125-137, 1999.~~
- 22 Schmid, H. P., Su, H. B., Vogel, C. S., and Curtis, P. S.: Ecosystem-atmosphere exchange of carbon dioxide over a
23 mixed hardwood forest in northern lower Michigan, *J Geophys Res-Atmos*, 108, 4417, doi:
24 10.1029/2002JD003011, 2003.
- 25 Stanturf, J. A. and Oosten, C. v.: Operational Poplar and Willow Culture. In: *Poplars and willows: trees for society
26 and the environment*, Isebrands, J. G. and Richardson, J. (Eds.), The Food and Agriculture Organization of the
27 United Nations and CABI, Available from: <http://www.fao.org/forestry/ipc/69946@158687/en/> (last access: April
28 3, 2015), 2014.
- 29 Stoy, P. C., Mauder, M., Foken, T., Marcolla, B., Boegh, E., Ibrom, A., Arain, M. A., Arneth, A., Aurela, M., Bernhofer,
30 C., Cescatti, A., Dellwik, E., Duce, P., Gianelle, D., van Gorsel, E., Kiely, G., Knohl, A., Margolis, H., McCaughey, H.,
31 Merbold, L., Montagnani, L., Papale, D., Reichstein, M., Saunders, M., Serrano-Ortiz, P., Sottocornola, M., Spano,
32 D., Vaccari, F., and Varlagin, A.: A data-driven analysis of energy balance closure across FLUXNET research sites:
33 The role of landscape scale heterogeneity, *Agr Forest Meteorol*, 171–172, 137-152, 2013.
- 34 Sun, G., Noormets, A., Gavazzi, M., McNulty, S., Chen, J., Domec, J.-C., King, J., Amatya, D., and Skaggs, R.: Energy
35 and water balance of two contrasting loblolly pine plantations on the lower coastal plain of North Carolina, USA,
36 *Forest Ecol Manag*, 259, 1299-1310, 2010.
- 37 Takagi, K., Kimura, R., and Şaylan, L.: Variability of surface characteristics and energy flux patterns of sunn hemp
38 (*Crotalaria juncea* L.) under well-watered conditions, *Theor Appl Climatol*, 96, 261-273, 2009.
- 39 Tang, Y., Wen, X., Sun, X., and Wang, H.: Interannual Variation of the Bowen Ratio in a Subtropical Coniferous
40 Plantation in Southeast China, 2003-2012, *Plos One*, 9, e88267, doi: 10.1371/journal.pone.0088267, 2014.
- 41 Tchebakova, N. M., Kolle, O., Zolotoukhine, D., Arneth, A., Styles, J. M., Vygodskaya, N. N., Schluze, E.-D.,
42 Shibistova, O., and Lloyd, J.: Inter-annual and seasonal variations of energy and water vapour fluxes above
43 a *Pinus sylvestris* forest in the Siberian middle taiga, *Tellus B*, 54, 537-551, 2002.
- 44 Wang, E., Yu, Q., Wu, D., and Xia, J.: Climate, agricultural production and hydrological balance in the North China
45 Plain, *Int J Climatol*, 28, 1959-1970, 2008.

- 1 ~~Watt, M. S., Coker, G., Clinton, P. W., Davis, M. R., Parfitt, R., Simcock, R., Garrett, L., Payn, T., Richardson, B., and~~
2 ~~Dunningham, A.: Defining sustainability of plantation forests through identification of site quality indicators~~
3 ~~influencing productivity—A national view for New Zealand, *Forest Ecol Manag*, 216, 51–63, 2005.~~
- 4 Webb, E. K., Pearman, G. I., and Leuning, R.: Correction of flux measurements for density effects due to heat and
5 water vapour transfer, *Q J Roy Meteor Soc*, 106, 85–100, 1980.
- 6 Wilczak, J., Oncley, S., and Stage, S.: Sonic Anemometer Tilt Correction Algorithms, *Bound-Lay Meteorol*, 99, 127–
7 150, 2001.
- 8 Wilske, B., Lu, N., Wei, L., Chen, S. P., Zha, T. G., Liu, C. F., Xu, W. T., Noormets, A., Huang, J. H., Wei, Y. F., Chen, J.,
9 Zhang, Z. Q., Ni, J., Sun, G., Guo, K., McNulty, S., John, R., Han, X. G., Lin, G. H., and Chen, J. Q.: Poplar plantation
10 has the potential to alter the water balance in semiarid Inner Mongolia, *J Environ Manage*, 90, 2762–2770, 2009.
- 11 Wilson, K., Goldstein, A., Falge, E., Aubinet, M., Baldocchi, D., Berbigier, P., Bernhofer, C., Ceulemans, R., Dolman,
12 H., Field, C., Grelle, A., Ibrom, A., Law, B. E., Kowalski, A., Meyers, T., Moncrieff, J., Monson, R., Oechel, W.,
13 Tenhunen, J., Valentini, R., and Verma, S.: Energy balance closure at FLUXNET sites, *Agr Forest Meteorol*, 113,
14 223–243, 2002a.
- 15 Wilson, K. B. and Baldocchi, D. D.: Seasonal and interannual variability of energy fluxes over a broadleaved
16 temperate deciduous forest in North America, *Agr Forest Meteorol*, 100, 1–18, 2000.
- 17 Wilson, K. B., Baldocchi, D. D., Aubinet, M., Berbigier, P., Bernhofer, C., Dolman, H., Falge, E., Field, C., Goldstein,
18 A., Granier, A., Grelle, A., Halldor, T., Hollinger, D., Katul, G., Law, B. E., Lindroth, A., Meyers, T., Moncrieff, J.,
19 Monson, R., Oechel, W., Tenhunen, J., Valentini, R., Verma, S., Vesala, T., and Wofsy, S.: Energy partitioning
20 between latent and sensible heat flux during the warm season at FLUXNET sites, *Water Resour Res*, 38, 1294,
21 doi:10.1029/2001WR000989, 2002b.
- 22 Wu, J. B., Guan, D. X., Han, S. J., Shi, T. L., Jin, C. J., Pell, T. F., and Yu, G. R.: Energy budget above a temperate mixed
23 forest in northeastern China, *Hydrol Process*, 21, 2425–2434, 2007.
- 24 Zha, T. S., Li, C. Y., Kellomaki, S., Peltola, H., Wang, K. Y., and Zhang, Y. Q.: Controls of Evapotranspiration and CO₂
25 Fluxes from Scots Pine by Surface Conductance and Abiotic Factors, *Plos One*, 8, e69027, doi:
26 10.1371/journal.pone.0069027, 2013.
- 27 Zhang, J. B., Shangguan, T., and Meng, Z. Q.: Changes in soil carbon flux and carbon stock over a rotation of poplar
28 plantations in northwest China, *Ecol Res*, 26, 153–161, 2011.
- 29 Zhang, Y., Kadota, T., Ohata, T., and Oyunbaatar, D.: Environmental controls on evapotranspiration from sparse
30 grassland in Mongolia, *Hydrol Process*, 21, 2016–2027, 2007.
- 31 Zhang, Y., Zhang, Z., Sun, G., Fang, X., Zha, T., Noormets, A., McNulty, S., Chen, J., Liu, c., and Chen, L.: Water
32 Balances of a Poplar Plantation Forest in Suburban Beijing, China In: *J Environ Manage*, 2014 (in review).
- 33 Zhao, Q., Guo, W., Ling, X., Liu, Y., Wang, G., and Xie, J.: Analysis of Evapotranspiration and Water Budget for
34 Various Land Use in Semi-Arid Areas of Tongyu, China, *Climatic and Environmental Research*, 18, 415–426, 2013
35 (in Chinese).
- 36 Zhou, J., Zhang, Z., Sun, G., Fang, X., Zha, T., McNulty, S., Chen, J., Jin, Y., and Noormets, A.: Response of ecosystem
37 carbon fluxes to drought events in a poplar plantation in Northern China, *Forest Ecol Manag*, 300, 33–42, 2013.
- 38 Zhou, L., Zhou, G. S., Liu, S. H., and Sui, X. H.: Seasonal contribution and interannual variation of
39 evapotranspiration over a reed marsh (*Phragmites australis*) in Northeast China from 3-year eddy covariation data,
40 *Hydrol Process*, 24, 1039–1047, 2010.
- 41 Zhu, G., Lu, L., Su, Y., Wang, X., Cui, X., Ma, J., He, J., Zhang, K., and Li, C.: Energy flux partitioning and
42 evapotranspiration in a sub-alpine spruce forest ecosystem, *Hydrol Process*, 28, 5093–5104, 2014.
- 43
44
45

1

2

3

4

5

6

7

8

1
2
3
4
5
6
7
8
9

Table 1. The stand characteristics of four years from 2006 to 2009, including the minimum, maximum and mean temperature (T), the annual precipitation (P), evapotranspiration (ET), irrigation (I), canopy height (H), breast height diameter (DBH), leaf area index (LAI). The error estimates are standard deviation (SD).

	Tmin	Tmax	Tmean	P	ET	I	H	DBH	LAI
	(°C)			(mm)	(mm)	(mm)	(m)	(cm)	(m ² m ⁻²)
2006	-10.6	29.7	12.5±0.73	482	599	86	11.5±1.1	10.8±1.5	1.6±0.3
2007	-9.8	29.5	13.0±0.55	667	560	-	13.0±1.3	12.2±1.8	2.1±0.4
2008	-7.4	28.8	13.3±0.54	662	653	-	14.8±1.2	13.8±1.8	2.2±0.7
2009	-10.2	30.5	12.5±0.60	428	511	195	16.2±1.6	14.5±1.6	2.9±0.4

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18

Table 2. Energy balance closure statistics using half-hourly and daytime totals during growing season from 2006 to 2009

	daytime				Daytime sum			
	2006	2007	2008	2009	2006	2007	2008	2009
Slope	0.92	0.87	0.92	0.82	1.07	0.91	1.04	0.84
Intercept	20.50	17.24	10.72	13.08	-0.63	-0.09	-0.79	-0.30
R^2	0.81	0.80	0.81	0.82	0.88	0.81	0.92	0.82

Daytime was defined as the period between the sunrise and sunset with $PAR > 4 \text{ umol m}^{-2} \text{ s}^{-1}$;
The unit of Intercept for Half-hourly value and Daytime sum value were $\text{W} \cdot \text{m}^{-2}$ and $\text{MJ} \cdot \text{m}^{-2}$, respectively.

1 Table 3. The value of the soil water supply (WS), energy partitioning ratios and biophysical variables in the different periods of the growing
 2 season during 2006-2009

Year	Periods(DOY)	WS (mm)	LE/(R _n -G)(%)	H/(R _n -G)(%)	β	R _s (s m ⁻¹)	R _i (s m ⁻¹)	R _a (s m ⁻¹)	α	Ω
2006	100-163	76.2+56	50.5(23.4)	45.9(19.7)	3.48(6.37)	418.7(528.7)	87.8(30.2)	20.0(6.3)	0.64(0.35)	0.25(0.13)
	164-192 ^d	127.8	68.0(13.3)	33.2(11.1)	0.66(0.35)	184.0(94.7)	94.9(45.2)	23.8(5.1)	0.79(0.19)	0.42(0.14)
	193-230	219.6	77.7(11.9)	13.8(6.7)	0.19(0.13)	50.4(29.9)	51.5(16.4)	27.8(8.6)	1.01(0.24)	0.70(0.12)
	231-300 ^d	43	51.9(12.7)	31.7(11.6)	0.94(0.52)	178.5(68.8)	77.4(27.5)	25.6(6.8)	0.69(0.23)	0.36(0.14)
2007	100-143 ^d	61.8	35.2(6.4)	57.8(8.3)	2.37(0.66)	426.9(148.8)	96.1(29.4)	18.1(5.4)	0.41(0.13)	0.16(0.07)
	151-200 ^d	146.8	49.5(18.2)	37.0(17.7)	1.41(1.06)	314.1(225.6)	91.7(42.8)	25.3(7.1)	0.58(0.23)	0.35(0.16)
	200-300	396.8	66.0(16.3)	15.5(8.5)	0.35(0.32)	74.1(27.3)	61.1(22.7)	30.4(9.2)	0.87(0.20)	0.60(0.15)
2008	100-117	53.4	16.3(14.1)	71.8(9.7)	1.86(1.12)	206.9(102.0)	60.7(22.9)	13.6(4.1)	0.59(0.35)	0.21(0.14)
	118-155 ^d	15.6	58.8(12.3)	39.5(10.7)	0.71(0.36)	130.8(48.6)	81.1(32.3)	14.7(4.2)	0.81(0.23)	0.31(0.11)
	156-188	212.7	68.1(14.6)	33.3(10.7)	0.35(0.23)	70.2(33.4)	56.1(20.6)	19.3(5.9)	0.94(0.23)	0.53(0.14)
	189-212 ^d	26	73.5(12.7)	20.4(7.5)	0.18(0.15)	59.3(27.1)	67.4(41.1)	27.8(6.8)	1.07(0.25)	0.68(0.11)
	213-239	173.4	74.8(11.9)	11.8(6.2)	0.24(0.16)	61.5(23.7)	55.8(14.3)	19.3(5.2)	0.92(0.14)	0.57(0.10)
	240-251 ^d	19.2	60.4(12.6)	23.4(9.9)	0.42(0.22)	88.7(34.6)	60.4(15.3)	18.0(4.1)	0.87(0.21)	0.46(0.10)
2009	252-300	116.2	47.2(5.7)	39.2(3.6)	0.41(0.22)	72.1(17.8)	57.3(28.9)	18.4(4.4)	0.85(0.23)	0.48(0.10)
	100-158 ^d	37.6+52	36.0(16.5)	48.8(13.4)	1.90(0.83)	298.9(150.8)	84.2(39.3)	18.2(3.8)	0.43(0.19)	0.21(0.08)

	165-186 ^d	1.2	47.8(15.6)	38.1(14.8)	1.32(0.78)	360.5(139.8)	137.4(43.8)	21.2(5.9)	0.53(0.28)	0.24(0.10)
	187-235	265+32	65.9(12.8)	12.4(6.7)	0.28(0.18)	61.2(30.9)	53.0(22.8)	27.4(6.6)	0.82(0.18)	0.66(0.13)
	236-300 ^d	20.4+20	50.4(20.5)	33.1(18.4)	1.28(1.31)	208.3(194.3)	72.3(26.5)	26.9(10.7)	0.64(0.28)	0.39(0.21)
2006	Growing season	466+86	59.1(18.9)	31.8(16.4)	1.60(3.94)	231.4(338.3)	77.9(33.6)	24.0(7.4)	0.76(0.30)	0.41(0.21)
2007	Growing season	630	56.6(19.5)	28.7(19.6)	0.93(0.98)	192.2(190.7)	75.4(34.0)	26.9(9.3)	0.73(0.44)	0.46(0.22)
2008	Growing season	630	66.1(15.2)	22.1(13.4)	0.73(1.04)	118.1(115.3)	68.3(44.9)	18.5(6.3)	0.89(0.59)	0.43(0.19)
2009	Growing season	400+195	48.5(21.9)	34.6(18.5)	1.54(2.19)	248.9(273.3)	77.1(39.1)	23.8(8.5)	0.63(0.38)	0.39(0.24)
dry years										
(2006, 2009)	Growing season	-	52.6(22.3)	33.0(18.4)	1.57(3.17)	240.3(306.9)	77.5(36.5)	23.9(8.0)	0.68(0.31)	0.40(0.22)
wet years										
(2007, 2008)	Growing season	-	61.5(18.1)	25.1(17.0)	0.83(1.01)	153.1(159.7)	71.6(40.3)	22.5(8.9)	0.81(0.29)	0.45(0.20)

WS: soil water supply of period (sum of precipitation and irrigation); β : Bowen ratio; R_s , the surface resistance; R_i , the climatological resistance; R_a , the aerodynamic resistance; α , the Priestley-Taylor coefficient; Ω , the decoupling coefficient;

^d indicate the drought stressed periods.

The value in table represents Mean (SD), the superscript uppercase letters (A, B, C) and lowercase letters (a, b, c) respectively indicate the significance at the 0.01 level and the 0.05 level.

Table 4. The correlation analysis between the Bowen ratio (β) and R_s , R_i and R_a .

		Partial correlation analysis*			Correlation analysis		
		<i>SOCC</i>	<i>p</i>	df	Pearson	<i>p</i>	df
dry year	β & R_s	0.965	<0.001		0.939	<0.001	
	β & R_i	-0.667	<0.001	347	-0.042	=0.436	349
	β & R_a	0.037	=0.496		-0.221	<0.001	
wet year	β & R_s	0.905	<0.001		0.85	<0.001	
	β & R_i	-0.614	<0.001	383	0.64	=0.006	385
	β & R_a	-0.217	<0.001		-0.286	<0.001	

*Partial correlation analysis was proceeded between Bowen ratio and each of three resistance parameters (R_s , R_i and R_a) with the other two as controlling variables.

SOCC: The abbreviation of *Second-order correlation coefficient*.

5

10




Figure 1. The cumulative precipitation (P) and periodic irrigation during 2006-2009, irrigation in 2006 and 2009 were separately represented by the solid and dotted bracebrackets, respectively.

15

20




Figure 2. The seasonal variation of environmental conditions during 2006-2009, a-d: the relative extractable water (REW) (drought periods longer than 20 days are shaded), daily sum of precipitation (P); e-h: daytime mean air temperature (T_a), daytime mean air vapor deficit (VPD).





Figure 3. Seasonal patterns of daytime energy components (5-day running average) during the growing season from 2006 to 2009, including net radiation (R_n), latent heat (LE), sensible heat (H) and soil heat flux (G) and heat storage term (S).

5

10



15

Figure 4. Seasonal and inter-annual variability of the midday (10:00-15:00 LST) mean Bowen ratio (β) (5-day running average) across the growing season, with detailed β during DOY 185-255 representing in small pane; ~~Midday means the time course from 10:00 a.m. to 15:00 p.m. at local standard time.~~

20




Figure 5. Seasonal dynamics of the midday (10:00-15:00 LST) mean surface resistance (R_s), climatological resistance (R_i), aerodynamic resistance (R_a), LE/LE_{eq} and decoupling coefficient (Ω) (5-day running average) across the growing season from 2006 to 2009. ~~Midday means the time course from 10:00 a.m. to 15:00 p.m. LST.~~

5

5




Figure 6. The relationship between leaf area index (LAI) and surface resistance (R_s) during growing season of the wet and dry year.

10

15




Figure 7. The relationships between surface resistance (R_s) and LE/LE_{eq} (Priestley-Taylor coefficient) during growing season of the wet (a) and dry (b) year.




Figure 8. The response of Bowen ratio and $LE/(R_n-G)$ on Water Supply (WS) (including precipitation (P) and irrigation (I) during individual period) of the different periods across the 10 four growing seasons.

5





Figure 9. Seasonal variations of monthly average LAI and R_s during the growing season in wet year 2007 and 2008.

15

20



5 Figure 10. Response of monthly average Bowen ratio (β) on surface resistance (R_s) in the wet and dry year.