

Quantifying wind and pressure effects on trace gas fluxes

K. R. Redeker et al.

Quantifying wind and pressure effects on trace gas fluxes across the soil–atmosphere interface

K. R. Redeker¹, A. J. Baird², and Y. A. Teh³

¹Department of Biology, University of York, Heslington, York, YO10 5DD, UK

²School of Geography, University of Leeds, UK

³Institute of Biological and Environmental Sciences, University of Aberdeen, UK

Received: 31 December 2014 – Accepted: 2 March 2015 – Published: 24 March 2015

Correspondence to: K. R. Redeker (kelly.redeker@york.ac.uk)

Published by Copernicus Publications on behalf of the European Geosciences Union.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Abstract

Large uncertainties persist in estimates of soil–atmosphere exchange of important trace gases. One significant source of uncertainty is the combined effect of wind and pressure on these fluxes. Wind and pressure effects are mediated by surface topography: few surfaces are uniform and over scales of tenths of a meter to tens of meters, air pressure and wind speed at the ground surface may be very variable. In this paper we consider how such spatial variability in air pressure and wind speed affects fluxes of trace gases. We used a novel nested wind tunnel design, comprising a toroidal wind tunnel in which wind speed and pressure may be controlled, set within a larger, linear wind tunnel. The effects of both wind speed and pressure differentials on fluxes of CO₂ and CH₄ within three different ecosystems (forest, grassland, peat bog) were quantified. We find that trace gas fluxes are positively correlated with both wind speed and pressure differential near the surface boundary. We argue that wind speed is the better proxy for trace gas fluxes because of its stronger correlation and because wind speed measurement is more easily accomplished and wind speed measurement methodology can be more easily standardized. Trace gas fluxes, whether into or out of the soil, increase with wind speed within the toroidal tunnel (+54 % flux per m s⁻¹), while faster, localized surface winds that are external to the toroidal wind tunnel reduce trace gas fluxes (–11 % flux per m s⁻¹). These results are consistent for both trace gases over all ecosystem soil types studied. Our findings support the need for a revised conceptualization of soil–atmosphere gas exchange. We propose a conceptual model of the soil profile that has a “mixed layer”, with fluxes controlled by wind speed, wind duration, porosity, water table, and gas production and consumption.

BGD

12, 4801–4832, 2015

Quantifying wind and pressure effects on trace gas fluxes

K. R. Redeker et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



1 Introduction

Soils play a key role in the production, sequestration, consumption and release of all climatically-important trace gases. Soils contribute greater than 25 % of surface fluxes of CO₂ to the atmosphere, while a substantial fraction of the sources (> 30 %) and sinks (> 5 %) of CH₄ are driven by soil microbial processes (Holmen and Jaffe, 2000; Wuebbles and Hayhoe, 2002).

The movement of gases within soils has been reviewed by, *inter alios*, Hillel (1998), Scanlon et al. (2000), Rolston and Moldrup (2012) and Monson and Baldocchi (2014). Gas movement may occur via diffusion and/or advection. Different types of diffusion can occur in a soil, although the most important is “ordinary” or molecular diffusion. Ordinary diffusion involves the transport of a gas along a gas concentration or mole fraction gradient. Ordinary diffusion of a mixture of two gases is usually modeled using Fick’s second law, while, for mixtures of three or more gases, the Stefan–Maxwell equations may be used (Rolston and Moldrup, 2012). Advective fluxes are typically modelled with Darcy’s law which is usually used in combination with the continuity equation.

Little empirical work has been done on the relative importance of gas diffusion and advection in soils. Despite the lack of substantial empirical evidence, Rolston and Moldrup (2012) suggest that diffusive flow is more important than advective flow. Their suggestion is also commonly assumed by scientists measuring trace gas fluxes using closed chambers. Static and dynamic flux chambers are widely employed to measure soil–atmosphere trace gas exchanges, but are usually set up such that diffusion-only conditions prevail (no or slow circulation of fan air) or under unrealistic conditions of within-chamber air flow (constant air flow generated by a single fan or set of fans) (see, e.g., Denmead, 2008; Rochette, 2011) which give an undefined combination of diffusion and advection. Gradient flux measurements also rely upon this basic assumption (Myklebust et al., 2008).

In general there is considerable uncertainty about the degree to which chambers provide reliable measurements, and problems with chamber use are discussed in the

BGD

12, 4801–4832, 2015

Quantifying wind and pressure effects on trace gas fluxes

K. R. Redeker et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



reviews by Denmead (2008) and Rochette (2011). The use of fans provides a good example of this uncertainty. Some authors, such as Davidson et al. (2002), suggest that chambers fitted with fans give unreliable readings. In contrast, Christiansen et al. (2011) found that, only in chambers in which the air was mixed by a fan, was the measured flux similar to reference fluxes (they introduced methane (CH_4) at controlled rates through the base of various laboratory sand beds – some dry and some wet – and used chambers to record the fluxes above the sand). Furthermore, Denmead (2008) notes that chambers without fans or with fixed wind speeds may give unrealistic flux estimates, especially during windy conditions in the environment outside of the chambers. To illustrate the problem, he cites Denmead and Reicosky (2003) who, in a study of a tilled soil, found that, while carbon dioxide (CO_2) fluxes within a chamber with a fixed-speed fan stayed steady, those in the area around the chamber (as measured using a micrometeorological dispersion method) increased with ambient wind speed.

Even if we assume that diffusive fluxes are an important form of gaseous movement in soils, such fluxes are highly sensitive to gradients in local soil gas concentrations. Variations in these local soil gas concentrations (and hence surface–atmosphere fluxes) can be caused by a range of mechanisms including (i) horizontal and vertical variations in abiotic and biotic processes (Segers, 1998) and (ii) advection through soil pore networks of gas mixtures that have different compositions from those they are replacing.

Spatial variation in soil trace gas profiles are determined by a complex set of biological, chemical and physical processes (Holmen and Jaffe, 2000; Montzka, Reimann et al., 2010). For instance, CO_2 is produced biologically in soils by respiration, contingent upon the vertical distribution of roots, hyphae and labile organic C, temperature, moisture, redox state and CO_2 concentration. Other trace gases, including CH_4 , are both produced and consumed by separate groups of microbes that reside in different locations (at different depths or different locations at the same depth) within soils. Local gas concentrations are also dependent upon the residence time of the trace gas in the soil profile, since first order chemical and biological consumption rates are time and con-

BGD

12, 4801–4832, 2015

Quantifying wind and pressure effects on trace gas fluxes

K. R. Redeker et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



centration dependent. Sufficiently high local concentrations can either lead to negative feedbacks (reduced root respiration rates; Qi et al., 1994) or greater consumption of the gas of interest (CH₄; Wuebbles and Hayhoe, 2002). Gas residence time will depend on the processes transporting gases through and within soils.

5 Advection may significantly affect local gas residence time. Advection of soil gases occurs when there is a pressure gradient between the air in the soil and that in the overlying atmosphere. Horizontal pressure gradients and horizontal advection may also occur. Pressure gradients form under a range of circumstances. Variations in wind speed at the soil surface both over time and spatially can lead to variations in pressure
10 within the soil profile. Percolation of water through the soil profile and spatial variations in soil temperature may also be the cause of within-soil pressure variations.

Empirical and modeling studies have shown that, through their effect on advection, soil–atmosphere pressure differentials can alter the direction and magnitude of gas fluxes substantially ($\pm \leq 1000\%$) (Yonemura et al., 2000; Takle et al., 2004; Xu et al., 2006; Flechard et al., 2007; Reicosky et al., 2008; Bowling and Massman, 2011; Schack-Kirchner et al., 2012; Rey et al., 2012). The suggested mechanism for this process is that localized pressure differentials, driven by spatially and temporally variable winds, create a push-pull mechanism by which soil pore spaces are mixed with neighboring pores and overlying air (Webster and Taylor, 1992, Massman, 2006). This
20 mechanism has been shown to be significantly more effective than diffusion alone in driving soil–atmosphere fluxes (Massman and Frank, 2006; Bowling and Massman, 2011; Schack-Kirchner et al., 2012) and is particularly affected by abrupt boundary transitions (e.g. – (1) a stone, or a fence, in a field or (2) the edge of a dense vegetation patch).

25 While these published studies note the importance of advective transport in surface–atmosphere fluxes, they have not systematically quantified its importance over a broad range of environments, soil types or wind states. Likewise, the majority of these studies have only observed one trace gas at a time, reducing our ability to generate broadly applicable rules for surface–atmosphere trace gas fluxes. Here we close this knowledge

BGD

12, 4801–4832, 2015

Quantifying wind and pressure effects on trace gas fluxes

K. R. Redeker et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Quantifying wind and pressure effects on trace gas fluxes

K. R. Redeker et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



gap by using a novel nested wind tunnel (Fig. 1) to investigate the role of advection in regulating soil–atmosphere gas exchange for two different trace gases, each of which is controlled by very different processes at different depths within the soil. Carbon dioxide, under dark conditions, is predominantly produced through plant, fungal and bacterial respiration and will have high soil concentrations (relative to the atmosphere) close to the soil surface. In contrast methane, whose biological response in soils is broadly insensitive to sunlight, is often consumed by aerobic soils and therefore has lower than atmospheric concentrations within the soil column. At greater depths within the soil profile, in anaerobic regions, methane can be produced by methanogenic archaea but much of this produced methane is consumed by methylotrophic bacteria in the regions directly above the production zone.

Using four sites, we investigated three different ecosystem types: peat bog (two sites), evergreen coniferous forest, and managed grassland. We use the empirical data that we collected to build upon the model proposed by Massman (2006) in which diffusive flow is enhanced by pressure-based mixing. Based on our measured flux data we propose two modifications, including (i) a “mixed layer” of soil pore spaces near the soil surface that, depending on wind speed, has a similar gas composition to the atmosphere immediately above the surface, and (ii) an inherent likelihood of horizontal gas flow through advective/diffusive mechanisms, which can affect observed trace gas fluxes.

2 Methods

2.1 The nested wind tunnel (NWT)

In order to quantify the impact of local (≤ 1 m) and microscale (in the meteorological sense; 1 m to 1 km) winds on trace gas fluxes from various ecosystem surfaces, we required an experimental design that allowed us to vary local (≤ 1 m) wind speeds and atmospheric pressures concurrently. We resolved this difficulty by nesting an isolated,

toroidal wind tunnel within a larger, straight-line wind tunnel. By increasing wind speeds in both wind tunnels we were able to maintain similar pressures in the local space of the toroidal wind tunnel (1 m^2), relative to zero-wind speed conditions, under higher wind speeds (up to 4 m s^{-1}) (Fig. 1).

Current flux measurement methodology relies, in many cases, on the assumption that diffusive flux is dominant within the system. The nested wind tunnel allowed us to test a number of different natural-world scenarios in which this assumption may not be valid. For instance, with fast winds both within the toroidal tunnel and externally, within the straight-line wind tunnel (similar to an entire region experiencing a windy day) we can examine whether faster winds drive more rapid mixing of air within soil pores. Alternately, if we keep the air flow within the toroid at zero and increase the wind speeds externally, within the straight-line wind tunnel (similar to a sheltered forest/field edge near open land), we can examine the influence of greater mixing within the soils external to the toroid and the impact of horizontal mixing within the soil column.

2.1.1 The inner, toroidal wind tunnel

The inner wind tunnel was a toroid, equipped with internal fans, which can generate wind speeds up to 6 m s^{-1} . The toroid was constructed from acrylic and was 40 cm high, 1 m in diameter, and had an internal chamber 30 cm in diameter (Fig. 1). These dimensions created a tunnel footprint of 1.015 m^2 , with an internal volume of 428 L. If the toroid is considered within a compass ordinate system there were two sets of three high-speed computer fans (5214 NH, EBM-Papst, Mulfingen, Germany) placed at North and South, 20 cm above the soil surface as well as two digital anemometers (ATP Instrumentation; Leicestershire, UK) placed at West and East, 22 cm above the soil surface. Anemometers were tested in various locations within the toroid, from near the inner, bottom edge to near the outer, top edge and were found to record similar wind speeds in all locations; therefore, the anemometers were ultimately placed for ease of access.

BGD

12, 4801–4832, 2015

Quantifying wind and pressure effects on trace gas fluxes

K. R. Redeker et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Four separate 30 cm diameter removable vents were located at each compass ordinate, although in practice only those located over the anemometers were covered (during measurements) or uncovered (during equilibration periods) (Fig. 1). During gas flux measurements each vent was covered and pressure sealed with silicone gaskets.

Internal air temperature probes (DT-612, Thermosense, Manchester, UK) and pressure differential gauges (264, Setra Systems Inc., Boxborough, MA, USA) were located at the top of the apparatus above the anemometers and penetrated 15 and 2 cm respectively into the toroidal tunnel.

The installation of the toroid at each site occurred within 48 h of tests being run. At our forest site, one of the two bog sites (Forsinard – see below) and the managed grassland it was sealed at the soil surface using wet sand, while at the second bog site (Cors Fochno – see below) its weight caused it to sink slightly into the peat so that its lower edge was below the water table. Sealing of the toroid was required to maintain/isolate its air mass over the course of each experiment.

2.1.2 The outer wind tunnel

The straight-line wind tunnel enclosing the toroid comprised a standard aluminium and wooden agricultural tunnel (FirstTunnels, Lancashire, UK) (3.5 m long × 2 m wide × 1.5 m high) with the option to be covered by PAR transparent or opaque plastic sheeting. This option allowed the combined wind tunnel system to be capable of examining the soil-plant-atmosphere system under either respiration- or photosynthesis-dominant conditions (Fig. 1). Only “dark” results are shown here. In terms of soil-atmosphere CO₂ exchanges, diffusion will almost always occur from soil to atmosphere because soil CO₂ concentrations are higher than those in the atmosphere above due to ongoing respiration by plants, fungi and bacteria. By using dark conditions, we were able to remove photosynthetic uptake of CO₂ and its assimilation into plant tissue as a confounding factor. That is, we were able to interpret a decrease in chamber CO₂ concentrations as due to advective transport processes without having to adjust our data for CO₂ fixation by plants which can vary greatly with small changes in incident irradiance.

BGD

12, 4801–4832, 2015

Quantifying wind and pressure effects on trace gas fluxes

K. R. Redeker et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Three high-volume drum fans (DF24S, Prem-I-Air, Manchester, UK) were placed at one end of the wind tunnel, each capable of moving 235 m³ of air per minute at the highest speed setting (for a maximum calculated wind speed of ~ 10 m s⁻¹).

The toroid and outer tunnel were in place at the respective field sites (see below) for one or two days. Between measurements, which typically took less than 10 min, the toroid was unshrouded (the available sunlight between measurements was similar to that of a regionally cloudy day) and its vents opened. Therefore, the effects of the apparatus on the soil being studied were kept to a minimum; i.e., gas concentrations in the air above the soil were not allowed to build over long time periods which would have affected gas concentrations in the soil and soil biochemical processes.

2.2 Field sites

To investigate wind and pressure effects on air flow into and out of soils, we selected four sites offering a broad range of soil porosities, pore water contents, and organic matter contents. The sites also differed in the processes affecting CO₂ and CH₄ production and consumption.

2.2.1 Wheldrake Forest

Investigations at Wheldrake Forest (53°54'36" N, 0°59'55" W) occurred on 20 April and from 4–6 December 2011. The site was within a lodgepole pine (*Pinus contorta* Douglas) plantation with a small, scattered population of silver birch (*Betula pendula* Roth) with little or no understory. The soil is a well-drained, fine, sandy podzol. CO₂ fluxes from the soil are likely to be dominated by tree roots and heterotrophic respiration (Heinemeyer et al., 2011). In contrast, relatively high rates of net CH₄ uptake have been observed previously within these soils, driven by methanotrophic bacteria (Heinemeyer et al., 2011).

BGD

12, 4801–4832, 2015

Quantifying wind and pressure effects on trace gas fluxes

K. R. Redeker et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.2.2 University of York managed grassland

Investigations of managed grassland on the University of York campus (53°56'50" N, 1°3'26" W) occurred on 21 April and from 18–19 August 2011. The sample site was a tended lawn surface. As grasses are not particularly symbiotic with either arbuscular or ectomycorrhizal fungi we expected limited fungal influence, limiting CO₂ production within the soil to primarily roots or bacterial respiration. Rainfall during the August measurements significantly affected the soil pore water content, and localized pools of standing water were observed on both sampling days, likely limiting further the biogenic production and consumption of trace gases.

2.2.3 Cors Fochno peat bog

Cors Fochno is an estuarine raised bog in west-central Wales (centered on 52°30'14" N, 4°00'47" W). Measurements at the site took place between 13 and 14 September 2011 on a wet "lawn" (sensu Belyea and Clymo, 2001) dominated by the moss *Sphagnum pulchrum* (Lindb. ex Braithw.) Warnst. with a scattering of the sedge *Rhynchospora alba* (L.) Vahl. CH₄ is produced throughout the soil profile at Cors Fochno, including the upper layers (e.g., Green and Baird, 2013; Comas et al., 2013). Respiration was expected to be primarily from surface peats and mosses, with some respiration from the sedge, *R. alba*. The water table across the lawn was at or close to the surface (within 2–3 cm of the top of the *Sphagnum* plants). Sections of wooden boardwalk were placed around the measurement area to minimize compression of the peat (soil) profile by observers.

The measurement period followed the landfall of a significant atmospheric depression. Wind gusts in excess of 50 mph were common on the 11th and 12th, and on the first day of sampling (the 13th) winds were often in excess of 20 mph. Winds had slowed considerably by the 14th to between 3.5 and 7 mph (1.5–3.0 ms⁻¹).

BGD

12, 4801–4832, 2015

Quantifying wind and pressure effects on trace gas fluxes

K. R. Redeker et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.2.4 Forsinard peat bog

Measurements at Forsinard Flows Reserve (58°21'25" N, 3°53'48" W) took place from 13–14 July 2012. The reserve is a low altitude blanket bog in Caithness and Sutherland in northern Scotland. It is protected for its nature conservation interest by the Royal Society for the Protection of Birds (RSPB), and some areas are actively managed having previously been damaged by afforestation. Measurements took place in an unmanaged area of bog containing a mixed assemblage of vascular plants and bryophytes, including *Trichophorum cespitosum* (L.) Hartm., *Erica tetralix* L., *Eriophorum vaginatum* L. and *Sphagnum papillosum* Lindb. (Bellamy et al., 2012). The water table depths at our sampling locations in Forsinard were significantly lower than Cors Fochno (> 10 cm).

2.3 Trace gas flux measurements

Trace gas fluxes from the footprint of the toroid were estimated in the same way as for a conventional flux chamber; i.e. by measuring gas concentrations within the toroid over time and using the rate of change in concentration to calculate a flux (cf. Denmead, 2008). Fluxes were measured across a 3 × 3 matrix of local (≤ 1 m radius; isolated toroidal wind tunnel) and microscale (≥ 1 m radius; straight line wind tunnel) wind speeds, denoted “zero”, “mid” and “high” (Table 1). Replicate measurements were made for each wind state, and the order of tested sample conditions was randomized to avoid conflating temporal effects.

During the experiments, trace gas concentrations in the toroidal wind tunnel were continuously measured using a Los Gatos Research Fast Greenhouse Gas Analyzer (FGGA; Los Gatos Research, Mountain View, CA, USA). The instrument is capable of measuring both methane (CH₄) and carbon dioxide (CO₂) simultaneously. The measurement interval at the forest, the managed grassland, and the Forsinard peat bog sites was every second (1 Hz) while at Cors Fochno peat bog it was every 5 s (0.2 Hz). At these sampling intervals instrumental precision is better than ±0.1 % for both gases. Air from within the toroid was drawn from the East vent lid, 16 cm above the anemome-

BGD

12, 4801–4832, 2015

Quantifying wind and pressure effects on trace gas fluxes

K. R. Redeker et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ters and 38 cm above the soil surface, into the FGGA, where it was analyzed, via off-axis Integrated Cavity Output Spectroscopy in a non-destructive manner (Baer et al., 2002), and returned to the toroid at the West vent lid.

For all but the first set of measurements on a given day (the first sampled combination of wind speeds and pressure), the measurements were only initiated after the straight line wind tunnel and toroid gas concentrations returned to approximately ambient concentrations (1.8–2.0 ppmv for CH₄; 385–400 ppmv for CO₂). These starting conditions were confirmed through continuous FGGA sampling and analysis of the toroidal and straight line wind tunnel concentrations between sampling measurements and occurred rapidly, within 2–3 min. Once the next sampling period was ready to begin, the fans in the toroid and straight-line tunnel were engaged at the appropriate settings; zero, mid or high. The toroid was then isolated from external air masses by placing the toroid's vent lids on silicone gaskets and weighing them down with lead-shot-filled tubing. The toroid remained isolated from exterior air masses, for ~ 6 min during sampling, after which the fans in the toroid and wind tunnel were powered down, the vent lids removed and the system left to re-equilibrate to ambient conditions. At Forsinard, and only Forsinard, 90 min gaps were allowed between each faster wind sampling state, and in these conditions flux measurements at zero-wind speed (both within and without the toroidal wind tunnel) were taken prior to further testing to ensure that fluxes had returned to their original zero-wind range (as described in the Results section). Care was taken at all sites to minimize the amount of pressure placed upon nearby soils prior to and during sampling.

Pressure differential, soil temperature, ambient air temperature and internal wind speeds were not measured within the isolated toroid and straight line wind tunnels during each measurement period. Wind speeds remained steady during each placement (Table 1) but differed significantly between sites. Lab-based wind speeds for the toroid (as determined by maximum measured wind speeds over smooth aluminum sheeting) were ~ 6 ms⁻¹ while estimated maximum wind speeds within the straight-line wind tunnel were predicted to be ~ 10 ms⁻¹. In situ wind speeds were significantly reduced,

BGD

12, 4801–4832, 2015

Quantifying wind and pressure effects on trace gas fluxes

K. R. Redeker et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



due to friction from the ground surface and, in the case of the straight line wind tunnel, alternative wind paths along the tunnel wall.

2.4 Flux estimates:

Trace gas fluxes in the toroid were calculated using:

$$F = \frac{\Delta[G] V}{\Delta t A} \quad (1)$$

where F is the mass flux per unit area ($\text{ML}^{-2}\text{T}^{-1}$), $[G]$ is mass concentration of CH_4 or CO_2 in the toroid (ML^{-3}), t is time (T), V is toroid internal volume (L^3), and A is the soil surface area or footprint of the toroid (L^2).

Data retrieved from the FGGA from each 6 min sampling period was manually analyzed. Up to the first 2 min of data were discarded due to pressure-based fluctuations that masked any linear response from the set wind states. The amplitude and duration of these initial fluctuations were compared to set wind speeds and no correlation was observed, so this data was discarded.

After the initial disturbance in concentration measurements both CO_2 and CH_4 proceeded to increase or decrease in a linear fashion for the duration of the remainder of the experiment (< 6 min). We utilized the earliest 120–180 s period during which both CO_2 and CH_4 gas concentrations either rose or fell in a linear fashion. A linear regression line was fitted to the data from this 120 to 180 s sampling period to estimate $\Delta[G]/\Delta t$ for use in Eq. (1). In nearly all cases the r^2 of the linear regression was greater than 0.9.

2.5 Relative flux calculation

Trace gas fluxes are likely to be significantly different for different trace gas species, both temporally and spatially. To compare trace gas fluxes across different dates and locations the average of measured fluxes from the zero-wind treatments, where wind

BGD

12, 4801–4832, 2015

Quantifying wind and pressure effects on trace gas fluxes

K. R. Redeker et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



speeds in both the toroid and straight-line outer wind tunnels were zero, was taken as a baseline condition, and set to represent a value of 1.0. All other treatments were then compared relative to this value so that the relative flux for each trace gas was equal to:

$$F_R = F_T / F_0 \quad (2)$$

where F_R is the relative flux for each gas under each set of conditions, F_T is the treatment flux and F_0 is the appropriate average baseline flux. Using these relative measures, trace gas fluxes can be compared across space (between and within ecosystems) and time.

3 Results

3.1 Wind speed differences vs. pressure differentials

Our data show that wind speeds were better at predicting trace gas fluxes than pressure differentials (Figs. 2–4). While the physical relationship between pressure and wind is well established, wind speed is not strongly correlated with pressure differences measured between the toroidal and straight line wind tunnels (Fig. 2; $r^2 = 0.63$). Of particular interest to the comparison of wind speed and pressure differential as explanatory variables are measurements taken during the managed grassland measurement campaign where warming within the toroid (from residual thermal energy from the soil surface) led to an increase in pressure within the instrument. The observed differences in pressure (+) were opposite to those expected due to ongoing higher wind speeds within the toroid (-). When pressure is higher within the toroid one might expect air within it to be driven into the soil, reducing gas fluxes from the soil to the atmosphere. However, CH_4 and CO_2 fluxes from soil to atmosphere were substantially higher within these treatments, suggesting that fluxes were more strongly influenced by measurable wind speed than by measured pressure changes (Tables 1–3).

BGD

12, 4801–4832, 2015

Quantifying wind and pressure effects on trace gas fluxes

K. R. Redeker et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3.2 Wind speed effects on trace gas fluxes

Wind speeds internal and external to the toroidal wind tunnel affected CH₄ and CO₂ fluxes in a planar fashion ($r^2 = 0.82$; Fig. 3a–c). CH₄ and CO₂ fluxes are enhanced as wind speeds directly above the soil surface increase (i.e., within the toroid) (+54 % flux relative to zero-wind conditions per ms⁻¹) but are reduced as wind speeds external to the toroid increase (i.e., within the straight line wind tunnel but outside the toroid) (–11 % flux relative to zero-wind conditions per ms⁻¹). Under open field wind conditions, where internal and external wind speeds are similar, trace gas fluxes increase by 42 % per ms⁻¹ wind speed relative to zero-wind conditions (Fig. 3a). Although fluxes increased linearly across the range of wind speeds (and wind speed differentials) considered here (Fig. 3a), it is important to note that they could exhibit a different functional form over a wider range of speeds. For example, trace gas fluxes may approach an asymptote at very high wind speeds.

The relationships identified above are irrespective of the initial flux direction (efflux or influx). When CH₄ is taken up by soils, increased wind speeds in the isolated toroid led to greater CH₄ uptake while faster wind speeds within the straight line wind tunnel reduced CH₄ uptake (Fig. 3c; Table 3). The observed wind speed-trace gas flux correlation was consistent for both gases measured over all ecosystems, and was reproducible both within and between sampling campaigns (Tables 2 and 3; Fig. 3a–c).

3.3 Abrupt flux transitions driven by high wind speeds

Data collected from the Forsinard peat bog site provides compelling evidence of abrupt flux transitions. During this campaign it became clear that, unlike other study sites, it was impossible to obtain replicable results while randomly selecting toroid and wind tunnel wind speed conditions. At this location surface soil pore spaces were purged under short exposure (< 10 min) to “high” wind speed conditions (~ 2.0 ms⁻¹ within the toroid) and required an hour to re-equilibrate to their original zero-wind fluxes (Fig. 5). The evergreen forest experiment showed a similarly abrupt transition in flux (a 30 %

BGD

12, 4801–4832, 2015

Quantifying wind and pressure effects on trace gas fluxes

K. R. Redeker et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



reduction in zero-wind fluxes after a single long term exposure to high winds within both the isolated toroid and the linear wind tunnel).

Increases in fluxes at higher wind speeds, followed by periods of lower fluxes have previously been reported for eddy correlation measurements (Sachs et al., 2008; Wille et al., 2008; Schrier-Uijl et al., 2012). Likewise, internal wind-speed effects on instantaneous chamber fluxes have been documented (e.g. – Denmead, 2008; Xu et al., 2006). These previous studies have allowed these effects to be measured, but mostly as a by-product of trying to reduce or evaluate poorly-constrained errors in measurement methods. Our study is the first to consider both wind and pressure effects simultaneously in a replicated study for realistic ranges of wind speeds and pressure differentials and is the first to quantify the duration of the wind-driven evacuation effect on fluxes. It is possible that high fluxes during high winds followed by low fluxes due to soil evacuation may help to reduce the overall discrepancy between eddy flux and chamber measurements; however, the precise scale of this discrepancy remains unknown since our empirical quantification of the wind driven pressure fluctuation mechanisms affecting trace gas fluxes is still in its early stages.

4 Discussion

4.1 Which is a more effective tool to understand trace gas fluxes, wind speed or pressure differential?

Our results demonstrate that both wind speed and pressure differential are correlated to surface fluxes of trace gases (Figs. 3a–c and 4). Wind speed, however, is consistently a better predictor than pressure differential. For instance, incursions of air masses from soils to the local atmosphere have been observed to be much more important to observed trace gas concentrations than incursions of surface air into the soil (Xu et al., 2006). Furthermore, soil pore spaces buffer, through expansion and contraction of soil pore air, local boundary layer air pressures (Xu et al., 2006).

BGD

12, 4801–4832, 2015

Quantifying wind and pressure effects on trace gas fluxes

K. R. Redeker et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Quantifying wind and pressure effects on trace gas fluxes

K. R. Redeker et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Our observations support the concept of pressure buffering. One of the aspects of the system that is not described explicitly by Xu et al. (2006) is the effect of temperature on the chamber pressure. In our experiments the internal temperature of the isolated toroid was, at times, 10 °C warmer than the air within the linear wind tunnel, due to transfer of residual heat from the soil surface to the enclosed air within the toroid. Using the Ideal Gas law we would expect the pressure differential (straight line wind tunnel minus isolated toroid) under these conditions to be –34 mbar but the observed pressure differential was much less, –0.18 mbar. To place this in context, it would require 80 ms⁻¹ wind speeds to generate the same pressure differential generated by a 10 °C temperature difference.

A further complication to common use of pressure differential measurements is the placement of the pressure gauge. We suggest that an aboveground placement is not particularly helpful, since it does not address the soil-boundary layer buffering previously described. However, sub-surface placements become problematic due to problems associated with standardization of depth and of disturbance. More broadly, the criteria for pressure differential gauge placement have not been standardized, which has significant implications for comparing published results from different studies. Therefore, it may be argued that obtaining data on pressure differentials for the purposes of trace gas flux measurements is not practical. A more tractable, plausible, measureable quantity is local wind speed, although some standardization of measurement heights and locations will be necessary; most published data to date have utilized measurement heights from 0.2 to 5.0 m from the soil surface.

4.2 Wind speed effects on soil–atmosphere exchange of trace gases: a revised conceptualization

Measurements taken under realistic surface wind speeds indicate that gas exchange rates are considerably influenced by both wind speed and spatial distribution of local winds. This implies that the commonly used conceptual model based on simple 1-dimensional diffusion is insufficient, and that a revised model of soil–atmosphere ex-

change is required. In particular we propose to build upon the Massman model (Massman, 2006), developed for soil and snow surfaces, by the inclusion of a near-surface mixed layer.

We propose that boundary layers develop at the near surface in soils, similar to that of plant canopies or the near-surface ocean. This mixed layer develops according to local solar radiation, wind speeds, and wind duration. In the case of soils, the mixed layer develops due to the interplay between abiotic aspects (i) wind speed: we have demonstrated that there exist positive correlations between local wind speed and trace gas fluxes (Fig. 3a), (ii) wind duration: bursts of high winds (maximum-to-zero toroidal wind conditions, Table 1, at one minute intervals) caused $\sim +40\%$ increase in CO_2 flux relative to zero-wind conditions (Table 2; $1.69 \pm 0.32 \mu\text{moles CO}_2 \text{ m}^{-2} \text{ s}^{-1}$) but was less than that of consistent high wind exposure (Table 2; $2.55 \pm 0.63 \mu\text{moles CO}_2 \text{ m}^{-2} \text{ s}^{-1}$), (iii) water table depth: our data suggests that, under fully saturated conditions, the proposed “mixed layer” does not develop (Tables 2 and 3), (iv) soil porosity and biological processes (production, consumption). This new model would explain the observed results through enhanced mixing of soil pore space air with overlying air and the development of horizontal concentration gradients within the soil profile.

Previous soil–atmosphere models cannot explain the full range of soil–atmosphere fluxes that we observed. In the simple diffusive model, CH_4 travels 70% faster than CO_2 , (Sahoo and Mayaa, 2010) which conflicts with our observed, similar response of CH_4 and CO_2 to increased winds over multiple soil types. External wind speed effects are particularly difficult to reconcile with this simple model since diffusion is a relatively slow process while the patterns we observed occurred rapidly (< 2 min).

Pressure differentials, leading to expansion or contraction of air within soil pores, leading to greater and more rapid mixing within pores, have been proposed as a mechanism by which air may be mixed between soil pore spaces and the overlying atmosphere (i.e., “pressure-pumping”) (Denmead, 1979; Yonemura et al., 2000; Takle et al., 2004; Xu et al., 2006; Flechard et al., 2007; Reicosky et al., 2008; Rey et al., 2012). However, the effects of high external winds within the linear wind tunnel on trace gas

BGD

12, 4801–4832, 2015

Quantifying wind and pressure effects on trace gas fluxes

K. R. Redeker et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Quantifying wind and pressure effects on trace gas fluxes

K. R. Redeker et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



fluxes from low or moderate wind environments within the isolated toroid (the isolated toroid in this scenario is similar to the real world scenarios of (i) a forest verge, nearby an open field, (ii) an open field surrounding a slight depression with deeper grass depth providing a protected canopy, or (iii) a hedgerow) cannot be explained through this pressure-pumping model. While pressure waves have been demonstrated to travel up to 50 cm within soils (Takle et al., 2004; Flechard et al., 2007; Reicosky et al., 2008) such waves, under high external wind conditions, would lead to lower relative pressure in the soil below the toroid. A lower pressure below the toroid would lead to a reduction in, or neutral impact on, fluxes of CO₂ (similar concentrations, but lower pressure, in soil pores would mean similar diffusive fluxes, but potential for atmosphere-to-soil transfer to maintain pressure equilibrium). Similarly, lower pressures in the soil would likely lead to greater CH₄ fluxes. Our observed results show neither an increase in methane uptake concurrent with decreases in CO₂ fluxes, nor do they demonstrate an overall neutral impact.

Furthermore, the correlation between pressure differential and flux is significantly weaker than the correlation observed for wind speeds (Fig. 4; $r^2 = 0.41$ for pressure differential vs. Fig. 3a; $r^2 = 0.82$ for wind speed). If neither the diffusion gradient model nor the pressure-pumping model is capable of explaining the available data then a revised model is needed.

Our proposed “mixed layer” conceptualization of the soil–atmosphere interface is described below. In the zero-wind condition (where there is no wind inside either the linear external wind tunnel or the isolated toroidal wind tunnel, and representative of long term, no-wind conditions on either side of a natural boundary), soil concentration gradients are identical on either side of the boundary and fulfill the smooth gradient expectations of the current 1-dimensional gradient diffusive soil model.

Alternately, when the nested wind tunnel is set so that faster winds are experienced within the toroid than in the linear wind tunnel (similar to an open soil surface nearby a rock-covered surface, or an open field near a forest verge), we hypothesize that a mixed layer develops in local soils under high surface winds (directly under the

Quantifying wind and pressure effects on trace gas fluxes

K. R. Redeker et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



toroidal wind tunnel) while soils, external to the isolated toroid and under zero surface winds, retain their diffusion-controlled soil gradient. In this scenario the developing mixed layer either “mines” the soil of high concentration gases or delivers higher concentration, atmospheric gases to consumption zones, leading to enhanced soil–atmosphere fluxes regardless of whether consumption or production processes dominate.

Under the opposite condition, where local surface winds are negligible and microscale surface winds are high (set within the NWT so that there are zero-, or low winds within the toroid and faster winds within the straight line wind tunnel) the mixed layer develops away from the site of interest (in this case, below the toroid) creating a horizontal concentration gradient within surface soils which competes with the vertical concentration gradient at the soil surface, lowering observed fluxes relative to zero-wind conditions.

When fast winds are experienced across an ecosystem equally (as in the case where both linear and toroidal wind tunnels are exposed to fast winds) fluxes are enhanced over zero-wind conditions despite the development of competitive horizontal gradients.

We found two conditions under which the observed relationship between wind speeds and trace gas fluxes break down, neither of which conflict with our proposed hypothesis that surface wind speeds affect the rate of greenhouse gas exchange between soils and the atmosphere through the development of a mixed layer. The first occurs when there is little or no concentration gradient between the atmosphere and the soil profile, leading to zero-wind-state fluxes that are essentially zero. In this situation the development of a mixed layer under elevated wind speeds merely mixes equivalent concentration fluxes between the soil and atmosphere, leading to zero net transfer. This condition was observed for CH₄ fluxes at the managed grassland and Forsinard peat bog sites. The second condition occurs when the water table is very close to the soil surface (< 2 cm), as occurred at Cors Fochno peat bog, where both CO₂ and CH₄ fluxes were affected (Tables 2 and 3). In this situation, it is likely that a mixed layer is unable to develop rapidly due to a combination of water acting as a diffusive barrier

within near-surface soils as well as increased hydrostatic pressure from the overlying water column.

The mixed layer model explicitly allows the disruption of smooth concentration gradients under moderate surface wind conditions and is better able to describe abrupt flux transitions over short timescales (Fig. 5).

This new mixed-layer model suggests that estimates of soil–atmosphere fluxes should be revisited, given that overall fluxes represent the net balance of multiple small, local fluxes. Indeed, all spatially distributed wind conditions described above exist in all ecosystems and each state contributes to the overall ecosystem flux. The concept of flux measurements using traditional techniques as accurate portrayals of soil–atmosphere exchange becomes, in this model, more relativistic.

A mixed layer in surface soils changes our understanding of gross budgets for many trace gases. For instance, up to 90 % of CH₄ generated within soils is consumed in situ (Segers, 1998). The mixed layer model implies that a significantly greater fraction of microbially-produced CH₄ will avoid in situ consumption through rapid mixing with overlying air under windy conditions. This effectively increases soil–atmosphere flux of CH₄ relative to no wind conditions, even if production rates are equal within the soil column.

Acknowledgements. We would like to acknowledge the manufacturers of the inner flux chamber, Mark Bentley and Steve Howarth from the University of York, Dept. of Biology Mechanical and Electronics workshops, respectively. Furthermore, we would like to acknowledge the Forestry Commission for access and aid at Wheldrake Forest, Mike Bailey and Natural Resources Wales for access and assistance at Cors Fochno, and Norrie Russell and the Royal Society for the Protection of Birds for access and aid at Forsinard. We would also like to thank Graham Hambley, James Robinson and Elizabeth Donkin for equipment preparation and sampling. Phil Ineson is thanked for the loan of essential equipment, site suggestions and accessible power supply. Funding was provided by the University of York, Dept. of Biology and by a grant to YAT by the UK Natural Environment Research Council (NE/H01182X/1).

BGD

12, 4801–4832, 2015

Quantifying wind and pressure effects on trace gas fluxes

K. R. Redeker et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



References

- Baer, D. S., Paul, J. B., Gupta, M., and O'Keefe, A.: Sensitive absorption measurements in the near-infrared region using off-axis integrated cavity-output spectroscopy, *Appl. Phys. B Lasers O.*, 75, 261–265, doi:10.1007/s00340-002-0971-z, 2002.
- 5 Bellamy, P. E., Stephen, L., Maclean, I. S., and Grant, M. C.: Response of blanket bog vegetation to drain-blocking, *Appl. Veg. Sci.*, 15, 129–135, 2012.
- Belyea, L. R. and Clymo, R. S.: Feedback control of the rate of peat formation, *P. R. Soc. B*, 268, 1315–1321, 2001.
- 10 Chapman, S. J., Bell, J., Donnelly, D., and Lilly, A.: Carbon stocks in Scottish peatlands, *Soil Use Manage.*, 25, 105–112, 2009.
- Christiansen, J. S., Korhonen, J. F. J., Juszczak, R., Giebels, M., and Pihlatie, M.: Assessing the effects of chamber placement, manual sampling and headspace mixing on CH₄ fluxes in a laboratory experiment, *Plant Soil*, 343, 171–185, doi:10.1007/s11104-010-0701-y, 2011.
- 15 Comas, X., Kettridge, N., Binley, A., Slater, Parsekian, L., A., Baird, A. J., Strack, M., and Waddington, J. M.: The effect of peat structure on the spatial distribution of biogenic gases within bogs, *Hydrol. Process.*, 28, 5483–5494, doi:10.1002/hyp.10056, 2013.
- Davidson, E. A., Savage, K., Verchot, L. V., and Navarro, R.: Minimizing artifacts and biases in chamber-based measurements of soil respiration, *Agr. Forest Meteorol.*, 113, 21–37, 2002.
- 20 Denmead, O. T.: Chamber systems for measuring nitrous oxide emission from soils in the field, *Soil Sci. Soc. Am. J.*, 43, 89–95, 1979.
- Denmead, O. T.: Approaches to measuring fluxes of methane and nitrous oxide between landscapes and the atmosphere, *Plant Soil*, 309, 5–24, doi:10.1007/s11104-008-9599-z, 2008.
- Denmead, O. T. and Reicosky, D. C.: Tillage-induced gas fluxes: comparison of meteorological and large chamber techniques, in: *Proceedings of the International Soil Tillage Research Organizations 2003 Conference*, Brisbane, Australia, 13–18 July 2003, 2003.
- 25 Flechard, C. R., Neftel, A., Jocher, M., Ammann, C., Leifeld, J., and Fuhrer, J.: Temporal changes in soil pore space CO₂ concentration and storage under permanent grassland, *Agr. Forest Meteorol.*, 142, 66–84, 2007.
- Green, S. M. and Baird, A. J.: The importance of episodic ebullition methane losses from three peatland microhabitats: a controlled-environment study, *Eur. J. Soil Sci.*, 64, 27–36, doi:10.1111/ejss.12015, 2013.
- 30 Hillel, D.: *Environmental Soil Physics*, Academic Press, San Diego, London, 800 pp., 1998.

Quantifying wind and pressure effects on trace gas fluxes

K. R. Redeker et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Quantifying wind and pressure effects on trace gas fluxes

K. R. Redeker et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Heinemeyer, A., Di Bene, C., Lloyd, A. R., Tortorella, D., Baxter, R., Huntley, B., Gelsomino, A., and Ineson, P.: Soil respiration: implications of the plant-soil continuum and respiration chamber collar-insertion depth on measurement and modeling of soil CO₂ efflux rates in three ecosystems, *Eur. J. Soil Sci.*, 62, 82–94, 2011.

5 Holmen, K. and Jaffe, D. A.: In *Earth System Science*, edited by: Butcher, S. S., Charlson, R. J., Rodhe, H., and Orians, G.H., Elsevier, London, chaps. 11–12, 282–340, 2000.

Lindsay, R. A., Charman, D. J., Everingham, F., O'Reilly, R. M., Palmer, M. A., Rowell, T. A., and Stroud, D. A.: *The Flow Country: the Peatlands of Caithness and Sutherland*, Interpretive Services Branch, Nature Conservancy Council, Peterborough, UK, 1988.

10 Massman, W. J. and Lee, X.: Eddy covariance flux corrections and uncertainties in long-term studies of carbon and energy exchanges, *Agr. Forest Meteorol.*, 113, 121–144, 2002.

Monson, R. and Baldocchi, D.: *Terrestrial Biosphere – Atmosphere Fluxes*, Cambridge University Press, Cambridge, 2014.

15 Montzka, S. A., Reimann, S. et al.: Ozone-Depleting Substances (ODSs) and related chemicals, in: *2010 WMO Ozone Assessment*, World Meteorological Organization, Geneva, 2010, chap. 1–108, 2010.

Myklebust, M. C., Hipps, L. E., and Ryel, R. J.: Comparison of eddy covariance, chamber and gradient methods of measuring soil CO₂ in an annual semi-arid grass, *Bromus tectorum*, *Agr. Forest Meteorol.*, 148, 1894–1907, 2008.

20 Qi, J., Marshall, J. D., and Mattson, K. G.: High soil carbon dioxide concentration inhibit root respiration of Douglas fir, *New Phytol.*, 128, 435–442, 1994.

Reicosky, D. C., Gesch, R. W., Wagner, S. W., Gilbert, R. A., Wentle, C. D., and Morris, D. R.: Tillage and wind effects on soil CO₂ concentrations in muck soils, *Soil Till. Res.*, 99, 221–231, 2008.

25 Rey, A., Belelli-Marchesini, L., Were, A., Serrano-Ortiz, P., Etiope, G., Papale, D., Domingo, F., and Pegoraro, E.: Wind as a main driver of net ecosystem carbon balance of a semiarid Mediterranean steppe in the South-East of Spain, *Glob. Change Biol.*, 18, 539–554, 2012.

Rochette, P.: Towards a standard non-steady-state chamber methodology for measuring soil N₂O emissions, *Anim. Feed Sci. Tech.*, 166, 141–146, doi:10.1016/j.anifeedsci.2011.04.063, 2011.

30 Rolston, D. E. and Moldrup, P.: Gas transport in soils, in: *Handbook of Soil Science and Processes*, Second Edition, edited by: Huang, P. M., Li, Y., and Sumner, M. E., CRC Press, Boca Raton, FL, 1422 pp., 8-1–8-20, 2012.

Quantifying wind and pressure effects on trace gas fluxes

K. R. Redeker et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Sachs, T., Wille, C., Boike, J., and Kutzbach, L.: Environmental controls on ecosystem-scale CH₄ emission from polygonal tundra in the Lena River Delta, Siberia *J. Geophys. Res.*, 113, G00A03, doi:10.1029/2007JG00505, 2008.

Sahoo, B. K. and Mayaa, Y. S.: Two dimensional diffusion theory of trace gas build up in soil chambers for flux measurements, *Agr. Forest Meteorol.*, 150, 1211–1224, 2010.

Scanlon, B. R., Nicot, J. P., and Massmann, J. W.: Soil gas movement in unsaturated systems, in: *Handbook of Soil Science*, edited by: Sumner, M. E., CRC Press, Boca Raton, 2148 pp., A277–A319, 2000.

Schrier-Uijl, A. P., Kroon, P. S., Hensen, A., Leffelaar, P. A., Berendse, F., and Veenendaal, E. M.: Comparison of chamber and eddy covariance-based CO₂ and CH₄ emission estimates in a heterogeneous grass ecosystem on peat, *Agr. Forest Meteorol.*, 150, 825–831, 2012.

Segers, R.: Methane production and methane consumption: a review of processes underlying wetland methane fluxes, *Biogeochemistry*, 41, 23–51, 1998.

Stamp, I.: Methane Emissions Variability From a Welsh Patterned Raised Bog, Ph.D. thesis, University of London, 2011.

Takle, E. S., Massman, W. J., Brandle, J. R., Schmidt, R. A., Zhou, X., Litvina, I. V., Garcia, R., Doyle, G., and Rice, C. W.: Influence of high frequency ambient pressure pumping on carbon dioxide efflux from soils, *Agr. Forest Meteorol.*, 124, 193–206, 2004.

Wille, C., Kutzbach, L., Sachs, T., Wagner, D., and Pfeiffer, E. M.: Methane emission from Siberian arctic polygonal tundra: eddy covariance measurements and modeling, *Glob. Change Biol.*, 14, 1395–1408, 2008.

Wuebbles, D. J. and Hayhoe, K.: Atmospheric methane and global change, *Earth-Sci. Rev.*, 57, 177–210, 2002.

Xu, L., Furtaw, M. D., Madsen, R. A., Garcia, R. L., Anderson, D. J., and McDermitt, D. K.: On maintaining pressure equilibrium between a soil CO₂ flux chamber and the ambient air, *J. Geophys. Res.*, 111, D08S10, doi:10.1029/2005JD006435, 2006.

Yonemura, S., Miyata, A., and Yokozawa, M.: Concentrations of carbon monoxide and methane at two heights above a grass field and their deposition onto the field, *Atmos. Environ.*, 34, 5007–5014, 2000.

Quantifying wind and pressure effects on trace gas fluxes

K. R. Redeker et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 1. Average chamber wind speed and pressure differential for various inner toroid – outer wind tunnel treatments. Italicized, top values are for data collected in April 2011 (Grassland and Forest) and July 2012 (Peat Bog) while non-italicized bottom values indicate data collected in December, August and September 2011 for Forest, Grassland and Peat Bog, respectively. Upper values are wind speeds (inner toroid; outer wind tunnel) and are listed in m s^{-1} . No SD are listed since wind speeds were consistent to $\pm 0.1 \text{ m s}^{-1}$ at each emplacement. Pressure differential (defined as outer wind tunnel pressure minus inner toroid pressure) is listed below wind speeds, and is shown in mbar.

| Isolated Toroid Wind Speed | Linear Wind Tunnel Wind Speed | | | | | | | | |
|----------------------------------|--|-------------------|-------------------|--|-------------------|-------------------|--|-----------------|--|
| | Forest Soils | | | Managed Grassland Soils | | | Peat Bog | | |
| | Zero | Mid | High | Zero | Mid | High | Zero | Mid | High |
| Zero | <i>0; 0</i> <i>-0.074</i> 0; 0.2 0.025 | 0; 2.4 -0.010 | 0; 3.2 0.001 | <i>0; 0</i> <i>-0.183</i> 0; 0 0.072 | 0; 2.0 0.038 | 0; 3.2 0.012 | 0; 0 <i>0.009</i> 0; 0 0.003 | | <i>0; 2.0</i> <i>0.003</i> |
| Mid | | | | | | | <i>1.1; 0</i> <i>0.054</i> 0.9; 0.2 0.070 | 0.8; 0 0.016 | |
| High | <i>0; 2.5</i> <i>0.160</i> 1.7; 0.2 0.187 | 1.7; 2.4 0.179 | 1.7; 3.2 0.171 | <i>0; 2.6</i> <i>0.169</i> 3.2; 0 0.151 | 3.2; 1.9 0.150 | 3.2; 3.6 0.215 | <i>2.0; 0</i> <i>0.132</i> 1.4; 0 0.049 | | <i>2.0; 1.4</i> <i>0.090</i> 1.4; 2.5 0.056 |

Quantifying wind and pressure effects on trace gas fluxes

K. R. Redeker et al.

Table 2. Average carbon dioxide fluxes (in $\mu\text{moles CO}_2 \text{ m}^{-2} \text{ s}^{-1}$) \pm SD for each component of the inner toroid-outer wind tunnel matrix. Italicized, top values are for data collected in April 2011 (Grassland and Forest) and July 2012 (Peat Bog) while non-italicized bottom values indicate data collected in December, August and September 2011 for Forest, Grassland and Peat Bog, respectively. By convention positive values indicate efflux of CO_2 from the soil surface into the atmosphere.

| Isolated Toroid Wind Speed | Linear Wind Tunnel Wind Speed | | | | | | | | |
|----------------------------------|-------------------------------|------------------------|------------------------|-------------------------------|------------------------|------------------------|-------------------------------|-----|--------------------------|
| | Forest Soils | | | Managed Grassland Soils | | | Peat Bog | | |
| | Zero | Mid | High | Zero | Mid | High | Zero | Mid | High |
| Zero | <i>0.35 ± 0.10</i> (n = 3) | | | <i>2.21 ± 0.36</i> (n = 3) | | | <i>1.09 ± 0.27</i> (n = 3) | | <i>1.18</i> (n = 1) |
| | 1.20 ± 0.33 (n = 6) | 0.81 ± 0.42 (n = 6) | 0.67 ± 0.31 (n = 6) | 2.19 ± 1.17 (n = 3) | 1.50 ± 0.67 (n = 3) | 1.35 ± 0.59 (n = 3) | 0.397 ± 0.160 (n = 3) | | |
| Mid | | | | | | | <i>1.39 ± 0.40</i> (n = 2) | | |
| | 2.13 ± 0.26 (n = 5) | 1.65 ± 0.33 (n = 6) | 1.43 ± 0.51 (n = 6) | 4.19 ± 1.24 (n = 3) | 3.90 ± 0.26 (n = 3) | 3.98 ± 0.75 (n = 3) | 0.392 ± 0.102 (n = 5) | | |
| High | <i>1.02 ± 0.51</i> (n = 4) | | | <i>7.35 ± 0.16</i> (n = 3) | | | <i>2.80 ± 0.60</i> (n = 4) | | <i>1.54</i> (n = 1) |
| | 2.55 ± 0.63 (n = 6) | 1.46 ± 0.61 (n = 6) | 1.48 ± 0.65 (n = 6) | 5.66 ± 0.26 (n = 3) | 5.29 ± 0.40 (n = 3) | 5.63 ± 0.38 (n = 3) | 0.401 ± 0.034 (n = 4) | | 0.357 ± 0.015 (n = 3) |

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Quantifying wind and pressure effects on trace gas fluxes

K. R. Redeker et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Table 3. Average methane fluxes (in $\mu\text{g CH}_4 \text{ m}^{-2} \text{ h}^{-1}$) \pm SD for each component of the inner toroid – outer wind tunnel matrix. Italicized, top values are for data collected in April 2011 (Grassland and Forest) and July 2012 (Peat Bog) while non-italicized bottom values indicate data collected in December, August and September 2011 for Forest, Grassland and Peat Bog, respectively. By convention positive values indicate efflux of CH_4 from the soil surface into the atmosphere.

| Isolated Toroid Wind Speed | Linear Wind Tunnel Wind Speed | | | | | | | | |
|----------------------------|-------------------------------|-----------------------|-----------------------|--------------------------------|------------------------|------------------------|--------------------------------|-----|-------------------------|
| | Forest Soils | | | Managed Grassland Soils | | | Peat Bog | | |
| | Zero | Mid | High | Zero | Mid | High | Low | Mid | High |
| Zero | <i>-103 ± 22</i> (n = 3) | | | <i>-29.8 ± 23.0</i> (n = 3) | | | <i>-19.8 ± 18.7</i> (n = 3) | | <i>-33.0</i> (n = 1) |
| | -402 ± 183 (n = 6) | -226 ± 150 (n = 5) | -271 ± 62 (n = 6) | -6.0 ± 8.1 (n = 3) | -16.8 ± 7.3 (n = 3) | -24.6 ± 6.9 (n = 3) | 3080 ± 860 (n = 3) | | |
| Mid | | | | | | | <i>8.3 ± 19.5</i> (n = 2) | | |
| | -498 ± 186 (n = 6) | -404 ± 179 (n = 6) | -381 ± 71 (n = 6) | -17.1 ± 5.0 (n = 3) | -18.7 ± 9.4 (n = 3) | 2.6 ± 6.2 (n = 3) | 2910 ± 530 (n = 5) | | |
| High | <i>-256 ± 110</i> (n = 3) | | | <i>-53.0 ± 9.8</i> (n = 3) | | | <i>30.3 ± 41.6</i> (n = 4) | | <i>4.40</i> (n = 1) |
| | -624 ± 350 (n = 6) | -616 ± 300 (n = 6) | -514 ± 260 (n = 6) | -9.0 ± 1.6 (n = 3) | -19.0 ± 4.9 (n = 3) | -4.8 ± 18.1 (n = 3) | 3050 ± 60 (n = 4) | | 3040 ± 60 (n = 3) |

BGD

12, 4801–4832, 2015

Quantifying wind and pressure effects on trace gas fluxes

K. R. Redeker et al.



Figure 1. The nested wind tunnel system. Note high speed fans within the toroid at East and West compass points, with anemometers measuring wind speeds at points North and South. Toroid vents are open at this point and all internal fans are off. Wind tunnel sides are PAR transparent in this picture and drum fans at the end of the agricultural tunnel are off. Pressure differential gauges can be seen above fan banks and the PAR sensor is front and center on the top of the flux chamber.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Figure 2. The relationship between wind speed and measured pressure differential (outer wind tunnel minus inner toroid). The planar surface indicates best fit to data ($r^2 = 0.63$, $z = -0.03 + 0.07 \times \text{IT WS (m s}^{-1}) + 0.01 \times \text{OWT WS (m s}^{-1})$). Note insensitivity of pressure to external wind speeds.

Quantifying wind and pressure effects on trace gas fluxes

K. R. Redeker et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Quantifying wind and pressure effects on trace gas fluxes

K. R. Redeker et al.




Figure 3. (a) Relationship between wind speeds internal and external to the toroid and relative fluxes for the pooled CO₂ ($n = 27$) and CH₄ ($n = 13$). Planar surface indicates best fit to data ($z = 0.98 + 0.54 \times \text{IT WS} - 0.11 \times \text{OWT WS}$; $r^2 = 0.82$). The shaded red line represents equal wind speeds inside and outside the flux chamber, where $z = 0.098 + 0.42 \times \text{IT/OWT WS}$ (m s^{-1}). (b) and (c) show disaggregated CO₂ and CH₄ data, with (b) showing the relationship between wind speeds internal and external to the toroid and relative CO₂ fluxes ($n = 27$). All details as shown in Fig. 3a. Planar surface indicates best fit to data ($z = 1.021 + 0.57 \times \text{IT WS} - 0.13 \times \text{OWT WS}$; $r^2 = 0.85$). For (c) we show the relationship between wind speeds internal and external to the toroid and relative CH₄ fluxes ($n = 13$). All details as shown in Fig. 3a. Planar surface indicates best fit to data ($z = 0.94 + 0.43 \times \text{Internal WS} - 0.09 \times \text{External WS}$; $r^2 = 0.80$).

| | |
|--------------------------|--------------|
| Title Page | |
| Abstract | Introduction |
| Conclusions | References |
| Tables | Figures |
| ◀ | ▶ |
| ◀ | ▶ |
| Back | Close |
| Full Screen / Esc | |
| Printer-friendly Version | |
| Interactive Discussion | |



Quantifying wind and pressure effects on trace gas fluxes

K. R. Redeker et al.




Figure 4. The relationship between measured pressure differential (outer wind tunnel minus inner toroid, in mbar) and flux relative to zero wind conditions (F_R). For direct comparison, only data included in Fig. 3 have been included in this figure. Solid fill symbols indicate CO₂ flux ratios while open symbols show CH₄ flux ratios (Tables 2, 3). Trend line indicates best fit for data ($r^2 = 0.41$, $y = 4.48 \times \text{Pressure differential (mbar)} + 1.15$).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Figure 5. Flux recovery from peat soils after high wind events. Open circle (o) represents fluxes taken under zero wind conditions prior to wind events. Gray closed squares represent fluxes measured under zero wind conditions after wind events. The logarithmic trend line indicates best fit to data ($y = 0.195 \times \ln(t(\text{min})) + 0.233$; $r^2 = 0.95$). Error bars indicate SD; $n \geq 3$.

Quantifying wind and pressure effects on trace gas fluxes

K. R. Redeker et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

