

This discussion paper is/has been under review for the journal Biogeosciences (BG).
Please refer to the corresponding final paper in BG if available.

Optical sampling of the flux tower footprint

J. A. Gamon

Department of Earth and Atmospheric Sciences, and Department of Biological Sciences,
University of Alberta, Edmonton, Alberta, Canada

Received: 27 January 2015 – Accepted: 17 February 2015 – Published: 30 March 2015

Correspondence to: J. A. Gamon (jgamon@gmail.com)

Published by Copernicus Publications on behalf of the European Geosciences Union.

BGD

12, 4973–5014, 2015

Optical sampling of the flux tower footprint

J. A. Gamon

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Abstract

The purpose of this review is to address the reasons and methods for conducting optical remote sensing within the flux tower footprint. Fundamental principles and conclusions gleaned from over two decades of proximal remote sensing at flux tower sites are reviewed. An organizing framework is the light-use efficiency (LUE) model, both because it is widely used, and because it provides a useful theoretical construct for integrating optical remote sensing with flux measurements. Multiple ways of driving this model, ranging from meteorological measurements to remote sensing, have emerged in recent years, making it a convenient conceptual framework for comparative experimental studies. New interpretations of established optical sampling methods, including the Photochemical Reflectance Index (PRI) and Solar-Induced Fluorescence (SIF), are discussed within the context of the LUE model. Multi-scale analysis across temporal and spatial axes is a central theme, because such scaling can provide links between ecophysiological mechanisms detectable at the level of individual organisms and broad patterns emerging at larger scales, enabling evaluation of emergent properties and extrapolation to the flux footprint and beyond. Proper analysis of sampling scale requires an awareness of sampling *context* that is often essential to the proper interpretation of optical signals. Additionally, the concept of optical types, vegetation exhibiting contrasting optical behavior in time and space, is explored as a way to frame our understanding of the controls on surface–atmosphere fluxes. Complementary NDVI and PRI patterns across ecosystems are offered as an example of this hypothesis, with the LUE model and light-response curve providing an integrating framework. We conclude that experimental approaches allowing systematic exploration of plant optical behavior in the context of the flux tower network provides a unique way to improve our understanding of environmental constraints and ecophysiological function. In addition to an enhanced mechanistic understanding of ecosystem processes, this integration of remote sensing with flux measurements offers many rich opportunities for upscaling, satellite valida-

Optical sampling of the flux tower footprint

J. A. Gamon

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



tion, and informing practical management objectives ranging from assessing ecosystem health and productivity to quantifying biospheric carbon sequestration.

1 Introduction

The global flux tower network has greatly expanded our understanding of the exchange of mass and energy between terrestrial ecosystems and the atmosphere (Baldocchi et al., 2001). Emerging from a long history of research in the micrometeorological community, eddy covariance has increasingly been integrated with ecophysiological perspectives to become an essential tool for understanding terrestrial ecosystem photosynthesis and productivity. Examples of this integration include eddy covariance studies that apply the concept of light response curves to the analysis of whole ecosystem fluxes (Wofsy et al., 1993; Goulden et al., 1997). Originally derived from the ecophysiological literature using steady-state gas exchange (Björkman, 1981), photosynthetic response curves enable powerful functional analyses of the physiological controls on gas exchange (VonCaemmerer and Farquhar, 1981; Long and Bernacchi, 2003). When applied to surface–atmosphere fluxes, response curves can reveal insights into the environmental and biological controls on these fluxes. They can also be used for partitioning net flux into gross (photosynthetic) and respiratory fluxes (Reichstein et al., 2012). The light-response-curve also provides a powerful theoretical framework for integrating optical and flux data within the framework of the Light-Use Efficiency (LUE) model, allowing a deeper, physiological examination of ecosystem function than possible from monitoring fluxes or optical signals alone.

Like eddy covariance, remote sensing emerged from a physical sciences background, and its history has often focused on mapping and monitoring broad vegetation types or a few key biophysical parameters (e.g. LAI, biomass, or fPAR) rather than providing a functional understanding of underlying ecosystem physiological responses. In part, this limitation derives from the relatively coarse scale of most global earth observation. Observations of large-scale phenology through broad spatial and temporal

BGD

12, 4973–5014, 2015

Optical sampling of the flux tower footprint

J. A. Gamon

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Optical sampling of the flux tower footprint

J. A. Gamon

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



patterns of greenness (NDVI, Tucker et al., 1985), and redefinition of terrestrial biomes through their spatially and temporally dynamic NDVI properties (Defries and Townshend, 1994) are classic studies that illustrate the power of remote sensing to map vegetation patterns over continental regions and yearly time scales. The development of the MODIS NPP products (Running et al., 2004) have moved beyond depictions of spatial and temporal patterns of vegetation to a more functional description of vegetation dynamics and productivity, but still at relatively coarse spatial and temporal scales that can be difficult to directly relate to flux tower sites. Biome-scale comparisons of aggregated optical properties to yearly-integrated net primary production (NPP, Goward et al., 1985) and gross primary production (GPP, Frankenberg et al., 2011) demonstrate the parallel abilities of optical remote sensing and flux measurements to characterize broad-scale patterns of ecosystem–atmosphere exchange and ecosystem production, even if the mechanistic details operating at finer temporal and spatial scales remain obscure. Our current challenge is to fill in those details, and proximal remote sensing within the flux tower footprint provides an important means to accomplish this.

In the past, remote sensing and flux measurements have often been seen as fundamentally different disciplines, and the integration of these methods to address ecosystem–atmosphere fluxes is a relatively new phenomenon. An important stimulus of this transition to an integrated approach has been NASA’s field campaigns, particularly FIFE (Sellers et al., 1992) and BOREAS (Sellers et al., 1997). These large studies began to integrate eddy covariance and remote sensing in a systematic way, and exposed entire generations of scientists to the kind of interdisciplinary thinking needed for ecosystem physiology and global ecology. Since that time, the advent of new tools – principally improved field spectrometers and sampling platforms – have enabled more detailed optical studies of the flux tower footprint. The emergence of optical sampling networks such as SpecNet (Gamon et al., 2006b, 2010), EuroSpec (this issue) and the Phenocam network (Richardson, 2009) have drawn attention to the need for more systematic integration of proximal remote sensing within the flux tower network.

Optical sampling of the flux tower footprint

J. A. Gamon

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



In this review, I will argue that direct integration of remote sensing with flux measurements within the context of the global flux tower network improves our understanding of both fluxes and remote sensing. This integration offers greater insights into the underlying controls on biospheric-atmospheric fluxes than possible with either method alone.

While optical sampling is readily scaleable, a key focus will be on proximal remote sensing explicitly designed to match the sampling scale of flux measurements as a means of understanding functional processes. Proximal sampling can also be readily linked to conventional remote sensing, providing critical validation (e.g. “ground truthing”) and extrapolation of fine-scale measurements to larger regions (“upscaling”). While optical methods will be the primary focus, other methods (e.g. LIDAR, thermal, microwave, or acoustic) can also be applied to a wide range of ecological questions within the flux footprint. A primary conceptual framework will be the LUE model, in part due to the rich history of this approach, but also because potential parameterizations of this model remain largely unexplored, providing rich opportunities for new advances. When properly applied, proximal remote sensing offers many prospects for a deeper, physiological examination of whole ecosystem function, and can be seen as an emerging ecophysiological frontier.

2 Sampling context

A well-designed proximal remote sensing campaign must consider sampling *context*, including the scale of biological responses in temporal and spatial dimensions. Attention to the temporal and spatial context is essential because optical and flux instrumentation, and the biological processes they detect, operate with different time constants over different spatial scales (Figs. 1 and 2). Context also includes the angular response of illumination and the measured signals. Attention to the spectral (wavelength) domain is also needed, as optical instruments and the biological phenomena they measure can be very particular when it comes to spectral responses.

BGD

12, 4973–5014, 2015

Optical sampling of
the flux tower
footprint

J. A. Gamon

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



We can illustrate the spectral-spatial-temporal domain defining the realm of optical sampling using ecosystem photosynthetic carbon uptake and its underlying biological controls as an example. Plants possess a myriad of ways to adjust the receipt and distribution of light energy, all of which can be detected optically (Fig. 1). These photosynthetic regulatory processes operate over a wide range of scales. At fine spatial and temporal scales, leaves adjust light energy distribution via energy dissipation mechanisms, including chlorophyll fluorescence (red text in Fig. 1), which responds to illumination in milliseconds and relaxes to steady-state levels over seconds to minutes (Jones, 2013). At slightly longer time scales (seconds to hours), non-radiative energy dissipation via the xanthophyll cycle (Demmig et al., 1987) provides further means of adjusting energy distribution (left side of Fig. 1), and these changes can be monitored via fluorescence and spectral reflectance (Gamon et al., 1990, 1997; Peñuelas et al., 1995). Critical biochemical aspects of photosynthetic regulation (e.g. Rubisco activation) occur over similarly short time scales, and this biochemical capacity, illustrated as carboxylation capacity (V_{cmax}) and electron transport capacity (J_{max}) in Fig. 1, can be assessed with optical methods (Serbin et al., 2011). In parallel to these photochemical and biochemical regulatory processes, leaf structure, ultrastructure, and pigmentation adjust over minutes (in the case of chloroplast or rapid leaf movement) to days or weeks (in the case of pigment pool sizes and leaf area index) (Björkman and Demmig-Adams, 1994). These adjustments affect leaf light absorption (blue text in Fig. 1), and can be detected via spectral transmittance (Brugnoli and Björkman, 1992) or reflectance (Zygielbaum et al., 2012; Wong and Gamon, 2015a).

At the scale of a flux footprint, vegetation's capacity to absorb light determines its potential photosynthetic rate and net primary productivity. This capacity varies with canopy and stand structure, often expressed as green leaf area index, leaf angle distribution, or percent cover, which can be measured as the fraction of PAR irradiance (PPFD) absorbed (F_{APAR}) using a number of vegetation reflectance indices, most notably the Normalized Difference Vegetation Index, NDVI (Gamon et al., 1995; Running et al., 2004).

can be functionally categorized according to its LUE response. If the slope is variable, then we must reconsider this assumption, as further discussed below.

The light-response curve provides a potent framework for analyzing physiological responses, and for deciding how to best integrate optical and flux measurements. In this analysis, because APAR reflects the amount of light absorbed by green canopy structure, it provides key information on *structural* control on fluxes associated with such factors as green canopy cover and leaf-area index, and depicts the maximum potential photosynthesis determined by light absorption. By contrast, the slope (efficiency) represents how much of that photosynthetic potential is actually realized due to variations in environmental conditions and physiological performance. In this way, a changing slope can provide additional information on the degree of *physiological* control associated with photosynthetic adjustment and downregulation under varying environmental conditions. This distinction between structural and physiological controls is somewhat artificial; in nature, structural and physiological factors interact in interesting and complex ways. Nevertheless, it provides a useful conceptual framework for analyzing ecosystem behavior and determining how best to parameterize the LUE model.

A strength of the LUE model is that APAR can be readily assessed from optical remote sensing. A bigger challenge than APAR assessment lies in determining the efficiency (ε) term in the LUE model, in part because it can vary with vegetation type and environmental conditions (Garbulsky et al., 2010). In some cases, for a given vegetation type and time frame, ε can be treated as a constant, greatly simplifying the LUE model (Monteith, 1977; Ruimy et al., 1994). This view is analogous to the observation that all unstressed leaves of a particular type (C_3 or C_4) tend towards a common photon yield (LUE expressed at the leaf level), representing optimal conversion efficiency (Björkman, 1981). However, Nature often grossly violates the assumption of a constant LUE, and recent studies have emphasized the large temporal variation in ε , particularly during conditions of chronic stress (Garbulsky et al., 2010). In these cases, environmental factors including temperature, moisture, and nutrients cause the efficiency term to vary on a variety of time scales, leading to large errors in LUE model estimates that assume

BGD

12, 4973–5014, 2015

Optical sampling of the flux tower footprint

J. A. Gamon

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



requirement for a saturating light pulse, PRI relies on reflected radiation and can be applied over a range of scales, but this also adds challenges to the interpretation. Over longer time scales, or when comparing different species or vegetation types, a primary cause of PRI variation is the adjustments in pigment pool sizes, notably chlorophyll:carotenoid ratios, that strongly influence the PRI wavelengths (Garrity et al., 2011; Wong and Gamon, 2015a). Several recent studies have explored these pool size effects and related them to photosynthetic activity, with promising results. For evergreen species PRI often varies strongly with season (Wong and Gamon, 2015a, b) or with canopy position (Gamon and Berry, 2012) indicating changing chlorophyll:carotenoid levels. This pigment ratio declines during periods of chronic stress, providing a readily detectable optical index of seasonally changing photosynthetic activity and ϵ , with further diurnal modulation in PRI driven by the xanthophyll cycle (Wong and Gamon, 2015a, b). These observations are consistent with studies illustrating the utility of pigments as optical indicators of photosynthetic activity and GPP (Peng et al., 2011), and are contributing to a changing view of the role of how we might best utilize PRI in the assessment of photosynthetic activity.

Much of the confusion in the recent PRI literature arises from the fact that multiple factors drive variation in PRI over different temporal scales (Barton and North, 2001), and relatively few studies have fully considered sampling context or attempted to distinguish these multiple causes. The importance of the temporal context for PRI assessment is illustrated in Fig. 4, showing short-term facultative changes due to xanthophyll cycle activity dominated by long-term constitutive pool size effects. The interaction of these two components yield complex spatial patterns of PRI within the flux footprint due to the complex light regimes within a stand (Hilker et al., 2008; Middleton et al., 2009), varying with canopy depth (Gamon and Berry, 2012) and aspect (Gamon and Bond, 2013). Direct assessment of photosynthetic activity and ϵ with PRI remains an area of ongoing research, requiring a careful consideration of sampling context affecting these PRI components. Some sampling methods allow separation of PRI into its facultative xanthophyll cycle and constitutive pigment pool components. These methods typically

BGD

12, 4973–5014, 2015

Optical sampling of the flux tower footprint

J. A. Gamon

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



particular context or question of interest, with different investigators pursuing different approaches. A key opportunity lies in comparing these different approaches to each other and across sites within the context of the flux tower network. As discussed below, it is likely that any single parameterization will not apply equally well across all ecosystems, biomes or seasons due to operational issues and contrasting controls on physiology and structure, but this is also an open question.

3.4 Solar-induced fluorescence

For years, studies have noted a close association between Solar-Induced chlorophyll Fluorescence (SIF) and photosynthetic activity (Carter et al., 1990). Recent studies, mostly from spaceborne platforms, have demonstrated a strong link between GPP and SIF (Frankenberg et al., 2011; Joiner et al., 2011, 2014). The technical breakthrough of the recent SIF studies arose from the recognition that atmospheric absorption bands (e.g. the Fraunhofer lines) contain a small fluorescence signal that can be detected with the appropriate narrow-band instruments, allowing quantification of this small signal against a larger background of solar radiation (Damm et al., 2010; Meroni et al., 2009). Some of these studies are the accidental consequence of satellites designed for other purposes, so the fluorescence measurements have not been optimized in terms of signal-to-noise. Because this fluorescence signal in these satellite instruments is relatively weak, data must be either aggregated over large regions, annual time scales, or both, providing coarse, biome-scale estimates instead of site-based measurements. This aggregation has demonstrated remarkable coarse-scale agreement between fluorescence and GPP for most biomes (Frankenberg et al., 2011; Joiner et al., 2011, 2014), but necessarily obscures the influence of fine-scale regulatory mechanisms, which have yet to be fully addressed in most SIF studies.

Despite the novelty of these recent spaceborne applications, basic fluorescence theory has been understood for decades (see reviews by Meroni et al., 2009; Porcar-Castell et al., 2014), and the chlorophyll fluorescence signal is known to be affected by both the amount of illuminated chlorophyll (related to APAR), and the degree of

Optical sampling of the flux tower footprint

J. A. Gamon

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Optical sampling of the flux tower footprint

J. A. Gamon

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



photosynthetic downregulation (related to ε) (Porcar-Castell et al., 2014). As a demonstration of the latter point, fluorescence yield ($\Delta F/Fm'$), typically derived using the pulse-amplitude modulated (PAM) method provide an excellent means of assessing (ε), closely matching short-term PRI dynamics, but far less scaleable, since the PAM technique requires a saturating light pulse, restricting the measurements to proximal measurements (Gamon et al., 1997). As discussed above, this downregulation occurs under a variety of stresses and can be related to meteorological conditions, xanthophyll cycle, pigment pools, and ε . Thus, it is plausible that SIF, because it is readily scaleable and sensitive to *both* terms of the LUE model, can provide an optical metric that scales with realized rates of photosynthesis or GPP, bypassing the need for independent assessment of the LUE model terms (Fig. 5).

Since several of these recent observations of SIF have been at a coarse biome scale, more work is needed to confirm this hypothesis, and to compare this method against other established optical methods over a range of temporal and spatial scales, with proper consideration of sampling context. As of this writing, few off-the-shelf commercial instruments have the sufficient narrow bandwidth, signal-to-noise, and stability to extract a reliable fluorescence signal. Consequently, there is a need for better instrumentation, and several research groups are actively developing and testing improved instruments for proximal SIF application. Currently, there is a rich opportunity for testing the SIF fluorescence approach using the flux tower network and LUE model as a validation framework. Ideally, such experiments would include not only fluxes, but also other optical measurements (e.g. irradiance and NDVI and PRI) to independently and simultaneously assess the LUE model components.

4 Sampling considerations: a question of scale

When conducting ecosystem flux measurements using eddy covariance, the challenges of integration with optical sampling across scales becomes clear. Many eddy covariance systems collect data at 10 to 20 Hz, and report summary data on half-

hourly time steps (Burba and Anderson, 2010). When expressing net primary production (NPP), yearly aggregated carbon uptake is a common expression. Aggregating flux data over such large temporal scales misses many of the fine-scale regulatory processes described above, each of which can provide a partial understanding of the environmental constraints and physiological responses that can contribute to the overall controls of photosynthesis or productivity. Thus, a key challenge in comparing optical and flux measurements lies in careful consideration of scale, and development of the most appropriate methods for sampling and subsequent data aggregation in the temporal or spatial dimensions. Because of the complexity of stand light regimes, angular scale (e.g. sun angle and view angle) can be considered an additional sampling dimension. The spectral scale represents an additional sampling dimension that must be carefully matched to the question at hand. Key examples of scaling issues as applied to optical measurements are further discussed below.

4.1 Spatial dimension

Due to the inherent variability of the flux tower footprint (Vesala et al., 2008), matching the full sampling region with optical sampling is a challenging task. From a practical perspective, it is difficult to obtain a meaningful, spatially representative sampling of the footprint without disturbing the vegetation or flux measurements themselves. While tractable in short-statured vegetation, this becomes far more difficult for vegetation exceeding 2–3 m in height. A brief discussion of instrumentation is warranted here. The choice of method and its proper application depends to a large extent on the particular sampling objective. To follow basic physiological processes, high-frequency leaf- or canopy-scale sampling is often used, and automated methods may be necessary. For fully characterizing the flux tower footprint, fine-grained sampling (ideally 1 m² on a daily time step or finer) is often optimal. Most publicly available satellite data are typically too coarse-grained to provide this level of spatially detailed footprint monitoring. Commercial satellite options, some of which sample at fine spatial scales, tend to be cost prohibitive for routine applications. Airborne sampling can provide a flexible plat-

Optical sampling of the flux tower footprint

J. A. Gamon

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Optical sampling of the flux tower footprint

J. A. Gamon

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



form to help fill this need, but typically at a high cost. Consequently, to sample in the footprint, many new applications of field spectrometers have been developed (Fig. 6). Most of these methods are either limited in height (Fig. 6a and c), or in spatial replication (Fig. 6b), making full footprint sampling difficult. The advent of inexpensive drones that can carry small spectrometers (Fig. 6d) now offer improved optical coverage of the entire flux footprint, as technical and legal issues (Elias, 2012) become resolved. Alternatively, new automated, portable sensors, including 2-band NDVI or PRI sensors (Eklundh et al., 2011; Harris et al., 2014; Gamon et al., 2015) allow greater flexibility in sampling, particularly in the time domain. Wireless sensor networks enable improved spatial coverage, but at added cost and complexity. Affordable, image-based methods, including multiband cameras, portable imaging spectrometers, and webcams (Fig. 6b) are now becoming widely available (e.g. Richardson et al., 2009) and offer the interpretive power of imagery, but with additional analytical challenges resulting from ill-defined sampling geometry, which can create difficulties when trying to conduct cross-site or multi-season analyses.

4.2 Angular dimension

Because vegetation optical signals have a strong angle-dependency, consideration and normalization of sampling geometry is needed if standard products are to be generated and compared (Barton and North, 2001; Hilker et al., 2008, 2011). Traditionally, optical remote sensing emphasizes the nadir (vertical) view, in part because it provides a standard that can be readily replicated and measured from above. Multi-angle measurements can help provide a deeper functional understanding of vegetation optical signals relevant to fluxes (Hilker et al., 2011), largely because vegetation physiology and resulting patterns of reflected radiation are so responsive to illumination. This can become especially relevant when evaluating diurnal responses or comparing measurements across latitudes, seasons, or contrasting stand structure (Hilker et al., 2010). Vegetation physiology and structure both affect the measured optical signals, which are also strongly influenced by sun angle or sky conditions (Barton and North, 2001).

spectral range of the solar spectrum, from roughly 300 nm (in the near-UV) to 2500 nm (the short-wave infrared). Many other bands and indices, including many pigment indices not discussed here offer additional ways to explore factors influencing fluxes. Depending upon the specific sampling goal, some methods use relatively few, discrete bands to assess particular features, while others may utilize the full spectral range, yielding a seemingly infinite number of sampling possibilities. Similarly, analytical approaches can range from specific targeted approaches using a few optimal formulations to more open-ended statistical methods to discover which bands or formulations can yield the best result for a given purpose or context (e.g. Serbin et al., 2011). Particularly for detecting specific physiological features (e.g. pigments, water content, or chlorophyll fluorescence), proper attention to waveband definition (both spectral position and bandwidth) is essential. Many narrow-band indices such as PRI are notoriously sensitive to formulation, requiring that users and metadata specify the exact band definitions used, and this may vary slightly with instrument manufacturer (Castro et al., 2006), or even among instruments from the same manufacturer (Harris et al., 2015). Furthermore, as more studies explore pigment pool sizes from different instrument platforms, new and improved expressions of this and other pigment indices will emerge, each requiring renewed attention to band definitions. Greater attention to standardization of instruments, calibration procedures, and field methods is clearly needed. Given the multiplicity of brands, calibration methods, and field protocols in use, improved attention to band definitions in reporting data and metadata is also needed, particularly if we are to compare results across instruments or studies (Gamon et al., 2010).

5 Vegetation optical types

Particularly when combined with flux measurements, variation in optical behavior can reveal contrasting vegetation light absorption (APAR), light-use efficiency (ϵ), and other functional differences reflecting differing environmental and evolutionary constraints. This introduces the hypothesis of vegetation *optical types*: functionally significant vari-

BGD

12, 4973–5014, 2015

Optical sampling of the flux tower footprint

J. A. Gamon

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ation in optical properties resulting from a combination of leaf traits, canopy structure, and phenology (Ustin and Gamon, 2010). This concept applies to remote sensing of biodiversity, but also provides a useful framework for analyzing the functional responses of different vegetation types, ecosystems or biomes having different environmental limitations on fluxes. While full-spectral methods are often used, we illustrate the concept and its application with a focus on NDVI and PRI, largely because of their relevance to the LUE model and photosynthetic function.

One example of optical types emerges from the global biome mapping effort based on NDVI phenology (Defries and Townshend, 1994). At the ecosystem scale, using the light-response and LUE model framework (Fig. 3b), optical and flux measurements can be combined to distinguish optical types from environments having different resource levels associated with contrasting light-use efficiencies (ϵ values) (Huemmrich et al., 2011), providing a convenient basis for upscaling fluxes (Huemmrich et al., 2013). At the leaf or canopy level, there is a well-developed history of vegetation functional categories, based on concepts of adaptation and resource theory (Grime, 1977; Bloom et al., 1985; Wright et al., 2004). Sampling efforts can address categorical distinctions such as evergreens, annuals, or deciduous perennials (Gamon et al., 1997), or can be represented as a continuous spectrum of trait variation (Wright et al., 2004). In both cases, optical measurements can provide useful additional plant trait information regarding evolutionary responses or resource limitations at a range of scales. There is a rich opportunity to further define this concept, to apply it to the flux tower footprint, and to use it to further clarify controls on surface–atmosphere fluxes of mass and energy and as a foundation for upscaling. Integrating optical sampling across the global flux tower network provides an ideal opportunity to further explore the optical type hypothesis.

In a recent, review and meta-analysis of the PRI literature (Garbulsky et al., 2011) proposed that different ecosystems having contrasting constraints on carbon flux should also have contrasting optical properties reflected in PRI and NDVI dynamics. According to this hypothesis, the relationship between these optical indices and

BGD

12, 4973–5014, 2015

Optical sampling of the flux tower footprint

J. A. Gamon

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Optical sampling of the flux tower footprint

J. A. Gamon

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



ecosystem carbon flux should vary across ecosystems, due to the contrasting environmental constraints and evolutionary responses to these constraints. Here, we restate this hypothesis to illustrate how NDVI and PRI signals might vary across functionally distinct vegetation types to yield complementary information on fluxes (Fig. 8). This hypothesis predicts that carbon fluxes in deciduous or annual vegetation (e.g. grasslands or most crops showing strong growth cycles) should be primarily driven by canopy structural changes, detectable via NDVI and related to LAI, F_{PAR} , and other structural parameters strongly influencing ecosystem fluxes (Fig. 8a). Accordingly, these ecosystems would exhibit less variation associated with changing efficiency (ε), detectable as PRI (Fig. 8c; e.g. Flanagan et al., 2015). On the other hand, evergreen vegetation having little seasonal variation in canopy structure should have a weak relationship between NDVI and carbon fluxes (Fig. 8b), but would have a strong correlation between PRI and fluxes (Fig. 8c), driven by the strong seasonal changes in chlorophyll:carotenoid ratios (Wong and Gamon, 2015a, b; Gamon et al., 2015).

This hypothesis of contrasting, complementary physiological and structural controls for different optical types has yet to be fully tested across a wide range of ecosystems and conditions, in part because the lack of standardized sampling protocols and appropriate databases have made it difficult to directly compare separate measurements from different studies, instruments, and ecosystems. Fully testing this hypothesis requires properly replicated instrumentation and experimental design, something that can best be achieved within the context of combined flux and optical networks. The benefit of this approach is that it can guide the application and inform the parameterization of the LUE model across contrasting ecosystems. In landscapes dominated by annual or deciduous vegetation, a good characterization of APAR via NDVI and irradiance measurements might be sufficient to define the annual course of gas exchange (Flanagan et al., 2015); in landscapes dominated by evergreens, the additional information on ε becomes necessary, and sampling PRI could improve our understanding of flux controls and phenology (Wong and Gamon, 2015a, b). Adding standardized NDVI and PRI sensors to the flux tower network, and comparing ground measure-

ments to satellite measurements, presents an ideal opportunity to test this hypothesis of complementary NDVI and PRI behavior associated with different vegetation types.

6 Conclusions: the path forward

In the past, remote sensing has provided useful estimates of biophysical parameters and a powerful means of extrapolating beyond the flux tower footprint. However, a newer crop of optical sampling methods has now emerged that allow deeper exploration of both physiological and structural controls on carbon fluxes. Two decades of proximal remote sensing within flux tower footprints have yielded many studies that relate optical signals to various components of ecosystem carbon fluxes. However, the literature largely remains a series of site-based observations, and we are still a long way from a comprehensive understanding of how best to integrate optical and flux measurements across biomes. Comparative, experimental approaches are needed both within and across sites. The goal of this review has been to consider accomplishments so far, and to suggest directions and hypotheses for future work in the context of the global flux tower network.

Clearly, the LUE model provides a useful framework that is readily integrated with photosynthetic flux measurements, but we are left with several questions. Is there a single, consistent way to parameterize the LUE model from remote sensing, or does this model have to be parameterized and calibrated individually for each ecosystem? Can the optical type concept be universally useful as a means of categorizing vegetation functional response and upscaling from sites to larger regions? Alternatively, are we better-served by continuous-field approaches that interpret optical measurements in terms of continuous responses to environmental conditions and resource gradients? How do newly emerging SIF methods compare to other more established optical methods based on reflectance? Experimental, cross-site approaches are needed to more fully address these questions.

Optical sampling of the flux tower footprint

J. A. Gamon

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Optical sampling of the flux tower footprint

J. A. Gamon

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A full functional characterization of the flux tower footprint requires more than the LUE model, and continued comparisons with other approaches are also warranted. In addition to the optical sampling methods, other methods not fully discussed here, including thermal, lidar, microwave, and acoustic remote sensing, can add to our understanding of functional processes within the footprint. To obtain a deeper mechanistic understanding, the LUE approach can be compared to other, more biochemically based models of photosynthetic carbon flux (e.g., Sellers et al., 1986). Presumably, integration of several different and complementary optical methods (e.g. reflectance with fluorescence and canopy absorptance) would provide a more complete assessment of ecosystem function than any single method alone (Fig. 5). Better integration of optical with structural (e.g. LIDAR) measurements, particularly using canopy radiative transfer models, would add further insight into the structural vs. physiological controls on carbon flux. Integration of thermal sampling methods could help integrate energy balance and water vapor fluxes with the LUE approach.

We need to expand our optical exploration of pigments beyond chlorophyll content and xanthophyll cycle pigment activity to include a wider range of plant pigments and their functional roles, with a particular focus on the rich world of carotenoid biochemistry and function. This would include the evaluation of carotenoid : chlorophyll pigment pools as a generalized stress response affecting isoprenoid emissions in addition to CO₂ fluxes. Currently, our understanding of the role of carotenoids as optical indicators of stress responses is in its infancy, but an area ripe for further exploration. As new instruments and platforms are developed, it is likely that new and improved pigment indices will also emerge.

To accomplish these goals, we need more systematic deployment of optical sensors with better attention to contextual information, as discussed above. Ideally, protocols would be sufficiently standardized or defined to allow ready comparisons across sites and studies. To match this deployment, we need a capable and flexible informatics approach that encompasses the large data volume represented by Fig. 2. This should accommodate not only flux and optical data, but also the essential ancillary data (e.g.

Optical sampling of the flux tower footprint

J. A. Gamon

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Barton, C. V. M. and North, P. R. J.: Remote sensing of canopy light use efficiency using the photochemical reflectance index – model and sensitivity analysis, *Remote Sens. Environ.*, 78, 264–273, doi:10.1016/s0034-4257(01)00224-3, 2001.

Björkman, O.: Responses to different quantum flux densities, in: *Encyclopedia of Plant Physiology. New Series. Volume 12 A. Physiological Plant Ecology I. Responses to the Physical Environment*, edited by: Lange, O. L., Nobel, P. S., Osmond, C. B., and Ziegler, H., 57–107, 1981.

Björkman, O. and Demmig-Adams, B.: Regulation of photosynthetic light energy capture, conversion, and dissipation in leaves of higher plants, in: *Ecophysiology of Photosynthesis*, edited by: Schulze, E.-D., Caldwell, M. M., and Pospisilova, J., Springer, Berlin, Heidelberg, New York, 27–47, 1994.

Bloom, A. J., Chapin, F. S., and Mooney, H. A.: Resource limitation in plants-an economic analogy, *Annu. Rev. Ecol. Syst.*, 16, 363–392, 1985.

Brugnoli, E. and Björkman, O.: Chloroplast movements in leaves – influence on chlorophyll fluorescence and measurements of light-induced absorbency changes related to delta-pH and zeaxanthin formation, *Photosynth. Res.*, 32, 23–35, doi:10.1007/bf00028795, 1992.

Burba, G. and Anderson, D.: *A Brief Practical Guide to Eddy Covariance Flux Measurements: Principles and Workflow Examples for Scientific and Industrial Applications*, Li-Cor Biosciences, 2010.

Carter, G., Theisen, A., and Mitchell, R.: Chlorophyll fluorescence measured using the Fraunhofer line – depth principle and relationship to photosynthetic rate in the field, *Plant Cell Environ.*, 13, 79–83, 1990.

Damm, A., Elbers, J., Erler, A., Gioli, B., Hamdi, K., Hutjes, R., Kosvancova, M., Meroni, M., Miglietta, F., Moersch, A., Moreno, J., Schickling, A., Sonnenschein, R., Udelhoven, T., van der Linden, S., Hostert, P., Rascher, U.: Remote sensing of sun-induced fluorescence to improve modeling of diurnal courses of gross primary production (GPP), *Glob. Change Biol.*, 16, 171–186, doi:10.1111/j.1365-2486.2009.01908.x, 2010.

Defries, R. S. and Townshend, J. R. G.: NDVI-derived land cover classifications at a global-scale, *Int. J. Remote Sens.*, 15, 3567–3586, 1994.

Demmig, B., Winter, K., Kruger, A., and Czygan, F.-C.: Photoinhibition and zeaxanthin formation in intact leaves – a possible role of the xanthophyll cycle in the dissipation of excess light energy, *Plant Physiol.*, 84, 218–224, 1987.

Optical sampling of the flux tower footprint

J. A. Gamon

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Drolet, G. G., Middleton, E. M., Huemmrich, K. F., Hall, F. G., Amiro, B. D., Barr, A. G., Black, T. A., McCaughey, J. H., and Margolis, H. A.: Regional mapping of gross light-use efficiency using MODIS spectral indices, *Remote Sens. Environ.*, 112, 3064–3078, doi:10.1016/j.rse.2008.03.002, 2008.
- 5 Eklundh, L., Jin, H., Schubert, P., Guzinski, R., and Heliasz, M.: An optical sensor network for vegetation phenology monitoring and satellite data calibration, *Sensors*, 11, 7678–7709, doi:10.3390/s110807678, 2011.
- Elias, B.: Pilotless drones: background and considerations for Congress regarding unmanned aircraft operations in the national airspace system, Congressional Research Service, Library of Congress, 2012.
- 10 Flanagan, L. B., Sharp, E. J., and Gamon, J. A.: Application of the photosynthetic light-use efficiency model in a northern Great Plains grassland, *Remote Sens. Environ.*, in review, 2015.
- Frankenberg, C., Fisher, J. B., Worden, J., Badgley, G., Saatchi, S. S., Lee, J.-E., Toon, G. C., Butz, A., Jung, M., Kuze, A., and Yokota, T.: New global observations of the terrestrial carbon cycle from GOSAT: patterns of plant fluorescence with gross primary productivity, *Geophys. Res. Lett.*, 38, L17706, doi:10.1029/2011gl048738, 2011.
- Gamon, J. A. and Berry, J. A.: Facultative and constitutive pigment effects on the Photochemical Reflectance Index (PRI) in sun and shade conifer needles, *Isr. J. Plant Sci.*, 60, 85–95, doi:10.1560/ijps.60.1-2.85, 2012.
- 20 Gamon, J. A. and Bond, B.: Effects of irradiance and photosynthetic downregulation on the photochemical reflectance index in Douglas-fir and ponderosa pine, *Remote Sens. Environ.*, 135, 141–149, doi:10.1016/j.rse.2013.03.032, 2013.
- Gamon, J. A. and Surfus, J. S.: Assessing leaf pigment content and activity with a reflectometer, *New Phytol.*, 143, 105–117, doi:10.1046/j.1469-8137.1999.00424.x, 1999.
- 25 Gamon, J. A., Field, C. B., Bilger, W., Bjorkman, O., Fredeen, A. L., and Peñuelas, J.: Remote-sensing of the xanthophyll cycle and chlorophyll fluorescence in sunflower leaves and canopies, *Oecologia*, 85, 1–7, doi:10.1007/bf00317336, 1990.
- Gamon, J. A., Peñuelas, J., and Field, C. B.: A narrow-waveband spectral index that tracks diurnal changes in photosynthetic efficiency, *Remote Sens. Environ.*, 41, 35–44, doi:10.1016/0034-4257(92)90059-s, 1992.
- 30

Optical sampling of the flux tower footprint

J. A. Gamon

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Gamon, J. A., Field, C. B., Goulden, M. L., Griffin, K. L., Hartley, A. E., Joel, G., Peñuelas, J., and Valentini, R.: Relationships between NDVI, canopy structure, and photosynthesis in three Californian vegetation types, *Ecol. Appl.*, 5, 28–41, doi:10.2307/1942049, 1995.

Gamon, J. A., Serrano, L., and Surfus, J. S.: The photochemical reflectance index: an optical indicator of photosynthetic radiation use efficiency across species, functional types, and nutrient levels, *Oecologia*, 112, 492–501, doi:10.1007/s004420050337, 1997.

Gamon, J. A., Cheng, Y. F., Claudio, H., MacKinney, L., and Sims, D. A.: A mobile tram system for systematic sampling of ecosystem optical properties, *Remote Sens. Environ.*, 103, 246–254, doi:10.1016/j.rse.2006.04.006, 2006a.

Gamon, J. A., Rahman, A. F., Dungan, J. L., Schildhauer, M., and Huemmrich, K. F.: Spectral Network (SpecNet) – what is it and why do we need it?, *Remote Sens. Environ.*, 103, 227–235, doi:10.1016/j.rse.2006.04.003, 2006b.

Gamon, J. A., Coburn, C., Flanagan, L. B., Huemmrich, K. F., Kiddle, C., Sanchez-Azofeifa, G. A., Thayer, D. R., Vescovo, L., Gianelle, D., Sims, D. A., Rahman, A. F., and Pastorello, G. Z.: SpecNet revisited: bridging flux and remote sensing communities, *Can. J. Remote Sens.*, 36, S376–S390, 2010.

Gamon, J. A., Kovalchuk, O., Wong, C. Y. S., Harris, A., and Garrity, S. R.: Monitoring seasonal and diurnal changes in photosynthetic pigments with automated PRI and NDVI sensors, *Biogeosciences Discuss.*, 12, 2947–2978, doi:10.5194/bgd-12-2947-2015, 2015.

Garbulsky, M. F., Peñuelas, J., Papale, D., Ardo, J., Goulden, M. L., Kiely, G., Richardson, A. D., Rotenberg, E., Veenendaal, E. M., and Filella, I.: Patterns and controls of the variability of radiation use efficiency and primary productivity across terrestrial ecosystems, *Global Ecol. Biogeogr.*, 19, 253–267, doi:10.1111/j.1466-8238.2009.00504.x, 2010.

Garbulsky, M. F., Peñuelas, J., Gamon, J., Inoue, Y., and Filella, I.: The photochemical reflectance index (PRI) and the remote sensing of leaf, canopy and ecosystem radiation use efficiencies: a review and meta-analysis, *Remote Sens. Environ.*, 115, 281–297, doi:10.1016/j.rse.2010.08.023, 2011.

Garrity, S. R., Eitel, J. U. H., and Vierling, L. A.: Disentangling the relationships between plant pigments and the photochemical reflectance index reveals a new approach for remote estimation of carotenoid content, *Remote Sens. Environ.*, 115, 628–635, doi:10.1016/j.rse.2010.10.007, 2011.

Optical sampling of the flux tower footprint

J. A. Gamon

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Gitelson, A. A. and Gamon, J. A.: The need for a common basis for defining light-use efficiency: implications for productivity estimation, *Remote Sens. Environ.*, 156, 196–201, doi:10.1016/j.rse.2014.09.017, 2015.

Goerner, A., Reichstein, M., Rambal, S.: Tracking the seasonal drought effects on ecosystem light use efficiency with satellite-based PRI in a Mediterranean forest, *Remote Sens. Environ.*, 113, 1101–1111, doi:10.1016/j.rse.2009.02.001, 2009.

Goulden, M. L., Daube, B. C., Fan, S.-M., Sutton, D. J., Bazzaz, A., Munger, J. W., and Wofsy, S. C.: Physiological responses of a black spruce forest to weather, *J. Geophys. Res.-Atmos.*, 102, 28987–28996, doi:10.1029/97jd01111, 1997.

Goward, S. N., Tucker, C. J., and Dye, D. G.: North-American vegetation patterns observed with the NOAA-7 Advanced Very High Resolution Radiometer, *Vegetatio*, 64, 3–14, doi:10.1007/bf00033449, 1985.

Grime, J. P.: Evidence for the existence of three primary strategies in plants and its relevance to ecological and evolutionary theory, *Am. Nat.*, 111, 1169–1194, doi:10.1086/283244, 1977.

Guarini, R., Nichol, C., Clement, R., Loizzo, R., Grace, J., and Borghetti, M.: The utility of MODIS-sPRI for investigating the photosynthetic light-use efficiency in a Mediterranean deciduous forest, *Int. J. Remote Sens.*, 35, 6157–6172, doi:10.1080/01431161.2014.950762, 2014.

Harris, A., Gamon, J. A., Pastorello, G. Z., and Wong, C. Y. S.: Retrieval of the photochemical reflectance index for assessing xanthophyll cycle activity: a comparison of near-surface optical sensors, *Biogeosciences*, 11, 6277–6292, doi:10.5194/bg-11-6277-2014, 2014.

Heinsch, F. A., Zhao, M., Running, S. W., Kimball, J. S., Nemani, R. R., Davis, K. J., Bolstad, P. V., Cook, B. D., Desai, A. R., Ricciuto, D. M., Law, B. E., Oechel, W. C., Kwon, H., Luo, H., Wofsy, S. C., Dunn, A. L., Munger, J. W., Baldocchi, D. D., Xu, L., Hollinger, D. Y., Richardson, A. D., Stoy, P. C., Siqueira, M. B. S., Monson, R. K., Burns, S. P., and Flanagan, L. B.: Evaluation of remote sensing based terrestrial productivity from MODIS using regional tower eddy flux network observations, *IEEE T. Geosci. Remote*, 44, 1908–1925, doi:10.1109/tgrs.2005.853936, 2006.

Hilker, T., Coops, N. C., Nestic, Z., Wulder, M. A., and Black, A. T.: Instrumentation and approach for unattended year round tower based measurements of spectral reflectance, *Comput. Electron. Agr.*, 56, 72–84, doi:10.1016/j.compag.2007.01.003, 2007.

**Optical sampling of
the flux tower
footprint**

J. A. Gamon

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Hilker, T., Coops, N. C., Hall, F. G., Black, T. A., Wulder, M. A., Nestic, Z., and Krishnan, P.: Separating physiologically and directionally induced changes in PRI using BRDF models, *Remote Sens. Environ.*, 112, 2777–2788, doi:10.1016/j.rse.2008.01.011, 2008.
- Hilker, T., Hall, F. G., Coops, N. C., Lyapustin, A., Wang, Y., Nestic, Z., Grant, N., Black, T. A., Wulder, M. A., Klijun, N., Hopkinson, C., and Chasmer, L.: Remote sensing of photosynthetic light-use efficiency across two forested biomes: spatial scaling, *Remote Sens. Environ.*, 114, 2863–2874, doi:10.1016/j.rse.2010.07.004, 2010.
- Hilker, T., Gitelson, A., Coops, N. C., Hall, F. G., and Black, T. A.: Tracking plant physiological properties from multi-angular tower-based remote sensing, *Oecologia*, 165, 865–876, doi:10.1007/s00442-010-1901-0, 2011.
- Huemmrich, K. F., Black, T. A., Jarvis, P. G., McCaughey, J. H., and Hall, F. G.: High temporal resolution NDVI phenology from micrometeorological radiation sensors, *J. Geophys. Res.-Atmos.*, 104, 27935–27944, doi:10.1029/1999jd900164, 1999.
- Huemmrich, K. F., Gamon, J. A., Tweedie, C. E., Oberbauer, S. F., Kinoshita, G., Houston, S., Kuchy, A., Hollister, R. D., Kwon, H., Mano, M., Harazono, Y., Webber, P. J., and Oechel, W. C.: Remote sensing of tundra gross ecosystem productivity and light use efficiency under varying temperature and moisture conditions, *Remote Sens. Environ.*, 114, 481–489, doi:10.1016/j.rse.2009.10.003, 2010.
- Huemmrich, K. F., Gamon, J. A., Tweedie, C. E., Campbell, P. K. E., Landis, D. R., and Middleton, E. M.: Arctic tundra vegetation functional types based on photosynthetic physiology and optical properties, *IEEE J. Sel. Top. Appl.*, 6, 265–275, doi:10.1109/jstars.2013.2253446, 2013.
- Joiner, J., Yoshida, Y., Vasilkov, A. P., Yoshida, Y., Corp, L. A., and Middleton, E. M.: First observations of global and seasonal terrestrial chlorophyll fluorescence from space, *Biogeosciences*, 8, 637–651, doi:10.5194/bg-8-637-2011, 2011.
- Joiner, J., Yoshida, Y., Vasilkov, A., Schaefer, K., Jung, M., Guanter, L., Zhang, Y., Garrity, S., Middleton, E. M., Huemmrich, K. F., Gu, L., and Marchesini, L. B.: The seasonal cycle of satellite chlorophyll fluorescence observations and its relationship to vegetation phenology and ecosystem atmosphere carbon exchange, *Remote Sens. Environ.*, 152, 375–391, doi:10.1016/j.rse.2014.06.022, 2014.
- Jones, H. G.: *Plants and Microclimate: a Quantitative Approach to Environmental Plant Physiology*, Cambridge University Press, 2013.

Optical sampling of the flux tower footprint

J. A. Gamon

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Knobl, A. and Baldocchi, D. D.: Effects of diffuse radiation on canopy gas exchange processes in a forest ecosystem, *J. Geophys. Res.-Biogeo.*, 113, G02023, doi:10.1029/2007jg000663, 2008.

Kumar, M. and Monteith, J. L.: Remote sensing of plant growth, in: *Plants and the Daylight Spectrum*, edited by: Smith, H., Academic Press, London, 133–144, 1981.

Long, S. P. and Bernacchi, C. J.: Gas exchange measurements, what can they tell us about the underlying limitations to photosynthesis? *Procedures and sources of error*, *J. Exp. Bot.*, 54, 2393–2401, doi:10.1093/jxb/erg262, 2003.

Meroni, M., Rossini, M., Guanter, L., Alonso, L., Rascher, U., Colombo, R., and Moreno, J.: Remote sensing of solar-induced chlorophyll fluorescence: review of methods and applications, *Remote Sens. Environ.*, 113, 2037–2051, doi:10.1016/j.rse.2009.05.003, 2009.

Michaletz, S. T., Cheng, D., Kerkhoff, A. J., and Enquist, B. J.: Convergence of terrestrial plant production across global climate gradients, *Nature*, 512, 39–43, doi:10.1038/nature13470, 2014.

Middleton, E. M., Cheng, Y.-B., Hilker, T., Black, T. A., Krishnan, P., Coops, N. C., and Huemmrich, K. F.: Linking foliage spectral responses to canopy-level ecosystem photosynthetic light-use efficiency at a Douglas-fir forest in Canada, *Can. J. Remote Sens.*, 35, 166–188, 2009.

Monteith, J. L.: Climate and efficiency of crop production in Britain, *Philos. T. Roy. Soc. B*, 281, 277–294, doi:10.1098/rstb.1977.0140, 1977.

Nestola, E., Calfapietra, C., Emmerton, C. E., Wong, C. Y. S., Thayer, D. R., and Gamon, J. A.: Monitoring grassland carbon dynamics: integrating optical and flux measurements, *Biogeosciences*, in review, 2015.

Norman, J. M. and Arkebauer, T. J.: Predicting canopy light-use efficiency from leaf characteristics, Chapter 5, in: *Modeling Crop Photosynthesis – from Biochemistry to Canopy*, edited by: Boote, K. J. and Loomis, R. S., CCSA Special Publication 19, 125–143, 1991.

Osmond, C. B. and Chow, W. S.: Ecology of photosynthesis in the sun and shade – summary and prognostications, *Aust. J. Plant Physiol.*, 15, 1–9, 1988.

Peng, Y., Gitelson, A. A., Keydan, G., Rundquist, D. C., and Moses, W.: Remote estimation of gross primary production in maize and support for a new paradigm based on total crop chlorophyll content, *Remote Sens. Environ.*, 115, 978–989, doi:10.1016/j.rse.2010.12.001, 2011.

Optical sampling of the flux tower footprint

J. A. Gamon

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Sellers, P. J., Hall, F. G., Kelly, R. D., Black, A., Baldocchi, D., Berry, J., Ryan, M., Ranson, K. J., Crill, P. M., Lettenmaier, D. P., Margolis, H., Cihlar, J., Newcomer, J., Fitzjarrald, D., Jarvis, P. G., Gower, S. T., Halliwell, D., Williams, D., Goodison, B., Wickland, D. E., and Guertin, F. E.: BOREAS in 1997: experiment overview, scientific results, and future directions, *J. Geophys. Res.-Atmos.*, 102, 28731–28769, doi:10.1029/97jd03300, 1997.
- Serbin, S. P., Dillaway, D. N., Kruger, E. L., and Townsend, P. A.: Leaf optical properties reflect variation in photosynthetic metabolism and its sensitivity to temperature, *J. Exp. Bot.*, 63, 489–502, doi:10.1093/jxb/err294, 2011.
- Sims, D. A., Luo, H. Y., Hastings, S., Oechel, W. C., Rahman, A. F., and Gamon, J. A.: Parallel adjustments in vegetation greenness and ecosystem CO₂ exchange in response to drought in a Southern California chaparral ecosystem, *Remote Sens. Environ.*, 103, 289–303, doi:10.1016/j.rse.2005.01.020, 2006.
- Soudani, K., Hmimina, G., Dufrêne, E., Berveiller, D., Delpierre, N., Ourcival, J.-M., Rambal, S., and Joffre, R.: Relationships between photochemical reflectance index and light-use efficiency in deciduous and evergreen broadleaf forests, *Remote Sens. Environ.*, 144, 73–84, doi:10.1016/j.rse.2014.01.017, 2014.
- Tucker, C. J., Townshend, J. R. G., and Goff, T. E.: African land-cover classification using satellite data, *Science*, 227, 369–375, 1985.
- Ustin, S. L. and Gamon, J. A.: Remote sensing of plant functional types, *New Phytol.*, 186, 795–816, doi:10.1111/j.1469-8137.2010.03284.x, 2010.
- Vesala, T., Kljun, N., Rannik, Ü., Rinne, J., Sogachev, A., Markkanen, T., Sabelfeld, K., Foken, T., and Leclerc, M.: Flux and concentration footprint modelling: state of the art, *Environ. Pollut.*, 152, 653–666, doi:10.1016/j.envpol.2007.06.070, 2008.
- VonCaemmerer, S. and Farquhar, G. D.: Some relationships between the biochemistry of photosynthesis and the gas-exchange of leaves, *Planta*, 153, 376–387, 1981.
- Wofsy, S. C., Goulden, M. L., Munger, J. W., Fan, S. M., Bakwin, P. S., Daube, B. C., Basow, S. L., and Bazzaz, F. A.: Net exchange of CO₂ in a midlatitude forest, *Science*, 260, 1314–1317, doi:10.1126/science.260.5112.1314, 1993.
- Wong, C. Y. S. and Gamon, J. A.: Three causes of variation in the Photochemical Reflectance Index (PRI) in evergreen conifers, *New Phytol.*, 206, 187–195, doi:10.1111/nph.13159, 2015a.

Optical sampling of the flux tower footprint

J. A. Gamon

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Wong, C. Y. S. and Gamon, J. A.: The photochemical reflectance index provides an optical indicator of spring photosynthetic activation in evergreen conifers, *New Phytol.*, 206, 196–208, doi:10.1111/nph.13251, 2015b.

Wright, I. J., Reich, P. B., Westoby, M., Ackerly, D. D., Baruch, Z., Bongers, F., Cavender-Bares, J., Chapin, T., Cornelissen, J. H. C., Diemer, M., Flexas, J., Garnier, E., Groom, P. K., Gulias, J., Hikosaka, K., Lamont, B. B., Lee, T., Lee, W., Lusk, C., Midgley, J. J., Navas, M.-L., Niinemets, Ü., Oleksyn, J., Osada, N., Poorter, H., Poot, P., Prior, L., Pyankov, V. I., Roumet, C., Thomas, S. C., Tjoelker, M. G., Veneklaas, E. J., and Villar, R.: The worldwide leaf economics spectrum, *Nature*, 428, 821–827, doi:10.1038/nature02403, 2004.

Zhang, L.-X., Zhou, D.-C., Fan, J.-W., and Hu, Z.-M.: Comparison of four light use efficiency models for estimating terrestrial gross primary production, *Ecol. Model.*, 300, 30–39, doi:10.1016/j.ecolmodel.2015.01.001, 2015.

Zygielbaum, A. I., Arkebauer, T. J., Walter-Shea, E. A., and Scoby, D. L.: Detection and measurement of vegetation photoprotection stress response using PAR reflectance, *Isr. J. Plant Sci.*, 60, 37–47, doi:10.1560/IJPS.60.1-2.37, 2012.

Optical sampling of the flux tower footprint

J. A. Gamon




Figure 1. Distribution of absorbed radiation within a leaf, showing processes of productive photosynthesis (blue text and arrows), energy dissipation (red text and arrows), regulatory processes associated with the xanthophyll cycle (black text and arrows within box on left), and carotenoid and chlorophyll pigment pools, all of which can be assessed with optical sampling, but at different temporal, spatial and spectral scales, as discussed in the text.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Figure 2. The sampling domain defined by time, space, and spectral space (Z dimension) axes, illustrating how the optical sampling scale varies with the process being measured. Time and space axes are typically log scales (Osmond and Chow, 1988). The spectral or wavelength axis (represented as the Z dimension), will also vary with the type and purpose of optical sampling. Similarly the appropriate degree of data aggregation in temporal, spatial, and spectral domains will also vary with the purpose at hand.

Optical sampling of the flux tower footprint

J. A. Gamon

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Figure 3. (a) Modeled photosynthetic light-response curves (rectangular hyperbola) for healthy and stressed leaves, illustrating the decline in light-use efficiency (LUE), typically termed photon (or quantum) yield. (b) Stand-level light-response curves of gross ecosystem production (GEP) for different vegetation types having different moisture regimes, LUEs and contrasting optical properties. (a) Modified from Björkman and Demmig-Adams (1994). (b) Modified from Huemmrich et al. (2010).




Figure 4. Illustration of constitutive (pigment pool size) and facultative (xanthophyll cycle) effects on PRI time series, representing two annual cycles of PRI for evergreen vegetation. In this case, the dominant cause of PRI variation is the constitutive adjustment in chlorophyll : carotenoid ratios across seasons, with less variation due to the diurnal xanthophyll cycle operation (Δ PRI), a facultative process that further modulates light energy distribution (Fig. 1). In year two, chronic stress (e.g. drought) causes reduction in the constitutive component (dark-state PRI) along with additional adjustments of the facultative component (Δ PRI). Adapted from Sims et al. (2006); Wong and Gamon (2014)

Optical sampling of the flux tower footprint

J. A. Gamon

[Title Page](#)

[Abstract](#) [Introduction](#)

[Conclusions](#) [References](#)

[Tables](#) [Figures](#)

[◀](#) [▶](#)

[◀](#) [▶](#)

[Back](#) [Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)






Figure 5. Representation of the LUE model (Eq. 2), showing optical measurements useful for model parameterization and validation (red text), including solar-induced fluorescence, the Normalized Difference Vegetation Index (NDVI), the Photochemical Reflectance Index (PRI), and fluorescence yield ($\Delta F/Fm'$). Alternatively, efficiency (ϵ) can be addressed using meteorological data (temperature and vapor pressure deficit) according to Eq. (3), or can be addressed with thermal remote sensing and modeling methods (Anderson et al., 2008; Schull et al., 2015). To most directly compare with eddy covariance, which measures net ecosystem exchange (NEE, analogous to NPP), additional modeling, or partitioning into gross ecosystem production (GEP, analogous to gross primary production, GPP) and respiration (R) is needed.

Optical sampling of the flux tower footprint

J. A. Gamon

Title Page

| | |
|-------------|--------------|
| Abstract | Introduction |
| Conclusions | References |
| Tables | Figures |

⏪ ⏩
◀ ▶
 Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Figure 6. Optical sampling methods applied to spatially extensive sampling within flux tower footprint. **(a)** Hand sampling along a boardwalk leading to a flux tower **(b)** automated sampling with an automated spectrometer (“AMSPEC”) and webcam mounted on a flux tower (adapted from Hilker et al., 2010, 2011), **(c)** automated sampling with a tram, a motorized cart on a track, and **(d)** airborne sampling with an octocopter drone (photo courtesy Richard Streeter and Alasdair Mac Arthur). **(a–c)** depict the “UniSpec DC” (PP Systems, Amesbury MA), a dual detector instrument that can be configured for manual or automatic sampling from a variety of platforms. The “Cinestar 8” drone **(d)** carries a dual field-of-view VNIR “Piccolo” spectrometer (developed by Alasdair MacArthur NERC FSF), comprised of a spectral detector (USB 2000, Ocean Optics, Dunedin FL, USA) with a shuttered, bifurcated fiber providing alternate upward and downward sampling. Note that all methods correct for changing sky conditions, either by simultaneous **(a–c)**, or alternating **(d)** radiance and irradiance sampling, allowing expanded sampling in both space and time to match the scale of eddy covariance measurements.

Optical sampling of the flux tower footprint

J. A. Gamon

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Figure 8. Hypothesized relationships between NDVI (**a** and **b**) and PRI (**c** and **d**) and daily Net Ecosystem Exchange (NEE) or Gross Ecosystem Exchange (GEE) for annual or deciduous (left) or evergreen (right) vegetation. In this example, more positive values indicate greater photosynthetic carbon uptake. Modified from Garbulsky et al. (2011).