### Author comment to Anonymous Referee #1

We thank anonymous Referee #1 for his/her constructive criticism and valuable comments. In the following we address the points raised, with referee comments in boldface and author responses in normal typeface.

#### **General comment:**

Frigstad et al. present data from the PAP site in the northeast Atlantic, with estimates of seasonal NCP and new production that will be useful for the scientific community. A particle tracking method is used to examine source regions of material within a sediment trap, and the manuscript demonstrates that the site shows high flux attenuation by calculating the export ratio and transfer efficiency. Time-series stations such as the PAP site are of great importance in evaluating changes in the environment and understanding these important biogeochemical processes. With further clarification and inclusion of the errors involved, and mixing influences this paper could be a valid contribution to productivity estimates in this region. If the following issues are addressed, I support the publication of this manuscript.

We thank the reviewer for the positive comment, and have addressed the errors involved with calculating NCP and the contribution from mixing in the responses below.

#### **Specific comments:**

# p.5176 line 25. The error associated with the calculation of DIC from calculated TA and measured CO2 should be quantified. The error quoted as $3 \,\mu$ mol kg<sup>-1</sup> is the measurement uncertainty if DIC were measured discretely, which is not the case in the method presented. Please quantify this error and calculate how this error propagates into the NCP estimates.

We have calculated the total error associated with the calculation of DIC from estimated TA and measured fCO2 of  $\pm 0.85$  mol C m<sup>-2</sup>. This was determined by propagation using the method described in Dickson and Riley (1978), together with the errors in the estimated TA values ( $\pm 6.4 \mu$ molkg<sup>-1</sup>; Lee et al. 2006) and measured pCO2 ( $\pm 2 \mu$ atm; Wanninkhof et al., 2013). The propagated probable error is larger than the uncertainty due to monthly variation in NCP<sub>MLD</sub> of  $\pm 0.27$  mol C m<sup>-2</sup>, and we have chosen to use the propagated error to give a conservative estimate of the uncertainty in our NCP estimates. We have therefore updated the uncertainty in NCP estimates in the results section.

#### The text has been revised as follows:

"A different source of uncertainty in NCP and new production estimates come from measurement uncertainty, which propagates into calculated values. However, for new production, the variability associated with measurement uncertainty is negligible compared to the natural variability. The measurement uncertainty for the NO3 sensor given by the manufacturer (Satlantic ISUS V3 sensor) was 2 µmolkg kg<sup>1</sup>. This gives a propagated probable error in the  $NO3_{MLD}$  estimates of < 0.05 mol N m<sup>-2</sup> (calculated from the square root of the sum of squared errors), which is considered negligible compared to the uncertainty associated with monthly variation of  $\pm 0.14$  mol N m<sup>-2</sup>. Therefore we present the uncertainty related to monthly variation in NO3<sub>MLD</sub>. Conversely, the total uncertainty associated with monthly variations in NCP<sub>MLD</sub> was  $\pm 0.27$  mol C m<sup>-2</sup>, which is smaller than the total error associated with the calculation of DIC from estimated TA and measured fCO2 of  $\pm 0.85$  mol C m<sup>-2</sup>. The latter was determined by propagation using the method described in Dickson and Riley (1978), together with the errors in the estimated TA values ( $\pm 6.4 \mu$ molkg<sup>-1</sup>; Lee et al. 2006) and measured pCO2 (±2 µatm; Wanninkhof et al., 2013). We have therefore chosen to present the propagated probable error as a conservative estimate of the uncertainty in our NCP estimates.

## p.5178 line 11 Please include justification/reference for why all particles are assumed to have a sinking speed of 100 m d<sup>-1</sup>.

Observations at the PAP observatory have estimated the particle sinking speed as between 60 and 180 m day<sup>-1</sup> (Riley et al., 2012; Villa-Alfageme et al., 2014). The value we choose of 100 m day-1 falls in the middle of this range. Faster particle sinking speeds would result in a smaller source region and vice versa for slower sinking particles.

The text has been revised as follows: "..., which is chosen because it falls in the middle of the range observed for particle sinking speeds at the PAP observatory of between 60 and 180 m day<sup>-1</sup> (Riley et al., 2012; Villa-Alfageme et al., 2014)..."

# p.5177 line 4 and figure 3 How can mixing be assumed negligible? Kortzinger et al. (2008) demonstrate that mixing plays an important role in increasing NO3 and DIC concentrations in the mixed layer during winter months at the PAP site. Using the method presented, the positive NCP and new production values in figure 3 are not necessarily solely due to biological drawdown, but are actually where the biological drawdown exceeds mixing and remineralisation. This assumption should be discussed further, and made clearer to the reader.

This is a valid comment, and an issue we discussed frequently in the preparation of this manuscript. It is stated in Sect. 2.3 that the monthly changes in DIC and NO<sub>3</sub> can be attributed to changes caused by air-sea gas exchange (for DIC), physical mixing processes and biological drawdown. The physical mixing processes, such as vertical entrainment, diffusion and advection are difficult to account for without proper measurements. In Kortzinger et al. (2008) they also assume that the contribution from these three mixing processes are small and negligible in the calculation of NCP, but acknowledges that a "full mixed layer budget cannot be constructed". Only a simplified budget is possible, under certain limitations and for restricted periods. We have followed the same rationale, calculating NCP and new production for the period when the MLD is stable and where biological drawdown is believed to play a dominating role in monthly changes in DIC and NO<sub>3</sub>. We do, however, acknowledge the limitations in this approach, and will elaborate on the uncertainties associated with mixing in the manuscript.

The manuscript has been revised as follows:

Physical mixing processes, such as vertical entrainment, diffusion and advection, will to some degree contribute to monthly DIC changes, however are difficult to quantify without information on vertical and horizontal gradients. Following the approach by Kortzinger et al. (2008) we have performed a simplified calculation of seasonal NCP and new production for the summer period when the mixed layer is relatively stable and the biological drawdown in DIC (and NO<sub>3</sub>) is strong. Therefore, the contribution of  $\Delta DIC_{mix}$  was assumed negligible, and  $\Delta DIC_{BP}$  was assumed to be largely determined by NCP (excluding the effect of calcification).

Figure 3. Why are the productive periods (grey shaded) different for NO3MLD and NCPMLD? Using the MLD it should be possible to determine if the mixed layer is deepening, and therefore give an indication of when high concentrations of DIC and nutrients are being entrained in to the mixed layer. The summer months before the mixed layer deepens may be a better period to calculate NCP and new production The period NO3<sub>MLD</sub> and NCP<sub>MLD</sub> are positive, is when there is a net decrease in NO<sub>3</sub> or DIC concentrations, respectively, in the mixed layer caused by biological drawdown. As can be seen in the monthly climatology in Fig. 2, DIC concentrations starts increasing from August, which causes the sign of the NCP<sub>MLD</sub> to change, from positive to negative. NO<sub>3</sub> concentration has a very small decrease from July to August, before there is a small increase from August to September. Therefore the NO3<sub>MLD</sub> and NCP<sub>MLD</sub> is negative in September, one month later that for NCP<sub>MLD</sub>. These months with very small changes in concentrations have error bars that are almost as large as the NO3<sub>MLD</sub> and NCP<sub>MLD</sub> itself, and it is therefore little use in speculating

into a reason for the difference in timing of negative values.

The MLD climatology is also shown in Fig. 2, and the MLD shallows rapidly from around 250m in March to 50m in May. The MLD remains fairly stable at 50m until October. We chose to use the difference in MLD between consecutive months, and not to integrate over a fixed time period. We find the strongest  $NCP_{MLD}$  and  $NO3_{MLD}$  before the onset of shallow stratification in summer, however believe it gives value to show the monthly changes, and the stabile MLD and nutrient depletion in summer indicates that there is not a large contribution from entrainment during these months.

## Figure 3. Please include a table of the different years used to calculate the monthly values and the inter-annual variability. This would be useful to the reader and might also explain why some of the error bars are so small in the figure.

We have compiled a table (se below) with monthly mean, SD and number of samples (n) for each year we have data (taking the years vs. months would mean that the table is very large). The means and SDs are identical to the values given in Fig. 3, and from this table it is clear that for most months we have data from 2-4 different years. This is also described on p. 5176 l. 3-16, where the method of calculating the NO3<sub>MLD</sub> and NCP<sub>MLD</sub> is explained. However, one small error was discovered in the compilation of this table. For NCP<sub>MLD</sub> there is only data from one year (2004) in March and April, and therefore we cannot calculate the SD, and consequently there is no error bar in Fig. 3. This is now explicitly described in the revised manuscript. In addition, we added which months are lacking error bars in Fig. 3 in parenthesis in the text:

"However, for most months of the year data from at least two to four different years were available, and NCP<sub>MLD</sub> and NO3<sub>MLD</sub> was estimated two to four times for each of these months. This allowed the quantification of the average seasonal NCP<sub>MLD</sub> and NO3<sub>MLD</sub> and variability (expressed as one standard deviation; SD) from the PAP time series. With the exception of March and April for NCP<sub>MLD</sub>, when there was only data for one year (2004) and consequently SD could not be calculated (error bars missing in Fig. 3). The monthly changes in equations 3 and 4 (below) were computed in a circular manner i.e. the change in the 12<sup>th</sup> month is the difference January minus December. For NO3<sub>MLD</sub> there was only data for one year in December, and consequently the SD could not be calculated. For NCP<sub>MLD</sub> there was data from two different years in December and three years in January, however they coincided only for 2004. Therefore, the variability could not be determined for December for either NCP<sub>MLD</sub> and NO3<sub>MLD</sub> (error bar missing for December in Fig. 3)."

In addition, we add which months are lacking error bars in the caption to Fig. 3: "Figure 3. Monthly changes in MLD integrated NO<sub>3</sub><sup>-</sup> (top) and NCP (bottom) with  $\pm$  1 SD (vertical bars). The shaded grey area indicates the months over which the seasonal new production (0.37  $\pm$  0.14 mol N m<sup>-2</sup>) and NCP (4.57  $\pm$  0.27 mol C m<sup>-2</sup>) were calculated. For NCP<sub>MLD</sub> no SD could be calculated in March and April, and SD is also lacking for December for both NCP<sub>MLD</sub> and NO3<sub>MLD</sub> (see Sect. 2.3 for calculations)."

Seeing as how the information in the table is identical to Fig. 3, and we have expanded on the which months there are no error bars due to missing data we feel that adding this table to the revised manuscript would not justify the extra space this would require.

	Jan	feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Okt	Nov	Des
NO3 <sub>MLD</sub>												
	-								-	-	-	-
Mean	0,179	0,126	0,107	0,064	0,031	0,029	0,016	0,002	0,013	0,161	0,070	0,142
SD	0,372	0,072	0,109	0,032	0,001	0,017	0,008	0,001	0,012	0,109	0,229	NA
n	3	3	2	2	2	2	2	2	3	3	3	1

NCPMLD												
	-							-	-	-	-	-
Mean	0,713	0,294	2,351	0,768	0,667	0,265	0,220	0,202	0,261	0,934	1,463	1,532
SD	0,822	0,267	NA	NA	0,010	0,041	0,027	0,153	0,244	0,404	1,072	1,065
n	3	3	1	1	2	3	3	4	4	4	4	2

## p.5192 figure 1. The MLD looks like it goes shallower than 30 m during the summer, would this influence your calculations of NCP and new production within the mixed layer as your sensors are at 30 m? Please clarify this and discuss if necessary.

It is the difference in MLD between consecutive months that is used to calculate the NCP<sub>MLD</sub> and NO3<sub>MLD</sub>, and from May-August this is shallower than 30m (23.7m, 17.1m, 16.3m and 21.3m, respectively). When the MLD is shallower than the sensors measuring NO<sub>3</sub><sup>-</sup> and DIC concentrations, this could imply that the drawdown from biological activity within the mixed layer is stronger (and thereby reaches lower concentrations) than what is actually measured by the sensors at 30m. However, the NO<sub>3</sub><sup>-</sup> concentrations in the mixed layer could become depleted, and thereby limit further biological drawdown. On the other side, the biological drawdown could be stronger at 30m than in the mixed layer due to more supply of nutrients than what is available within the mixed layer. Which of these processes would be dominating at the PAP observatory is impossible to determine without the nutrient data. The former process could imply that our estimates of NCP and new production are on the low side (because additional drawdown takes place within the mixed layer due to more readily available nutrients. This is an additional source of uncertainty, and we have added a short description of this in the discussion in Sect 4.1:

"... during this period of rapid stratification. From May to August the MLD is slightly shallower than the depth of the sensors at 30m, which could influence our estimates of NCP and new production. This uncertainty is impossible to quantify without measurements from within the mixed layer, although the effect is believed to be minor. It could potentially overestimate biological production due to more readily available nutrients at 30m than within the mixed layer, or lead to an underestimate due to additional drawdown within the mixed layer. ..."

Figure 3 and figure 4. It would be beneficial to the reader if the units of CO2 flux, NCP, new production, Satellite derived NPP were all in mmol  $m^{-2} d^{-1}$  so that they are more easily comparable. Please make units consistent throughout the manuscript. We agree with the comment by the reviewer and have converted the units in Fig. 4 to mmol  $m^{-2} d^{-1}$ .

The revised Fig 4 is shown below; here we have also included the other issues brought up by the reviewer regarding this figure:



# p.5199 Figure 4. Please add a time dimension to this plot, it is difficult to trace the particle with time without knowing which month the coloured dots are representing. It may also be useful to include a 100km box around the PAP site for reference, and to demonstrate that the source region is outside of the box in some years.

Time labels have been added to mark the start and end of data for each year in Fig. 4 (se above). We did experiment with adding a 100 km box around the PAP site, however there is all ready a lot of information on this plot and unfortunately the box made the figure too cluttered and difficult to interpret for the reader.

The caption to Fig. 4 has been revised as follows: "... The start and end dates of the data for each year is indicated in the panels".

#### **Technical corrections:**

## p.5171 line 1 Is nitrate being measured, or nitrate + nitrite as it is in Hartman et al. 2012. On first use of NO3 please clarify this within the manuscript.

It is nitrate+nitrite, and this has been clarified in the revised text.

#### p.5171 line 15 closing brackets missing

The text has been revised accordingly.

#### p.5196 Figure 2. Add '(black dots)' to legend

The text has been revised accordingly, and now reads: "Monthly climatology (black dots)..."

## p.5199 Figure 4. Make text on axes bigger, and add label to the colour bar including the units

See the revised Fig. 4 above. The font size for the axis labels has been increased, and a label has been added to the top of the figure with variable name and units (not under the color bar due to space constraints).

### Author comment to Anonymous Referee #2

We thank anonymous Referee #1 for his/her constructive criticism and valuable comments. In the following we address the points raised, with referee comments in boldface and author responses in normal typeface.

#### **General comment:**

Frigstad et al. present observations from the Porcupine Abyssal Plain (PAP) time series station / observatory and related interpretation. The dataset presented covers up to a decade of data of observations from automated devices in the surface mixed layer (euphotic zone) and from sediment traps at 3000m. The PAP site is one of the very few open ocean time series sites outside oligotrophic waters. Though parts of the data have been published earlier this is a timely overview, providing also interesting additional analysis of the combined dataset. Using data from Argo, remote sensing and ocean circulation models this, by nature, spatially limited data are set into a wider context. Running such a site and putting such a dataset together is clearly a significant effort of the group lead by the senior author, Richard Lampitt. I recommend publication with minor corrections as indicated below. My largest concern is related to the interpretation of particle tracking, NPP and flux. See the comments on section 2.4, 4.2 below. We thank the reviewer for the positive comment, and have addressed the issue regarding particle tracking in the response below.

#### **Specific comments:**

Abstract

p5170: The Redfield C:N ratio is 6.6. The observed C:N ratio of NCP was 12, I suggest to not confuse younger readers and the speak about 'C:N-ratio of 12' and not to refer to Redfield here.

We agree with the reviewer, and have revised the text accordingly: "The C:N ratio was high  $(12) \dots$ "

Introduction

**p 5170, l 26 (and elsewhere): delete 'full depth'** We have deleted "full depth" throughout the revised text.

p5171, l 9-10: The Sabine reference for the phrase 'biological carbon pump is key to understanding the global carbon cycle' is not justified. In the last paragraph Sabine speaks about potential biological feedbacks to OA in a very general sense. Hence you shouldn't use that reference here. You should refer to papers that give evidence to this statement and not just use such a phrase themselves in either the intro or the outlook! The Falkowski reference is well suited here, perhaps refer to Volk and Hofferts centennial paper in addition, or some significant post-1998 overview paper. We have deleted the Sabine reference in the revised text, and the sentence now reads: "and therefore quantifying the biological carbon pump is key in understanding the global carbon cycle (Falkowski et al. 1998)."

122-24: Is it really the 'multitude of methods ..." that lead to a poor understanding of NCP...? I understand Quay rather in the following sence 'Unfortunately, there are only a few sites where multiple NCP methods have been compared (e.g., JGOFS study sites, BATS and ALOHA time series sites).' (p2). Multiple methods may be rather an advantage, in the absence of a gold standard'.

We agree that this sentence can be misunderstood, and have revised the text to: "It can be challenging to compare between techniques and there is ..."

#### p5171, l24 - p 5172, l6 may be shortened

We go into a bit of detail explaining the differences in NCP and export flux in this paragraph. We felt it was important to explain these differences, because the study computes both NCP and export flux, and compare the two variables in the results and discussion sections. We therefore believe that it is important to give the reader some background, and that these sentences can be justified.

## p 5172, l 13-15: Is the correct reference for Lampitt et al 2018 given? From my memory (but the senior author should know better), the Royal Society paper from the same year is referred to here, right?

The reviewer is correct, and we have inserted the correct reference and updated the reference list.

# I 21-24: suggest to write: 'transfer efficiency has often been used to describe the efficiency'. Rational: a) the POC based metric ignores DOM, b) see papers by Marinov and co-authors Overall, DOM as a pathway is ignored completely in the paper. Its role in sequestration may be less understood, but you might want to mention this pathway (and your ignorance of it in the analysis) at least once in the intro.

The manuscript has been revised as suggested: "... and has often been used to describe the efficiency of the biological pump ..."

It is noted in p 5171, 127 - p 5172, 1 1 that one of the differences between NCP (as calculated in this study) and export flux (calculated as POC flux at a nominal depth), is that the NCP estimate will include the contribution of DOC. However, have added to the revised manuscript that the POC-based metric of calculating export flux and transport efficiency does not include the DOM pathway. Revised text on p. 5172, 1 23: "It should be noted that the POC-based metric of calculating export flux and transfer efficiency does not include the contribution from DOC."

Data and methods **p 5173, l24: rewrite: "(2010). Briefly …"** The text has been revised accordingly.

# Same paragraph: Perhaps mention at least two more details: a) why is formaline addition not an issue for POC measurements (with reference to a study that gives respective evidence). b) what about losses of POC to the supernatant in the cup until splitting of samples, losses e.g. to DOC (see e.g. Kähler and Bauerfeind, L&O, 2001). This is (evidently ?) no issue in your traps?

a) We use analytical grade (AnalR or NORMAOUR) formalin as directed by the JGOFS protocols for preservation. There are numerous unpublished studies that show that somewhat surprisingly formalin does not affect POC. The most cited paper (Knauer et al 1984) indicated some losses.

b) Kähler and Bauerfeind worked with shallow sediment traps, which are much more susceptible to swimmer contamination. At PAP traps are at 3000m and 100mab and have few swimmers therefore negligible DOC leaching.

p5175, l23-24: I am not really sure about the meaningfulness of Lee's T-S to Alk relationship. In particular the T-part. See e.g. Friis et al., 2003, GRL. In your data, (Fig. 2) what is driving the seaonality of ALK? T or S? How sensitive is your DIC seasonality to the computed ALK. What if you assume no seasonality of ALK, e.g. by taking the annual mean of your computed ALK together with your seasonally varying pCO2 data? Hopefully, that gives almost identical DIC values, compared to the presented ones. Please check into this.

This is a very relevant issue raised by the reviewer. In Lee's T-S to Alk relashionship, the

salinity coefficients are several orders larger than those of the temperature-part. Thus, the seasonality is driven mainly by salinity. To verify this, we have recalculated Alk from salinity alone (following Nondal et al. (2009)), and there was still seasonal variation in calculated Alk values. Furthermore, this change in the calculation of Alk had only a negligible effect on the seasonal variation in DIC values (figure below).



For this reason, and since Lee originally included temperature to account for the nutrient cycle, we chose to keep Alk values obtained with Lee's relationship. We also note that the main issue in Friis et al. (2003) concerned the use of zero intercept during the normalization of Alk values to constant salinity. This normalization is not used in this manuscript.

# p5176, l 6 (and elsewhere!!) Kon"rzinger has an o-Umlaut, also Kan"hler has an a-Umlaut. Please check the ms carefully for correct spelling of authors! Go back to the original papers to check, if needed.

The text has been revised accordingly.

## p5176, l 24-27: You should include the error from ALK-S-T in your error budget, and refer to what I proposed above in the text.

This is the same issue raised by reviewer#1, and the text has been revised to include the error from calculated Alk (from temperature and salinity) instead of measured Alk as in the previous version of the manuscript:

"... which is smaller than the total error associated with the calculation of DIC from estimated TA and measured fCO2 of  $\pm 0.85$  mol C m<sup>-2</sup>. The latter was determined by propagation using the method described in Dickson and Riley (1978), together with the errors in the estimated TA values ( $\pm 6.4 \mu$ molkg<sup>-1</sup>; Lee et al. 2006) and measured pCO2 ( $\pm 2 \mu$ atm; Wanninkhof et al., 2013)."

p 5177, 14: 'contribution from delDICmix was assumed negligible': can it be? If I recall correctly Gruber et al. 1998 (DSR ?) used 13C-CO2-data at Bermuda to constrain the role of mixing to a seasonal surface ocean DIC budged. There it was important, I think. I suggest, that you at least discuss this limitation of your estimate briefly, in particular in terms of sign for C:N – NCP estimate, and mention the Gruber and related studies. This is the same issue as raised by reviewer#1, and we repeat the response given on this issue: It is stated in Sect. 2.3 that the monthly changes in DIC and NO<sub>3</sub> can be attributed to changes caused by air-sea gas exchange (for DIC), physical mixing processes and biological drawdown. The physical mixing processes, such as vertical entrainment, diffusion and

advection are difficult to account for without proper measurements. In Kortzinger et al. (2008) they also assume that the contribution from these three mixing processes are small and negligible in the calculation of NCP, but acknowledges that a "full mixed layer budget cannot be constructed". Only a simplified budget is possible, under certain limitations and for restricted periods. We have followed the same rationale, calculating NCP and new production for the period when the MLD is stable and where biological drawdown is believed to play a dominating role in monthly changes in DIC and NO<sub>3</sub>. We do, however, acknowledge the limitations in this approach and will elaborate on the uncertainties associated with mixing in the manuscript.

#### The manuscript has been revised as follows:

Physical mixing processes, such as vertical entrainment, diffusion and advection, will to some degree contribute to monthly DIC changes, however are difficult to quantify without information on vertical and horizontal gradients. Following the approach by Kortzinger et al. (2008) we have performed a simplified budget calculation for the summer period when the mixed layer is relatively stable and the biological drawdown in DIC (and NO<sub>3</sub>) is strong. Therefore the contribution of  $\Delta DIC_{mix}$  was assumed negligible, and  $\Delta DIC_{BP}$  was assumed to be largely determined by NCP (excluding the effect of calcification).

## 17 vs. p 5176128-29. This is a little unclear. I suggest you rewrite Equ3 to explicitly include the gas exchange term.

The changes in DIC concentrations caused by air-sea gas exchange are calculated from the air-sea flux and MLD in Eq. 2. We have revised the text to show how this is included in the terms of Eq. 3 (the contribution of air-sea gas exchange is added to the DIC concentrations because the flux is positive throughout the year):

"... corrected for the effects of air-sea gas exchange  $(\Delta DIC^{GasCorr} = \Delta DIC_{obs} + \Delta DIC_{gas})$ "

## l 21-23: Please clarify in the text whether you computed NPP, or downloaded it from the web site.

The NPP data were downloaded from the website listed in Table 1, and this is already stated in the text. We have clearified this sentence:

"The NPP data were downloaded ... ".

Section 2.4: The particle tracking analysis is done here much better than in some older papers of the senior author which used moored current meters. This is acknowledged by the reviewer. However, here and also later in the paper, you seem to take the transports in the model to be fully consistent with the real ocean patterns and distribution of NPP as seen by the satellite. Why should that be the case? The best you may hope for, I think, is that the applied physical model has the right statistics of transports compared to the real ocean. Whether the eddies (etc.) are at the right place at the right time in the model vs. the real ocean is not known. Hence, the combination of particle tracking, remote sensing, and the deep traps stands on somewhat slippery ground. You need to mention and discuss that – unless you can provide hard remote sensing evidence (e.g altimetry, sst, sss patterns) supporting that your model behaves perfectly in that sense. Recently, Jamie Wilson and co-authors had a very nice paper (currently in review in BGD, I think) demonstrating how deficient similar combinations of models and reality can be. To demonstrate that the NEMO model produces currents that are consistent with observations, we have plotted below the surface currents (geostrophic + Ekman) derived from satellite altimetry and wind data (downloaded from http://www.oscar.noaa.gov/). The mean observed current vectors (top panels) and speed (lower panels) for 2008 are plotted alongside the modelled currents in the figure below. The model reproduces well the main features of the circulation in the region, i.e. the band of strong northeastward currents in the NW quadrant of the domain and the relatively quiescent SE quadrant. The magnitude of the currents are also in a very similar range in the model and observations. Note that satellite observations can necessarily only supply information on the surface currents and so the

analysis we present in the manuscript would not be possible without relying on modelled currents. The paper mentioned by the reviewer by Wilson et al. discusses the issues of accurate model circulation in the context of thousand year spin-ups and attempts to reproduce the 3-D distribution of phosphate in the oceans, i.e. very different time and space scales than we consider here.



## p 5179, l 24-25: Could you explain a little further from your data why there is no seasonal signal in CO2-fluxes?

The air-sea CO<sub>2</sub> flux does not show corresponding winter to summer variations as for example pCO<sub>2</sub> in Fig. 2, largely because of the balancing effect of the seasonal cycles in pCO<sub>2</sub> and wind speed (U<sub>10</sub>). During spring and summer the reduction in wind speed is compensated by the effect of increased  $\Delta fCO_2$  leading to overall small variations in air-sea flux of CO<sub>2</sub> throughout the year.

## 1 25: delete: 'also'. There is no (causal) relationship between the two issues, I think. p 5181, l 11: rewrite: 'for surface sea-water pCO2 ...'

The text has been revised accordingly: "The sediment fluxes had high..."

#### p 5183, l8: Schneider was not the first to report this. Please check for example papers of Cindy Lee from the early 80s, e.g. Lee and Cronin, 1984 and Wakeham et al. 1984. Please do not cite only convenient references, but also the original literature.

The text has been updated to include the Lee and Cronin (1984) reference: "Studies have shown an increasing C:N of sinking material due to preferential remineralization of nutrients (Schneider et al. 2003; Lee and Cronin 1984) ..."

#### 19: rewrite 'ratio may influence ... by about 20 ppm ...'

The text has been revised accordingly: "ratio may influence atmospheric CO<sub>2</sub> concentrations by about 20 ppm"

## l 17: 'basin was too complex' is awk and not to the point. I guess GS's mixing model did not resolve more than 2 or 3 endmembers? Please check and present carefully.

We agree with reviewer, and deleted this part of the sentence as it is not essential to the argument. The revised text now reads: "However, the deep ocean remineralization rates of Anderson and Sarmiento (1994) did not include the Atlantic Ocean."

## p 5183 l 22 to p 5184, l 9. I think this paragraph can be deleted. You follow an idea that does not work out well, for reasons published by others decades ago.

We agree with reviewer that this paragraph does not give added value to the manuscript, and have deleted it in the revised text.

## p 5184, l 17; '64 and 207' is a little awk, explain with one more sentence why the difference is so large.

The Lampitt et al (2008) is an integrated estimate based on the deployment of the Pelagra sediment trap during summers of 2003 to 2006, while the estimate from Tomalla et al. (2008) is based on a simple sample close to the PAP-site (48.6N;where material was collected using a pump). The samples therefore represent different time (and also spatial scales, due to the location not being identical), and therefore represent the natural variability in POC flux. The different time scales is emphasized in the revised text:

"determined to be in the range between 64 and 207 mg C m<sup>-2</sup> d<sup>-1</sup> (based on measurements from a single cruise and long-time trap data; Lampitt et al. 2008, Thomalla et al. 2008)."

## 121: is De La Rocha and Passow the appropriate reference for 'export ratio'. This term is much longer in use! Also the reference of that paper given reference section is not complete!

We have changed the reference in this sentence to Dugdale and Goering (1967), which is the reference given in the introduction where the term is introduced. The revised text now reads: "... divided by the NPP (Dugdale & Goering 1967)."

#### 1 22: '115', please give error bar of your mean value

The standard deviation of the literature values given in the text is  $\pm$  61 mg C m<sup>-2</sup> d<sup>-1</sup>. This uncertainty is added to the revised text:

"Using an average of the above values for POC flux out of the surface layer of  $115 \pm 61$  mg C m<sup>-2</sup> d<sup>-1</sup>".

## Section 4.2 & Conclusion. See my comments on section 2.4. You need to discuss the issue stated above.

Se response to Sect. 2.4 above.

## References: See my comment of on" and an" for Korzinger, Kahler, (but Koeve is correct,

:-)

These references have been corrected in the reference list as well

#### Overall, I enjoyed reading the paper.

Thank you!

#### **References (not cited in article):**

Kähler P, Bauerfeind E. Organic particles in a shallow sediment trap: substantial loss to the dissolved phase. Limnol Oceanogr 2001, 46:719–723

Knauer, George A.; Karl, David M.; Martin, John H.; Hunter, Craig N. In situ effects of selected preservatives on total carbon, nitrogen and metals collected in sediment traps. Journal of Marine Research, Volume 42, Number 2, May 1984, pp. 445-462(18) Nondal, Gisle; Bellerby, Richard G. J.; Olsen, A; Johannessen, T. Optimal evaluation of the surface ocean CO<sub>2</sub> system in the northern Atlantic using data from voluntary observing ships. Limnology and Oceanography: Methods. 7, 2009, 109-118

Author comment to Anonymous Referee #3

We thank anonymous Referee #3 for his/her constructive criticism and valuable comments. In the following we address the points raised, with referee comments in boldface and author responses in normal typeface.

#### **General comments**

This study provides a decade long time series data set on hydrography, biogeochemistry and sediment traps for the PAP site in the North Atlantic. Results are valuable for understanding the present and future functioning of the biological pump in the PAP site. They also suggest that there might be changes in export as the PAP site transitions into subpolar conditions. However there is little discussion on the actual implications of the results in that context. Although I just have some minor comments on the ms, I recommend the authors to further develop the implications of their study. We thank the reviewer for the positive comment, and have expanded on the potential changes in export regime at the PAP site in the revised manuscript.

#### Abstract

Line 10 please specify that a Redfield ratio of 12 is higher than the expected value of 6.6. The text has been revised accordingly: "The C:N ratio was high (12) compared to the Redfield ratio (6.6), ..."

#### Introduction

I am surely not well informed on the dynamics of the PAP site but I have trouble with the sentence "It is at or near the boundary between the sub-polar and sub-tropical gyres of the North Atlantic". Is it or is it not? And if both are true how is it explained? This is important for data interpretation.

The text has been changed to: "The PAP observatory is near the boundary between the subpolar and sub-tropical gyres of the North Atlantic". Please also see figure 8 of Henson et al. (2009).

## Slide 5171 line 1 (and throughout the text) please include the correct formula of nitrate (NO3-).

The formula has been corrected to NO<sub>3</sub><sup>-</sup> throughout the manuscript.

# Lines 10 to 15: These are valuable motivations for the study but I though you already named those before. Please regroup your arguments at the end of the introduction section for clarity.

We agree with the reviewer, and have deleted the sentence in the revised manuscript, as this argument is already given at the end of the introduction section. The revised text now reads: "global carbon cycle (Falkowski et al. 1998). These factors are currently not fully understood, so it is therefore difficult to predict how they will respond to climate change (e.g. Passow & Carlson 2012)."

#### Line 23: "multitude" is vague and you cite only one reference. Please detail or rephrase.

This sentence has been revised in response to a comment made by reviewer#2, and now reads: "It can be challenging to compare between these techniques, and there is ... ".

## Slide 5172: consider giving acronyms to concepts such as transfer efficiency, new production, sequestration flux and export flux.

We considered doing this in the preparation of the manuscript, however we believe that not using too many acronyms in the text makes it easier to follow for the reader.

#### **Materials and Methods**

Section 2.1: I think details are missing here. For instance, what sensors for nitrate were

# used during the last 10 years? What is the detection limit? I understand that this information can be found in a variety of papers but it is best not to force the reader to look for sparse information. A Table sumarising limits of detection, techniques and periods of use can be useful here.

We feel that adding a table containing this information would take up a lot of space, however agree that more information could be beneficial, especially on  $NO_3^-$  measurements (for which the instruments changed over time). We have added the text below to the manuscript. Regarding detection limits, both the methods below claim to measure zero  $NO_3^-$ . However, PAP is not an oligotrophic site, so this should not be an issue in our data.

Text added to the revised manuscript: "NAS nutrient analysers (Envirotech LLC) were used in the period from 2002 until 2012, which measures  $NO_3^-$  plus  $NO_2^-$  (hereafter referred to as  $NO_3^-$ ) with a precision of ±0.05 umol  $I^{-1}$ . The NAS was used alongside an optical nitrate sensor from 2009 to 2012, after that time all  $NO_3^-$  measurements were made with optical sensors (Satlantic ISUS), with a precision of 2umol  $I^{-1}$ . For the remaining variables a Seabird MicroCAT was used for temperature and salinity ...".

## Slide 5176 line 9 "an explanation for December to be followed shortly". Delete. One supposes you are going to explain this later.

This parenthesis has been deleted from the revised text.

#### Slide 5177: Please justify your assumption of negligible mixing.

This issue was raised by the other reviewers, and we have expanded on the contribution from mixing in the manuscript.

It is stated in Sect. 2.3 that the monthly changes in DIC and NO<sub>3</sub> can be attributed to changes caused by air-sea gas exchange (for DIC), physical mixing processes and biological drawdown. The physical mixing processes, such as vertical entrainment, diffusion and advection are difficult to account for without proper measurements. In Kortzinger et al. (2008) they also assume that the contribution from these three mixing processes are small and negligible in the calculation of NCP, but acknowledges that a "full mixed layer budget cannot be constructed". Only a simplified budget is possible, under certain limitations and for restricted periods. We have followed the same rationale, calculating NCP and new production for the period when the MLD is stable and where biological drawdown is believed to play a dominating role in monthly changes in DIC and NO<sub>3</sub>. We do, however, acknowledge the limitations in this approach and will elaborate on the uncertainties associated with mixing in the manuscript.

#### The manuscript has been revised as follows:

Physical mixing processes, such as vertical entrainment, diffusion and advection, will to some degree contribute to monthly DIC changes, however are difficult to quantify without information on vertical and horizontal gradients. Following the approach by Kortzinger et al. (2008) we have performed a simplified budget calculation for the summer period when the mixed layer is relatively stable and the biological drawdown in DIC (and NO<sub>3</sub>) is strong. Therefore the contribution of  $\Delta DIC_{mix}$  was assumed negligible, and  $\Delta DIC_{BP}$  was assumed to be largely determined by NCP (excluding the effect of calcification).

#### Slide 5178 line 11: Why did you use 100 m d-1 for all particles?

This same issue was raised by reviewer#1. Observations at the PAP site have estimated the particle sinking speed as between 60 and 180 m day<sup>-1</sup> (Riley et al., 2012; Villa-Alfageme et al., 2014). The value we choose of 100 m day-1 falls squarely in the middle of this range. Faster particle sinking speeds would result in a smaller source region and vice versa for slower sinking particles. The manuscript has been revised: "..., which is chosen because it

falls in the middle of the range observed for particle sinking speeds at the PAP observatory of between 60 and 180 m day<sup>-1</sup> (Riley et al., 2012; Villa-Alfageme et al., 2014)..."

## Results. Slide 5180 line 14. Particle tracking shows considerable interannual variability. You don't mentioned mesoscale variability as a cause of this variability. Please comment.

This is a valid comment, and mesoscale variability could contribute to the variability in source location of material arriving at the sediment trap, both between and within years. This has been added to the discussion, in Sect. 4.2

Revised text at p. 5185, 19: "... reflecting the seasonal cycle. Mesoscale variability will also contribute to the variability in source location of particles, both between and within years."

#### Discussion

Slide 5182 lines 20-23. Maybe a table compiling previous N:C estimate would help here. We believe a table giving these values would take up too much space, especially since comparing with previous C:N estimates is not a central part of this study.

# Line 23: N2 fixation could be (and references will show you that it is) important in the North Atlantic. Assuming the opposite can be misleading. Please discuss its possible effect on your observed C:N. Also discuss your choice of assuming negligible N2 fixation.

Although  $N_2$  fixation is known to be significant, especially in the subtropical North Atlantic gyres, many studies show that it is not likely to be so at the latitude of the PAP observatory. Figure 2a of Moore et al. (2009) has a transect over the PAP observatory (unfortunately, the plot is cut off at 40N), and shows that  $N_2$  fixation is close to zero from 30-40N, a pattern that is likely to continue at higher latitudes. Reynolds et al. (2007) suggest that N2 fixation plays a minor role in supplying N to the "northern subtropical gyre", the northern boundary of which they call 40N.

The text has been revised as follows: "Studies have found  $N_2$  fixation to be low or zero at latitudes >40°N (PAP observatory is at 49°N) in the North Atlantic (Moore et al. 2009, Reynolds et al. 2007), therefore  $N_2$  fixation is not thought to be an important factor for C overconsumption. Therefore ... ".

## Slide 5185 line 20. Transfer efficiency is already defined in the text. Also, please discuss a little bit more the implications of the 4% you report here.

We have deleted the definition of transfer efficiency in this sentence.

We have also added more description of what the relatively low transfer efficiency found at the PAP observatory implies in the revised text: "region by Henson et al. (2012), indicative of regions where the available nitrate is not fully consumed and with strong seasonal mixing."

Slide 5186 last paragraph of discussion. This is a speculative statement and I don't think it's a good way to finish your paper. Either you discuss this point in more detail or you replace it with a more conclusive statement that is actually supported by the data. We believe it is relevant to mention that the export regime at the PAP observatory (with high export ratio and low transfer efficiency) could alter as a result of climate change. We have elaborated on the causes of the changes and potential effects.

The final paragraph of the discussion now reads: "The PAP observatory currently sits near the boundary between the sub-polar and sub-tropical gyres of the North Atlantic. Seasonably variable areas, like the sub-polar region, are thought to export a higher fraction of labile material than sub-tropical regions (Lutz et al. 2007). As climate change is predicted to result in the oligotrophic gyres expanding over the next century (Sarmiento et al. 2004), the PAP observatory will likely transition into more sub-tropical conditions. This could result in more refractory material being exported at the PAP observatory, potentially reversing the pattern we report here with a high export ratio and low transfer efficiency. In addition, a more strongly positive NAO index is predicted due to climate change (Gillett 2003), which is expected to increase diatom abundance at the PAP observatory, and result in reduced organic carbon flux to the deep ocean (Henson et al., 2012a). Although the precise response of the biological carbon pump to climate change is as yet unclear, transition-zone regions between gyres (such as the PAP observatory), could be among the systems that are most strongly affected by climate change (Henson et al. 2013)."

#### Slide 5186, Line 10: Change how?

Into a sub-tropical export regime with more refractory material exported out of the euphotic zone, see above.

#### Figure 5: Please clarify what the axes are.

The caption to Fig. 5 has been revised as follows: "Cross-correlations between sediment trap data (top: volume flux, middle: dry weight, bottom: POC) and NPP in the source regions defined by particle tracking (left) or in a 100 km box around the PAP observatory (right). See Sect. 2.4 for calculations."

- Links between surface productivity and deep ocean particle flux at the 1 2 Porcupine Abyssal Plain (PAP) sustained observatory. 3 H. Frigstad<sup>1,\*</sup>, S. A. Henson<sup>2</sup>, S. E. Hartman<sup>2</sup>, A. M. Omar<sup>1,3</sup>, E. Jeansson<sup>1</sup>, H. Cole<sup>4</sup>, 4 C. Pebody<sup>2</sup> & R. S. Lampitt<sup>2</sup> 5 6 7 [1] {Uni Research Climate, Allégaten 55, 5007 Bergen, Norway} 8 [2] {National Oceanography Centre, Waterfront Campus, European Way, 9 Southampton SO14 3ZH, UK} 10 [3] {Geophysical Institute, University of Bergen, Allégaten 79, 5007 Bergen, 11 Norway} 12 [4] {Ocean and Earth Science, University of Southampton Waterfront Campus, 13 European Way, Southampton, SO14 3ZH, UK} 14 15 [\*] {Now at the Norwegian Environment Agency, Grensesvingen 7, 0661 Oslo} 16 17 Correspondence to: H. Frigstad (helene.frigstad@miljodir.no)
- 19 Abstract

18

- 20 In this study we present hydrography, biogeochemistry and sediment trap
- 21 observations between 2003 and 2012 at Porcupine Abyssal Plain (PAP) sustained
- 22 observatory in the northeast Atlantic. The time series is valuable as it allows for
- 23 investigation of the link between surface productivity and deep ocean carbon flux.
- 24 The region is a perennial sink for CO<sub>2</sub>, with an average uptake of around 1.5 mmol m
- 25 <sup>2</sup> day<sup>-1</sup>. The average monthly drawdowns of inorganic carbon and nitrogen were used
- 26 to quantify the net community production (NCP) and new production, respectively.
- 27 Seasonal NCP and new production were found to be  $4.57 \pm 0.27$  mol C m<sup>-2</sup> and  $0.37 \pm$
- 28 0.14 mol N m<sup>-2</sup>. The <u>C:N</u> ratio was high (12) compared to the Redfield ratio (6.6), and
- 29 the production calculated from carbon was higher than production calculated from
- 30 nitrogen, which is indicative of carbon overconsumption. The export ratio and transfer
- efficiency were 16% and 4%, respectively, and the site thereby showed high flux attenuation. Particle tracking was used to examine the source region of material in t
- 32 attenuation. Particle tracking was used to examine the source region of material in the 33 sediment trap, and there was large variation in source regions, both between and
- within years. There were higher correlations between surface productivity and export
- flux when using the particle-tracking approach, than by comparing with the mean
- productivity in a 100km box around the PAP site. However, the differences in
- 37 correlation coefficients were not significant, and a longer time series is needed to
- draw conclusions on applying particle tracking in sediment trap analyses.
- 39

#### 40 Introduction

- 41 The Porcupine Abyssal Plain (PAP) sustained observatory is situated in the northeast
- 42 Atlantic Ocean (49°N 16.5°W) in a water depth of 4800 m. It is near the boundary
- 43 between the sub-polar and sub-tropical gyres of the North Atlantic (Henson et al.
- 44 2009). A time series of particle flux measurements at 3000 m depth are available back
- to the early 1990's (Lampitt et al. 2010), and since 2003 there has also been a

Field Code Changed

Unknown Field Code Changed

Unknown

Field Code Changed

Helene Frigstad 25.8.15 18.26 Deleted: Redfield

Helene Frigstad 25.8.15 19.48 Deleted: at or near

1 multidisciplinary mooring with instruments at approximately 30 m depth recording hydrography, nitrate (NO3<sup>-</sup>), partial pressure of CO2 (pCO2) and Chlorophyll a (Chl a) Helene Frigstad 25.8.15 18.29 2 Deleted: full depth 3 (Hartman et al. 2012). The simultaneous observations of surface and deep ocean 4 biogeochemistry make this time series ideal to study linkages between surface ocean 5 productivity and deep ocean particle flux. The pathway by which a small fraction (<1%; Martin et al. 1987) of the 6 7 carbon fixed by photosynthesis in the sunlit upper ocean is exported to great depths, 8 thereby constituting a sink for atmospheric  $CO_2$ , is referred to as the biological carbon 9 pump. Carbon sinking to the deep ocean is sequestered on long time scales (100s to 10 1000s of years), and therefore quantifying the biological carbon pump is key in 11 understanding the global carbon cycle (Falkowski et al. 1998). These factors are Helene Frigstad 25.8.15 18.30 12 currently not fully understood, so it is therefore difficult to predict how they will Deleted: Sabine et al. 2004 13 respond to climate change (e.g. Passow & Carlson 2012). Helene Frigstad 25.8.15 20.05 14 Primary production in the surface ocean can be measured by several Deleted: The PAP time series allows an 15 techniques (broadly separated into vitro incubations or changes in bulk properties; investigation of the factors controlling the flux Platt et al. (1989)), however, from the perspective of the oceanic carbon cycle the 16 of carbon. 17 most important rate is the net community production (NCP). NCP is the net primary 18 production (NPP) minus heterotrophic respiration, and represents the sum of the 19 particulate and dissolved organic carbon available for export or utilization by higher 20 trophic levels. NCP is traditionally measured by bottle O2 incubations (Gaarder & Gran 1927), but has also been estimated from oxygen or carbon budgets, <sup>234</sup>Thorium, 21 22 sediment traps and  $O_2/Ar$  ratios. It can be challenging to compare between techniques 23 and there is a poor understanding of NCP rates in many regions of the ocean (Quay et Helene Frigstad 25.8.15 18.31 24 al. 2010). In many studies, the steady-state NCP is equated with the export flux at the Deleted: The multitude of methods and 25 base of the euphotic zone (Platt et al. 1989, Lee 2001, Long et al. 2011, Nevison et al. difficulty in comparing techniques has lead to 26 2012), based on the rationale that NCP is the organic material available for export out 27 of the mixed layer. However, it is not directly comparable to the most common 28 definition of export flux (i.e. the downward flux of POC at a nominal depth) as NCP 29 represents a bulk measurement integrated over the mixed layer and long time scales 30 (and also includes the contribution of DOC to export). The export ratio (ie. Dugdale & 31 Goering 1967) is used to quantify the proportion of the organic material produced that 32 is exported below the euphotic zone, and is often calculated as the flux of POC at 100 33 m divided by the NPP (Henson et al. 2012b). Global estimates of the export ratio range from 10% (Henson et al. 2011) to 40% (Eppley & Peterson 1979), and is well 34 correlated with temperature, and thereby also latitude (Laws et al. 2000). 35 36 New production (Dugdale & Goering (1967) is the production supported by Helene Frigstad 25.8.15 22 37 the input of new nitrogen into the euphotic zone through upwelling and horizontal Deleted: [ 38 mixing, but also by processes such as atmospheric deposition and nitrogen fixation Helene Frig ad 25.8.15 23.18 39 (Sarmiento & Gruber 2006, Gruber 2008). On an annual basis, assuming the system is Deleted: and 40 in steady state, export production is considered equivalent to new production (Eppley 41 & Peterson 1979). 42 From a climate change perspective, the long-term (> 100 years) removal of 43 carbon from the atmosphere is important to quantify, which is often defined as the flux of carbon below 1000 meters (Lampitt et al. 2008a), known as the sequestration 44 45 flux. The sequestration flux is smaller than the export flux out of the euphotic zone or mixed layer, and is ~ 6-25% of the new production based on sediment trap data 46 47 (Berelson 2001, Francois et al. 2002). This large reduction in carbon flux with 48 increasing depth is caused by intensive remineralization of organic material as it sinks 49 through the mesopelagic zone, which is often referred to as flux attenuation (Martin et 50 al. 1987, Steinberg et al. 2008). The ratio of deep POC flux to export flux (POC flux

- 1 at 2000m / POC flux at 100 m), is known as the transfer efficiency and has often been 2 used to describe the efficiency of the biological pump (Francois et al. 2002, De La 3 Rocha & Passow 2012, Henson et al. 2012b). It should be noted that the POC-based 4 metric of calculating export flux and transfer efficiency does not include the 5 contribution from DOC. In this study we present a time series of surface ocean measurements and 6 7 particle trap data from the PAP observatory station from 2003 to 2012. Our aim is 8 twofold; firstly, we will quantify NCP and new production from the average monthly 9 drawdown of dissolved inorganic carbon (DIC) and NO<sub>3</sub><sup>-</sup>, respectively. This allows us 10 to compare these two estimates of surface productivity, and derive export ratios by comparison with satellite NPP estimates and published values of shallow POC flux at 11 12 the PAP observatory. Secondly, we will investigate the link between the production at 13 the surface and particle flux at 3000 meters depth, both by investigating the transfer 14 efficiency and by examining the source location of exported material using particle-
- 15 tracking techniques.
- 16

#### 17 **1. Data and methods**

#### 18 2.1 PAP surface mooring

19 Hydrographical and biogeochemical parameters were measured using data from

- 20 instruments at a nominal 30m depth on a mooring at the PAP observatory (49N,
- 16.5W). The surface mooring was first deployed in July 2003, and more information
- about the time series can be found in Hartman et al. (2012). Due to problems with
- damage to the mooring and/or failure of sensors, there is no or little data between
- 24 2005 and 2010, however, after May 2010 there is good temporal coverage for all
   25 biogeochemical parameters. The data from 2003-2005 have previously been published
- 26 in Körtzinger et al. (2008) and Hartman et al. (2010). The sensors used after 2005 are
- described in Table 1 in Hartman et al. (2012). NAS nutrient analysers (Envirotech
- 28 LLC) were used in the period from 2002 until 2012, which measures  $NO_3^-$  plus  $NO_2^-$
- 29 (hereafter referred to as NO<sub>3</sub>) with a precision of  $\pm 0.05$  umol l<sup>-1</sup>. The NAS was used
- 30 alongside an optical nitrate sensor from 2009 to 2012, after that time all NO<sub>3</sub>=
- 31 measurements were made with optical sensors (Satlantic ISUS), with a precision of
- 32 <u>2umol 1<sup>-1</sup>. For the remaining variables a</u> Seabird MicroCAT was used for temperature
- and salinity, Wetlabs (FLNTUSB) for Chl *a* and PRO-OCEANUS for pCO<sub>2</sub>. The
- 34  $NO_3^-$ , Chl *a* and pCO<sub>2</sub> data have all been quality controlled and calibrated against
- 35 CTD data on cruises to the PAP observatory at deployment and/or recovery of the
- 36 mooring, while this has not always been achieved for temperature and salinity.

#### 37 2.2 Sediment trap

- 38 The sediment trap mooring at the PAP observatory was deployed in the depth range
- 39 3000 to 3200 m, which is around 1800 m above the seabed. The methodology is
- 40 described in Lampitt et al. (2010). Briefly a Parflux sediment trap was used with
- 41 mouth area  $0.5 \text{ m}^2$ , prefilled with hypersaline buffered formalin, following the JGOFS
- 42 protocols. The collection period varied between 2 and 8 weeks depending on the time
- 43 of year and anticipated flux. All fluxes are temporally and spatially integrated, and 44 given in either ml m<sup>-2</sup> d<sup>-1</sup> (volume flux) or mg m<sup>-2</sup> d<sup>-1</sup> (dry weight and Particulate
- 45 Organic Carbon (POC)).

Helene Frigstad 25.8.15 18.38

**Deleted:** describes the efficiency of the biological pump

Unknown Field Code Changed Unknown

Field Code Changed

Helene Frigstad 25.8.15 18.28 Deleted: full depth

#### Helene Frigstad 25.8.15 23.04

#### Deleted: o

Helene Frigstad 25.8.15 20.07

**Deleted:** and the sensors and calibrations used during these deployments will not be discussed further here

Helene Frigstad 25.8.15 20.08

Deleted: , and are

Helene Frigstad 25.8.15 20.10 **Deleted:** Satlantic ISUS for NO<sub>3</sub> (+ nitrite),

Helene Frigstad 25.8.15 18.40 Deleted: , but Helene Frigstad 25.8.15 18.40 Deleted: b

#### 1 2.3 Ancillary data and calculated parameters

2 To interpret and expand on the data from the PAP observatory, the following

3 parameters were used from external data sources (see Table 1): temperature and

4 salinity profiles from Argo floats, atmospheric CO<sub>2</sub> concentration, sea level

5 barometric pressure (SLP), wind speed at 10 m height and satellite derived net

6 primary production (NPP).

7 Temperature and salinity profiles were extracted from the global fields for the 8 PAP observatory (49°N 16.5°W), made available by the Coriolis project 9 (http://www.coriolis.eu.org). The gridded fields use temperature and salinity profiles 10 collected by Argo floats, XBTs, CTD/XCTDs and moorings, and the irregularly sampled data are gridded onto a regularly spaced grid by the statistical objective 11 12 analysis method (Gaillard et al. 2009). The North Atlantic is the region most 13 frequently sampled by Argo floats and has good coverage in both time and space 14 (Gaillard et al. 2009). Here we use monthly averaged temperature and salinity fields 15 for 2002-2009 (delayed mode data), however, after 2010 only near real-time data 16 were available, which has undergone less rigorous quality control than the delayed 17 mode data. In the calculation of carbon parameters (below) the Argo float temperature 18 and salinity at 30 m were used, because of data gaps in the temperature and salinity 19 data from the PAP sensor (referred to as Argo temperature and salinity in text and 20 Figs. 1 and 2) and the lack of consistent calibration with CTD data. Density profiles were calculated using the recently updated standard for seawater properties (TEOS-21 22 10; www.teos-10.org). The mixed layer depth (MLD) was calculated from density 23 profiles using the same global gridded fields used for the temperature and salinity data 24 at 30m. The depth of the mixed layer was defined by a density difference of 0.03 kg25 m<sup>-3</sup> from the density at a reference depth (in this case 10 m to avoid diurnal changes 26 in temperature and salinity at the surface). We followed the algorithm developed by 27 Holte and Talley (2009), which incorporates linear interpolation to estimate the exact 28 depth at which the density difference is crossed. 29 The atmospheric CO<sub>2</sub> concentration measurements were obtained from the

observatory closest to the PAP observatory, the Mace Head land station in Ireland
(53.33N 9.90W) from the Cooperative Atmospheric Data Integration Projects
(GLOBALVIEW-CO2 2012). There were no measurements available after 2010, and
so the annual averaged growth rate in atmospheric CO<sub>2</sub> for marine sites from the
NOAA ESRL data for 2011 (1.69 µmol mol<sup>-1</sup>) and 2012 (2.59 µmol mol<sup>-1</sup>) was added

to the seasonal cycle in CO<sub>2</sub> concentrations at the Mace Head station for 2010
 (http://www.esrl.noaa.gov/gmd/ccgg/trends/global.html). The atmospheric pCO<sub>2</sub> was

 $(http://www.csrinbaa.gov/gnid.cegg/itchas/grobal.ntml). The annosphere <math>peo_2$  was calculated using the Mace Head station  $CO_2$  measurements at barometric pressure (6-

hourly Sea Level Pressure; Table 1) and equilibrium water vapour pressure (from

39 Argo temperature and salinity at 30 m; Table 1).

40 The air-sea  $CO_2$  flux (in mmol m<sup>-2</sup> d<sup>-1</sup>) was calculated from the air-sea p $CO_2$ 41 difference, Argo temperature and salinity (30 m) and wind speed at 10 m height, using 42 the following equation:

43

### 44 $F_{CO2} = k \times K_0 \times (pCO_{2sea} - pCO_{2air})$ 45

(1)

46 where k is the transfer coefficient based on the wind speed-dependent formulation of

47 Nightingale et al. (2000) scaled to the temperature-dependent Schmidt number

48 according to Wanninkhof (1992),  $K_0$  is the CO<sub>2</sub> solubility at in situ temperature and

49 salinity (Argo temperature and salinity at 30 m) after Weiss (1974), while  $pCO_{2sea}$ 

50 and  $pCO_{2air}$  are the CO<sub>2</sub> partial pressures of seawater and air, respectively.

The alkalinity (Alk) was calculated from Argo temperature and salinity (30 1 2 m), following the relationship for the North Atlantic developed by Lee et al. (2006). 3 The dissolved inorganic carbon (DIC) was calculated from Alk and measured pCO<sub>2</sub> 4 using the "seacarb" package (Lavigne & Gattuso 2011), developed for R (R 5 Development Core Team, 2012), using Argo temperature and salinity (30 m) and nutrient concentrations set to zero. The chosen constants were Lueker et al. (2000) for 6 7 K1 and K2, Perez and Fraga (1987) for Kf and the Dickson (1990) constant for Ks, as 8 recommended by Dickson et al. (2007). 9 The seasonal drawdown of DIC and NO<sub>3</sub><sup>-</sup> were used to quantify the 10 NCP and new production, respectively. The gaps in the PAP time series did not allow 11 for examination of the seasonal drawdown on an annual basis (expect for the year of 12 2004, as published by Körtzinger et al. (2008)). However, for most months of the 13 year data from at least two to four different years were available, and NCP<sub>MLD</sub> and 14 NO3<sub>MLD</sub> was estimated two to four times for each of these months. This allowed the quantification of the average seasonal  $NCP_{MLD}$  and  $NO3_{MLD}$  and variability 15 16 (expressed as one standard deviation; SD) from the PAP time series. With the 17 exception of March and April for NCP<sub>MLD</sub>, when there was only data for one year 18 (2004) and consequently SD could not be calculated (error bars missing in Fig. 3). 19 The monthly changes in equations 3 and 4 (below) were computed in a circular manner i.e. the change in the 12<sup>th</sup> month is the difference January minus December. 20 For NO3<sub>MLD</sub> there was only data for one year in December, and consequently the SD 21 22 could not be calculated. For NCP<sub>MLD</sub> there was data from two different years in 23 December and three years in January, however they coincided only for 2004. 24 Therefore, the variability could not be determined for December for either NCP<sub>MLD</sub> and NO3<sub>MLD</sub> (error bar missing for December in Fig. 3), 25 26 A different source of uncertainty in NCP and new production estimates come 27 from measurement uncertainty, which propagates into calculated values. However, for 28 new production, the variability associated with measurement uncertainty is negligible compared to the natural variability. The measurement uncertainty for the NO3 sensor 29 30 given by the manufacturer (Satlantic ISUS V3 sensor) was 2 µmolkg kg<sup>1</sup>. This gives a propagated probable error in the  $NO3_{MLD}$  estimates of  $< 0.05 \text{ mol N m}^{-2}$  (calculated 31 32 from the square root of the sum of squared errors), which is considered negligible compared to the uncertainty associated with monthly variation of  $\pm 0.14$  mol N m<sup>-2</sup> 33 Therefore we present the uncertainty related to monthly variation in  $NO3_{MLD.}$ 34 35 Conversely, the total uncertainty associated with monthly variations in NCP<sub>MLD</sub> was 36  $\pm 0.27$  mol C m<sup>-2</sup>, which is smaller than the total error associated with the calculation of DIC from estimated TA and measured fCO2 of  $\pm 0.85$  mol C m<sup>-2</sup>. The latter was 37 38 determined by propagation using the method described in Dickson and Riley (1978), together with the errors in the estimated TA values ( $\pm 6.4 \mu molkg^{-1}$ ; Lee et al. 2006) 39 40 and measured pCO2 (±2 µatm; Wanninkhof et al., 2013). We have therefore chosen 41 to present the propagated probable error as a conservative estimate of the uncertainty 42 in our NCP estimates, 43 The monthly changes in DIC concentrations  $(\Delta DIC_{obs})$  can be attributed to

44 changes caused by air-sea gas exchange ( $\Delta DIC_{gas}$ ), physical mixing processes  $(\Delta DIC_{mix})$  and biological production  $(\Delta DIC_{BP})$ . The monthly  $(\Delta DIC_{gas})$  (in µmol 45 kg<sup>-1</sup>) can be estimated from the air-sea CO<sub>2</sub> flux ( $F_{co2}$ ) and MLD by the following 46 formulation:

47

48

 $\Delta DIC_{gas} = \frac{F_{co2}}{MLD} \times \frac{365}{12}$ 49

#### Helene Frigstad 25.8.15 23.0 Deleted: o

Helene Frigstad 25.8.15 22.39 Formatted: Default Paragraph Font, Font:Cambria, Norwegian Bokmal

Helene Frigstad 25.8.15 18.00 **Deleted:** 

#### Helene Frigstad 25.8.15 18.00

Deleted: However, for each month of the year data from at least two to four different years were available. Therefore, NCP and new production were estimated two to four times for each of the months January - November (an explanation for December to be followed shortly). This allowed the quantification of the average seasonal NCP and new production and variability (expressed as one standard deviation) from the PAP time series. The monthly changes in equations 3 and 4 (below) were computed in a circular manner i.e. the change in the 12<sup>th</sup> month is the difference January minus December. Data from three different years were available for both of these months, but they coincided only for 2004. Therefore, NCP and new production in December could be estimated only once (based on the 2004 data) and, thus, the variability could not be estimated for this month

#### Helene Frigstad 25.8.15 2<sup>-</sup>

#### Formatted: Not Highlight

Helene Frigstad 25.8.15 17.41

Deleted: An additional source of variability in NCP and new production estimates come from measurement uncertainty, which propagates into calculated values. However, the variability associated with measurement uncertainty is negligible compared to the natural variability. For instance, the measurement uncertainty for the NO3 sensor given by the manufacturer (Satlantic ISUS V3 sensor) was  $\pm 2 \text{ µmol kg}^{-1}$ . This gives a propagated probable error in the NO3<sub>MLD</sub> estimates of < 0.001 mol N m<sup>-2</sup> (calculated from the square root of the sum of squared errors), which is considered negligible compared to the uncertainty associated with monthly variation (see Fig 3). Similarly, using  $\pm 3 \mu$ mol kg<sup>-1</sup> as the uncertainty in DIC and Alk values (e.g., Millero 1995) does not include errors by using Alk calculated from salinity) gives a probable propagated error of < 0.05 mol C m<sup>-2</sup>, which is small compared to the uncertainty from monthly variation in Fig.

5

(2)

10

11

1 2

3

4

Physical mixing processes, such as vertical entrainment, diffusion and advection, will to some degree contribute to monthly DIC changes, however are difficult to quantify without information on vertical and horizontal gradients. Following the approach by Körtzinger et al. (2008) we have performed a simplified calculation of seasonal NCP and new production for the summer period when the mixed layer is relatively stable and the biological drawdown in DIC (and NO3<sup>-</sup>) is strong. Therefore, the contribution of  $\Delta DIC_{mix}$  was assumed negligible, and  $\Delta DIC_{BP}$  was assumed to be largely determined by NCP (excluding the effect of calcification). The monthly NCP integrated over the MLD (NCP<sub>MLD</sub> in mol C m<sup>-2</sup>) was calculated from the monthly changes in DIC corrected for the effects of air-sea gas exchange  $(\Delta DIC^{GasCorr} =$ 12  $\Delta DIC_{obs} + \Delta DIC_{gas})$ 13

16

 $NCP_{MLD} = (\Delta DIC_{m+1}^{GasCorr} - \Delta DIC_{m}^{GasCorr}) \times \frac{(MLD_{m+1} + MLD_{m})}{2}$ (3)

where  $\Delta DIC_{m+1}^{GasCorr} - \Delta DIC_m^{GasCorr}$  is the difference in  $\Delta DIC^{GasCorr}$  between two 17 consecutive months (m and m + 1) and the last term gives the average MLD of the 18 19 two months. Positive values of NCP<sub>MLD</sub> represents net autotrophy (i.e. the months 20 where the biological drawdown of DIC exceeds the DIC released by heterotrophic 21 processes), and the seasonal  $NCP_{MLD}$  can then be calculated as the sum of months 22 with a positive NCP. The same rationale can be applied to the monthly changes in  $NO_3^{-1}$ 23 24 concentrations ( $\Delta NO3$ ), naturally without having to consider the effect of air-sea exchange. The monthly MLD-integrated NO<sub>3</sub><sup>-</sup> changes ( $\Delta NO3$  in mol N m<sup>-2</sup>) were 25 calculated as:

26

27

28 29

30 Summing up the months with a net drawdown in NO<sub>3</sub><sup>2</sup> gives the seasonal new 31 production.

 $NO3_{MLD} = (\Delta NO3_{m+1} - \Delta NO3_m) \times \frac{(MLD_{m+1} + MLD_m)}{2}$ 

32 Net primary production (NPP) was estimated from satellite data using the 33 Vertically Generalised Production Model (Behrenfeld & Falkowski 1997), which 34 requires inputs of chlorophyll concentration, sea surface temperature and

35 photosynthetically available radiation data, here taken from NASA's MODIS Aqua

36 satellite (reprocessing R2012.0). The NPP data were downloaded from the Ocean

Productivity website (see Table 1). 37

#### 38 2.4 Particle tracking and cross-correlations

In addition to estimating the surface origin of particles sinking to the sediment trap 39

40 using a simple 100 km box around the PAP observatory, we also used modelled

41 velocity fields to determine the likely source region. The velocity field (u and v

42 components) was taken from the NEMO model (Madec 2008) run at NOC at 5-day,

43 1/4° resolution for the period 2002-2011. The model has 75 depth levels increasing in

44 thickness with depth, ranging from 1 m near surface to 200 m at 6000 m depth. All

particles reaching the PAP sediment trap at 3000 m depth are assumed to have a 45

sinking speed of 100 m day<sup>-1</sup>, which is chosen because it falls in the middle of the 46 range observed for particle sinking speeds at the PAP observatory of between 60 and 47

48 180 m day<sup>-1</sup> (Riley et al., 2012; Villa-Alfageme et al., 2014). Particles are tracked

Helene Frigstad 25.8.15 18.53 Deleted: D Helene Frigstad 25.8.15 17.4 Deleted: and Helene Frigstad 25.8.15 17.47 Deleted: p

Helene Frigstad 25.8.15 17

Helene Frigstad 25.8.15 18.50

Deleted: (\DIC GasCorr)

Deleted: T

6

(4)

1 backwards in time in 3 dimensions by linear interpolation of the gridded velocity field

2 to the local position of the particle, until they reach the surface (30 days after release).

3 The cross-correlation between the sediment trap data and either NPP in a 100

4 km box around the PAP observatory or in source locations identified by particle

5 tracking, were calculated using the ccf function in R (R Development Core Team,

6 2012). The cross-correlations were performed on monthly anomalies (monthly

7 climatology - observed monthly value), to avoid possible inflation of p-values due to

8 auto-correlation. To test for significant differences between the correlation

9 coefficients the Fisher r-to-z transformation was used (two-tailed test, with two

10 dependent correlations sharing one variable), from the R library "Psych" (Fisher

11 1915, Revelle 2012).

12

#### 13 2. Results

Time series data from 2003 to 2012 from the PAP surface mooring and sediment trap are shown in Fig 1. The temperature and salinity (both PAP sensors and Argo 30 m) varied in the range 12-18 °C and 35.4-35.8, respectively. The mixed layer depths (MLD) were fairly consistent between years, although the winter mixed layer only extended down to ~100 m in 2010. There was a pronounced seasonal drawdown in

pCO<sub>2</sub> (similarly for DIC), with summer values as low as 300  $\mu$ atm during August

2004 and typical winter values between 360-380 µatm. Corresponding seasonal trends

2004 and typical which values between 500-580 path. Corresponding seasonal if 21 were seen for  $NO_3^-$ , with a winter maximum of 10 µmol kg<sup>-1</sup> in March 2004 and

values close to detection limit during summer. The strongest bloom was observed in

June 2011 with Chl *a* concentrations between 3 and 5  $\mu$ g l<sup>-1</sup>, with higher than typical

summer values of around  $2 \ \mu g \ I^{-1}$ . The air-sea CO<sub>2</sub> flux was negative (i.e. oceanic

uptake of  $CO_2$ ) throughout the time period, with an average uptake of around 1.5

26 mmol  $m^{-2}$  day<sup>-1</sup>. There were three years with unusually high sediment fluxes, with

short bursts of high flux during summer in 2004, 2009 and 2012.

28 The monthly climatology (or average seasonal cycle) for temperature showed 29 a seasonal warming of around 5 °C, with very good overlap between the temperatures 30 measured by the PAP sensor and the Argo floats at 30 m (see monthly climatologies 31 in Fig. 2). There was little seasonal variation in salinity, although the Argo float data 32 is generally around 0.05 lower than the salinity measured by the sensors at the PAP 33 observatory. The summer MLD was around 30 m (usually between May and 34 October), and mixing extended down to 250 m depth in winter. The pCO<sub>2</sub> decreased 35 by around 30 µatm from winter values to the summer minimum in August (reduction of around 35 µmol kg<sup>-1</sup> for DIC), while NO<sub>3</sub><sup>-</sup> decreased by around 5 µmol kg<sup>-1</sup> to the 36 37 summer minimum typically found in September. There was a gradual build-up of Chl 38 a from February, with highest values typically found between May and July with 39 large standard deviations reflecting the high interannual variability in Fig.1. There was no clear seasonal signal in air-sea CO<sub>2</sub> flux, with high variability throughout the 40 41 year. The sediment fluxes had high interannual variability, however, the highest 42 volume flux was typically found in June, while an autumn peak was often found for 43 dry weight and POC in September or October. 44 The monthly MLD-integrated NO<sub>3</sub><sup>-</sup> changes (NO3<sub>MLD</sub>) were positive from 45 February to August (Fig. 3), which means that during these months there was a net 46 decrease in NO<sub>3</sub><sup>-</sup> concentrations in the mixed layer caused by biological drawdown. 47 Conversely, there were negative NO3<sub>MLD</sub> from September to February, meaning that

48 during these months the NO<sub>3</sub><sup>-</sup> concentrations increased due to remineralization and

Helene Frigstad 25.8.15 18.56 Deleted: also

1 entrainment of new nutrients through winter mixing. This corresponds to a MLD-2 integrated seasonal new production (from February to August, see 2.3 for calculation) of  $0.37 \pm 0.14$  mol N m<sup>-2</sup>. Note that all uncertainties given for production estimates in 3 this work are due to interannual variability, as explained in section 2.3. The monthly 4 5 MLD-integrated changes in NCP (NCP<sub>MLD</sub>) showed the same seasonal trend, with a positive NCP<sub>MLD</sub> (i.e. decrease in DIC concentrations) from February to July. In 6 7 addition, there was a much higher NCP<sub>MLD</sub> in March compared to the other 8 spring/summer months. The MLD-integrated seasonal NCP (from February to July) 9 was  $4.57 \pm 0.27$  mol C m<sup>-2</sup>. 10 Tracking of the particles arriving at the sediment trap at 3000 m at the PAP 11 observatory (see Sect. 2.4), revealed that the source locations of particles could vary 12 substantially between years, and also on an annual time scale (Fig. 4). The satellite NPP in these source regions also varied markedly, and the highest NPP of around 210 13 mmol C m<sup>-2</sup> day<sup>-1</sup> was found in 2009. 14 15 There was a high cross-correlation between the seasonal anomalies of NPP in 16 source locations identified by the particle tracking and the volume flux in the sediment trap (+0.62; Fig. 5) at lag = 0 months. The corresponding cross-correlation 17 18 for NPP averaged over a 100 km box around the PAP observatory and volume flux 19 was considerably lower (+0.48; Fig 5), however the difference between the two cross-20 correlations was not significant (Fisher transformation, n = 111, z = 1.58, p = 0.11). 21 The correlation coefficients between either dry weight or POC and the two different 22 NPP estimates were lower, and similarly showed no significant difference between 23 either using the NPP identified by particle tracking or a 100 km box. The highest 24 cross-correlations between NPP and dry weight was found at lag = -1 month (i.e. dry 25 weight lagged NPP by one month), while it was at lag = -3 months for NPP and POC. 26 We also tested the cross-correlations using the NPP at the exact latitude and longitude 27 of the PAP observatory (49°N 16°W) and extending the box to 200 km around the 28 observatory, but there were no significant differences in cross-correlations using these 29 instead of the mean of a 100 km box around the PAP observatory as above (results not 30 shown).

31

#### 32 **3. Discussion**

33

#### 34 4.1 Estimates of surface productivity

The seasonal cycles of carbon (pCO<sub>2</sub>/DIC) and NO<sub>3</sub><sup>-</sup> at the PAP observatory 35 36 are characteristic of highly productive sub-polar regions, where cooling, convection 37 and remineralization cause a winter maximum, while drawdown from biological 38 production during spring and summer causes a minimum during late summer 39 (Takahashi et al. 2002). This is true also for pCO<sub>2</sub>, because the decreasing effect of 40 production is stronger than the opposing effect of warming. The same seasonal effects 41 are seen in the MLD integrated changes in NO3<sup>-</sup> and DIC concentrations used to calculate new production and NCP (i.e. Fig 3). There were positive monthly changes 42 in NO3<sub>MLD</sub> and NCP<sub>MLD</sub> during spring/summer, due to decreasing concentrations 43 44 related to biological production within a shallowing or fairly stable mixed layer. 45 During winter the combined effects of cooling, deep mixing and remineralization 46 caused negative NO3<sub>MLD</sub> and NCP<sub>MLD</sub> (i.e. increasing NO<sub>3</sub><sup>-</sup> and DIC concentrations).

Helene Frigstad 25.8.15 18.19 Deleted: 2500 Helene Frigstad 25.8.15 18.19 Deleted: g

1 The MLD-integrated seasonal NCP (from February to July) was  $4.57 \pm 0.27$ 2 mol C m<sup>-2</sup>, which is comparable to, but lower than, the NCP calculated for the PAP observatory by Körtzinger et al. (2008) for 2004 of  $6.4 \pm 1.1$  mol C m<sup>-2</sup> from March to 3 early August. This is natural given that Körtzinger et al. (2008) integrated over a 4 5 much deeper fixed mixed layer (238 m) from March until mid May, while here we 6 used the mean MLD between consecutive months. As can be seen in Fig. 2, the MLD 7 shallows rapidly from around 250 m in March to 50 m in May, and we believe using the actual depth of the mixed layer gives a more realistic estimate of NCP during this 8 9 period of rapid stratification. From May to August the MLD is slightly shallower than 10 the depth of the sensors at 30m, which could influence our estimates of NCP and new 11 production. This uncertainty is impossible to quantify without measurements from 12 within the mixed layer, although the effect is believed to be minor. It could potentially 13 overestimate biological production due to more readily available nutrients at 30m than within the mixed layer, or lead to an underestimate due to additional drawdown within 14 15 the mixed layer. Consistent with Körtzinger et al. (2008), we find that the strongest 16 NCP occurs before the onset of shallow stratification in summer, with a NCP 17 maximum in March. The MLD-integrated seasonal new production (from February to August) was  $0.37 \pm 0.14$  mol N m<sup>-2</sup>. This is within the range of new production 18 19 estimates, using different approaches, from 0.23 - 1.1 mol N m<sup>-2</sup> from the northeast 20 Atlantic Ocean and Icelandic Sea (Fernández I et al. 2005, Hartman et al. 2010, 21 Jeansson et al. 2014). 22 Converting the new production in terms of nitrogen to carbon units using the 23 Redfield ratio of 6.6 (Redfield 1958), gives a value of 2.5 mol C m<sup>-2</sup>, which is 24 substantially smaller than the NCP calculated from DIC changes. However, there is 25 not necessarily any basis for assuming that new production and NCP should be equal, 26 even in a steady state system (cf. Laws 1991). This would imply that carbon is 27 assimilated and recycled by heterotrophs (as respiration is included in NCP) in the 28 same ratio as nitrogen is assimilated by autotrophs during new production, which 29 need not be the case (i.e. variable stoichiometry, see review by Sterner and Elser 30 (2002)). The fact that the production calculated from carbon is higher than the 31 production calculated from nitrogen, is referred to as carbon overconsumption 32 (Toggweiler 1993), and has been demonstrated repeatedly in the North Atlantic 33 (Sambrotto et al. 1993, Körtzinger et al. 2001, Koeve 2006). It has also been shown 34 specifically for the PAP observatory (Körtzinger et al. 2008, Painter et al. 2010). The 35 seasonal C:N ratio for the PAP observatory from this study (i.e. NCP/new production) 36 would be  $\approx 12$ , which is greatly exceeding the Redfield ratio, although within the 37 range of C:N ratios previously found in this region (Koeve 2006, Körtzinger et al. 38 2008, Painter et al. 2010). 39 Studies have found N<sub>2</sub> fixation to be low or zero at latitudes >40°N (PAP 40 observatory is at 49°N) in the North Atlantic (Moore et al. 2009, Reynolds et al. 41 2007), therefore N<sub>2</sub> fixation is not thought to be an important factor for C 42 overconsumption. Therefore the C overconsumption must be sustained by preferential remineralization of nutrients, either in slowly sinking detritus (Sambrotto 43 et al. 1993, Thomas et al. 1999, Körtzinger et al. 2001) or a build up of C-rich (and N-44 45 poor) DOM in the euphotic zone (Williams 1995, Kähler & Koeve 2001, Falck & 46 Anderson 2005). Additionally transparent exopolymer particles (TEP; Alldredge et al. 47 1993), which have been shown to have high C:N ratios (Engel & Passow 2001) could 48 represent a route for the C overconsumed in the euphotic zone to reach the deep ocean 49 (Koeve 2005). The regenerated nutrients can fuel additional production in the 50 euphotic zone, and consequently estimates of new production based on nitrate might

Helene Frigstad 25.8.15 22.21 Deleted: o Helene Frigstad 25.8.15 22.21 Deleted: o

Helene Frigstad 25.8.15 22.22 Deleted: o

Helene Frigstad 25.8.15 23.04 **Deleted:** o Helene Frigstad 25.8.15 23.04 **Deleted:** o Helene Frigstad 25.8.15 23.05 **Deleted:** o Helene Frigstad 25.8.15 20.22 **Deleted:** I

#### Helene Frigstad 25.8.15 20.22

**Deleted:** n regions where N<sub>2</sub> fixation is not thought to be an important factor, Helene Frigstad 25.8.15 23.05

Deleted: o

Jeleteu. 0

Deleted: a

1 underestimate production rates (Thomas et al. 1999).

2 The degree to which the C overconsumed in the surface waters reaches the 3 deep ocean, and thus is sequestered on long time scales is important, because it 4 represents a potential negative feedback on atmospheric CO2. Studies have shown an 5 increasing C:N of sinking material due to preferential remineralization of nutrients 6 (Schneider et al. 2003, Lee & Cronin 1984), and the potential feedback of a depth 7 dependent C:N ratio may influence atmospheric CO<sub>2</sub> concentrations by about 20 ppm 8 (Schneider et al. 2004). However, the deep ocean remineralization ratio of C:N has 9 been shown to be close to the Redfield ratio (Anderson & Sarmiento 1994), and if the 10 C overconsumption is mainly during summer (Koeve 2004, Jiang et al. 2013) and 11 remineralized above the depth of the winter mixed layer, it could be questioned 12 whether the "extra-Redfield" C is sequestered in the deeper ocean, and can therefore 13 influence the oceanic C-budget on longer time scales (Koeve 2006). However, the 14 deep ocean remineralization rates of Anderson and Sarmiento (1994) did not include 15 the Atlantic Ocean, A study on the remineralization ratios in the North Atlantic Ocean 16 specifically showed higher than Redfield C:nutrient ratios in the remineralized 17 material in the deeper waters, and thereby a higher C drawdown by the biological 18 carbon pump than would be expected from applying Redfield ratios in the formation 19 of organic matter (Thomas 2002). The export flux of POC around the PAP observatory has been quantified in 20

21 several studies using different techniques (see overview in Fig. 4 in Riley et al. (2012)). The average POC flux in the upper 170 m obtained from PELAGRA drifting 22 23 sediment trap deployments for short periods of time (3-5 days) between 2003 and 2005 was 72 mg C m<sup>-2</sup> d<sup>-1</sup> (Lampitt et al. 2008). During a cruise in August 2009 the 24 flux was found to be 84 and 146 mg C m<sup>-2</sup> d<sup>-1</sup> at 50 m, using PELAGRA and a marine 25 snow catcher, respectively (Riley et al. 2012). Using the <sup>234</sup>Th technique the flux of 26 27 POC at 100 m in the vicinity of the PAP observatory was determined to be in the range between 64 and 207 mg C m<sup>-2</sup> d (based on measurements from a single cruise 28 and long-time trap data; Lampitt et al. 2008, Thomalla et al. 2008). The export ratio 29 30 describes the efficiency of nutrient utilization in the euphotic zone, and is often 31 calculated as the POC flux at the base of the euphotic zone or a fixed depth (typically 32 100 m), divided by the NPP (Dugdale & Goering (1967), Using an average of the 33 above values for POC flux out of the surface layer of  $115 \pm 61$  mg C m<sup>-2</sup> d<sup>-1</sup> and the 34 March-July average NPP in the 100 km box around the PAP observatory of 772 mg C 35  $m^{-2} d^{-1}$  gives an export ratio of 0.15. This is identical to the estimate by Lampitt et al. 36 (2008) for the PAP observatory during post-bloom conditions from 2003-2005, and 37 consistent with the estimate by Henson et al. (2011) of between 10 and 30% for 38 temperate and sub-polar waters, respectively. 39

#### 40 4.2 Links between surface production and deep ocean flux of POC

Using particle tracking to identify the source location of material arriving in the
sediment trap at 3000 meters at the PAP observatory showed that the particles could
originate up to 140 km away (in 2007; Fig. 4). There was large variation in the source

- 44 location of particles between years, depending on the prevailing current conditions in
- 45 the given year. There was also large variation within individual years, but the satellite
- 46 NPP generally increased during spring and decreased during autumn along the
- 47 trajectory of the particles reflecting the seasonal cycle. <u>Mesoscale variability will also</u>
- 48 <u>contribute to the variability in source location of particles, both between and within</u>
- 49 years. The highest NPP was found in 2009 (around 210 mmol, C m<sup>-2</sup> day<sup>-1</sup>), which

#### Helene Frigstad 25.8.15 19.26

Deleted: Compilation of existing data sets Helene Frigstad 25.8.15 19.28 Deleted: can

Deleteu. can

Helene Frigstad 25.8.15 19.29 Deleted: parts per million

Deletered: parts per minion

**Deleted:**, because this basin was too complex for the chosen method

Helene Frigstad 25.8.15 22.46 Formatted: Default Paragraph Font, Font:Cambria, Norwegian Bokmal, Check spelling and grammar

#### lelene Frigstad 25.8.15 19.31

Deleted: Comparing our estimates of NCP and new production with satellite-derived estimates of NPP could shed some light on the balance between autotrophy and heterotrophy in this region. Summing the average monthly satellite NPP in a 100 km box around the PAP observatory from the period March to July gives ~10 mol C m<sup>-2</sup>, which is about double the estimated seasonal NCP at the PAP observatory. This would imply approximately equal contributions from NCP and heterotrophic respiration to NPP (as NCP = NPP - heterotrophic respiration). However, it is questionable whether estimates of surface productivity (as given in this study) can be directly compared with satellite NPP. The NCP and new production are estimated based on changes in bulk properties over fairly long timescales, and are integrated over the mixed layer (assuming that the measurements made at ~30 m depth are representative for the entire mixed layer). While the satellite-derived estimates of NPP are often based on shorter timescales and different depth profiles, and therefore the two estimates of surface productivity can involve fundamentally different timescales (cf. Platt et al. 1989). An additional complicating factor is that NCP theoretically includes all the carbon contained in organic material, both the particulate ... [1] Helene Frigstad 25.8.15 19.33

**Deleted:** determined to be 64 and 207 mg C  $m^{-2} d^{-1}$ 

Helene Frigstad 25.8.15 21.56
Deleted: (
Helene Frigstad 25.8.15 23.17
Deleted:
Helene Frigstad 25.8.15 23.17
<b>Deleted:</b> (De La Rocha & Passow 2012, Henson et al. 2012)
Helene Frigstad 25.8.15 18.20
Deleted: >
Helene Frigstad 25.8.15 18.21
Deleted: 000
Helene Frigstad 25.8.15 18.21
Deleted: a

1 corresponds to very high fluxes in both volume flux and dry weight in the sediment 2 trap at the PAP observatory (Fig. 1). Interestingly, there was not a strong bloom at the 3 PAP observatory according to the in situ Chl a observations at 30 m depth (sensor 4 data available from May to late July; Fig. 1), while the satellite NPP showed high 5 correlations with the volume flux and dry weight in the sediment trap. The correlations were highest between the NPP in source locations as identified by particle 6 7 tracking, compared to the mean NPP in a fixed 100 km box around the PAP 8 observatory. However, the differences in correlation coefficients were not statistically 9 different, and more observations (n = 111 in present analysis) would be needed to 10 determine if using a particle tracking approach when examining the origin of particles in sediment traps indeed gives higher correlations. 11 12 The transfer efficiency is a useful metric to describe the long-term removal of 13 carbon (> 100 years) from the atmosphere (cf. De La Rocha & Passow 2012). Using 14 the same average POC flux of the surface layer (0-170 m) as in the calculation of the 15 export flux above and the average flux between March and July of POC at 3000 m 16 from the sediment trap at the PAP observatory (5.1 mg C  $m^{-2} d^{-1}$ ), the transfer 17 efficiency was calculated to be 4%. This corresponds well with the transfer efficiency 18 between 5 and 10% found for the 50°N region by Henson et al. (2012b), indicative of 19 regions where the available nitrate is not fully consumed and with strong seasonal 20 mixing. The fairly high export ratio (15%) and low transfer efficiency (4%) fits the 21 description of the general trends in high latitude ecosystems in the above-cited study 22 well. This dichotomy in efficiencies implies that although a large proportion of the 23 primary production is exported below the euphotic zone, this material is relatively labile and is efficiently remineralized in the mesopelagic zone, so that only a very 24 25 small fraction of the exported organic matter reaches the deep ocean and is stored on 26 long time scales. 27 The PAP observatory currently sits near the boundary between the sub-polar and sub-tropical gyres of the North Atlantic. Seasonably variable areas, like the sub-28 29 polar region, are thought to export a higher fraction of labile material than subtropical regions (Lutz et al. 2007). As climate change is predicted to result in the 30 oligotrophic gyres expanding over the next century (Sarmiento et al. 2004), the PAP 31 32 observatory will likely transition into more sub-tropical conditions. This could result 33 in more refractory material being exported at the PAP observatory, potentially 34 reversing the pattern we report here with a high export ratio and low transfer 35 efficiency. In addition, a more strongly positive NAO index is predicted due to 36 climate change (Gillett 2003), which is expected to increase diatom abundance at the 37 PAP observatory, and result in reduced organic carbon flux to the deep ocean (Henson 38 et al., 2012a). Although the precise response of the biological carbon pump to climate 39 change is as yet unclear, transition-zone regions between gyres (such as the PAP 40 observatory), could be among the systems that are most strongly affected by climate 41 change (Henson et al. 2013), 42

#### 43 5. Conclusions

- 44 The PAP observatory is characterized by strong interannual variability in
- 45 hydrography, biogeochemistry, and especially sediment fluxes. The seasonal cycles of
- 46 carbon and nitrogen show a winter maximum and summer minimum, characteristic of
- 47 highly productive sub-polar regions. The MLD-integrated seasonal NCP (from
- 48 February to July) was  $4.57 \pm 0.27$  mol C m<sup>-2</sup>, which is consistent, with but slightly
- 49 lower than the estimate by Körtzinger et al. (2008) for 2004. The MLD-integrated

#### Helene Frigstad 25.8.15 20.25

**Deleted:** is used to describe the efficiency of the biological carbon pump, and

Helene Frigstad 25.8.15 20.28 Deleted: is Helene Frigstad 25.8.15 20.28 Deleted:

#### Helene Frigstad 25.8.15 20.28

**Deleted:** at or close to the boundary between the sub-polar and sub-tropical gyres of the North Atlantic, and will likely transition into more sub-tropical conditions as the gyres expand over the next century (<u>Sarmiento et al.</u> 2004). It has been shown that seasonably variable areas, like the sub-polar region, export a higher fraction of labile material than subtropical regions (Lutz et al. 2007), which could imply that the current export regime of the PAP observatory is likely to change as a result of climate change.

Helene Frigstad 25.8.15 23.05 Deleted: o

- 1 seasonal new production (from February to August) was  $0.37 \pm 0.14$  mol N m<sup>-2</sup>,
- 2 which gives a Redfield ratio (NCP/new production) of 12, corroborating other reports
- 3 of carbon overconsumption for the North Atlantic (Sambrotto et al. 1993, Körtzinger

4 <u>et al. 2001, Koeve 2006</u> and the PAP observatory specifically (Körtzinger et al. 2008, 5 Painter et al. 2010).

6 The export ratio was 15%, while the transfer efficiency was 4%, which is

- 7 typical of high latitude ecosystems where, although a large proportion of the primary
- 8 production is exported out of the euphotic zone, this material is relatively labile and
- 9 therefore remineralized before it reaches the deep ocean. It is hypothesized that the 10 export regime at the PAP observatory could change with climate change, as the region
- will probably transition into more sub-tropical conditions over the next century
- 12 (Sarmiento et al. 2004, Lutz et al. 2007).
- 13 Using particle tracking to identify the source regions of material reaching the
- 14 sediment trap at the PAP observatory, revealed higher correlations between NPP in
- 15 the identified source regions and export flux than other methods. However, more
- 16 observations are needed to establish if a particle-tracking approach indeed gives
- 17 added value in sediment trap analyses.

#### 18 Acknowledgements

- 19
- 20 We would like to acknowledge the various ship crew, engineers and
- 21 scientists involved in preparation, deployment and recovery of the PAP sustained
- 22 observatory moorings, especially Jon Campbell and Mark Hartman. We wish to thank
- 23 Maureen Pagnani, Athanos Gkritzalis-Papadopoulos, Zong-Pei Jiang and Andres
- 24 Cianca for compilation, quality control and calibration of PAP data. Mooring data and
- 25 support for this research was provided by the European research projects ANIMATE
- 26 (Atlantic Network of Interdisciplinary Moorings and Time- Series for Europe),

27 MERSEA (Marine Environment and Security for the European Sea), EUR- OCEANS

- 28 (European Network of Excellence for Ocean Ecosystems Analysis) and EuroSITES
- 29 grant agreement EU 202955. The work was also supported through the Natural
- 30 Environment Research Council (NERC), UK, project Oceans 2025 and National
- 31 Capability. H. F. was supported by EU FP7, through projects MEECE (212085),
- 32 EURO-BASIN (264933) and GreenSeas (265294).
- 33
- 34

Helene Frigstad 25.8.15 22.50 Moved (insertion) [1] Unknown Field Code Changed Helene Frigstad 25.8.15 23.05 Deleted: o Helene Frigstad 25.8.15 22.50 Moved up [1]: (Sambrotto et al. 1993, Kortzinger et al. 2001, Koeve 2006) Helene Frigstad 25.8.15 22.51 Deleted: ;

1	References	
2		
3	Alldredge AL, Passow U, Logan BE (1993) The Abundance and Significance of a	
4	Class of Large, Transparent Organic Particles in the Ocean. Deep-Sea Res Pt I	
5	40:1131-1140	
6	Anderson LA, Sarmiento JL (1994) Redfield Ratios of Remineralization Determined	
7	by Nutrient Data-Analysis. Global Biogeochem Cy 8:65-80	
8	Behrenfeld MJ, Falkowski PG (1997) Photosynthetic rates derived from satellite-	
9	based chlorophyll concentration. Limnol Oceanogr 42:1-20	
10	Berelson WM (2001) The flux of partuculate organic carbon into the ocean interior. A	
11	comparison of four U.S. JGOFS regional studies. Oceanography 14:59-67	
12	De La Rocha CL, Passow U (eds) (2012) The biological pump, Vol 6. Elsevier,	
13	Uxford $D(1, D(1, 0, 0))$ The first first state of the last first first state of the last first state of the last state	
14	Dickson AG, Kiley JP (1978). The effect of analytical error on the evaluation of the	
15	<u>Components of the aquatic carbon-dioxide system. Mar Chem 6: //-85</u>	
10	Dickson AG (1990) Standard Potential of the Reaction - Agci(S)+ $1/2n$ -	
1/	2(G)=Ag(S)+Hcl(Aq) and the Standard Acidity Constant of the Ion Hso4- in	
10	Synthetic Sea-water from 2/3.15-K to 318.15-K. J Chem Thermodyn 22.115-	
20	Dickson AG Sabine CL. Chrsitian IR (2007) Guide to best practices for ocean CO2	
21	measurements. PICES Special Publication. Book 3	
22	Dugdale RC, Goering JJ (1967) Uptake of New and Regenerated Forms of Nitrogen	
23	in Primary Productivity. Limnol Oceanogr 12:196-&	
24	Engel A, Passow U (2001) Carbon and nitrogen content of transparent exopolymer	
25	particles (TEP) in relation to their Alcian Blue adsorption. Mar Ecol-Prog Ser	
26	219:1-10	
27	Eppley RW, Peterson BJ (1979) Particulate Organic-Matter Flux and Planktonic New	
28	Production in the Deep Ocean. Nature 282:677-680	
29	Falck E, Anderson LG (2005) The dynamics of the carbon cycle in the surface water	
30	of the Norwegian Sea. Mar Chem 94:43-53	
31	Falkowski PG, Barber RT, Smetacek V (1998) Biogeochemical controls and	
32	feedbacks on ocean primary production. Science 281:200-206	
33	Fernández I C, Raimbault P, Garcia N, Rimmelin P, Caniaux G (2005) An estimation	
34	of annual new production and carbon fluxes in the northeast Atlantic Ocean	
35	during 2001. Journal of Geophysical Research: Oceans 110:C07S13	
36	Fisher RA (1915) Frequency distribution of the values of the correlation coefficient in	
37	samples of an indefinitely large population. Biometrika 10:507-521	
38	Francois R, Honjo S, Krishfield R, Manganini S (2002) Factors controlling the flux of	
39	organic carbon to the bathypelagic zone of the ocean. Global Biogeochem Cy	
40	16	
41	Gaillard F, Autret E, Thierry V, Galaup P, Coatanoan C, Loubrieu T (2009) Quality	
42	Control of Large Argo Datasets. J Atmos Ocean Technol 26:337-351	
43	GLUBALVIEW-CU2 (2012) Cooperative Atmospheric Data Integration Project –	
44	Carbon Dioxide. NOAA ESRL, Boulder, Colorado [Available at	
45	<u>nttp://www.esri.noaa.gov/gma/ccgg/globalview/%5D</u> .	
40	Druber in (2008) The marine mulogen cycle. Overview and challenges. In: Capone	
4/ 10	Environment Rock 2 Elsevier Ameterdem	
40 40	Environment, DOUK 2. Elsevier, Amsterdam Georder T. Gran HH (1027) Investigation of the production of phytoplanktan in the	
47 50	Oslo Fiord Papp DV Cons Int Explor Mar 42:1 49	
50	Usio I Joid. Rapp I V Cons in Exploi Mici 42.1-40	

1	Hartman SE, Lampitt RS, Larkin KE, Pagnani M, Campbell J, Gkritzalis T, Jiang ZP,
2	Pebody CA, Ruhl HA, Gooday AJ, Bett BJ, Billett DSM, Provost P,
3	McLachlan R, Turton JD, Lankester S (2012) The Porcupine Abyssal Plain
4	fixed-point sustained observatory (PAP-SO): variations and trends from the
5	Northeast Atlantic fixed-point time-series. Ices J Mar Sci 69:776-783
6	Hartman SE, Larkin KE, Lampitt RS, Lankhorst M, Hydes DJ (2010) Seasonal and
7	inter-annual biogeochemical variations in the Porcupine Abyssal Plain 2003-
8	2005 associated with winter mixing and surface circulation. Deep-Sea Res Pt
9	li 57:1303-1312
10	Henson SA, Dunne JP, Sarmiento JL (2009) Decadal variability in North Atlantic
11	phytoplankton blooms. J Geophys Res-Oceans 114
12	Henson SA, Sanders R, Madsen E, Morris PJ, Le Moigne F, Quartly GD (2011) A
13	reduced estimate of the strength of the ocean's biological carbon pump.
14	Geophys Res Lett 38
15	Henson S, Lampitt R, Johns D (2012a) Variability in phytoplankton community
16	structure in response to the North Atlantic Oscillation and implications for
17	organic carbon flux, Limnol Oceanogr 57:1591-1601
18	Henson SA, Sanders R, Madsen E (2012b) Global patterns in efficiency of particulate
19	organic carbon export and transfer to the deep ocean. Global Biogeochem Cy
20	26
21	Henson S, Cole H, Beaulieu C, Yool, A (2013) The impact of global warming on
22	seasonality of ocean primary production, Biogeosciences 10: 4357-4369
23	Holte J, Talley L (2009) A New Algorithm for Finding Mixed Layer Depths with
24	Applications to Argo Data and Subantarctic Mode Water Formation. J Atmos
25	Ocean Tech 26:1920-1939
26	Jeansson E, Bellerby RGJ, Skjelvan I, Frigstad H, Ólafsdóttir SR, Ólafsson J (2014)
27	Fluxes of carbon and nutrients to the Iceland Sea surface layer and inferred
28	primary productivity and stoichiometry Biogeosciences In review
29	Jiang Z-P, Hydes DJ, Tyrrell T, Hartman SE, Hartman MC, Dumousseaud C, Padin
30	XA, Skjelvan I, González-Pola C (2013) Key controls on the seasonal and
31	interannual variations of the carbonate system and air-sea CO2 flux in the
32	Northeast Atlantic (Bay of Biscay). Journal of Geophysical Research: Oceans
33	118:785-800
34	Kähler P, Koeve W (2001) Marine dissolved organic matter: can its C : N ratio
35	explain carbon overconsumption? Deep-Sea Res Pt I 48:49-62
36	Koeve W (2004) Spring bloom carbon to nitrogen ratio of net community production
37	in the temperate N. Atlantic. Deep-Sea Res Pt I 51:1579-1600
38	Koeve W (2005) Magnitude of excess carbon sequestration into the deep ocean and
39	the possible role of TEP. Mar Ecol-Prog Ser 291:53-64
40	Koeve W (2006) C : N stoichiometry of the biological pump in the North Atlantic:
41	Constraints from climatological data. Global Biogeochem Cy 20
42	Körtzinger A, Koeve W, Kähler P, Mintrop L (2001) C : N ratios in the mixed layer
43	during the productive season in the northeast Atlantic Ocean. Deep-Sea Res Pt
44	I 48:661-688
45	Körtzinger A, Send U, Lampitt RS, Hartman S, Wallace DWR, Karstensen J,
46	Villagarcia MG, Llinas O, DeGrandpre MD (2008) The seasonal pCO(2) cycle
47	at 49 degrees N/16.5 degrees W in the northeastern Atlantic Ocean and what it
48	tells us about biological productivity. J Geophys Res-Oceans 113
49	Lampitt RS, Achterberg EP, Anderson TR, Hughes JA, Iglesias-Rodriguez MD,
50	Kelly-Gerreyn BA, Lucas M, Popova EE, Sanders R, Shepherd JG, Smythe-

Helene Frigstad 25.8.15 23.08 Deleted: a

1	Wright D, Yool A (2008a) Ocean fertilization: a potential means of
2	geoengineering? Phil. Trans. R. Soc. A 366: 3919-3945
3	Lampitt RS, Boorman B, Brown L, Lucas M, Salter I, Sanders R, Saw K, Seeyave S,
4	Thomalla SJ, Turnewitsch R (2008b) Particle export from the euphotic zone:
5	Estimates using a novel drifting sediment trap, Th-234 and new production.
6	Deep-Sea Res Pt I 55:1484-1502
7	Lampitt RS, Salter I, de Cuevas BA, Hartman S, Larkin KE, Pebody CA (2010)
8	Long-term variability of downward particle flux in the deep northeast Atlantic:
9	Causes and trends. Deep-Sea Res Pt Ii 57:1346-1361
10	Lavigne H, Gattuso J-P (2011) Seacarb: seawater carbonate chemistry with R. R
11	package version 2.4. <u>http://CRAN.R-project.org/package=seacarb</u> .
12	Laws EA (1991) Photosynthetic Quotients, New Production and Net Community
13	Production in the Open Ocean. Deep-Sea Res 38:143-167
14	Laws EA, Falkowski PG, Smith WO, Ducklow H, McCarthy JJ (2000) Temperature
15	effects on export production in the open ocean. Global Biogeochem Cy
16	14:1231-1246
17	Lee C, Cronin C (1984) Particulate amino acids in the sea: effects of primary
18	productivity and biological decomposition. J. Mar. Res 42:1075–1097.
19	Lee K (2001) Global net community production estimated from the annual cycle of
20	surface water total dissolved inorganic carbon. Limnol Oceanogr 46:1287-
21	1297
22	Lee K, Tong LT, Millero FJ, Sabine CL, Dickson AG, Goyet C, Park G-H,
23	Wanninkhof R, Feely RA, Key RM (2006) Global relationships of total
24	alkalinity with salinity and temperature in surface waters of the world's
25	oceans. Geophys Res Lett 33
26	Long MC, Dunbar RB, Tortell PD, Smith WO, Mucciarone DA, DiTullio GR (2011)
27	Vertical structure, seasonal drawdown, and net community production in the
28	Ross Sea, Antarctica. J Geophys Res-Oceans 116
29	Lueker TJ, Dickson AG, Keeling CD (2000) Ocean pCO(2) calculated from dissolved
30	inorganic carbon, alkalinity, and equations for K-1 and K-2: validation based
31	on laboratory measurements of CO2 in gas and seawater at equilibrium. Mar
32	Chem 70:105-119
33	Lutz MJ, Caldeira K, Dunbar RB, Behrenfeld MJ (2007) Seasonal rhythms of net
34	primary production and particulate organic carbon flux to depth describe the
35	efficiency of biological pump in the global ocean. Journal of Geophysical
36	Research: Oceans 112:C10011
37	Madec G (2008) NEMO ocean engine. Note du Pole de modélisation, Institut Pierre-
38	Simon Laplace (IPSL), France, No 27 ISSN No 1288-1619
39	Martin JH, Knauer GA, Karl DM, Broenkow WW (1987) Vertex - Carbon Cycling in
40	the Northeast Pacific. Deep-Sea Res 34:267-285
41	Millero FJ (1995) The thermodynamics of the carbonic acid
42	system in the oceans. Geochimica et Cosmochimica Acta 59:661-667
43	Moore C, Mills M, Matthew M, Achterberg EP, Geider RJ, LaRoche J, Lucas MI,
44	McDonagh EL, Pan X, Poulton AJ, Rijkenberg, MJA, Suggett, DJ, Ussher SJ,
45	Woodward E, Malcolm S (2009) Large-scale distribution of Atlantic nitrogen
46	tixation controlled by iron availability.Nature Geoscience 2:867-871.
47	Nevison CD, Keeling RF, Kahru M, Manizza M, Mitchell BG, Cassar N (2012)
48	Estimating net community production in the Southern Ocean based on
49	atmospheric potential oxygen and satellite ocean color data. Global
50	Biogeochem Cy 26

- 1 Nightingale PD, Malin G, Law CS, Watson AJ, Liss PS, Liddicoat MI, Boutin J, 2 Upstill-Goddard RC (2000) In situ evaluation of air-sea gas exchange 3 parameterizations using novel conservative and volatile tracers. Global 4 Biogeochem Cy 14:373-387 5 Painter SC, Lucas MI, Stinchcombe MC, Bibby TS, Poulton AJ (2010) Summertime trends in pelagic biogeochemistry at the Porcupine Abyssal Plain study site in 6 7 the northeast Atlantic. Deep-Sea Res Pt Ii 57:1313-1323 8 Passow U, Carlson CA (2012) The biological pump in a high CO2 world. Mar Ecol 9 Prog Ser 470:249-271 10 Perez FF, Fraga F (1987) Association Constant of Fluoride and Hydrogen-Ions in 11 Seawater. Mar Chem 21:161-168 12 Platt T, Harrison WG, Lewis MR, Li WKW, Sathyendranath S, Smith RE, Vezina AF 13 (1989) Biological Production of the Oceans - the Case for a Consensus. Mar 14 Ecol Prog Ser 52:77-88 15 Quay PD, Peacock C, Bjorkman K, Karl DM (2010) Measuring primary production rates in the ocean: Enigmatic results between incubation and non-incubation 16 17 methods at Station ALOHA. Global Biogeochem Cy 24 18 Redfield AC (1958) The Biological Control of Chemical Factors in the Environment. 19 Am Sci 46:205-221 20 Revelle W (2012) Psych: Procedures for Personality and Psychological Research 21 Northwestern University, Evanston, http://personality-22 project.org/r/psych.manual.pdf, 1.2.1. 23 Reynolds, SE, Mather RL, Wolff GA, Williams RG, Landolfi A, Sanders R, Woodward EMS (2007) How widespread and important is N2 fixation in the 24 North Atlantic Ocean?, Global Biogeochem Cy 21 Riley JS, Sanders R, Marsay C, Le Moigne FAC, Achterberg EP, Poulton AJ (2012) 25 26 27 The relative contribution of fast and slow sinking particles to ocean carbon 28 export. Global Biogeochem Cy 26 29 Sabine CL, Feely RA, Gruber N, Key RM, Lee K, Bullister JL, Wanninkhof R, Wong 30 CS, Wallace DWR, Tilbrook B, Millero FJ, Peng TH, Kozyr A, Ono T, Rios AF (2004) The oceanic sink for anthropogenic CO2. Science 305:367-371 31 32 Sambrotto RN, Savidge G, Robinson C, Boyd P, Takahashi T, Karl DM, Langdon C, 33 Chipman D, Marra J, Codispoti L (1993) Elevated Consumption of Carbon 34 Relative to Nitrogen in the Surface Ocean. Nature 363:248-250 35 Sarmiento JL, Gruber N (2006) Ocean Biogeochemical Dyamics. Princeton 36 University Press, Princeton, New Jersey, USA 37 Sarmiento JL, Slater R, Barber R, Bopp L, Doney SC, Hirst AC, Kleypas J, Matear R, 38 Mikolajewicz U, Monfray P, Soldatov V, Spall SA, Stouffer R (2004) 39 Response of ocean ecosystems to climate warming. Global Biogeochem Cy 40 18:GB3003 41 Schneider B, Engel A, Schlitzer R (2004) Effects of depth- and CO2-dependent C : N 42 ratios of particulate organic matter (POM) on the marine carbon cycle. Global 43 **Biogeochem Cy 18** Schneider B, Schlitzer R, Fischer G, Nothig EM (2003) Depth-dependent elemental 44 45 compositions of particulate organic matter (POM) in the ocean. Global 46 Biogeochem Cy 17 47 Steinberg DK, Cope JS, Wilson SE, Kobari T (2008) A comparison of mesopelagic 48 mesozooplankton community structure in the subtropical and subarctic North 49 Pacific Ocean. Deep-Sea Res Pt Ii 55:1615-1635
  - 16

1	Sterner RW, Elser JJ (2002) Ecological Stoichiometry: the biology of elements from
2	Takahashi T. Sutharland SC. Swaanay C. Daisson A. Matzl N. Tilbraak P. Datas N.
3 4	Wanninkhof R Eeely RA Sabine C Olafsson I Nojiri V (2002) Global sea
т 5	air $\Omega(2)$ flux based on climatological surface ocean $p(\Omega(2))$ and seasonal
6	hiological and temperature effects. Deen-Sea Res Pt Ii 49:1601-1622
7	Team RDC (2012) R: A language and environment for statistical computing R
8	Foundation for Statistical Computing Vienna Austria
9	Thomalla SJ. Poulton AJ. Sanders R. Turnewitsch R. Holligan PM. Lucas MI (2008)
10	Variable export fluxes and efficiencies for calcite, opal, and organic carbon in
11	the Atlantic Ocean: A ballast effect in action? Global Biogeochem Cy 22
12	Thomas H (2002) Remineralization ratios of carbon, nutrients, and oxygen in the
13	North Atlantic Ocean: A field databased assessment. Global Biogeochem Cy
14	16
15	Thomas H, Ittekkot V, Osterroht C, Schneider B (1999) Preferential recycling of
16	nutrients - the ocean's way to increase new production and to pass nutrient
17	limitation? Limnol Oceanogr 44:1999-2004
18	Toggweiler JR (1993) Oceanography - Carbon Overconsumption. Nature 363:210-
19	211
20	Villa-Alfageme M, De Soto Borrero F, Le Moigne F, Giering SL, Sanders R (2014)
21	Observations and modeling of slow sinking particles in the Twilight Zone.
22	Global Biogeochem Cy 28:1327-1342
23	Wanninkhof R (1992) Relationship between Wind-Speed and Gas-Exchange over the
24	Ocean. J Geophys Res-Oceans 97:7373-7382
25	Wanninkhof R, D Bakker, N Bates, A Olsen, T Steinhoff, A Sutton (2013)
26	Incorporation of Alternative Sensors in the SOCAT Database and Adjustments
27	to Dataset Quality Control Flags. Carbon Dioxide Information Analysis
28	Center, Oak Ridge National Laboratory, US Department of Energy, Oak
29	Ridge, Tennessee.
30	Weiss RF (1974) Carbon dioxide in water and seawater: The solubility of a non-ideal
31	gas. Mar Chem 2:203-215
32	Williams PJL (1995) Evidence for the Seasonal Accumulation of Carbon-Rich
33	Dissolved Organic Material, Its Scale in Comparison with Changes in
34	Particulate Material and the Consequential Effect on Net C/N Assimilation
35	Ratios. Mar Chem 51:17-29
36	

### 1 Tables

4	
3 Table 1. Overview	of ancillary data
Parameter	Source
ARGO Temperature and salinity	Temperature and salinity fields: 1° by 1°, monthly temporal resolution. Depth in profiles had resolution of 5 metres between 10 and 100m and 10m between 100
Atmospheric CO <sub>2</sub>	and 800m. Available through Coriolis project; http://www.coriolis.eu.org/ Mace Head land station in Ireland (53.33N 9.90W) from the Cooperative Atmospheric Data Integration Projects (GLOBALVIEW-CO2 2012);
Sea Level Pressure (SLP) Wind speed	http://www.esrl.noaa.gov/gmd/ccgg/globalview/co2/co2_download.html FNMOC 1° by 1°, 6-hourly; http://las.pfeg.noaa.gov/las6_5/servlets/constrain?var=6 FNMOC 1° by 1°, 6-hourly at 10 m height; http://las.pfeg.noaa.gov/las6_5/servlets/constrain?var=82
Net primary production (NPP)	Vertically Generalised Production Model; http://www.science.oregonstate.edu/ocean.productivity/index.php

#### 1 Figure captions 2

13

8
9 Figure 2. Monthly climatology (black dots) with ± 1 SD (vertical bars) of available
data from PAP surface mooring (30 m) and sediment trap. Temperature and salinity
calculated from the Argo float data (grey dots in first two panels) is also shown, along
with derived mixed layer depth estimates.

14 Figure 3. Monthly changes in MLD integrated  $NO_3^{-}$  (top) and NCP (bottom) with  $\pm 1$ 15 SD (vertical bars). The shaded grey area indicates the months over which the seasonal

- 16 new production  $(0.37 \pm 0.14 \text{ mol N m}^2)$  and NCP  $(4.57 \pm 0.27 \text{ mol C m}^2)$  were
- 17 calculated. For NCP<sub>MLD</sub> no SD could be calculated in March and April, and SD is also
- $\frac{\text{Lacking for December for both NCP_{MLD} and NO3_{MLD} (see Sect. 2.3 for calculations)}{\text{Lacking for December for both NCP_{MLD} and NO3_{MLD} (see Sect. 2.3 for calculations)}}$

18 1 lacking for December for both INCP<sub>MLD</sub> and NOS<sub>MLD</sub> (see Sect. 2.3 for carculations), 19

# Figure 4. Satellite NPP (mmol C m<sup>-2</sup> d<sup>-1</sup>) in the source regions for the sediment trap as identified by particle tracking. The star shows the position of the PAP mooring and unfilled circles indicate that no NPP estimate was available from satellite data (most often during January and February). The start and end dates of the data for each year

often during January and February). The start and end dates of the data for each year
 are indicated in the panels.

Figure 5. Cross-correlations between <u>sediment trap data (top: volume flux, middle:</u>
dry weight, bottom: POC) and NPP in the source regions defined by particle tracking

dry weight, bottom: POC) and NPP in the source regions defined by particle tracking
 (left) or in a 100 km box around the PAP observatory (right), The dashed lines show

29 the 95% confidence intervals. The unit of the lags is months.

#### Helene Frigstad 25.8.15 18.01

**Deleted:** Monthly changes in MLD integrated NO<sub>3</sub> (top) and NCP (bottom). The shaded grey area indicates the months over which the seasonal new production ( $NO3_{MLD}$ :  $0.37 \pm 0.14$  mol N m<sup>-2</sup>) and seasonal NCP ( $NCP_{MLD}$ :  $4.57 \pm 0.27$  mol C m<sup>-2</sup>) were calculated. See Sect. 2.3 for calculations. Helene Frigstad 25.8.15 18.18

Deleted: g

#### Helene Frigstad 25.8.15 20.32

**Deleted:** and sediment fluxes