

This discussion paper is/has been under review for the journal Biogeosciences (BG).
Please refer to the corresponding final paper in BG if available.

Nutrient and mercury deposition and storage in an alpine snowpack of the Sierra Nevada, USA

C. Pearson^{1,2,3}, R. Schumer^{1,3}, B. D. Trustman^{2,3}, K. Rittger⁴, D. W. Johnson^{3,5},
and D. Obrist^{2,3}

¹Desert Research Institute, Division of Hydrologic Sciences, Reno, NV, USA

²Desert Research Institute, Division of Atmospheric Sciences, Reno, NV, USA

³University of Nevada-Reno, Graduate Program of Hydrologic Sciences, Reno, NV, USA

⁴National Snow and Ice Data Center, Boulder, CO, USA

⁵University of Nevada-Reno, Department of Natural Resources and Environmental Science,
Reno, NV, USA

Received: 29 September 2014 – Accepted: 5 December 2014 – Published: 12 January 2015

Correspondence to: C. Pearson (chrispearson86@gmail.com)

Published by Copernicus Publications on behalf of the European Geosciences Union.

BGD

12, 593–636, 2015

Nutrient and mercury
deposition and
storage in an alpine
snowpack

C. Pearson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Bi-weekly snowpack core samples were collected at seven sites along two elevation gradients in the Tahoe Basin during two consecutive snow years to evaluate total wintertime snowpack accumulation of nutrients and pollutants in a high elevation watershed of the Sierra Nevada. Additional sampling of wet deposition and detailed snow pit profiles was conducted the following year to compare wet deposition to snowpack storage and assess the vertical dynamics of snowpack chemicals. Results show that on average organic N comprised 48 % of all snowpack N, while nitrate (NO_3^- -N) and TAN (total ammonia nitrogen) made up 25 and 27 %, respectively. Snowpack NO_3^- -N concentrations were relatively uniform across sampling sites over the sampling seasons and showed little difference between seasonal wet deposition and integrated snow pit concentrations in agreement with previous studies that identify wet deposition as the dominant source of wintertime NO_3^- -N deposition. However, vertical snow pit profiles showed highly variable concentrations of NO_3^- -N within the snowpack indicative of additional deposition and in snowpack dynamics. Unlike NO_3^- -N, snowpack TAN doubled towards the end of winter and in addition to wet deposition, had a strong dry deposition component. Organic N concentrations in snowpack were highly variable (from 35 to 70 %) and showed no clear temporal or spatial dependence throughout the season. Integrated snowpack organic N concentrations were up to 2.5 times higher than seasonal wet deposition, likely due to microbial immobilization of inorganic N as evident by coinciding increases of organic N and decreases of inorganic N, in deeper, aged snowpack. Spatial and temporal deposition patterns of snowpack P were consistent with particulate-bound dry deposition inputs and strong impacts from in-basin sources causing up to 6 times enrichment at urban locations compared to remote sites. Snowpack Hg showed little temporal variability and was dominated by particulate-bound forms (78 % on average). Dissolved Hg concentrations were consistently lower in snowpack than in wet deposition which we attribute to photochemical-driven gaseous re-emission. In agreement with this pattern is a significant positive relationship between

BGD

12, 593–636, 2015

Nutrient and mercury deposition and storage in an alpine snowpack

C. Pearson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



snowpack Hg and elevation, attributed to a combination of increased snow accumulation at higher elevations causing limited light penetration and lower photochemical re-emission losses in deeper, higher elevation snowpack. Finally, estimates of basin-wide loading based on spatially extrapolated concentrations and a satellite-based snow water equivalent reconstruction model identify snowpack chemical loading from atmospheric deposition as a substantial source of nutrients and pollutants to the Lake Tahoe basin, accounting for 113 t of N, 9.3 t of P, and 1.2 kg of Hg each year.

1 Introduction

Atmospheric deposition accounts for significant nutrient and pollutant input to high elevation watersheds such as the Sierra Nevada (Dolislager, 2006; Fain et al., 2011; McDaniel, 2013; Sickman et al., 2003; TERC, 2011; Vicars and Sickman, 2011; Williams and Melack, 1991a, b). Sierra Nevada snowpack supplies the majority of water to downstream communities as well as to some of the nation's largest agricultural areas. Quantifying atmospheric deposition in alpine watersheds is challenging because of large spatial variability in deposition rates caused by complex terrain, precipitation gradients, and varied origins of atmospheric constituents (i.e. local vs. regional and global, natural vs. anthropogenic; Jassby et al., 1994; Rohrbough et al., 2003). Single-site measurements, therefore, do not allow for accurate extrapolation of nutrient or pollutant deposition in alpine regions and broader temporal and spatial data is needed to assess the mass and dynamics of atmospheric inputs.

In this study, we used multiple and repeated sampling of full depth snowpack cores (integrated snowpack sampling) across the Lake Tahoe basin to quantify atmospheric deposition loads and patterns from the first snowfall until the end of melting. Snowpack acts as an integrating reservoir for water, solutes, and particulates that deposit throughout winter and spring (Turk et al., 2001). Wet deposition, in the form of snowfall and rain, directly accumulates in developing snowpack throughout the snow season (Kuhn, 2001). Additionally, during storm-free periods, snowpack also collects dry de-

BGD

12, 593–636, 2015

Nutrient and mercury deposition and storage in an alpine snowpack

C. Pearson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



across mountain ranges (Fain et al., 2011; NADP, 2012). Assessing spatial deposition patterns using snowpack sampling at multiple locations across a watershed should allow for better characterization of basin-wide deposition patterns as well as assessment of impacts of nearby urban areas vs. regional and global sources of atmospheric deposition (Brown et al., 2011; Kuhn, 2001; Morales-Baquero et al., 2006; Vicars and Sickman, 2011).

The main goal of this study was to quantify N, P, and Hg concentrations and loads in Sierra Nevada snowpack in order to characterize the magnitude, origin, and fate of atmospheric deposition of nutrients and pollutants that accumulate throughout the winter and spring in this mountain range. We quantified chemical loading at seven sites in the Lake Tahoe basin, along two elevation transects, throughout the duration of two full snow seasons. Sampling included bi-weekly snowpack cores (full profile; integrated snowpack samples) representing an integrated load of constituents in the developing snowpack collected throughout the 2011–2012 and 2012–2013 snow years. In addition, volume-weighted wet deposition measured at two sites in 2013–2014 was compared to snowpack accumulation and detailed vertical snow pit profiles in that year to compare snowpack accumulation to wet deposition and to further study in-snowpack chemical dynamics. Finally, basin-wide loading estimates (mass area⁻¹) were calculated by spatially extrapolating nutrient and pollutant measurements across the basin combined with a satellite-based snowpack reconstruction model.

2 Materials and methods

2.1 Study site

The Lake Tahoe watershed lies in the northern portion of the Sierra Nevada range along the border of Nevada and California. Renowned for its intense blue color and water clarity, this lake has become a national landmark and tourism hotspot. Lake clarity measurements have decreased, however, from approximately 30.5 to 21.3 m since the

BGD

12, 593–636, 2015

Nutrient and mercury deposition and storage in an alpine snowpack

C. Pearson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



1960s due to eutrophication from increased input of N and P, as well as additional input of light scattering particulates (TERC, 2011). Directly west and upwind of the basin lies the central valley of California and cities of Sacramento and San Francisco, California, which are suspected of contributing significant amounts of nutrients and pollutants to the basin through agricultural and industrial emissions.

Including all drainages, the Lake Tahoe watershed has an area of 1310 km² (Fig. 1). The lake is 19 km wide and 35 km long with a total surface area of 495 km². The lake lies at 1897 m a.s.l. and is on average 300 m deep. Surrounding the lake on all sides are mountains up to elevations of 3068 m. At the lake's surface, summer temperatures on average reach 27 °C and wintertime lows reach -9 °C. Precipitation patterns in the watershed are highly dependent on elevation with an average annual precipitation of 0.76 m at lake level and an average of 2.03 m falling at higher elevations in the surrounding mountains (Fram, 2011). Extreme snow events in this area are common and often produce snowpack depths greater than 4.5 m at high elevations. Rain shadow effects typically lead to decreased snow loading on the downwind, eastern side of the basin. Approximately two-thirds of Lake Tahoe basin parent material is granitic and one-third is volcanic (LTTMDL, 2008). Vegetation, consisting of mixed coniferous forest and montane-subalpine species, cover approximately 80 % of the basin (LTTMDL, 2010). Areas of dense urban development occur along the shoreline at South Lake Tahoe, Tahoe City, and Incline Village. Large portions of the northern and western shores are occupied by seasonal cabins, while much of the eastern shore is undeveloped.

2.2 Sample collection

2.2.1 Integrated snowpack sampling: 2011–2012 and 2012–2013 snow years

During the 2011–2012 and 2012–2013 water years, full snowpack bi-weekly core samples were collected at seven sites in the basin starting from the first measureable snowpack until the majority of spring melting occurred (2011–2012: $n = 49$; 2012–2013: $n = 56$). This included mid-January through mid-April in 2012 and December to early-

BGD

12, 593–636, 2015

Nutrient and mercury deposition and storage in an alpine snowpack

C. Pearson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Nutrient and mercury deposition and storage in an alpine snowpack

C. Pearson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



April in 2012–2013. The seven sites were distributed along eastern and western elevation transects (Fig. 1). Three of the sites were located at lake level (one remote site; two sites in urban areas; elevation approximately 1900 m); two sites were at mid-mountain elevation (approximately 2200 m); and two sites were at high elevation close to the mountain ridges (elevation approximately 2500 m). To minimize throughfall input and allow better comparison among sites, we selected areas that were open, free of canopy coverage, and had minimal snowpack disturbance (i.e., away from congested areas).

Samples were collected using a Mt. Rose Federal Sampler and were immediately transferred to Whirl-pack[®] clean bags and a cooler with blue ice packs. Samples were transported within four hours to the Desert Research Institute in Reno, NV for storage at -20°C until laboratory analysis could be completed. Depth and snow water equivalent (SWE) were measured for each core using the Mt. Rose Federal Sampler. In cases of low snow accumulation, multiple cores were collected and homogenized to provide sufficient sample for all analyses. During collection, sterile gloves were worn, and soil contact and contamination were avoided in order to capture only constituents stored within the snowpack. While sampling, the first core taken at each site was discarded in order to avoid carryover from previous sampling. Between each sampling campaign, the Federal Sampler was cleaned with Milli-Q deionized water ($< 18.2\text{ M}\Omega$) and a chelating soap in accordance with trace metal sampling procedures (EPA, 2002). Field blanks were measured by rinsing the sampler with Milli-Q water prior to each sampling campaign.

2.2.2 Wet deposition sampling and snow pit collection: 2013–2014 snow year

In order to differentiate between snowpack storage and wet deposition and further assess dynamics in snowpack, additional sampling of full snow pit profiles and wet deposition was completed during the 2013–2014 snow year. Bi-weekly wet deposition sample collection following National Atmospheric Deposition Program protocol (<http://nadp.sws.uiuc.edu/>) was conducted at the two high altitude sites by N-Con dual

port trace metal samplers (Model TM 00-127; N-Con Inc., Crawford, GA, USA). These samplers allowed for collection of real-time wet deposition samples of both nutrients (N, P, and S) and Hg without cross contamination. The sample trains consists of NADP standard (19-128) glass funnel, glass anti-evaporation capillary, glass sample bottle (2 L) for collection of Hg, and a (19-130) polyethylene funnel with connector and 1.5 L HDPE sample container for nutrients. The glass sample bottle was pre-charged with 20 mL of deionized water and 0.5 mL of 12 M HCl (EMD Omnitrace HX0607) to act as a preservative for Hg. Sample bottles were collected in the field and kept in a cooler during transport back to the Desert Research Institute in Reno, NV. Sample bottles were then weighed in the lab and decanted into 250 mL HDPE bottles for nutrient samples and glass containers for Hg samples. All samples were stored in refrigerators until processing.

Three snow pit analyses were conducted at the high elevation sites, two near the Mt. Rose site (1 March 2014 and 4 April 2014) and one at the Homewood High site (28 February 2014). The snow pit measured a minimum of a 1.5 m² and was dug from the snow surface to the ground. Measuring sticks were placed on either side of the pit face. A measurement of height, layer density, and crystal form was noted. Snow samples were collected vertically every 10 cm using a 1000 cm³ Kelly wedge cutter (Model: RIP 1 Cutter; Snowmetrics, Fort Collins, CO, USA). Prior to collection, the acid washed wedge was inserted into the snow adjacent to the sample wall two to three times at each layer before sampling to avoid carry over. Duplicate samples were collected at each height and analyzed separately. All samples were double bagged in Whirl-pack[®] clean bags and weighed for density. Samples were then transferred to -20 °C storage at the Desert Research Institute in Reno, NV until analysis. Reported concentrations and densities are averages of the duplicate samples.

BGD

12, 593–636, 2015

Nutrient and mercury deposition and storage in an alpine snowpack

C. Pearson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.3 Laboratory analysis

Samples were analyzed for nitrite (NO_2^- -N), nitrate (NO_3^- -N), total ammonia nitrogen (TAN; $\text{NH}_3 + \text{NH}_4^+$), total Kjeldhal nitrogen (TKN), orthophosphate (o-PO_4), total phosphorus (TP), total Hg (THg, no filtration), and dissolved Hg (DHg, filtration). Prior to analysis, all samples were removed from the freezer and placed in a dark cabinet at room temperature for approximately 18 h to melt. Once fully melted, the samples were thoroughly mixed and dispensed into various aliquots for each analysis. Subsamples of NO_2^- -N, NO_3^- -N, TAN, SO_4^{2-} , and o-PO_4 were filtered through 0.45 μm filters (Pall Supor[®]) prior to analysis. Laboratory filter blanks were approximately $< 2 \mu\text{g L}^{-1}$ for NO_2^- -N, $6 \mu\text{g L}^{-1}$ for NO_3^- -N, $5 \mu\text{g L}^{-1}$ for TAN, $< 20 \mu\text{g L}^{-1}$ for SO_4^{2-} , and $2 \mu\text{g L}^{-1}$ for o-PO_4 .

Ortho-phosphate and TP were measured according to EPA Standard Method (SM) 365.1 and SM 365.1/USGS I-4600-85, respectively (EPA, 1993; USGS, 1985). Method detection limits (MDL) for these techniques were 0.60 and $0.63 \mu\text{g L}^{-1}$, respectively. Both techniques employed colorimetric measurement with ascorbic acid. Prior to measurement of TP, samples were digested with persulfate. Absorbance was then measured through flow injection analysis (FIA; Rapid Flow Analyzer 300 equipped with an Astoria-Pacific 305D high sensitivity photometer detector; Alpkem, College Station, TX).

Nitrite, NO_3^- -N, and TAN analyses followed EPA SM 353.2 and SM 353.1 (EPA, 1979, 1993). Nitrite and NO_3^- -N MDL were $0.84 \mu\text{g L}^{-1}$, and the TAN MDL was $0.77 \mu\text{g L}^{-1}$. Nitrite and NO_3^- -N were measured by automated colorimetric analysis with cadmium reduction being applied for the nitrate samples. Each sample was then measured by FIA (Rapid Flow Analyzer 300 equipped with an Astoria-Pacific 305D high sensitivity photometer detector; Alpkem, College Station, TX). TAN samples were analyzed using automated phenate colorimetric techniques. Total Kjeldhal Nitrogen was analyzed using automated phenate block digestion according to EPA method 351.2. The MDL for TKN was $11.3 \mu\text{g L}^{-1}$. Organic N (bulk) was calculated as the difference between

BGD

12, 593–636, 2015

Nutrient and mercury deposition and storage in an alpine snowpack

C. Pearson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



TKN and TAN. All nitrogen species were reported as $[\mu\text{g L}^{-1}]$ -N with total N calculated as the sum of organic N, TAN, and NO_3^- -N. All snow sample NO_2^- concentrations were below the DL.

Sulfate was determined using a chromatography system (ICS 2000 with Chromeleon Software version 6.6 and AS14A Column; Dionex Inc., Sunnyville, CA) by EPA Method 300.0 (EPA, 1979). The MDL for SO_4^{2-} was $19 \mu\text{g L}^{-1}$.

Total Hg and DHg were measured using a water analyzer (Model 2600; Tekran Inc., Toronto, Canada) according to EPA method 1631 revision E (EPA, 2002). For DHg samples, approximately 50 mL of sample were filtered through a $0.45 \mu\text{m}$ filter (Acrodisc syringe filter with Supor[®] Membrane; Pall Corporation, Port Washington, NY) while for THg, 50 mL of sample were poured directly into a vial for analysis. Laboratory filter blanks were below the detection limit (DL) of the system ($< 0.3 \text{ ng L}^{-1}$). Samples were preserved with 10 % bromine chloride (BrCl) solution for storage until analysis the next day. Before analysis, excess BrCl was neutralized with pre-purified hydroxylamine hydrochloride. During analysis, samples were automatically mixed with stannous chloride (SnCl_2) in a phase separator; reducing oxidized Hg to elemental Hg. Elemental Hg is then loaded onto two sequential gold traps by an argon carrier gas. The Hg is then released through thermal desorption and detected using atomic fluorescence spectrometry. The Tekran Model 2600 was calibrated using a NIST SRM-3133 Hg standard (with concentrations of 0, 0.5, 1.0, 5.0, 10.0, 25.0, and 50.0 ng L^{-1} Hg). System reliability was checked using ongoing precision recovery injections of 5 ng L^{-1} throughout each run and ranged between 87 to 112 % recovery. Reagent blanks measured regularly throughout each run ensured no contamination of the system. DLs calculated as three times the SD of the calibration blanks, averaged 0.3 ng L^{-1} for all runs. Particulate Hg was calculated as the difference between THg and DHg.

BGD

12, 593–636, 2015

Nutrient and mercury deposition and storage in an alpine snowpack

C. Pearson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.4 Statistics

We performed analysis of variance (ANOVA) for all chemical species using the following independent variables: (i) year ($n = 2$, 2011–2012 and 2012–2013), (ii) site elevation ($n = 3$; low, mid- and high elevation site), (iii) location ($n = 2$; eastern and western basin); and season ($n = 2$; early season [December through February] and late season [March and April]). ANOVAs attribute variance of dependent variables to these various independent variables and test their significance against the residual variance. All relationships were considered statistically significant when p values were ≤ 0.05 .

Integrated snowpack concentrations were calculated by weighting each 10 cm snow pit layer by its density. Seasonal wet deposition was calculated by weighting all wet deposition samples by their volume up to the date of sampling. Linear regression analyses were performed to test for correlations between snowpack chemical concentrations, SWE, and elevation. All error bars in figures represent standard error.

2.5 Basin-wide modeling with SWE reconstruction

Basin-wide loads and distribution were assessed using chemical concentrations and loads measured throughout the 2011–2012 and 2012–2013 snow seasons as well as basin-wide mean peak SWE estimates from SWE reconstruction for the Sierra Nevada from 2000 to 2011 (Rittger, 2012). SWE reconstruction uses estimates of snow energy balance with areal snow cover depletion from MODIS Snow Covered Area and Grain size (MODSCAG) (Rittger, 2011). MODSCAG calculates fractional snow cover area and grain size from Moderate Resolution Imaging Spectroradiometer (MODIS) data (Painter et al., 2009). Compared with previous methods, MODSCAG has proven to give reliable depletion rates throughout the spring season when snowmelt is highest (Rittger et al., 2013). Finally, the spatially refined MODSCAG data set was combined with energy balance and temperature data to give accurate reconstructed estimates of SWE throughout the Sierra Nevada, and specifically the Lake Tahoe Basin. At the time of our study, SWE reconstruction data were only available for 2000 to 2011, with

BGD

12, 593–636, 2015

Nutrient and mercury deposition and storage in an alpine snowpack

C. Pearson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Nutrient and mercury deposition and storage in an alpine snowpack

C. Pearson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



no information from our sampling seasons, 2011–2012 or 2012–2013. The 2000 to 2011 data set includes both high and low accumulation snow years and gives a reasonable representation of average snowpack accumulation in the Lake Tahoe basin. In order to give an estimate of average annual snowpack chemical storage, we applied the decadal average peak SWE for 2000 to 2011 to our data (Fig. 3a). Estimates made during this study were to establish relationships to previous estimates of the Lake Tahoe nutrient budget and were not meant to represent a completely accurate distribution or load stored within the basin's snowpack each year.

Snowpack sampling throughout the Lake Tahoe basin during 2011–2012 and 2012–2013 allowed for assessment of spatial and temporal chemical deposition patterns. Specifically, relationships to wet or dry deposition, in-basin or out-of-basin sources, and early or late season increases were identified. These deposition and source controls were then related to orographic characteristics to estimate chemical concentrations throughout the basin in unknown areas. A GIS land-use layer of the Tahoe Basin (LTTMDL, 2010) was applied in order to separate urban and non-urban locations with similar orographic characteristics for urban influenced species (i.e. TP). These scaled concentrations were then applied to SWE reconstruction estimates to determine total snowpack chemical loading throughout the entire basin.

Snowpack sampling occurred in open areas free of canopy coverage, but it is possible that tree and plant particulate matter still were incorporated in the snowpack. Litterfall contributions represent a form of chemical recycling and will cause an overestimate of atmospheric contributions made during this study. Visual inspection of snow samples, however, showed low contributions of plant detritus in samples, and due to consistent forest types present across the basin we would expect any additional plant-derived inputs to be random and unbiased across sites.

3 Results and discussion

3.1 Spatial and temporal trends of snow accumulation and SWE

In the Lake Tahoe basin, approximately 70 % of annual precipitation falls during the winter and spring as snow (Fram, 2011). The 2011–2012, 2012–2013, and 2013–2014 winter seasons were marked by relatively low snow accumulation. Peak basin average snowpack storage (1 April) for the central Sierra Nevada during 2011–2012, 2012–2013, and 2013–2014 was approximately 50, 53, and 41 % of the historical average (1951–present), respectively (CADWR, 2014). Although peak SWE was similar in each season, the temporal trends in snow accumulation and spatial distributions differed (Fig. 2). In 2011–2012, the Lake Tahoe Basin experienced low snowpack accumulation until the middle of January, when a series of storms led to solid snow cover throughout the basin. January storms were followed by a hiatus until late February and March when a series of storms brought peak basin average SWE up to approximately 625 mm. The 2012–2013 snow year started earlier, with late December storms bringing nearly 750 mm of SWE. Similar snowpack loading and timing occurred across the Lake Tahoe basin at sites with similar elevations (e.g., Mt. Rose/Squaw Valley, Marlette Lake/Rubicon). Early season storms were dominated by northerly wind patterns contributing substantial snowfall in the northeastern areas of the Lake Tahoe basin and reducing the typical pattern of lower snow accumulation on the eastern side of the basin due to the rain shadow effect of the Sierra Nevada crest (e.g., 2012–2013 Mt. Rose/Squaw Valley SNOTEL data). These early storms were followed by three dry months with very little accumulation for the rest of winter. The 2013–2014 snow year experienced the lowest snow accumulation of all three study years with minimal snowpack development occurring until late season storms in March and April brought peak SWE storage up to approximately 575 mm. Minimal snowpack development occurred at lower lake level elevations (e.g. Tahoe City SNOTEL data) throughout the entire 2013–2014 season.

Nutrient and mercury deposition and storage in an alpine snowpack

C. Pearson et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Nutrient and mercury deposition and storage in an alpine snowpack

C. Pearson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



However, vertical snow pit profile patterns show large variability in NO_3^- -N concentrations with depth, e.g. decreasing concentrations below the top 30–40 cm (Fig. 6). This variability suggests pronounced in snowpack dynamics possibly driven by conversion, vertical transport, or elution. In addition, several studies have shown significant wintertime dry deposition of NO_3^- -N, in particular close to highways and urban areas (Cape et al., 2004; Dasch and Cadle, 1986; Kirchner et al., 2005). Therefore, the fact that wet deposition concentrations were very similar to snowpack concentrations could be merely a coincidence and we do not feel confident to include or exclude significant dry deposition processes.

Finally, previous studies have observed parallel concentration declines of SO_4^{2-} and NO_3^- -N during snowpack melt events due to similar early elution characteristics (Stottlmyer and Rutkowski, 1990; Williams and Melack, 1991b). Comparing volume-weighted seasonal wet deposition concentrations of SO_4^{2-} to snowpack concentrations showed no elution losses through our sampling period and that SO_4^{2-} was also not subject to additional increases (Fig. 5). Our results therefore suggest that Sierra Nevada snowpack is subject to multiple inputs and complex in snowpack processes.

3.2.2 Total ammonia nitrogen (TAN)

Snowpack concentrations of TAN ranged from 16 to $104 \mu\text{g L}^{-1}$ ($n = 49$ cores), 10 to $77 \mu\text{g L}^{-1}$ ($n = 56$ cores), and 28 to $85 \mu\text{g L}^{-1}$ ($n = 3$ integrated snow pits) during 2011–2012, 2012–2013, and 2013–2014, respectively. Snowpack TAN concentrations are within the range of previous measurements made in the Emerald Lake Watershed of California, where the amount of TAN deposited within the seasonal snowpack accounted for approximately 90 % of annual loading (Williams et al., 1995).

Unlike NO_3^- -N, TAN is known to deposit through both wet and dry pathways during winter (Clow et al., 2002; Ingersoll et al., 2008). In our study, strong evidence for an important role of TAN dry deposition can be inferred from the fact that snowpack TAN concentrations doubled from early (December–February) to the late (March–April) sea-

son in both 2011–2012 and 2012–2013 (Fig. 4). ANOVA results confirmed significant differences in snowpack TAN concentrations between early and late season snowpack sampling (Table 1, $p = 0.01$). Increased late season TAN concentration in snowpack is consistent with similar observations in the Rocky Mountains and the Stubai Alps (Bowman, 1992; Kuhn, 2001). These increases were attributed to the onset of agricultural production in upwind valleys, as well as increased dry deposition due to decreased atmospheric stability and increased convection. Importantly, the late season increase in snowpack TAN occurred in both years, even though no significant late season snowfall occurred in 2012–2013 (Fig. 4). The patterns of increasing TAN concentration in late season snowpack with no significant snowfall agree with previous research showing dry deposition as the significant source of TAN deposition in the Sierra Nevada (Bytnerowicz and Fenn, 1996).

Large increases in NH_3 emissions from winter to spring have been measured upwind of the Sierra Nevada in the San Joaquin Valley, CA and were attributed to increased agricultural and livestock activities (Battye et al., 2003). Further support of snowpack TAN sourcing in the San Joaquin Valley, was higher concentrations at west basin sites than east basin sites during both 2011–2012 and 2012–2013. ANOVA results revealed a significant difference between the east and west basin snowpack TAN concentrations (Table 1, $p = 0.03$). This increase is likely due to the west basin sites being closer in proximity to San Joaquin Valley agricultural activity allowing for increased transport and deposition.

During the 2013–2014 snow year, TAN concentrations were consistently higher (up to a factor of 3) in volume-weighted wet deposition than integrated snow pit samples (Fig. 5; $p = 0.08$, note low replicate of $n = 3$). This increase of TAN further emphasizes the importance of dry deposition of TAN to Tahoe Basin snowpack. During snowpack storage, TAN is known to elute relatively late during melt events (Kuhn, 2001); however, other transformations such as microbial conversion can lead to decreases and losses throughout the season. Snow pit depth profile sampling shows a decrease in TAN concentrations with depth and therefore age (Fig. 6). This decrease coincides

BGD

12, 593–636, 2015

Nutrient and mercury deposition and storage in an alpine snowpack

C. Pearson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



with increases in organic N suggesting microbial conversion of inorganic N to organic N. Despite these possible losses; the increase we observe between wet deposition and snow pit concentrations indicates that the additional input of TAN from dry deposition is large enough to exceed transformations that occur during snowpack storage.

Late season deposition doubled TAN snowpack loads prior to end-of-season melt. The fate of snowpack TAN has been studied extensively through both watershed mass balance and tracer-based research. For example, less than 1 % of TAN stored in snowpack at Emerald Lake, California reached the lake as TAN during melt and runoff (Williams and Melack, 1991b). During a later study, however, snowmelt with isotopically labeled NH_3 was retained in the soils during melt making it a possible contributor to future NO_3^- stream pulses after nitrification (Williams et al., 1996). Current predictions show an increase in total N emissions during the next half-century in the western United States due to large increases in agricultural and livestock NH_3 emissions (Fenn et al., 2003). Such increased emissions could result in significant additional deposition loads of TAN to snowpack in the Sierra Nevada with the potential to alter ecosystem nutrient dynamics.

3.2.3 Organic nitrogen

Integrated snowpack core organic N concentrations ranged from 30 to $280 \mu\text{gL}^{-1}$ in 2011–2012 ($n = 49$ cores), 30 to $180 \mu\text{gL}^{-1}$ in 2012–2013 ($n = 56$ cores), and 120 to $260 \mu\text{gL}^{-1}$ in 2013–2014 ($n = 3$ integrated snow pit). No dominant spatial or temporal patterns were observed in snowpack organic N concentration or load for either 2011–2012 or 2012–2013 (Fig. 4). ANOVA results supported this finding with no significant effects of location, elevation, or early/late season on organic N concentrations (Table 1). A previous study found large variation in wintertime deposition of organic N throughout the Rocky Mountain Range; accounting for 40, 3, and 50 % of total N in wet deposition during January, February, and March, respectively (Benedict et al., 2013). Deposition rates and patterns of organic N are difficult to quantify due to the large number of compounds – including gaseous, particulate, and dissolved phases – origi-

BGD

12, 593–636, 2015

Nutrient and mercury deposition and storage in an alpine snowpack

C. Pearson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



organic to organic forms), and different mobilization during elution sequences leading to large fluctuations in both the concentration and spatial–temporal patterns of snowpack total N throughout the season. Total N accumulation in Sierra Nevada snowpack shows strong interannual variability as well as different representation of various N species and tracing N speciation throughout the snowpack season may give insight into microbial-driven snowpack chemical cycling.

3.3 Phosphorus

Snowpack TP concentrations ranged from 3 to 109 $\mu\text{g L}^{-1}$ in 2011–2012 ($n = 49$ cores), 3 to 59 $\mu\text{g L}^{-1}$ in 2012–2013 ($n = 56$ cores), and 10 to 41 $\mu\text{g L}^{-1}$ in 2013–2014 ($n = 3$ integrated snow pits). Figure 8 shows that the urban site in Incline Village at lake level had by far the highest snowpack TP concentrations, ranging up to six times higher than any other snowpack concentration at similar elevation (i.e. lake level). In comparison, the Thunderbird site also at lake level, located in a very remote setting just 10 km from Incline, had much lower P concentrations. Sources such as fugitive dust from plowing, forest and agriculture biomass burning, and diesel engine combustion have been identified as major sources of particulate-phase atmospheric P in California (Alexis, 2001). Specifically in the Lake Tahoe basin, road dust has been identified as a primary contributor of P input into Lake Tahoe (Dolislager et al., 2012), while another study found significant P emissions from urban biomass burning (Zhang et al., 2013). Our patterns suggest that urban areas in the Lake Tahoe basin are a major source area for P deposition to snowpack during winter and spring.

Local and regional emissions are also relevant at larger scales, as evident in 2011–2012, where remote sites at eastern locations in the basin showed higher TP concentrations than western sites. We propose that the large concentration of urban source sites at lake level combined with the dominant west to east wind pattern led to increased deposition on the east side of the basin. During 2012–2013, no west-to-east increase in TP concentration was observed; however, the strong influence of urban activity remained. It is unlikely that sources of P in the basin changed between 2011–

BGD

12, 593–636, 2015

Nutrient and mercury deposition and storage in an alpine snowpack

C. Pearson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



terial is granitic and one-third is volcanic (LTTMDL, 2008), making soil adsorption potentials of atmospherically deposited P throughout the watershed highly variable with location. Along with N, P levels directly control algal production within aquatic ecosystems, and algal production is a key reason for declining clarity in Lake Tahoe (Dolinger, 2006). In particular, the high snowpack concentrations at urban locations near the lake may cause a significant influx of P into Lake Tahoe during melt.

3.4 Mercury

Snowpack THg concentrations ranged from 0.81 to 7.58 ngL⁻¹ in 2011–2012 ($n = 49$ cores), 0.97 to 5.96 ngL⁻¹ in 2012–2013 ($n = 56$ cores), and 3.28 to 7.56 ngL⁻¹ in 2013–2014 ($n = 3$ integrated snow pits). Tahoe Basin average snowpack core THg concentration for 2011–2012 and 2012–2013 was 2.56 ± 1.3 ngL⁻¹. Observed THg concentrations are slightly lower, but within range of the end-of-season average snowpack concentration measured during a watershed Hg balance study in 2009 at Sagehen Creek, CA (i.e. 3.3 ngL⁻¹; Fain et al., 2011), a remote watershed located only 32 km north of the Tahoe Basin. Particulate Hg was the dominant form of Hg within Tahoe snowpack accounting for 76.1 ± 8.7 , 70.3 ± 13.4 , and 87.1 ± 4.7 of THg on average during 2011–2012, 2012–2013 and 2013–2014, respectively. The large percentage of particulate Hg in snowpack agrees with previous findings from a study in Canada that saw a post-depositional increase in particulate associated Hg from approximately 50 to 70% (Poulain et al., 2007). This study attributed particulate throughfall and photochemical induced emission as the main causes of the speciation shift and also noted strong differences in snowpack Hg concentrations between open and forested areas which were attributed to throughfall contributions from tree canopies as well as shading reducing photochemical evasion.

Snowpack coring revealed no dominant temporal or spatial patterns in THg or DHg deposition with ANOVA results showing no significant effects of season (i.e., early vs. late) or location (i.e., east vs. west; Table 1). The lack of either temporal or spatial trends suggests that global background atmospheric pollution, rather than specific

BGD

12, 593–636, 2015

Nutrient and mercury deposition and storage in an alpine snowpack

C. Pearson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Nutrient and mercury deposition and storage in an alpine snowpack

C. Pearson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Further support of photochemical reemission of Hg during snowpack storage can be inferred from the consistently lower DHg concentrations (up to a factor of 4.5) in integrated snow pit samples than volume-weighted wet deposition samples (Fig. 5). Photochemical reduction and volatile reemission of gaseous Hg during snowpack storage has been widely studied and is known to account for losses of up to 50 % from that measured in initial deposition (Fain et al., 2007, 2011; Lalonde et al., 2002; Mann et al., 2011; Poulain et al., 2007). In addition to the declines of DHg during storage, increase in particulate Hg was observed in two of the three comparisons of snow pit and wet deposition samples (Fig. 5). This increase in particulate Hg in snow pit samples is likely due to gaseous-dry deposition and particulate throughfall during storm free periods.

After photochemical losses, there is still a substantial amount of Hg left in the snowpack that will be subject to melt and infiltration into the watershed. The study at the nearby Sagehen Creek, California watershed quantified that only 4 % of total annual Hg wet deposition was exported from the watershed in stream water and identified soil uptake and storage as well as photochemical re-emission as the major sinks of atmospherically deposited Hg (Fain et al., 2011). While soil uptake serves as a buffer delaying the transport of upland wet deposition to streams, sediment core analyses still showed that upland watershed contributions (i.e., through soil erosion and sediment flux) are significant contributors of Hg input to lakes even under relatively low watershed to lake area ratios as in the Lake Tahoe basin (extrapolated to 42 % contributions when using relationships presented by Lorey and Driscoll, 1999). Snowpack-based Hg input to the watershed, therefore, is expected to contribute to lake water quality through erosion and sediment-based influx, albeit delayed in time and closely linked to soil Hg pools and mobilization.

3.5 Basin-wide loading estimates

Declines in Lake Tahoe water quality have been observed during the last 50 years (Sahoo et al., 2010; Schuster and Grismer, 2004). Specifically, secchi depths, a measure of lake transparency, have decreased from approximately 30.5 to 21.3 m since the 1960s

(TERC, 2011). Eutrophication from atmospheric and terrestrial nitrogen (N) and phosphorus (P) inputs as well as light scattering by particulate inputs are the main causes of this decline (Jassby et al., 2003; Swift et al., 2006). Most previous studies in the Lake Tahoe basin have focused on direct atmospheric deposition to the lake surface (Dolislager et al., 2012; NADP, 2012), and little information is available on snowpack-based loading for the surrounding upland watershed. The surrounding land surface covers 814 km² of the 1310 km² Lake Tahoe watershed. Direct atmospheric inputs to the lake surface are estimated to contribute 55 and 15 % of total N and P, respectively (TERC, 2011). Stream monitoring data show that upon snowmelt, Lake Tahoe receives large pulses of N and P (Goldman et al., 1989; Hatch et al., 1999), which together control algal production within the basin's aquatic ecosystems contributing to the decline in clarity in Lake Tahoe during the last 50 years (Dolislager, 2006). Although much of snowpack-based chemical loads may not directly enter Lake Tahoe upon melt, snowpack loads are important for terrestrial chemical budgets. For example, nutrient rich O-horizon runoff – measuring as high as 87.2 mg L⁻¹ NH₄-N, 95.4 mg L⁻¹ NO₃-N, and 24.4 mg L⁻¹ PO₄-P – has been observed in Lake Tahoe forests during snowmelt events due to leaching from the forest litter layer (Miller et al., 2005). In order to relate peak snowpack nutrient and pollutant loading to previous terrestrial and lake chemical budgets, we here estimate average peak basin-wide snowpack chemical storage using the peak SWE decadal average from 2000–2011 (Fig. 3a).

3.5.1 Nitrogen

Snowpack NO₃⁻-N loading was highly dependent on snow accumulation, but concentrations showed little temporal or spatial trends throughout the Lake Tahoe basin (Table 1). To calculate basin-wide NO₃⁻-N loads, we therefore multiplied the two-year seasonal average concentration (47.1 µg L⁻¹) by the decadal average reconstructed SWE. Basin-wide NO₃⁻-N loading estimates (mass area⁻¹) thus reflect snowpack accumulation patterns (i.e., SWE) with the highest loading occurring on the west-side of the

BGD

12, 593–636, 2015

Nutrient and mercury deposition and storage in an alpine snowpack

C. Pearson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



basin at high elevations, up to approximately 1 kg ha^{-1} , and decreasing toward the east and with lower elevations due to lower SWE accumulation. Average annual snowpack NO_3^- -N storage for the Lake Tahoe basin is estimated at 28.7 t (metric tons).

Unlike NO_3^- -N, snowpack TAN loading in the Lake Tahoe basin showed strong spatial and temporal trends. Late season deposition effectively doubled snowpack TAN concentrations with significantly higher concentrations on the west side of the basin than the east side (Table 1). Due to these relationships, we applied the March and April (peak SWE generally occurs during March and April in the Lake Tahoe basin) average snowpack TAN concentration from the east and west basin sites to the reconstructed SWE data (western sites: $57.9 \mu\text{g L}^{-1}$, eastern sites: $41.6 \mu\text{g L}^{-1}$) to scale up snowpack TAN loading to the whole basin. Modeled estimates, therefore, show greater TAN accumulation on the western side of the basin with highest loading occurring at high elevations in the west (up to approximately 1.2 kg ha^{-1}) due to the combination of both large snow accumulation and proximity to upwind sources. Our estimate of average annual basin-wide accumulation of TAN within the basin's snowpack is 30.5 t.

Snowpack organic N concentrations throughout each sampling season were variable and showed no clear temporal or spatial trends (Table 1). Applying the average concentration of $88.7 \mu\text{g L}^{-1}$ from all snowpack samples throughout both years produced an annual estimate of 54.1 t of organic N stored within the basin's snowpack.

Average annual snowpack N storage for the Lake Tahoe watershed – calculated as the sum of NO_3^- -N, TAN, and organic N – totaled 113 t (Fig. 3b). Inorganic and organic forms made up 52 and 48 % of total N, respectively. TAN and NO_3^- -N accounted for 27 and 25 % of total snowpack N, respectively. Annual N loading estimates for Lake Tahoe (from terrestrial runoff and direct atmospheric deposition) were previously estimated to be 397 t yr^{-1} , with 218 t yr^{-1} originating from atmospheric sources depositing directly on the lake's surface (LTTMDL, 2010). With the caveat that estimation methods differed, snowpack N storage estimates from our study represent approximately 28 % of the lake's total N budget. Comparing our estimates to the 179 t of N that origi-

BGD

12, 593–636, 2015

Nutrient and mercury deposition and storage in an alpine snowpack

C. Pearson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



nates from runoff and terrestrial sources, annual snowpack N storage would replenish approximately 63% of this flux.

3.5.2 Phosphorus

Snowpack P accumulation in the Lake Tahoe basin was strongly related to proximity to urban sources, as well as transport along the dominant westerly winds throughout the basin. This dependence caused the highest P concentration in snowpack to occur in developed areas and higher concentrations across east basin sites than remote west basin sites (Table 1). Applying different P concentrations based on degree of urbanization (see Sect. 2.5), highest P loading (up to approximately 0.4 kg ha^{-1}) occurs, therefore, at high elevations with significant impacts of urban emissions (i.e., north-eastern and southern locations influenced by Incline Village, Nevada and South Lake Tahoe, California). The basin-wide average TP storage estimated during this study of 0.11 kg ha^{-1} is over double the average snowpack storage reported for the ELW watershed (0.04 kg ha^{-1} ; Sickman et al., 2003) and reflects increased urbanization within the Tahoe Basin. Homyak et al. (2014) estimate that atmospheric deposition has contributed up to 31% of P accumulation and loss in soils and runoff since deglaciation of the Emerald Lake Watershed. The higher snowpack loading rates estimated during this study indicate that atmospheric deposition could be the primary supplier of excess P input to the Tahoe Basin.

Overall, we estimate a peak P load of approximately 9.3 t of P stored annually in Lake Tahoe basin snowpack (Fig. 3c). Previous pollutant loading studies for Lake Tahoe estimated that approximately 46 t of P enters the lake each year with approximately 39 t of the annual budget originating from land-based sources (LTTMDL, 2010). Annual snowpack TP storage estimates, therefore, could represent approximately 20% of total P input into Lake Tahoe each year.

BGD

12, 593–636, 2015

Nutrient and mercury deposition and storage in an alpine snowpack

C. Pearson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3.5.3 Mercury

Similar to NO_3^- -N, snowpack THg concentrations showed little temporal or east to west variation (Table 1). However, THg concentrations were positively related to total SWE (slope: $0.00201 \text{ [ngL}^{-1} \text{ mm}^{-1}]$; p value: 0.016). Applying this relationship to reconstructed SWE data produced the following THg distribution throughout the Lake Tahoe basin (Fig. 3d); THg loading throughout the basin followed strong elevation gradients, with the uppermost areas of the basin receiving the highest concentrations and total loading (up to approximately 125 mg ha^{-1}) due to increased snow accumulation. Average annual snowpack THg concentration and loading for the Lake Tahoe watershed was 3.6 ng L^{-1} and 30 mg ha^{-1} , respectively, based on the decadal SWE accumulation average of 750 mm. We do not have any previous data on Hg deposition to this basin, but these values are comparable to the 3.3 ng L^{-1} average snowpack Hg concentration and 13 mg ha^{-1} peak snowpack loading from the Sagehen Creek watershed in 2009 when snowpack accumulation was approximately 400 mm (Fain et al., 2011). The basin-wide estimate of THg stored within the annual snowpack was 1166.2 g. Snow-based Hg fluxes estimated during this study fall within range of measurements (3.36 to $36 \text{ mg ha}^{-1} \text{ yr}^{-1}$) taken at seven national parks throughout western North America during the Western Airborne Contaminants Assessment Project (WACAP), which found fish Hg levels above the human consumption threshold even at sites with relatively low Hg deposition (Landers et al., 2008).

4 Conclusions

In summary, spatial and temporal pattern analyses suggest that out-of-basin sources were important for Hg and TAN, while in-basin sources controlled P deposition, with the highest concentrations measured near urban areas, exceeding remote location concentrations by up to a factor of 6. Snowpack NO_3^- -N concentrations were relatively uniform throughout the basin indicating out-of-basin sourced wet deposition as a pri-

BGD

12, 593–636, 2015

Nutrient and mercury deposition and storage in an alpine snowpack

C. Pearson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Nutrient and mercury deposition and storage in an alpine snowpack

C. Pearson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



5
10
15
20
25

mary input; however high variability in snow pit vertical concentrations suggests additional inputs and in snowpack transport and conversion processes. Second, increased NH_3 emissions from the San Joaquin Valley and increased atmospheric vertical mixing during the onset of spring likely led to dry deposition-based increases of snowpack TAN during March and April, effectively doubling snowpack TAN concentrations prior to melt. Third, chemical speciation showed that organic N in Lake Tahoe snowpack accounted for 48 % of total N on average with possible microbial conversion leading to higher enhanced organic N levels in deeper older snowpack. Fourth, particulate Hg was the dominant form of Hg (78 % on average) within Tahoe snowpack and concentrations of both THg and DHg increased with elevation and SWE likely due to decreased light penetration and reduced photochemical reemission in deeper snowpack. Finally, basin-wide modeling estimates indicated that Lake Tahoe basin snowpack acts as a substantial reservoir in which atmospheric nutrients and pollutants accumulated throughout winter and spring. Estimates of basin-wide annual snowpack mass loading showed accumulation of N, P, and Hg yielding 113 t of N, 9.3 t of P, and 1166.2 g of Hg. Further research should focus on quantifying the relationship between snowmelt processes and stream and groundwater input, and address the substantial amount of organic N stored within the basin's snowpack.

Acknowledgements. Special thanks to Kevin Mitchell and Ted Tank of Homewood Ski Resort and Scott Hackley of the Tahoe Environmental Research Center for providing access for sampling, Mary Miller for running all nutrient analysis, Chris Moore for Hg laboratory support, Tracy Backes and Hal Voepel for assistance with data analysis, as well as Scott Kobs and Ashley Pierce for help with sample collection.

The project described in this publication was supported by Grant/Cooperative Agreement Number G11AP20092 from the United States Geological Survey (USGS). Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the USGS. Additional funding and support were provided by DRI's Divisions of Atmospheric and Hydrologic Sciences as well as the National Institutes for Water Resources.

References

- Alexis, A., A. Delao, C. Garcia, M. Nystrom, and Rosenkranz, K.: The 2001 California Almanac of Emissions and Air Quality, California Air Resources Board, Callifornia Environmental Protection Agency, 2001.
- 5 Batty, W., Aneja, V. P., and Roelle, P. A.: Evaluation and improvement of ammonia emissions inventories, *Atmos. Environ.*, 37, 3873–3883, 2003.
- Benedict, K. B., Carrico, C. M., Kreidenweis, S. M., Schichtel, B., Malm, W. C., and Collett, J. L.: A seasonal nitrogen deposition budget for Rocky Mountain National Park, *Ecol. Appl.*, 23, 1156–1169, 2013.
- 10 Berg, N. H.: Ion elution and release from deep snowpacks in the central Sierra-Nevada, California, *Water Air Soil Poll.*, 61, 139–168, 1992.
- Bowman, W. D.: Inputs and storage of nitrogen in winter snowpack in an alpine ecosystem, *Arctic Alpine Res.*, 24, 211–215, 1992.
- Brooks, P., Williams, M., and Schmidt, S.: Microbial activity under alpine snowpacks, Niwot Ridge, Colorado, *Biogeochemistry*, 32, 93–113, 1996.
- 15 Brooks, P. D. and Williams, M. W.: Snowpack controls on nitrogen cycling and export in seasonally snow-covered catchments, *Hydrol. Process.*, 13, 2177–2190, 1999.
- Brown, L. J., Taleban, V., Gharabaghi, B., and Weiss, L.: Seasonal and spatial distribution patterns of atmospheric phosphorus deposition to Lake Simcoe, ON, *J. Great Lakes Res.*, 37, 15–25, 2011.
- 20 Bytnerowicz, A. and Fenn, M. E.: Nitrogen deposition in California forests: a review, *Environ. Pollut.*, 92, 127–146, 1996.
- CADWR: available at: <http://cdec.water.ca.gov/snow/current/snow/index.html> (last access: 9 November 2014), 2014.
- 25 Cape, J. N., Tang, Y. S., van Dijk, N., Love, L., Sutton, M. A., and Palmer, S. C. F.: Concentrations of ammonia and nitrogen dioxide at roadside verges, and their contribution to nitrogen deposition, *Environ. Pollut.*, 132, 469–478, 2004.
- Cape, J. N., Cornell, S. E., Jickells, T. D., and Nemitz, E.: Organic nitrogen in the atmosphere – where does it come from? A review of sources and methods, *Atmos. Res.*, 102, 30–48, 30 2011.

Nutrient and mercury deposition and storage in an alpine snowpack

C. Pearson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Nutrient and mercury deposition and storage in an alpine snowpack

C. Pearson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Clement, J. C., Robson, T. M., Guillemin, R., Saccone, P., Lochet, J., Aubert, S., and Lavorel, S.: The effects of snow-N deposition and snowmelt dynamics on soil-N cycling in marginal terraced grasslands in the French Alps, *Biogeochemistry*, 108, 297–315, 2012.
- Clow, D. W., Ingersoll, G. P., Mast, M. A., Turk, J. T., and Campbell, D. H.: Comparison of snowpack and winter wet-deposition chemistry in the Rocky Mountains, USA: implications for winter dry deposition, *Atmos. Environ.*, 36, 2337–2348, 2002.
- Dasch, J. M. and Cadle, S. H.: Dry deposition to snow in an urban area, *Water Air Soil Poll.*, 29, 297–308, 1986.
- Dodds, W. K.: Misuse of inorganic N and soluble reactive P concentrations to indicate nutrient status of surface waters, *J. N. Am. Benthol. Soc.*, 22, 171–181, 2003.
- Dolislager, L., Lashgari, A., Pederson, J., and VanCuren, J.: Lake Tahoe Atmospheric Deposition Study (LTADS), Atmospheric Processes Research Section Research Division, Air Resources Board, California Environmental Protection Agency, 2006.
- Dolislager, L. J., VanCuren, R., Pederson, J. R., Lashgari, A., and McCauley, E.: A summary of the Lake Tahoe Atmospheric Deposition Study (LTADS), *Atmos. Environ.*, 46, 618–630, 2012.
- EPA: Methods for Chemical Analysis of Water and Wastes. Environmental Monitoring and Support Laboratory, U.S. Environmental Protection Agency, Cincinnati, Ohio 45468, Revised 3/1983, 1979.
- EPA: Methods for the Determination of Inorganic Substances in Environmental Samples. Environmental Monitoring Systems Laboratory, U.S. Environmental Protection Agency, Cincinnati, Ohio 45268, 1993.
- EPA: Method 1631, Revision E: Mercury in Water by Oxidation, Purge and Trap, and Cold Vapor Atomic Fluorescence Spectrometry, U.S. Environmental Protection Agency, Washington DC 20460, 2002.
- Fain, X., Grangeon, S., Bahlmann, E., Fritsche, J., Obrist, D., Dommergue, A., Ferrari, C. P., Cairns, W., Ebinghaus, R., Barbante, C., Cescon, P., and Boutron, C.: Diurnal production of gaseous mercury in the alpine snowpack before snowmelt, *J. Geophys. Res.-Atmos.*, 112, doi:10.1029/2007JD008520, 2007.
- Fain, X., Obrist, D., Pierce, A., Barth, C., Gustin, M. S., and Boyle, D. P.: Whole-watershed mercury balance at Sagehen Creek, Sierra Nevada, CA, *Geochim. Cosmochim. Ac.*, 75, 2379–2392, 2011.

Nutrient and mercury deposition and storage in an alpine snowpack

C. Pearson et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



- Fenn, M. E., Haeuber, R., Tonnesen, G. S., Baron, J. S., Grossman-Clarke, S., Hope, D., Jaffe, D. A., Copeland, S., Geiser, L., Rueth, H. M., and Sickman, J. O.: Nitrogen emissions, deposition, and monitoring in the western United States, *Bioscience*, 53, 391–403, 2003.
- Fierz, C., Armstrong, R. L., Durand, Y., Etchevers, P., Greene, E., McClung, D. M., Nishimura, K., Satyawali, P. K., and Sokratov, S. A.: The International Classification for Seasonal Snow on the Ground (ICSSG), International Association of Cryospheric Sciences, Paris, 90 pp., 2009.
- Fram, M. S. and Belitz, K.: Groundwater Quality in the Tahoe and Martis Basins, California, US Geologic Survey Fact Sheet, 4, 2011.
- Galbavy, E. S., Anastasio, C., Lefer, B., and Hall, S.: Light penetration in the snowpack at Summit, Greenland: part 2 nitrate photolysis, *Atmos. Environ.*, 41, 5091–5100, 2007.
- Goldman, C. R., Jassby, A., and Powell, T.: Interannual fluctuations in primary productions – meteorological forcing at 2 subalpine lakes, *Limnol. Oceanogr.*, 34, 310–323, 1989.
- Gunda, T. and Scanlon, T. M.: Topographical influences on the spatial distribution of soil mercury at the catchment scale, *Water Air Soil Poll.*, 224, 2013.
- Halsall, C. J.: Investigating the occurrence of persistent organic pollutants (POPs) in the arctic: their atmospheric behaviour and interaction with the seasonal snow pack, *Environ. Pollut. (Barking, Essex: 1987)*, 128, 163–175, 2004.
- Hatch, L. K., Reuter, J. E., and Goldman, C. R.: Daily phosphorus variation in a mountain stream, *Water Resour. Res.*, 35, 3783–3791, 1999.
- Homyak, P. M., Sickman, J. O., and Melack, J. M.: Pools, transformations, and sources of P in high-elevation soils: implications for nutrient transfer to Sierra Nevada lakes, *Geoderma*, 217–218, 65–73, 2014.
- Ingersoll, G. P., Mast, M. A., Campbell, D. H., Clow, D. W., Nanus, L., and Turk, J. T.: Trends in snowpack chemistry and comparison to National Atmospheric Deposition Program results for the Rocky Mountains, US, 1993–2004, *Atmos. Environ.*, 42, 6098–6113, 2008.
- Jacobi, H. W. and Hilker, B.: A mechanism for the photochemical transformation of nitrate in snow, *J. Photoch. Photobio. A*, 185, 371–382, 2007.
- Jassby, A. D., Reuter, J. E., Axler, R. P., Goldman, C. R., and Hackley, S. H.: Atmospheric deposition of nitrogen and phosphorus in the annual nutrient load of Lake Tahoe (California Nevada), *Water Resour. Res.*, 30, 2207–2216, 1994.

Nutrient and mercury deposition and storage in an alpine snowpack

C. Pearson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Jassby, A. D., Reuter, J. E., and Goldman, C. R.: Determining long-term water quality change in the presence of climate variability: Lake Tahoe (USA), *Can. J. Fish. Aquat. Sci.*, 60, 1452–1461, 2003.

Jones, H. G.: The ecology of snow-covered systems: a brief overview of nutrient cycling and life in the cold, *Hydrol. Process.*, 13, 2135–2147, 1999.

King, M. D. and Simpson, W. R.: Extinction of UV radiation in Arctic snow at Alert, Canada (82° N), *J. Geophys. Res.-Atmos.*, 106, 12499–12507, 2001.

Kirchner, M., Jakobi, G., Feicht, E., Bernhardt, M., and Fischer, A.: Elevated NH₃ and NO₂ air concentrations and nitrogen deposition rates in the vicinity of a highway in southern Bavaria, *Atmos. Environ.*, 39, 4531–4542, 2005.

Kuhn, M.: The nutrient cycle through snow and ice, a review, *Aquat. Sci.*, 63, 150–167, 2001.

Lalonde, J. D., Poulain, A. J., and Amyot, M.: The role of mercury redox reactions in snow on snow-to-air mercury transfer, *Environ. Sci. Technol.*, 36, 174–178, 2002.

Landers, D. H., Simonich, S. L., Jaffe, D. A., Geiser, L. H., Campbell, D. H., Schwindt, A. R., Schreck, C. B., M. L. K., Hafner, W. D., Taylor, H. E., Hageman, K. J., Usenko, S., L. K., Ackerman, J. E. S., Rose, N. L., Blett, T. F., and Erway, M. M.: The Fate, Transport, and Ecological Impacts of Airborne Contaminants in Western Nation Parks (USA), EPA/600/R-07/138, US Environmental Protection Agency, Office of Research and Development, NHEERL, Western Ecology Division, Corvallis, Oregon, 2008.

Larose, C., Dommergue, A., and Vogel, T. M.: Microbial nitrogen cycling in Arctic snowpacks, *Environ. Res. Lett.*, 8, 035004, 2013.

Lorey, P. and Driscoll, C. T.: Historical trends of mercury deposition in Adirondack lakes, *Environ. Sci. Technol.*, 33, 718–722, 1999.

LTTMDL: Lake Tahoe TMDL Pollutant Reduction Opportunity Report, California Water Boards, Nevada Division of Environmental Protection, 2008.

LTTMDL: Lake Tahoe Total Maximum Daily Load Report 2010, Nevada Division of Environmental Protection, 2010.

Mann, E., Meyer, T., Mitchell, C. P. J., and Wania, F.: Mercury fate in ageing and melting snow: development and testing of a controlled laboratory system, *J. Environ. Monitor.*, 13, 2695–2702, 2011.

McDaniel, M. R.: Semivolatile Organic Compounds in Snowmobile Emissions and in the Snowpack and Surface Water in Blackwood Canyon, Lake Tahoe, CA, 3566275, Ph.D., University of Nevada, Reno, Ann Arbor, 137 pp., 2013.

Nutrient and mercury deposition and storage in an alpine snowpack

C. Pearson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Melack, J. M. and Stoddard, J. L.: Sierra Nevada in: Acidic Deposition and Aquatic Ecosystems: Regional Case Studies, edited by: Charles, D. F., Springer-Verlag, New York, 503–530, 1991.
- Miller, W. W., Johnson, D. W., Denton, C., Verburg, P. S. J., Dana, G. L., and Walker, R. F.: Inconspicuous nutrient laden surface runoff from mature forest Sierran watersheds, *Water Air Soil Poll.*, 163, 3–17, 2005.
- Morales-Baquero, R., Pulido-Villena, E., and Reche, I.: Atmospheric inputs of phosphorus and nitrogen to the southwest Mediterranean region: biogeochemical responses of high mountain lakes, *Limnol. Oceanogr.*, 51, 830–837, 2006.
- NADP: National Atmospheric Deposition Program (NADP), National Trends Network, 2012.
- Neff, J., Holland, E., Dentener, F., McDowell, W., and Russell, K.: The origin, composition and rates of organic nitrogen deposition: a missing piece of the nitrogen cycle?, *Biogeochemistry*, 57–58, 99–136, 2002.
- NRCS: available at: <http://www.wcc.nrcs.usda.gov/snow/> (last access: 9 November 2014), 2013.
- O'Hara, B. F., Kaplan, M. L., and Underwood, S. J.: Synoptic climatological analyses of extreme snowfalls in the Sierra Nevada, *Weather Forecast.*, 24, 1610–1624, 2009.
- Obrist, D., Johnson, D. W., and Lindberg, S. E.: Mercury concentrations and pools in four Sierra Nevada forest sites, and relationships to organic carbon and nitrogen, *Biogeosciences*, 6, 765–777, doi:10.5194/bg-6-765-2009, 2009.
- Obrist, D., Johnson, D. W., Lindberg, S. E., Luo, Y., Hararuk, O., Bracho, R., Battles, J. J., Dail, D. B., Edmonds, R. L., Monson, R. K., Ollinger, S. V., Pallardy, S. G., Pregitzer, K. S., and Todd, D. E.: Mercury distribution across 14 US forests. Part I: Spatial patterns of concentrations in biomass, litter, and soils, *Environ. Sci. Technol.*, 45, 3974–3981, 2011.
- Painter, T. H., Rittger, K., McKenzie, C., Slaughter, P., Davis, R. E., and Dozier, J.: Retrieval of subpixel snow covered area, grain size, and albedo from MODIS, *Remote Sens. Environ.*, 113, 868–879, 2009.
- Poissant, L. and Pilote, M.: Mercury concentrations in single event precipitation in southern Quebec, *Sci. Total Environ.*, 213, 65–72, 1998.
- Poulain, A. J., Roy, V., and Amyot, M.: Influence of temperate mixed and deciduous tree covers on Hg concentrations and photoredox transformations in snow, *Geochim. Cosmochim. Ac.*, 71, 2448–2462, 2007.
- Raison, R. J., Khanna, P. K., and Woods, P. V.: Mechanisms of element transfer to the atmosphere during vegetation fires, *Can. J. Forest Res.*, 15, 132–140, 1985.

Nutrient and mercury deposition and storage in an alpine snowpack

C. Pearson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Rittger, K.: Spatial Estimates of Snow Water Equivalent in the Sierra Nevada, Ph. D. Thesis, University of California, Santa Barbara, 225, 2012.
- Rittger, K., Kahl, A., and Dozier, J.: Topographic Distribution of Snow Water Equivalent in the Sierra Nevada, Proceedings, Western Snow Conference 79, 37–46, 2011.
- 5 Rittger, K., Painter, T. H., and Dozier, J.: Assessment of methods for mapping snow cover from MODIS, *Adv. Water Resour.*, 51, 367–380, 2013.
- Rohrbough, J. A., Davis, D. R., and Bales, R. C.: Spatial variability of snow chemistry in an alpine snowpack, southern Wyoming, *Water Resour. Res.*, 39, doi:10.1029/2003WR002067, 2003.
- 10 Rothlisberger, R., Hutterli, M. A., Wolff, E. W., Mulvaney, R., Fischer, H., Bigler, M., Goto-Azuma, K., Hansson, M. E., Ruth, U., Siggaard-Andersen, M. L., and Steffensen, J. P.: Nitrate in Greenland and Antarctic ice cores: a detailed description of post-depositional processes, in: *Annals of Glaciology*, Vol. 35, *Annals of Glaciology-Series*, edited by: Wolff, E. W., Int. Glaciological Soc., Cambridge, 206–216, 2002.
- 15 Sahoo, G. B., Schladow, S. G., and Reuter, J. E.: Effect of sediment and nutrient loading on Lake Tahoe optical conditions and restoration opportunities using a newly developed lake clarity model, *Water Resour. Res.*, 46, doi:10.1029/2009WR008447, 2010.
- Schroeder, W. H. and Munthe, J.: Atmospheric mercury – an overview, *Atmos. Environ.*, 32, 809–822, 1998.
- 20 Schuster, S. and Grismer, M. E.: Evaluation of water quality projects in the Lake Tahoe basin, *Environ. Monit. Assess.*, 90, 225–242, 2004.
- Sickman, J. O., Leydecker, A., and Melack, J. M.: Nitrogen mass balances and abiotic controls on N retention and yield in high-elevation catchments of the Sierra Nevada, California, United States, *Water Resour. Res.*, 37, 1445–1461, 2001.
- 25 Sickman, J. O., Melack, J. M., and Clow, D. W.: Evidence for nutrient enrichment of high-elevation lakes in the Sierra Nevada, California, *Limnol. Oceanogr.*, 48, 1885–1892, 2003.
- Stottlmyer, R. and Rutkowski, D.: Multiyear trends in snowpack ion accumulation and loss, northern Michigan, *Water Resour. Res.*, 26, 721–737, 1990.
- Swift, T. J., Perez-Losada, J., Schladow, S. G., Reuter, J. E., Jassby, A. D., and Goldman, C. R.:
 30 Water clarity modeling in Lake Tahoe: linking suspended matter characteristics to Secchi depth, *Aquat. Sci.*, 68, 1–15, 2006.
- TERC: State of the Lake Report 2011, UC Davis Tahoe Environmental Reserach Center, 2011.

Nutrient and mercury deposition and storage in an alpine snowpack

C. Pearson et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Turk, J. T., Taylor, H. E., Ingersoll, G. P., Tonnessen, K. A., Clow, D. W., Mast, M. A., Campbell, D. H., and Melack, J. M.: Major-ion chemistry of the Rocky Mountain snowpack, USA, *Atmos. Environ.*, 35, 3957–3966, 2001.

USGS: Methods for the Determination of Inorganic Substances in Water and Fluvial Sediments, US Government Printing Office, Washington DC, 20402, 1985.

Vicars, W. C. and Sickman, J. O.: Mineral dust transport to the Sierra Nevada, California: loading rates and potential source areas, *J. Geophys. Res.-Biogeosci.*, 116, doi:10.1029/2010JG001394, 2011.

Vicars, W. C., Sickman, J. O., and Ziemann, P. J.: Atmospheric phosphorus deposition at a montane site: size distribution, effects of wildfire, and ecological implications, *Atmos. Environ.*, 44, 2813–2821, 2010.

White, A. F., Bullen, T. D., Vivit, D. V., Schulz, M. S., and Clow, D. W.: The role of disseminated calcite in the chemical weathering of granitoid rocks, *Geochim. Cosmochim. Ac.*, 63, 1939–1953, 1999.

Williams, M. W. and Melack, J. M.: Precipitation chemistry in and ionic loading to an alpine basin, Sierra-Nevada, *Water Resour. Res.*, 27, 1563–1574, 1991a.

Williams, M. W. and Melack, J. M.: Solute chemistry of snowmelt and runoff in an alpine basin, Sierra-Nevada, *Water Resour. Res.*, 27, 1575–1588, 1991b.

Williams, M. W., Bales, R. C., Brown, A. D., and Melack, J. M.: Fluxes and transformations of nitrogen in a high-elevation catchment, Sierra-Nevada, *Biogeochemistry*, 28, 1–31, 1995.

Williams, M. W., Brooks, P. D., Mosier, A., and Tonnessen, K. A.: Mineral nitrogen transformations in and under seasonal snow in a high-elevation catchment in the Rocky Mountains, United States, *Water Resour. Res.*, 32, 3161–3171, 1996.

Williams, M. W., Hood, E., and Caine, N.: Role of organic nitrogen in the nitrogen cycle of a high-elevation catchment, Colorado Front Range, *Water Resour. Res.*, 37, 2569–2581, 2001.

Zhang, Y. Y., Obrist, D., Zielinska, B., and Gertler, A.: Particulate emissions from different types of biomass burning, *Atmos. Environ.*, 72, 27–35, 2013.




Figure 1. Lake Tahoe watershed map with bi-weekly sampling sites located along east and west basin elevation gradients for spatial and temporal sampling campaigns in 2011–2012 and 2012–2013. Additional wet deposition and snow pit profile samples were collected near the Homewood High and Mt. Rose sites during the 2013–2014 snow year.

Nutrient and mercury deposition and storage in an alpine snowpack

C. Pearson et al.

[Title Page](#)

[Abstract](#) | [Introduction](#)

[Conclusions](#) | [References](#)

[Tables](#) | [Figures](#)

[◀](#) | [▶](#)

[◀](#) | [▶](#)

[Back](#) | [Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



BGD

12, 593–636, 2015

Nutrient and mercury deposition and storage in an alpine snowpack

C. Pearson et al.




Figure 2. Snow water equivalent measured in 2011–2012, 2012–2013, and 2013–2014 at select SNOTEL sites (NRCS, 2013) throughout the Lake Tahoe Basin.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)






Figure 3. (a) Decadal average (2000–2011) peak SWE for the Tahoe Basin from SWE reconstruction for the Sierra Nevada; basin-wide peak snowpack chemical loading estimates for (b) nitrogen, (c) total phosphorus, and (d) total Hg.

Nutrient and mercury deposition and storage in an alpine snowpack

C. Pearson et al.




Figure 4. Average snowpack core concentrations during 2011–2012 (left) and 2012–2013 (right) snow seasons along with average SWE estimated from six SNOTEL sites located within the Tahoe Basin.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Figure 5. Comparison of seasonal average volume-weighted wet deposition concentrations with integrated snow pit samples from the 2013–2014 snow year.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Nutrient and mercury deposition and storage in an alpine snowpack

C. Pearson et al.




Figure 6. 2013–2014 snow pit profiles for nitrogen species concentration, snow density, and crystal form. Crystal classifications are based on the ICSI classification for seasonal snow on the ground (Fierz, 2009).

[Title Page](#)
[Abstract](#) [Introduction](#)
[Conclusions](#) [References](#)
[Tables](#) [Figures](#)
⏪ ⏩
◀ ▶
[Back](#) [Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)



Nutrient and mercury deposition and storage in an alpine snowpack

C. Pearson et al.




Figure 7. Snowpack total N distribution for 2011–2012 (left) and 2012–2013 (right).

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)






Figure 8. Average snowpack total phosphorus concentrations at all lake-level sites. The Incline and Thunderbird sites are located on the east side of the basin in urban and remote settings, respectively, and the Homewood Low site is located on the west side of the basin in an urban setting.

Nutrient and mercury deposition and storage in an alpine snowpack

C. Pearson et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

