

Characterization of active and total fungal communities in the atmosphere over the Amazon rainforest

A. M. Womack, P. E. Artaxo, F. Y. Ishida, R. C. Mueller, S. R. Saleska, K. T. Wiedemann, B. J. M. Bohannan¹ and J. L. Green

Responses to editor comments

Detailed responses and revisions are given below. Editor text is in black and our responses are in blue.

Responses to Editor Comments

I find the manuscript as written to be methodologically sound and a nice contribution to an understudied area of Earth science. I would prefer that the authors provide references for the following recently-added statement: Research has shown, for example, that concentrations of fungal spores in the atmosphere vary diurnally and seasonally.

We added a citation for Ingold 1971.

I would also remove the distinction between wet and dry season in this case; rainfall is abundant during both seasons but dry season storms tend to be more intermittent and energetic. This counters the previous statement about convective instability before thunderstorms as they influence atmospheric spore concentration and composition.

We agree and have removed the text discussing potential changes in Ascomycota abundance during the wet season.

Likewise, the following statement needs references and less speculation: 'This result makes sense in light of the natural histories of many of the Ascomycota, which have single-celled or filamentous vegetative growth forms that are small enough to become aerosolized, whereas many of the Basidiomycota are too large to be easily aerosolized, other than in the form of metabolically inactive spores. It is possible that if we had sampled at night rather than during the day, we would have observed a higher relative abundance of Basidiomycota in the active community. The abundance of vegetative Ascomycota fragments may peak during the day when wind speeds are high, assuming they are passively dispersed by wind and convection (as opposed to active mechanisms many fungi use to disperse spores).' Such dynamics depend on the ability of the canopy to be flushed by turbulence and is the topic of an in review paper for the ZF2 tower: Gerken, T., Chamecki, M., Fuentes J.D.: Air parcel residence times and chemical processing of biogenic hydrocarbons within forest canopies. In review for: Agricultural and Forest Meteorology

We have edited the second paragraph of results section 3.2 to reduce speculative language. As suggested, we added text to acknowledge that temporal dynamics of fungi in the atmosphere above the canopy likely depend on the residence times of air parcels within and below the canopy as well as the ability of fungi to disperse through the canopy.

1 **Characterization of active and total fungal communities in the**
2 **atmosphere over the Amazon rainforest**

3

4 **A. M. Womack¹, P. E. Artaxo², F. Y. Ishida^{3,4}, R. C. Mueller^{1,5}, S. R. Saleska⁶, K. T.**
5 **Wiedemann^{2,7}, B. J. M. Bohannan¹, and J. L. Green^{1,8}**

6

7 [1]{Institute of Ecology and Evolution, University of Oregon, Eugene, OR USA}

8 [2]{Institute of Physics, University of São Paulo, São Paulo, Brazil}

9 [3]{Instituto Nacional de Pesquisas da Amazonia, Manaus, Brazil}

10 [4]{School of Marine and Tropical Biology, James Cook University, Cairns, Qld, Australia}

11 [5]{Los Alamos National Laboratory, Los Alamos, NM, USA}

12 [6]{Department of Ecology and Evolutionary Biology, University of Arizona, Tucson, AZ, USA}

13 [7]{School of Engineering and Applied Sciences, Harvard University, Cambridge, MA, USA}

14 [8]{Santa Fe Institute, Santa Fe, NM, USA}

15 Correspondence to: Ann M. Womack (womack.ann@gmail.com)

16

17 **Abstract**

18 Fungi are ubiquitous in the atmosphere and may play an important role in atmospheric processes.
19 We investigated the composition and diversity of fungal communities over the Amazon rainforest
20 canopy and compared these communities to fungal communities found in terrestrial environments.
21 We characterized the total fungal community and the metabolically active portion of the
22 community using high-throughout DNA and RNA sequencing and compared these data to
23 predictions generated by a mass-balance model. We found that the total community was primarily
24 comprised of fungi from the phylum Basidiomycota. In contrast, the active community was
25 primarily composed of members of the phylum Ascomycota and included a high relative abundance

1 of lichen fungi, which were not detected in the total community. The relative abundance of
2 Basidiomycota and Ascomycota in the total and active communities was consistent with our model
3 predictions, suggesting that this result was driven by the relative size and number of spores
4 produced by these groups. When compared to other environments, fungal communities in the
5 atmosphere were most similar to communities found in tropical soils and leaf surfaces. Our results
6 demonstrate that there are significant differences in the composition of the total and active fungal
7 communities in the atmosphere, and that lichen fungi, which have been shown to be efficient ice
8 nucleators, may be abundant members of active atmospheric fungal communities over the forest
9 canopy.

10

11 **1 Introduction**

12 Fungi are critical to the functioning of terrestrial ecosystems and may also play an important role
13 in the functioning of the atmosphere. Fungi are abundant and ubiquitous in the atmosphere, with
14 an estimated global land surface emission rate of 50 Tg/year for fungal spores alone (Elbert et al,
15 2007). Fungal bioaerosols are not only abundant but also affect physical and chemical processes in
16 the atmosphere. Fungal spores, cellular fragments, and cell-free biological particles have the
17 potential to affect precipitation by acting as ice and cloud condensation nuclei (Després et al., 2012;
18 Morris et al., 2013; Pouleur et al., 1992; Richard et al., 1996), and metabolically active fungi
19 sampled from the atmosphere are capable of transforming compounds known to play a major role
20 in atmospheric chemistry, including carboxylic acids (Ariya, 2002; Côté et al., 2008; Vaïtilingom
21 et al., 2013), formaldehyde, and hydrogen peroxide (Vaïtilingom et al., 2013).

22 The *in situ* function of airborne fungi will depend on the physiological state of fungal cells.

23 Metabolically active vegetative cells have the potential to transform atmospheric compounds and
24 ultimately alter atmospheric chemistry, whereas for dormant spores this metabolic capability is
25 greatly reduced (Sussman and Douthit, 1973). The ice nucleation efficiency of fungal cells also
26 likely depends on their physiological state; vegetative cells derived from potentially active fungi
27 are more efficient ice nucleators than spores. Vegetative forms of *Fusarium* (a filamentous fungi)
28 as well as several lichen fungi have been shown to nucleate ice at temperatures as warm as -1°C
29 (Després et al., 2012) (Supplement figure 1), and ice nucleation by hyphae has been observed at -

1 2.5°C. In contrast, dormant spores – particularly those with surface hydrophobins – are generally
2 poor ice nucleators. For example, ice nucleation of rust (*Puccinia*) spores requires temperatures
3 lower than -10°C (Morris et al., 2013), and *Cladsporium* spores nucleate ice at temperatures of
4 approximately -28.4°C (Iannone et al., 2011).

5 Despite its importance, we know relatively little about the physiological state of fungal cells in the
6 atmosphere. Specifically, we know little about the taxonomic composition of metabolically active
7 airborne fungi and how this compares to the composition of the total fungal community. One way
8 to survey the total and active communities is to measure community composition from rDNA (i.e.
9 rRNA genes) and rRNA in ribosomes. Sequencing rDNA provides information about the total
10 community, which includes both active and dormant individuals, whereas rRNA sequences provide
11 information about the potentially active community, because ribosomes are more abundant in
12 active cells than dormant cells (Prosser, 2002). This approach has been applied to study active
13 fungal communities in soils and on decaying plant material (Baldrian et al., 2012; Barnard et al.,
14 2013, 2014; Rajala et al., 2011) but has not been applied to fungal communities in the atmosphere.

15 Information about the taxonomic composition of airborne fungi that are present in different
16 physiological states can be used to advance atmospheric science. For example, such data can be
17 used to improve estimates of the ice nucleating capacity of fungal bioaerosols. Historically, the
18 composition of fungal communities in the atmosphere has been measured using culture-based
19 approaches such as the abundance of colony forming units of specific taxa. This has led some
20 scientists to conclude that fungal communities in the atmosphere have a low capacity for ice
21 nucleation because taxa that appear abundant using plate counts have a low ice nucleation
22 efficiency (Iannone et al., 2011, but see Pummer et al., 2013). This data may be misleading, as the
23 vast majority of fungi require identification using culture-independent approaches (Borneman and
24 Hartin, 2000). Today, culture-independent identification of active fungal taxa sampled from the
25 atmosphere can be used to direct selective culturing of potentially important fungi in the laboratory,
26 where their ice nucleation efficiencies and their metabolic capabilities can be further tested.

27 In this study, we used culture-independent approaches to measure the composition of total and
28 active atmospheric fungal communities *in situ* and a mass-balance model to aid in the interpretation
29 of our results. Our study system is the atmosphere above the Amazon rainforest canopy. We chose
30 this system because fungal bioaerosols make up a substantial proportion of aerosol particulate

1 matter over the Amazon (Elbert et al., 2007; Heald and Spracklen, 2009) and are estimated to be a
2 dominant force responsible for cloud formation over the Amazon (Pöschl et al., 2010). We used a
3 combined approach of DNA and RNA sequencing to address the following questions: 1) What is
4 the composition of total airborne fungal communities? 2) What is the composition of active
5 airborne fungal communities? 3) What likely drives differences in the composition of the total and
6 active airborne fungal communities? 4) Is the diversity and structure of fungal communities in the
7 atmosphere similar to that found in terrestrial environments?

8

9 **2 Methods**

10 **2.1 Sample collection**

11 Sampling was conducted on the ZF2 K34 flux tower (S -2.60907, W -60.20917, 67 m a.s.l.) in the
12 Reserva Biológica do Cueiras in central Amazonia, about 60 km NNW of Manaus, Brazil. The site
13 is operated by the Instituto Nacional de Pesquisas da Amazonia (INPA) under the Large Scale
14 Biosphere-Atmosphere Experiment in Amazonia (LBA) program (Martin et al., 2010). Tower
15 height is approximately 54 m. Surrounding vegetation is undisturbed, mature, terra firme rainforest,
16 with a leaf area index of 5–6 and an average canopy height of 30 m. Samples were collected at the
17 end of dry season over four days, December 8-11, 2010, from a height of 48m above the forest
18 floor. Environmental conditions during the four-day sampling period were typical for the location
19 in early December with partial clouds and temperatures ranging from approximately 28.5°C to
20 32.1°C. Heavy rain and thunderstorms occurred on 12/8 and 12/11. Aerosol samples were collected
21 using SKC Biosamplers (BioSampler SKC Inc.). Samplers were filled with 20 mL of a water-based
22 preservation solution (LifeGuard Soil Preservation Solution, MO BIO Laboratories, Inc) to prevent
23 DNase and RNase activity and maintain cells in stasis to allow accurate community profiling of the
24 total and active fungal community. Twelve impingers were operated at 12.5 L/min from
25 approximately 9:00 am – 4:00 pm each day. At the end of each day, the sampling liquid from all
26 impingers was pooled and stored at -20°C. Impingers were cleaned each day by rinsing in 70%
27 ethanol followed by sterilization using a portable pressure cooker.

28 **2.2 Nucleic acid isolation and cDNA synthesis**

1 Samples were transported on ice to the University of Oregon where the liquid sample from each
2 day was separated into two aliquots, one to be used for DNA extraction and the other for RNA
3 extraction. The divided samples were filtered through sterile, individually wrapped, 0.22 µm
4 cellulose nitrate filters (Nalgene Analytical Test Filter Funnels, Thermo Fisher Scientific). DNA
5 was extracted from filters using the MO BIO PowerWater DNA Isolation Kit according to the
6 manufacturer's instructions with a 100 µl elution volume. RNA was extracted from filters using
7 the MO BIO PowerWater RNA Isolation Kit with the following modifications. The DNase steps
8 included in the kit were omitted. RNA was eluted in 50 µl. The extracted RNA was treated with
9 DNase I (RNase-free) (Fermentas International, Inc) according to the manufacturer's instructions.
10 DNase reactions were cleaned (Zymo Research Clean and Concentrate-5) and eluted into 50 µl.
11 cDNA was synthesized from the total RNA extract using the SuperScript II First-Strand Synthesis
12 System (Invitrogen, Life Technologies Corporation) with random hexamers. All RNA was
13 converted into cDNA in six synthesis reactions and one reverse transcriptase negative control
14 reaction. Three field blanks were generated by filtering unused LifeGuard Solution through new,
15 sterile filters. Blanks were processed in parallel to the RNA and DNA samples including extraction,
16 PCR amplification, and library preparation. Following library preparation, blank samples were
17 visualized on an agarose gel and no visible bands were observed.

18 **2.3 Library preparation and sequencing**

19 To increase the concentration of cDNA to levels required for sequencing, we used multiple
20 displacement amplification (GenomiPhi V2, GE Healthcare) according to the protocol described in
21 Gilbert *et al.* (2010) including second-stand synthesis, amplification, and de-branching of
22 amplification products. The fully de-branched products were sheared by sonication (24 cycles, 30
23 seconds each) using the Bioruptor sonication system (Diagenode). cDNA fragments were end-
24 repaired (End-It DNA End-Repair Kit, Epicentre Biotechnologies), cleaned and concentrated
25 (Zymo Research Clean and Concentrate-5) and eluted in 40 µl. A-overhangs were added to the
26 end-repaired fragments using Klenow exo(-) (Epicentre Biotechnologies) in a 50 µl reaction.
27 Reaction products were cleaned and concentrated (Zymo Research Clean and Concentrate-5).
28 Standard paired-end, barcoded Illumina adaptors (Supplement table 1) were ligated to the
29 fragments using T4 ligase (Fermentas). Reaction products were cleaned and concentrated (Zymo

1 Research Clean and Concentrate-5) and eluted in 12 μ l. To enrich fragments with ligated adaptors,
2 PCR amplification was performed using primers containing the flowcell adaptor and
3 complementary to the Illumina sequencing primer (Supplement Table 1). PCR reactions were
4 performed using Phusion DNA polymerase (New England Biolabs) with 12 μ l template, 10 μ l 5x
5 HF buffer, 1 μ l 10 mM dNTPs, 2 μ l 10 mM primer mix, 0,5 μ l enzyme and 25.5 μ l water for a final
6 reaction volume of 50 μ l. PCR cycling conditions were as follows: 30 seconds denaturation at 98°C
7 followed by 18 cycles of 98°C for 30 seconds, 65°C for 30 seconds and 72°C for 30 seconds
8 following by a final extension at 72°C for 5 minutes. PCR products were size fractionated by gel
9 electrophoresis (2.5%, low-melt agarose). Products in the range of 150-500 bp were excised, and
10 DNA from the excised gel pieces was extracted (QiagenMinElute Gel Extraction) and eluted into
11 20 μ l. DNA was quantitated using a Qubit 2.0 Fluorometer (Invitrogen, Life Technologies
12 Corporation) and combined in equal molar concentrations. Shotgun metatranscriptome libraries
13 were sequenced (150 base pairs, paired-end) on the Illumina HiSeq 2000 (Illumina, Inc.) platform
14 at the University of Oregon Genomics Core Facility. LSU rDNA amplicons were sequenced (250
15 base pairs, paired-end) on the Illumina MiSeq platform at the Dana-Farber Cancer Institute
16 Molecular Biology Core.

17 The D1-D2 region of the large subunit (LSU) rRNA gene was targeted using PCR with the primers
18 LR0R (5'-ACCCGCTGAACTTAAGC-3') and LR3 (5'-CCGTGTTTCAAGACGGG-3')
19 (<http://sites.biology.duke.edu/fungi/mycolab/primers.htm>). LSU amplicon libraries were prepared
20 using a two-stage PCR procedure as described in (Kembel and Mueller, 2014) using unique
21 combinatorial barcodes (Gloor et al., 2010) to identify samples (Supplement table 2).

22 **2.4 Sequence pre-processing**

23 **2.4.1 Metatranscriptome**

24 Overlapping paired end reads were aligned and joined using fastq-join (<https://code.google.com/p/ea-utils/wiki/FastqJoin>). Joined reads and non-overlapping single-end reads were trimmed and
25 filtered using PrinSeq (Schmieder and Edwards, 2011). Sequences <75 bp, > 2% Ns, and/or mean
26 quality score <20 were removed. Sequence artifacts defined as exact duplicates with >5,000
27 sequences were removed. Sequences in the Dec. 10 sample were primarily artifacts, so this
28

1 metatranscriptome sample was excluded from further analysis. Putative rRNAs in the remaining
2 sequences were identified using SortMeRNA (Kopylova et al., 2012) with the non-redundant
3 version of the following databases: rfam 5.8S (version 11.0) (Burge et al., 2013); Unite (November
4 2011 version) (Kõljalg et al., 2013), and Silva 18S and Silva 28S (Release 115) (Quast et al., 2013).
5 Of 5,165,185 quality-filtered reads, 1,915,994 with an average length of 137.5 bp were identified
6 as putative rRNAs (Supplement table 3).

7 **2.4.2 LSU amplicons**

8 Forward and reverse barcodes were combined to make a 12 bp barcode on the forward read. Only
9 forward reads derived from the LR3 region were used for analysis. This region has been shown to
10 have high species-level resolution even with short read lengths (Liu et al., 2012).

11 **2.4.3 Multi-environment sequences**

12 LSU sequences from four soil studies (Barnard et al., 2013; Kerekes et al., 2013; Penton et al.,
13 2013, 2014) and one phyllosphere study (Kembel and Mueller, 2014) were compared to air samples
14 collected for this study (Supplement table 4). Raw sequence data and associated metadata were
15 downloaded from publically available databases. 12 bp barcodes were added to all sequences to
16 identify each sample in downstream analysis.

17 **2.5 LSU amplicon and metatranscriptome sequence processing**

18 All sequences were processed in QIIME version 1.7 (Caporaso et al., 2010). Briefly, libraries were
19 individually demultiplexed and filtered for quality. Sequences with an average quality score less
20 than 20, shorter than 150 bp and with greater than 2 primer mismatches were discarded. The same
21 parameters were used across all samples except the metatranscriptome rRNAs were a size cut off
22 of greater than 75 bp was used. In order to decrease computation time, sequences from Kembel and
23 Mueller (2014) and Penton *et al.* (2014) were randomly subsampled to 25% and 60% of the total
24 number of sequences, respectively. Sequences were clustered into operational taxonomic units
25 (OTUs) at 97% sequence similarity using closed reference BLAST (Altschul et al., 1990) against
26 the Ribosomal Database Project Fungal LSU training set 1 (Cole et al., 2014). Taxonomy was
27 assigned to each OTU was that of the most similar representative in the RDP database.

1 Following sequence processing and quality filtering, a total of 55,414 amplicon and 1,915,994
2 metatranscriptome LSU sequences generated for this study and 1,577,458 LSU sequences from soil
3 and phyllosphere studies were retained (Supplement table 3). For analyses using only samples from
4 this study, the data were rarefied to 5,300 sequences per sample. For analyses that compare samples
5 in this study to samples from other studies, the data were rarefied to 500 sequences per sample.

6 **2.6 Statistical analyses and data availability**

7 All statistical analyses were conducted in R (R Core Team, 2014) primarily using the `vegan`
8 (Oksanen et al., 2013) package for ecological statistics and the `ggplot2` (Wickham, 2009)
9 package for visualizations.

10 Sequence files and metadata have been deposited in Figshare
11 (<http://dx.doi.org/10.6084/m9.figshare.1335851>). Data from other studies used for cross
12 environment analyses are available using the databases and identifiers referenced in the respective
13 manuscripts.

14 **2.7 Mass-balance model**

15 We use a global, well-mixed, one-box material-balance model to predict the relative abundances
16 of fungal cells measured as gene copies sampled in the active and total portions of atmospheric
17 bioaerosols. Model description and details are available in Appendix A.

18

19 **3 Results & Discussion**

20 **3.1 Basidiomycota dominate total airborne fungal communities**

21 Measurements of airborne fungi using culture-based methods such as quantifying spore and colony-
22 forming unit counts have been conducted for centuries (Després et al., 2012). In comparison, there
23 have been few culture-independent studies of the fungal composition of atmospheric samples (e.g.
24 Boreson et al., 2004; Bowers et al., 2013; Fierer et al., 2008; Fröhlich-Nowoisky et al., 2009, 2012;
25 Pashley et al., 2012; Yamamoto et al., 2012). Using a culture-independent approach, we found the
26 composition of total airborne fungal communities primarily included taxa belonging to the phyla

1 Ascomycota and Basidiomycota (Figure 1). This result is similar to what is observed in
2 environments on the Earth's surface (James et al., 2006) and what has been reported in other studies
3 of fungi in the atmosphere (Bowers et al., 2013; Fröhlich-Nowoisky et al., 2009, 2012; Yamamoto
4 et al., 2012).

5 Basidiomycota dominated the total airborne community in our air samples (mean relative
6 abundance = $90.2 \pm 6.9\%$) (Figure 1). Within the phylum Basidiomycota, Agaricomycetes were the
7 most abundant class in our samples. Agaricomycetes have been previously detected in air samples
8 (Fröhlich-Nowoisky et al., 2012; Woo et al., 2013; Yamamoto et al., 2012) and are common in
9 tropical soils (Tedersoo et al., 2014) and leaf surfaces (Kembel and Mueller, 2014). Within the
10 Agaricomycetes, the most abundant order was the Polyporales (mean = $55.7 \pm 2.3\%$). Polyporales
11 have been detected in culture-independent studies of urban aerosols (Yamamoto et al., 2012) and
12 culturable representatives have been isolated from cloud water (Amato et al., 2007). At the genus
13 level, there were several taxa detect in the total community with ice nucleation activity including
14 *Acremonium*, *Cladosporium*, *Fusarium*, and *Rhizopus* (Table S4).

15 The presence of Agaricomycetes may have implications for atmospheric processes. Ice nucleation
16 efficiency within the Agaricomycetes is variable, with some taxa capable of nucleating ice at
17 temperatures as warm as -17°C (Haga et al., 2014) (Supplement figure 1). These temperatures are
18 warmer than what has been measured for *Penicillium* spores (Iannone et al., 2011) although not as
19 warm as what has been measured for other biological particles including other spore types (Morris
20 et al., 2013), suspensions of *Fusarium* cultures (containing spores and hyphae) (Pouleur et al.,
21 1992), and lichen fungi (Després et al., 2012). Despite the low ice nucleation efficiency of some
22 taxa in this group, given the high abundance of Agaricomycetes over the forest canopy, this group
23 could still have a significant impact on cloud formation and precipitation in the tropics.

24 The patterns we report reflect a snapshot in space and time. As in other environmental systems, the
25 composition of total fungal communities in the atmosphere will vary across different spatial and
26 temporal scales. Research has shown, for example, that concentrations of fungal spores in the
27 atmosphere vary diurnally and seasonally (Ingold, 1971). This variation is driven by complex
28 interactions between fungal dispersal mechanisms and environmental conditions, particularly
29 moisture and wind speed. (Lacey, 1996). Our samples were collected during the day, and spores
30 released by mechanical disturbances often peak in abundance in the air during midday when wind

Deleted: .

1 speeds are highest (Lacey, 1996). Taxa that require dry conditions for dispersal also tend to release
2 spores during the day, and taxa that require high relative humidity, including many Basidiomycota,
3 tend to release spores at night when humidity is highest (Elbert et al., 2007; Lacey, 1996). In
4 addition to humidity, precipitation events can also affect the dispersal of fungi. Overall
5 concentrations of spores have been shown to increase in the atmosphere due to convective
6 instability preceding thunderstorms (Burch and Levetin, 2002), and Ascomycota concentrations
7 increase during and immediately after rainstorms (Elbert et al., 2007).

Deleted: Our samples were collected at the end of the dry season. If we had sampled during the wet season, it is possible we would have observed a higher relative abundance of Ascomycota in the total community since the dispersal of Ascospores has been shown to increase before and after rain storms.

8 **3.2 Ascomycota dominate active airborne fungal communities**

9 The composition of total and active fungal communities over the Amazon rainforest canopy
10 significantly differed (ADONIS, $R^2 = 0.342$, $p = 0.029$). The active community in the atmosphere
11 over the forest canopy was dominated by Ascomycota (mean relative abundance = $80.4 \pm 20\%$)
12 (Figure 1). Basidiomycota comprised a smaller fraction of the sampled genes (mean = $7.3 \pm 6.8\%$)
13 with the remainder of identified sequences belonging to the phyla Chytridiomycota and
14 Glomeromycota. This result makes sense in light of the natural histories of many of the
15 Ascomycota, which have single-celled or filamentous vegetative growth forms that are small
16 enough to become aerosolized, whereas many of the Basidiomycota are too large to be easily
17 aerosolized (Moore et al., 2011), other than in the form of metabolically inactive spores.

Deleted: ,

18 Similar to the total community, we expect the compositions of active fungal communities in the
19 atmosphere likely vary over different time scales. For example, had we sampled at night rather than
20 during the day, we may have observed a higher relative abundance of Basidiomycota in the active
21 community. This could be due to an increase in the concentration of basidiospores combined with
22 a decrease in vegetative Ascomycota at night because basidiospores are abundant in the Amazon
23 atmosphere at night (Elbert et al., 2007) and the detection of ribosomes within spores could lead to
24 an increase in the observed relative abundance of Basidiomycota in the active community. We
25 would also expect the relative abundance of Ascomycota to decrease at night when wind speeds
26 typically decrease, particularly considering that many vegetative Ascomycota fragments are
27 passively dispersed by wind and convection (as opposed to active mechanisms many fungi use to
28 disperse spores). However, these patterns will depend on the relative abilities of spores (Gilbert &
29 Reynolds, 2005) and fragments to disperse beyond the understory as well as the residence times of

Deleted: It is possible that if

Deleted: had

Deleted: ,

Deleted: would

Deleted: The

Deleted: may peak during the day when wind speeds are high, assuming they

1 [air parcels below and within the canopy. It will be fruitful for future studies to dynamically sample](#)
2 [both above and below the canopy to elucidate the mechanisms driving temporal variation in fungal](#)
3 [communities in the atmosphere.](#)

4 The most abundant classes of Ascomycota detected were Sordariomycetes (mean = 27.1±6.6%),
5 and Lecanoromycetes (mean = 17.5±7.6%). Sordariomycetes have been previously detected in
6 culture-independent air samples (Fröhlich-Nowoisky et al., 2009, 2012; Yamamoto et al., 2012)
7 and have been shown to be abundant on tropical tree leaves (Kembel and Mueller, 2014) and
8 tropical soils (Peay et al., 2013). In most ecosystems, Sordariomycetes are endophytes, pathogens,
9 and saprotrophs (Zhang et al., 2007). Xylariales, which includes both endophytes and plant
10 pathogens (Zhang et al., 2007), was the most abundant order within the Sordariomycetes in our
11 samples. Several genera with known ice nucleation capability were detected in the active
12 community including *Agaricus*, *Amanita*, *Aspergillus*, *Boletus*, *Lepsita*, *Mortierella*, *Puccinia*,
13 *Rhizopus*, and the lichen fungus *Cladonia* (Table S4). Below we focus our discussion on the class
14 Lecanoromycetes, an understudied but potentially important group of fungi in the atmosphere.

15 Lecanoromycetes were the second most abundant class of Ascomycota detected over the rainforest
16 canopy. This group has been detected in other culture-independent studies of fungi in the
17 atmosphere (Fröhlich-Nowoisky et al., 2012; Yamamoto et al., 2012). The Lecanoromycetes
18 contain 90% of the lichen-associated fungi (Miadlikowska et al., 2007). Lichens are a symbiosis
19 between a fungus and a photosynthetic partner such as eukaryotic algae or cyanobacteria. Lichens
20 are known to be hardy and may be particularly well-adapted for long distance transport and
21 metabolic activity in the atmosphere. Lichens are often the dominant life forms in environments
22 that have conditions similar to those found in the atmosphere, including low water (Kranter et al.,
23 2008) and nutrient availability, wide temperature variations, and high UV irradiance (e.g. Solhaug,
24 Gauslaa, Nybakken, & Bilger, 2003) (Onofri et al., 2004).

25 Another notable trait of lichens is their efficient ice nucleation capacity. Although there have been
26 no investigations specifically on the most abundant lichen species detected in this study, *Physcia*
27 *stellaris* (mean = 8.3±3.8%) and *Rinodina milvina* (mean = 4.8±3.4%), there have been multiple
28 studies of the ice nucleation efficiency of many other lichen fungi species. Ice nucleation activity
29 of lichens has been measured at temperatures warmer than -8°C, including 13 of 15 taxa tested by
30 Henderson-Begg and colleagues (Henderson-Begg et al., 2009) and 9 of 15 taxa tested by Kieft

Deleted: And, Basidiospores have been shown to be particularly abundant in the Amazon atmosphere at night (Elbert et al., 2007). As with the total community, we expect that the composition of active fungal communities in the atmosphere will vary across different spatial and temporal scales.

1 (Kieft, 1988). These studies have demonstrated that lichens are among the most efficient biological
2 ice nucleators. Therefore, their presence in the atmosphere may have a significant impact on cloud
3 formation and precipitation. This ice nucleation capacity may also enable lichens to control the
4 extent of their dispersal through the atmosphere. It is possible that lichens achieve this by nucleating
5 ice formation, which leads to precipitation and ultimately deposition. This phenomenon has been
6 shown to occur in some phytopathogenic bacteria (Morris et al., 2008, 2010) and may occur in
7 fungi as well (Morris et al., 2013).

8 **3.3 Dominance of Basidiomycota in total communities and Ascomycota in active** 9 **communities is consistent with mass-balance predictions**

10 Our mass balance model (Appendix A) predicted Basidiomycota would dominate the total
11 community because they produce orders of magnitude more spores and have smaller aerodynamic
12 diameters compared to Ascomycota. Consistent with this prediction, the total airborne community
13 was dominated by Basidiomycota in our air samples (mean relative abundance = $90.2 \pm 6.9\%$)
14 (Figure 1). There have been some empirical studies reporting the opposite pattern, with a higher
15 relative abundance of Ascomycota compared to Basidiomycota (Bowers et al., 2013; Fierer et al.,
16 2008; Pashley et al., 2012). There has been one study focused on airborne fungal communities in
17 the Amazon Basin (Fröhlich-Nowoisky et al., 2012). Although the site of this study was the
18 atmosphere above a rural pasture (versus a tropical rainforest, as in our study) these investigators
19 also found that Basidiomycota dominate airborne fungal communities

20 Our mass-balance model explains the differences in composition between the total and active air
21 communities. However, some of the differences we observed may be partially attributable to the
22 use of different approaches in characterizing the total and active communities. In this study, the
23 total community was characterized by PCR-based amplification and sequencing of LSU genes,
24 whereas the active community was characterized through random sequencing of all the RNA
25 present in the samples. Shotgun metatranscriptome sequencing and PCR-based community
26 characterization approaches each have their own biases (Hong et al., 2009; Morgan et al., 2010).
27 Our data suggest that the selection of LSU primers led to biased results. For example, the high
28 relative abundance of lichen fungi (class Lecanoromycetes) in the active community was
29 unexpected because this group was not detected in the total community and has only been detected

1 in low abundance in other PCR-based studies of fungi in the atmosphere (Fröhlich-Nowoisky et
2 al., 2012). We tested the primer pair used in this study (LR0R-LR3) using the SILVA TestPrime
3 tool (Klindworth et al., 2013) and found coverage of the Lecanoromycetes with this primer pair
4 was 71.4%. Within the class Lecanoromycetes, the order Teloschistales, which contains the most
5 abundant species detected in the active community, would not have been detected with this primer
6 pair. However, the general pattern that Ascomycota were much less abundant than Basidiomycota
7 in the total community is not likely due to primer bias as overage of the phylum Ascomycota by
8 the LR0R-LR3 primer pair is 85.5% according to TestPrime. Our findings underscore the value of
9 using a combination of PCR-based and shotgun-based sequencing approaches, particularly in
10 environments that are understudied and where little is known about microbiome structure and
11 function.

12 **3.4 Fungal air communities above the forest canopy are most similar in** 13 **composition to tropical phyllosphere and soil communities**

14 We compared total and active fungal air communities to communities from tropical, temperate, and
15 tundra soils and from the surfaces of tropical tree leaves. Community composition significantly
16 differed across environment types (ADONIS, $R^2 = 0.167$, $p = 0.001$), and fungal communities in
17 the atmosphere were compositionally distinct from communities in other environments (Figure 2).
18 Ascomycota was the most abundant phylum across all soil and phyllosphere samples (soil mean
19 relative abundance = $78.4 \pm 14.9\%$, phyllosphere = $90.9 \pm 4.9\%$) followed by Basidiomycota (soil
20 mean relative abundance = $19.0 \pm 14.9\%$, phyllosphere = $7.4 \pm 4.5\%$) (Figure 3). We expected
21 communities to be distinct across habitat types because environmental conditions may differ across
22 the habitat types and select for different communities. However, in the atmosphere, dispersal and
23 mixing of fungi from multiple habitat types may be driving the observed community composition
24 differences instead of environmental selection.

25 The diversity of fungal communities in the atmosphere is within the range of diversities reported
26 for terrestrial environments, including those of tropical leaf surfaces, tropical soils, temperate
27 grassland soils, and tundra soils. Overall taxonomic richness, defined as the number of OTUs,
28 significantly varied among environment types (ANOVA, $F(5,237) = 66.89$, $p < 0.001$) (Supplement
29 figure 2). Tukey's HSD post-hoc comparisons indicated that the richness of air communities, both

1 total and active, was greater than tundra soil communities and did not significantly differ from
2 temperate grassland soil communities. In general, air communities were less diverse than tropical
3 forest phyllosphere and soil communities with the exception of tropical forest soils and active air
4 communities, which did not significantly differ. Similar patterns have been observed in soil
5 communities where taxonomic richness in arctic soils was significantly lower than soils from
6 temperate and tropical ecosystems (Fierer et al., 2012).

7 Total air communities were most similar to tropical phyllosphere communities (mean Sørensen
8 similarity = 0.015 ± 0.009 ; Tukey's HSD, $p < 0.001$) and active air communities were most similar
9 to tropical soil communities (mean Sørensen similarity = 0.010 ± 0.007 , Tukey's HSD, $p < 0.001$)
10 (Supplement figure 3). This suggestion makes sense since fungal spores and hyphae are relatively
11 large aerosol particles with short residence times in the atmosphere, limiting opportunities for long-
12 distance dispersal. While these results are suggestive, detailed information is lacking regarding the
13 potential influence of terrestrial source environments and their role in structuring airborne fungal
14 communities.

15

16 **4 Conclusion**

17 Fungi in the atmosphere play an important role in atmospheric processes including precipitation
18 development through ice nucleation. This is of particular significance in the atmosphere over the
19 Amazon rainforest canopy where fungi constitute a large fraction of the total aerosol content (Elbert
20 et al., 2007; Heald and Spracklen, 2009) and precipitation is aerosol-limited (Pöschl et al., 2010).
21 Our study represents the first culture-independent survey of fungal communities over the Amazon
22 rainforest canopy. It is also the first to measure metabolically active microbial communities in the
23 atmosphere using an RNA-based approach. Using this RNA-based approach, we found evidence
24 for the presence of potentially active fungi in the atmosphere, including lichen fungi (class
25 Lecanonomycetes) and the following genera: Agaricus; Amanita; Aspergillus; Boletus; Cladonia;
26 Lepsita; Mortierella; Puccinia; and Rhizopus. While an understanding of the structure of fungal
27 communities in the atmosphere is beginning to emerge, studies on the function of these
28 communities have lagged behind. We suggest that future research focus on understanding the
29 functional capacity of airborne microbes with traits particularly well-suited for survival and

1 metabolic activity in extreme environments. As with any environment, understanding both the
2 structure and function of microbial communities in the atmosphere is needed to assess their
3 potential impact on ecosystem processes such as water and carbon cycling. This study opens the
4 door for future investigations of the diversity and function of fungal communities in the
5 atmosphere.

6

7 **Author contributions**

8 A. M. Womack conceived and designed the experiments, analyzed the data, wrote the paper,
9 prepared figures and/or tables, and reviewed drafts of the paper. P. E. Artaxo conceived and
10 designed the experiments. F. Y. Ishida collected the samples and reviewed drafts of the paper. R.
11 C. Mueller conceived and designed the experiments, reviewed drafts of the paper and contributed
12 reagents/materials/analysis tools. S. R. Saleska conceived and designed the experiments. K. T.
13 Wiedemann collected the samples. B. J. M. Bohannan conceived and designed the experiments,
14 collected the samples, and reviewed drafts of the paper. J. L. Green conceived and designed the
15 experiments, wrote the paper, reviewed drafts of the paper, and contributed
16 reagents/materials/analysis tools.

17

18 **Acknowledgements**

19 This research was funded by the University of Oregon and the Alfred P. Sloan Foundation. We
20 thank Jonas Frankel-Bricker for his work in preparing the LSU libraries for sequencing. We also
21 thank members of the Green and Bohannan labs for their constructive feedback during the
22 preparation of this manuscript.

23

24 **References**

- 25 Altschul, S. F., Gish, W., Miller, W., Myers, E. W. and Lipman, D. J.: Basic local alignment search
26 tool., *J. Mol. Biol.*, 215(3), 403–10, doi:10.1016/S0022-2836(05)80360-2, 1990.
- 27 Amato, P., Parazols, M., Sancelme, M., Laj, P., Mailhot, G. and Delort, A.-M.: Microorganisms
28 isolated from the water phase of tropospheric clouds at the Puy de Dôme: major groups and growth

- 1 abilities at low temperatures., *FEMS Microbiol. Ecol.*, 59(2), 242–54, doi:10.1111/j.1574-
2 6941.2006.00199.x, 2007.
- 3 Ariya, P. A.: Microbiological degradation of atmospheric organic compounds, *Geophys. Res. Lett.*,
4 29(22), 2–5, doi:10.1029/2002GL015637, 2002.
- 5 Baldrian, P., Kolařík, M., Stursová, M., Kopecký, J., Valášková, V., Větrovský, T., Zifčáková, L.,
6 Snajdr, J., Řídl, J., Vlček, C. and Voříšková, J.: Active and total microbial communities in forest
7 soil are largely different and highly stratified during decomposition., *ISME J.*, 6(2), 248–58,
8 doi:10.1038/ismej.2011.95, 2012.
- 9 Barnard, R. L., Osborne, C. a and Firestone, M. K.: Responses of soil bacterial and fungal
10 communities to extreme desiccation and rewetting., *ISME J.*, 7(11), 2229–41,
11 doi:10.1038/ismej.2013.104, 2013.
- 12 Barnard, R. L., Osborne, C. A. and Firestone, M. K.: Changing precipitation pattern alters soil
13 microbial community response to wet-up under a Mediterranean-type climate, *ISME J.*,
14 doi:10.1038/ismej.2014.192, 2014.
- 15 Binder, M., Justo, A., Riley, R., Salamov, A., Lopez-Giraldez, F., Sjökvist, E., Copeland, A.,
16 Foster, B., Sun, H., Larsson, E., Larsson, K.-H., Townsend, J., Grigoriev, I. V and Hibbett, D. S.:
17 Phylogenetic and phylogenomic overview of the Polyporales., *Mycologia*, 105(6), 1350–73,
18 doi:10.3852/13-003, 2013.
- 19 Boreson, J., Dillner, A. and Peccia, J.: Correlating bioaerosol load with PM2.5 and PM10cf
20 concentrations: a comparison between natural desert and urban-fringe aerosols, *Atmos. Environ.*,
21 38(35), 6029–6041, doi:10.1016/j.atmosenv.2004.06.040, 2004.
- 22 Borneman, J. and Hartin, R. J.: PCR Primers That Amplify Fungal rRNA Genes from
23 Environmental Samples, *Appl. Environ. Microbiol.*, 66(10), 4356–4360,
24 doi:10.1128/AEM.66.10.4356-4360.2000, 2000.
- 25 Bowers, R. M., Clements, N., Emerson, J. B., Wiedinmyer, C., Hannigan, M. P. and Fierer, N.:
26 Seasonal variability in bacterial and fungal diversity of the near-surface atmosphere., *Environ. Sci.*
27 *Technol.*, 47(21), 12097–106, doi:10.1021/es402970s, 2013.
- 28 Burge, S. W., Daub, J., Eberhardt, R., Tate, J., Barquist, L., Nawrocki, E. P., Eddy, S. R., Gardner,
29 P. P. and Bateman, A.: Rfam 11.0: 10 years of RNA families., *Nucleic Acids Res.*, 41(Database
30 issue), D226–32, doi:10.1093/nar/gks1005, 2013.
- 31 Burch, M. and Levetin, E.: Effects of meteorological conditions on spore plumes., *Int. J.*
32 *Biometeorol.*, 46(3), 107–17, doi:10.1007/s00484-002-0127-1, 2002.
- 33 Caporaso, J. G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F. D., Costello, E. K., Fierer,
34 N., Peña, A. G., Goodrich, J. K., Gordon, J. I., Huttley, G. A., Kelley, S. T., Knights, D., Koenig,

- 1 J. E., Ley, R. E., Lozupone, C. A., McDonald, D., Muegge, B. D., Pirrung, M., Reeder, J., Sevinsky,
2 J. R., Turnbaugh, P. J., Walters, W. A., Widmann, J., Yatsunenko, T., Zaneveld, J. and Knight, R.:
3 QIIME allows analysis of high-throughput community sequencing data., *Nat. Methods*, 7, 335–
4 336, doi:10.1038/nmeth.f.303, 2010.
- 5 Cole, J. R., Wang, Q., Fish, J. A., Chai, B., McGarrell, D. M., Sun, Y., Brown, C. T., Porras-Alfaro,
6 A., Kuske, C. R. and Tiedje, J. M.: Ribosomal Database Project: data and tools for high throughput
7 rRNA analysis., *Nucleic Acids Res.*, 42(Database issue), D633–42, doi:10.1093/nar/gkt1244,
8 2014.
- 9 Côté, V., Kos, G., Mortazavi, R. and Ariya, P. A.: Microbial and “de novo” transformation of
10 dicarboxylic acids by three airborne fungi., *Sci. Total Environ.*, 390(2-3), 530–7,
11 doi:10.1016/j.scitotenv.2007.10.035, 2008.
- 12 Després, V. R., Huffman, J. A., Burrows, S. M., Hoose, C., Safatov, A. S., Buryak, G., Fröhlich-
13 Nowoisky, J., Elbert, W., Andreae, M. O., Pöschl, U. and Jaenicke, R.: Primary biological aerosol
14 particles in the atmosphere: a review, *Tellus B*, 64, doi:10.3402/tellusb.v64i0.15598, 2012.
- 15 Elbert, W., Taylor, P. E., Andreae, M. O. and Pöschl, U.: Contribution of fungi to primary biogenic
16 aerosols in the atmosphere: wet and dry discharged spores, carbohydrates, and inorganic ions,
17 *Atmos. Chem. Phys.*, 7(17), 4569–4588, doi:doi:10.5194/acp-7-4569-2007, 2007.
- 18 Fierer, N., Leff, J. W., Adams, B. J., Nielsen, U. N., Bates, S. T., Lauber, C. L., Owens, S., Gilbert,
19 J. a, Wall, D. H. and Caporaso, J. G.: Cross-biome metagenomic analyses of soil microbial
20 communities and their functional attributes., *Proc. Natl. Acad. Sci. U. S. A.*, 109(52), 21390–5,
21 doi:10.1073/pnas.1215210110, 2012.
- 22 Fierer, N., Liu, Z., Rodríguez-Hernández, M., Knight, R., Henn, M. and Hernandez, M. T.: Short-
23 term temporal variability in airborne bacterial and fungal populations., *Appl. Environ. Microbiol.*,
24 74(1), 200–7, doi:10.1128/AEM.01467-07, 2008.
- 25 Fröhlich-Nowoisky, J., Burrows, S. M. S. M., Xie, Z., Engling, G., Solomon, P. a., Fraser, M. P.,
26 Mayol-Bracero, O. L., Artaxo, P., Begerow, D., Conrad, R., Andreae, M. O., Després, V. R. and
27 Pöschl, U.: Biogeography in the air: fungal diversity over land and oceans, *Biogeosciences*, 9(3),
28 1125–1136, doi:10.5194/bg-9-1125-2012, 2012.
- 29 Fröhlich-Nowoisky, J., Pickersgill, D. A., Després, V. R. and Pöschl, U.: High diversity of fungi
30 in air particulate matter, *Proc. Natl. Acad. Sci. U. S. A.*, 106(31), 12814–9,
31 doi:10.1073/pnas.0811003106, 2009.
- 32 Fröhlich-Nowoisky, J., Hill, T. C. J., Pummer, B. G., Yordanova, P., Franc, G. D. and Pöschl, U.:
33 Ice nucleation activity in the widespread soil fungus *Mortierella alpina*, *Biogeosciences*, 12(4),
34 1057–1071,doi:10.5194/bg-12-1057-2015, 2015.

- 1 [Gilbert, G. S., & Reynolds, D. R.: Nocturnal Fungi: Airborne Spores in the Canopy and](#)
- 2 [Understory of a Tropical Rain Forest. *Biotropica*, 37\(3\), 462–464. \[http://doi.org/10.1111/j.1744-\]\(http://doi.org/10.1111/j.1744-7429.2005.00061.x\)](#)
- 3 [7429.2005.00061.x](#)

- 4 [Gilbert, J. A., Zhang, K. and Neufeld, J. D.: Handbook of Hydrocarbon and Lipid Microbiology,](#)
- 5 [in Handbook of Hydrocarbon and Lipid Microbiology, edited by K. N. Timmis, pp. 4256–4262,](#)
- 6 [Springer Berlin Heidelberg, Berlin, Heidelberg., 2010.](#)

- 7 Gloor, G. B., Hummelen, R., Macklaim, J. M., Dickson, R. J., Fernandes, A. D., MacPhee, R. and
- 8 Reid, G.: Microbiome profiling by illumina sequencing of combinatorial sequence-tagged PCR
- 9 products., *PLoS One*, 5(10), e15406, doi:10.1371/journal.pone.0015406, 2010.

- 10 Green, B., Schmechel, D. and Summerbell, R.: Aerosolized fungal fragments, in *Fundamentals of*
- 11 *mold growth in indoor environments and strategies for healthy living*, edited by O. C. Adan and R.
- 12 A. Samson, pp. 211–243, Wageningen Academic Publishers., 2011.

- 13 Haga, D. I., Burrows, S. M., Iannone, R., Wheeler, M. J., Mason, R. H., Chen, J., Polishchuk, E.
- 14 A., Pöschl, U. and Bertram, A. K.: Ice nucleation by fungal spores from the classes
- 15 Agaricomycetes, Ustilaginomycetes, and Eurotiomycetes, and the effect on the atmospheric
- 16 transport of these spores, *Atmos. Chem. Phys.*, 14(16), 8611–8630, doi:10.5194/acp-14-8611-
- 17 2014, 2014.

- 18 Heald, C. L. and Spracklen, D. V.: Atmospheric budget of primary biological aerosol particles from
- 19 fungal spores, *Geophys. Res. Lett.*, 36(9), L09806, doi:10.1029/2009GL037493, 2009.

- 20 Henderson-Begg, S. K., Hill, T., Thyrrhaug, R., Khan, M. and Moffett, B. F.: Terrestrial and
- 21 airborne non-bacterial ice nuclei, *Atmos. Sci. Lett.*, n/a–n/a, doi:10.1002/asl.241, 2009.

- 22 Hong, S., Bunge, J., Leslin, C., Jeon, S. and Epstein, S. S.: Polymerase chain reaction primers miss
- 23 half of rRNA microbial diversity., *ISME J.*, 3(12), 1365–73, doi:10.1038/ismej.2009.89, 2009.

- 24 Huffman, J. A., Prenni, A. J., DeMott, P. J., Pöhlker, C., Mason, R. H., Robinson, N. H., Fröhlich-
- 25 Nowoisky, J., Tobo, Y., Després, V. R., Garcia, E., Gochis, D. J., Harris, E., Müller-Germann, I.,
- 26 Ruzene, C., Schmer, B., Sinha, B., Day, D. A., Andreae, M. O., Jimenez, J. L., Gallagher, M.,
- 27 Kreidenweis, S. M., Bertram, A. K. and Pöschl, U.: High concentrations of biological aerosol
- 28 particles and ice nuclei during and after rain, *Atmos. Chem. Phys.*, 13(13), 6151–6164,
- 29 doi:10.5194/acp-13-6151-2013, 2013.

- 1 Iannone, R., Chernoff, D. I., Pringle, A., Martin, S. T. and Bertram, K.: The ice nucleation ability
2 of one of the most abundant types of fungal spores found in the atmosphere, *Atmos. Chem. Phys.*,
3 11(3), 1191–1201, doi:10.5194/acp-11-1191-2011, 2011.
- 4 Ingold, C. T.: (1971). *Fungal spores: Their liberation and dispersal*. Oxford: Clarendon Press.
- 5 Ingold, C. T.: Range in size and form of basidiospores and ascospores, *Mycologist*, 15(4), 165–
6 166, doi:10.1016/S0269-915X(01)80010-0, 2001.
- 7 James, T. Y., Kauff, F., Schoch, C. L., Matheny, P. B., Hofstetter, V., Cox, C. J., Celio, G.,
8 Gueidan, C., Fraker, E., Miadlikowska, J., Lumbsch, H. T., Rauhut, A., Reeb, V., Arnold, A. E.,
9 Amtoft, A., Stajich, J. E., Hosaka, K., Sung, G.-H., Johnson, D., O'Rourke, B., Crockett, M.,
10 Binder, M., Curtis, J. M., Slot, J. C., Wang, Z., Wilson, A. W., Schüssler, A., Longcore, J. E.,
11 O'Donnell, K., Mozley-Standridge, S., Porter, D., Letcher, P. M., Powell, M. J., Taylor, J. W.,
12 White, M. M., Griffith, G. W., Davies, D. R., Humber, R. A., Morton, J. B., Sugiyama, J., Rossmann,
13 A. Y., Rogers, J. D., Pfister, D. H., Hewitt, D., Hansen, K., Hambleton, S., Shoemaker, R. A.,
14 Kohlmeyer, J., Volkmann-Kohlmeyer, B., Spotts, R. A., Serdani, M., Crous, P. W., Hughes, K. W.,
15 Matsuura, K., Langer, E., Langer, G., Untereiner, W. A., Lücking, R., Büdel, B., Geiser, D. M.,
16 Aptroot, A., Diederich, P., Schmitt, I., Schultz, M., Yahr, R., Hibbett, D. S., Lutzoni, F.,
17 McLaughlin, D. J., Spatafora, J. W. and Vilgalys, R.: Reconstructing the early evolution of Fungi
18 using a six-gene phylogeny., *Nature*, 443(7113), 818–22, doi:10.1038/nature05110, 2006.
- 19 Kembel, S. W. and Mueller, R. C.: Plant traits and taxonomy drive host associations in tropical
20 phyllosphere fungal communities, *Botany*, 140210143428007, doi:10.1139/cjb-2013-0194, 2014.
- 21 Kerekes, J., Kaspari, M., Stevenson, B., Nilsson, R. H., Hartmann, M., Amend, A. and Bruns, T.
22 D.: Nutrient enrichment increased species richness of leaf litter fungal assemblages in a tropical
23 forest., *Mol. Ecol.*, 22(10), 2827–38, doi:10.1111/mec.12259, 2013.
- 24 Kieft, T. L.: Ice Nucleation Activity in Lichens, *Appl. Environ. Microbiol.*, 54(7), 1678–1681,
25 1988.
- 26 Klindworth, A., Pruesse, E., Schweer, T., Peplies, J., Quast, C., Horn, M. and Glöckner, F. O.:
27 Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-generation
28 sequencing-based diversity studies., *Nucleic Acids Res.*, 41(1), e1, doi:10.1093/nar/gks808, 2013.
- 29 Kõljalg, U., Nilsson, R. H., Abarenkov, K., Tedersoo, L., Taylor, A. F. S., Bahram, M., Bates, S.
30 T., Bruns, T. D., Bengtsson-Palme, J., Callaghan, T. M., Douglas, B., Drenkhan, T., Eberhardt, U.,
31 Dueñas, M., Grebenc, T., Griffith, G. W., Hartmann, M., Kirk, P. M., Kohout, P., Larsson, E.,
32 Lindahl, B. D., Lücking, R., Martín, M. P., Matheny, P. B., Nguyen, N. H., Niskanen, T., Oja, J.,
33 Peay, K. G., Peintner, U., Peterson, M., Põldmaa, K., Saag, L., Saar, I., Schübler, A., Scott, J. A.,
34 Senés, C., Smith, M. E., Suija, A., Taylor, D. L., Telleria, M. T., Weiss, M. and Larsson, K.-H.:
35 Towards a unified paradigm for sequence-based identification of fungi., *Mol. Ecol.*, 22(21), 5271–
36 7, doi:10.1111/mec.12481, 2013.

Deleted: .

- 1 Kopylova, E., Noé, L. and Touzet, H.: SortMeRNA: fast and accurate filtering of ribosomal RNAs
2 in metatranscriptomic data., *Bioinformatics*, 28, 3211–7, doi:10.1093/bioinformatics/bts611, 2012.
- 3 Kranner, I., Beckett, R., Hochman, A. and Nash, T. H.: Desiccation-Tolerance in Lichens: A
4 Review, *Bryologist*, 111(4), 576–593, doi:10.1639/0007-2745-111.4.576, 2008.
- 5 Lacey, J.: Spore dispersal — its role in ecology and disease: the British contribution to fungal
6 aerobiology, *Mycol. Res.*, 100(6), 641–660, doi:10.1016/S0953-7562(96)80194-8, 1996.
- 7 Larsson, K.-H., Parmasto, E., Fischer, M., Langer, E., Nakasone, K. K. and Redhead, S. A.:
8 Hymenochaetales: a molecular phylogeny for the hymenochaetoid clade, *Mycologia*, 98(6), 926–
9 936, doi:10.3852/mycologia.98.6.926, 2007.
- 10 Liu, K.-L., Porras-Alfaro, A., Kuske, C. R., Eichorst, S. A. and Xie, G.: Accurate, rapid taxonomic
11 classification of fungal large-subunit rRNA genes., *Appl. Environ. Microbiol.*, 78(5), 1523–33,
12 doi:10.1128/AEM.06826-11, 2012.
- 13 Martin, S. T., Andreae, M. O., Althausen, D., Artaxo, P., Baars, H., Borrmann, S., Chen, Q.,
14 Farmer, D. K., Guenther, A., Gunthe, S. S., Jimenez, J. L., Karl, T., Longo, K., Manzi, A., Müller,
15 T., Pauliquevis, T., Petters, M. D., Prenni, A. J., Pöschl, U., Rizzo, L. V., Schneider, J., Smith, J.
16 N., Swietlicki, E., Tota, J., Wang, J., Wiedensohler, A. and Zorn, S. R.: An overview of the
17 Amazonian Aerosol Characterization Experiment 2008 (AMAZE-08), *Atmos. Chem. Phys.*,
18 10(23), 11415–11438, doi:10.5194/acp-10-11415-2010, 2010.
- 19 Miadlikowska, J., Kauff, F., Hofstetter, V., Fraker, E., Grube, M., Hafellner, J., Reeb, V.,
20 Hodkinson, B. P., Kukwa, M., Lucking, R., Hestmark, G., Otalora, M. G., Rauhut, A., Budel, B.,
21 Scheidegger, C., Timdal, E., Stenroos, S., Brodo, I., Perlmutter, G. B., Ertz, D., Diederich, P.,
22 Lendemer, J. C., May, P., Schoch, C. L., Arnold, A. E., Gueidan, C., Tripp, E., Yahr, R., Robertson,
23 C. and Lutzoni, F.: New insights into classification and evolution of the Lecanoromycetes
24 (Pezizomycotina, Ascomycota) from phylogenetic analyses of three ribosomal RNA- and two
25 protein-coding genes, *Mycologia*, 98(6), 1088–1103, doi:10.3852/mycologia.98.6.1088, 2007.
- 26 Moore, D., Robson, G. D. and Trinci, A. P. J.: 21st Century Guidebook to Fungi, Cambridge
27 University Press., 2011.
- 28 Morgan, J. L., Darling, A. E. and Eisen, J. A.: Metagenomic sequencing of an in vitro-simulated
29 microbial community., *PLoS One*, 5(4), e10209, doi:10.1371/journal.pone.0010209, 2010.
- 30 Morris, C. E., Sands, D. C., Glaux, C., Samsatly, J., Asaad, S., Moukahel, A. R., Gonçalves, F. L.
31 T. and Bigg, E. K.: Urediospores of rust fungi are ice nucleation active at > –10 °C and harbor ice
32 nucleation active bacteria, *Atmos. Chem. Phys.*, 13(8), 4223–4233, doi:10.5194/acp-13-4223-
33 2013, 2013.
- 34 Morris, C. E., Sands, D. C., Vanneste, J. L., Montarry, J., Oakley, B., Guilbaud, C. and Glaux, C.:
35 Inferring the evolutionary history of the plant pathogen *Pseudomonas syringae* from its

- 1 biogeography in headwaters of rivers in North America, Europe, and New Zealand., *MBio*, 1(3),
2 e00107–10–, doi:10.1128/mBio.00107-10, 2010.
- 3 Morris, C. E., Sands, D. C., Vinatzer, B. A., Glaux, C., Guilbaud, C., Buffière, A., Yan, S.,
4 Dominguez, H. and Thompson, B. M.: The life history of the plant pathogen *Pseudomonas syringae*
5 is linked to the water cycle., *ISME J.*, 2(3), 321–34, doi:10.1038/ismej.2007.113, 2008.
- 6 Oksanen, J., Blanchet, F. G., Kindt, R., Legendre, P., Minchin, P. R., O’Hara, R. B., Simpson, G.
7 L., Solymos, P., Stevens, M. H. H. and Wagner, H.: *vegan*: Community Ecology Package, [online]
8 Available from: <http://cran.r-project.org/package=vegan>, 2013.
- 9 Onofri, S., Selbmann, L., Zucconi, L. and Pagano, S.: Antarctic microfungi as models for
10 exobiology, *Planet. Space Sci.*, 52(1-3), 229–237, doi:10.1016/j.pss.2003.08.019, 2004.
- 11 Pashley, C. H., Fairs, A., Free, R. C. and Wardlaw, A. J.: DNA analysis of outdoor air reveals a
12 high degree of fungal diversity, temporal variability, and genera not seen by spore morphology.,
13 *Fungal Biol.*, 116(2), 214–24, doi:10.1016/j.funbio.2011.11.004, 2012.
- 14 Peay, K. G., Baraloto, C. and Fine, P. V. A.: Strong coupling of plant and fungal community
15 structure across western Amazonian rainforests., *ISME J.*, 7(9), 1852–61,
16 doi:10.1038/ismej.2013.66, 2013.
- 17 Penton, C. R., Gupta, V. V. S. R., Tiedje, J. M., Neate, S. M., Ophel-Keller, K., Gillings, M.,
18 Harvey, P., Pham, A. and Roget, D. K.: Fungal Community Structure in Disease Suppressive Soils
19 Assessed by 28S LSU Gene Sequencing, *PLoS One*, 9(4), e93893,
20 doi:10.1371/journal.pone.0093893, 2014.
- 21 Penton, C. R., St Louis, D., Cole, J. R., Luo, Y., Wu, L., Schuur, E. A. G., Zhou, J. and Tiedje, J.
22 M.: Fungal diversity in permafrost and tallgrass prairie soils under experimental warming
23 conditions, *Appl. Environ. Microbiol.*, 79(22), 7063–72, doi:10.1128/AEM.01702-13, 2013.
- 24 Pöschl, U., Martin, S. T., Sinha, B., Chen, Q., Gunthe, S. S., Huffman, J. A., Borrmann, S., Farmer,
25 D. K., Garland, R. M., Helas, G. and others: Rainforest aerosols as biogenic nuclei of clouds and
26 precipitation in the Amazon, *Science* (80-), 329(5998), 1513–1516,
27 doi:10.1126/science.1191056, 2010.
- 28 Pouleur, S. S., Richard, C., Martin, J. G. and Antoun, H.: Ice nucleation activity in *Fusarium*
29 *acuminatum* and *Fusarium avenaceum*, *Appl. Environ. Microbiol.*, 58(9), 2960–4, 1992.
- 30 Pringle, A.: Asthma and the diversity of fungal spores in air., *PLoS Pathog.*, 9(6), e1003371,
31 doi:10.1371/journal.ppat.1003371, 2013.
- 32 Prosser, J. I.: Molecular and functional diversity in soil micro-organisms, *Plant Soil*, 244(1-2), 9–
33 17, doi:10.1023/A:1020208100281, 2002.

- 1 Pummer, B. G., Atanasova, L., Bauer, H., Bernardi, J., Druzhinina, I. S., Fröhlich-Nowoisky, J.
2 and Grothe, H.: Spores of many common airborne fungi reveal no ice nucleation activity in oil
3 immersion freezing experiments, *Biogeosciences*, 10(12), 8083–8091, doi:10.5194/bg-10-8083-
4 2013, 2013.
- 5 Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J. and Glöckner, F.
6 O.: The SILVA ribosomal RNA gene database project: improved data processing and web-based
7 tools., *Nucleic Acids Res.*, 41(Database issue), D590–6, doi:10.1093/nar/gks1219, 2013.
- 8 R Core Team: R: A Language and Environment for Statistical Computing, [online] Available from:
9 <http://www.r-project.org>, 2014.
- 10 Rajala, T., Peltoniemi, M., Hantula, J., Mäkipää, R. and Pennanen, T.: RNA reveals a succession
11 of active fungi during the decay of Norway spruce logs, *Fungal Ecol.*, 4(6), 437–448,
12 doi:10.1016/j.funeco.2011.05.005, 2011.
- 13 Richard, C., Martin, J.-G. and Pouleur, S.: Ice nucleation activity identified in some
14 phytopathogenic *Fusarium* species, *Phytoprotection*, 77(2), 83, doi:10.7202/706104ar, 1996.
- 15 Schmieder, R. and Edwards, R.: Quality control and preprocessing of metagenomic datasets,
16 *Bioinformatics*, 27(6), 863–864, doi:10.1093/bioinformatics/btr026, 2011.
- 17 Solhaug, K. A., Gauslaa, Y., Nybakken, L. and Bilger, W.: UV-induction of sun-screening
18 pigments in lichens, *New Phytol.*, 158(1), 91–100, doi:10.1046/j.1469-8137.2003.00708.x, 2003.
- 19 Sussman, A. S. and Douthit, H. A.: Dormancy in microbial spores, *Annu. Rev. Plant Physiol.*,
20 24(1), 311–352, 1973.
- 21 Tedersoo, L., Bahram, M., Polme, S., Koljalg, U., Yorou, N. S., Wijesundera, R., Ruiz, L. V.,
22 Vasco-Palacios, A. M., Thu, P. Q., Suija, A., Smith, M. E., Sharp, C., Saluveer, E., Saitta, A.,
23 Rosas, M., Riit, T., Ratkowsky, D., Pritsch, K., Poldmaa, K., Piepenbring, M., Phosri, C., Peterson,
24 M., Parts, K., Partel, K., Otsing, E., Nouhra, E., Njouonkou, A. L., Nilsson, R. H., Morgado, L. N.,
25 Mayor, J., May, T. W., Majuakim, L., Lodge, D. J., Lee, S. S., Larsson, K.-H., Kohout, P., Hosaka,
26 K., Hiiesalu, I., Henkel, T. W., Harend, H., Guo, L. -d., Greslebin, A., Grelet, G., Geml, J., Gates,
27 G., Dunstan, W., Dunk, C., Drenkhan, R., Dearnaley, J., De Kesel, A., Dang, T., Chen, X., Buegger,
28 F., Brearley, F. Q., Bonito, G., Anslan, S., Abell, S. and Abarenkov, K.: Global diversity and
29 geography of soil fungi, *Science* (80-.), 346(6213), 1256688–1256688,
30 doi:10.1126/science.1256688, 2014.
- 31 Vaitilingom, M., Deguillaume, L., Vinatier, V., Sancelme, M., Amato, P., Chaumerliac, N. and
32 Delort, A.-M.: Potential impact of microbial activity on the oxidant capacity and organic carbon
33 budget in clouds., *Proc. Natl. Acad. Sci. U. S. A.*, 110(2), 559–64, doi:10.1073/pnas.1205743110,
34 2013.
- 35 Wickham, H.: *ggplot2: elegant graphics for data analysis*, Springer New York., 2009.

1 Woo, A. C., Brar, M. S., Chan, Y., Lau, M. C. Y. Y., Leung, F. C. C. C., Scott, J. A., Vrijmoed, L.
 2 L. P. P., Zavar-Reza, P. and Pointing, S. B.: Temporal variation in airborne microbial populations
 3 and microbially-derived allergens in a tropical urban landscape, *Atmos. Environ.*, 74, 291–300,
 4 doi:http://dx.doi.org/10.1016/j.atmosenv.2013.03.047, 2013.


5 Yamamoto, N., Bibby, K., Qian, J., Hospodsky, D., Rismani-Yazdi, H., Nazaroff, W. W. and
 6 Peccia, J.: Particle-size distributions and seasonal diversity of allergenic and pathogenic fungi in
 7 outdoor air, *ISME J.*, 6(10), 1801–11, doi:10.1038/ismej.2012.30, 2012.

8 Yamamoto, N., Nazaroff, W. W. and Peccia, J.: Assessing the aerodynamic diameters of taxon-
 9 specific fungal bioaerosols by quantitative PCR and next-generation DNA sequencing, *J. Aerosol*
 10 *Sci.*, doi:10.1016/j.jaerosci.2014.08.007, 2014.

11 Zhang, N., Castlebury, L. A., Miller, A. N., Huhndorf, S. M., Schoch, C. L., Seifert, K. A.,
 12 Rossman, A. Y., Rogers, J. D., Kohlmeyer, J., Volkmann-Kohlmeyer, B. and Sung, G.-H.: An
 13 overview of the systematics of the Sordariomycetes based on a four-gene phylogeny, *Mycologia*,
 14 98(6), 1076–1087, doi:10.3852/mycologia.98.6.1076, 2007.


15

16 **Figures**




17

18 Figure 1. Basidiomycota dominate the total fungal community (mean relative abundance =
 19 90.2±6.9%). Bars are colored according to class-level taxonomic assignments. Taxonomy was
 20 assigned to representative sequences from each OTU.



1
 2 Figure 2. OTU-based community composition significantly differed across environment types
 3 (ADONIS, $R^2 = 0.167$, $p = 0.001$). Total and active communities in air samples (upper left)
 4 clustered together and separate from other environments indicating these communities are distinct
 5 from communities found in soils and on leaf surfaces. Sørensen similarities are depicted, ordinated
 6 via NMDS.



1
 2 Figure 3. Relative abundances of fungal phyla across environment types. The active atmospheric
 3 fungal community over the Amazon rainforest was more similar in phylum-level composition to
 4 fungal communities found in tropical soils and on plant leaves than was the total community. Error
 5 bar represent standard deviations.

6

7 **Appendix A**

8 **Mass-balance model**

9 We use a global, well-mixed, one-box material-balance model to explain the relative abundances
 10 of fungal cells measured as gene copies sampled in the active and total portions of atmospheric
 11 bioaerosols. By material-balance, for any taxon i within a biological community, the change in time
 12 in the abundance of fungal gene copies, N_i , must be equal to the difference in source and sinks:

13
$$\frac{dN_i}{dt} = \sum sources - \sum sinks \tag{A1}$$

14 Here we assume sources are equal to the emission of fungal gene copies from the Earth's surface
 15 into the atmosphere, E_i (gene copies/hour). We assume sinks are equal to deposition of fungal gene

1 copies out of the atmosphere back to the Earth's surface, $D_i = N_i k_i$ (gene copies/hour), where k_i
2 (1/hour) represents a first order deposition coefficient. We can rewrite Equation (A1) as:

$$3 \quad \frac{dN_i}{dt} = E_i - N_i k_i$$

4 We expect the terms E_i and k_i to vary as a function of life history traits including the method of cell
5 release into the atmosphere, the physiological state of sampled cells, and the aerodynamic diameter
6 of fungal taxa. In this case, Equation (A2) does not directly represent the entire airborne fungal
7 gene copy abundance. We assume that a first order approximation of fungal bioaerosol behavior
8 may be obtained by subdividing the particle distribution into two modes: vegetative cells, $N_{i,veg}$,
9 and spores, $N_{i,spores}$. We thus model fungal gene copy abundance as:

$$10 \quad N_i = N_{i,veg} + N_{i,spores}$$

11 We can then write and solve parallel versions of Equation (A2) for each mode. At steady state, the
12 expected gene copy abundance taxa i in each mode is:

$$13 \quad N_{i,veg} = \frac{E_{i,veg}}{k_{i,veg}}$$

$$14 \quad N_{i,spore} = \frac{E_{i,spore}}{k_{i,spore}}$$

15 Our interest lies in the two most common fungal phyla sampled in the atmosphere: Ascomycota,
16 N_A , and Basidiomycota, N_B . To make predictions about the expected relative abundance of gene
17 copies in these two groups, we make informed assumptions about the relative magnitude of their
18 respective emission and deposition rates. We begin by considering fungal spores. Although a few
19 empirical studies have suggested that Ascomycota are more abundant than Basidiomycota in likely
20 source environments including tropical soils (Kerekes et al., 2013) and leaf surfaces (Kembel and
21 Mueller, 2014), Basidiomycota (e.g. Agaricomycetes, the most abundant class of Basidiomycota
22 in our samples) produce orders of magnitude more spores per individual than Ascomycota (Elbert
23 et al., 2007; Pringle, 2013). For this reason, we assume the emission rate of Basidiomycota spores
24 is much greater than that of Ascomycota spores:

$$25 \quad E_{A,spores} \ll E_{B,spores}$$

1 Culture-based microscopy data suggests that spores of Ascomycota are typically larger than spores
2 of Basidiomycota (Elbert et al., 2007; Ingold, 2001; Yamamoto et al., 2014). Owing to the
3 difference in spore size, we expect deposition rate of Ascomycota spores to be greater than that of
4 Basidiomycota spores:

$$5 \quad k_{d,A,spores} > k_{d,B,spores}$$

6 Based on these assumptions, it follows that the expected number of Ascomycota spores in the
7 atmosphere will be less than the number of Basidiomycota spores:

$$8 \quad \frac{E_{A,spore}}{k_{A,spore}} \ll \frac{E_{B,spore}}{k_{B,spore}}$$

9 or

$$10 \quad N_{A,spores} \ll N_{B,spores}$$

11 We next consider fungal vegetative cells. Vegetative forms of Ascomycota are generally smaller
12 than vegetative forms of Basidiomycota (Moore et al., 2011). Many Ascomycota grow as filaments
13 or single cells which are small enough to be aerosolized (Després et al., 2012). In contrast, many
14 Basidiomycota grow as mushrooms, which are too large to be aerosolized (although debris from
15 mushrooms and their mycelia can be aerosolized). Due to this difference in the vegetative forms of
16 each group, we expect emission rate of vegetative Ascomycota to be greater than Basidiomycota:

$$17 \quad E_{A,veg} > E_{B,veg}$$

18 No comparative data currently exists on the relative deposition rate of vegetative cells across fungal
19 taxa. Research has shown that at the phylum level, the aerodynamic diameter of Ascomycota is
20 greater than that of Basidiomycota, resulting in a greater deposition rate overall for Ascomycota
21 (Yamamoto et al. 2014). However, this work did not differentiate between vegetative cells and
22 spores, and there is no *a priori* reason to assume that the deposition rate of Ascomycota vegetative
23 cells are less than or greater to that of Basidiomycota cells. For this reason, we make the null
24 assumption that the deposition rate of each group is equal:

$$25 \quad k_{d,A,veg} = k_{d,B,veg}$$

1 Based on these assumptions, we expect the number of vegetative Ascomycota genes to be greater
2 than the number of vegetative Basidiomycota genes:

$$3 \quad \frac{E_{A,veg}}{k_{A,veg}} > \frac{E_{B,veg}}{k_{B,veg}}$$

4 or

$$5 \quad N_{A,veg} > N_{B,veg} \quad (A3)$$

6 Equation (A3) predicts that Ascomycota will dominate the active fungal community in the
7 atmosphere.

8 Finally, we relate the abundance of Ascomycota and Basidiomycota gene copies in their totality
9 to ask if $N_A < N_B$ or $N_A \geq N_B$. $N_A < N_B$ if and only if:

$$10 \quad N_{A,veg} + N_{A,spores} < N_{B,veg} + N_{B,spores}$$

11 Rearranging terms and dividing both sides of the equation by $N_{B,spores}$ yields the inequality:

$$12 \quad \frac{N_{A,veg} - N_{B,veg}}{N_{B,spores}} + \frac{N_{A,spores}}{N_{B,spores}} < 1$$

13 or

$$14 \quad \frac{N_{A,veg} - N_{B,veg}}{N_{B,spores}} < 1 - \varepsilon$$

15 where $\varepsilon = \frac{N_{A,spores}}{N_{B,spores}}$. Empirical data on the discharge of Ascomycota and Basidiomycota spores
16 from fruiting bodies suggests that $\varepsilon \leq 0.01$ (Elbert et al. 2007). In this case $N_A < N_B$ if and only
17 if:

$$18 \quad \frac{N_{A,veg} - N_{B,veg}}{N_{B,spores}} < 0.99 \quad (A4)$$

19 We expect Equation A4 to hold due to the likelihood that spores greatly outnumber vegetative
20 cells in the atmosphere in both phyla. Spores can be actively discharged into the air, whereas
21 vegetative cells are not actively propelled into the atmosphere and require aerosolization by
22 mechanical forces like wind. Furthermore, empirical data suggests that vegetative cell fragments

1 constitute a small fraction (0.2-16% (Green et al., 2011)) of the total fungal biomass in the
2 atmosphere. For these reasons, we predict that

$$3 \quad N_A < N_B$$

4 Based on the conclusions of this model, we expect Basidiomycota will dominate the total
5 community, and Ascomycota will dominate the active community in the atmosphere. We note there
6 are many limitations to our model. First, we model fungal gene copy abundances assuming a well-
7 mixed atmosphere at steady state. Yet the atmosphere is a highly heterogeneous and dynamic
8 environment; the sampled air volume was likely neither well mixed nor at steady state over the
9 time intervals we measured. Second, we use a global model with emission and deposition as the
10 sole input and output, whereas a local model that incorporated site-specific environmental fate and
11 transport terms would likely provide more accurate expectations. Third, due to a paucity of data,
12 our estimates of fungal gene abundance levels rely on assumptions about the emission and
13 deposition rates of vegetative cells and spores across fungal taxonomic groups. Empirically derived
14 estimates of these model parameters would significantly improve our approach. Fourth, we do not
15 know to what extent vegetative cells and spores are associated with other particulate matter and
16 how this affects their deposition and emission rates. Adopting an approach to empirically estimate
17 the aerodynamic diameter of these fungal cell types across taxonomic groups would allow for
18 improved estimates of deposition rates (Yamamoto et al., 2014).