

Derivation of Greenhouse Gas emission factors for peatlands managed for extraction in the Republic of Ireland and the United Kingdom

D. Wilson¹, S.D. Dixon², R.R.E. Artz³, T. E.L. Smith⁴, C. D. Evans⁵, H.J.F. Owen⁴, E. Archer² and F. Renou-Wilson⁶

¹ Earthy Matters Environmental Consultants, Glenvar, Co. Donegal, Ireland. + 353 74 9177613, david.wilson@earthymatters.ie

² Department of Earth Sciences, University of Durham, Durham, United Kingdom +44 191 334 2356

³ The James Hutton Institute, Aberdeen, Scotland. + 44 844 928 5428

⁴ King's College London, Department of Geography, Strand, London, United Kingdom, + 44 207 8482525

⁵ Centre for Ecology and Hydrology, Bangor, Wales + 44 124 8374500

⁶ School of Biology & Environmental Science, University College Dublin, Ireland + 353 1 716 2440

1 **Abstract**

2 Drained peatlands are significant hotspots of carbon dioxide (CO₂) emissions and may also
3 be more vulnerable to fire with its associated gaseous emissions. Under the United Nations
4 Framework Convention on Climate Change (UNFCCC) and the Kyoto Protocol, greenhouse
5 gas (GHG) emissions from peatlands managed for extraction are reported on an annual basis.
6 However, the Tier 1 (default) emission factors (EFs) provided in the IPCC 2013 Wetlands
7 Supplement for this land use category may not be representative in all cases and countries are
8 encouraged to move to higher Tier reporting levels with reduced uncertainty levels based on
9 country or regional specific data. In this study, we quantified (1) CO₂-C emissions from
10 peat extraction sites in the Republic of Ireland and the United Kingdom, which were initially
11 disaggregated by land use type (industrial versus domestic peat extraction), and (2) a range of
12 GHGs that are released to the atmosphere with the burning of peat. Drainage related methane
13 (CH₄) and nitrous oxide (N₂O) emissions, as well as CO₂-C emissions associated with the off-
14 site decomposition of horticultural peat were not included here. Our results show that net
15 CO₂-C emissions were strongly controlled by soil temperature at the industrial sites (bare
16 peat), and by soil temperature and leaf area index at the vegetated domestic sites. Our derived
17 EFs of 1.70 (±0.47) and 1.64 (±0.44) t CO₂-C ha⁻¹ yr⁻¹ for the industrial and domestic sites
18 respectively, are considerably lower than the Tier 1 EF (2.8±1.7 t CO₂-C ha⁻¹ yr⁻¹) provided
19 in the Wetlands Supplement. We propose that the difference between our derived values and
20 the Wetlands Supplement value is due to differences in peat quality and, consequently,
21 decomposition rates. Emissions from burning of the peat (g kg⁻¹ dry fuel burned) were
22 estimated to be approximately 1346 (CO₂), 8.35 (methane, CH₄), 218 (carbon monoxide,
23 CO), 1.53 (ethane, C₂H₆), 1.74 (ethylene, C₂H₄), 0.60 (methanol, CH₃OH), 2.21 (hydrogen
24 cyanide, HCN) and 0.73 (ammonia, NH₃) and emphasises the importance of understanding
25 the full suite of trace gas emissions from biomass burning. Our results highlight the
26 importance of generating reliable Tier 2 values for different regions and land-use categories.
27 Furthermore, given that the IPCC Tier 1 EF was only based on 20 sites (all from
28 Canada/Fenno-Scandia) we suggest that data from another 9 sites significantly expands the
29 global dataset, as well as adding a new region.

30

31

32

1 **1 Introduction**

2 Greenhouse gas (GHG) emissions to the atmosphere have increased significantly since pre-
3 industrial times as a direct result of human activities, such as fossil fuel burning, cement
4 production and land use changes (IPCC, 2013). The Intergovernmental Panel on Climate
5 Change (IPCC) have estimated in their Fifth Assessment Report (AR5) that around one third
6 of all anthropogenic emissions of carbon dioxide (CO₂) for the period 1750-2011, were
7 caused by land use changes (IPCC, 2013). From 2000-2009, the Agriculture, Forestry and
8 Other Land-Use (AFOLU) sector accounted for 24% of all global GHG emissions (around 10
9 Gt CO₂-eq yr⁻¹), with emissions from peatland drainage and burning alone estimated at
10 around 0.9 Gt CO₂-eq yr⁻¹.

11 Natural (i.e. undrained) peatlands function as long term carbon (C) stores as the sequestration
12 of CO₂ over time is greater than the amount of C that is emitted from the peatland as methane
13 (CH₄) and leached in waterborne exports (Roulet et al., 2007; Nilsson et al., 2008; Koehler et
14 al., 2011; Gažovič et al., 2013). Key to this role is the position of the water table, which
15 largely dictates the rate of decomposition within the peatland. When the water table is
16 positioned close to the peat surface, the breakdown and degradation of organic matter
17 typically proceeds very slowly in the absence of oxygen. As a consequence, there is an
18 accumulation of peat (and C within) (Dise, 2009).

19 In the Republic of Ireland (ROI) and the United Kingdom (UK), peat has been extracted for
20 energy use for many centuries (Chapman et al., 2003; Renou et al., 2006). Traditionally, this
21 involved the manual removal of the peat i.e. hand cutting, however this has been largely
22 superseded by highly mechanised methods to extract the peat for both energy and horticulture
23 requirements. In the ROI, over 4 million tonnes of peat per annum are industrially extracted
24 from approximately 50,000 ha to provide ca. 5.5% of primary energy requirements (Howley
25 et al., 2012) and for use in horticulture. A further 0.4 million tonnes per year is likely burned
26 for domestic heating (Duffy et al., 2014) and may impact as much as 600,000 ha of peatlands
27 (Wilson et al., 2013b). Although peat extraction areas in the UK have generally declined over
28 the last few decades, approximately 0.8 million tonnes of peat is still extracted each year in
29 England and Scotland (Webb et al., 2014), although it is UK Government policy to phase out
30 peat extraction in England by 2030 (Department of Environment Food and Rural Affairs,
31 2011). Peat extraction areas in Wales are small (482 ha) and have remained unchanged in the
32 1991-2010 period (Webb et al., 2014). In Northern Ireland, the area of peatland utilised for
33 fuel (mechanical and hand cutting) has declined considerably in the 1990-2008 period,

1 although a slight increase in the areas used for horticulture have been recorded (Tomlinson,
2 2010).

3 In industrial peatlands, the extraction of peat is facilitated by the installation of drainage
4 ditches at regular (typically 15-30m) intervals across the peatland. For peat used for
5 horticultural purposes, the more fibrous upper layers (e.g. *Sphagnum* peat) are extracted and
6 utilised. If the peat is to be used for energy production the more highly decomposed peat is
7 milled, dried in the production fields and removed for immediate use or stockpiled for later
8 requirements. Peat extraction ceases for energy production when either the sub-peat mineral
9 soil is reached, large quantities of fossilised timber are encountered or drainage is no longer
10 practical (Farrell and Doyle, 2003). For peatlands used for the provision of domestic heating,
11 the peat is either removed by a digger from the margins of peatlands, placed in a tractor
12 mounted hopper and extruded onto the surface of the peatland, or the peat is extruded onto
13 the surface of the peatland from openings made in the peat by a chain cutter. Over a period of
14 weeks the peat is dried *in situ* and removed from the site. The effect of peat extraction on the
15 hydrological functioning is marked by a large fall in the water level either throughout the
16 peatland (industrial) or at the margins of the peatland (domestic). In the latter, significant
17 water level drawdown is also experienced further inward towards the centre of the peatland
18 (Schouten, 2002).

19 The impact of drainage on C cycling in peatlands has been widely documented. In general, a
20 lowering of the water table leads to increased CO₂ emissions (Silvola et al., 1996; Salm et al.,
21 2012; Haddaway et al., 2014) as the aerobic layer is deepened and mineralisation rates are
22 accentuated. Concurrently, CH₄ emissions (with the exception of ditches) may decrease or
23 cease (Salm et al., 2012; Turetsky et al., 2014), waterborne C exports may increase (Strack et
24 al., 2008; Evans et al., 2015) and there may be a heightened risk of C loss through fire
25 (Turetsky et al., 2015). In the case of peat extraction, C cycling may be further altered by the
26 removal of vegetation (Waddington and Price, 2000), and losses of windblown particulate
27 organic carbon (POC) may be exacerbated from the bare peat surfaces (Lindsay, 2010).

28 Under the United Nations Framework Convention on Climate Change (UNFCCC) and the
29 Kyoto Protocol, “Annex 1” countries (i.e. countries that have committed to targets that limit
30 or reduce emissions) are obligated to prepare annual National Inventory Reports (NIR) and
31 up-to-date annual inventories, detailing GHG emissions and removals from six different
32 sectors. Emissions associated with off-site peat combustion are reported under the Energy

1 sector and are not considered further here. The recent IPCC Wetlands Supplement (IPCC,
2 2014) to the 2006 Good Practice Guidance (GPG) (IPCC, 2006) derived new Tier 1 emission
3 factors (EFs) for drained organic soils that differentiated between on-site emissions (e.g.
4 $\text{CO}_2\text{-C}_{\text{on-site}}$, fire) and off-site losses (e.g. leaching of waterborne C). In the case of peatlands
5 managed for extraction in the temperate climate zone, the $\text{CO}_2\text{-C}_{\text{on-site}}$ values have increased
6 from 0.2 (nutrient poor/bogs) and 1.1 (nutrient rich/fens) $\text{t CO}_2\text{-C ha}^{-1} \text{ yr}^{-1}$ in the 2006 GPG to
7 a single higher EF of 2.8 $\text{t CO}_2\text{-C ha}^{-1} \text{ yr}^{-1}$ (covering the entire boreal and temperate regions)
8 in the Wetlands Supplement. On-site burning directly consumes aboveground C stocks
9 (prescribed and wildfire burning) and the underlying peat C store (wildfire burning), and
10 rapidly releases both gases (e.g. CO_2 , CH_4) and particulates (e.g. black carbon) to the
11 atmosphere. In the Wetlands Supplement, an EF for GHG emissions from prescribed fire on
12 drained peatlands is not provided due to a paucity of published data at present. However,
13 emissions from wildfires are addressed and EFs of 362, 9 and 207 g kg^{-1} dry fuel burned is
14 provided for $\text{CO}_2\text{-C}$, CH_4 and CO respectively with a proviso that they were derived from a
15 very small dataset.

16 Given the relatively large areas under peat extraction in both the ROI and the UK, a move
17 from Tier 1 to higher reporting levels is desirable, particularly as (a) a wide range in
18 uncertainty is associated with the IPCC Tier 1 values (1.1 - 4.2 $\text{t CO}_2\text{-C ha}^{-1} \text{ yr}^{-1}$), which
19 reflects the disparity in emissions from drained peatlands from different climate zones and
20 nutrient composition, (b) the most recently published annual CO_2 flux estimates (not included
21 in the derivation of IPCC Tier 1 values) also display a very wide amplitude (cf. Järveoja et
22 al., 2012; Mander et al., 2012; Salm et al., 2012; Strack et al., 2014), (c) no data from ROI or
23 UK peatlands were included in the IPCC derivation, which might mean that the Tier 1 value
24 may not be appropriate for these countries, and (d) no distinction is made between industrial
25 or domestic extraction sites, despite large differences in their drainage, vegetation cover and
26 management characteristics. In addition, previous studies of peatland fire EFs have focused
27 on the boreal peatlands of Alaska (Yokelson et al., 1997) and Canada (Stockwell et al.,
28 2014); and the temperate peatlands of Minnesota (Yokelson et al., 1997) and North Carolina
29 (Stockwell et al., 2014). These studies found that the smouldering combustion of peats
30 associated with low combustion efficiency leads to relatively lower CO_2 emissions
31 (compared with other ecosystems), and much higher carbon monoxide (CO), CH_4 , and other
32 non- CH_4 hydrocarbon emissions. Therefore, it is important to quantify emissions of these

1 gases as they include strong GHGs (e.g. CH₄) and reactive gases responsible for tropospheric
2 ozone formation and poor air quality (e.g. CO, ammonia (NH₃), hydrogen cyanide (HCN)).

3 The objectives of the study are (1) to provide estimates of the annual CO₂-C exchange (i.e.
4 CO₂-C_{on-site}) for 9 peat extraction sites in the ROI and the UK, (2) to derive regional specific
5 CO₂-C EFs for drained peat extraction areas that would permit ROI and the UK to progress to
6 the Tier 2 reporting level, (3) analyse the factors that influence CO₂-C dynamics in this
7 region (i.e. land use, climate etc.), and (4) to report GHG emissions associated with the
8 burning of Irish *Sphagnum* moss peat in the first laboratory study to investigate fire emissions
9 from European temperate peats.

10

11 **2 Materials and Methods**

12 **2.1 Study sites**

13 The study sites were located at 9 peat extraction areas in the ROI and the UK with a history
14 of either industrial peat (IP) or domestic peat (DP) extraction (Table 1). Boora (IP1),
15 Blackwater (IP2), Bellacorick (IP3), Turraun (IP4), Middlemuir Moss (IP5) and Little
16 Woolden Hall Moss (IP6) are industrial cutaway peatlands where significant areas of bare
17 peat (i.e. unvegetated microsites) have remained following the cessation of milled peat
18 extraction. At IP6, milled peat is currently extracted from areas close (<150m) to the study
19 site. The IP sites are former raised bogs with the exception of IP3, which is a former Atlantic
20 blanket bog. At all sites, the drainage ditches have remained functional. Here we define
21 “drained” as a mean annual water table position deeper than -20cm (Couwenberg and Fritze,
22 2012; Strack et al., 2014). Physico-chemical characteristics of all the sites are detailed in
23 Table 1.

24 At Clara (DP1), Glenlahan (DP2) and Moyarwood (DP3) the peat has been extracted from
25 the margins of the sites for use in domestic heating. In the case of Clara, peat extraction was
26 an ongoing activity at the time of our study despite the designation of the site as a Special
27 Area of Conservation (SAC). DP1 and DP3 are raised bogs and DP2 is a mountain blanket
28 bog. The vegetation component at all the sites is species poor and is composed mainly of ling
29 heather (*Calluna vulgaris*), cross leaved heather (*Erica tetralix*) and lichens (*Cladonia* spp.)
30 A continuous water table level was not observed at DP2, as the relatively shallow peat

1 deposit (~40cm) over bedrock at that site was prone to drying out at various times throughout
2 the study.

3 **2.2 Climatic conditions**

4 All the sites are located within the temperate zone as defined by IPCC (2006), and are
5 characterised by an oceanic climate with prevailing south-west winds, mild mean annual air
6 temperatures (8 to 10.3°C) and moderate to high annual rainfall (804 to 1245 mm) (Table 1).

7 **2.3 Environmental monitoring**

8 At each site, 3-9 aluminium square collars (60 x 60 cm) were inserted to a depth of 30cm into
9 the peat. At IP6, smaller circular plastic collars were used (15cm diameter) to facilitate the
10 use of the CPY-4 chamber (PP Systems, UK) at that site. Soil loggers (μ logger; Zeta-tec,
11 UK, Hobo External Data Loggers; Onset Computer Corporation, MA, USA or Comark
12 N2012 Diligence Loggers, Norwich, UK) were established in all the IP sites and recorded soil
13 temperatures (°C) at hourly intervals. Weather stations were installed at all the DP sites and
14 recorded photosynthetic photon flux density (PPFD; $\mu\text{mol m}^{-2} \text{s}^{-1}$) and soil temperatures (5
15 and 10cm depths) at 10 minute intervals. At DP3, soil volumetric moisture content (VMC, %)
16 was also recorded (at 10 min intervals) by the weather station at that site. At sites IP5 and
17 IP6, soil temperature was only measured manually during CO₂ flux measurements. In order
18 to estimate soil temperature at times where data was lacking at these two sites, a regression
19 based approach between manually recorded T_{5cm} and air temperature recorded at 15 min
20 intervals by a logger on the site was used to gap fill the data ($r^2 = 88.7\%$). Water table level
21 (WT) was manually measured from dipwells (internal diameter 2 cm) inserted adjacent to
22 each collar. Wooden boardwalks were established at each site (exception IP6).

23 **2.4 Leaf area index (LAI)**

24 At the IP sites, the vegetation had been removed prior to the commencement of peat
25 extraction and virtually no natural recolonization has taken place following cessation of peat
26 extraction. However, at the DP sites a vegetation component was present and in order to
27 incorporate the seasonal dynamics of the plants into CO₂-C exchange models, the leaf area
28 index (LAI) was estimated for each of the collars. This involved accounting for the green
29 photosynthetic area of all vascular plants (leaves and stems) within the collar at monthly
30 intervals. In short, the number of leaves and stems were counted from five subplots (8 x 8cm)

1 within each collar. The size (length, width) of the leaves was measured from sample plants
2 outside the collars. The LAI was then calculated by multiplying the estimated number of
3 leaves by an area estimate of the leaf. Moss and lichen % cover was estimated at the same
4 time. Species-specific model curves were applied to describe the phenological dynamics of
5 the vegetation of each collar, and the models (vascular plants and moss) were summed to
6 produce a plot-specific LAI. For a detailed description of the method see Wilson et al. (2007).
7 At site DP1 only, the vegetation was removed by regular clipping from one third of the
8 collars, in order to provide an estimate of the heterotrophic contribution (R_H) to ecosystem
9 respiration (R_{eco}).

10 **2.5 On site carbon dioxide flux estimation**

11 **2.5.1 Field measurements**

12 At sites IP1-5 and DP1-3, R_{eco} was measured with a static polycarbonate chamber (60 x 60 x
13 33 cm) equipped with two internal fans to ensure mixing of the air within the chamber, and a
14 cooling system (submerged ice packs, and pumped water to a radiator located within the
15 chamber) to maintain the temperature within the chamber close to the ambient air
16 temperature (for a more detailed description see Alm et al., 2007b). At IP6, R_{eco} was
17 measured with a CPY-4 (PP Systems, UK) clear acrylic chamber (14.6 cm diameter, 14.5 cm
18 height). The CPY-4 chamber was equipped with an internal fan, PPFd sensor and thermistor.
19 Sampling was carried out at fortnightly or monthly (winter) intervals (2-4 measurements per
20 collar per measurement day). For each R_{eco} flux measurement, the chamber was placed in a
21 water-filled channel at the top of the collar or connected with a rubber gasket (IP5), covered
22 with an opaque cover and the CO_2 concentration (ppmv) in the chamber headspace was
23 measured at 15-second (5-second at IP6) intervals over a period of 60-180 seconds using a
24 portable CO_2 analyser (EGM-4; PP Systems, UK). Concurrently, air temperature ($^{\circ}C$) within
25 the chamber and soil temperatures at 5, 10 and 20 cm depths were recorded at each collar
26 (soil temperature probe; ELE International, UK). The WT position relative to the soil surface
27 was manually measured with a water level probe (Eijkelkamp Agrisearch Equipment, The
28 Netherlands). At the DP sites, net ecosystem exchange (NEE) was measured with the same
29 polycarbonate chambers described above under a range of ambient light levels (PPFD; μmol
30 $m^{-2} s^{-1}$) prior to R_{eco} measurements. NEE measurements were carried out between 8 am and
31 6pm in the summer and between 9am and 3pm in the winter (3 to 8 measurements per collar
32 per measurement day) to ensure that the maximum PPFd was reached at each measurement

1 date. Artificial shading was used in the early morning to obtain low PPFD levels (<100 μmol
2 $\text{m}^{-2} \text{s}^{-1}$). PPFD was recorded from a sensor (PAR-1. PP Systems) located within the chamber.
3 The portable CO_2 analysers were regularly calibrated with a CO_2 standard gas.

4 **2.5.2 Flux calculations**

5 Flux rates ($\text{mg CO}_2\text{-C m}^{-2} \text{h}^{-1}$) were calculated as the linear slope of the CO_2 concentration in
6 the chamber headspace over time, with respect to the chamber volume, collar area and air
7 temperature. A flux was accepted if the coefficient of determination (r^2) was at least 0.90. An
8 exception was made in cases where the flux was close to zero (mainly in winter time where
9 soil processes are typically slower) and the r^2 is always low (Alm et al., 2007b). In these
10 cases the flux data were examined graphically and fluxes with obvious non-linearity (due to
11 chamber leakage, fan malfunction etc.) were discarded. The remainder were accepted
12 provided that some of the environmental variables measured at the same time (e.g. soil
13 temperature) were sufficiently low to account for the low flux values (Wilson et al., 2013a).
14 In this study, we followed the sign convention whereby positive values indicated a $\text{CO}_2\text{-C}$
15 flux from the peatland to the atmosphere (source) and negative values indicated a flux from
16 the atmosphere to the peatland (sink). Gross primary production (GPP) was calculated as
17 NEE minus R_{eco} (Alm et al., 2007b), and the closest R_{eco} flux value in time to a NEE flux
18 value was used.

19 **2.5.3 Modelling**

20 Statistical and physiological response models (Alm et al., 2007b) were constructed and
21 parameterised for each study site. Model evaluation was based on the following criteria; (a)
22 statistically significant model parameters ($p < 0.05$), (b) lowest possible standard error of the
23 model parameters and (c) highest possible coefficient of determination (adjusted r^2) (see
24 Laine et al., 2009). The basic R_{eco} models, based upon the Arrhenius equation (Lloyd and
25 Taylor, 1994), are non-linear models related to soil temperature. GPP was related to PPFD
26 using the Michaelis–Menten type relationship that describes the saturating response of
27 photosynthesis to light (Tuittila et al., 1999). GPP model coefficients and associated standard
28 errors were estimated using the Levenberg-Marquardt multiple non-linear regression
29 technique (IBM SPSS Statistics for Windows, Version 21.0. Armonk, NY, USA). During
30 model construction, the relationship between R_{eco} or GPP and a range of independent
31 environmental variables (recorded in conjunction with flux measurements) was tested. Only

1 variables that increased the explanatory power of the model (i.e. improved r^2 values) were
2 included. The models were accepted if the residuals were evenly scattered around zero.

3 **2.5.4 Annual CO₂–C balance**

4 The response functions estimated for R_{eco} and GPP were used for the reconstruction of the
5 annual CO₂-C balance. R_{eco} fluxes were reconstructed for each collar in combination with an
6 hourly time series of (1) $T_{5\text{cm}}$, (2) VMC (at DP3) recorded by the data loggers or (3) WT
7 depths linearly interpolated from weekly measurements. The annual CO₂-C balance (g C m^{-2}
8 yr^{-1}) was calculated for each sample plot by integrating the hourly R_{eco} values over each 12-
9 month period. (Note: integration periods vary between study sites; see Table 1). At the DP
10 sites, GPP was reconstructed in combination with (1) PPFD values recorded by the weather
11 station, (2) plot specific modelled LAI and (3) an hourly time series of $T_{5\text{cm}}$ (DP1 only). At
12 the DP sites, annual NEE was calculated as annual GPP + annual R_{eco} .

13 **2.5.5 Statistical analysis**

14 The CO₂-C flux data (R_{eco} for the IP sites, and R_{eco} and GPP for the DP sites) had a non-
15 normal distribution, so the non-parametric Kruskal-Wallis ($p=0.05$) and Mann-Whitney tests
16 were used to test for differences between sites. Uncertainty in reconstructed annual R_{eco} and
17 GPP was calculated by summing up the maximum and minimum standard errors associated
18 with each of the model parameters (e.g. Drösler, 2005; Elsgaard et al., 2012; Renou-Wilson
19 et al., 2014). Uncertainty in the annual R_{eco} or NEE estimate was calculated following the law
20 of error propagation as the square root of the sum of the squared standard errors of GPP and
21 R_{eco} (IPCC, 2006).

22 **2.6 Peat fire emissions**

23 Around 5 kg (dry mass) of loose Irish *Sphagnum* moss peat (H2-H3 on the von post
24 decomposition scale) was used for measuring fire EFs. Subsamples of the peat were taken
25 and placed into a 22 x 12 x 10 cm open-topped insulated chamber. The chamber was
26 constructed from lightweight Celcon insulation blocks and was used to replicate natural
27 surface combustion conditions, leaving only one surface of the peat exposed to open air
28 thereby reducing heat loss and oxygen exchange from the other surfaces, in accordance with
29 the suggested peat combustion methodology of Rein et al. (2009). Each sample was dried in
30 an oven overnight at 60°C. In order to produce comparable replicates, the samples for the
31 burning experiment had to be dried to an absolute dry base to increase ignition probability

1 (Frandsen, 1997) and encourage pyrolysis (Rein et al., 2009). Following drying, the chamber
2 and sample were placed in a fume cupboard under controlled air flow conditions and the peat
3 was ignited using a coiled nichrome wire heated to ~600°C and placed in contact with the
4 surface of the peat. This also best represents natural ignition conditions (e.g. from a surface
5 shrub fire), also in accordance with the methodology of Rein et al. (2009). Once ignited, each
6 1 kg sample proceeded to burn for ~90 minutes. The resulting smoke was continuously
7 sampled using a pump and a 90 cm sample line with a funnel held ~12 cm above the
8 smouldering peat. The smoke was sampled into an 8.5 litre infrared White (multipass) cell
9 (Infrared Analysis, Inc.) where infrared spectra were collected using a Fourier Transform
10 Infrared (FTIR) spectrometer. Analysis of the FTIR spectra was performed using the Multi
11 Atmospheric Layer Transmission (MALT) software (Griffith, 1996), yielding trace gas mole
12 fractions inside the White cell, from which emissions factors may be calculated. A full
13 description of how EFs may be calculated from FTIR measurements of gas mole fractions is
14 given in Paton-Walsh et al. (2014) and Smith et al. (2014). Here we use the C mass balance
15 approach to calculate EFs for CO₂ and CO (Eq. 1 in Paton-Walsh et al., 2014). The C content
16 of the peat (required for calculating EFs via the C mass balance approach) is assumed to be
17 53.3%, as measured in Scottish sphagnum moss peat (Cancellieri et al., 2012). For all other
18 gas species considered in the study; CH₄, ethylene (C₂H₄), ethane (C₂H₆), methanol
19 (CH₃OH), HCN, NH₃), we use their respective emission ratios to CO and the EF for CO to
20 calculate EFs (via Eq. 5 in Paton-Walsh et al., 2014).

21 Combustion efficiency is a measure of the amount of fuel carbon released as CO₂, and may
22 be approximated using the Modified Combustion Efficiency (MCE) formula, which requires
23 only a measurement of CO and CO₂ rather than all the C containing gases (Yokelson et al.,
24 2008):

$$25 \quad MCE = \frac{\Delta CO_2}{\Delta CO_2 + \Delta CO} \quad (1)$$

26 where ΔCO_2 and ΔCO represent the elevated mixing ratios of these gases (the difference
27 between mixing ratios measured in biomass burn emissions and those in the ambient air).
28 MCE is often expressed as a percentage. Generally, an MCE lower than 0.9 (90%) is
29 considered a low combustion efficiency burn (Lobert et al., 1991; Yokelson et al., 1996).

30

31 **3 Results**

1 **3.1 Environmental variables**

2 Annual rainfall varied between sites and between years (Fig. 1). The wettest site was DP3
3 (1390 mm), and the driest was IP6 (746 mm) in the first year of measurements at that site. All
4 multi-year sites displayed inter-annual variation in rainfall with the largest differences
5 observed in IP4 (210 mm difference in annual rainfall between years). Annual rainfall at IP2,
6 IP5, DP1, DP2 and DP3 was above the long-term average in all years. IP1 and IP4 were
7 wetter than the long-term average in one of the years and drier in the other. IP3 and IP6 were
8 drier than the long-term average. The mean annual water table was below -20cm at all sites in
9 all years (Fig. 1). The deepest mean annual values were at IP1 (-60cm) and the shallowest at
10 IP3, 4 and 5 (-25cm). Mean water table position tracked annual rainfall (i.e. higher rainfall
11 resulted in higher water table positions) in all multi-year sites with the exception of IP1.

12 The highest mean annual soil temperature ($T_{5\text{cm}}$) value (12.7°C) was recorded at IP4 and the
13 lowest at IP5 (6.7°C) and inter-annual variation was evident in the multi-year sites (Fig. 1).
14 The lowest hourly $T_{5\text{cm}}$ value (-12.9°C) was recorded at IP5 and the highest (28.4°C) at IP4
15 (Fig. 2). The proportion of hourly $T_{5\text{cm}}$ values less than 0°C ranged from 0% (IP3) to 13.8%
16 (IP5), and the proportion of values greater than 20°C ranged from 0.2% (IP5) to 5.3% (IP2)
17 (Fig.2).

18 **3.2 On-site carbon dioxide fluxes**

19 At the IP sites, R_{eco} fluxes ranged from 0 to $133 \text{ mg CO}_2\text{-C m}^{-2} \text{ hr}^{-1}$ and differed significantly
20 between sites (Fig. 3a Kruskal-Wallis, $H=98.59$). Site IP4 had significantly higher R_{eco} flux
21 values than all the other IP sites (Mann Whitney $p<0.001$) and IP5 had significantly lower
22 flux values than IP2, IP4 and IP6 (Mann Whitney $p<0.001$) but not IP1 and IP3 (Mann
23 Whitney $p=0.31$). At the DP sites, R_{eco} fluxes ranged from 12 to $200 \text{ mg CO}_2\text{-C m}^{-2} \text{ hr}^{-1}$ and
24 there was a significant difference in R_{eco} fluxes between the DP sites (Fig. 3b Kruskal-Wallis,
25 $H=37.52$) but no significant difference between DP1 and DP2 (Mann Whitney $p=0.075$). R_{eco}
26 values differed significantly between the IP and DP sites (Kruskal-Wallis, $H=395.22$).
27 Measured NEE (at $\text{PPFD}>1000 \mu\text{mol m}^{-2} \text{ s}^{-1}$) ranged from 60 to $-325 \text{ mg CO}_2\text{-C m}^{-2} \text{ hr}^{-1}$ at
28 the DP sites and values differed significantly between sites (Fig. 3c Kruskal-Wallis,
29 $H=90.82$).

30 **3.2.1 Modelling**

1 At sites IP6 and DP2, T_{5cm} was the sole explanatory variable in the R_{eco} models (Eq.2) and
 2 explained 32% and 42% respectively of the variability in fluxes. The addition of water table
 3 to the R_{eco} model (Eq.3) slightly improved the explanatory power and the model explained
 4 between 55% and 85% of the variability at IP1-4 and 69% at DP1. No relationship between
 5 R_{eco} and WT was observed at DP3, but the addition of VMC (Eq. 4) also slightly improved
 6 the explanatory power of the model (78%). At IP5, the data were too limited (n=22) to
 7 construct a reliable model that satisfied the criteria outlined in section 2.5.3. Instead, we
 8 calculated monthly mean values and integrated these values over the 12 month study period.

$$9 \quad R_{eco} = a * \exp * \left[b \left(\frac{1}{T_{REF}-T_0} - \frac{1}{T-T_0} \right) \right] \quad (2)$$

$$10 \quad R_{eco} = a * \exp * \left[b \left(\frac{1}{T_{REF}-T_0} - \frac{1}{T-T_0} \right) \right] * WT \quad (3)$$

$$11 \quad R_{eco} = a * \exp * \left[b \left(\frac{1}{T_{REF}-T_0} - \frac{1}{T-T_0} \right) \right] * VMC \quad (4)$$

12 where R_{eco} is ecosystem respiration, T_{REF} is reference temperature set at 283.15 K, T_0 is the
 13 (minimum) temperature at which respiration reaches zero and is set here at 227.13 K, T is the
 14 soil temperature at 5 cm depth, WT is water table depth, VMC is volumetric moisture
 15 content, a and b are fitted model parameters.

16 A strong relationship was observed between GPP and PPFD at the DP sites. It was the sole
 17 explaining variable at DP2 (Eq. 5) where it accounted for 70% of the variation. The addition
 18 of LAI (Eq. 6) increased the explanatory power of the GPP model at DP3 (59%) and the
 19 addition of LAI and T_{5cm} resulted in 62% of the variation explained at DP1.

$$20 \quad GPP = P_{max} \left(\frac{PPFD}{PPFD + k_{PPFD}} \right) \quad (5)$$

$$21 \quad GPP = P_{max} \left(\frac{PPFD}{PPFD + k_{PPFD}} \right) * LAI \quad (6)$$

$$22 \quad GPP = P_{max} \left(\frac{PPFD}{PPFD + k_{PPFD}} \right) * LAI * T_{5cm} \quad (7)$$

23 where GPP is gross primary productivity, P_{max} is maximum photosynthesis, PPFD is
 24 photosynthetic photon flux density, k_{PPFD} is the PPFD value at which GPP reaches half its
 25 maximum, LAI is leaf area index, T_{5cm} is soil temperature at depth of 5 cm.

26 **3.2.2 Annual CO₂-C balance**

1 The annual CO₂-C balance varied both spatially (between sites) and temporally (multi-year
2 sites) (Figs. 4 and 5). In the IP sites, emissions ranged from 93 g CO₂-C m⁻² yr⁻¹ (IP5) to 304
3 g CO₂-C m⁻² yr⁻¹ (IP4). Annual emissions varied considerably within the multi-year sites,
4 where coefficient of variation values ranged from 4% (IP1) to 20% (IP2). As would be
5 expected given the close relationship observed between soil temperature and CO₂-C fluxes, a
6 noticeable increase in modelled CO₂-C emissions was observed during the summer months at
7 all sites (Fig. 4), although the rate of the increase varied somewhat in strength between years
8 in the multi-year sites as a function of measured T_{5cm} and WT (where applicable). In the DP
9 sites (Fig. 5), annual GPP and R_{eco} were highest in DP1 (-526 and 702 g CO₂-C m⁻² yr⁻¹
10 respectively), intermediate in DP2 (-484 and 687 g CO₂-C m⁻² yr⁻¹ respectively) and lowest in
11 DP3 (-319 and 434 g CO₂-C m⁻² yr⁻¹ respectively). The DP sites were a net annual CO₂-C
12 source with the highest emissions observed at DP2 (203 g CO₂-C m⁻² yr⁻¹), intermediate at
13 DP1 (176 g CO₂-C m⁻² yr⁻¹) and lowest at DP3 (114 g CO₂-C m⁻² yr⁻¹). Estimated emissions
14 from heterotrophic respiration (R_H) at DP1 were 344 g CO₂-C m⁻² yr⁻¹, which equates to 49%
15 of R_{eco} at that site. Applying this proportional value to the other DP sites, we estimate that R_H
16 emissions to be 337 and 213 g CO₂-C m⁻² yr⁻¹ at DP2 and DP3 respectively.

17 **3.3 Drivers of annual CO₂-C_{on site}**

18 No relationships were observed between annual CO₂-C balances (NEE) and nutrient
19 concentrations, water table levels (average, maximum or minimum) or the von Post scale at
20 either the IP or DP (p>0.05) sites. A strong relationship (r²=0.63) between average soil
21 temperature at 5cm depth and R_{eco} was very evident across the IP sites (Fig. 6); the highest
22 annual emissions and highest average soil temperatures were associated with IP4 and the
23 lowest at IP5. The variation in NEE between the DP sites appeared to be related to
24 differences in LAI (Fig. 6), however the number of sites was very small (n=3) and some
25 caution must be used in this regard.

26 **3.4 Emission factors**

27 Using a single mean value for each multi-year site and for its associated uncertainty (IPCC,
28 2014), an EF was calculated for each land use category. The derived EFs for the IP and DP
29 sites were 1.70 (±0.47) and 1.64 (±0.44) t CO₂-C ha⁻¹ yr⁻¹ respectively (Table 2). The 95%
30 confidence intervals associated with the derived EFs were ±28% and ±26% for the IP and the

1 DP sites respectively. There was no significant difference in the EF values between the IP
2 and DP sites ($p=0.90$).

3 **3.5 Peat fire emission factors**

4 Mean modified combustion efficiency (MCE) and EFs with their standard deviations for
5 eight trace gas species were calculated from measurements of five Irish sphagnum moss peat
6 samples (Table 3). The peat burned with a mean MCE of 0.837 (± 0.019) typical of
7 smouldering combustion (e.g. Yokelson et al., 1996; Bertschi et al., 2003). Emissions of CO₂
8 amounted to 1,346 (± 31) g CO₂ kg⁻¹ of dry fuel burned or 342 (± 8) g CO₂-C. Other
9 carbonaceous emissions amounted to 218 g CO kg⁻¹; 8.35 g CH₄ kg⁻¹; 1.74 g C₂H₄ kg⁻¹; 1.53
10 g C₂H₆ kg⁻¹; and 0.60 g CH₃OH kg⁻¹ of dry fuel burned. Emissions of the nitrogenous
11 compounds amounted to 2.21 g HCN kg⁻¹; and 0.73 g NH₃ kg⁻¹.

12

13 **4 Discussion**

14 There is a very wide range in reported CO₂ emissions from both active and abandoned peat
15 extraction areas in the literature (Figure 7). Much of this variation can be attributed to
16 differences in climate, drainage level, peat type, peat extraction methods and the end use of
17 the peat and, as such, provides a useful framework to examine the variations in this study.

18 **4.1 Effects of climate**

19 While the study sites in this paper are all located within the temperate zone, considerable
20 variation in CO₂-C emissions was evident. Given that all the sites are drained to a similar
21 depth (Fig. 1), it is not surprising that the variation in emissions appeared to be controlled
22 largely by differences in soil temperatures between the sites (Fig. 6). The coldest site in terms
23 of mean soil temperatures and lowest in terms of annual emissions was Muirhead Moss (IP5)
24 in North-Eastern Scotland. Although rainfall and site water table levels were similar to the
25 other sites, soil temperatures at this site remained below 0°C for a high proportion (~14%) of
26 the year, and are likely to have resulted in a slowdown of extracellular enzymatic diffusion
27 (Davidson and Janssens, 2006), reduced microbial activity (Fenner et al., 2005) and
28 consequently lower rates of CO₂ production (Basiliko et al., 2007). Indeed, it is likely that our
29 value of 0.93 t CO₂-C ha⁻¹ yr⁻¹ at this site may be an overestimation given that it was
30 calculated from monthly mean values that were measured during day time hours (highest

1 daily temperatures). As much of the peatlands in Scotland fall within the same temperature
2 regime (Chapman and Thurlow, 1998), CO₂-C emissions data from a wider range of peat
3 extraction sites in this region might significantly refine our EF derivation.

4 At the other end of the spectrum, the highest emissions and soil temperatures were observed
5 at Turraun (IP4) in the Irish Midlands. Data from this site had been previously reported by
6 Wilson et al. (2007). In this study, we only utilised CO₂-C flux data from plots where the
7 mean annual water table position was deeper than -20cm. This resulted in a higher mean
8 value (taken over two years) in this current study. Three of IP sites in the ROI are located in
9 the Midlands where more “extremes” in climate are generally experienced (lower winter
10 temperatures, higher summer temperatures) than along the Western coast (IP3). However,
11 during this study, winter temperatures at all the ROI sites seldom decreased below 0°C (Fig.
12 3) and the proportion of hourly temperatures higher than 20°C were somewhat similar
13 between the sites. Although, Little Woolden Moss (IP6) received the lowest annual rainfall of
14 all sites in year 1 of the study at that site (Fig.1), mean annual soil temperatures were in the
15 mid-range of the 9 study sites, hourly T_{5cm} values were normally distributed (Fig.3) and CO₂-
16 C_{on site} emissions were close to the derived EF value of 1.70 t CO₂-C ha⁻¹ yr⁻¹ (Table 2).

17 The DP sites are all located in the ROI and within a 35km radius, but considerable variation
18 in annual rainfall was apparent during this study (Fig. 1), with DP3 (the furthest west)
19 receiving the highest rainfall of all sites in the study (on average 34% more rainfall than the
20 other DP sites). The east-west rainfall gradient in the ROI is well documented and coincides
21 with a change in peatland types (i.e. raised bogs to Atlantic blanket bogs). This climatic
22 variation is reflected in the annual R_{eco} values, which were similar between DP1 and DP2 but
23 much lower in DP3 (Fig. 5). There is an established relationship between rainfall amount and
24 the moisture content of peat (Price and Schlotzhauer, 1999; Strack and Price, 2009). For the
25 sites located in high rainfall areas, such as DP3, there may be a suppression of aerobic
26 microbial activity within the peat matrix, and as a consequence R_{eco} values may be lower than
27 would be expected for a drained peat soil. Indeed, at some of these sites, occult precipitation
28 (e.g. dew and fog droplets) may also contribute significantly to higher levels of soil moisture
29 (Lindsay et al., 2014). During the growing season, the transpiration process is also likely to
30 play a role in determining the moisture content of the peat within the rooting zone (~20cm
31 depth) at these vegetated sites. Moisture losses are likely to be accentuated on sunny days
32 when air and soil temperatures are high, when LAI values are highest (mid-summer) and
33 when vapour pressure deficit is not a limiting factor. As CO₂ emissions were closely

1 correlated to soil temperature at 5 cm depth, reduced moisture content in this zone is likely to
2 stimulate aerobic microbial activity. Annual GPP showed a similar trend to annual R_{eco} in
3 these vegetated DP sites. GPP is strongly controlled by the amount of light received by the
4 plants (i.e. PPFD levels and LAI) and the efficiency with which the plants use it. PPFD
5 values (data not shown) and the vegetation composition were broadly similar during the
6 sampling periods, which would seem to indicate that LAI is the driver of both productivity
7 and therefore NEE at these sites (Fig. 6). However, variations in LAI are likely to be the
8 result of subtle differences in a number of other variables (e.g. nutrient status, site
9 management) that were not captured in our measurements.

10 **4.2 Effects of drainage level**

11 While a close relationship between WT position and $\text{CO}_2\text{-C}$ emissions has been established in
12 some peatland studies (Silvola et al., 1996; Blodau and Moore, 2003; Blodau et al., 2004),
13 soil temperature proved to be the strongest determinant of $\text{CO}_2\text{-C}_{\text{on-site}}$ emissions at our sites
14 and this relationship has also been observed by other studies in peat extraction areas (e.g.
15 Shurpali et al., 2008; Mander et al., 2012; Salm et al., 2012). While the addition of WT or
16 VMC improved the performance of the R_{eco} models at some of the sites, the improvement
17 was only slight and this is likely due to the fairly narrow range of WT/VMC values recorded
18 over the course of the 12 month study (e.g. the range in VMC values at DP3 only ranged
19 between 56-64%). Therefore, optimum WT /VMC levels for respiration may not have been
20 encountered. The R_{eco} models used here are only valid for the data that was measured over
21 the course of the study at each site and cannot be readily extrapolated beyond the range of
22 that data. For those sites where water table did not appear to influence R_{eco} dynamics it may
23 be that fluctuations in WT level were missed with the interpolation approach and $\text{CO}_2\text{-C}$ flux
24 measurement regimes that we employed here, although these methodologies have been
25 widely used elsewhere (Riutta et al., 2007; Soini et al., 2010; Renou-Wilson et al., 2014).
26 Instead, it is probable that our results reflect the complexity of the relationship between R_{eco}
27 and WT in very dry soils as outlined by Lafleur et al.(2005), where factors such as a stable,
28 low surface soil moisture content, and decreased porosity (i.e. limited oxygen availability) at
29 the depths that the WT is mainly located, ensure that when $\text{CO}_2\text{-C}$ fluxes are measured, the
30 WT is deeper than the zone where it has a discernible impact on R_{eco} (Juszczak et al., 2013).
31 As such, the soil temperature regime in these sites may act as a “proxy” for drainage level

1 (i.e. higher soil temperatures are likely to occur in conjunction with deeper water table levels
2 and vice versa) (Mäkiranta et al., 2009).

3 **4.3 Peat characteristics**

4 Industrial peat extraction involves the removal of surface vegetation and results in the
5 exposure of decomposed peat at the surface. The level of decomposition in the peat is related
6 to depth and as extraction proceeds, the more highly decomposed peat is exposed. The peat in
7 industrial extraction sites tends to have a lower aerobic CO₂ production potential than natural
8 sites for example, due to differences in substrate and nutrient availability, a more extreme
9 physical environment (Glatzel et al., 2004) and reduced labile organic matter supply in the
10 absence of plant communities (i.e. priming). In our study, the C content (with the exception
11 of DP2) was similar across all sites (Table 1). Although, Glatzel et al. (2004) noted that CO₂
12 production was negatively correlated with the von Post scale of decomposition, no correlation
13 with annual CO₂-C emissions was evident in our study ($p>0.05$). Similarly, despite obvious
14 difference in nitrogen content and pH values between IP sites, no relationships with CO₂
15 fluxes were discerned. However, the residual peat at IP4 is strongly influenced by the close
16 proximity of limestone parent material, as evidenced by high pH values and the lowest C:N
17 ratio (Table 1), and is highly minerotrophic. Given the high CO₂-C emissions associated with
18 this site, consideration should be given to disaggregation by nutrient type should more data
19 become available in the future.

20 Organic matter quality has been closely linked to the soil respiration rate, with lower
21 emission rates associated with the poorer quality organic matter found at depth in drained
22 peatlands (Leifeld et al., 2012). The lowest emissions at our sites occurred where the residual
23 peat was either of Cyperaceous (IP3) or *Sphagnum* / Cyperaceous (IP5) origin. However,
24 while the slow decomposition rate of *Sphagnum* litter in comparison to other plant litter has
25 been well documented (Verhoeven and Toth, 1995; Bragazza et al., 2007), there is
26 insufficient data from our study sites to determine whether the limited relationship observed
27 here between peat type and CO₂-C emissions in our study sites is coincidental rather than
28 causal.

29 **4.4 Effects of peat extraction methods and peat end use**

30 For peat utilised for horticulture, the more fibrous peat layers nearer the surface are extracted.
31 This may result in the oxidation of more labile organic matter and may account for the very

1 high emissions associated with Canadian peatlands for example (Fig. 7) in comparison to
2 countries where the deeper peat layers are extracted (Mander et al., 2012). However, the IP
3 sites in this study are highly decomposed peat and have been abandoned for 30 years or more
4 in some cases (e.g. IP4) and have remained unvegetated. It is possible that CO₂-C emissions
5 from active extraction areas may be higher than those derived in this study given that over the
6 summer period the surface of the peat is regularly scarified and aerated. However, Salm et al.
7 (2012) reported higher emissions from abandoned areas in comparison to active areas,
8 although colonisation by vegetation in the former may have accentuated respiration losses.
9 High annual CO₂-C emissions following abandonment and recolonization have also been
10 reported by Strack and Zuback (2013) and are in close agreement with the R_{eco} values
11 reported here for the DP sites (Fig. 5).

12 We have estimated the contribution of heterotrophic respiration (R_H) to R_{eco} at 49%.
13 Although, this is based on measurements at a single site (DP1), it is within the range reported
14 by other studies (Frolking et al., 2002; Moore et al., 2002; Shurpali et al., 2008). The R_H
15 values measured at DP1 (Fig.5) and estimated at DP2 are higher than the R_{eco} values at the IP
16 sites, which would indicate that decomposition of the belowground biomass (following
17 clipping) and subsequent “priming” effects may contribute significantly to CO₂-C dynamics
18 at vegetated extraction sites. Furthermore, the methods employed to extract the peat at some
19 of the DP sites (the peat is extruded onto the surface of the peatland from narrow openings
20 made in the peat by a chain cutter) has led to the formation of deep fissures (ca. 4 cm wide
21 and > 2m deep) within the peat that may enhance oxidation throughout the peat profile.
22 Nonetheless, fissures (ca. 10 cm wide and > 1m deep) formed in the peat during climatically
23 dry years and that were partially filled in during wetter/windier years were also observed at
24 IP5 where the lowest annual emissions were observed.

25 **4.5 Fire emission factors**

26 The mean MCE reported here (0.837) is typical of smouldering combustion (e.g. Yokelson et
27 al., 1996; Bertschi et al., 2003) and comparable with the reported range of MCE in other
28 studies of high latitude peats (Yokelson et al., 1997; Stockwell et al., 2014). Emission factors
29 for CO₂ and CO are also typical of smouldering combustion and similar to those from other
30 peat studies, particularly Yokelson et al. (1997). As found in other studies of peat fire
31 emissions, our measurements confirm that the CH₄ EF for Irish peat is particularly high (8.35
32 g kg⁻¹ dry fuel burned) when compared with other forms of biomass burning. Given the high

1 Global Warming Potential, where each gram of emitted CH₄ is equivalent to 34 g of CO₂
2 (100 year time horizon, IPCC, 2013), the CH₄ emissions from Irish peat fires may account for
3 over 12% of the CO₂-equivalent emissions. This result emphasises the importance of
4 understanding the full suite of trace gas emissions from biomass burning, rather than
5 focussing solely on CO₂ and CH₄ emissions. In general, the other EFs reported here lie within
6 the range of variability observed by other peat burning studies, with the exception of NH₃,
7 which is particularly low, possibly as a result of the nitrogen-poor soils that are typical of
8 Irish and UK blanket bogs. Here, we also report the first C₂H₆ EF for peat ($1.53 \pm 0.17 \text{ g kg}^{-1}$
9 dry fuel burned), similar in magnitude to C₂H₆ emissions from boreal forests (1.77 g kg^{-1} dry
10 fuel burned) according to Akagi et al. (2011). However, the use of prescribed fire in the UK
11 to burn off old heather growth to encourage new growth (e.g. the muirburn practice) may not
12 impact the underlying peat to any great extent, given that the practice is restricted to the
13 October-April period when soil moisture conditions are highest. Emissions result from the
14 burning of the woody aboveground biomass, and the underlying peat is generally unaffected.
15 In contrast, wildfires typically occur during the summer months when temperatures are
16 highest and moisture levels are low, resulting in burning of both the vegetation and the peat
17 itself. Indeed, recent work by Kettridge et al. (2015) has highlighted the vulnerability of
18 drained peatlands, even at high latitudes, to increased risk of wildfire and subsequent
19 vegetation changes.

20 **4.6 Implications for National Inventory reporting**

21 The ROI currently employs the 2006 GPG default value of 0.2 t CO₂-C ha⁻¹ (nutrient poor) in
22 reporting of all peat extraction areas, and estimated emissions for 2012 (the most recent
23 assessment year) were 9,312 t CO₂-C yr⁻¹ (Table 4). In contrast, the approach in the UK has
24 been to differentiate between peat extracted for fuel and horticulture and then applying the
25 default EFs for nutrient rich (1.1 t CO₂-C ha⁻¹) and nutrient poor peat (0.2 t CO₂-C ha⁻¹)
26 respectively. For 2012, CO₂-C emissions from UK extraction peatlands were estimated at
27 2,118 t CO₂-C yr⁻¹ (Table 4).

28 Reported annual emissions are likely to increase considerably if the Tier 1 values in the IPCC
29 Wetlands Supplement are adopted by inventory compilers. We estimate that emissions from
30 peatlands managed for extraction will be approximately 16 and 10 times higher for the ROI
31 and UK respectively (Table 4). The EFs derived in this study for CO₂-C_{on site} for both
32 industrial and domestic peatlands (Table 2) are considerably lower than the Tier 1 value of

1 2.8 tonnes CO₂-C ha⁻¹ yr⁻¹ provided in the IPCC Wetlands Supplement (2014). Although the
2 EFs derived in this study fall within the lower confidence margin of the Tier 1 range, our new
3 EFs have a marked reduction in associated uncertainty. As the Tier 1 is a generic value based
4 on published literature rather than a targeted measurement programme, it is naturally subject
5 to a certain level of bias, which result when the underlying studies are not representative of
6 management practices, climatic zones, or soil types in a particular region (Ogle et al., 2004),
7 and may lead to either an over- or underestimation of CO₂-C emissions. Given that no
8 significant difference exists between the EFs derived for the IP and DP sites in this study, we
9 propose a single EF for CO₂-C_{on-site} of 1.68 t CO₂-C ha⁻¹ yr⁻¹ to be applied to peatlands
10 managed for extraction in the ROI and UK regardless of peat type. This EF value could be
11 further disaggregated by regional climate, domestic peat extraction intensity (based on
12 extraction rates) or by end use of the peat (horticulture or energy) if more data becomes
13 available. For the latter, it would be highly useful to determine quantitatively whether CO₂-
14 C_{on-site} emissions vary between the less decomposed residual peat utilised for horticulture and
15 the more decomposed residual peat used for energy production. As the EFs derived in this
16 study have come from sites located within the same “climatic” region, we feel that they are
17 more appropriate for the ROI and the UK inventory purposes than either the 2006 GPG or the
18 2013 Wetlands Supplement. If the CO₂-C_{on site} EFs derived from this study are used in annual
19 NIRs, we estimate that annual emissions would be 9.5 and 6 times higher for the ROI and UK
20 respectively, in comparison to the emissions calculated with the 2006 GPG Tier 1 value, and
21 40% lower than emissions calculated with the Wetlands Supplement EF.

22 As reported CO₂-C_{on-site} emissions are henceforth likely to be much higher for any country
23 that moves from the 2006 GPG to the 2013 Wetlands Supplement, some consideration of
24 potential mitigation measures is required. Wetland Drainage and Rewetting is a new elective
25 activity under Article 3.4 of the Kyoto Protocol (second commitment period) and applies to
26 all lands that have been drained since 1990 and to all lands that have been rewetted since
27 1990. Countries that elect to report under this activity will also be able to claim C benefits
28 from the rewetting of drained peatlands. In theory, this should provide an impetus for the
29 rewetting of high emitting land use categories such as peatlands managed for extraction,
30 particularly as these areas will remain persistent long term emission hotspots in the absence
31 of rewetting actions (Waddington et al., 2002).

32 **4.7 Information gaps**

1 Greenhouse gas emissions from peatlands used for extraction are composed of (a) on-site
2 emissions (i.e. from peat extraction areas, ditches and stockpiles) and (b) off-site emissions
3 associated with water borne losses and the use of the peat for energy or horticulture. In this
4 paper, we have focused solely on the on-site CO₂-C emissions from the peat extraction areas,
5 and GHG emissions from fire. However, C losses from other pathways may also be
6 substantial. Research has shown that GHG emissions from on-site peat stockpiles and ditches
7 are considerable (Alm et al., 2007a and references therein). Currently, emissions data from
8 stockpiles in the temperate zone are not available and the IPCC Wetlands Supplement does
9 not provide a Tier 1 value, and instead encourages countries to move to higher Tiers in terms
10 of reporting (IPCC, 2014). However, countries such as Finland have developed a Tier 2
11 approach in which EFs (incl. CH₄ and N₂O) depend on regional weather and in which
12 emissions from ditches and stockpiles are taken into account (Alm et al., 2007a;
13 Lapveteläinen et al., 2007). The IPCC Wetlands Supplement provides Tier 1 EFs for CH₄
14 emissions from both peat extraction areas and from ditches. The value for the latter is
15 particularly high (542 kg CH₄ ha⁻¹ yr⁻¹, expressed per unit area of ditch surface) and indicates
16 the importance of this pathway in the full GHG balance (Evans et al., 2015). Similarly, N₂O
17 emissions have been shown to be significant from drained peatlands (Regina et al., 1996) yet
18 despite this, there are only a small number of published studies and more research is critical
19 in order to provide regional specific EFs. While CH₄ and N₂O fluxes have been quantified at
20 some of the sites, the data is currently being processed with a view to publication in the
21 future. In terms of the fire study, N₂O is a difficult gas to measure using the FTIR setup
22 employed in this study, as it can only be determined from spectra with very large
23 enhancements of trace gases. This is because the N₂O absorption occurs in a similar wave
24 number region to both the CO₂ and CO absorption bands (Paton-Walsh et al., 2014). Paton-
25 Walsh et al. (2014) could only determine N₂O from two of their five open fires, whilst Smith
26 et al. (2014), who used a similar setup, failed to determine N₂O from any of their 21 fires
27 studied. In our study, we found that excess mole fractions of N₂O could not be correlated to
28 either CO₂ or CO for the determination of emission ratios, precluding the calculation of EFs.
29 One explanation for this is that N₂O is predominantly a product of flaming combustion and is
30 strongly correlated to CO₂ (Paton-Walsh et al., 2014). The lack of flaming combustion in our
31 peat burns probably explains our inability to detect significant excess N₂O mole fractions.

32 Other pathways may be of equal importance. For example, the loss of POC from bare peat
33 surfaces may be considerable where the surface is exposed and subject to wind or water

1 erosion (Evans et al., 2006; Lindsay, 2010). While some of the windborne POC is likely to be
2 deposited within the extraction field itself, a proportion undoubtedly leaves the peatland,
3 although there are currently few data available to quantify losses from either wind or water
4 erosion, or the extent to which POC is converted to CO₂ (IPCC, 2014). In addition, high
5 losses of DOC from drained peatlands have been reported (Evans et al., 2015 and references
6 therein). Although a Tier 1 EF value for DOC is provided in the IPCC Wetlands Supplement,
7 disaggregated by climate zone, with the assumption that 90% of the exported DOC is
8 converted to CO₂, there is an obvious need to quantify these losses on a regional basis given
9 the high precipitation loads experienced by the ROI and the UK, and associated differences in
10 peat type (Evans et al., 2015). Emissions from burning are not currently reported in either the
11 ROI or UK inventory reports. The EF provided in the IPCC Wetlands Supplement for CO₂
12 emissions associated with wildfire burning is similar to our value here (Table 3).
13 Furthermore, given the high CH₄ emissions associated with the burning of the peat that we
14 have reported here (Table 3), and taking cognisance of the strong GWP of CH₄, more
15 research is urgently required to quantify this emission pathway, particularly under field
16 conditions.

17 The provision of activity data for inventory reporting varies between the ROI and the UK,
18 with the peat extraction industry the source of data in the former (Duffy et al., 2014), and a
19 multi-source approach (Directory of Mines and Quarries point locations with Google Earth
20 imagery, scientific reports/papers) used in the latter (Webb et al., 2014). However, CO₂
21 emissions from domestic peat extraction in the ROI are not currently reported due to a lack of
22 activity data and could potentially be very high (Wilson et al., 2013b). In the UK, areas under
23 domestic extraction are included in the Grassland category but may be moved as the UK
24 considers changes post-Wetlands Supplement. Determining to what degree that peatlands
25 have been affected by domestic peat extraction and how far those impacts extend into the
26 main peatland area are obvious challenges facing future research. The use of remote sensing
27 platforms could provide high resolution data that will be able to differentiate between
28 domestic peat extraction and other types of disturbed peatlands. In particular, the use of
29 Unmanned Aerial Vehicles (i.e. drones), which have been used to map individual peatlands
30 at a very high resolution (e.g. Knoth et al., 2013) offer considerable potential for more
31 detailed mapping of domestic peatlands at the national scale.

32

33 **5 Conclusion**

1 Peatlands managed for extraction are a substantial CO₂-C emissions hotspot at the landscape
2 scale and further contribute to climate change through significant GHG emissions when the
3 peat is burned or utilised in horticulture. This study, which measured and modelled emissions
4 from a range of sites across the ROI and the UK, has highlighted the importance of
5 generating robust Tier 2 values for different regions and land-use categories. Given that the
6 IPCC Tier 1 EF was only based on 20 sites (all from Canada/Fenno-Scandia) we suggest that
7 data from another 9 sites significantly expands the global dataset, as well as adding a new
8 region.

9

10 **Acknowledgements**

11 Funding to DW and FRW by the Environment Protection Agency (Ireland) and by Bord na
12 Móna is acknowledged. Thanks to Ignatius Kelly, Bord na Móna for rainfall data. Funding to
13 SDD and EA (Site IP6) was provided by the Department for Environment, Food and Rural
14 Affairs (UK) under research grant number SP1210. Thanks to Chris Millar, Lancashire
15 Wildlife Trust and Phil Jepson for access to IP6, and to Cat Moody for fieldwork assistance.
16 Experimental work at IP5 was carried out as part of the RECIPE project, with funding
17 through the European Union Fifth Framework Programme. We would like to thank Dr
18 Christopher MacLellan and Dr Alasdair MacArthur of NERC FSF for their wide ranging
19 support, including the loan of the FTIR spectrometer and infrared White cell used here. We
20 thank Dr Bruce Main and Bill Luckhurst of King's College London for their technical
21 support at the laboratory burns. Grateful thanks to Eeva-Stiina Tuittila and Anna Laine for
22 useful modelling discussions. We thank the two anonymous reviewers for their insightful
23 comments and suggestions on the earlier draft of the manuscript.

24

25 **References**

- 26 Akagi, S. K., Yokelson, R. J., Wiedinmyer, C., Alvarado, M. J., Reid, J. S., Karl, T., Crouse,
27 J. D., and Wennberg, P. O.: Emission factors for open and domestic biomass burning for
28 use in atmospheric models, *Atmos. Chem. Phys.*, 11, 4039-4072, 10.5194/acp-11-4039-
29 2011, 2011.
- 30 Alm, J., Shurpali, N. J., Minkinen, K., Aro, L., Hytönen, J., Laurila, T., Lohila, A.,
31 Maljanen, M., Martikainen, P. J., Mäkiranta, P., Penttilä, T., Saarnio, S., Silvan, N.,
32 Tuittila, E.-S., and Laine, J.: Emission factors and their uncertainty for the exchange of

- 1 CO₂, CH₄ and N₂O in Finnish managed peatlands., *Boreal Environment Research*, 12,
2 191-209, 2007a.
- 3 Alm, J., Shurpali, N. J., Tuittila, E.-S., Laurila, T., Maljanen, M., Saarnio, S., and Minkkinen,
4 K.: Methods for determining emission factors for the use of peat and peatlands - flux
5 measurements and modelling, *Boreal Environment Research*, 12, 85-100, 2007b.
- 6 Basiliko, N., Blodau, C., Roehm, C., Bengtson, P., and Moore, T.: Regulation of
7 decomposition and methane dynamics across natural, commercially mined, and restored
8 northern peatlands, *Ecosystems*, 10, 1148-1165, 10.1007/s10021-007-9083-2, 2007.
- 9 Bertschi, I., Yokelson, R. J., Ward, D. E., Babbitt, R. E., Susott, R. A., Goode, J. G., and Hao,
10 W. M.: Trace gas and particle emissions from fires in large diameter and belowground
11 biomass fuels, *Journal of Geophysical Research: Atmospheres*, 108,
12 10.1029/2002jd002100, 2003.
- 13 Blodau, C., and Moore, T. R.: Experimental response of peatland carbon dynamics to a water
14 table fluctuation., *Aquatic Sciences*, 65, 47-62, 2003.
- 15 Blodau, C., Basiliko, N., and Moore, T.: Carbon turnover in peatland mesocosms exposed to
16 different water table levels., *Biogeochemistry*, 67, 331-351, 2004.
- 17 Bragazza, L., Siffi, C., Iacumin, P., and Gerdol, R.: Mass loss and nutrient release during
18 litter decay in peatland: The role of microbial adaptability to litter chemistry, *Soil Biology
19 & Biochemistry*, 39, 257-267, 2007.
- 20 Cancellieri, D., Leroy-Cancellieri, V., Leoni, E., Simeoni, A., Kuzin, A. Y., Filkov, A. I., and
21 Rein, G.: Kinetic investigation on the smouldering combustion of boreal peat, *Fuel*, 93,
22 479-485, <http://dx.doi.org/10.1016/j.fuel.2011.09.052>, 2012.
- 23 Chapman, S., Buttler, A., Francez, A.-J., Laggoun - Défarge, F., Vasander, H., Schloter, M.,
24 Combe, J., Grosvernier, P., Harms, H., Epron, D., Gilbert, D., and Mitchell, E.:
25 Exploitation of northern peatlands and biodiversity maintenance: a conflict between
26 economy and ecology., *Front Ecol Environ*, 1, 525-532, 2003.
- 27 Chapman, S. J., and Thurlow, M.: Peat respiration at low temperatures., *Soil Biology and
28 Biochemistry*, 30, 1013-1021, 1998.
- 29 Cleary, J., Roulet, N. T., and Moore, T. R.: Greenhouse gas emissions from Canadian peat
30 extraction, 1990-2000: a life cycle analysis., *Ambio*, 34, 456-461, 2005.
- 31 Couwenberg, J., and Fritze, C.: Towards developing IPCC methane 'emission factors' for
32 peatlands (organic soils) available at: [http://www.mires-and-
33 peat.net/pages/volumes/map10/map1003.php](http://www.mires-and-peat.net/pages/volumes/map10/map1003.php) (last access: 18 May 2015), *Mires and Peat*,
34 10, 1-17, 2012.
- 35
- 36 Davidson, E., and Janssens, I. A.: Temperature sensitivity of soil carbon decomposition and
37 feedbacks to climate change., *Nature*, 440, 165-173, 2006.
- 38 Department of Environment Food and Rural Affairs: The Natural Choice: securing the value
39 of nature, www.official-documents.gov.uk, (last access: 18 May 2015), 2011.

- 1 Dise, N. B.: Peatland response to global change, *Science*, 326, 810-811, 2009.
- 2 Drösler, M.: Trace gas exchange and climatic relevance of bog ecosystems, southern
3 Germany., Department für Ökologie, Universität München, Munich, Germany , 182 pp.,
4 2005.
- 5 Duffy, P., Hanley, E., Hyde, B., O'Brien, P., Ponzi, J., Cotter, E., and Black, K.: National
6 Inventory Report 2014. Greenhouse gas emissions 1990–2012 reported to the United
7 Nations Framework Convention on Climate Change, 469, Environmental Protection
8 Agency (Ireland), Co. Wexford, Ireland, 2014.
- 9
- 10 Elsgaard, L., Gorres, C.-M., Hoffmann, C. C., Blicher-Mathiesen, G., Schelde, K., and
11 Petersen, S. O.: Net ecosystem exchange of CO₂ and carbon balance for eight temperate
12 organic soils under agricultural management, *Agriculture, Ecosystems & Environment*,
13 162, 52-67, 2012.
- 14 Evans, C., Renou-Wilson, F., and Strack, M.: The role of waterborne carbon in the
15 greenhouse gas balance of drained and re-wetted peatlands, *Aquatic Sciences*, In press,
16 2015.
- 17 Evans, M., Warburton, J., and Yang, J.: Eroding blanket peat catchments: Global and local
18 implications of upland organic sediment budgets., *Geomorphology*, 79, 45-57, 2006.
- 19 Farrell, C. A., and Doyle, G. J.: Rehabilitation of industrial cutaway Atlantic blanket bog in
20 County Mayo, North-West Ireland, *Wetlands Ecology and Management*, 11, 21-35, 2003.
- 21 Fenner, N., Freeman, C., and Reynolds, B.: Observations of a seasonally shifting thermal
22 optimum in peatland carbon-cycling processes; implications for the global carbon cycle
23 and soil enzyme methodologies *Soil Biology and Biochemistry*, 37, 1814-1821, 2005.
- 24 Frandsen, W. H.: Ignition probability of organic soils, *Canadian Journal of Forest Research*,
25 27, 1471-1477, 10.1139/x97-106, 1997.
- 26 Frolking, S., Roulet, N. T., Moore, T. R., Lafleur, P. M., Bubier, J. L., and Crill, P. M.:
27 Modeling seasonal to annual carbon balance of Mer Bleue Bog, Ontario, Canada, *Global
28 Biogeochemical Cycles*, 16, 4-1-4-21, 10.1029/2001gb001457, 2002.
- 29 Gažovič, M., Forbrich, I., Jager, D. F., Kutzbach, L., Wille, C., and Wilmking, M.:
30 Hydrology-driven ecosystem respiration determines the carbon balance of a boreal
31 peatland, *Science of the Total Environment*, 463–464, 675-682,
32 <http://dx.doi.org/10.1016/j.scitotenv.2013.06.077>, 2013.
- 33 Glatzel, S., Kalbitz, K., Dalva, M., and Moore, T.: Dissolved organic matter properties and
34 their relationship to carbon dioxide efflux from restored peat bogs., *Geoderma*, 113, 397-
35 411, 2003.
- 36 Glatzel, S., Basiliko, N., and Moore, T.: Carbon dioxide and methane production potentials of
37 peats from natural, harvested and restored sites, eastern Québec, Canada, *Wetlands.*, 24,
38 261-267, 2004.
- 39 Griffith, D. W. T.: Synthetic calibration and quantitative analysis of gas-phase FT-IR spectra,
40 *Appl. Spectrosc.*, 50, 59-70, 1996.

- 1 Haddaway, N. R., Burden, A., Evans, C. D., Healey, J. R., Jones, D. L., Dalrymple, S. E.,
2 and Pullin, A. S.: Evaluating effects of land management on greenhouse gas fluxes and
3 carbon balances in boreo-temperate lowland peatland systems, available at:
4 <http://www.environmentalevidencejournal.org/content/3/1/5> (last access: 18 May 2015),
5 *Environmental Evidence*, 3, 30, doi:10.1186/2047-2382-3-5, 2014.
6
- 7 Howley, M., Dennehy, E., O'Gallachoir, B., and Holland, M.: *Energy in Ireland 1990-2011*,
8 Sustainable Energy Authority of Ireland, Dublin, 92, 2012.
- 9 IPCC: *IPCC Guidelines for National Greenhouse Gas Inventories*, Prepared by the National
10 Greenhouse Gas Inventories programme, IGES, Published by the Institute for Global
11 Environmental Strategies (IGES), Hayama, Japan, 2006.
12
- 13 IPCC: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I*
14 *to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*,
15 Cambridge, United Kingdom and New York, NY, USA, 1535, 2013.
- 16 IPCC: *2013 Supplement to the 2006 IPCC Guidelines for National Greenhouse Gas*
17 *Inventories: Wetlands*, edited by: Hiraishi, T., Krug, T., Tanabe, K., Srivastava, N.,
18 Baasansuren, J., Fukuda, M., and Troxler, T. G., IPCC, Switzerland, 2014.
- 19 Järveoja, J., Laht, J., Maddison, M., Soosaar, K., Ostonen, I., and Mander, Ü.: Mitigation of
20 greenhouse gas emissions from an abandoned Baltic peat extraction area by growing reed
21 canary grass: life-cycle assessment, *Reg. Environ. Change*, 13, 781–795,
22 doi:10.1007/s10113-012-0355-9, 2012.
23
- 24 Juszczak, R., Humphreys, E., Acosta, M., Michalak-Galczevska, M., Kayzer, D., and
25 Olejnik, J.: Ecosystem respiration in a heterogeneous temperate peatland and its sensitivity
26 to peat temperature and water table depth, *Plant and Soil*, 366, 505-520, 10.1007/s11104-
27 012-1441-y, 2013.
- 28 Kettridge, N., Turetsky, M. R., Sherwood, J. H., Thompson, D. K., Miller, C. A., Benscoter,
29 B. W., Flannigan, M. D., Wotton, B. M., and Waddington, J. M.: Moderate drop in
30 water table increases peatland vulnerability to post-fire regime shift, *Sci. Rep.*, 5, 8063,
31 doi:10.1038/srep08063, 2015.
- 32 Knoth, C., Klein, B., Prinz, T., and Kleinebecker, T.: Unmanned aerial vehicles as innovative
33 remote sensing platforms for high-resolution infrared imagery to support restoration
34 monitoring in cut-over bogs, *Applied Vegetation Science*, 16, 509-517,
35 10.1111/avsc.12024, 2013.
- 36 Koehler, A.-K., Sottocornola, M., and Kiely, G.: How strong is the current carbon
37 sequestration of an Atlantic blanket bog?, *Global Change Biology*, 17, 309-319, 2011.
- 38 Lafleur, P. M., Moore, T. R., Roulet, N. T., and Frohling, S.: Ecosystem respiration in a cool
39 temperate bog depends on peat temperature but not water table., *Ecosystems*, 8, 619-629,
40 2005.
- 41 Laine, A., Riutta, T., Juutinen, S., Väiliranta, M., and Tuittila, E.-S.: Acknowledging the
42 spatial heterogeneity in modelling / reconstructing carbon dioxide exchange in a northern
43 aapa mire, *Ecological Modelling*, 220, 2646-2655, 2009.

- 1 Lapveteläinen, T., Regina, K., and Perälä, P.: Peat-based emissions in Finland's national
2 greenhouse gas inventory, *Boreal Environment Research*, 12, 225-236, 2007.
- 3 Leifeld, J., Steffens, M., and Galego-Sala, A.: Sensitivity of peatland carbon loss to organic
4 matter quality, *Geophys. Res. Lett.*, 39, L14704, doi:10.1029/2012GL051856, 2012.
- 5 Lindsay, R.: *Peatbogs and Carbon - A critical synthesis*, University of East London and
6 RSPB Scotland, London, 2010.
- 7 Lloyd, J., and Taylor, J. A.: On the temperature dependence of soil respiration., *Functional*
8 *Ecology*, 8, 315-323, 1994.
- 9 Lobert, J. M., Scharffe, D. H., Hao, W. M., Kuhlbusch, T. A., Seuwen, R., Warneck, P., and
10 Crutzen, P. J.: Experimental evaluation of biomass burning emissions: Nitrogen and
11 carbon containing compounds., in: *Global Biomass Burning: Atmospheric, Climatic and*
12 *Biospheric Implications*, edited by: Levine, J. S., MIT Press, Cambridge, Massachusetts,
13 USA, 1991.
- 14 Mäkiranta, P., Laiho, R., Fritze, H., Hytönen, J., Laine, J., and Minkkinen, K.: Indirect
15 regulation of heterotrophic peat soil respiration by water level via microbial community
16 structure and temperature sensitivity, *Soil Biology and Biochemistry*, 41, 695-703,
17 <http://dx.doi.org/10.1016/j.soilbio.2009.01.004>, 2009.
- 18 Mander, Ü., Järveoja, J., Maddison, M., Soosaar, K., Aavola, R., Ostonen, I., and Salm, J.-O.:
19 Reed canary grass cultivation mitigates greenhouse gas emissions from abandoned peat
20 extraction areas, *GCB Bioenergy*, 4, 462-474, 10.1111/j.1757-1707.2011.01138.x, 2012.
- 21 McNeil, P., and Waddington, J. M.: Moisture controls on *Sphagnum* growth and CO₂
22 exchange on a cutover bog., *Journal of Applied Ecology*, 40, 354-367, 2003.
- 23 Moore, T. R., Bubier, J. L., Froking, S. E., Lafleur, P. M., and Roulet, N. T.: Plant biomass
24 and production and CO₂ exchange in an ombrotrophic bog, *Journal of Ecology*, 90, 25-26,
25 2002.
- 26 Nilsson, M., Sagerfors, J., Buffam, I., Laudon, H., Eriksson, T., Grelle, A., Klemetsson, L.,
27 Weslien, P. E. R., and Lindroth, A.: Contemporary carbon accumulation in a boreal
28 oligotrophic minerogenic mire – a significant sink after accounting for all C-fluxes, *Global*
29 *Change Biology*, 14, 2317-2332, 10.1111/j.1365-2486.2008.01654.x, 2008.
- 30 Ogle, S. M., Conant, R. T., and Paustian, K.: Deriving grassland management factors for a
31 carbon accounting method developed by the Intergovernmental Panel on Climate Change,
32 *Environmental Management*, 33, 474-484, 2004.
- 33 Paton-Walsh, C., Smith, T. E. L., Young, E. L., Griffith, D. W. T., and Guérette, É.-A.: New
34 emission factors for Australian vegetation fires measured using open-path Fourier
35 transform infrared spectroscopy – Part 1: Methods and Australian temperate forest fires,
36 *Atmos. Chem. Phys.*, 14, 11313–11333, doi:10.5194/acp-14-11313-2014, 2014.
37
- 38 Price, J. S., and Schlotzhauer, S. M.: Importance of shrinkage and compression in
39 determining water storage changes in peat: the case of a mined peatland, *Hydrological*
40 *Processes*, 13, 2591-2601, 1999.

- 1 Regina, K., Nykänen, H., Silvola, J., and Martikainen, P.: Fluxes of nitrous oxide from boreal
2 peatlands as affected by peatland type, water table level and nitrification capacity,
3 *Biogeochemistry*, 35, 401-418, 10.1007/bf02183033, 1996.
- 4 Rein, G., Cohen, S., and Simeoni, A.: Carbon emissions from smouldering peat in shallow
5 and strong fronts, *Proceedings of the Combustion Institute* 32 2489-2496, 2009.
- 6 Renou-Wilson, F., Barry, C., Müller, C., and Wilson, D.: The impacts of drainage, nutrient
7 status and management practice on the full carbon balance of grasslands on organic soils
8 in a maritime temperate zone, *Biogeosciences*, 11, 4361-4379, doi:10.5194/bg-11-4361-
9 2014, 2014.
- 10 Renou, F., Egan, T., and Wilson, D.: Tomorrow's landscapes: studies in the after-uses of
11 industrial cutaway peatlands in Ireland, *Suo*, 57, 97-107, 2006.
- 12 Riutta, T., Laine, J., and Tuittila, E.-S.: Sensitivity of CO₂ exchange of fen ecosystem
13 components to water level variation, *Ecosystems*, 10, 718-733, 10.1007/s10021-007-9046-
14 7, 2007.
- 15 Roulet, N. T., Lafleur, P. M., Richard, P. J. H., Moore, T., Humphreys, E. R., and Bubier, J.:
16 Contemporary carbon balance and late Holocene carbon accumulation in a northern
17 peatland, *Global Change Biol.*, 13, 397-411, doi:10.1111/j.1365-2486.2006.01292.x,
18 2007.
- 19
- 20 Saarnio, S., Morero, M., Shurpali, N. J., Tuittila, E.-S., Mäkilä, M., and Alm, J.: Annual CO₂
21 and CH₄ fluxes of pristine boreal mires as a background for the lifecycle analyses of peat
22 energy, *Boreal Environment Research*, 12, 101-113, 2007.
- 23 Salm, J.-O., Maddison, M., Tammik, S., Soosaar, K., Truu, J., and Mander, Ü.: Emissions of
24 CO₂, CH₄ and N₂O from undisturbed, drained and mined peatlands in Estonia,
25 *Hydrobiologia*, 692, 41-55, 10.1007/s10750-011-0934-7, 2012.
- 26 Schouten, M. G. C.: Conservation and restoration of raised bogs: geological, hydrological
27 and ecological studies, Department of the Environment and Local Government /
28 Staatsbosbeheer, Dublin, Ireland, 2002.
- 29
- 30 Shurpali, N. J., Hyvönen, N., Huttunen, J. T., Biasi, C., Nykanen, H., Pekkarinen, N., and
31 Martikainen, P. J.: Bare soil and reed canary grass ecosystem respiration in peat extraction
32 sites in Eastern Finland, *Tellus*, 60B, 200-209, 2008.
- 33 Silvola, J., Alm, J., Ahlholm, U., Nykänen, H., and Martikainen, P. J.: CO₂ fluxes from peat
34 in boreal mires under varying temperature and moisture conditions, *Journal of Ecology*,
35 84, 219-228, 1996.
- 36 Smith, T. E. L., Paton-Walsh, C., Meyer, C. P., Cook, G. D., Maier, S. W., Russell-Smith, J.,
37 Wooster, M. J., and Yates, C. P.: New emission factors for Australian vegetation fires
38 measured using open-path Fourier transform infrared spectroscopy – Part 2: Australian
39 tropical savanna fires, *Atmos. Chem. Phys.*, 14, 11335-11352, 10.5194/acp-14-11335-
40 2014, 2014.

- 1 Soini, P., Riutta, T., Yli-Petäys, M., and Vasander, H.: Comparison of vegetation and CO₂
2 dynamics between a restored cut-way peatland and a pristine fen: evaluation of the
3 restoration success, *Restoration Ecology*, 18, 894-903, 2010.
- 4 Stockwell, C. E., Yokelson, R. J., Kreidenweis, S. M., Robinson, A. L., DeMott, P. J.,
5 Sullivan, R. C., Reardon, J., Ryan, K. C., Griffith, D. W. T., and Stevens, L.: Trace gas
6 emissions from combustion of peat, crop residue, domestic biofuels, grasses, and other
7 fuels: configuration and Fourier transform infrared (FTIR) component of the fourth Fire
8 Lab at Missoula Experiment (FLAME-4), *Atmos. Chem. Phys.*, 14, 9727-9754,
9 10.5194/acp-14-9727-2014, 2014.
- 10 Strack, M., Waddington, J. A., Bourbonniere, R. A., Buckton, E. L., Shaw, K., Whittington,
11 P., and Price, J. S.: Effect of water table drawdown on peatland dissolved organic carbon
12 export and dynamics, *Hydrological Processes*, 22, 3373-3385, DOI: 3310.1002/hyp.6931,
13 2008.
- 14 Strack, M., and Price, J. S.: Moisture controls on carbon dioxide dynamics of peat-Sphagnum
15 monoliths, *Ecohydrology*, 2, 34-41, 10.1002/eco.36, 2009.
- 16 Strack, M., and Zuback, Y. C. A.: Annual carbon balance of a peatland 10 yr following
17 restoration, *Biogeosciences* 10, 2885-2896, 2013.
- 18 Strack, M., Keith, A. M., and Zu, B.: Growing season carbon dioxide and methane exchange
19 at a restored peatland on the Western Boreal Plain, *Ecological Engineering*, 64, 231-239,
20 2014.
- 21 Sundh, I., Nilsson, M., Mikkela, C., Granberg, G., and Svensson, B. H.: Fluxes of methane
22 and carbon dioxide on peat-mining areas in Sweden, *Ambio*, 29, 499-503, 2000.
- 23 Tomlinson, R. W.: Changes in the extent of peat extraction in Northern Ireland 1990-2008 and
24 associated changes in carbon loss, *Applied Geography*, 30, 294-301, 2010.
- 25 Tuittila, E.-S., Komulainen, V.-M., Vasander, H., and Laine, J.: Restored cut-away peatland
26 as a sink for atmospheric CO₂, *Oecologia*, 120, 563 - 574, 1999.
- 27 Tuittila, E.-S., Vasander, H., and Laine, J.: Sensitivity of carbon sequestration in reintroduced
28 *Sphagnum* to water-level variation in a cutaway peatland, *Restoration Ecology*, 12, 482-
29 492, 2004.
- 30 Tuittila, E. S., and Komulainen, V. M.: Vegetation and CO₂ balance in an abandoned
31 harvested peatland in Aitoneva, southern Finland, *Suo*, 46, 69-80, 1995.
- 32 Turetsky, M. R., Kotowska, A., Bubier, J., Dise, N. B., Crill, P., Hornibrook, E. R. C.,
33 Minkinen, K., Moore, T. R., Myers-Smith, I. H., Nykänen, H., Olefeldt, D., Rinne, J.,
34 Saarnio, S., Shurpali, N., Tuittila, E.-S., Waddington, J. M., White, J. R., Wickland, K. P.,
35 and Wilmsking, M.: A synthesis of methane emissions from 71 northern, temperate, and
36 subtropical wetlands, *Glob. Change Biol.*, 20, 2183-2197, doi:10.1111/gcb.12580, 2014.
- 37 Turetsky, M. R., Benscoter, B., Page, S., Rein, G., van der Werf, G. R., and Watts, A.: Global
38 vulnerability of peatlands to fire and carbon loss, *Nature Geoscience*, 8, 11-14,
39 10.1038/ngeo2325.[http://www.nature.com/ngeo/journal/v8/n1/abs/ngeo2325.html#supple](http://www.nature.com/ngeo/journal/v8/n1/abs/ngeo2325.html#supplementary-information)
40 [mentary-information](http://www.nature.com/ngeo/journal/v8/n1/abs/ngeo2325.html#supplementary-information), 2015.

- 1 Verhoeven, J. T. A., and Toth, E.: Decomposition of *Carex* and *Sphagnum* litter in fens:
2 Effect of litter quality and inhibition by living tissue homogenates, *Soil Biology and*
3 *Biochemistry*, 27, 271-275, [http://dx.doi.org/10.1016/0038-0717\(94\)00183-2](http://dx.doi.org/10.1016/0038-0717(94)00183-2), 1995.
- 4 Waddington, J. M., and Price, J. S.: Effect of peatland drainage, harvesting and restoration on
5 atmospheric water and carbon exchange, *Physical Geography*, 21, 433-451, 2000.
- 6 Waddington, J. M., Warner, K. D., and Kennedy, G. W.: Cutover peatlands: a persistent
7 source of atmospheric CO₂, *Global Biogeochemical Cycles*, 16, 21-27, 2002.
- 8 Waddington, J. M., Strack, M., and Greenwood, M. J.: Toward restoring the net carbon sink
9 function of degraded peatlands: Short-term response in CO₂ exchange to ecosystem-scale
10 restoration, *J. Geophys. Res.*, 115, G01008, doi:10.1029/2009JG001090, 2010.
- 11 Webb, N., Broomfield, M., Brown, P., Buys, G., Cardenas, L., Murrells, T., Pang, Y.,
12 Passant, N., Thistlewaite, G., and Watterson, J.: UK Greenhouse Gas Inventory, 1990 to
13 2012. Annual Report for Submission under the Framework Convention on Climate
14 Change, Compiled on Behalf of the UK Department of Energy and Climate Change
15 (DECC) Science Division by Ricardo-AEA, 594, Ricardo-AEA, Oxfordshire, UK, 2014.
16
- 17 Wilson, D., Tuittila, E.-S., Alm, J., Laine, J., Farrell, E. P., and Byrne, K. A.: Carbon dioxide
18 dynamics of a restored maritime peatland, *Ecoscience*, 14, 71-80, 2007.
- 19 Wilson, D., Farrell, C., A., Müller, C., Hepp, S., and Renou-Wilson, F.: Rewetted industrial
20 cutaway peatlands in western Ireland: prime location for climate change mitigation?,
21 available at: <http://mires-and-peat.net/pages/volumes/map11/map1101.php> (last access: 18
22 May 2015), *Mires and Peat*, 11, 1–22, 2013a.
23
- 24 Wilson, D., Müller, C., and Renou-Wilson, F.: Carbon emissions and removals from Irish
25 peatlands: current trends and future mitigation measures, *Irish Geography*, 46, 1-23,
26 2013b.
- 27 Yokelson, R. J., Griffith, D. W. T., and Ward, D. E.: Open-path Fourier transform infrared
28 studies of large-scale laboratory biomass fires, *Journal of Geophysical Research:*
29 *Atmospheres*, 101, 21067-21080, 10.1029/96jd01800, 1996.
- 30 Yokelson, R. J., Susott, R., Ward, D. E., Reardon, J., and Griffith, D. W. T.: Emissions from
31 smoldering combustion of biomass measured by open-path Fourier transform infrared
32 spectroscopy, *Journal of Geophysical Research: Atmospheres*, 102, 18865-18877,
33 10.1029/97jd00852, 1997.
- 34 Yokelson, R. J., Susott, R., Ward, D. E., Reardon, J., and Griffith, D. W.: Emissions from
35 smoldering combustion of biomass measured by open-path Fourier transform infrared
36 spectroscopy, *Journal of Geophysical Research: Atmospheres* (1984-2012), 102 (D15),
37 18865-18877, 2008.

38

39

40

1

2

1 Table 1. Site characteristics. Mean annual air temperature ($^{\circ}\text{C}$) and mean annual rainfall (mm yr^{-1}) are long-term values (1981-2010); Met
 2 Éireann <http://www.met.ie/> and Met Office UK; <http://www.metoffice.gov.uk>). *Time between cessation of peat extraction and the study period.

Site name	Boora	Blackwater	Bellacorrick	Turraun	Middlemuir Moss	Little Woolden	Clara	Glenlahan	Moyarwood
Site code	IP1	IP2	IP3	IP4	IP5	IP6	DP1	DP2	DP3
Time since last extraction*	>20 years	>25 years	>10 years	>30 years	>10 years	ca. 1	0	>20 years	>20 years
Study period	1/9/2007: 30/8/2009	1/5/2011: 30/4/2014	1/1/2012: 31/12/2013	1/1/2002: 31/12/2003	1/1/2003: 31/10/2004	1/1/2013: 31/12/2014	1/4/2006: 31/3/2007	1/4/2006: 31/3/2007	1/4/2013: 31/3/2014
Latitude	53.203	53.297	54.128	53.260	57.60	53.451	53.316	53.103	53.346
Longitude	-7.726	-7.965	-9.556	-7.720	-2.15	-2.468	-7.647	-7.538	-8.514
Sub-region	Irish Midlands	Irish Midlands	North-West Ireland	Irish Midlands	North-East Scotland	Northern England	Irish Midlands	Irish Midlands	Western Ireland
Mean annual air temperature ($^{\circ}\text{C}$)	9.3	9.8	10.3	9.3	8.0	10.2	9.3	9.3	10.0
Mean annual rainfall (mm yr^{-1})	970	907	1245	807	851	867	970	804	1193
Vegetation									
Peat type	Phragmites	Phragmites	Cyperaceous	Phragmites	Sphagnum/ Cyperaceous	Sphagnum/ Cyperaceous	Sphagnum	Ericaceous	Sphagnum
von Post scale	H7	H7	H5 to 6	H7	H8	H6 to 7	H6	H6	H6
Parent material	Limestone	Limestone	Shale	Limestone	Granite drifts and rocks	Triassic Sandstone	Limestone	Old Red Sandstone	Limestone
Peat depth (m)	1.0	1.5	0.5	0.5-1.8	0.7-3.1	0.5-1.75	4	0.4	4.4
pH	4.3	4.9	3.8	6.3	3.6-4.1	2.9	4.0	3.8	4.4
C (%)	50	52.4	56	52	52	49.1	49.8	29.1	51.5
N (%)	1.09	2.14	0.97	2.1	1.4	1.34	1.46	0.69	1.32
C:N	45.9	24.5	57.7	24.8	37	36.6	34.1	42.2	39

Calluna vulgaris, *Erica tetralix*, *Cladonia* sp.

1 Table 2. Emission factors (tonnes CO₂-C ha⁻¹ yr⁻¹) for sites IP1-6 and DP1-3.

2 Uncertainties are 95% confidence intervals.

Site	CO ₂ -C (t ha ⁻¹ yr ⁻¹)	95% confidence interval (t ha ⁻¹ yr ⁻¹)	
IP1	1.82	1.75	1.89
IP2	1.53	1.37	1.60
IP3	1.38	1.25	1.52
IP4	2.86	2.65	3.06
IP5	0.93	0.59	1.27
IP6	1.70	1.43	1.98
Emission factor	1.70	1.23	2.17
DP1	1.76	1.59	1.99
DP2	2.03	1.73	2.30
DP3	1.14	0.85	1.41
Emission factor	1.64	1.22	2.06

11

12

13

1 Table 3. Mean modified combustion efficiency (MCE) and emission factors (g kg^{-1} dry fuel
 2 burned) reported by this study and those for the same trace gases reported by previous studies
 3 of temperate or boreal peat (Yokelson et al. 1997; Stockwell et al. 2014). The mean and
 4 standard deviation of the emission factor is calculated from individual sample burns. *nr*=not
 5 reported.

Trace Gas	Emission Factor (g kg^{-1} dry fuel burned)			
	Irish sphagnum moss peat (this study)	Canadian boreal peat (Stockwell et al. 2014)	North Carolina temperate peat (Stockwell et al. 2014)	Alaska/Minnesota peat (Yokelson et al. 1997)
MCE	0.837 ± 0.019	0.805 ± 0.009	0.726 ± 0.009	0.809 ± 0.033
CO ₂	1346 ± 31	1274 ± 19	1066 ± 287	1395 ± 52
CO	218 ± 22	197 ± 9	276 ± 139	209 ± 68
CH ₄	8.35 ± 1.3	6.25 ± 2.17	10.9 ± 5.3	6.85 ± 5.66
C ₂ H ₄	1.74 ± 0.23	0.81 ± 0.29	1.27 ± 0.51	1.37 ± 0.51
C ₂ H ₆	1.53 ± 0.17	<i>nr</i>	<i>nr</i>	<i>nr</i>
CH ₃ OH	0.60 ± 0.87	0.75 ± 0.35	2.83 ± 2.87	4.04 ± 3.43
HCN	2.21 ± 0.35	1.77 ± 0.55	4.45 ± 3.02	5.09 ± 5.64
NH ₃	0.73 ± 0.50	2.21 ± 0.24	1.87 ± 0.37	8.76 ± 13.76

6
7
8
9
10
11
12
13
14
15
16

1

2 Table 4. Annual CO₂-C emissions (tonnes CO₂-C yr⁻¹) from peatlands managed for extraction
3 in the ROI and UK calculated using the IPCC 2006 Good Practice Guidance (Tier 1 value:
4 0.2 and 1.1 t CO₂-C ha⁻¹ yr⁻¹ for nutrient poor and nutrient rich peatlands respectively), the
5 IPCC 2013 Wetlands Supplement (Tier 1 value: 2.8 t CO₂-C ha⁻¹ yr⁻¹) and the Emission
6 Factors derived in this study (Table 2). Areas (ha) and CO₂-C emissions using the IPCC 2006
7 Good Practice Guidance values are taken from the 2014 National Inventory Reports (NIR)
8 for the ROI (Duffy et al., 2014) and the UK (Webb et al., 2014).

9

Country	Area (ha)	Emissions (tonnes CO ₂ -C yr ⁻¹)		
		IPCC 2006	IPCC 2013	This study
ROI	52,422	9,312	146,782	88,069
England	4,790	960	13,412	8,047
Scotland	1,610	545	4,508	2,705
Wales	482	95	1,350	810
N. Ireland	1,030	518	2,884	1,730
UK	7,912	2,118	22,154	13,292

10

11

12

13

14

15

16


17

18

19

20

1



2

3

4 Figure 1. Annual rainfall (mm), mean annual water tables (cm), mean annual temperature

5 (°C) at 5 cm depths (T_{5cm}) at sites IP1 (two years), IP2 (three years), IP3 (two years), IP4

6 (two years), IP5 (one year), IP6 (two years), DP1 (one year), DP2 (one year) and DP3 (one

7 year). Dotted horizontal line indicates 30 year mean rainfall at each site (1981-2010; Met


8 Éireann <http://www.met.ie/> and Met Office UK; <http://www.metoffice.gov.uk>). Error bars are

9 standard deviations. Negative water table values indicate water level below the soil surface.

10

11

1




2

3 Figure 2. Frequency distribution of soil temperature at 5cm depth (T_{5cm}) at sites IP1-6 shown
4 as a percentage (%) of total count.


5

1




2

3 Figure 3. (a) Ecosystem respiration (R_{eco} ; $\text{mg CO}_2\text{-C m}^{-2} \text{hr}^{-1}$) at sites IP1-6, (b) R_{eco} (mg
4 $\text{CO}_2\text{-C m}^{-2} \text{hr}^{-1}$) at sites DP1-3 and (c) net ecosystem exchange (NEE; $\text{mg CO}_2\text{-C m}^{-2} \text{hr}^{-1}$)
5 when $\text{PPFD} > 1000 \mu\text{mol m}^{-2} \text{s}^{-1}$ at sites DP1-3. Positive values indicate $\text{CO}_2\text{-C}$ flux from the
6 peatland to the atmosphere (source) and negative values indicate $\text{CO}_2\text{-C}$ flux from the
7 atmosphere to the peatland (sink). The 10th and 90th percentile are indicated by the bars, the
8 25th and 75th percentiles with the top and bottom of the box and the median value by the
9 centre line. Different letters indicate significant differences in the *post-hoc* test for multiple
10 comparisons.



1
2
3
4
5
6
7
8
9


Figure 4. Annual cumulative ecosystem respiration (R_{eco} : $\text{g CO}_2\text{-C m}^{-2}$) at sites IP1-6. Positive values indicate $\text{CO}_2\text{-C}$ flux from the peatland to the atmosphere (source). Value at end of the curve indicates the total annual R_{eco} value. Brown line indicates year 1, black line year 2 and grey line year 3 of the study at the individual sites. Note the differences in integration period between sites (x axis).



1

2

3 Figure 5. Annual cumulative gross primary productivity (GPP: $\text{g CO}_2\text{-C m}^{-2}$), ecosystem
 4 respiration (R_{eco} : $\text{g CO}_2\text{-C m}^{-2}$), heterotrophic respiration (R_{H} : Site DP1 only) and net
 5 ecosystem exchange (NEE: $\text{g CO}_2\text{-C m}^{-2}$) at sites DP1-3. Positive values indicate $\text{CO}_2\text{-C}$ flux
 6 from the peatland to the atmosphere (source) and negative values indicate $\text{CO}_2\text{-C}$ flux from
 7 the atmosphere to the peatland (sink). Value at end of the curve indicates the total annual
 8 value for each component. Note the differences in integration period between sites (x axis).



1

2 Figure 6. Relationship between (a) ecosystem respiration (R_{eco} : t CO₂-C ha⁻¹ yr⁻¹) and mean
 3 soil temperature (°C) at 5 cm depth at the IP sites and (b) net ecosystem exchange (NEE: t
 4 CO₂-C ha⁻¹ yr⁻¹) and leaf area index (LAI: m² m⁻²). Circles indicate an annual value.


5

6

7

8

9



1

2 Figure 7. Carbon dioxide emissions ($t\ CO_2-C\ ha^{-1}\ yr^{-1}$) from peatlands managed for extraction
 3 in Canada, ROI/UK (this study) and Fenno-scandinavia. The 10th and 90th percentile are
 4 indicated by the bars, the 25th and 75th percentiles with the top and bottom of the box and
 5 the median value by the centre line.

6 (Data for Canada and Fenno-Scandia taken from the following studies; Tuittila and Komulainen, 1995; Sundh et
 7 al., 2000; Waddington et al., 2002; Glatzel et al., 2003; McNeil and Waddington, 2003; Tuittila et al., 2004;
 8 Cleary et al., 2005; Alm et al., 2007a; Shurpali et al., 2008; Waddington et al., 2010; Järveoja et al., 2012;
 9 Mander et al., 2012; Salm et al., 2012; Strack et al., 2014). Where studies reported seasonal fluxes (typically
 10 May to October), these were converted to annual fluxes by assuming that 15% of the flux occurs in the non-
 11 growing season (Saarnio et al., 2007).