



This discussion paper is/has been under review for the journal Biogeosciences (BG).
Please refer to the corresponding final paper in BG if available.

Human land uses enhance sediment denitrification and N₂O production in Yangtze lakes primarily by influencing lake water quality

W. Liu¹, L. Yao^{1,2}, Z. Wang^{1,2}, Z. Xiong^{1,2}, and G. Liu¹

¹Key Laboratory of Aquatic Botany and Watershed Ecology, Wuhan Botanical Garden, Chinese Academy of Sciences, Wuhan 430074, China

²University of Chinese Academy of Sciences, Beijing 100049, China

Received: 23 April 2015 – Accepted: 6 May 2015 – Published: 22 May 2015

Correspondence to: G. Liu (liuwz@wbgcas.cn)

Published by Copernicus Publications on behalf of the European Geosciences Union.

BGD

12, 7815–7844, 2015

**Human land uses
enhance sediment
denitrification**

W. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Sediment denitrification in lakes alleviates the effects of eutrophication through removal of nitrogen to the atmosphere as N_2O and N_2 . However, N_2O contributes notably to the greenhouse effect and global warming. Human land uses (e.g., agricultural and urban areas) strongly affect lake water quality and sediment characteristics, which, in turn, may regulate lake sediment denitrification and N_2O production. In this study, we investigated sediment denitrification and N_2O production and their relationships to within-lake variables and watershed land uses in 20 lakes from the Yangtze River basin in China. The results indicated that both lake water quality and sediment characteristics were significantly influenced by watershed land uses. Increased background denitrification rate would result in increased N_2O production rate. Background denitrification and N_2O production rates were positively related to water nitrogen concentrations but were not significantly correlated with sediment characteristics and plant community structure. A significant positive relationship was observed between background denitrification rate and percentage of human-dominated land uses (HDL) in watersheds. Structural equation modelling revealed that the indirect effects of HDL on sediment denitrification and N_2O production in Yangtze lakes were mediated primarily through lake water quality. Our findings also suggest that although sediments in Yangtze lakes can remove large quantities of nitrogen through denitrification, they may also be an important source of N_2O , especially in lakes with high nitrogen content.

1 Introduction

As a global environmental issue, land use change has numerous consequences, ranging from changes in global climate to local water quality (Foley et al., 2005). Conversion of natural land use (e.g., forests and wetlands) to human land use (e.g., cropland and urban areas) releases large quantities of pollutants, including nitrogen, and has widespread effects on biodiversity and ecological function of lakes, rivers and other

BGD

12, 7815–7844, 2015

Human land uses enhance sediment denitrification

W. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



water bodies (Müller et al., 1998). Lake water quality has deteriorated worldwide due to increasing point- and nonpoint-source pollution in watersheds (Nielsen et al., 2012). For instance, the percentage of eutrophic lakes in China has rapidly increased from 41 % in 1980 to 85 % in 2005 (Liu et al., 2010). The effects of watershed land uses on lake water quality are well-documented (Liu et al., 2012; Nielsen et al., 2012). However, relatively few studies have reported the relationship between watershed land uses and lake sediment characteristics, and their results are inconsistent (Müller et al., 1998; Bruesewitz et al., 2011).

Denitrification plays an important role in lake ecosystems by removing excess nitrogen input from watersheds (McCrackin and Elser, 2010). The denitrification process predominantly occurs under anaerobic or anoxic conditions and involves the successive reduction of nitrate (NO_3^-) to nitrite (NO_2^-), nitric oxide (NO), nitrous oxide (N_2O) and, finally, dinitrogen gas (N_2) (Sirivedhin and Gray, 2006). In lakes, denitrification capacity is greatest in sediments due to the favourable environment for denitrifying bacteria and fungi (McCrackin and Elser, 2010). Sediment denitrification can account for approximately 82 % of the total nitrate loss in lakes (Kreiling et al., 2011). Despite the significant increases in nitrogen loading in lake ecosystems and the growing attention to water quality degradation and toxic algal blooms, our understanding of large-scale sediment denitrification in lakes remains limited (McCrackin and Elser, 2010; Bruesewitz et al., 2011; Liu et al., 2015).

N_2O is an intermediate product of sediment denitrification and may be released into the atmosphere in substantial amounts under certain conditions (Hefting et al., 2006). In Lake Taihu of China, both the littoral and pelagic zones can release large quantities of N_2O , especially during the summer algal bloom period (Wang et al., 2007). However, N_2O fluxes in the pelagic regions of shallow boreal lakes are reported to be negligible (Huttunen et al., 2003), which indicates that lakes are only moderate sources of N_2O (Mengis et al., 1997). N_2O is a powerful greenhouse gas that is approximately 300 times stronger than CO_2 and is responsible for approximately 6 % of global warming (IPCC, 2007). Thus, it is important to examine the N_2O production during sediment

BGD

12, 7815–7844, 2015

Human land uses enhance sediment denitrification

W. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



denitrification. A wide range of relative N_2O production during denitrification has been reported for wetlands, forests and cultivated soils (e.g., Garc  -Ruiz et al., 1998; Ullah et al., 2005). Seitzinger and Kroeze (1998) have indicated that the relative N_2O production for heavily polluted river and estuary sediments is approximately 0.03. By contrast, the relative N_2O production for lakes is still poorly understood (McCrackin and Elser, 2010).

At the within-lake scale, overlying water quality and sediment characteristics are recognised as the primary determinants of sediment denitrification and N_2O production (Saunders and Kalff, 2001; Hasegawa and Okino, 2004; Zhong et al., 2010; Rissanen et al., 2011; Liu et al., 2015). Recently, several studies have found that aquatic plant communities can also greatly influence sediment denitrification in fresh waters (e.g., Forshay and Dodson, 2011; Wang et al., 2013). Plants may affect denitrification primarily through the modification of carbon inputs and soil redox conditions via oxygen excretion from roots (Sutton-Grier et al., 2013). At the watershed scale, few studies have addressed the indirect effects of watershed land uses on lake sediment denitrification and N_2O production (but see McCrackin and Elser, 2010; Bruesewitz et al., 2011; Liu et al., 2015), and no studies have examined whether the indirect effects are mainly mediated through lake water quality or sediment characteristics. A better understanding of multi-scale determinants of sediment denitrification in lakes is necessary to achieve high nitrogen removal efficiency and low N_2O production rates.

In this work, we investigated sediment denitrification and N_2O production and their relationships to within-lake variables and watershed land uses in 20 lakes from the Yangtze River basin in China. We hypothesised that the human land uses in watersheds would influence sediment denitrification and N_2O production in lakes, primarily via effects on lake water quality. The objectives of this study were (1) to examine if watershed land uses could influence lake sediment characteristics, as well as water quality, (2) to determine the effects of water quality, sediment characteristics and plant community structure on lake sediment denitrification and N_2O production, and (3) to re-

BGD

12, 7815–7844, 2015

Human land uses enhance sediment denitrification

W. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



veal the mechanisms of lake sediment denitrification and N₂O production in response to human land uses in watersheds for the first time.

2 Methods

2.1 Study sites

The Yangtze River, originating from the Qinghai-Tibet Plateau and flowing into the Pacific Ocean in the city of Shanghai, is the largest river in China and has a drainage area of approximately 1.8 million km² and a total length of 6300 km (Fig. 1). There are 648 lakes with areas greater than 1 km² in the Yangtze River basin (Wang and Dou, 1998). Most of these lakes are concentrated in the middle and lower Yangtze River basin (i.e., the Yangtze floodplain; Fig. 1), where alluvium predominates. Lakes in the Yangtze floodplain are generally shallow (mean depth < 5 m) with short hydraulic retention times (Liu et al., 2011).

Over the past decades, the watershed land uses of many Yangtze lakes have undergone substantial changes (Liu et al., 2012). Large areas of forests, wetlands and other natural landscapes have been transformed into agricultural, industrial, and urban lands. For instance, the percentage of built-up lands in the Chaohu Lake watershed has increased from 6.59 to 9.58 % during the period of 1979–2008 (Wu, 2011). As a result of watershed land use changes, water quality in many lakes has deteriorated considerably in recent years, and up to 86 % of the Yangtze lakes are eutrophic or even hypereutrophic (Yang et al., 2010).

2.2 Field sampling

During the summer (from 30 July to 7 August) of 2012, twenty shallow lakes in the middle and lower Yangtze River basin of China were selected non-randomly on the basis of ease of access (Fig. 1). These selected lakes covered a wide range of physical and chemical conditions and vegetation characteristics (Wang and Dou, 1998). The name,

BGD

12, 7815–7844, 2015

Human land uses enhance sediment denitrification

W. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Human land uses
enhance sediment
denitrification

W. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



geographic location and morphology characteristics of these studied lakes are listed in Table S1. To obtain representative samples, one transect from the littoral zone to the lake centre was randomly established in each lake and 3–4 sampling sites were chosen along this transect at regular intervals based on lake area (Yang et al., 2008). Littoral sampling sites were generally located at a water depth less than 1.5 m, approximately 50–100 m away from the lake shore.

At each sampling site, three replicate surface sediments were randomly collected within an approximately 50 m² area from a boat using a Peterson dredge and then mixed and homogenised to form a composite sample. For each site, approximately 1 kg of sediment was placed in a sealed plastic bag and stored at approximately 5 °C in a refrigerator until return to the laboratory. At each sampling site, a 200 mL unfiltered water sample was collected at a depth of approximately 1 m for use in the denitrification and N₂O production assays. At the same time, we collected an additional 200 mL water sample at the same depth for water quality analyses in the laboratory. Water sampling was performed before sediment collection to prevent sediments from being resuspended and thereby contaminating the water. At each sampling site, submerged macrophytes were sampled using a pronged grab (25 cm × 35 cm) with three replicates, and the species richness (i.e., mean species number recorded in a sampling site) and fresh plant biomass were calculated.

2.3 Denitrification and N₂O production assays

The most common method for measuring sediment denitrification is based on the ability of acetylene to inhibit the reduction of N₂O to N₂ during denitrification (Groffman et al., 1999). Although the acetylene blockage technique has a number of limitations (Yu et al., 2010; Felber et al., 2012), it is still amenable to large-scale comparisons of denitrification, especially for systems with moderate or high NO₃⁻ concentrations (Groffman et al., 2006). In this study, we measured potential and background denitrification rates of lake sediments using the acetylene blockage technique. The potential denitrification rate was measured under optimal conditions (by supplying an excess of NO₃⁻

and organic carbon and ensuring anoxia) and thus provided an upper-bound estimate of in situ denitrification. Background denitrification (i.e., unamended denitrification) rate was a conservative estimate of in situ denitrification without carbon and NO_3^- amendments because acetylene also inhibited nitrification (NO_3^- production). The increase in N_2O concentrations was always linear throughout the short assay duration, therefore, short-term incubation (2–4 h) was recommended for measuring sediment denitrification of lakes (Bruesewitz et al., 2012).

For potential denitrification rate assays, 50 g of homogenised sediments from each sampling site were slurried with 30 mL of incubation solution (final concentrations: 0.1 g L^{-1} KNO_3 , 0.18 g L^{-1} glucose and 1 g L^{-1} chloramphenicol) in a 250 mL serum bottle. Each bottle was then sealed and purged with N_2 gas for 2 min to induce anoxic conditions. Approximately 10 % of the bottle headspace was replaced with acetylene to block the conversion of N_2O to N_2 during denitrification. We measured background denitrification and N_2O production rates using a similar procedure, but with the addition of 30 mL of unfiltered lake water instead of the incubation solution (McCrackin and Elser, 2012). Parallel incubations with and without acetylene (10 % vol/vol in the bottle headspace) were used to differentiate between background denitrification and N_2O production (McCrackin and Elser, 2010). All bottles were then incubated in the dark for 4 h at 25°C (the approximate in situ water temperature). At the beginning and end of incubation, 5 mL of headspace gas samples were collected from each bottle (after shaking vigorously) using a syringe. The N_2O concentrations were measured using a gas chromatograph (Agilent 7890, Santa Clara, CA, USA) equipped with an electron capture detector.

Potential denitrification, background denitrification and N_2O production rates were calculated as the difference between the initial and final headspace N_2O concentrations (corrected for N_2O dissolved in water, Bunsen coefficient = 0.544) divided by the incubation time (Hayakawa et al., 2012), and was expressed on the basis of dry matter of sediment ($\text{ng N g}^{-1} \text{ h}^{-1}$). The relative N_2O production was calculated by dividing the

BGD

12, 7815–7844, 2015

Human land uses enhance sediment denitrification

W. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



N₂O production rate by the background denitrification rate (García-Ruiz et al., 1998; McCrackin and Elser, 2010).

2.4 Sediment characteristics and water quality measurements

Sediment pH was measured in a soil to water ratio of 1 : 5 (v/v) using a pH meter and bulk density was determined by weighing soil cores of known volume after drying for 24 h at 105 °C. Sediment moisture was measured gravimetrically (24 h at 105 °C) from 30 g sediment samples. Sediment total nitrogen (STN) was measured using the Kjeldahl method after digesting samples in a digester using a sulfuric acid/mercuric oxide catalyst. Sediment total carbon (STC) content of air-dried samples was analysed by a TOC analyser (Vario TOC cube, Elementar, Germany).

Conductivity (Cond), oxidation–reduction potential (ORP), dissolved oxygen (DO), nitrate (NO₃⁻) and ammonium (NH₄⁺) were measured on site at the sampling depth using a YSI 6920 multiparameter water quality probe (YSI Inc., Yellow Springs, Ohio, USA). Total nitrogen (TN) concentration was analysed using the micro-Kjeldahl method. The concentrations of total carbon (TC) and total organic carbon (TOC) in waters were measured with a TOC analyser (Vario TOC cube, Elementar, Germany).

2.5 Watershed land use calculation

After delineating the watershed boundaries of the 20 studied lakes using a 1 km resolution digital elevation model, overlay functions were used to extract a land use map of each lake’s watershed from 100 m resolution national land cover data in ArcGIS 10 software (ESRI, Redlands, California, USA). The national land cover data, interpreted from recent Landsat TM images, were obtained from the Data Sharing Infrastructure of Earth System Science in China (http://www.geodata.cn/).

The original land use classes were further grouped into four main categories: (1) vegetation, including forest and grassland; (2) cropland, including dry land and paddy fields; (3) built-up land, including urban areas, rural settlements and others such as



industrial areas, roads, and airports; and (4) water bodies, including lakes, rivers, streams, reservoirs, ponds, and wetlands (Fig. 1). Bare land was not included in our analysis because its area in watersheds was very small. Cropland and built-up land can be considered as human-dominated land uses (HDL; Abdullah and Nakagoshi, 2006). The percentages of land use types in each lake's watershed were calculated in ArcGIS 10 software. We used the HDL here, because the correlations between in-lake characteristics (i.e., water quality and sediment characteristics) and the percentage of cropland or percentage of built-up land were weak.

To quantify the vegetation cover, we calculated the normalised difference vegetation index (NDVI) of each watershed from the Moderate-resolution Imaging Spectroradiometer (MODIS) red and near-infrared bands. The 250 m resolution MODIS data for the years 2010 and 2011 was obtained from the USGS EROS Data Center. The NDVI is a reflection of the biophysical condition of a watershed's vegetation cover, which, in turn, may affect the water runoff and water quality (Griffith et al., 2002).

2.6 Statistical analyses

For each studied lake, we calculated the mean values for sediment denitrification, N_2O production, water quality, sediment characteristics and plant community structure (Tables S2 and S3). Before statistical analyses, we tested data for normal distribution using a Shapiro–Wilk test. When possible, non-normally distributed data were natural log or square root transformed to reach a normal distribution. To examine the relationships among watershed land uses, water chemistry, sediment characteristics, plant community structure, sediment denitrification and N_2O production at the lake level, we performed Pearson correlation and regression analyses using PASW Statistics 18 software (IBM SPSS Inc., Chicago, USA).

Structural equation modelling (SEM) with Bayesian estimation was used to study the indirect effects of watershed human land use on lake sediment denitrification and N_2O production. In contrast to regression analysis, SEM can test indirect effects between two variables (i.e., effects mediated by other intermediary variables). We used results

Human land uses enhance sediment denitrification

W. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Human land uses
enhance sediment
denitrification

W. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



from our correlation analysis to select the promising explanatory variables to include in the SEM models. Given the positive correlations between NO_3^- and TN ($r = 0.86$, $P < 0.01$) and between NO_3^- and NH_4^+ ($r = 0.47$, $P < 0.05$), we only included the NO_3^- in the SEM analyse to simplify the final models. The software Amos 20 (IBM SPSS Inc., Chicago, USA) was used to design the SEM models and calculate path coefficients, squared multiple correlations, direct and indirect effects. Indirect effects, i.e., effects mediated by other variables, were calculated by multiplying the standardised path coefficients (i.e., estimates) of the direct effects of the variables involved in the total pathway. We used the chi-square (χ^2) test and the comparative fit index (CFI) to evaluate the overall fit of the SEM models. According to Kline (1998), a good fit was indicated by an insignificant χ^2 statistic ($P > 0.05$) and a CFI value > 0.90 . The χ^2 provides an estimate of how closely the proposed SEM model matches the structure of the actual data. Thus, an insignificant χ^2 statistic ($P > 0.05$) indicates that the structure of the proposed model is no different than the structure of the data.

3 Results

3.1 Within-lake characteristics and watershed land uses

The highest TC and TN concentrations in water were found in Lake Makouhu (31.40 mg L^{-1}) and Lake Yiaihu (2.87 mg L^{-1}), respectively (Table S2). The mean TC and TN contents in sediments were 25.18 and 0.51 mg g^{-1} , respectively (Table 1). Submerged plants were only found in 5 lakes, with maximum biomass (2438 g m^{-2}) recorded in Lake Haikouhu (Table S2).

The percentage of HDL in watersheds varied from 33.60 to 80 % with a mean of 60.91 % (Table 1). Among water quality and sediment characteristics, DO, NO_3^- , NH_4^+ , TN and STN were significantly correlated with the percentage of HDL (Table 2). There was no significant relationship between plant community structure and watershed land uses (Table 2).

3.2 Denitrification and N₂O production

Potential denitrification rates varied widely across the 20 lakes and ranged from 11.24 ng N g⁻¹ h⁻¹ in Lake Chaohu to 125.37 ng N g⁻¹ h⁻¹ in Lake Makouhu (Table S3). Background denitrification rates were lowest in Lake Huangdahu at 0.67 μg N g⁻¹ h⁻¹, and highest in Lake Linghu at 18.95 ng N g⁻¹ h⁻¹. N₂O production rates ranged from 0.05 to 0.71 ng N g⁻¹ h⁻¹, with the largest value recorded in Lake Caizihu where the relative N₂O production was 0.13 (Table S3).

There was no significant relationship between potential denitrification rate and background denitrification rates, N₂O production rate or relative N₂O production (figures not shown). N₂O production rates increased with increasing background denitrification rates ($R^2 = 0.61$, $P < 0.01$; Fig. 2).

3.3 Relationship of denitrification and N₂O production to within-lake characteristics

Denitrification and N₂O production rates were found to be significantly related to several water quality parameters (Table 3). However, sediment characteristics and plant community structure had no significant relationships with denitrification and N₂O production rates. Potential denitrification rate was negatively correlated with DO (Table 3). We determined that both background denitrification rate and N₂O production rate were enhanced at higher NO₃⁻ and TN concentrations and lower ORP and DO concentrations (Table 3).

3.4 Relationship of denitrification and N₂O production to watershed land uses

Watershed land uses were not significantly related to potential denitrification rates or relative N₂O production (figures not shown). However, background denitrification rate was found to be positively correlated with the percentage of HDL in watersheds

BGD

12, 7815–7844, 2015

**Human land uses
enhance sediment
denitrification**

W. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



($R^2 = 0.29$, $P < 0.05$; Fig. 3b). We also found that N_2O production rate was significantly related to the percentage of water bodies in watersheds ($R^2 = 0.27$, $P < 0.05$; Fig. 3a).

3.5 Indirect effects of human land uses on denitrification and N_2O production

5 The results of the SEM analysis indicated that the fit of the data to the models was acceptable (Fig. 4). In the potential denitrification model ($\chi^2 = 3.063$, $df = 4$, $P = 0.547$, CFI = 1.000), the total indirect effect of watershed HDL on potential denitrification (0.26) was small and statistically insignificant (Table 4, Fig. 4a). In the background denitrifica-
10 tion model ($\chi^2 = 1.281$, $df = 4$, $P = 0.865$, CFI = 1.000), 100 % of the positive indirect effect of HDL on background denitrification (0.55) was mediated through water quality (principally via NO_3^- ; Table 4, Fig. 4b). In both the N_2O production model ($\chi^2 = 1.920$, $df = 4$, $P = 0.750$, CFI = 1.000) and relative N_2O production model ($\chi^2 = 2.483$, $df = 4$, $P = 0.648$, CFI = 1.000), HDL indirect effects were largely transmitted via water quality (Table 4).

4 Discussion

15 4.1 Effects of watershed human land use on in-lake characteristics

Numerous studies have revealed that human modification of the land uses plays a key role in determining the water quality of adjacent aquatic ecosystems (e.g., generation of point and non-point source pollutants and expansion of impervious surface area) (Crosbie and Chow-Fraser, 1999; Arbuckle and Downing, 2001; Taranu and Gregory-
20 Eaves, 2008). In this study, human land use in watersheds was strongly and positively related to water N concentrations in lakes (Table 2). Built-up land, including urban, industrial areas and rural settlements, is the leading source of point source pollution in the Yangtze River basin (Liu et al., 2012). Chaohu Lake, one of the lakes included in the present study, received 2.17×10^8 t of industrial wastewater and domestic sewage

BGD

12, 7815–7844, 2015

Human land uses enhance sediment denitrification

W. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Human land uses enhance sediment denitrification

W. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



discharged from adjacent Hefei City every year (Shang and Shang, 2005). Cropland is recognised as the most important source of non-point source pollution affecting lakes in many countries. Approximately half of the area of the middle and lower Yangtze River basin is covered by agriculture land, especially paddy fields (Liu et al., 2011). In the Yangtze River basin, the annual consumption of nitrogen fertiliser on agricultural lands increased from 94 kg ha⁻¹ in 1980 to 509 kg ha⁻¹ in 2000 (Liu et al., 2008). Moreover, the nitrogen use efficiency decreased considerably from 0.31 to 0.23 between 1980 and 2000. Most of the excess nitrogen in paddy fields was discharged into aquatic environments through surface runoff.

Relatively few studies have reported on the relationship between watershed landscapes and lake sediment characteristics and their results are inconsistent (Müller et al., 1998; Bruesewitz et al., 2011). Bruesewitz et al. (2011) found that the percentage of catchment agriculture was positively related to sediment carbon but not nitrogen. These researchers found that agricultural landscapes can increase carbon inputs to aquatic ecosystems as a result of increased soil erosion and organic carbon transport or through increased lake productivity resulting in higher rates of organic matter deposition in sediments. However, Müller et al. (1998) indicated that TN concentration in lake sediments strongly increased with increasing agricultural land use in watersheds. Our study also found a significant relationship between watershed human landscapes and STN. It is noteworthy that the correlation between human landscapes and water TN was greater than that between human land uses and STN (Table 2). One possible explanation is that water TN level in lakes primarily reflects recent land use patterns in watersheds. Sediment TN content, however, reflects sedimentation and deposition of nutrients that have occurred over several decades. Any modelling of water and sediment nutrient contents using recent land use data is likely to show stronger correlations with water than sediment nutrient levels (Houlahan and Findlay, 2004).

4.2 The role of sediment denitrification in nitrogen removal in Yangtze lakes

Sediment denitrification is recognised as the most important pathway of nitrogen removal in lakes, followed by nitrogen sedimentation and uptake by aquatic macrophytes (Saunders and Kalff, 2001; Lindau et al., 2009). Scaled on an areal basis using the bulk density of the top 7 cm of sediment (Liu et al., 2015), background denitrification rate ranged from 354 to 10 968 kg N km⁻² year⁻¹, with a mean of 1065 kg N km⁻² year⁻¹. This mean rate is smaller than the global average of lakes (3421 kg N km⁻² year⁻¹) estimated by a NiRReLa model (Harrison et al., 2009). Background denitrification rate can be considered as a conservative estimate of actual in situ denitrification. Therefore, according to the areal background denitrification rate and the surface area of the 20 studied Yangtze lakes (Table S1), we find that Lake Chaohu and Lake Caizihu can at least remove 601 880 and 553 812 kg of N every year, respectively.

It is hard to quantify total nitrogen inputs to a lake. To evaluate the proportion of annual nitrogen inputs that are removed via denitrification, Seitzinger et al. (2006) established a relationship between the percentages of N removed by denitrification and water residence time (WRT), detailed below:

$$\% \text{ N removed by denitrification} = 23.4(\text{WRT})^{0.204} \quad (1)$$

WRT for most of the lakes we sampled are unknown. Because almost all lakes in the Yangtze floodplain are shallow (mean depth < 5 m), the WRT of these lakes is estimated to be between 1 and 12 months (Wang and Dou, 1998). Therefore, we estimated that 23.40–38.85 % of annual N inputs to the Yangtze lakes can be removed via sediment denitrification.

Net N₂O production during sediment N cycles is gaining increased attention because N₂O is a powerful greenhouse gas (Hefting et al., 2006). The mean net N₂O production rate (i.e., N₂O production in the absence of acetylene) of the studied Yangtze lakes was 0.25 ng N g⁻¹ h⁻¹ or 142.86 kg N km⁻² year⁻¹. This value is approximately 2.1 times greater than the average rate in Norwegian lakes (68.60 kg N km⁻² year⁻¹) calculated

BGD

12, 7815–7844, 2015

**Human land uses
enhance sediment
denitrification**

W. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



by McCrackin and Elser (2010). It should be noted that N_2O production in the absence of acetylene can be derived from both denitrification and nitrification processes, especially under incomplete anaerobic conditions (García-Ruiz et al., 1998; Xu et al., 2008). In the present study, for N_2O production assays, sediments were slurried with unfiltered lake water, which was always saturated with oxygen.

The present study found that the average of relative N_2O production was 0.17, similar to the value (0.18) reported by García-Ruiz et al. (1998). However, our results contrast with those found in other investigations, the majority of which give values less than 0.05 (e.g., Seitzinger, 1988; Beaulieu et al., 2011). This result might be explained by the fact that oxygen in overlying water of shallow lakes could suppress the N_2O reductase enzyme, which reduces N_2O to N_2 during denitrification (García-Ruiz et al., 1998). It is known that this enzyme is the most oxygen sensitive of all enzymes involved in denitrification process, and slurries in the present study were made with unfiltered lake water. The relative N_2O production > 1 implies that the production of N_2O through nitrification must have occurred, and nitrification processes is the major source of N_2O (Xu et al., 2008).

4.3 Local and regional determinants of sediment denitrification in lakes

Environmental variables that affect sediment denitrification can be categorised as proximal or distal regulators (Saggar et al., 2013). Proximal regulators, including NO_3^- concentration, oxygen supply or water content, carbon availability and temperature in sediment and overlying water, affect the instantaneous rate of sediment denitrification. Consistent with previous studies conducted in streams and rivers (García-Ruiz et al., 1998; Inwood et al., 2005), we found that only overlying water quality had significant relationships with sediment denitrification and N_2O production in Yangtze lakes. Our study suggests that nitrogen availability in water is the primary factor limiting sediment denitrification in Yangtze lakes. NO_3^- concentration in water explained approximately 70 % of the variability in denitrification rates in a meta-analysis (Piña-Ochoa and Álvarez-Cobelas, 2006). Some studies also reported a significant relationship be-

BGD

12, 7815–7844, 2015

Human land uses enhance sediment denitrification

W. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



tween denitrification and sediment carbon concentration (e.g., Bruesewitz et al., 2011). However, this relationship is not significant in the Yangtze lakes, most likely because sediment TC concentration was high (25.18 mg g⁻¹) and was not limiting to sediment denitrification.

Distal regulators affect sediment denitrification indirectly by acting on the proximal controls (Saggar et al., 2013). These regulators include factors such as land use and climate. The indirect effects of human land uses on stream sediment denitrification and N₂O emission are mainly mediated through stream water quality or sediment characteristics (Inwood et al., 2007). Only one study has examined the effects of watershed human land uses on sediment denitrification in lakes (Bruesewitz et al., 2011). This study found that potential denitrification but not background denitrification was positively related to the proportion of catchment agriculture. However, consistent with previous observations regarding stream denitrification (e.g., Arango and Tank, 2008), we found a significant positive relationship between background denitrification and human land uses. As we hypothesised, the human land use in watersheds affects the sediment denitrification and N₂O production in Yangtze lakes indirectly, mainly via effects on lake water quality. The positive relationships between watershed human land uses and sediment denitrification were primarily driven by nitrogen loading.

5 Conclusions

Increased human land uses in watersheds have led to elevated nitrogen concentration in both water and sediments in Yangtze lakes. Among the water quality parameters, ORP, DO, NO₃⁻ and TN were significantly related to sediment background denitrification and N₂O production rates. Only the background denitrification rate was found to be positively correlated with the percentage of human land uses in the watersheds. The results of SEM analysis supported the hypothesis that sediment denitrification and N₂O production in the Yangtze lakes were indirectly associated with watershed human land-

BGD

12, 7815–7844, 2015

Human land uses enhance sediment denitrification

W. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



scapes, primarily through the effect of land use on lake water quality, but not sediment characteristics.

**The Supplement related to this article is available online at
doi:10.5194/bgd-12-7815-2015-supplement.**

- 5 *Acknowledgements.* The authors would like to thank Guomin Li and Nanhua Liu for their assistance with field sampling. Funding was provided by the Special S and T Project on Treatment and Control of Water Pollution (2013ZX07102–005) and the National Science Foundation of China (31270583).

References

- 10 Abdullah, S. A. and Nakagoshi, N.: Changes in landscape spatial pattern in the highly developing state of Selangor, peninsular Malaysia, *Landscape Urban Plan.*, 77, 263–275, 2006.
- Arango, C. P. and Tank, J. L.: Land use influences the spatiotemporal controls on nitrification and denitrification in headwater streams, *J. N. Am. Benthol. Soc.*, 27, 90–107, 2008.
- Arbuckle, K. E. and Downing, J. A.: The influence of watershed land use on lake N: P in a predominantly agricultural landscape, *Limnol. Oceanogr.*, 46, 970–975, 2001.
- 15 Beaulieu, J. J., Tank, J. L., Hamilton S, K, Wollheim, W. M., Hall, R. O., Mulholland, P. J., Peterson, B. J., Ashkenas, L. R., Cooper, L. W., Dahm, C. N., Dodds, W. K., Grimm, N. B., Johnson, S. L., McDowell, W. H., Poole, G. C., Valett, H. M., Arango, C. P., Bernot, M. J., Burgin, A. J., Crenshaw, C. L., Helton, A. M., Johnson, L. T., O'Brien, J. M., Potter, J. D., Sheibley, R. W., Sobota, D. J., and Thomas, S. M.: Nitrous oxide emission from denitrification in stream and river networks, *P. Natl. Acad. Sci. USA*, 108, 214–219, 2011.
- 20 Bruesewitz, D. A., Hamilton, D. P., and Schipper, L. A.: Denitrification potential in lake sediment increases across a gradient of catchment agriculture, *Ecosystems*, 14, 341–352, 2011.
- Bruesewitz, D. A., Tank, J. L., and Hamilton, S. K.: Incorporating spatial variation of nitrification and denitrification rates into whole-lake nitrogen dynamics, *J. Geophys. Res.*, 117, G00N07, doi:10.1029/2012JG002006, 2012.
- 25

Human land uses enhance sediment denitrification

W. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Human land uses enhance sediment denitrification

W. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Crosbie, B. and Chow-Fraser, P.: Percentage land use in the watershed determines the water and sediment quality of 22 marshes in the Great Lakes basin, *Can. J. Fish. Aquat. Sci.*, 56, 1781–1791, 1999.
- Felber, R., Conen, F., Flechard, C. R., and Neftel, A.: Theoretical and practical limitations of the acetylene inhibition technique to determine total denitrification losses, *Biogeosciences*, 9, 4125–4138, doi:10.5194/bg-9-4125-2012, 2012.
- Foley, J. A., DeFries, R., Asner, G. P., Barford, C., Bonan, G., Carpenter, S. R., Chapin, F. S., Coe, M. T., Daily, G. C., Gibbs, H. K., Helkowski, J. H., Holloway, T., Howard, E. A., Kucharik, C. J., Monfreda, C., Patz, J. A., Prentice, I. C., Ramankutty, N., and Snyder, P. K.: Global consequences of land use, *Science*, 309, 570–574, 2005.
- Forshay, K. J. and Dodson, S. I.: Macrophyte presence is an indicator of enhanced denitrification and nitrification in sediments of a temperate restored agricultural stream, *Hydrobiologia*, 668, 21–34, 2011.
- García-Ruiz, R., Pattinson, S. N., and Whitton, B. A.: Denitrification in river sediments: relationship between process rate and properties of water and sediment, *Freshwater Biol.*, 39, 467–476, 1998.
- Griffith, J. A., Martinko, E. A., Whistler, J. L., and Price, K. P.: Interrelationships among landscape, NDVI, and stream water quality in the US Central Plains, *Ecol. Appl.*, 12, 1702–1718, 2002.
- Groffman, P. M., Holland, E. A., Myrold, D. D., Robertson, G. P., and Zou, X.: Denitrification, in: *Standard Soil Methods for Long Term Ecological Research*, edited by: Robertson, G. P., Coleman, D. C., Bledsoe, C. C., and Sollins, P., Oxford University Press, New York, 1999.
- Groffman, P. M., Altabet, M. A., Böhlke, J. K., Butterbach-Bahl, K., David, M. B., Firestone, M. K., Giblin, A. E., Kana, T. M., Nielsen, L. P., and Voytek, M. A.: Methods for measuring denitrification: diverse approaches to a difficult problem, *Ecol. Appl.*, 16, 2091–2122, 2006.
- Harrison, J. A., Maranger, R. J., Alexander, R. B., Giblin, A. E., Jacinthe, P., Mayorga, E., Seitzinger, S. P., Sobota, D. J., and Wollheim, W. M.: The regional and global significance of nitrogen removal in lakes and reservoirs, *Biogeochemistry*, 93, 143–157, 2009.
- Hasegawa, T. and Okino, T.: Seasonal variation of denitrification rate in Lake Suwa sediment, *Limnology*, 5, 33–39, 2004.
- Hayakawa, A., Nakata, M., Jiang, R., Kuramochi, K., and Hatano, R.: Spatial variation of denitrification potential of grassland, windbreak forest, and riparian forest soils in an agricultural catchment in eastern Hokkaido, Japan, *Ecol. Eng.*, 47, 92–100, 2012.

Human land uses enhance sediment denitrification

W. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Hefting, M. M., Bobbink, R., and Janssens, M. P.: Spatial variation in denitrification and N_2O emission in relation to nitrate removal efficiency in a N-stressed riparian buffer zone, *Ecosystems*, 9, 550–563, 2006.

Houlahan, J. E. and Findlay, C. S.: Estimating the “critical” distance at which adjacent land-use degrades wetland water and sediment quality, *Landscape Ecol.*, 19, 677–690, 2004.

Huttunen, J. T., Alm, J., Liikanen, A., Juutinen, S., Larmola, T., Hammar, T., Silvola, J., and Martikainen, P. J.: Fluxes of methane, carbon dioxide and nitrous oxide in boreal lakes and potential anthropogenic effects on the aquatic greenhouse gas emissions, *Chemosphere*, 52, 609–621, 2003.

Inwood, S. E., Tank, J. L., and Bernot, M. J.: Patterns of denitrification associated with land use in 9 midwestern headwater streams, *J. N. Am. Benthol. Soc.*, 24, 227–245, 2005.

Inwood, S. E., Tank, J. L., and Bernot, M. J.: Factors controlling sediment denitrification in midwestern streams of varying land use, *Microb. Ecol.*, 53, 247–258, 2007.

IPCC: Summary for Policymakers, in: *Climate Change 2007: Mitigation, Contribution of Working Group III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge University Press, Cambridge, UK and New York, NY, USA, 2007.

Kline, R. B.: *Principles and Practices of Structural Equation Modeling*, The Guilford Press, New York, 1998.

Kreiling, R. M., Richardson, W. B., Cavanaugh, J. C., and Bartsch, L. A.: Summer nitrate uptake and denitrification in an upper Mississippi River backwater lake: the role of rooted aquatic vegetation, *Biogeochemistry*, 104, 309–324, 2011.

Lindau, C. W., Iwai, R. R., Delaune, R. D., and Gambrell, R. P.: Denitrification potential of lake sediment at the Davis Pond Mississippi River freshwater diversion site, Louisiana, USA, *J. Freshwater Ecol.*, 24, 181–189, 2009.

Liu, C., Watanabe, M., and Wang, Q.: Changes in nitrogen budgets and nitrogen use efficiency in the agroecosystems of the Changjiang River basin between 1980 and 2000, *Nutr. Cycl. Agroecosys.*, 80, 19–37, 2008.

Liu, W., Zhang, Q., and Liu, G.: Lake eutrophication associated with geographic location, lake morphology and climate in China, *Hydrobiologia*, 644, 289–299, 2010.

Liu, W., Zhang, Q., and Liu, G.: Effects of watershed land use and lake morphometry on trophic state of Chinese lakes: implications for eutrophication control, *Clean-Soil, Air, Water*, 39, 35–42, 2011.

Human land uses enhance sediment denitrification

W. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Liu, W., Zhang, Q., and Liu, G.: Influences of watershed landscape composition and configuration on lake water quality in the Yangtze River basin of China, *Hydrol. Process.*, 26, 570–578, 2012.
- Liu, W., Wang, Z., Zhang, Q., Cheng, X., Lu, J., and Liu, G.: Sediment denitrification and nitrous oxide production in Chinese plateau lakes with varying watershed land uses, *Biogeochemistry*, 123, 379–390, 2015.
- McCrackin, M. L. and Elser, J. J.: Atmospheric nitrogen deposition influences denitrification and nitrous oxide production in lakes, *Ecology*, 91, 529–539, 2010.
- McCrackin, M. L. and Elser, J. J.: Denitrification kinetics and denitrifier abundances in sediments of lakes receiving atmospheric nitrogen deposition (Colorado, USA), *Biogeochemistry*, 108, 39–54, 2012.
- Mengis, M., Gachter, R., and Wehrli, B.: Sources and sinks of nitrous oxide (N_2O) in deep lakes, *Biogeochemistry*, 38, 281–301, 1997.
- Müller, B., Lotter, A. F., Sturm, M., and Ammann, A.: Influence of catchment quality and altitude on the water and sediment composition of 68 small lakes in central Europe, *Aquat. Sci.*, 60, 316–337, 1998.
- Nielsen, A., Trolle, D., Søndergaard, M., Lauridsen, T. L., Bjerring, R., Olesen, J. E., and Jeppesen, E.: Watershed land use effects on lake water quality in Denmark, *Ecol. Appl.*, 22, 1187–1200, 2012.
- Piña-Ochoa, E. and Álvarez-Cobelas, M.: Denitrification in aquatic environments: a cross-system analysis, *Biogeochemistry*, 81, 111–130, 2006.
- Rissanen, A. J., Tirola, M., and Ojala, A.: Spatial and temporal variation in denitrification and in the denitrifier community in a boreal lake, *Aquat. Microb. Ecol.*, 64, 27–40, 2011.
- Saggar, S., Jha, N., Deslippe, J., Bolan, N. S., Luo, J., Giltrap, D. L., Kim, D. G., Zaman, M., and Tillman, R. W.: Denitrification and N_2O : N_2 production in temperate grasslands: processes, measurements, modelling and mitigating negative impacts, *Sci. Total Environ.*, 465, 173–195, 2013.
- Saunders, D. L. and Kalff, J.: Denitrification rates in the sediments of Lake Memphremagog, Canada–USA, *Water Res.*, 35, 1897–1904, 2001.
- Seitzinger, S. P.: Denitrification in freshwater and coastal marine ecosystems: ecological and geochemical significance, *Limnol. Oceanogr.*, 33, 702–724, 1988.
- Seitzinger, S. P. and Kroeze, C.: Global distribution of nitrous oxide production and N inputs in freshwater and coastal marine ecosystems, *Global Biogeochem. Cy.*, 12, 93–113, 1998.

Human land uses enhance sediment denitrification

W. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Seitzinger, S. P., Harrison, J. A., Böhlke, J. K., Bouwman, A. F., Lowrance, R., Peterson, B., Tobias, C., and Van Drecht, G.: Denitrification across landscapes and waterscapes: a synthesis, *Ecol. Appl.*, 16, 2064–2090, 2006.

Shang, G. P. and Shang, J. C.: Causes and control countermeasures of eutrophication in Chaohu Lake, China, *Chinese Geogr. Sci.*, 15, 348–354, 2005.

Sirivedhin, T. and Gray, K. A.: Factors affecting denitrification rates in experimental wetlands: field and laboratory studies, *Ecol. Eng.*, 26, 167–181, 2006.

Sutton-Grier, A. E., Wright, J. P., and Richardson, C. J.: Different plant traits affect two pathways of riparian nitrogen removal in a restored freshwater wetland, *Plant Soil*, 365, 41–57, 2013.

Taranu, Z. E. and Gregory-Eaves, I.: Quantifying relationships among phosphorus, agriculture and lake depth at an inter-regional scale, *Ecosystems*, 11, 715–725, 2008.

Ullah, S., Breitenbeck, G. A., and Faulkner, S. P.: Denitrification and N₂O emission from forested and cultivated alluvial clay soil, *Biogeochemistry*, 73, 499–513, 2005.

Wang, H., Yang, L., Wang, W., Lu, J., and Yin, C.: Nitrous oxide (N₂O) fluxes and their relationships with water-sediment characteristics in a hyper-eutrophic shallow lake, China, *J. Geophys. Res.*, 112, G01005, doi:10.1029/2005JG000129, 2007.

Wang, S. and Dou, H.: Chinese Lake Catalogue, Science Press, Beijing, 1998 (in Chinese).

Wang, Y., Li, Z., Zhou, L., Feng, L., Fan, N., and Shen, J.: Effects of macrophyte-associated nitrogen cycling bacteria on denitrification in the sediments of the eutrophic Gonghu Bay, Taihu Lake, *Hydrobiologia*, 700, 329–341, 2013.

Wu, L.: Study of LUCC and its driving forces in Chaohu Lake basin during the past 30 years, *Chin. J. Soil Sci.*, 42, 1293–1296, 2011. (In Chinese with English abstract)

Xu, Y., Wan, S., Cheng, W., and Li, L.: Impacts of grazing intensity on denitrification and N₂O production in a semi-arid grassland ecosystem, *Biogeochemistry*, 88, 103–115, 2008.

Yang, G., Ma, R., Zhang, L., Jiang, J., Yao, S., Zhang, M., and Zeng, H.: Lake status, major problems and protection strategy in China, *J. Lake. Sci.*, 22, 799–810, 2010 (in Chinese with English abstract).

Yang, X., Anderson, N. J., Dong, X., and Shen, J.: Surface sediment diatom assemblages and epilimnetic total phosphorus in large, shallow lakes of the Yangtze floodplain: their relationships and implications for assessing long-term eutrophication, *Freshwater Biol.*, 53, 1273–1290, 2008.

Yu, K., Seo, D. C., and DeLanue, R. D.: Incomplete acetylene inhibition of nitrous oxide reduction in potential denitrification assay as revealed by using ^{15}N -nitrate tracer, Commun. Soil Sci. Plan., 41, 2201–221, 2010.

- 5 Zhong, J., Fan, C., Liu, G., Zhang, L., Shang, J., and Gu, X.: Seasonal variation of potential denitrification rates of surface sediment from Meiliang Bay, Taihu Lake, China, J. Environ. Sci., 22, 961–967, 2010.

BGD

12, 7815–7844, 2015

Human land uses enhance sediment denitrification

W. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Human land uses
enhance sediment
denitrification

W. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Table 1.** Summary statistics for sediment denitrification variables, within-lake characteristics and watershed land uses ($N = 20$).

	Unit	Min.	Max.	Mean	S.E.
Denitrification variables					
Potential denitrification rate	ng N g ⁻¹ h ⁻¹	11.24	125.37	46.65	6.52
Background denitrification rate	ng N g ⁻¹ h ⁻¹	0.67	18.95	2.99	0.90
N ₂ O production rate	ng N g ⁻¹ h ⁻¹	0.05	0.71	0.27	0.04
Relative N ₂ O production		0.02	0.62	0.17	0.03
Water quality					
Cond	μS cm ⁻¹	127.67	359.67	252.71	16.76
ORP	mv	-80.67	153.33	101.16	11.90
DO	mg L ⁻¹	4.51	8.13	5.98	0.27
NO ₃ ⁻	mg L ⁻¹	0.09	1.50	0.53	0.08
NH ₄ ⁺	mg L ⁻¹	0.02	0.50	0.19	0.02
TN	mg L ⁻¹	0.31	2.87	1.10	0.17
TC	mg L ⁻¹	8.47	31.40	19.45	1.26
TOC	mg L ⁻¹	2.47	6.60	4.24	0.29
Sediment characteristics					
pH		5.52	8.19	7.21	0.18
Bulk density	g cm ⁻³	0.64	0.94	0.86	0.01
Moisture	%	26.58	71.33	60.76	2.10
STN	mg g ⁻¹	0.18	1.74	0.51	0.10
STC	mg g ⁻¹	5.39	53.88	25.18	2.91
Plant community structure					
Species richness		0.00	1.00	0.15	0.07
Fresh biomass	g m ⁻²	0.00	2438.10	465.67	190.87
Watershed land uses					
Percent of vegetation	%	1.35	56.47	22.05	3.16
Percent of water body	%	8.85	42.51	16.95	1.93
Percent of HDL	%	33.60	80.00	60.91	2.47
NDVI		0.34	0.58	0.47	0.02

HDL: Human-dominated land uses;
NDVI: Normalised difference vegetation index.

Human land uses
enhance sediment
denitrification

W. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Table 2.** Pearson correlation coefficients between watershed land uses and within-lake characteristics ($N = 20$).

	Vegetation (%)	Watershed land uses		
		Water body (%)	HDL (%)	NDVI
Water quality				
Cond	−0.27	0.17	0.33	−0.49 ^a
ORP	0.07	0.07	−0.11	0.02
DO	0.13	0.30	−0.45 ^a	−0.11
NO ₃ [−]	−0.46 ^a	−0.11	0.72 ^b	−0.18
NH ₄ ⁺	−0.61 ^b	0.38	0.59 ^b	−0.48 ^a
TN	−0.47 ^a	−0.13	0.75 ^b	−0.17
TC	−0.17	0.26	0.08	−0.31
TOC	−0.34	0.10	0.38	−0.09
Sediment characteristics				
pH	−0.02	0.50 ^a	−0.34	−0.48 ^a
Bulk density	−0.06	0.17	−0.03	0.01
Moisture	0.11	0.10	−0.15	0.10
STN	−0.61 ^b	0.47 ^a	0.46 ^a	−0.30
STC	0.03	0.06	−0.06	−0.14
Plant community structure				
Species richness	−0.01	0.34	−0.31	−0.18
Fresh biomass	0.09	0.12	−0.26	−0.01

^a Significant at the 0.05 level;^b Significant at the 0.01 level.

HDL: Human-dominated land uses;

NDVI: Normalised difference vegetation index.

Human land uses
enhance sediment
denitrification

W. Liu et al.

Table 3. Pearson correlation coefficients between sediment denitrification variables and within-lake characteristics ($N = 20$).

	Potential denitrification rate	Background denitrification rate	N ₂ O production rate	Relative N ₂ O production
Water quality				
Cond	0.32	−0.01	0.01	0.04
ORP	−0.10	−0.60 ^b	−0.50 ^a	0.37
DO	−0.48 ^a	−0.52 ^a	−0.56 ^a	0.15
NO ₃ [−]	0.34	0.79 ^b	0.62 ^a	−0.32
NH ₄ ⁺	0.05	0.38	0.03	−0.45 ^a
TN	0.23	0.72 ^b	0.56 ^a	−0.33
TC	0.36	−0.02	0.12	0.32
TOC	0.32	0.16	0.29	0.14
Sediment characteristics				
pH	−0.01	−0.31	−0.35	0.26
Bulk density	0.30	0.16	0.12	−0.05
Moisture	0.39	0.01	0.10	0.11
STN	0.18	0.30	−0.11	−0.36
STC	0.38	−0.07	−0.08	0.14
Plant community structure				
Species richness	−0.28	−0.34	−0.33	0.10
Fresh biomass	−0.32	−0.17	−0.10	0.05

^a Significant at the 0.05 level;^b Significant at the 0.01 level.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Human land uses
enhance sediment
denitrification

W. Liu et al.

Table 4. Relative effects of human-dominated land uses on sediment denitrification in lakes mediated by lake water quality and sediment characteristics ($N = 20$).

	Potential denitrification rate		Background denitrification rate		N ₂ O production rate		Relative N ₂ O production	
	Indirect effects	Contribution (%)	Indirect effects	Contribution (%)	Indirect effects	Contribution (%)	Indirect effects	Contribution (%)
Water quality								
ORP	−0.07	17.50	0.04	7.02	0.09	13.64	−0.08	25.00
DO	0.21	52.50	−0.01	1.75	0.11	16.67	0.03	9.38
NO ₃ [−]	0.06	15.00	0.52	91.23	0.29	43.94	−0.09	28.13
Sediment characteristics								
STN	0.06	15.00	0.00	0.00	−0.17	25.76	−0.12	37.50
Total Indirect effects	0.26	100	0.55	100	0.32	100	−0.26	100

Note: indirect effects were calculated by multiplying the standardized path coefficients of the direct effects of the variables involved in the total pathway. Percentage of contribution was calculated using the sum of absolute values for contributing effects.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Human land uses enhance sediment denitrification

W. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Figure 1. Locations of the Yangtze River basin and 20 studied lakes in China.




Figure 2. Relationships between background denitrification rate and N₂O production rate of the Yangtze lakes ($N = 20$).

Human land uses
enhance sediment
denitrification

W. Liu et al.




Figure 3. Relationships between watershed land uses and sediment denitrification and between watershed land uses and N₂O production of the Yangtze lakes ($N = 20$).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Human land uses enhance sediment denitrification

W. Liu et al.




Figure 4. Structural equation models depicting the indirect effects of human-dominated land uses (HDL) in watersheds on the sediment denitrification and N_2O production of the Yangtze lakes ($N = 20$). Solid lines indicate significant ($P < 0.05$) positive or negative effects, and dashed lines indicate insignificant positive or negative effects. Numbers adjacent to the lines are standardised path coefficients.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

