

This discussion paper is/has been under review for the journal Biogeosciences (BG).
Please refer to the corresponding final paper in BG if available.

Calculating the global contribution of coralline algae to carbon burial

L. H. van der Heijden^{1,2} and N. A. Kamenos¹

¹School of Geographical and Earth Sciences, University of Glasgow, Glasgow, Scotland

²Department of Earth Sciences, Faculty of Geosciences, Utrecht University,
Utrecht, the Netherlands

Received: 7 April 2015 – Accepted: 3 May 2015 – Published: 26 May 2015

Correspondence to: L. H. van der Heijden (luukvdheijden88@gmail.com)

Published by Copernicus Publications on behalf of the European Geosciences Union.

BGD

12, 7845–7877, 2015

Calculating the
global contribution of
coralline algae to
carbon burial

L. H. van der Heijden and
N. A. Kamenos

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Abstract

The ongoing increase in anthropogenic carbon dioxide (CO_2) emissions is changing the global marine environment and is causing warming and acidification of the oceans. Reduction of CO_2 to a sustainable level is required to avoid further marine change.

- 5 Many studies investigate the potential of marine carbon sinks (e.g. seagrass) to mitigate anthropogenic emissions, however, information on storage by coralline algae and the beds they create is scant. Calcifying photosynthetic organisms, including coralline algae, can act as a CO_2 sink via photosynthesis and CaCO_3 dissolution and act as a CO_2 source during respiration and CaCO_3 production on short-term time scales. Long-
10 term carbon storage potential might come from the accumulation of coralline algae deposits over geological time scales. Here, the carbon storage potential of coralline algae is assessed using meta-analysis of their global organic and inorganic carbon production and the processes involved in this metabolism. Organic and inorganic production were estimated at $330 \text{ g C m}^{-2} \text{ yr}^{-1}$ and $880 \text{ g CaCO}_3 \text{ m}^{-2} \text{ yr}^{-1}$ respectively giving
15 global organic/inorganic C production of $0.7/1.8 \times 10^9 \text{ t C yr}^{-1}$. Calcium carbonate production by free-living/crustose coralline algae (CCA) corresponded to a sediment accretion of $70/450 \text{ mm kyr}^{-1}$. Using this potential carbon storage by coralline algae, the global production of free-living algae/CCA was $0.4/1.2 \times 10^9 \text{ t C yr}^{-1}$ suggesting a total potential carbon sink of $1.6 \times 10^9 \text{ t C yr}^{-1}$. Coralline algae therefore have production
20 rates similar to mangroves, saltmarshes and seagrasses representing an as yet unquantified but significant carbon store, however, further empirical investigations are needed to determine the dynamics and stability of that store.

1 Carbon storage and coralline algae

An increase in exploitation of fossil fuels since the mid-18th century caused a rise in the partial pressure of carbon dioxide in both atmospheric (CO_2) and oceanic ($p\text{CO}_2$) reservoirs (Sabine et al., 2004; Meehl, 2007). Atmospheric CO_2 has risen from
25

BGD

12, 7845–7877, 2015

Calculating the global contribution of coralline algae to carbon burial

L. H. van der Heijden and N. A. Kamenos

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



280 ppm in 1750 to nearly 380 ppm in 2005 at a rate unprecedented in geological history (Denman and Brasseur, 2007). The marine environment has been changing rapidly in the last few centuries too (Cubasch et al., 2013), with increasing CO₂ causing warming and acidification of the Earth's oceans (Caldeira and Wickett, 2005).

5 Concentrations of atmospheric CO₂ simulated by coupled climate-carbon cycle models range between 730 and 1200 ppm by 2100 (Meehl, 2007). Therefore, a reduction of atmospheric CO₂ to a sustainable level is needed to avoid further environmental damage (Collins et al., 2013; Kirtman et al., 2013).

The oceans are a major sink of anthropogenic CO₂ emissions, accounting for ~ 48 %
10 of emissions absorption since the Industrial Revolution (Sabine et al., 2004). Significantly, around 50 % of the global primary production (which uses pCO₂) is by marine organisms (Beardall and Raven, 2004) with marine microalgae and bacteria being the dominant source of primary production and respiration (Duarte and Cebrian, 1996; del Giorgio and Duarte, 2002; Duarte et al., 2005). Vegetated marine habitats, including
15 macroalgae and seagrasses, are often neglected from accounts of the global ocean carbon cycle because of their limited extent (< 2 %) (Duarte and Cebrian, 1996). However, vegetated coastal habitats have a great carbon storage capacity (Duarte et al., 2005) and the potential of marine coastal vegetation as a sink for anthropogenic carbon emissions (blue carbon) is becoming of interest (Nellemann et al., 2009). These marine macrophyte ecosystems have slow turnover rates and are therefore more effective
20 carbon sinks than planktonic ecosystems (Smith, 1981).

Red coralline algae are present from the tropics to polar regions (Johansen, 1981; Steneck, 1986; Foster, 2001; Wilson, 2004). Coralline algae are important for ecosystems due to their role in carbon cycling, creating and maintaining habitats, and reef
25 building/structuring roles (Nelson, 2009). They are divided in two morpho-functional groups; geniculated (articulated) and non-geniculated (non-articulated) (Johansen, 1981). The morphological states range from totally adherent to having nonadherent margins (leafy) to totally nonadherent (free-living, e.g. rhodoliths, maerls and nodules) (Steneck, 1986; Cabioch, 1988). The calcium carbonate skeleton of coralline algae pre-

Calculating the global contribution of coralline algae to carbon burial

L. H. van der Heijden and
N. A. Kamenos

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

vents them from breaking down quickly compared to fleshy algae (Borowitzka, 1982; Wilson, 2004). Coralline algal species have been observed in the fossil record since the early Cretaceous (Aguirre et al., 2000) and coralline algal communities reach 500–800 years (Adey and Macintyre, 1973; Kamenos, 2010) with ~ 8000 year old free-living 5 coralline algal beds present in France (Birkett et al., 1998).

Coralline algae are important contributors to the marine calcium carbonate (CaCO_3) deposited in the coral reef sediments (Goreau, 1963; Adey and Macintyre, 1973) and account for approximately 25 % of CaCO_3 accumulation within coastal regions (Martin et al., 2007). Calcifying photosynthesisers are both a sink and a source of CO_2 10 (Frankignoulle, 1994). Coralline algae act as a CO_2 sink in the processes of photosynthesis and CaCO_3 dissolution and act as a CO_2 source in the processes of respiration and CaCO_3 production (Martin et al., 2005, 2006, 2007, 2013; Barron et al., 2006; Kamenos et al., 2013). We aim to estimate the global distribution of coralline algae, and from that, determine their potential role in long-term carbon burial.

15 2 Coralline algal succession and small-scale distribution

The distribution and abundance of coralline algae is determined by ecological processes including growth, succession and competition (Steneck, 1986; McCoy and Kamenos, 2015) as well by environmental conditions such as disturbance, temperature and irradiance (Adey and Macintyre, 1973; Kamenos et al., 2004; Gattuso et al., 2006).

20 Coralline algae grow both laterally to increase area and vertically to increase thickness (Steneck, 1986). Coralline algal vertical accretion rates vary widely from 0.1 to 80 mm yr^{-1} (Adey and McKibbin, 1970; Steneck and Adey, 1976; Edyvean and Ford, 1987). Succession in coralline algae is for thick and/or branched crusts to replace thinner unbranched crusts (Adey and Vassar, 1975; Steneck, 1986). Succession seems 25 most rapid in the tropics, where colonization and succession takes ~ 1 year, compared to 6–7 years in the boreal North Pacific and > 10 years in the subarctic North Atlantic (Steneck, 1986; McCoy and Ragazzola, 2014). In shallow productive zones coralline al-

Calculating the global contribution of coralline algae to carbon burial

L. H. van der Heijden and N. A. Kamenos

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



5 gae require disturbances, mainly herbivory as well as water motion, to remain clear of fleshy algae and invertebrates (Steneck, 1986). However, towed fishing gear (e.g. trawling) can easily damage rhodoliths (maerl) (Hall-Spencer and Moore, 2000; Kamenos and Moore, 2003). Overall, coralline algal distribution is likely primarily determined by
irradiance and temperature (Adey and McKibbin, 1970; Adey and Adey, 1973; Gattuso et al., 2006).

2.1 Global distribution

10 Coralline algae are ecosystem engineers (Nelson, 2009), major framework builders and carbonate producers, especially in temperate and cold water benthic ecosystems (Nelson, 1988; Freiwald and Henrich, 1994; Foster, 2001; Gherardi, 2004; Bracchi and Basso, 2012; Savini et al., 2012; Basso, 2012) resulting in a wide distribution. Coralline algae are found from the low intertidal to the infralittoral and circalittoral zones (> 200 m depth) (Steneck, 1986; Basso, 1998; Foster, 2001) and have a worldwide spatial distribution (Fig. 1; Supplement, Table S2). While crustose coralline algae (CCA) grow exclusively on hard surfaces, free-living coralline algae are able to form rhodoliths when they settle on non-cohesive particulate substrates or are detached from existing hard substrates by fragmentation (Bosence, 1983).

2.2 Surface covered by coralline algae

20 The surface of the coastal zone covered by coralline algae varies spatiotemporally and differs for free-living algae, geniculate and CCA (Supplement, Table S1). The average coralline algal sea bed coverage from published studies is 52.5 % for CCA, 45.0 % for rhodoliths and 45.0 % for coralline algae overall. Figueiredo et al. (2008) indicate that the surface covered by CCA on the Abrolhos Bank ($20\,900\text{ km}^2$) in Brazil ranges from 5–40 % on the reef flats, 30–80 % on the reef crests and 10–50 % on the reef walls. Coverage varies due to differences in the abundance of turf algae and herbivory pressure (Figueiredo et al., 2008). On coral reefs CCA (e.g. *Porolithon onkodes*) can

Calculating the global contribution of coralline algae to carbon burial

L. H. van der Heijden and N. A. Kamenos

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)


cover ~ 40 % of the reef slope (Littler and Doty, 1975; Stearn et al., 1977), 60 % of the reef flat and 5 % of lagoon sites (Atkinson and Grigg, 1984) with rhodoliths covering up 90 % of the reef crest (Sheveiko, 1981) and 90 % of the seaward shallow reef slope (Chisholm, 1988). Importantly, the area covered by coralline algae is not necessarily lower in regions dominated by other algal forms, because of their ability to occur on the primary substratum (up to 90 %) or as epiphytes on larger algae (Johansen, 1981).

3 Processes and metabolism

While coralline algae are slow growing their high abundance and spatial distribution indicates their production is likely important (Johansen, 1981) and they are major contributors to the carbon- and carbonate cycles of coastal environments (Martin et al., 2013). Organic production relates to the photosynthetic capacity of coralline algae, while inorganic production relates to the calcium carbonate production (Johansen, 1981).

3.1 Organic production

Organic production of coralline algae is low compared to other marine plants (Johansen, 1981; Steneck, 1986). However, because of their high abundance and worldwide dispersal, corallines can contribute significantly to the total marine primary production (Roberts et al., 2002). Production of 1 mole of organic material (photosynthesis) decreases dissolved inorganic carbon (DIC) by approximately 1 mole:



Primary production also decreases $p\text{CO}_2$, however the magnitude of these changes depends on the equilibrium constants (Johansen, 1981). Respiration increases both DIC and $p\text{CO}_2$ (Johansen, 1981). Coralline algal respiration is between 20–60 % of gross primary production (Marsh, 1970; Littler, 1973; Littler and Murray, 1974; Souria, 1976; Wanders, 1976). Net community production for coralline algae is induced

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	

or limited by environmental parameters including light reaching the communities (Gattuso et al., 2006; Martin et al., 2006; Burdett et al., 2014), temperature (Martin et al., 2006; Kamenos and Law, 2010) and nutrient availability (Smith et al., 2001). For example, Chisholm (2003) suggested that the high rates of productivity measured in situ at Lizard Island, Australia, came from the coralline algae that derive nutrients from the underlying reef.

3.2 Inorganic production and accumulation

Photosynthesis also plays a crucial role in the production of inorganic material as it creates the environment in which calcification occurs (Johansen, 1981). The ratio of inorganic–organic production is high in coralline algae, compared to non-coralline seaweeds (Johansen, 1981). Precipitation of CaCO_3 decreases DIC by 1 mole and total alkalinity by two equivalents for each mole precipitated:



For calcium carbonate to be deposited an alkaline environment is required, as well as high concentrations of calcium and carbonate (Johansen, 1981). Calcification of coralline algae occurs internally, compared to external calcification in corals and other invertebrates (Chisholm, 2003). The cell-walls of coralline algae are composed of calcium carbonate, and are mainly consist of high Mg-calcite (HMC: > 4 % wt of MgCO_3) (Moberly, 1968; Kamenos et al., 2008; Basso, 2012). The preservation of shallow-water carbonates is about 60 % and even up to 70–80 % if export is not taken into account (Milliman, 1993).

Coralligenous algal-dominated rocky bottoms and rhodolith beds are among the highest algal carbonate producers when compared with *Posidonia oceanica* meadows, sandy bottom communities, *Caulerpa-Cymodocea* meadows, coralligenous animal-dominated, photophilic algae and hemisciaephili algal communities (Canals and Ballessteros, 1997). The quantity of calcite production by coralline algae depends on their morphology (e.g., geniculate or non-geniculate, thick or thin crusts), growth rate and

the environmental conditions (Basso, 2012). Calcification is indirectly limited by temperature where warmer waters are supersaturated with CaCO_3 (Littler, 1976) often over a season cycle (Martin et al., 2006). Martin et al. (2006) also observed a light limitation for the calcification of coralline algae.

5 4 Potential global contribution of coralline algae to carbon burial

The shallow-water ocean environment (i.e. bays, estuaries, lagoons, banks, and continental shelves) accounts for 14–30 % of the oceanic primary production, 80 % of organic material burial and ~ 50 % of CaCO_3 deposition (Gattuso et al., 1998). The total surface area of the coastal zone, thus the potential habitat for benthic coralline algae, is estimated between $0.45\text{--}49.4 \times 10^{12} \text{ m}^2$ (Ryther, 1969; Koblentz-Mishke et al., 1970; Whittaker and Likens, 1973; Bunt, 1975; Platt and Subba Rao, 1975; Smith, 1981). The coastal area, that has depths ranging between 0 and 200 m covers 7.49 % of the world ocean, which correspond to $27.123 \times 10^{12} \text{ m}^2$ (Menard and Smith, 1966). Charpy-Roubaud and Sournia (1990) suggest an area of $6.8 \times 10^{12} \text{ m}^2$, because the average benthic photic zone of the world is smaller than 200 m. Here we will use 33 % of the coastal zone which is the part of that receives enough light for photosynthesis (Gattuso et al., 2006) and thus assuming the production mainly occurs in the top 66 m of the coastal zone. Because coralline algae usually attach to harder substrata (Bosence, 1983) the surface covered by coralline algae (Supplement, Table S1) has to be taken into account. However, as there are substrates that have 0 % surface coverage of coralline algae (e.g. on sandy substrata), to be conservative, we have assumed only half of the estimated surface coverage percentages estimated above contain coralline algae (CCA = 26.25 %, rhodoliths = 22.5 %, coralline algae = 22.5 %). At present we have an incomplete knowledge of the real distribution of coralline algae, so we estimate a global production based on the following parameters: the production of coralline algae (median), the top 66 m global coastal zone and the surface of this coastal zone

BGD

12, 7845–7877, 2015

Calculating the global contribution of coralline algae to carbon burial

L. H. van der Heijden and
N. A. Kamenos

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	

covered by coralline algae (22.5 %). We use the median in/organic C production for coralline algae due to skewed data distribution (Zar, 1999) across available studies.

4.1 Global coralline algal organic C production

Primary production by coralline algae ranges widely from $10\text{ g C m}^{-2}\text{ yr}^{-1}$ by *Lithothamnion coralloides* in the Bay of Brest, France (Martin et al., 2006) to $2391\text{ g C m}^{-2}\text{ yr}^{-1}$ by *Hydrolithon onkodes* at Lizard Island, Australia (Chisholm, 2003), giving a median production of $329\text{ g C m}^{-2}\text{ yr}^{-1}$ ($n = 39$) (Table 1) across depths and locations. Global C production may thus be as high as $0.7 \times 10^9\text{ t C yr}^{-1}$. The daily production of coralline algae corresponds with the range of production of benthic fleshy algae, turf algae, sand algae, phytoplankton, seagrasses and zooxanthellae (Table 2) and estimated yearly coralline algal production rate ($329\text{ g C m}^{-2}\text{ yr}^{-1}$) is in the range of production by mangroves, salt marshes and seagrasses and appears more productive than coastal phytoplankton, benthic diatoms and coral reefs (Table 2). Payri (2000) observed that the annual production of a coralline algal communities corresponds to approximately one third of the production of seagrass beds, which was also observed on the west-coast of France with a production ratio of 3.12 (Martin et al., 2005). A production ratio of 1.5–3.7 is observed in this study when compared to seagrass production rate studies (Table 2).

The estimated production of free-living coralline algae ($0.35 \times 10^9\text{ t C yr}^{-1}$) is in the range determined by other studies while the production for CCA ($0.88 \times 10^9\text{ t C yr}^{-1}$) is slightly higher (Table 3). Thus, with a global oceanic production estimated at $48.5 \times 10^9\text{ t C yr}^{-1}$ (Field et al., 1998) coralline algal production represent a measurable component.

4.2 Global inorganic coralline algal C production and accumulation

Studies focusing on coralline algae and calcium carbonate indicate a production range of $8\text{--}7400\text{ g CaCO}_3\text{ m}^{-2}\text{ yr}^{-1}$ and a median of $880\text{ g CaCO}_3\text{ m}^{-2}\text{ yr}^{-1}$ (Table 4). The global calcium carbonate production using the previously estimated surface coverage

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

was $1.8 \times 10^9 \text{ tCaCO}_3 \text{ yr}^{-1}$ for coralline algae. Thus CaCO_3 production by coralline algae of $880 \text{ g CaCO}_3 \text{ m}^{-2} \text{ yr}^{-1}$ lies within the range of coral reef calcite production of $75\text{--}4000 \text{ g CaCO}_3 \text{ m}^{-2} \text{ yr}^{-1}$ (Canals and Ballesteros, 1997) and is comparable with the production rate in the Late Holocene ocean for coral reefs ($1500 \text{ g CaCO}_3 \text{ m}^{-2} \text{ yr}^{-1}$ (Milliman, 1993)). Basso (2012) estimated an average production rate of $5 \text{ g CaCO}_3 \text{ m}^{-2} \text{ yr}^{-1}$ for the coralline algae in the Mediterranean sea, however this included coralline algae occurring below 100 m. Gattuso et al. (1998) suggested that communities in the coastal zone are responsible for more than 40 % ($23 \times 10^9 \text{ tCaCO}_3 \text{ yr}^{-1}$) of the total marine calcium carbonate production. Thus the estimated calcite production by coralline algae is similar to the production of other coastal communities (e.g. coral reefs, banks and non/carbonate shelves) and might represent a large fraction of the coastal- and total ocean calcite production (Gattuso et al., 1998).

Using average production rates for free-living algae and CCA an inorganic production was estimated for these two groups. The inorganic production for free-living algae was $22 \text{ g C-inorganic m}^{-2} \text{ yr}^{-1}$ and $147 \text{ g C-inorganic m}^{-2} \text{ yr}^{-1}$ for CCA. Thus inorganic production by coralline algae of $105 \text{ g C-inorganic m}^{-2} \text{ yr}^{-1}$ and organic production of $330 \text{ g C-organic m}^{-2} \text{ yr}^{-1}$ gives a PIC : POC ratio of 0.32 (PIC is the particular inorganic carbon and POC the particular organic carbon). The PIC : POC ratio for free-living algae was 0.13 and 0.40 for the CCA. Significantly, a similar PIC : POC range of ratios of 0.23–0.29 was also observed for coccolithophores (Engel et al., 2005).

4.3 Global carbon accumulation

The long-term removal of C requires the fixed carbon to remain stored for 100–1000 years (Gattuso et al., 1998). The global long-term deposition rate of free-living coralline algae is 500 mm kyr^{-1} (Table 5) and the accumulation rates range from 80 to 1400 mm kyr^{-1} for temperate (Orkney Island, Scotland) to polar (Tromsø district, Norway). The calcium carbonate production by free-living algae ($187 \text{ g CaCO}_3 \text{ m}^{-2} \text{ yr}^{-1}$) with a calcite density of 2.71 g cm^{-3} (DeFoe and Compton, 1925) corresponds to a sed-

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

iment accretion of 70 mm kyr⁻¹, while for CCA this corresponds to a sediment accretion of 450 mm kyr⁻¹. Given the accretion rate of 500 mm kyr⁻¹, the preservation potential of coralline algae would be 64 %. This is consistent with the empirically calculated calcium carbonate preservation of 60 % (Milliman, 1993). However, if the preservation of CCA is excluded because of the lack of available accretion rates, and heavy grazing (Steneck, 1986), the preservation potential for this morphotype would be 14 %. As the complete preservation potential for coralline algae still required further refining, the potential carbon burial is estimated based on the sum of total organic production and the inorganic production. The estimated potential burial for the free-living algae was 0.4×10^9 and 1.2×10^9 tCyr⁻¹ for CCA giving a potential carbon burial of 1.6×10^9 tCyr⁻¹ for coralline algae.

5 Future prospects: ocean acidification and rising temperature

Coralline algae may be vulnerable to the warming and lowering pH of sea water, caused by the currently occurring increase in anthropogenic CO₂ (Kleypas et al., 2006; McCoy and Kamenos, 2015). High pCO₂ conditions negatively affect the community growth (Jokiel et al., 2008), recruitment (Kuffner et al., 2008), calcification (Anthony et al., 2008) and area covered by coralline algae (Kuffner et al., 2008). The high-Mg calcite (HMC) cell-walls of coralline algae, containing 7.7–28.8 % MgCO₃, play a crucial role in their response to the risen temperature and acidification of seawater (Basso, 2012; Kamenos et al., 2013). Biogenic HMC cell-walls, containing > 8–12 % MgCO₃, have a high solubility and sensitive response to ocean acidification (Andersson et al., 2008). The decreasing abundance and growth of coralline algae could have dramatic consequences for worldwide coastal ecosystems (Johansen, 1981; Martin and Gattuso, 2009; Basso, 2012). However, there is evidence that coralline algae will continue to calcify with increasing pCO₂ due to their high structural plasticity, but may be structurally weakened (Kamenos et al., 2013; Ragazzola et al., 2013). Additionally, the ongoing rise in atmospheric pCO₂ is expected to have a positive impact on the organic production

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

and growth of algae due to increased $p\text{CO}_2$ availability (Hendriks et al., 2010). For example, Semesi et al. (2009) observed an increase in photosynthetic rates of coralline algae with a rising $p\text{CO}_2$ of seawater, however, whether this also translates to their accretion at longer time scales is still not clear.

5 6 Conclusions

The ongoing increase of anthropogenic CO_2 is causing warming and acidification of world's ocean. Reduction of CO_2 to a sustainable level is required to avoid further environmental damage and various solutions have already been proposed. We calculate coralline algae to have a global average primary of $0.7 \times 10^9 \text{ tCyr}^{-1}$ and an estimated 10 total global CaCO_3 production of $1.8 \times 10^9 \text{ tCaCO}_3 \text{ yr}^{-1}$ which corresponds to an inorganic production of $0.2 \times 10^9 \text{ tinorganicCyr}^{-1}$. With their substantial preservation potential and the longevity of the deposits they create, coralline algae have a significant capacity to store carbon. However, we are still uncertain of the impact future global change is likely to have on that capacity. Given their storage potential, empirical studies 15 are now needed to refine the calculations.

**The Supplement related to this article is available online at
doi:10.5194/bgd-12-7845-2015-supplement.**

Acknowledgements. Nicholas A. Kamenos was in receipt of NERC (UK) grant (NE/H017305/1). This paper is a contribution to the UK Ocean Acidification Research Programme. This review 20 profited greatly from discussions with Appy Sluijs and Jack Middelburg and their constructive feedback and criticism of the thesis. We are grateful to both of them.

BGD

12, 7845–7877, 2015

Calculating the
global contribution of
coralline algae to
carbon burial

L. H. van der Heijden and
N. A. Kamenos

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	

References

Calculating the global contribution of coralline algae to carbon burial

L. H. van der Heijden and N. A. Kamenos

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Calculating the global contribution of coralline algae to carbon burial

L. H. van der Heijden and
N. A. Kamenos

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

1

1

▶

Back

Close

Full Screen / Esc

Printer friendly Version

Interactive Discussion



Calculating the global contribution of coralline algae to carbon burial

L. H. van der Heijden and
N. A. Kamenos

- Cabioch, J.: Morphogenesis and generic concepts in coralline algae – a reappraisal, *Helgolander Meeresun.*, 42, 493–509, doi:10.1007/BF02365623, 1988.
- Cadee, G. C. and Hegeman, J.: Primary production of the benthic microflora living on tidal flats in the Dutch Wadden Sea, *Netherlands J. Sea Res.*, 8, 260–291, 1974.
- Caldeira, K. and Wickett, M. E.: Ocean model predictions of chemistry changes from carbon dioxide emissions to the atmosphere and ocean, *J. Geophys. Res.*, 110, 1–12, doi:10.1029/2004JC002671, 2005.
- Canals, M. and Ballesteros, E.: Production of carbonate particles by phytobenthic communities on the Mallorca-Menorca shelf, northwestern Mediterranean Sea, *Deep-Sea Res. Pt. II*, 44, 611–629, doi:10.1016/S0967-0645(96)00095-1, 1997.
- Charpy-Roubaud, C. and Sournia, A.: The comparative estimation of phytoplanktonic, micro-phytobenthic and macrophytobenthic primary production in the oceans, *Mar. Microb. Food Webs*, 4, 31–57, 1990.
- Chisholm, J. R. M.: Photosynthesis, calcification, and photoadaptation in reef-building crustose coralline algae on the Great Barrier Reef, PhD thesis, James Cook Univ., available at: <http://researchonline.jcu.edu.au/27501/>, 1988.
- Chisholm, J. R. M.: Calcification by crustose coralline algae on the northern Great Barrier Reef, Australia, *Limnol. Oceanogr.*, 45, 1476–1484, doi:10.4319/lo.2000.45.7.1476, 2000.
- Chisholm, J. R. M.: Primary productivity of reef-building crustose coralline algae, *Limnol. Oceanogr.*, 48, 1376–1387, 2003.
- Clarke, L., Jiang, K., Akimoto, K., Babiker, M., Blanford, G., Fisher-Vanden, K., Hourcade, J.-C., Krey, V., Kriegler, E., Loscher, A., McCollum, D., Paltsev, S., Rose, S., Shukla, P. R., Tavoni, M., van der Zwaan, B., van Vuuren, D. P., Bottcher, H., Calvin, K., Daenzer, K., den Elzen, M., Dhar, S., Eom, J., Hoeller, S., Hohne, N., Hultman, N., Irvine, P., Jewell, J., Johnson, N., Kanudia, A., Kelemen, A., Keller, K., Kolp, P., Lawrence, M., Longden, T., Lowe, J., Lucena, A., Luderer, G., Marangoni, G., Moore, N., Mouratiadou, I., Petermann, N., Rasch, P., Riahi, K., Rogelj, J., Schaeffer, M., Schafer, S., Sedlacek, J., Sokka, L., von Storchow, C., Wing, I. S., Vaughan, N., Wiertz, T., Zwickel, T., Chen, W., and Weyant, J.: Assessing transformation pathways, in: *Climate Change 2013: Mitigation of Climate Change. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Edenhofer, O., Pichs-Madruga, R., Sokona, Y., Minx, J. C., Farahani, E., Kadner, S., Seyboth, K., Adler, A., Baum, I., Brunner, S., Eickemeier, P.,

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Calculating the
global contribution of
coralline algae to
carbon burial

L. H. van der Heijden and
N. A. Kamenos

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
	
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	

- Kriemann, B., Savolainen, J., Schlömer, S., von Stechow, C., and Zwickel, T., Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 1–141, 2013.
- Collins, M., Knutti, R., Arblaster, J., Dufresne, J.-L., Fichefet, T., Friedlingstein, P., Gao, X., Gutowski Jr, W. J., Johns, T., Krinner, G., Shongwe, M., Tebaldi, C., Weaver, A. J., and Wehner, M.: Long-term climate change: projections, commitments and irreversibility, in: Climate Change 2013: The Physical Basis, Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, edited by: Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M., Cambridge University Press, Cambridge, UK, New York, NY, USA, 1031–1106, 2013.
- Cubasch, U., Wuebbles, D., Chen, D., Facchini, M. C., Frame, D., Mahowald, N., and Winther, J.-G.: Introduction, in: Climate Change 2013: The Physical Basis, Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, edited by: Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M., Cambridge University Press, Cambridge, UK, New York, NY, USA, 121–155, 2013.
- Cumani, F., Bradassi, F., Di Pascoli, A., and Bressan, G.: Marine acidification effects on reproduction and growth rates of Corallinaceae spores (Rhodophyta), *Rapp. Comm. Int. Mer Medit.*, 39, 735, 2010.
- DeFoe, O. K. and Compton, A. H.: The density of rock salt and calcite, *Phys. Rev.*, 25, 5, 1925.
- Del Giorgio, P. and Duarte, C. M.: Respiration in the open ocean, *Nature*, 420, 379–384, doi:10.1038/nature01165, 2002.
- Denman, K. L., Brasseur, G., Chidtaisong, A., Clais, P., Cox, P. M., Dickinson, R. E., Hauglustaine, D., Heinze, C., Holland, E., Jacob, D., Lohmann, U., Ramachandran, S., da Silva Dias, P. L., Wofsy, S. C., and Zhang, X.: Couplings between changes in the climate system and biogeochemistry, in: Climate Change 2007: The Physical Science Basis, Contributing to Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change, edited by: Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K. B., Tignor, M., and Miller, H. L., Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 501–568, 2007.
- Duarte, C. M. and Cebrian, J.: The fate of marine autotrophic production, *Limnol. Oceanogr.*, 41, 1758–1766, 1996.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	

- Duarte, C. M., Middelburg, J. J., and Caraco, N.: Major role of marine vegetation on the oceanic carbon cycle, *Biogeosciences*, 2, 1–8, doi:10.5194/bg-2-1-2005, 2005.
- Eakin, C. M.: Where have all the carbonates gone? A model comparison of calcium carbonate budgets before and after the 1982–1983 El Niño at Uva Island in the eastern Pacific, *Coral Reefs*, 15, 109–119, 1996.
- Edyvean, R. G. J. and Ford, H.: Growth rates of *Lithophyllum incrustans* (Corallinales, Rhodophyta) from South West Wales, *Brit. Phycol. J.*, 22, 139–146, 1987.
- El Haïkali, B., Bensoussan, N., Romano, J.-C., and Bousquet, V.: Estimation of photosynthesis and calcification rates of *Corallina elongata* Ellis and Solander, 1786, by measurements of dissolved oxygen, pH and total alkalinity, *Sci. Mar.*, 68, 45–56, 2004.
- Engel, A., Zondervan, I., Aerts, K., Beaufort, L., Benthién, A., Chou, L., Delille, B., Gattuso, J.-P., Harlay, J., Heemann, C., Hoffmann, L., Nejstgaard, J., Pizay, M.-D., Rochelle-Newall, E., Schneider, U., Terbrueggen, A., and Riebesell, U.: Testing the direct effect of CO₂ concentration on a bloom of the coccolithophorid *Emiliania huxleyi* in mesocosm experiments, *Limnol. Oceanogr.*, 50, 493–507, 2005.
- Farrow, G. E., Allen, N. H., and Ekpan, E. B.: Bioclastic carbonate sedimentation on a high-latitude, tide-dominated shelf: northeast Orkney Islands, Scotland, *J. Sediment. Petrol.*, 54, 373–393, 1984.
- Field, C. B., Behrenfeld, M. J., Randerson, J. T., and Falkowski, P.: Primary production of the biosphere: integrating terrestrial and oceanic components, *Science*, 281, 237–240, doi:10.1126/science.281.5374.237, 1998.
- Figueiredo, M. A. O., Horta, P. A., Pedrini, A. G., and Nunes, J. M. C.: Benthic marine algae of the coral reefs of Brazil: a literature review, *Oecologia Bras.*, 12, 258–269, 2008.
- Foster, M. S.: Rhodoliths: between rocks and soft places, *J. Phycol.*, 37, 659–667, 2001.
- Frankignoulle, M.: A complete set of buffer factors for acid/base CO₂ system in seawater, *J. Marine Syst.*, 5, 111–118, 1994.
- Freiwald, A.: Modern nearshore cold-temperate calcareous sediments in the Troms districts, northern Norway, *J. Sediment. Res.*, 68, 763–776, 1998.
- Freiwald, A. and Henrich, R.: Reefal corallina algal build-ups within the Arctic Circle: morphology and sedimentary dynamics under extreme environmental seasonality, *Sedimentology*, 41, 963–984, 1994.

Calculating the global contribution of coralline algae to carbon burial

L. H. van der Heijden and
N. A. Kamenos

- Gao, K., Aruga, Y., Asada, K., Ishihara, T., Akano, T., and Kiyohara, M.: Calcification in the articulated coralline alga *Corallinapilulifera*, with special reference to the effect of elevated CO₂ concentration, *Mar. Biol.*, 117, 129–132, 1993.
- Gattuso, J.-P., Frankignoulle, M., and Wollast, R.: Carbon and carbonate deposits, *Annu. Rev. Ecol. Syst.*, 29, 405–434, 1998.
- Gattuso, J.-P., Gentili, B., Duarte, C. M., Kleypas, J. A., Middelburg, J. J., and Antoine, D.: Light availability in the coastal ocean: impact on the distribution of benthic photosynthetic organisms and their contribution to primary production, *Biogeosciences*, 3, 489–513, doi:10.5194/bg-3-489-2006, 2006.
- Gherardi, D. F. M.: Community structure and carbonate production of a temperate Rhodolith Bank from Arvoredo Island, Southern Brazil, *Braz. J. Oceanogr.*, 52, 207–224, 2004.
- Gieskes, W. W. C. and Kraay, G. W.: The phytoplankton spring bloom in Dutch coastal waters of the North Sea, *Netherlands J. Sea Res.*, 9, 166–196, 1975.
- Goreau, T. F.: Calcium carbonate deposition by coralline algae and corals in relation to their roles as reef-builders, *Ann. NY Acad. Sci.*, 109, 127–168, 1963.
- Hall-Spencer, J. M. and Moore, P. G.: Scallop dredging has profound, long-term impacts on maerl habitats, *ICES J. Mar. Sci.*, 57, 1407–1415, doi:10.1006/jmsc.2000.0918, 2000.
- Hall-Spencer, J. M., Rodolfo-Metalpa, R., Martin, S., Ransome, E., Fine, M., Turner, S. M., Rowley, S. J., Tedesco, D., and Buia, M.-C.: Volcanic carbon dioxide vents show ecosystem effects of ocean acidification, *Nature*, 454, 96–99, doi:10.1038/nature07051, 2008.
- Hart, D. E. and Kench, P. S.: Carbonate production of an emergent reef platform, Warraber Island, Torres Strait, Australia, *Coral Reefs*, 26, 53–68, doi:10.1007/s00338-006-0168-8, 2006.
- Hendriks, I. E., Duarte, C. M., and Álvarez, M.: Vulnerability of marine biodiversity to ocean acidification: a meta-analysis, *Estuar. Coast. Shelf S.*, 86, 157–164, doi:10.1016/j.ecss.2009.11.022, 2010.
- Johansen, H. W.: Coralline algae, a first synthesis, Boca Raton, Florida, CRC Press, 1–239, 1981.
- Jokiel, P. L., Rodgers, K. S., Kuffner, I. B., Andersson, A. J., Cox, E. F., and Mackenzie, F. T.: Ocean acidification and calcifying reef organisms: a mesocosm investigation, *Coral Reefs*, 27, 473–483, doi:10.1007/s00338-008-0380-9, 2008.

Discussion Paper | Discussion Paper

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Calculating the global contribution of coralline algae to carbon burial

L. H. van der Heijden and
N. A. Kamenos

- Kamenos, N. A.: North Atlantic summers have warmed more than winters since 1353, and the response of marine zooplankton, *P. Natl. Acad. Sci. USA*, 107, 1–6, doi:10.1073/pnas.1006141107, 2010.
- Kamenos, N. A. and Law, A.: Temperature controls on coralline algal skeletal growth, *J. Phycol.*, 46, 331–335, doi:10.1111/j.1529-8817.2009.00780.x, 2010.
- Kamenos, N. A. and Moore, P. G.: Substratum heterogeneity of dredged vs. un-dredged maerl grounds, *J. Mar. Biol. Assoc. UK*, 83, 411–413, doi:10.1017/S0025315403007264h, 2003.
- Kamenos, N. A., Moore, P. G., and Hall-Spencer, J. M.: Maerl grounds provide both refuge and high growth potential for juvenile queen scallops (*Aequipecten opercularis* L.), *J. Exp. Mar. Biol. Ecol.*, 313, 241–254, doi:10.1016/j.jembe.2004.08.007, 2004.
- Kamenos, N. A., Cusack, M., and Moore, P. G.: Coralline algae are global palaeothermometers with bi-weekly resolution, *Geochim. Cosmochim. Ac.*, 72, 771–779, 2008.
- Kamenos, N. A., Burdett, H. L., Aloisio, E., Findlay, H. S., Martin, S., Longbone, C., Dunn, J., Widdicombe, S., and Calosi, P.: Coralline algal structure is more sensitive to rate, rather than the magnitude, of ocean acidification, *Global Change Biol.*, 19, 3621–3628, doi:10.1111/gcb.12351, 2013.
- Kinsey, D. W.: Metabolism, calcification and carbon production, I: System level studies, *Proc. 5th Int. Coral Reef Symp.*, Tahiti, 27 May–1 June, 5, 505–526, 1985.
- Kirby, C. J. and Gosselink, J. G.: Primary production in a Louisiana Gulf Coast Spartina alterniflora marsh, *Ecology*, 57, 1052–1059, 1976.
- Kirtman, B., Power, S. B., Adedoyin, A. J., Boer, G. J., Bojariu, R., Camilloni, I., Doblas-Reyes, F., Fiore, A. M., Kimoto, M., Meehl, G., Prather, M., Sarr, A., Schar, C., Sutton, R., van Oldenburgh, G. J., Vecchi, G., and Wang, H.-J.: Near-term climate change: projections and predictability, in: *Climate Change 2013: The Physical Basis, Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M., Cambridge University Press, Cambridge, UK, New York, NY, USA, 955–1008, 2013.
- Kleypas, J. A., Feely, R. A. A., Fabry, V. J., Langdon, C., Sabine, C. L., and Robbins, L. L.: Impacts of ocean acidification on coral reefs and other marine calcifiers: a guide for future research, in: *Workshop Report*, St. Petersburg, Florida, USA, 18–20, 2006.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Calculating the global contribution of coralline algae to carbon burial

L. H. van der Heijden and
N. A. Kamenos

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Koblenz-Mishke, O. J., Volkovinsky, V. V., and Kabanova, J. G.: Plankton primary production of the world ocean, in: *Scientific Exploration of the South Pacific*, edited by: Wooster, W. S., National Academy of Science, Washington, 183–193, 1970.

Kuffner, I. B., Andersson, A. J., Jokiel, P. L., Rodgers, K. S., and Mackenzie, F. T.: Decreased abundance of crustose coralline algae due to ocean acidification, *Nat. Geosci.*, 1, 114–117, doi:10.1038/ngeo100, 2008.

Langdon, C., Broecker, W. S., Hammond, D. E., Glenn, E., Fitzsimmons, K., Nelson, S. G., Peng, T.-H., Hajdas, I., and Bonani, G.: Effect of elevated CO₂ on the community metabolism of an experimental coral reef, *Global Biogeochem. Cy.*, 17, 1–14, doi:10.1029/2002GB001941, 2003.

Larkum, A. W. D.: The primary productivity of plant communities on coral reefs, in: *Perspectives on Coral Reefs*, edited by: Barnes, D. J., Australian Institute of Marine Science, Townsville, 221–230, 1983.

Le Quéré, C., Raupach, M. R., Canadell, J. G., Marland, G., Le Quéré, C., Marland, G., Bopp, L., Ciais, P., Conway, T. J., Doney, S. C., Feely, R. A., Foster, P., Friedlingstein, P., Gurney, K., Houghton, R. A., House, J. I., Huntingford, C., Levy, P. E., Lomas, M. R., Majkut, J., Metzl, N., Ometto, J. P., Peters, G. P., Prentice, I. C., Randerson, J. T., Running, S. W., Sarmiento, J. L., Schuster, U., Sitch, S., Takahashi, T., Viovy, N., van der Werf, G. R., and Woodward, F. I.: Trends in the sources and sinks of carbon dioxide, *Nat. Geosci.*, 2, 831–836, doi:10.1038/ngeo689, 2009.

Littler, M. M.: The productivity of Hawaiian fringing-reef crustose Corallinaceae and an experimental evaluation of production methodology, *Limnol. Oceanogr.*, 18, 946–952, 1973.

Littler, M. M.: Calcification and its role among the macroalgae, *Micronesica*, 12, 27–41, 1976.

Littler, M. M. and Doty, M. S.: Ecological components structuring the seaward edges of tropical Pacific reefs: the distribution, communities and productivity of Porolithon, *J. Ecol.*, 63, 117–129, 1975.

Littler, M. M. and Murray, S. N.: The primary productivity of marine macrophytes from a rocky intertidal community, *Mar. Biol.*, 27, 131–135, 1974.

Littler, M. M., Littler, D. S., Blair, S. M., and Norris, J. N.: Deep-water plant communities from an uncharted seamount off San Salvador Island, Bahamas: distribution, abundance, and primary productivity, *Deep. Res.*, 33, 881–892, 1986.

- 5 Littler, M. M., Littler, D. S., and Dennis Hanisak, M.: Deep-water rhodolith distribution, productivity, and growth history at sites of formation and subsequent degradation, *J. Exp. Mar. Biol. Ecol.*, 150, 163–182, doi:10.1016/0022-0981(91)90066-6, 1991.
- Marsh, J. A.: Primary productivity of reef-building calcareous Red Algae, *Ecology*, 51, 255–263, 1970.
- Martin, S. and Gattuso, J.-P.: Response of Mediterranean coralline algae to ocean acidification and elevated temperature, *Global Change Biol.*, 15, 2089–2100, doi:10.1111/j.1365-2486.2009.01874.x, 2009.
- 10 Martin, S., Clavier, J., Guarini, J.-M., Chauvaud, L., Hily, C., Grall, J., Thouzeau, G., Jean, F., and Richard, J.: Comparison of *Zostera marina* and maerl community metabolism, *Aquat. Bot.*, 83, 161–174, doi:10.1016/j.aquabot.2005.06.002, 2005.
- Martin, S., Castets, M.-D., and Clavier, J.: Primary production, respiration and calcification of the temperate free-living coralline alga *Lithothamnion coralliooides*, *Aquat. Bot.*, 85, 121–128, doi:10.1016/j.aquabot.2006.02.005, 2006.
- 15 Martin, S., Clavier, J., Chauvaud, L., and Thouzeau, G.: Community metabolism in temperate maerl beds, I. Carbon and carbonate fluxes, *Mar. Ecol.-Prog. Ser.*, 335, 19–29, doi:10.3354/meps335019, 2007.
- Martin, S., Rodolfo-Metalpa, R., Ransome, E., Rowley, S., Buia, M.-C., Gattuso, J.-P., and Hall-Spencer, J.: Effects of naturally acidified seawater on seagrass calcareous epibionts, *Biol. Lett.-UK*, 4, 689–92, doi:10.1098/rsbl.2008.0412, 2008.
- 20 Martin, S., Charnoz, A., and Gattuso, J.-P.: Photosynthesis, respiration and calcification in the Mediterranean crustose coralline alga *Lithophyllum cabiochae* (Corallinales, Rhodophyta), *Eur. J. Phycol.*, 48, 163–172, doi:10.1080/09670262.2013.786790, 2013.
- Matsuda, S.: Succession and growth rates of encrusting crustose coralline algae (Rhodophyta, Cryptonemiales) in the upper fore-reef environment off Ishigaki Island, Ryukyu Islands, *Coral Reefs*, 7, 185–195, 1989.
- 25 McCoy, S. J. and Kamenos, N. A.: Coralline algae (Rhodophyta) in a changing world: integrating ecological, physiological, and geochemical responses to global change, *J. Phycol.*, 51, 6–24, 2015.
- 30 McCoy, S. J. and Ragazolla, F.: Skeletal trade-offs in coralline algae in response to ocean acidification, *Nature Climate Change*, 4, 719–723, 2014.
- Meehl, G. A., Stocker, T. F., Collins, W. D., Friedlingstein, P., Gaye, A. T., Gregory, J. M., Kitoh, A., Knutti, R., Murphy, J. M., Noda, A., Raper, S. C. B., Watterson, I. G., Weaver, A. J.,

Calculating the global contribution of coralline algae to carbon burial

L. H. van der Heijden and N. A. Kamenos

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Calculating the global contribution of coralline algae to carbon burial

L. H. van der Heijden and
N. A. Kamenos

- and Zhao, Z.-C.: Global climate projections, in: Climate Change 2007: The Physical Science Basis, Contributing to Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change, edited by: Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K. B., Tignor, M., and Miller, H. L., Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 749–832, 2007.
- Menard, H. W. and Smith, S. M.: Hypsometry of ocean basin provinces, *J. Geophys. Res.*, 71, 4305–4325, 1966.
- Milliman, J. D.: Production and accumulation of calcium carbonate in the ocean: budget of a nonsteady state, *Global Biogeochem. Cy.*, 7, 927–957, 1993.
- Moberly Jr, R.: Composition of magnesian calcites of Algae and Pelecypods by electron microprobe analysis, *Sedimentology*, 11, 61–82, 1968.
- Nellemann, C., Corcoran, E., Duarte, C. M., Valdes, L., De Young, C., Fonseca, L., and Grimsditch, G.: Blue carbon, The role of oceans as carbon sinks, in: Blue Carbon, The Role of Healthy Oceans in Binding Carbon: a Rapid Response Assessment, United Nations Environment Programme, GRID-Arendal, Arendal, Norway, 1–80, 2009.
- Nelson, C. S.: An introductory perspective on non-tropical shelf carbonates, *Sediment. Geol.*, 60, 3–12, 1988.
- Nelson, W. A.: Calcified macroalgae – critical to coastal ecosystems and vulnerable to change: a review, *Mar. Freshwater Res.*, 60, 787–801, 2009.
- Nienhuis, P. H. and Bree, B. H. H.: Production and ecology of eelgrass (*Zostera marina* L.) in the Grevelingen estuary, the Netherlands, before and after the closure, *Hydrobiologia*, 52, 55–66, 1977.
- Odum, E. P.: Energetics and ecosystems, in: Ecology of Halophytes, edited by: Reimold, R. J. and Queen, W. H., Academic Press, London, New York, San Francisco, 599–602, 1974.
- Payri, C. E.: Production primaire et calcification des algues benthiques en milieu corallien, *Oceanis*, 26, 427–463, 2000.
- Platt, T. and Subba Rao, D. V.: Primary production of marine microphytes, in: Photosynthesis and Productivity in Different Environments, edited by: Cooper, J., Cambridge University Press, Cambridge, UK, 249–280, 1975.
- Porzio, L., Buia, M. C., and Hall-Spencer, J. M.: Effects of ocean acidification on macroalgal communities, *J. Exp. Mar. Biol. Ecol.*, 400, 278–287, doi:10.1016/j.jembe.2011.02.011, 2011.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Calculating the global contribution of coralline algae to carbon burial

L. H. van der Heijden and
N. A. Kamenos

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Potin, P., Floc, J. Y., Augris, C., and Cabioch, J.: Annual growth rate of the calcareous red alga *Lithothamnion coralliooides* (Corallinales, Rhodophyta) in the Bay of Brest, France, *Hydrobiologia*, 204, 263–267, 1990.

Ragazzola, F., Foster, L. C., Form, A. U., Büscher, J., Hansteen, T. H., and Fietzke, J.: Phenotypic plasticity of coralline algae in a high CO₂ world, *Ecol. Evol.*, 3, 3436–3446, doi:10.1002/ece3.723, 2013.

Ranwell, S. D.: World resources of *Spartina townsendii* (Sensu lato) and economic use of Spartina marshland, *J. Appl. Ecol.*, 4, 239–256, 1966.

Ries, J. B., Cohen, A. L., and McCorkle, D. C.: Marine calcifiers exhibit mixed responses to CO₂-induced ocean acidification, *Geology*, 37, 1131–1134, doi:10.1130/G30210A.1, 2009.

Roberts, R. D., Kühl, M., and Glud, R. N.: Primary production of crustose coralline red algae in a high Arctic fjord, *J. Phycol.*, 38, 273–283, 2002.

Ryther, J. H.: Photosynthesis and fish production in the sea, *Science*, 166, 72–76, 1969.

Sabine, C. L., Feely, R. A., Gruber, N., Key, R. M., Bullister, J. L., Wanninkhof, R., Wong, C. S., Wallace, D. W. R., Millero, F. J., Peng, T., Kozyr, A., Ono, T., and F, R. A.: The oceanic sink for anthropogenic CO₂, *Science*, 305, 367–371, 2004.

Savini, A., Basso, D., Bracchi, V. A., Corselli, C., and Penneta, M.: Maerl-bed mapping and carbonate quantification on submerged terraces offshore the Cilento peninsula (Tyrrhenian Sea, Italy), *Geodiversitas*, 34, 77–98, 2012.

Semesi, I. S., Kangwe, J., and Björk, M.: Alterations in seawater pH and CO₂ affect calcification and photosynthesis in the tropical coralline alga, *Hydrolithon* sp. (Rhodophyta), *Estuar. Coast. Shelf S.*, 84, 337–341, doi:10.1016/j.ecss.2009.03.038, 2009.

Sheveiko, S. V: Life forms of coralline algae as an environmental indicator on reefs, in: Proc. 4th Int. Cor. Reef. Symp., Manila, 18–22 May, Vol. 2, edited by: Gomez, E. D., Birkeland, C. E., Buddemeier, R. W., Johannes, R. E., Marsh, J. A., and Tsuda, R. Y., Marine Science Centre, Univ. Phillipines, Quezon City, 461, 1981.

Smith, J. E., Smith, C. M., and Hunter, C. L.: An experimental analysis of the effects of herbivory and nutrient enrichment on benthic community dynamics on a Hawaiian reef, *Coral Reefs*, 19, 332–342, doi:10.1007/s003380000124, 2001.

Smith, S. V.: Marine macrophytes as a global carbon sink, *Science*, 211, 838–840, doi:10.1126/science.211.4484.838, 1981.

Sournia, A.: Oxygen metabolism of a fringing reef in French polynesia, *Helgoland. Wiss. Meer.*, 28, 401–410, doi:10.1007/BF01610589, 1976.

- Stearn, C. W., Scoffin, T. P., and Martindale, W.: Calcium carbonate budget of a fringing reef on the west coast of Barbados, I: Zonation and productivity, *Bull. Mar. Sci.*, 27, 479–510, 1977.
- Steneck, R. S.: The ecology of coralline algal crusts: convergent patterns and adaptative strategies, *Annu. Rev. Ecol. Syst.*, 17, 273–303, 1986.
- 5 Steneck, R. S. and Adey, W. H.: The role of environment in control of morphology in *Lithophylum congestum*, a Caribbean algal ridge builder, *Bot. Mar.*, 19, 197–215, 1976.
- Thayer, G. W. and Adams, S. M.: Structural and functional aspects of a recently established *Zostera marina* community, in: *Estuarine Research, Vol. 1: Chemistry, Biology and the Estuarine System*, edited by: Cronin, L. E., Academic Press, London, New York, San Francisco, 10 518–540, 1975.
- Turner, R. E.: Geographic variations in salt marsh macrophyte production: a review, *Contr. Mar. Sci.*, 20, 47–68, 1976.
- Vooren, C. M.: Photosynthetic rates of benthic Algae from the deep coral reef of Curacao, *Aquat. Bot.*, 10, 143–159, 1981.
- 15 Walker, D. I. and Woelkerling, W. J.: Quantitative study of sediment contribution by epiphytic coralline red algae in seagrass meadows in Shark Bay, western Australia, *Mar. Ecol.-Prog. Ser.*, 43, 71–77, 1988.
- Wanders, J. B. W.: The role of benthic Algae in the shallow reef of Curacao (Netherlands Antilles), I: Primary productivity in the coral reef, *Aquat. Bot.*, 2, 235–270, 1976.
- 20 Ware, J. R., Smith, S. V., and Reaka-Kudla, M. L.: Coral Reefs: sources or sinks of atmospheric CO₂?, *Coral Reefs*, 11, 127–130, 1991.
- Whittaker, R. H. and Likens, G. E.: Primary production: the biosphere and man, *Hum. Ecol.*, 1, 357–369, doi:10.1007/BF01536732, 1973.
- 25 Wilson, S.: Environmental tolerances of free-living coralline algae (maerl): implications for European marine conservation, *Biol. Conserv.*, 120, 279–289, doi:10.1016/j.biocon.2004.03.001, 2004.
- Woodwell, G. M., Rich, P. H., and Hall, C. A.: Carbon in estuaries, in: *Carbon and the Biosphere*, edited by: Woodwell, G. M. and Pecan, E. V., AEC Symposium Series 30, NTIS U.S. Dept. of Commerce, Springfield, Virginia, 221–239, 1973.
- 30 Zar, J. H.: Measures of central tendency, in: *Biostatistical Analysis*, edited by Ryu, T., Prentice Hall, New Jersey, 20–31, 1999.

Calculating the global contribution of coralline algae to carbon burial

L. H. van der Heijden and
N. A. Kamenos

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Calculating the
global contribution of
coralline algae to
carbon burial

L. H. van der Heijden and
N. A. Kamenos

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	

Table 1. Primary production (daily and annual) of coralline algae (communities) from different depths and locations. Yearly primary production indicated in Italic are an estimate of the yearly production by taking a daily production and modifying this to a yearly production (x 365). The median production for crustose coralline algae and free-living algae is indicated.

Structure or species	Location	Depth	Primary production (g C m ⁻² d ⁻¹)	Primary production (g C m ⁻² yr ⁻¹)	Reference
Crustose coralline algae				370	this study (<i>n</i> = 35)
<i>Crustose coralline algae</i>	San Salvador Island, Bahamas	81 m	0.07	26	Littler et al. (1986)
<i>Hydrolithon</i> spp.	Klein Piscadera, Curacao	25 m	0.21	77	Vooren (1981)
<i>Sporolithon ptychoides</i>	Klein Piscadera, Curacao	25 m	0.21	78	Vooren (1981)
<i>Pseudolithodermma nigrum</i>	Wilson Cove, California, USA		0.40	146	Littler and Murray (1974)
<i>Sporolithon erythraeum</i>	Waikiki reef, Hawaii, USA		0.50	183	Littler (1973)
<i>Porolithon onkodes</i>	Waikiki reef, Hawaii, USA		0.50	183	Littler (1973)
<i>Porolithon gardneri</i>	Waikiki reef, Hawaii, USA		0.50	183	Littler (1973)
<i>Hydrolithon decipiens</i>	Wilson Cove, California, USA		0.50	183	Littler and Murray (1974)
<i>Phymatolithon foecundum</i> + <i>P. Tenue</i>	Young Sound, NE Greenland	17–36 m		70–300	Roberts et al. (2002)
Reef building coralline algae	Eniwetok Atoll, Hawaii, USA	2 m	0.66	240	Marsh (1970)
<i>Porolithon conicum</i>	Lizard Island, Australia	0–18 m	0.18–1.16	66–423	Chisholm (1988)
<i>Lithophyllum</i> spp.	Coral reef, Curacao	0.5–3 m	0.70	256	Wanders (1976)
<i>Neogoniolithon folsiei</i>	Lizard Island, Australia	0–6 m	0.46–0.95	168–347	Chisholm (1988)
<i>Porolithon onkodes</i>	Lizard Island, Australia	0–6 m	0.37–1.35	135–493	Chisholm (1988)
<i>Hydrolithon reinboldii</i>	Lizard Island, Australia	3–6 m	0.86–0.90	314–329	Chisholm (1988)
<i>Lithophyllum intermedium</i>	Coral reef, Curacao	0.5–3 m	0.90	329	Wanders (1976)
<i>Lithophyllum congestum</i>	Coral reef, Curacao	0.5–3 m	1.00	365	Wanders (1976)
Crustose coralline algae	Coral reef, Curacao	0.5–3 m	1.00	370	Wanders (1976)
<i>Porolithon pachydermum</i>	Coral reef, Curacao	0.5–3 m	1.10	402	Wanders (1976)
<i>Lithophyllum</i> spp.	Coral reef, Curacao	0.5–3 m	1.10	402	Wanders (1976)
<i>Neogoniolithon solubile</i>	Coral reef, Curacao	0.5–3 m	1.40	511	Wanders (1976)
Melobesioid species	Waikiki reef, Hawaii, USA		1.50	548	Littler (1973)
Mainly <i>Neogoniolithon frutescens</i>	Coral reef, Moorea, Tahiti	0.75 m	2.00	730	Sournia (1976)
<i>Porolithon onkodes</i>	Hawaiian Reef, USA	5 m	2.20	803	Littler and Doty (1975)
<i>Porolithon gardneri</i>	Hawaiian Reef, USA	5 m	2.40	876	Littler and Doty (1975)
<i>Corallina elongata</i>	Marseille, France	5 m	2.50	912	Ei Haikali et al. (2004)
<i>Hydrolithon reinboldii</i>	Waikiki reef, Hawaii, USA		2.60	949	Littler (1973)
<i>Neogoniolithon conicum</i> Lab.	Lizard Island, Australia	0–18 m	0.6–4.65	219–1697	Chisholm (2003)
<i>Hydrolithon reinboldii</i> Lab.	Lizard Island, Australia	0–6 m	1.6–3.8	584–1387	Chisholm (2003)
<i>Neogoniolithon brassica-florida</i> Lab.	Lizard Island, Australia	0–6 m	2.45–3.35	894–1223	Chisholm (2003)
<i>Neogoniolithon conicum</i> In situ	Lizard Island, Australia	0–18 m	0.85–5.9	310–2154	Chisholm (2003)
<i>Neogoniolithon brassica-florida</i> In situ	Lizard Island, Australia	0–6 m	2.15–4.7	785–1716	Chisholm (2003)
<i>Hydrolithon onkodes</i> In situ	Lizard Island, Australia	0–3 m	1.75–6.55	639–2391	Chisholm (2003)
<i>Hydrolithon reinboldii</i> In situ	Lizard Island, Australia	3–6 m	4.15–4.35	1515–1588	Chisholm (2003)
<i>Hydrolithon onkodes</i> Lab.	Lizard Island, Australia	0–3 m	4.01–6.05	1464–2208	Chisholm (2003)
Free-living algae				173	This study (<i>n</i> = 4)
Nongeniculate corallines	San Salvador Island, Bahamas	76 m	0.15	55	Littler et al. (1991)
Maerl beds	Bay of Brest, France	0.3–7.9 m	0.38	138	Martin et al. (2005)
<i>Lithophyllum</i> spp.	San Salvador Island, Bahamas	76 m	0.57	208	Littler et al. (1991)
<i>Lithothamnion coralloides</i>	Bay of Brest, France	1–10 m		10–600	Martin et al. (2006)

Calculating the global contribution of coralline algae to carbon burial

L. H. van der Heijden and
N. A. Kamenos

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	

Printer-friendly Version
Interactive Discussion

Table 2. The global average production rates of autotrophic coastal communities. Macroalgae in Gattuso et al. (1998) were macrophyte-dominated. Macrophytobenthic communities in Charpy-Roubaud and Sournia (1990) included brown algae, seagrasses, mangroves and salt marshes.

Community	Production rate		References
	(g C m ⁻² d ⁻¹)	(g C m ⁻² yr ⁻¹)	
Coralline algae (average)	0.9	329	This study (<i>n</i> = 39)
Free-living algae	0.15–0.83	173	This study (<i>n</i> = 4)
Crustose coralline algae	0.07–5	370	This study (<i>n</i> = 35) Chisholm (2003)
	0.9–5		Larkum (1983)
Benthic fleshy algae	0.1–4		Turf algae Larkum (1983)
Mangroves	1–6		Duarte et al. (2005)
		221	Gattuso et al. (1998)
		1081	
Salt marshes		1585	Duarte et al. (2005)
		210	Gattuso et al. (1998)
Seagrasses	1–7 ^L	1211 ^D	^L = Larkum (1983) ^D = Duarte et al. (2005)
		502	Ranwell (1966), Kirby and Gosselink (1976); Odum (1974), Turner (1976), Thayer and Adams (1975), Nienhuis and Bree (1977), Zieman (1975)
Macroalgae		1587	Duarte et al. (2005)
		222	Gattuso et al. (1998)
Benthic diatoms		123	Cadee and Hegeman (1974)
Coastal phytoplankton	0.1–0.5 ^L	196 ^W	^L = Larkum (1983) ^W = Woodwell et al. (1973); Cadee and Hegeman (1974), Gieskes and Kraay (1975)
Coral reefs		148	Duarte et al. (2005)
		120	Gattuso et al. (1998)
Macrophytobenthos		375	Charpy-Roubaud and Sournia (1990)

Calculating the global contribution of coralline algae to carbon burial

L. H. van der Heijden and
N. A. Kamenos

Table 3. The total global production of different coastal communities compared to the total marine oceanic production. The macrophytobenthic community in Charpy-Roubaud and Sournia (1990) included brown algae, seagrasses, mangroves and salt marshes.

Community	Total global production (in 10^9 tCyr^{-1})	References
Coralline algae	0.7	This study ($n = 39$)
Microphytobenthic community	0.34	Charpy-Roubaud and Sournia (1990)
Algal beds and reefs community	1.2	Whittaker and Likens (1973)
Macrophytobenthic community	2.55	Charpy-Roubaud and Sournia (1990)
Phytoplankton community	≥ 30	Charpy-Roubaud and Sournia (1990)
Marine community	48.5	Field et al. (1998)

Table 4. Median calcium carbonate production by coralline algae. Bracchi and Basso (2012) includes Lithophylloids, Canals and Ballesteros (1997) includes *Peysonellia*.

Species	Location	Depth	CaCO ₃ production (g CaCO ₃ m ⁻² yr ⁻¹)	Reference
Crustose coralline algae			1225	This study (<i>n</i> = 25)
Epiphyte corallines on seagrass	Mallorca-Menorca shelf, Mediterranean	0–35 m	68	Canals and Ballesteros (1997)
<i>Mesophyllum</i>	Barbados	fringing reef	167	Stearn et al. (1977)
Coralligenous build-ups + coralline species	Mallorca-Menorca shelf, Mediterranean	70–90 m	170	Canals and Ballesteros (1997)
Crustose coralline algae	Uva Island, Panama	reef flat	190	Eakin (1996)
<i>Neogoniolithon brassica-florida</i> + geniculate	Mallorca-Menorca shelf, Mediterranean	0–10 m	289	Canals and Ballesteros (1997)
<i>Lithophyllum cabiochae</i>	NW Mediterranean	25 m	292	Martin and Gattuso (2009)
Crustose coralline algae	Warraber Island, Australia	reef flat	299	Hart and Kench (2006)
<i>Lithophyllum incrassans</i>	South West Wales, UK	intertidal pools	379	Edyvean and Ford (1987)
Epiphyte corallines on seagrass	Shark Bay, western Australia	10 m	500	Walker and Woelkerling (1988)
<i>Neogoniolithon conicum</i>	Lizard Island, Australia	0–18 m	300–1550	Chisholm (2000)
<i>Hydrolithon reinboldii</i>	Lizard Island, Australia	3–6 m	910–1240	Chisholm (2000)
<i>Porolithon conicum</i>	Lizard Island, Australia	0–18 m	318–1862	Chisholm (1988)
<i>Neogoniolithon</i>	Barbados	fringing reef	1225	Stearn et al. (1977)
<i>Hydrolithon reinboldii</i>	Lizard Island, Australia	3–6 m	1035–1512	Chisholm (1988)
<i>Lithophyllum</i>	Barbados	fringing reef	1355	Stearn et al. (1977)
<i>Neogoniolithon brassica-florida</i>	Lizard Island, Australia	0–6 m	1200–2070	Chisholm (2000)
<i>Hydrolithon onkodes</i>	Ishigaki Is (Ryukyu Is)	upper fore reef	2044	Matsuda (1989)
<i>Hydrolithon onkodes</i>	Lizard Island, Australia	0–3 m	820–3310	Chisholm (2000)
<i>Porolithon onkodes</i>	Penguin Bank, Hawaii	40–100 m	2100	Agegian et al. (1988)
<i>Neogoniolithon fosliei</i>	Lizard Island, Australia	0–6 m	1542–2815	Chisholm (1988)
<i>Porolithon onkodes</i>	Lizard Island, Australia	0–6 m	947–3599	Chisholm (1988)
<i>Porolithon</i>	Barbados	fringing reef	2378	Stearn et al. (1977)
Coraline pavement	One Tree Island, Australia	0–1 m	4000	Kinsey (1985)
<i>Corallina elongata</i>	Marseille, France	0.5–1 m	5037	El Haïkali et al. (2004)
<i>Porolithon onkodes</i>	Rangiroa, Polynesia	reef flat	7400	Payri (2000)
Free-living algae			187	This study (<i>n</i> = 14)
mainly <i>Lithothamnion</i> spp.	Pontian Islands shelf, west Mediterra.	70–100 m	8	Bracchi and Basso (2012)
mainly <i>Lithothamnion</i> spp.	Pontian Islands shelf, west Mediterra.	40–70 m	32	Bracchi and Basso (2012)
<i>Lithothamnion corallicoides</i>	Cilento shelf, west Mediterranean	47 m	91	Savini et al. (2012)
Rhodolith bed	Arvoredo Island, southern Brazil	7–20 m	55–136	Gherardi (2004)
<i>Lithothamnion corallicoides</i>	Mannin Bay, Ireland	0–10 m	29–164	Bosence and Wilson (2003)
<i>Lithothamnion corallicoides</i>	Galway, Ireland	< 10 m	88–164	Bosence (1980)
<i>Phymatolithon calcareum</i>	Mannin Bay, Ireland	0–10 m	79–249	Bosence and Wilson (2003)
<i>Phymatolithon calcareum maerl</i>	Mallorca-Menorca shelf, Mediterranean	40–85 m	210	Canals and Ballesteros (1997)
<i>Phymatolithon calcareum</i>	Galway, Ireland	< 10 m	79–422	Bosence (1980)
<i>Lithothamnion glaciale</i>	Troms, Norway	18 m	420–630	Freiwald and Henrich (1994)
<i>Lithothamnion corallicoides</i>	Bay of Brest, France	0–10 m	876	Potin et al. (1990)
Rhodolith bed	Abrolhos shelf, Brazil	20–110 m	1000	Amado-Filho et al. (2012)
<i>Lithothamnion glaciale</i>	Troms, Norway	7 m	895–1432	Freiwald and Henrich (1994)
<i>Lithothamnion corallicoides</i>	Bay of Brest, France	1–10 m	145–3100	Martin et al. (2006)

Calculating the global contribution of coralline algae to carbon burial

L. H. van der Heijden and N. A. Kamenos

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Calculating the global contribution of coralline algae to carbon burial

L. H. van der Heijden and
N. A. Kamenos

Table 5. Accumulation rates of free-living coralline algae. Coralline algae in Bosence (1985) were predominantly *Neogoniolithon* species.


Species	Location	CaCO ₃ accumulation (mm kyr ⁻¹)	Reference
Rhodolith (maerl)	Troms district, Norway	1400	Freiwald (1998)
Mixed coralline algae	Troms district, Norway	900	Freiwald (1998)
Coralline algae	Orkney Islands, Scotland	80	Farrow et al. (1984)
Branched coralline algae	Tavernier Key, Florida, USA	450	Bosence (1985)
Rhodolith (maerl)	St Mawes Bank, Falmouth, UK	500	Bosence (1980)

- [Title Page](#)
- [Abstract](#) [Introduction](#)
- [Conclusions](#) [References](#)
- [Tables](#) [Figures](#)
- [◀](#) [▶](#)
- [◀](#) [▶](#)
- [Back](#) [Close](#)
- [Full Screen / Esc](#)

- [Printer-friendly Version](#)
- [Interactive Discussion](#)

Calculating the global contribution of coralline algae to carbon burial

L. H. van der Heijden and
N. A. Kamenos



7875

- [Title Page](#)
- [Abstract](#) [Introduction](#)
- [Conclusions](#) [References](#)
- [Tables](#) [Figures](#)
- [◀](#) [▶](#)
- [◀](#) [▶](#)
- [Back](#) [Close](#)
- [Full Screen / Esc](#)
- [Printer-friendly Version](#)
- [Interactive Discussion](#)

Calculating the global contribution of coralline algae to carbon burial

L. H. van der Heijden and
N. A. Kamenos

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Figure 1. The global distribution of the three coralline algae Families (Corallinaceae, Hapalidaceae and Sporolithaceae; for species list per country/region see Supplement, Table S2). The numbers indicate: 1. Spitsbergen, 2. Iceland, 3. Greenland, east, 4. Greenland, 5. Canada, Arctic, 6. Canada, Labrador, 7. Canada, Newfoundland, 8. Canada, New Brunswick, 9. Canada, Nova Scotia, 10. USA, Aleutian Islands, Alaska, 11. USA, Alaska, 12. Revillagigedo Islands, USA, 13. Canada, British Columbia, 14. Canada, Queen Charlotte Islands, 15. USA, Washington, 16. USA, Oregon, 17. USA, California, 18. USA, Channel Islands, California, 19. Mexico, Baja California, 20. Mexico, Isla Guadalupe, 21. USA, Gulf of California, 22. USA, Maine, 23. USA, New Hampshire, 24. USA, Connecticut, 25. USA, Virginia, 26. USA, North Carolina, 27. USA, South Carolina, 28. USA, Florida, 29. USA, Texas, 30. Mexico, 31. Belize, 32. Honduras, 33. El Salvador, 34. Nicaragua, 35. Costa Rica, 36. Panama, 37. Cuba, 38. Bahamas, 39. Caicos Islands, 40. Jamaica, 41. Hispaniola, Dominican Republic, 42. Puerto Rico, 43. Virgin Islands, USA, 44. Saints Kitts, 45. Martinique, 46. Barbados, 47. Saint Thomas, Barbados, 48. Lesser Antilles, 49. Trinidad, 50. Tobago, 51. Trinidad and Tobago, 52. Curacao, 53. Netherlands Antilles, 54. Tropical and Subtropical Western Atlantic, 55. Guyana, 56. Aves, island of Venezuela, 57. Venezuela, 58. Colombia, 59. Ecuador, 60. Galapagos Islands, 61. Peru, 62. Chile, 63. Brazil, 64. Uruguay, 65. Argentina, 66. Falkland Islands, 67. Gough Island, 68. Saint Helena, 69. Ascension, 70. Cape Verde Islands, 71. Canary Islands, 72. Portugal, Salvage Islands, 73. Madeira, 74. Azores, 75. Bermuda, 76. Norway, 77. Sweden, 78. Scandinavia, 79. Baltic Sea, 80. Faroe Islands, 81. Great-Britain, 82. Ireland, 83. Netherlands, 84. France, 85. Spain, 86. Portugal, 87. Gibraltar, 88. Spain, Isla de Alboran, 89. Balearic Islands, Spain, 90. Monaco, 91. Corsica, 92. Sardinia, 93. Italy, 94. Sicily, 95. Malta, 96. Italy, Pelagian Islands, 97. Italy, Adriatic Sea, 98. Slovenia, 99. Croatia, 100. Albania, 101. Greece, 102. Bulgaria, 103. Romania, 104. Black Sea, 105. Turkey, 106. Cyprus, 107. Syria, 108. Israel, 109. Saudi Arabia, 110. Red Sea, 111. Yemen, 112. Oman, 113. Dubai, 114. Abu Dhabi, 115. Qatar, 116. Bahrain, 117. Kuwait, 118. Iran, 119. Persian Gulf, 120. Djibouti, 121. Eritrea, 122. Sudan, 123. Egypt, Red Sea, 124. Egypt, 125. Libya, 126. Tunisia, 127. Algeria, 128. Morocco, 129. Western Sahara, 130. Mauritania, 131. Senegal, 132. Gambia, 133. Sierra Leone, 134. Liberia, 135. Côte d'Ivoire, 136. Ghana, 137. Nigeria, 138. Cameroon, 139. Equatorial Guinea, 140. São Tomé and Príncipe, 141. Gabon, 142. Congo, 143. Angola, 144. Namibia, 145. South Africa, 146. Mozambique, 147. Madagascar, 148. Tanzania, 149. Kenya, 150. Somalia, 151. Ethiopia, 152. Pakistan, 153. India, 154. Sri Lanka, 155. Bangladesh, 156. Comoros and Mayotte, 157. Aldabra Islands, 158. Réunion, 159. Mauritius, 160. Seychelles, 161. Amirante Islands, 162.

Saya de Malha Bank, 163. Cargados Carajos, 164. Rodrigues Island, 165. India, Laccadive Islands, 166. Maldives, 167. Chagos Archipelago, 168. Diego Garcia Atoll, 169. Amsterdam Island, 170. Cocos (Keeling) Islands, 171. Andaman Islands, India, 172. Nicobar Islands, India, 173. Indian Ocean Islands, 174. Myanmar, 175. Thailand, 176. Malaysia, 177. Vietnam, 178. Singapore, 179. Philippines, 180. Indonesia, 181. Indonesia, New Guinea, 182. Taiwan, 183. China, 184. Hong Kong, 185. Japan, 186. Korea, 187. Russia, east, 188. Russia, Kamchatka, 189. Russia, Commander Islands, 190. Saint Paul Island, 191. Easter Island, 192. Northwestern Hawaiian Islands, USA, 193. Hawaiian Islands, USA, 194. Wake Atoll, 195. Ryukyu Islands, Japan, 196. Mariana Islands, 197. Guam, 198. Republic of Palau, 199. Federated States of Micronesia, 200. Marshall Islands, 201. Tuvalu, 202. Samoan Archipelago, 203. American Samoa, 204. Central Polynesia, 205. French Polynesia, 206. Tahiti, 207. Fiji, 208. Solomon Islands, 209. Papua New Guinea, 210. Christmas Island, Australia, 211. Australia, western, 212. Australia, Houtman Abrolhos, 213. Australia, Northern Territory, 214. Australia, Queensland, 215. Australia, New South Wales, 216. Australia, Lord Howe Island, 217. Australia, Norfolk Island, 218. Australia, Victoria, 219. Australia, Bass Strait, 220. Australia, South, 221. Tasmania, 222. New Zealand, 223. New Zealand, Stewart Islands/Rakiura, 224. New Zealand, Snares Islands/Tini Heke, 225. New Zealand, Auckland Islands, 226. New Zealand, Kermadec Islands, 227. New Zealand, Chatman Islands, 228. New Zealand, Bounty Island, 229. New Zealand, Antipodes Islands, 230. Antarctica, Campbell Islands, 231. Antarctica, Macquarie Island, 232. Antarctica, Heard Island, 233. Antarctica, Kerguelen, 234. Antarctica, Crozet Islands, 235. Antarctica, South Georgia, 236. Antarctica, South Orkney Islands, 237. Antarctica, South Shetland Islands, 238. Antarctica, Fuegia, 239. Antarctica, Tierra del Fuego, 240. Antarctica, Peninsula, and 241. Antarctica, Subantarctic Islands.

Calculating the global contribution of coralline algae to carbon burial

L. H. van der Heijden and
N. A. Kamenos

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

