

# 1. Title

## GREENHOUSE GAS BALANCE OF CROPLAND CONVERSION TO BIOENERGY POPLAR SHORT ROTATION COPPICE

*Running title: From cropland to bioenergy SRC: a GHG balance*

*Simone Sabbatini<sup>1</sup>, Nicola Arriga<sup>2</sup>, Teresa Bertolini<sup>3</sup>, Simona Castaldi<sup>3</sup>, Tommaso Chiti<sup>1</sup>, Claudia Consalvo<sup>1</sup>, Sylvestre Njakou Djomo<sup>4,5</sup>, Beniamino Gioli<sup>6</sup>, Giorgio Matteucci<sup>7</sup>, Dario Papale<sup>1</sup>*

*<sup>1</sup> University of Tuscia, Department for Innovation in Biological, Agro-food and Forest systems, Via S. Camillo de Lellis snc, 01100 Viterbo, Italy.*

*<sup>2</sup> University of Antwerp, Department of Biology, Research Group of Plant and Vegetation Ecology, Universiteitsplein 1, B-2610 Wilrijk, Belgium*

*<sup>3</sup> Second University of Naples, Department of Environmental, Biological, Pharmaceutical Sciences and Technologies, Via Vivaldi 43, 81100 Caserta (CE), Italy*

*<sup>4</sup> Hasselt University, Department of Economic, Research Group of Environmental Economics, Martelarenlaan 42, 3500 Hasselt, Belgium*

*<sup>5</sup> Aarhus University, Department of Agroecology, Blichers Alle 20, 8830, Tjele, Denmark*

*<sup>6</sup> Institute of Biometeorology, National Research Council, Via G. Caproni 8, 50145 Firenze, Italy*

*<sup>7</sup> Institute for Agricultural and Forestry Systems in the Mediterranean, National Research Council, Via Cavour 4-6, I-87036 Rende (CS), Italy*

*Corresponding author: Simone Sabbatini, tel. +393396521486, fax +39 0761 357389 email: [simone.sabbatini@unitus.it](mailto:simone.sabbatini@unitus.it)*

*Keywords: land use change, GHG budget, climate mitigation, bioenergy SRC, eddy covariance, SOC change, GHG offset*

*Paper type: Original Research*

## 2. Abstract

The production of bioenergy in Europe is one of the strategies conceived to reduce greenhouse gas (GHG) emissions. The suitability of the land use change from a cropland (REF site) to a short rotation coppice plantation of hybrid poplar (SRC site) was investigated by comparing the GHG budgets of these two systems over 24 months in Viterbo, Italy. This period corresponded to a single rotation of the SRC site. The REF site was a crop rotation between grassland and winter wheat, i.e. the same management of the SRC site before the conversion to short rotation coppice. Eddy covariance measurements were carried out to quantify the net ecosystem exchange of CO<sub>2</sub> ( $F_{CO_2}$ ), whereas chambers were used to measure N<sub>2</sub>O and CH<sub>4</sub> emissions from soil. The measurements began two years after the conversion of arable land to SRC: for that an older poplar plantation was used to estimate the soil organic carbon (SOC) loss due to SRC establishment, and to estimate SOC recovery over time. Emissions from tractors and from production and transport of agricultural inputs ( $F_{MAN}$ ) were modelled. The GHG emission offset due to the substitution of natural gas with SRC biomass for production of heat was credited to the SRC site. Emissions generated by the use of biomass ( $F_{EXP}$ ) were also considered. Suitability was finally assessed by comparing the GHG budgets of the two sites. CO<sub>2</sub> uptake was  $3512 \pm 224$  gCO<sub>2</sub> m<sup>-2</sup> in the SRC site in two years, and  $1838 \pm 107$  gCO<sub>2</sub> m<sup>-2</sup> in the REF site.  $F_{EXP}$  was equal to  $1858 \pm 240$  gCO<sub>2</sub> m<sup>-2</sup> in the REF site, thus basically compensating  $F_{CO_2}$ , while it was  $1118 \pm 521$  gCO<sub>2</sub> m<sup>-2</sup> in the SRC site. This latter could offset  $379.7 \pm 175.1$  gCO<sub>2</sub>eq m<sup>-2</sup> from fossil fuel displacement. Soil CH<sub>4</sub> and N<sub>2</sub>O fluxes were negligible.  $F_{MAN}$  weighed 2% and 4% in the GHG budgets of SRC and REF sites respectively, while the SOC loss weighed  $455 \pm 524$  gCO<sub>2</sub> m<sup>-2</sup> in two years. Overall, the REF site was close to neutrality in a GHG perspective ( $156 \pm 264$  gCO<sub>2</sub>eq m<sup>-2</sup>), while the SRC site was a net sink of  $2202 \pm 792$  gCO<sub>2</sub>eq m<sup>-2</sup>. In conclusion the experiment led to a positive evaluation from a GHG viewpoint of the conversion of cropland to bioenergy SRC.

### 3. Introduction

In the articulated regulation concerning energy and climate change policies, the European Union (EU) established two targets for 2020: (i) reduction of 20% of greenhouse gas (GHG) emissions relative to the levels of 1990, and (ii) share of 20% renewable energy use in gross final energy consumption (European Commission, 2007; European Commission, 2008). For Italy the latter is modulated to 17% (European Commission, 2009).

In the context of climate mitigation, bioenergy crops are expected to play a key role in the renewable energy supply in the EU in the next coming decades (Djomo et al., 2013). Short rotation coppice (SRC) of fast growing trees, and especially of poplar (*Populus* spp.), is a promising culture in this context. SRC has the potential to reduce GHG emissions to the atmosphere during both its production (by capturing CO<sub>2</sub> from the atmosphere and storing it in aboveground biomass and soil) and use (by avoiding CO<sub>2</sub> emissions from fossil fuel burning). However, the management of SRC requires energy inputs, and converting land for SRC production (i.e. land use change, LUC) may alter the equilibrium of the existing ecosystems, causing an impact that in some cases can counterbalance the positive effects on climate mitigation of the SRC (Zona et al., 2013; see also Crutzen et al., 2008; Fargione et al., 2008 for bioenergy crops in general). The LUC to SRC may imply losses of soil organic carbon (SOC) at the installation (Don et al., 2012), especially in C-rich soils, and the management of SRC requires the use of fossil fuels which in some cases can outweigh part of the benefits of the supposed carbon neutral SRC systems (Abbasi and Abbasi, 2009). A recent study (Djomo et al., 2011), however, showed that poplar and willow SRC biomass use can save up to 80%-90% of GHG emissions compared to using coal for energy production. Studies on the climate mitigation potential of poplar cultivations constitute an important tool in supporting energy and environmental policies at different scales. In recent years researchers approached poplar SRCs from different perspectives: ecological (Jaoudé et al., 2010; Zhou et

76 al., 2013), economic (Strauss and Grado, 1997; Mitchell et al., 1999; El Kasmoui and  
77 Ceulemans, 2012, 2013), energetic and environmental (Jungmeier and Spitzer, 2001;  
78 Cherubini et al., 2009; Devis et al., 2009; Nasso et al., 2010; Arevalo et al., 2011;  
79 Don et al., 2012; Dillen et al., 2013; Djomo et al., 2013). However, these studies often used  
80 different approaches making it difficult to compare their results (Migliavacca et al., 2009;  
81 Djomo et al., 2011). Furthermore, emphasis was mainly given to emissions from fossil fuels  
82 compared with the biogenic emissions due to the LUC (Djomo et al., 2013). Including the  
83 different contributions of the LUC in the assessments of emission savings related to energy  
84 crops is crucial (Davis et al., 2009). A full GHG budget based on long-term measurements of  
85 CO<sub>2</sub> and non-CO<sub>2</sub> GHGs via eddy covariance (EC) methodology (Aubinet et al., 2012) and  
86 soil chambers measurements (Allard et al., 2007), can be used to assess the GHG mitigation  
87 potential of land conversion to SRC (Byrne et al., 2007; Ceschia et al., 2010). Several authors  
88 (e.g. Ceschia et al., 2010; Osborne et al., 2010) highlighted the need for a more consistent  
89 number of studies on GHG budgets, including different types of management practices,  
90 climate conditions, and soil characteristics, in order to reduce the uncertainty in GHG budgets  
91 at large scale (Smith et al., 2010). A GHG budget approach was used by Gelfand et al., 2011  
92 in a conversion of unmanaged lands to herbaceous biofuel crops in the US. In Europe, Zona et  
93 al., 2013 estimated the GHG balance in the first year after the conversion from agricultural  
94 lands to a poplar SRC in Belgium, focusing on biogenic contributions. The present study  
95 considered a conversion of a cropland (hereafter indicated as “REF site”) to a poplar SRC  
96 (hereafter indicated as “SRC site”) for bioenergy production in the Mediterranean area  
97 (Viterbo, Central Italy). The aim was to extend the GHG balance to emissions generated by  
98 field management and to the offset of GHG from fossil fuels substitution. The number of  
99 studies on SRC systems cultivated in Mediterranean areas is limited, where water availability  
100 can constitute a limiting factor for biomass yield and thus climate mitigation (Cherubini et al.,

101 2009). Given that the Climate Change mitigation potential of energy crops is the main reason  
102 for subsidies to arable land conversion, our study aimed to assess the suitability of the LUC to  
103 SRC in terms of mitigation of GHG emissions.

104

## 4. Materials and methods

### 4.1 GHG budgets assessment

The GHG budgets were calculated for the SRC and for the REF sites on a temporal basis of two years (24 months), corresponding to the second rotation cycle of the SRC site. They included several positive and negative GHG contributions, with the following sign convention: a positive flux indicates a release into the atmosphere, while a negative flux represents an uptake from the atmosphere. In both cases the boundary of the system was set to the farm level. For the SRC site, the net GHG budget ( $B_{SRC}$ ) was calculated as the algebraic sum of all GHG contributions as indicated in Eq. (1):

$$B_{SRC} = F_{CO_2} + F_{CH_4} + F_{N_2O} + F_{MAN} + F_{SOC} + F_{SAV} + F_{EXP} \quad (1)$$

In equation (1),  $F_{CO_2}$  represents the flux of  $CO_2$ , i.e. the net ecosystem exchange (NEE) of  $CO_2$ , while  $F_{CH_4}$  and  $F_{N_2O}$  represent the biogenic methane and nitrous oxide soil-atmosphere exchanges.  $F_{MAN}$  includes the GHG emissions related to the management of the SRC site, and  $F_{SOC}$  is the loss of soil organic carbon content due to the installation of the cuttings.  $F_{SAV}$  represents the GHG offsets, i.e. avoided GHG emissions due to the substitution of natural gas by biomass in the heat production, and  $F_{EXP}$  the biomass exported from the site at the end of the cycle and reemitted as  $CO_2$  at burning.

Similarly, the net GHG budget of the REF site ( $B_{REF}$ ) was estimated with the algebraic sum indicated in Eq. (2), where in respect to Eq. (1) there is not  $F_{SOC}$  and  $F_{SAV}$ , and  $F_{EXP}$  is the portion of the exported biomass that returns to the atmosphere as  $CO_2$  or  $CH_4$ :

$$B_{REF} = F_{CO_2} + F_{CH_4} + F_{N_2O} + F_{MAN} + F_{EXP} \quad (2)$$

All the contributions of  $B_{SRC}$  and  $B_{REF}$  were expressed as  $CO_2$ -equivalent ( $CO_2eq$ ) fluxes per unit of surface, as the functional unit of the study was one square meter of land. Finally, the net GHG cost or benefit of converting the cropland to a SRC plantation was calculated by

comparing  $B_{SRC}$  and  $B_{REF}$ . Displacement of food and feed production related to SRC cultivation on cropland was beyond the scope of this study.

#### 4.2 Site description

Two sites close to each other located in a private farm (*Gisella ed Elena Ascenzi S.A.A.S.*) in Castel d'Asso, Viterbo, Italy (coordinates: 42°22' N, 12°01' E), were selected during the summer 2011. Two EC towers were installed in the two sites to measure the exchanges of CO<sub>2</sub> and H<sub>2</sub>O between the ecosystem and the atmosphere following the methodology reported in Aubinet et al., 2000. The climate of the area is Mediterranean, with a yearly average rainfall of 766 mm, mean temperature of 13.76 °C and weak summer aridity in July-August (Blasi, 1993). The SRC site was a 2-year rotation cycle managed poplar plantation of 11 ha planted in 2010 to produce biomass for energy (heat). Poplar cultivar was *Populus x canadensis* – clone AF2 selected in *Alasia Franco Vivai*'s nurseries. According to the regional law (Rural Development Programme of Latium 2007-2013, Latium Region, 2015) 12 years is the maximum period to get subsidies for SRC, and corresponded to the time the farmer decided to cultivate the SRC site (personal communication). For that reason the calculations of the present study will be referred to a 12-years lifespan. The site was previously managed with a 2-year rotation between a clover grassland (*Trifolium incarnatum* L.) in mixture with ryegrass (*Lolium multiflorum* Lam) and winter wheat (*Triticum aestivum* L. emend. Fiori et Paol). The REF site was a 9 ha grassland-winter wheat rotation located at a short distance (300 m). Having the identical land use and management of the SRC site before the installation of the poplars, it was selected to assess the GHG effects of the LUC. GHG balances were calculated over 24 months in both sites. However, these 24-month periods did not completely overlap, as the two cultivations had different beginning times: for the SRC site the GHG budget estimation went from 12 January 2012 (immediately after the first harvest of

the SRC site) to 11 January 2014, corresponding to the second cycle of cultivation. The period of calculation of the GHG budget for the REF site went instead from 1 September 2011 until 31 August 2013. However, manual chamber measurements of CH<sub>4</sub> and N<sub>2</sub>O in the REF site started at the beginning of April 2012. The 24 months considered for the SRC site corresponded to the second cycle of the short rotation coppice, and thus did not include the period right after the conversion of agricultural land. This rotation was supposed to terminate with the harvest. However, due to unfavourable climate conditions (a strong drought during summer), the harvest of the SRC site planned for 2014 was postponed to 2015.

The SRC site had a planting density of around 5300 cuttings per hectare, that were planted in rows spaced 2.5 m at a distance of 0.75 m between each other. The first harvest occurred in January 2012. The SRC site was irrigated during the driest periods in summer using a system of tubes installed 35 cm belowground on alternate inter-rows, summing up to about 210 mm in 2012 and 80 mm in 2013 of equivalent precipitation added to the soil. No fertiliser was provided to the SRC site in 2012, while 40 kg per hectare of urea were dissolved in the irrigation water in a single event in 2013. Insecticide (DECIS) was used in May 2012 against *Chrysomela populi* L. In the REF site a shallow tillage (15 cm) was performed in September 2011 with a rotary harrow, and clover and ryegrass were sown. At the end of April 2012 half of the crop was converted to sorghum (*Sorghum vulgare Pers.*) after a period of aridity in spring time. Both the clover and the sorghum were grazed during the growing season, with grazing removing all the above-ground biomass from the sorghum, while the clover was harvested at the end of the cycle. At the end of October 2012 the land was tilled at 40 cm depth, and winter wheat was sown in November. In April 2013 herbicide was distributed over wheat (Buctril at a rate of 1 l ha<sup>-1</sup>), which was harvested at the beginning of July 2013 and no other operation was performed until the end of August. Sorghum was irrigated in several days in summer using a sprinkler with a total amount of 275 mm of equivalent precipitation, while



no irrigation was applied to the winter wheat. Sorghum was also fertilised twice with 150 kg ha<sup>-1</sup> of ammonium nitrate, while 200 kg ha<sup>-1</sup> of the same fertiliser were provided once to the wheat. Apart from irrigation and fertigation, all the operations described above were performed using two different types of tractors, generating different diesel consumptions associated to each operation (Table 3).

An older SRC site (indicated hereafter as O\_SRC site) located alongside the other one and subjected to the same type of management, but planted in 2007, was used in the estimation of SOC content loss caused by the LUC. This was necessary as the expected SOC loss following the conversion (i.e. during the first rotation) was not measured.

In the 24 months considered for the GHG budget of the SRC site, precipitations summed up to 1078 mm, with an average temperature of 14.72 °C, while in the 24 months used for the REF site precipitations were 1157 mm and average temperatures 15.31 °C. In both cases yearly values of precipitation were lower than the long-term average of 766 mm (Blasi, 1993).

An intense drought occurred in summer 2012, with no rain from the beginning of June until the end of August, in contrast to the long-period average of cumulate rainfall in these months (110 mm, Blasi, 1993). Soils were classified as *Chromic Luvisol* according to the World Reference Base classification (USS 2014), with a clay-loam texture. Values of pH ranged between 5.88 in the REF site, 6.66 in the O\_SRC site and 6.69 in the SRC site. The stock of nitrogen (N) up to 70 cm was not significantly different between sites, ranging from 3.16 ± 1.60 MgN ha<sup>-1</sup> to 3.19 ± 1.47 MgN ha<sup>-1</sup> and 3.25 ± 1.47 MgN ha<sup>-1</sup> respectively for SRC, O\_SRC and REF sites. See Fig. 1 for a schematic representation of land cover and management events of the two sites.

#### 4.3 F<sub>CO2</sub>: eddy covariance measurements

203 The EC technique was used to determine the turbulent vertical fluxes of momentum, CO<sub>2</sub>,  
 204 latent and sensible heat. A 3-D sonic anemometer was installed in each site for high-  
 205 frequency measurements of wind speed, wind direction and sonic temperature. CO<sub>2</sub> and water  
 206 vapour densities were collected using a fast-response open-path infrared gas analyser (see  
 207 Table 1 for models and manufacturers). These instruments were mounted on towers located in  
 208 the centre of the fetches. On the REF site the mast was 3 m high, while an extendible  
 209 telescopic pole was used in the SRC site in order to always measure turbulences above the  
 210 roughness layer (Foken, 2008). Measurement heights ranged between 5 and 8 m, and the  
 211 distances of the measuring system over the  $d + z_0$  plane between 2 and 5 m ( $d$  = displacement  
 212 height;  $z_0$  = roughness length). Several meteorological variables above and belowground were  
 213 continuously measured on a 30 min basis to properly calculate fluxes and characterise the two  
 214 sites. In Table 1 the complete instruments setup is described, including both meteorological  
 215 and high-frequency variables.

216 Half-hourly fluxes were calculated with EddyPro® software (LI-COR, Lincoln, NE, USA).  
 217 Several corrections to the time series (Aubinet et al., 2012) were applied as reported in Table  
 218 2. Post-processing included spike removal and friction velocity ( $u_*$ ) filtering (Papale et al.,  
 219 2006), gap-filling using the marginal distribution sampling (MDS) approach and partitioning  
 220 of  $F_{CO2}$  into gross primary production (GPP) and ecosystem respiration ( $R_{eco}$ ) components  
 221 (Reichstein et al., 2005). The gap-filled  $F_{CO2}$  and its components were then cumulated along  
 222 the 24-month period considered.

223 Uncertainty in  $F_{CO2}$  was calculated on the basis of the uncertainty in the  $u_*$  filtering, assuming  
 224 that the main potential systematic error is due to advection and thus linked to the  $u_*$  filtering.  
 225 One hundred thresholds were calculated using a bootstrapping technique and then applied to  
 226 filter the data. For each half-hour, the median of the distribution of  $F_{CO2}$  obtained using the

100 thresholds was used for the GHG budget (Gielen et al., 2013), and the range of uncertainty was derived as half the range 16<sup>th</sup>-84<sup>th</sup> percentile.

#### 4.4 Soil characteristics and SOC stock and changes

To better characterize the soil properties and to quantify the changes in SOC stocks due to the installation of the poplar plantation, a number of soil analyses were performed in the three sites in two different periods. In a first phase on February 2012 three soil trenches 150 cm wide were opened randomly in each site and the soil sampled by depth (0-5, 5-15, 13-30, 30-50, 50-70, 70-100 cm) at both the opposite sides of the profiles to have six replicate samples per depth. The bottom layer (70 cm – 100 cm) was absent in the REF site due to the presence of the bedrock at 80 cm, rather than 100 cm as in both the SRC sites. Samples were collected using a cylinder to determine also the bulk density. Main goals of this first sampling campaign were to describe the soil characteristics and to determine the number of replicates necessary to detect with statistical significance a change in SOC content of 0.5 gC kg<sup>-1</sup> soil (Conen et al., 2003). In the SRC and O\_SRC sites ten samples of the organic layer were also taken removing all the material present over the mineral surface within a squared frame with an area of 361 cm<sup>2</sup>. In the REF site this sampling was not performed because a permanent organic layer was missing. All samples were air-dried at room temperature and then sieved at 2 mm to separate the coarse fraction, and the analyses performed on the fine earth. The pH was measured in deionised water with a sure-flow electrode, using a ratio soil-solution of 1:2.5 (w/w), and texture was determined after destruction of the cement using sodium hypochlorite adjusted at pH 9 (Mikutta et al., 2005). The sand fraction was separated by wet sieving at 53 µm while the silt and the clay fractions were separated by time sedimentation according to the Stokes law. Total carbon (C) and nitrogen concentrations were measured on finely ground samples by dry combustion (ThermoFinnigan Flash EA112 CHN), while SOC

and N stocks were determined taking into account soil C and N concentrations and a weighed mean of bulk density, depth of sampling and stoniness (Boone et al., 1999). During the second phase in March 2014 a new sampling was performed in the REF, SRC and O\_SRC sites. The number of samples necessary to detect statistically a SOC change was 50, as derived from the first phase. Samples were taken from the first 15 cm of soil, as most of the changes in a short period occur in the shallower layers. C concentration was measured and SOC stocks re-calculated. The normality of the distributions was checked using a Chi-squared test (Pearson, 1900). An ANOVA test (Fisher, 1919), combined with a Tukey multiple comparison test were used to check if SOC stocks were different between the sites. As data of  $F_{CO_2}$  from the beginning of the cultivation are missing, SOC changes due to the installation of the poplar cuttings were calculated building a linear regression between SOC content of the SRC site (4 years old) and the O\_SRC site (7 years old), then estimating the SOC at the time of plantation (year “0”). Following the “free-intercept model” described by Anderson-Teixeira et al., 2009, the SOC content change due to the plantation of the SRC was then extrapolated considering the difference between the SOC content at year 0 and the one measured in the REF site, assuming the SOC content in the REF site in equilibrium, as this type of land use was constant in the last 30 years. Uncertainties in SOC concentration and stock were calculated as standard deviations from the mean values of each repeated measure, while errors were estimated using the law of error propagation as reported by Goodman, 1960.

#### 4.5 Soil $CH_4$ and $N_2O$ fluxes

On-site measurements of  $CH_4$  and  $N_2O$  soil fluxes were combined with laboratory incubation analyses, where soil samples were tested at different water contents and N addition levels. Field measurements of soil  $N_2O$  and  $CH_4$  fluxes were carried out in the two sites using nine

277 manual, dark, static PVC chambers (15 cm diameter, 20 cm height, and total volume  
278 0.0039 m<sup>3</sup>) per site, placed over as many PVC collars (7 cm height, 15 cm diameter)  
279 permanently inserted into the soil at 5 cm depth for the period of observation. At the SRC site,  
280 three collars were distributed along the tree line (between two trees), three along the irrigated  
281 inter-rows and three along the non-irrigated inter-rows, while in the REF site collars were  
282 placed in three different blocks of three collars each. Gas samples were collected from each  
283 chamber at the closure time, and 30 and 60 minutes after closure. Samples were stored in  
284 glass vials provided with butyl rubber air tight septum (20 ml) and concentration of N<sub>2</sub>O and  
285 CH<sub>4</sub> was measured using a trace Ultra gas chromatograph (GC) (Thermo Scientific, Rodano,  
286 IT). The flux detection limit due to the concentration measurement was in the order of about  
287 0.1 mg of CH<sub>4</sub> or N<sub>2</sub>O m<sup>-2</sup> day<sup>-1</sup>, and the analytical precision of the GC for standards at  
288 ambient concentration was approximately 3-5 %, using one standard deviation as a measure  
289 of mean error. Further details on GC set in Castaldi et al., 2013. Measurements started two  
290 weeks after collar insertion and samples were collected every 2-4 weeks, depending on land  
291 management practices and weather conditions, for a total of 30 dates in the SRC site and 24  
292 for the REF site. Similar frequencies were used in previous studies (e.g. Pihlatie et al., 2007;  
293 Weslien et al., 2009), and considered pertinent on the basis of the low magnitude of the  
294 measured fluxes. To test if fertilisation could trigger a peak of N<sub>2</sub>O emission as found in  
295 previous studies (e.g. Gauder et al., 2012), measurements in both sites were carried out more  
296 frequently in occasion of fertilisation events (on average every two days), starting from the  
297 day before the application of fertiliser and for a week. Measurements also covered different  
298 soil and meteorological conditions, including periods of drought and rewetting. Measured  
299 average daily soil CH<sub>4</sub> and N<sub>2</sub>O fluxes were cumulated over the 24 months by linear  
300 interpolation as described by Marble et al., 2013, and uncertainty calculated propagating the  
301 standard deviations of the replicates. Intergovernmental Panel on Climate Changes (IPCC)

100-year global warming potential (GWP) weighed estimates of GHGs (Forster et al., 2007) were used to convert  $F_{N_2O}$  and  $F_{CH_4}$  into CO<sub>2</sub> equivalents: factors 298 and 25 respectively.

#### 4.5.1 Laboratory incubations

Due to the fact that we don't have continuous measurements of non-CO<sub>2</sub> fluxes from soil, we performed a laboratory analysis to verify the accuracy of field campaigns. Laboratory incubations were carried out to assess the GHG emission rates under controlled laboratory conditions in soil treated with both water and nitrogen addition, and to quantify the rates of soil mineralization and nitrification. The rational of the incubation was to assess whether the fluxes were driven by limiting conditions like water and/or nitrogen, or slow rate of organic N mineralization, as found in a Mediterranean coppice site in the same region (Castaldi et al., 2009; Gundersen et al., 2012). Addition of N allowed to check if short-time peaks of emissions occurred that could escape due to the selected frequency of sampling. Soil cores (7 cm diameter, 10 cm height) sampled in the two ecosystems were incubated at 20 °C. Water was then added to reach three different ranges of Water Filled Pore Space (WFPS%): 20% (i.e. the value estimated at sampling), 50% and 90%, each of them replicated five times. The sample at the highest WFPS% was also replicated with or without nitrogen supply (100 kgN ha<sup>-1</sup> of NH<sub>4</sub>NO<sub>3</sub>). Cores were placed in gas-tight 1-litre jars and 6 ml air samples were collected immediately after closure and after 3 hours of incubation for N<sub>2</sub>O production determination. Gas concentration was determined by gas chromatography the day after the treatment and in the following 5 days, leaving the jars open during this period and closing them only when N<sub>2</sub>O production needed to be determined, in order to avoid developing of liquid oxygen tension conditions. Net mineralization and nitrification, and net potential nitrification rate were determined on sieved (2 mm mesh) soil samples over 14 days of incubation, while for the determination of potential nitrification soil was amended with

ammonium sulphate  $(\text{NH}_4)_2\text{SO}_4$  ( $100 \mu\text{gN g}^{-1}$  dry soil). A modified method (Kandeler, 1996; Castaldi and Aragosa, 2002) was used to extract  $\text{NH}_4^+$  and  $\text{NO}_3^-$  from the soil at  $T_0$  and  $T_{14}$  days for further concentration determination with calibrated specific electrodes after the addition of a pH and ionic buffer 0.4 ml di ISA (Ionic Strength Adjustor; Orion cat. No. 951211 e Orion cat No. 930711). Mineralization rates were calculated as the total soil mineral N ( $\mu\text{g}$  of  $\text{N-NH}_4^+ + \text{N-NO}_3^-$  per gram of dry soil) measured after 14 days of incubation ( $T_{14}$ ) minus total mineral N measured at incubation start ( $T_0$ ) divided by the number of days of incubation. Nitrification rates were calculated similarly, considering only the amount of  $\text{N-NO}_3^-$  produced at  $T_{14}$  minus the amount of  $\text{N-NO}_3^-$  present at  $T_0$ .

In order to compare results obtained with soil cores to field conditions, in situ *WFPS%* was calculated for the whole period of field monitoring:

$$WFPS\% = \frac{M_{SOIL}}{1 - (\rho_{BULK}/\rho_{PART})} * 100 \quad (3)$$

where  $M_{SOIL}$  is the volumetric soil moisture ( $\text{m}^3 \text{m}^{-3}$ ),  $\rho_{BULK}$  is the bulk density ( $\text{Mg m}^{-3}$ ) and  $\rho_{PART}$  is the particle density ( $\text{Mg m}^{-3}$ ). For mineral soil  $\rho_{PART}$  is approximated to that of common silicate materials ( $2.65 \text{ Mg m}^{-3}$ , Chesworth, 2008).

#### 4.6 Emissions due to management

Life cycle inventory (LCI) was used to estimate the anthropogenic GHG emissions due to farming operations (Robertson et al., 2000) in both sites (Table 3), and the GHG emissions due to grazing in the REF site (Table 4). The present study is not a full LCA, but the LCA approach was used to estimate emissions caused by field management as described in the following. In particular, indirect land use change (iLUC) was not taken into account. iLUC includes modifications in land use elsewhere in the world triggered by the local substitution of an arable land with an energy crop (Djomo et al., 2013). iLUC occurred outside the boundary

351 of the system we used for this analysis, i.e. the farm. Fossil fuel emissions associated with the  
 352 cultivation of the SRC and REF sites included on-site emissions from tractors (used to carry  
 353 out all the main operations: ploughing, seeding, solid fertilisation, harvesting) and irrigation,  
 354 as well off-site emissions from the production and transport of agricultural inputs (fertiliser,  
 355 insecticide, herbicide). Emissions due to the production of tractors were considered negligible  
 356 as in Budsberg et al., 2012 and Caputo et al., 2014. On-site GHG emissions due to diesel  
 357 consumption were calculated as the product of the amount of fuel diesel consumed to carry  
 358 out a given farm activity (e.g. harvesting) and the emissions factor of diesel,  $90 \text{ gCO}_2\text{eq MJ}^{-1}$   
 359 (Table 3). This factor includes emission costs due to the combustion of diesel  
 360 ( $74 \text{ gCO}_2\text{eq MJ}^{-1}$ ), and emissions due to its production and transportation ( $16 \text{ gCO}_2\text{eq MJ}^{-1}$ )  
 361 (Edwards et al., 2007). Considering energy density of diesel to be  $38.6 \text{ MJ l}^{-1}$  (Alternative  
 362 Fuel Data Center, 2014), producing, transporting and burning 1 l of diesel emitted  
 363  $3474 \text{ gCO}_2\text{eq}$ . An exception was made for harvesting in the SRC site, for which emissions for  
 364 diesel consumption relative to the previous harvest (2012) were considered, as the harvest at  
 365 the end of the cycle was postponed. Emissions due to irrigation were calculated by  
 366 multiplying the electricity consumed in powering the pumps by an emissions factor of  
 367  $750 \text{ gCO}_2 \text{ kWh}^{-1}$ , calculated as the average of different emission factors for different sources  
 368 of electricity (Bechis and Marangon, 2011) weighted on the Italian electricity grid mix,  
 369 derived from the Italian energetic balance 2012 (Italian Ministry of Interior, 2013). Off-site  
 370 emission costs for fertilisers and insecticides were estimated as the product of applied amount  
 371 of fertiliser or insecticide and the emission factors for manufacturing 1 kg of  
 372 fertiliser/insecticide:  $4018.9 \text{ gCO}_2 \text{ kg}^{-1} \text{ N}$  for urea (NPK rating 40-0-0),  $4812 \text{ gCO}_2 \text{ kg}^{-1} \text{ N}$  for  
 373 diammonium phosphate (NPK 18-46-0)<sup>1</sup>,  $7030.8 \text{ gCO}_2 \text{ kg}^{-1} \text{ N}$  for ammonium nitrate (NPK  
 374 33-0-0) and  $7481.9 \text{ gCO}_2 \text{ kg}^{-1} \text{ N}$  for calcium ammonium nitrate (NPK 27-0-0) (Wood and

---

<sup>1</sup> This includes production and transport costs of the overall fertiliser, including P.



Cowie, 2004). Although emission factors differ among insecticide types, in this analysis we assumed that the difference is negligible as the use of insecticides was limited, and thus considered the emission factor of insecticide (active ingredient: deltamethrin) as the product of energy required to produce 1 kg of insecticide ( $310 \text{ MJ kg}^{-1}$ ) and the emission rate of insecticide ( $60 \text{ gCO}_2 \text{ MJ}^{-1}$ ) (Barber, 2004; Liu et al., 2010). The emission factor of herbicide was taken from literature (Ceschia et al., 2010):  $3.92 \text{ kgC per kg of product}$ . Fuel used for the application of chemical products was included in the on-site calculations described above. All the contributions listed above were converted on a surface basis (Table 3).

#### 4.7 *Biomass use and GHG offset*

During the first year of cultivation the REF site was grazed by sheep, which were brought to the field in defined periods (Table 4). Hence, the aboveground biomass (AGB) from the REF site was rather grazed by sheep or provided as hay to other livestock, destined to meat and milk production, or in the case of wheat used in food (grains) and feed (foliage) production. Due to the different species cultivated throughout the two years and to the different uses of the biomass,  $F_{EXP}$  of the REF site (Eq. (2)) includes the following:

$$F_{EXP} = E_{CH_4,on} + E_{CO_2,on} + E_{CH_4,off} + E_{CO_2,off} \quad (4)$$

where the first subscript indicates whether the exported C is reemitted to the atmosphere as  $\text{CO}_2$  or  $\text{CH}_4$ , and the second subscript distinguishes between emissions occurring on-site (*on*) and off-site (*off*). In fact, the percentage of AGB ingested by herbivores on grassland varies with the intensity of management (Soussana et al., 2010). In the present study, however, what was left in the field by the sheep was then harvested and provided them off-site. We assumed then that, apart from the grains in wheat ears, all the AGB was ingested by sheep or other livestock, and that the digestible portion of the organic C ingested was respired back to the atmosphere as  $\text{CO}_2$  or emitted as  $\text{CH}_4$  via enteric fermentation (Eq. (4)) (Soussana et al.,

2007). Biomass in the REF site was sampled every 2-3 weeks in five plots (0.5 m x 0.5 m) randomly selected within the field. In three dates samples were collected immediately after grazing in a grazed area and in an undisturbed area to quantify the intensity of mowing (68%) and identify the C ingested on-site and off-site. Biomass samples were oven-dried at 70 °C to constant mass and weighed. Total AGB was obtained cumulating dry weights measured immediately before each grazing event, subtracting each time the 32% of the dry weight of the previous sample to consider mowing intensity. IPCC methodology (Dong et al., 2006) was then used to estimate  $E_{CH_4,on}$  (Eq. (4)), adjusting the methane emission factor per animal considering the average weight (55 kg) of sheep (19 gCH<sub>4</sub> head<sup>-1</sup> day<sup>-1</sup>), and multiplying it by the daily number of sheep present on-site. The method in Soussana et al., 2007 (their Eq. (4)) was then adapted to estimate the other three components in Eq. (4):  $E_{CH_4,off}$  was estimated applying to the C ingested off-site the ratio between the C weight in  $E_{CH_4,on}$  and the C ingested on-site. The C emitted as CH<sub>4</sub> was subtracted from the digestible portion of the C ingested, assumed to be 65%, and the remaining converted in CO<sub>2</sub> as to estimate  $E_{CO_2,on}$  and  $E_{CO_2,off}$ . The remaining, non-digestible C (35%) was assumed to be returned to the SOC of the grassland (for the on-site part) or of other systems (for the off-site part) as faeces, thus not contributing to the GHG balance. The portion of C that was stock in the body mass of animals was considered negligible (Soussana et al., 2007). For the sake of simplicity, we assumed that also the C content of wheat ears will be shortly respired back to the atmosphere as CO<sub>2</sub>, and thus included in  $E_{CO_2,off}$  (Eq. (4)).

At the end of the cycle, poplar aboveground woody biomass (AGWB) of the SRC site was supposed to be harvested and burnt, thus from one side releasing C back to the atmosphere, and from the other offsetting GHG emissions for fossil fuels displacement. To estimate poplar AGWB diameters were measured at the end of the cycle, after the leaves fall. Three rows of trees were selected inside the plantation and the diameters of these trees were measured

(minimum threshold 0.5 cm) at 1 m height. A simple model considering the regression between individual shoot dry weight ( $W_D$ ) and 1 m diameter ( $D$ ) was used:

$$W_D = b * D^c \quad (5)$$

where  $b$  and  $c$  are empirical parameters,  $W_D$  is in kg DM (kg of Dry Mass), and  $D$  is in cm. Parameters were set as  $b = 0.0847$  and  $c = 2.112$  following Mareschi, 2008 (see also Paris et al., 2011) for the second rotation cycle of clone AF2 of the plantation located in Bigarello (Mantua province), as the one with the more similar climatic and soil characteristics, and also with the same root and shoot age. Dry combustion (1108EA, Carlo Erba, Milan, IT) was used to determine the C concentration for both sites. Regarding the GHG emissions offset, it was assumed that heat produced from SRC biomass will substitute heat produced from natural gas. The GHG offset ( $F_{SAV}$ ) was estimated based on the yield of the SRC site, the energy density of poplar, the conversion efficiency of typical biomass boiler in Italy, and the emission rate of heat production from natural gas in Italy:

$$F_{SAV} = Y * H_L * \eta_{CONV} * I_{NG} \quad (6)$$

where  $Y$  is the biomass yield ( $\text{kg m}^{-2}$ ),  $H_L$  is the low heating value of poplar ( $13 \text{ MJ kg}^{-1}$  at 30% moisture content, Boundy et al., 2011),  $\eta_{CONV}$  is the efficiency of conversion of poplar chips to heat, assumed in this study to be 84% (Saidur et al., 2011), and  $I_{NG}$  is the carbon emission rate (intensity) of heat produced from natural gas (i.e.  $55.862 \text{ gCO}_2\text{eq MJ}^{-1}$ ) for Italy (Romano et al., 2014).

## 5. Results

### 5.1 Biogenic fluxes of CO<sub>2</sub>

The cumulative  $F_{CO_2}$  in the REF site for the two years considered was  $-1838 \pm 107 \text{ gCO}_2 \text{ m}^{-2}$ , partitioned in  $8032 \pm 313 \text{ gCO}_2 \text{ m}^{-2}$  absorbed through photosynthesis (GPP) and  $6216 \pm 338 \text{ gCO}_2 \text{ m}^{-2}$  emitted by total  $R_{eco}$ . In the SRC site cumulative  $F_{CO_2}$  summed up to  $-3512 \pm 224 \text{ gCO}_2 \text{ m}^{-2}$ , with GPP equal to  $8717 \pm 298 \text{ gCO}_2 \text{ m}^{-2}$  and  $R_{eco}$  equal to  $5205 \pm 425 \text{ gCO}_2 \text{ m}^{-2}$  (Fig. 2). Hence, the SRC site was a larger CO<sub>2</sub> sink compared to the REF site over the measuring period, due to both the higher GPP and the lower ecosystem respiration of the SRC site relative to the REF site.

Seasonal differences between the sites in the net flux of CO<sub>2</sub> were observed (Fig. 3). The main difference was the timing of the peak of CO<sub>2</sub> uptake, during the spring in the REF site and in summer for the SRC site. In both sites, peaks of CO<sub>2</sub> uptake were higher in 2013 than in 2012. In the latter, however, a minor peak of uptake were observed in early fall in the SRC site. Periods with positive net fluxes of CO<sub>2</sub> appeared longer and with higher values in the REF site (Fig. 3, top). Air temperatures registered in the two sites were similar, but higher in summer 2012, while the SWC at 30 cm depth was higher in the REF than in the SRC site (Fig. 3, bottom).

### 5.2 Soil CH<sub>4</sub> and N<sub>2</sub>O fluxes

Daily average of both  $F_{N_2O}$  and  $F_{CH_4}$  were very low in almost every measurement (Fig. 4), leading to low total cumulative soil  $F_{N_2O}$  and  $F_{CH_4}$  for both the sites: overall soil non-CO<sub>2</sub> fluxes were  $15.5 \pm 4.7 \text{ gCO}_2\text{eq m}^{-2}$  in two years for the SRC site and  $0.5 \pm 1.6 \text{ gCO}_2\text{eq m}^{-2}$  in two years for the REF site. Both sites were small sources of N<sub>2</sub>O and small sinks of CH<sub>4</sub>. CH<sub>4</sub> sink at the SRC site was not significantly different from the one at the REF site, although on

average slightly higher, and significantly higher N<sub>2</sub>O emissions were observed at the SRC site, although still very low. Measurements carried out in occasion of fertilisation events showed no significant increase in the emission rates of N<sub>2</sub>O in respect to non-fertilisation periods: fluxes in the SRC site in the period of the unique fertilisation occurred in the two years of study remained low, and in the REF site none of the four measurements taken in the period of the fertilisation event of June 2012 exceeded the detection limit of the GC.

### 5.2.1 Laboratory incubations

N<sub>2</sub>O emissions determined in laboratory incubations confirmed that over most of the analysed WFPS% values both soils were producing little N<sub>2</sub>O in absence of N addition, even at WFPS% normally considered to trigger N<sub>2</sub>O emission (WFPS% 60-80%) (Fig. 5). Addition of N did not seem sufficient to stimulate N<sub>2</sub>O production. In contrast, very high WFPS%, close to saturation, was able to trigger a strong increase of N<sub>2</sub>O production in the soil of the REF site. Comparing the data reported in Fig. 5 with the field data of WFPS% for the REF site (Fig. 6), it can be seen that most of the time WFPS% was significantly below 70% in the whole profile and that at 5 cm, where most of the interaction with added fertilizer might have occurred, the WFPS% never exceeded 50%. Mineralization and nitrification rates were quite low in both sites, with slightly positive mineralization rates in the SRC site ( $0.28 \pm 0.05 \mu\text{gN g}^{-1} \text{d}^{-1}$ ) and a very small net immobilization in the REF samples ( $-0.2 \pm 0.2 \mu\text{gN g}^{-1} \text{d}^{-1}$ ). Net nitrification rates calculated in the control (no N addition) were also quite low and varied between  $0.5 \pm 0.05 \mu\text{gN g}^{-1} \text{d}^{-1}$  and  $-0.1 \pm 0.2 \mu\text{gN g}^{-1} \text{d}^{-1}$  in the REF site, that might suggest either a quite slow ammonification phase as a limiting step of the nitrification or a slow nitrification rate. However, when ammonium sulphate was added to soil samples the potential nitrification rates significantly increased reaching  $1.8 \pm 0.1 \mu\text{gN g}^{-1} \text{d}^{-1}$  and  $1.4 \pm 0.3 \mu\text{gN g}^{-1} \text{d}^{-1}$  in the SRC and the REF sites respectively, suggesting that

mineralization might be the limiting step of subsequent nitrification and denitrification processes in the field.

### 5.3 Emissions due to management

The GHG emissions due to management practices were in total  $100.9 \pm 20 \text{ gCO}_2\text{eq m}^{-2}$  for the SRC site and  $135.7 \pm 27.1 \text{ gCO}_2\text{eq m}^{-2}$  for the REF site. Analysing the single contributions, differences arose between the two sites (Fig. 7): fertilisation was the main source of emission of GHGs in the REF site, while its contribution to GHG emissions of the SRC site was limited. Irrigation constituted a big portion of the GHG emissions from management operations in the SRC site, while in the REF site, despite similar amounts of water provided, irrigation played a smaller role, similar to harvesting and tillage. Emissions due to the latter were more relevant in the REF site than in the SRC site.

### 5.4 SOC content changes

In the first 15 cm of soil total C stocks were  $1603 \pm 376 \text{ gC m}^{-2}$  in the REF site,  $1169 \pm 442 \text{ gC m}^{-2}$  in the SRC site and  $1403 \pm 279 \text{ MgC ha}^{-1}$  in the O\_SRC site. The statistical analysis performed on the SOC stocks showed that there were statistically significant differences between SOC data of the three sites (Table 5;  $p\text{-value} = 2.05 \cdot 10^{-7}$ ). The linear regression between SOC content of SRC and O\_SRC sites led to the relation:

$$SOC(t) = 78 * t + 857 \quad (7)$$

where  $t$  are the years from plantation and  $SOC$  is the soil organic carbon content expressed in  $\text{gC m}^{-2}$ . Estimated uncertainty was  $25 \text{ gC m}^{-2}$  for the slope value, and  $139 \text{ gC m}^{-2}$  for the intercept (Fig. 8), meaning that the yearly SOC accumulation after poplar plantation was  $78 \pm 25 \text{ gC m}^{-2}$  and the initial value ( $t=0$ ) was  $857 \pm 139 \text{ gC m}^{-2}$ ,  $746 \pm 858 \text{ gC m}^{-2}$  lower than

the REF value, corresponding to the SOC content loss due to the installation of the SRC. As this loss was a positive flux occurring only once in a LUC at the installation of the cuttings (Arevalo et al., 2011), and that the expected lifespan of the SRC site was 12 years, the value considered for the 24-month GHG budget was  $1/6$ , corresponding to  $124 \pm 143 \text{ gC m}^{-2}$  ( $455 \pm 524 \text{ gCO}_2 \text{ m}^{-2}$ ).

### 5.5 Biomass use and GHG offset

The dry weight of AGB in the REF site summed up to  $0.72 \pm 0.18 \text{ kg m}^{-2}$  for the grassland, of which  $0.35 \pm 0.07 \text{ kg m}^{-2}$  due to the clover in mixture and  $0.37 \pm 0.17 \text{ kg m}^{-2}$  from the sorghum, while the winter wheat totalled  $0.63 \pm 0.09 \text{ kg m}^{-2}$ , of which  $0.36 \pm 0.05 \text{ kg m}^{-2}$  in the ears. The C content measured was 46% for all species, leading to a total of  $621.0 \pm 93.2 \text{ gC m}^{-2}$  in AGB, of which  $265.5 \pm 79.2 \text{ gC m}^{-2}$  ingested by sheep on-site,  $191.2 \pm 49.8 \text{ gC m}^{-2}$  used by livestock off-site, and  $163.9 \pm 21.9 \text{ gC m}^{-2}$  converted to food. The estimated emissions of  $\text{CH}_4$  due to enteric fermentation was  $4.3 \pm 1.3 \text{ gCH}_4 \text{ m}^{-2}$ , equal to  $3.3 \pm 1.0 \text{ gC m}^{-2}$  emitted as  $\text{CH}_4$ , and thus corresponding to  $109 \pm 33 \text{ gCO}_2\text{eq m}^{-2}$  ( $E_{\text{CH}_4,\text{on}}$ , Eq. (4)). Hence, about 1.25% of the ingested C became  $\text{CH}_4$  in the digestive process. Using this ratio led to estimate other  $2.4 \pm 0.6 \text{ gC m}^{-2}$  emitted as  $\text{CH}_4$  off-site, i.e.  $3.2 \pm 0.8 \text{ gCH}_4 \text{ m}^{-2}$ , or  $80 \pm 20 \text{ gCO}_2\text{eq m}^{-2}$  ( $E_{\text{CH}_4,\text{off}}$ ). Subtracting the C emitted as  $\text{CH}_4$  on- and off-site to the respective digestible C ingested by sheep and other livestock led to  $621 \pm 189 \text{ gCO}_2\text{eq m}^{-2}$  emitted on-site ( $E_{\text{CO}_2,\text{on}}$ ) and  $447 \pm 118 \text{ gCO}_2\text{eq m}^{-2}$  offsite. Adding to this latter the emissions expected from wheat ears use (i.e.  $601 \pm 80 \text{ gCO}_2\text{eq m}^{-2}$ ) gave  $1048 \pm 143 \text{ gCO}_2\text{eq m}^{-2}$  ( $E_{\text{CO}_2,\text{off}}$ ): in total  $1858 \pm 240 \text{ gCO}_2\text{eq m}^{-2}$  in two years ( $F_{\text{EXP}}$ , Eq.(4)).

For the SRC site, applying Eq. (5) with the diameters distribution led to estimate AGWB (dry matter) in  $0.62 \pm 0.29 \text{ kg m}^{-2}$ , which with a C content of 49%, corresponded to a  $F_{\text{EXP}}$  of

1118  $\pm$  521 gCO<sub>2</sub>eq m<sup>-2</sup> per two years that are expected to be reemitted to the atmosphere at the combustion. This value of AGWB then corresponded to 8.1  $\pm$  3.7 MJ m<sup>-2</sup> of gross energy from biomass chips, which decreased to 6.8  $\pm$  3.1 MJ m<sup>-2</sup> of final heat obtainable from burning biomass chips when the conversion efficiency is considered. This could offset about 379.7  $\pm$  175.1 gCO<sub>2</sub>eq m<sup>-2</sup> from final heat produced using natural gas.

## 5.6 GHG budgets

All the contributions reported in the previous sections were summed up to calculate the GHG budgets of the two sites. The net GHG budget of the REF site ( $B_{REF}$ , Eq. (2)) amounted to 156  $\pm$  264 gCO<sub>2</sub>eq m<sup>-2</sup>, indicating that the REF site was close to neutrality from a GHG perspective, while for the SRC site the  $B_{SRC}$  (Eq. (1)) resulted in a cumulative sequestration of -2202  $\pm$  792 gCO<sub>2</sub>eq m<sup>-2</sup>. The different components of the GHG budget of the two sites are summarized in Fig. 9. In the REF site the  $F_{CO_2}$ , weighing about 48% in the GHG budget, was completely compensated by the emissions of CO<sub>2</sub> and CH<sub>4</sub> due to the biomass utilisation (about 44% and 5% respectively), while the other components had a minor role ( $F_{MAN}$  around 4%, soil non-CO<sub>2</sub> <1%).  $F_{CO_2}$  was the main contribution also in the SRC site, where it represented the 63% of  $B_{SRC}$ , while  $F_{EXP}$  represented the 20%, SOC loss (8%) and the GHG offset for the fossil fuel substitution (7%) had a similar weight, and the other contributions played a minor role. As  $B_{REF}$  was almost neutral and the SRC site a sink of GHGs, the difference between the two GHG budgets was favourable to the SRC site (2358  $\pm$  835 gCO<sub>2</sub>eq m<sup>-2</sup> saved), highlighting the advantages in terms of GHGs of the LUC from common agricultural to SRC of poplar in the study area.



## 6. Discussion

The two ecosystems behaved differently in the measuring period: they were both characterised by a seasonal uptake of CO<sub>2</sub> (Fig. 3), driven by the timing and duration of the growing season, occurring in spring at the REF site and in summer at the SRC site. The peak of CO<sub>2</sub> uptake was similar at both sites in 2012, while it was higher at the REF site in 2013. Periods with positive CO<sub>2</sub> fluxes were longer in the REF site, and often higher in magnitude, likely as a consequence of the shorter growing season of grasses and winter wheat compared to the poplars trees of the SRC site. Also the land cover of the two sites during the dormant periods and the shift in time between them might have played a role on this difference: some herbaceous vegetation kept growing in the SRC site in wintertime, while harvesting and ploughing in the REF site in late summer/early fall might have enhanced ecosystem respiration. Inter-annual differences were also observed at both sites. Both the higher air temperature and the more extended period of low SWC proved the strong aridity of summer 2012, responsible for the autumnal increase of CO<sub>2</sub> uptake at the SRC site, corresponding to the rewetting of the soil. At the REF site, autumn uptake was higher in 2011, while the springtime uptake was much higher in 2013 than in 2012 (Fig. 3). This different behaviour was mostly ascribable to the different cultivations (grassland – winter wheat), and to some extent to the different climate conditions in springtime. All these differences in ecosystems responses resulted in a net sink of GHGs from the SRC site and in a neutral GHG balance for the REF site.

A GHG balance not significantly different from zero is in agreement with the average results for a set of sites in Soussana et al., 2007, where however management costs were not considered, and on-site CO<sub>2</sub> emissions from grazing animals were measured with EC. C sequestered by the SRC site in our study was higher than that of the Belgian site in the study of Zona et al., 2013. In the latter the net budget was positive (in a time span of one year and a

half) with a net emission of  $280 \pm 80 \text{ gCO}_2\text{eq m}^{-2}$ , due to both the higher emission rates of  $\text{CH}_4$  and  $\text{N}_2\text{O}$  fluxes from soil ( $350 \pm 50 \text{ gCO}_2\text{eq m}^{-2}$ ), and to the lower  $\text{CO}_2$  sink ( $-80 \pm 60 \text{ gCO}_2\text{eq m}^{-2}$ ) as compared to the present study. Also Jassal et al., 2013 found lower  $F_{\text{CO}_2}$  in a 3-year-old poplar SRC in Canada ( $-293 \text{ gCO}_2 \text{ m}^{-2} \text{ year}^{-1}$ ) as compared to  $F_{\text{CO}_2}$  of the SRC site of the present study (root age: 4 years), likely due to the lower stem density of their site. All these values lied in the range found by Arevalo et al., 2011, i.e.  $-77 \text{ gCO}_2 \text{ m}^{-2} \text{ year}^{-1}$  and  $-4756 \text{ gCO}_2 \text{ m}^{-2} \text{ year}^{-1}$  relative to a 2-year-old and 9-year-old poplar SRC respectively. These results show that even in a Mediterranean area, where plants are subjected to drought stress, with a proper use of irrigation there is the potential for a positive effect on climate mitigation.

Several studies (Grigal and Berguson, 1998; Price et al., 2009) confirmed that converting agricultural land to SRC resulted in an initial release of SOC due to SRC establishment, and then in a slow and continuous accumulation of SOC due to vegetation activity and wood encroachment (Arevalo et al., 2011). Despite the deep tillage at the SRC establishment, and the fact that the REF site was ploughed every year at different depths, a gradient decreasing with depth in the C distribution of the vertical profile was evident in the three sites (not shown). This suggests that the changes in SOC were attributable only to the plantation of the SRC due to the effects of tillage (Anderson-Teixeira et al., 2009), and not to the mechanical redistribution of SOC. This study indicates a SOC loss of 47% in respect to the value measured in the REF site, due to the installation of poplar cuttings. This loss was not measured at the time it occurred, i.e. right after the conversion of arable land to poplar short rotation coppice, but was estimated with data from the O\_SRC site. The reported value was close to the range maximum reported in the review by Post and Kwon, 2000 (20%-50%), but was higher than the results found by Arevalo et al., 2011 (7%). The absolute value, however, was close to the one of this latter study ( $8 \text{ MgC ha}^{-1}$ ), where though the initial SOC was one

614 order of magnitude higher ( $114.7 \text{ MgC ha}^{-1}$ ). To correctly interpret this rapid loss of SOC for  
 615 a conversion of a cropland to a SRC the low degree of disturbance that characterises the REF  
 616 site must be taken into account. Furthermore the loss of SOC found in the present study has to  
 617 be considered along with its own uncertainty that was as large as the estimated value: in the  
 618 purposes of the GHG balance, where the uncertainty of the single components are propagated  
 619 to the net budget, this result is correctly interpreted as a range. We highlight that a loss of  
 620 SOC close to the minimum of the abovementioned range by Post and Kwon, 2000, e.g.  
 621  $321 \text{ gC m}^{-2}$ , would have changed  $B_{SRC}$  ( $-2202 \pm 792 \text{ gCO}_2\text{eq m}^{-2}$ ) by only  $-259 \text{ gCO}_2\text{eq m}^{-2}$ .  
 622 Thus, even if a measured value would have probably been more accurate, the sensitivity of the  
 623 total GHG budget to this loss was shown to be relatively low. The estimated annual SOC  
 624 accumulation rate was in the range reported by Don et al., 2011 for SRCs  
 625 ( $0.44 \pm 0.43 \text{ MgC ha}^{-1} \text{ y}^{-1}$ ), which explained how the frequent harvest of above ground  
 626 biomass was likely to facilitate the die off of the roots that contributes to SOC accumulation.  
 627 In our study, the low biomass yield supports the hypothesis that a big fraction of C taken up  
 628 via photosynthesis was transferred to roots and soil. In our study the break-even point, where  
 629 the initial SOC content would be restored and a net SOC accumulation would start, was 10  
 630 years, in agreement with findings from other studies (e.g. Hansen, 1993; Arevalo et al., 2011  
 631 found a value of 7 years, while Grigal and Berguson, 1998, calculated a break-even point of  
 632 15 years). This result, not directly involved in the 24-month GHG budget, is relevant  
 633 considering that the SRC of the present study is expected to be used for 12 years, thus enough  
 634 to allow the complete recovery of the SOC loss occurred at the plantation. Different previous  
 635 land uses, soil types (in particular clay content), climate conditions, fertilisation rates may be  
 636 the main causes of differences between studies, as shown in a meta-analysis by Laganière et  
 637 al., 2010.

638 Our results showed that CH<sub>4</sub> and N<sub>2</sub>O soil fluxes were not relevant in the GHG budgets due to  
639 the combination of soil characteristics and climatic trends at both sites. Low values are  
640 reported in other studies for SRCs: e.g. Gauder et al., 2012 found that the soil of different  
641 energy crops acted as weak sink of CH<sub>4</sub> even in case of fertilisation, while emissions of N<sub>2</sub>O  
642 turned out to be higher for annual than perennial (willow) crops, which showed no significant  
643 effect of fertilisation on N<sub>2</sub>O fluxes. Agricultural sites usually have higher N<sub>2</sub>O effluxes from  
644 soil, though their magnitude depends on cultivations and on management practices, as shown  
645 by Ceschia et al., 2012. The SRC site as a perennial woody crop was subjected to low soil  
646 disturbance during its lifespan, while the REF site was ploughed once per year, with an  
647 impact on the ecosystem respiration. Zona et al., 2012 found high N<sub>2</sub>O emissions in the first  
648 growing season of a poplar SRC in Belgium:  $197 \pm 49 \text{ gCO}_2\text{eq m}^{-2}$  in six months, which  
649 drastically decreased to  $42 \pm 17 \text{ gCO}_2\text{eq m}^{-2}$  for the whole following year. This suggested an  
650 influence of soil disturbance during land conversion on the stock of N in soil, which was  
651 almost 1/3 lower in our study sites than in the one of Zona ( $9.1 \pm 2.1 \text{ MgN ha}^{-1}$ ). In the  
652 present experiment however, N<sub>2</sub>O fluxes were low both in the SRC and REF sites, even  
653 during periods of fertilisation, with no clear patterns. The low N<sub>2</sub>O fluxes were confirmed by  
654 laboratory analyses, as the presence of extra N did not affect the emission rates of N<sub>2</sub>O, and  
655 only very high WFPS% could trigger significant N<sub>2</sub>O fluxes. The needed conditions of soil  
656 humidity were never reached in the REF site and persisted only for a few days at 35 cm depth  
657 in the SRC site (Fig. 6). At this depth fertilizer was added as fertigation in the SRC site: we  
658 hypothesize that the very low porosity, the compaction and strength of the soil might have  
659 favoured slow gas release and further N<sub>2</sub>O reduction, thus leaving little N<sub>2</sub>O to escape to the  
660 atmosphere from soil surface. In the REF site, winter fertilisation was also associated with  
661 low temperatures, a further constraint to microbial activity. These results provide further  
662 evidence of how the simple application of the IPCC N<sub>2</sub>O emission factor to the analysed

systems might have led to an overestimation of the field GHG contribution to the overall GWP in both sites. Laboratory estimates of mineralization and nitrification rates suggested that N mineralization might be the limiting process of the chains of mineral N microbial transformations, that contributed to maintain N<sub>2</sub>O emissions low even during events of intense rainfall and soil saturation. The clay content and compaction of the analysed soils might be an important factor in limiting oxygen and substrate diffusion that are both necessary to have optimal rates of soil organic matter mineralization. From a methodological point of view, the low emissions of both CH<sub>4</sub> and N<sub>2</sub>O from soil also suggest that using 4 samples of gas concentration per chamber instead of 3 would have not dramatically improved the accuracy of the calculated fluxes, as a slight variation in the slope would have not induced significant changes in the results. The relevance of this result lies in the fact that fertilising a poplar SRC in a Mediterranean area and in this kind of soil does not necessarily lead to increased emissions of N<sub>2</sub>O, with the requirement that the correct equilibrium is found between irrigation and WFPS%. Thus it is possible to maximise yield and GHG mitigation with the right management practices (Nassi o Di Nasso et al., 2010). CH<sub>4</sub> and N<sub>2</sub>O fluxes might have been enhanced by the land conversion in the first period of cultivation of the SRC site, as found for CO<sub>2</sub>. However, measurements carried out in the REF site, ploughed every year, and the incubation experiment showed very low fluxes, mostly related to soil characteristics and not to management activities. Thus, a low sensitivity of the total GHG budget to these components can be expected.

Other components of the GHG budget related to N compounds (e.g. aerosol NH<sub>4</sub>NO<sub>3</sub>, N deposition and leaching) were considered negligible in this study as compared to the role of N<sub>2</sub>O emissions from soil and related to fertiliser production.

Regarding the use of biomass, comparisons with other studies for the REF site are complicated because half of the field was converted to sorghum in spring for the low

688 productivity experienced during the drought. However, the productivity of the clover in  
 689 mixture was found highly variable by Martiniello, 1999, and the results of the present study  
 690 are comparable with the lower values found by this author in non-irrigated stands in  
 691 Mediterranean climate ( $0.39 \text{ kg m}^{-2}$ ). Sorghum productivity was lower than that reported by  
 692 Nassi o Di Nasso et al., 2011 (around  $0.75 \text{ kg m}^{-2}$ ) in a similar climate, likely due to the short  
 693 period of cultivation and to grazing. The productivity of winter wheat was similar to that of  
 694 Anthoni et al., 2004 ( $0.32 \pm 0.03 \text{ kg m}^{-2}$ ). The drought in summer 2012 had an important  
 695 influence on the AGWB of the SRC site, which was lower compared to other studies (e.g.  
 696 Scholz and Ellerbrock, 2002,  $0.4$  to  $0.7 \text{ kg m}^{-2} \text{ year}^{-1}$ ), and to the  $F_{CO_2}$  values found with EC.  
 697 Our hypothesis is that the period of drought had influenced the aboveground/belowground  
 698 ratio, and that the herbaceous vegetation contributed to increase the  $F_{CO_2}$ . In terms of C, the  
 699 difference  $F_{CO_2} - F_{EXP}$  represents to a first approximation the C stocked by each ecosystem  
 700 that does not return shortly to the atmosphere after utilisation, minus heterotrophic respiration  
 701 (Rh). While in the SRC site that difference was negative (C sink of  $650 \text{ gC m}^{-2}$ ), the REF site  
 702 acted like a small source of C ( $120 \pm 98 \text{ gC m}^{-2}$ ). Small sources were also found by Anthoni et  
 703 al., 2004 (between  $50 \text{ gC m}^{-2}$  and  $100 \text{ gC m}^{-2}$ ), while Aubinet et al., 2009 reported a 4-year  
 704 rotation crop being a source of  $340 \text{ gC m}^{-2}$ . For poplar, Deckmyn et al., 2004 found a similar  
 705 behaviour in a poplar SRC in Belgium. Concerning the fraction of the exports that is emitted  
 706 as  $\text{CH}_4$  from enteric fermentation, our estimates were in agreement with those of Dengel et  
 707 al., 2011. Several studies (e.g. Gilmanov et al., 2007) used EC to measure  $\text{CO}_2$  and  $\text{CH}_4$   
 708 fluxes from grazed systems. Some included in the GHG budget only  $F_{CO_2}$ ,  $F_{CH_4}$  and  $F_{N_2O}$ , and  
 709 made a C budget for lateral fluxes like biomass export (e.g. Allard et al., 2007). However, the  
 710 EC method is not capable of measuring point sources of trace gases moving inside and  
 711 outside the footprint (data discarded by QA/QC procedures: see also Baldocchi et al., 2012).  
 712 Thus we adapted the method described in Soussana et al., 2007 for off-site emissions,

extending it also to on-site emissions, to include the effects of aboveground biomass use in the GHG budget.

Different studies (e.g. Cherubini et al., 2009; Djomo et al., 2013) confirmed the advantages of using biomass from SRC over fossil fuels in mitigating the increase of atmospheric GHG concentrations, while Abbasi and Abbasi, 2010 found that the SRC management led to GHG emissions that compensate the gain due to the fossil substitution. The low yield of the SRC site led to lower GHG savings compared to those found by Cherubini et al., 2009 for production of heat from woody products ( $379.7 \pm 175.1 \text{ gCO}_2\text{eq m}^{-2}$  in two years against  $600 \text{ gCO}_2\text{eq m}^{-2}$  per year). These latter found GHG mitigation to be directly proportional to crop yield for dedicated bioenergy crops. In a GHG budget perspective, however, the yield is also proportional to C emissions from combustion, and correlated with  $F_{CO_2}$ . The same study reported GHG savings of other bioenergy systems, showing that the performance of wood-based systems is lower in terms of GHG offset than the one of other bioenergy crops, e.g. switchgrass ( $1300 \text{ gCO}_2\text{eq m}^{-2} \text{ year}^{-1}$ ), Miscanthus ( $1600 \text{ gCO}_2\text{eq m}^{-2} \text{ year}^{-1}$ ) and fibre sorghum ( $1800 \text{ gCO}_2\text{eq m}^{-2} \text{ year}^{-1}$ ). In the present study the role of GHG offset was relevant in the GHG balance; however it's important to consider that natural gas, while being the most used fossil fuel for heating systems in Italy, has also a lower carbon intensity for heat production ( $55.862 \text{ gCO}_2\text{eq MJ}^{-1}$ ) as compared to coal ( $76.188 \text{ gCO}_2\text{eq MJ}^{-1}$ ) and oil ( $73.693 \text{ gCO}_2\text{eq MJ}^{-1}$ ) (Romano et al., 2014). A different scenario, where biomass would substitute the use of other energy sources with higher emission factors (like coal) would lead to a higher GHG offset.

Our study confirmed that farming operations have only a limited importance in the overall GHG budget when conditions of relevant  $\text{CO}_2$  uptake by vegetation are met, and the values we found were similar to the ones found by Gelfand et al., 2011. In the SRC site irrigation was more important than other contributions and caused more emissions than irrigation in the

REF site. This suggests that belowground irrigation was less efficient in terms of GHG emissions than the sprinkler. Fertilisers and other chemical products often have a higher impact on the GHG balance as compared to other field operations due to the off-site GHG emissions (Ceschia et al., 2010). At the study sites the amount and frequency of applications were relatively small, and this justifies the minor role of fertilisation in the total GHG budget. Thus the importance of farming operations can vary from year to year, depending on climate conditions and on farmer decisions.

This study reports on the GHG budget of poplar SRC in Mediterranean areas. However, when considering the implications of SRC in a wider perspective, other factors should also be considered to assess the overall sustainability of this type of LUC. Among them, irrigation is one of the most important (Dougherty and Hall, 1995), as poplar cultivations in Mediterranean climate require considerable amounts of water. In the LUC presented here, both the SRC and the REF sites were irrigated with similar amounts of water, using a less efficient technique at the REF site (sprinkler system) than at the SRC site (belowground drip system; e.g. Camp, 1998). The impact of the LUC on the local water balance is thus expected to be small in this particular case, but not in general. An appropriate design of these systems is also crucial to avoid water dispersion: in the present study we observed that irrigation could not compensate the drought stress experienced by the SRC site in 2012, thus concerns arise on the proper location of the belowground tubes and on the amounts of water applied. The aim of this study was to analyse the LUC from a GHG perspective at farm level: the boundary of our system constituted the main difference with a full LCA analysis, where the iLUC is considered in addition to the direct land use change. iLUC can cause GHG emissions elsewhere, thus reducing the mitigation potential of the studied SRC on a global scale.



## 7. Conclusions

This study analysed a land use change (LUC) for biomass production from a greenhouse gas (GHG) perspective. The conversion of a traditional cropland (REF site) to a short rotation coppice (SRC site) of poplar hybrids in a Mediterranean climate (Central Italy) was considered. Different fluxes were included in a GHG budget calculated for both sites: to and from vegetation and soil; caused by management; due to the loss of soil organic Carbon (SOC) at the installation of the cuttings; caused by the use of the biomass and the displacement of GHG emissions from fossil fuel. Other environmental aspects than the GHG balance were not considered, such as the water balance, the minor components of the nitrogen cycle and the contribution of the indirect land use change (iLUC: Djomo et al., 2013). The direct LUC contributions, like the SOC content loss at the installation of the SRC and the corresponding disadvantages in terms of GHG, were instead included in the calculation.

Our study showed how poplar SRC cultivation for biomass production was overall suitable from the point of view of the climate mitigation. The most important components of the GHG budgets were the net ecosystem exchange ( $F_{CO_2}$ ) and the C export ( $F_{EXP}$ ) at the end of the cultivation cycle. Interestingly, fluxes of  $CH_4$  and  $N_2O$  from soil were not relevant, likely due to physical soil characteristics. No significant effects were observed even in case of fertilisation, irrigation or rain events, in contrast with findings from other studies (e.g. Zona et al., 2012). Results demonstrated that poplar clones have the ability to stock high rates of C from the atmosphere even on Mediterranean climate, where the cultivation of poplar SRC may be critical for its dependence on water availability. Estimated uncertainty was however quite large, confirming the need of large efforts in terms of data collection to correctly estimate the different components. Benefits of the LUC from common agriculture to SRC derived from the interaction between the diverse components of the budget. Climate conditions and farmer needs are the most important factors controlling the single

787 contributions of the GHG budget. An equilibrate combination of clones selection, irrigation  
788 and management activities, depending also on soil properties, is thus crucial to achieve an  
789 efficient contribution to Climate Change mitigation by LUC for bioenergy crops.

790

## 7. Acknowledgments

This research was supported by the GHG Europe Project (EU Seventh Framework Programme, FP7) and the MIUR-PRIN project “Climate change mitigation strategies in tree crops and forestry in Italy” (CARBOTREES). The authors wish to acknowledge the technical employees of DIBAF (A. Baiocco, A. Boschi, M. Tomassucci), CNR IBAF (G. Santarelli) and CNR IBIMET (A. Zaldei) for their irreplaceable support and logistic help during the field campaign, and the database staff (E. Canfora, A. Ribeca, D. Polidori, C. Trotta) for their support in the data post-processing. S. Sabbatini is supported by the ICOS-Inwire EU project and S. Njakou Djomo was supported by the ERC Advanced Grant 233366 POPFULL. We further thank G. Parisi for gas chromatography analysis. We are also grateful to the farm *Gisella ed Elena Ascenzi S.A.A.S.*, and in particular to A. Trani, the farmer that kindly allowed us to install our instruments on his fields and to access his farm whenever we needed it and that supported our scientific activities with his interest and availability. All the data collected are available at the European Ecosystem Fluxes Database at [www.europe-fluxdata.eu](http://www.europe-fluxdata.eu), site codes IT-CA1, IT-CA2 and IT-CA3.

## 8. References

- Abbasi, T., and Abbasi, S.A.: Biomass energy and the environmental impacts associated with its production and utilization. *Renew Sust Energ Rev*, 14(3), 919-937, 2010.
- Allard, V., Soussana, J.F., Falcimagne, R., Berbigier, P., Bonnefond, J.M., Ceschia, E., D'hour, P., Hénault, C., Laville, P., Martin, C., and Pinarès-Patino, C.: The role of grazing management for the net biome productivity and greenhouse gas budget (CO<sub>2</sub>, N<sub>2</sub>O and CH<sub>4</sub>) of semi-natural grassland. *Agr Ecosyst Environ*, 121(1), 47-58, 2007.
- Alternative Fuels Data Center: Fuel properties comparison. Available at: [http://www.afdc.energy.gov/fuels/fuel\\_comparison\\_chart.pdf](http://www.afdc.energy.gov/fuels/fuel_comparison_chart.pdf) (last access: 10 March 2015), 2014
- Anderson-Teixeira, K.J., Davis, S.C., Masters, M.D., and Delucia, E.H.: Changes in soil organic carbon under biofuel crops. *Global Change Biology Bioenergy*, 1(1), 75–96, doi:10.1111/j.1757-1707.2008.01001.x, 2009
- Anthoni, P., Freibauer, A., Kolle, O., and Schulze, E.: Winter wheat carbon exchange in Thuringia, Germany. *Agr Forest Meteorol*, 121, 55-67, 2004.
- Arevalo, C., Bhatti, J.S., Chang, S.X., and Sidders, D.: Land use change effects on ecosystem carbon balance: from agricultural to hybrid poplar plantation. *Agr Ecosyst Environ*, 141(3), 342-349, 2011.
- Aubinet, M., Grelle, A., Ibrom, A., Rannik, Ü., Moncrieff, J., Foken, T., Kowalski, A.S., Martin, P.H., Berbigier, P., Bernhofer, C., Clement, R., Elbers, J., Granier, A., Grünwald, T., Morgenstern, K., Pilegaard, K., Rebmann, C., Snijders, W., Valentini, R., and Vesala, T.: Estimates of the annual net carbon and water exchange of forests: the EUROFLUX methodology. *Adv Ecol Res*, 30, 113-175, 2000.
- Aubinet, M., Moureaux, C., Bodson, B., Dufranne, D., Heinesch, B., Suleau, M., Dufranne, D., Heinesch, B., Suleau, M., Vancutsem, F., and Vilret, A.: Carbon sequestration by a crop

832 over a 4-year sugar beet/winter wheat/seed potato/winter wheat rotation cycle. *Agr Forest*  
833 *Meteorol*, 149(3), 407-418, 2009.

834 Aubinet, M., Vesala, T., and Papale, D. (Eds.): *Eddy covariance: a practical guide to*  
835 *measurement and data analysis*, Springer, New York, USA, 2012.

836 Baldocchi, D., Detto, M., Sonnentag, O., Verfaillie, J., Teh, Y.A., Silver, W., and Kelly,  
837 N.M.: The challenges of measuring methane fluxes and concentrations over a peatland  
838 pasture. *Agr Forest Meteorol*, 153, 177-187, 2012.

839 Barber, A.: Seven case study farms: total energy & carbon indicators for New Zealand arable  
840 & outdoor vegetable production. *AgriLINK New Zealand Ltd*, 288, 2004.

841 Bechis, S., and Marangon, F.: *Analisi delle emissioni di CO<sub>2</sub> nelle diverse fonti energetiche,*  
842 *report*, Department of Economy, Agricultural, Forest and Environmental Engineering,  
843 University of Turin, Italy, 2011.

844 Blasi, C.: *Carta del fitoclima del Lazio. Regionalizzazione e caratterizzazione climatica*, map,  
845 Regione Lazio, Assessorato Agricoltura e Foreste, Caccia e Pesca, Usi civici. Sapienza  
846 University, Dept of Vegetal Biology, Rome, 1993.

847 Boone, R.D., Grigal, D.F., Sollins, P., Ahrens, R.J., and Armstrong, D.E.: Soil sampling,  
848 preparation, archiving, and quality control, in: *Standard Soil methods for Long-term*  
849 *Ecological Research* (eds. Robertson, G.P., Coleman, D.C., Bledsoe, C.S., and Sollins, P.),  
850 Oxford University Press, New York, pp. 3-28, 1999.

851 Boundy, B., Diegel, S.W., Wright, S.W.L., and Davis, S.C.: *Biomass energy data book:*  
852 *edition 4. Report #ORNL/TM-2011/446 by the Oak Ridge National Laboratory for the Office*  
853 *of the Biomass Program*, US Department of Energy, Oak Ridge, Tennessee, 2011

854 Byrne, K.A., Kiely, G., and Leahy, P.: Carbon sequestration determined using farm scale  
855 carbon balance and eddy covariance. *Agr Ecosyst Environ*, 121(4), 357-364, 2007.

856 Budsberg, E., Rastogi, M., Puettmann, M.E., Caputo, J., Balogh, S., Volk, T.A., Gustafson,  
857 R., and Johnson, L.: Life-cycle assessment for the production of bioethanol from willow  
858 biomass crops via biochemical conversion. *Forest Prod J*, 62(4), 305, 2012.

859 Caputo, J., Balogh, S.B., Volk, T.A., Johnson, L., Puettmann, M., Lippke, B., and Oneil, E.:  
860 Incorporating uncertainty into a life cycle assessment (LCA) model of short-rotation willow  
861 biomass (*Salix* spp.) crops. *BioEnergy Research*, 7(1), 48-59, 2014.

862 Castaldi, S., and Aragosa, D.: Factors influencing nitrification and denitrification variability  
863 in a natural and fire-disturbed Mediterranean shrubland. *Biol Fert Soils*, 36(6), 418-425, 2002.

864 Castaldi, S., Carfora, A., Fiorentino, A., Natale, A., Messere, A., Miglietta, F., and Cotrufo,  
865 M.F.: Inhibition of net nitrification activity in a Mediterranean woodland: possible role of  
866 chemicals produced by *Arbutus unedo*. *Plant Soil*, 315, 273-283, 2009.

867 Castaldi, S., Bertolini, T., Valente, A., Chiti, T., and Valentini, R.: Nitrous oxide emissions  
868 from soil of an African rain forest in Ghana. *Biogeosciences*, 10, 4179-4187, 2013.

869 Ceschia, E., Béziat, P., Dejoux, J.F., Aubinet, M., Bernhofer, C., Bodson, B., Buchmann, N.,  
870 Carrara, A., Cellier, P., Di Tommasi, P., Elbers, J.A., Eugster, W., Grünwald, T., Jacobs,  
871 C.M.J., Jans, W.W.P., Jones, M., Kutsch, W., Lanigan, G., Magliulo, E., Marloie, O., Moors,  
872 E.J., Moureaux, C., Oliso, A., Osborne, B., Sanz, M.J., Saunders, M., Smith, P., Soegaard,  
873 H., and Wattenbach, M.: Management effects on net ecosystem carbon and GHG budgets at  
874 European crop sites. *Agr Ecosyst Environ*, 139(3), 363-383, 2010.

875 Cherubini, F., Bird, N.D., Cowie, A., Jungmeier, G., Schlamadinger, B., and Woess-Gallasch,  
876 S.: Energy-and greenhouse gas-based LCA of biofuel and bioenergy systems: Key issues,  
877 ranges and recommendations. *Resour Conserv Recy*, 53(8), 434-447, 2009.

878 Chesworth, W.: Encyclopedia of soil science. *Encyclopedia of Earth Sciences Series*,  
879 Springer, Dordrecht, The Netherland, 2008.

880 Conen, F., Yakutin, V., and Sambuu, A.D.: Potential for detecting changes in soil organic  
881 carbon concentrations resulting from climate change. *Glob Change Biol*, 9, 1515–1520, 2003

882 Crutzen, P.J., Mosier, A.R., Smith, K.A., and Winiwarter, W.: N<sub>2</sub>O release from agro-biofuel  
883 production negates global warming reduction by replacing fossil fuels. *Atmos Chem Phys*, 8,  
884 389–395, 2008.

885 Davis, S.C., Anderson-Teixeira, K.J., and DeLucia, E.H.: Life-cycle analysis and the ecology  
886 of biofuels. *Trends Plant Sci*, 14(3), 140-146, 2009.

887 Dengel, S., Levy, P.E., Grace, J., Jones, S.K., and Skiba, U.M.: Methane emissions from  
888 sheep pasture, measured with an open-path eddy covariance system. *Glob Change Biol*,  
889 17(12), 3524-3533, 2011.

890 Deckmyn, G., Muys, B., Garcia Quijano, J., and Ceulemans, R.: Carbon sequestration  
891 following afforestation of agricultural soils: comparing oak/beech forest to short-rotation  
892 poplar coppice combining a process and a carbon accounting model. *Glob Change Biol*,  
893 10(9), 1482-1491, 2004.

894 Dillen, S.Y., Djomo, S.N., Al Afas, N., Vanbeveren, S., and Ceulemans, R.: Biomass yield  
895 and energy balance of a short-rotation poplar coppice with multiple clones on degraded land  
896 during 16 years. *Biomass Bioenerg*, 56, 157-165, 2013.

897 Djomo, S.N., El Kasmioui, O., and Ceulemans, R.: Energy and greenhouse gas balance of  
898 bioenergy production from poplar and willow: a review. *Global Change Biology Bioenergy*,  
899 3(3), 181–197, doi:10.1111/j.1757-1707.2010.01073.x, 2011

900 Djomo, S.N., El Kasmioui, O., De Groote, T., Broeckx, L.S., Verlinden, M.S., Berhongaray,  
901 G., Fichot, R., Zona, D., Dillen, S.Y., King, J.S., Janssens, I.A., and Ceulemans, R.: Energy  
902 and climate benefits of bioelectricity from low-input short rotation woody crops on  
903 agricultural land over a two-year rotation. *Appl Energ*, 111, 862-870, 2013.

904 Don, A., Osborne, B., Hastings, A., Skiba, U., Carter, M.S., Drewer, J., Flessa, H., Freibauer,  
905 A., Hyvönen, N., Jones, M.B., Lanigan, G.J., Mander, Ü., Monti, A., Djomo, S.N., Valentine,  
906 J., Walter, K., Zegada-Lizarazu, W., and Zenone, T.: Land-use change to bioenergy  
907 production in Europe: implications for the greenhouse gas balance and soil carbon. *Global*  
908 *Change Biology Bioenergy*, 4(4), 372-391, 2012.

909 Dong, H., Mangino, J., and McAllister, T.A.: Emissions from livestock and manure  
910 management, in: 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Prepared  
911 by the National Greenhouse Gas Inventories Programme (eds. Eggleston, H.S., Buendia, L.,  
912 Miwa, K., Ngara, T., and Tanabe, K.), Hayama, Japan, PP. 10.1-10.89., 2006.

913 Dougherty, T.C., and Hall, A. W.: Environmental impact assessment of irrigation and  
914 drainage projects. *Irrigation and Drainage Paper 53*, FAO, Rome, 1995.

915 Edwards, R., Larivé, J.F., Mahieu, V., and Rouveiolles, P.: Well-to-wheels analysis of future  
916 automotive fuels and powertrains in the European context, *Tank-to-Wake Report Version 2c*,  
917 EUCAR, CONCAWE, & JRC, Ispra, Italy, 2007.

918 El Kasmoui, O., and Ceulemans, R.: Financial analysis of the cultivation of poplar and  
919 willow for bioenergy. *Biomass Bioenerg*, 43, 52-64, 2012.

920 El Kasmoui, O., and Ceulemans, R.: Financial Analysis of the Cultivation of Short Rotation  
921 Woody Crops for Bioenergy in Belgium: Barriers and Opportunities. *BioEnergy Research*, 6,  
922 336-350, 2013

923 European Commission: Renewable energy road map—renewable energies in the 21st century:  
924 building a more sustainable future. COM(2006) 848 final, Brussels, Belgium, 2007.

925 European Commission: 20 20 by 2020—Europe’s climate change opportunity. COM(2008)  
926 30, Brussels, Belgium, 2008

927 European Commission: Directive 2009/28/Ec Of The European Parliament And Of The  
928 Council of 23 April 2009 on the promotion of the use of energy from renewable sources and



929 amending and subsequently repealing Directives 2001/77/EC and 2003/30/EC Official  
 930 Journal of the European Union, L 140/16-62, EU, Brussels, Belgium, 2009.

931 Fargione, J., Hill, J., Tilman, D., Polasky, S., and Hawthorne, P.: Land clearing and the  
 932 biofuel carbon debt. *Science*, 319(5867), 1235-1238, 2008.

933 Fisher, R.: The Correlation between Relatives on the Supposition of Mendelian Inheritance. *T*  
 934 *Rse Earth*, 52(02), 399-433, 1919.

935 Foken, T.: *Micrometeorology*, Eds. Nappo C.J., Springer Verlag, Berlin-Heidelberg,  
 936 Germany, 308 pp., 2008.

937 Forster, P., Ramaswamy, V., Artaxo, P., Berntsen, T., Betts, R., Fahey, D.W., Haywood, J.,  
 938 Lean, J., Lowe, D.C., Myhre, G., Nganga, J., Prinn, R., Raga, G., Schulz, M., and Van  
 939 Dorland, R.: Changes in Atmospheric Constituents and in Radiative Forcing, in: *Climate*  
 940 *Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth*  
 941 *Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Solomon,  
 942 S., Qin, D., Manning, M., Chen. Z., Marquis, M., Averyt, K. B., Tignor, M., and Miller, H. L.,  
 943 Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, PP.  
 944 129-234, 2007.

945 Gash, J.H.C., and Culf, A.D.: Applying a linear detrend to eddy correlation data in realtime.  
 946 *Bound-Lay Meteorol*, 79(3), 301-306, 1996.

947 Gasol, C.M., Gabarrell, X., Anton, A., Rigola, M., Carrasco, J., Ciria, P., and Rieradevall, J.:  
 948 LCA of poplar bioenergy system compared with Brassica carinata energy crop and natural gas  
 949 in regional scenario. *Biomass Bioenerg*, 33(1), 119–129, 2009.

950 Gauder, M., Butterbach-Bahl, K., Graeff-Hönninger, S., Claupein, W., and Wiegel, R.: Soil-  
 951 derived trace gas fluxes from different energy crops—results from a field experiment in  
 952 Southwest Germany. *Global Change Biology Bioenergy*, 4(3), 289-301, 2012.

953 Gelfand, I., Zenone, T., Jasrotia, P., Chen, J., Hamilton, S.K., and Robertson, G.P.: Carbon  
 954 debt of Conservation Reserve Program (CRP) grasslands converted to bioenergy production.  
 955 P Natl Acad Sci Usa, 108, 13864-13869, 2011.

956 Gielen, B., De Vos, B., Campioli, M., Neirynck, J., Papale, D., Verstraeten, A., Ceulemans,  
 957 R., and Janssens, I.A.: Biometric and eddy covariance-based assessment of decadal carbon  
 958 sequestration of a temperate Scots pine forest. Agr Forest Meteorol, 174, 135-143, 2013.

959 Gilmanov, T.G., Soussana, J.F., Aires, L., Allard, V., Ammann, C., Balzarolo, M., Barcza, Z.,  
 960 Bernhofer, C., Campbell, C.L., Cernusca, A., Cescatti, A., Clifton-Brown, J., Dirks, B.O.M.,  
 961 Dore, S., Eugster, W., Fuhrer, J., Gimeno, C., Gruenwald, T., Haszpra, L., Hensen, A., Ibrom,  
 962 A., Jacobs, A.F.G., Jones, M.B., Lanigan, G., Laurila, T., Lohila, A., Manca, G., Marcolla, B.,  
 963 Nagy, Z., Pilegaard, K., Pinter, K., Pio, C., Raschi, A., Rogiers, N., Sanz, M.J., Stefani, P.,  
 964 Sutton, M., Tuba, Z., Valentini, R., Williams, M.L., and Wohlfahrt, G.: Partitioning  
 965 European grassland net ecosystem CO<sub>2</sub> exchange into gross primary productivity and  
 966 ecosystem respiration using light response function analysis. Agr Ecosyst Environ, 121(1),  
 967 93-120, 2007.

968 Goodman, L.A.: On the exact variance of products. J Am Stat Assoc, 55(292), 708-713, 1960.

969 Grigal, D.F., and Berguson, W.E.: Soil carbon changes associated with short-rotation systems.  
 970 Biomass Bioenerg, 14(4), 371-377, 1998.

971 Gundersen, P., Christiansen, J.R., Alberti, G., Brüggemann, N., Castaldi, S., Gasche, R.,  
 972 Kitzler, B., Klemedtsson, L., Lobo-do-Vale, R., Moldan, F., Rütting, T., Schleppi, P.,  
 973 Weslien, P., and Zechmeister-Boltenstern, S.: The response of methane and nitrous oxide  
 974 fluxes to forest change in Europe. Biogeosciences, 9, 3999-4012, 2012.

975 Italian Ministry of Interior: Bilancio energetico nazionale 2012. Dipartimento per l'energia,  
 976 Direzione generale per la sicurezza dell'approvvigionamento e le infrastrutture energetiche,  
 977 available at: <http://dgerm.sviluppoeconomico.gov.it/dgerm/ben.asp>, (last access: 25 February

2015), 2013.

Jassal, R.S., Black, T.A., Arevalo, C., Jones, H., Bhatti, J.S., and Sidders, D.: Carbon sequestration and water use of a young hybrid poplar plantation in north-central Alberta. *Biomass Bioenerg*, 56, 323-333, 2013.

Jaoudé, R.A., Lagomarsino, A., and De Angelis, P.: Impacts of nitrogen fertilisation and coppicing on total and heterotrophic soil CO<sub>2</sub> efflux in a short rotation poplar plantation. *Plant Soil*, 339(1-2), 219-230, 2011.

Jungmeier, G., and Spitzer, J.: Greenhouse gas emissions of bioenergy from agriculture compared to fossil energy for heat and electricity supply. *Nutr Cycl Agroecosys*, 60(1-3), 267-273, 2001.

Kandeler, E.: Nitrification during long term incubation, in: *Methods in soil biology* (eds. Schinner, F., Kandeler, E., Ohlinger, R., and Margesin, R.), Springer, Berlin Heidelberg New York, USA, pp. 149-151, 1996.

Laganiere, J., Angers, D.A., and Pare, D.: Carbon accumulation in agricultural soils after afforestation: a meta-analysis. *Glob Change Biol*. 16, 439–453, 2010.

Latium Region, Regional Direction of Agriculture and Rural Development, Hunting and Fishing: Rural Development Program of Latium 2007- 2013. Reg. (CE) n. 1698/05, Rome, available at: <http://www.agricoltura.regione.lazio.it/psr/programma/?vms=5&ref=Il-testo-integrale-psr> (Last access: 20 October 2015), 2015.

Liu, Y., Langer, V., Høgh-Jensen, H., and Egelyng, H.: Life cycle assessment of fossil energy use and greenhouse gas emissions in Chinese pear production. *J Clean Prod*, 18(14), 1423-1430, 2010.

Mareschi, L.: Piantagioni da bioenergia: sostenibilit  produttiva ed ambientale di nuovi cloni ibridi di pioppo (*Populus* spp.), Ph.D. thesis, University of Tuscia, Italy, 2008.

1002 Martiniello, P.: Effects of irrigation and harvest management on dry-matter yield and seed  
 1003 yield of annual clovers grown in pure stand and in mixtures with graminaceous species in a  
 1004 Mediterranean environment. *Grass Forage Sci*, 54(1), 52-61, 1999.

1005 Migliavacca, M., Meroni, M., Manca, G., Matteucci, G., Montagnani, L., Grassi, G., Zenone,  
 1006 T., Teobaldelli, M., Goded, I., Colombo, R., and Seufert, G.: Seasonal and interannual  
 1007 patterns of carbon and water fluxes of a poplar plantation under peculiar eco-climatic  
 1008 conditions. *Agr Forest Meteorol*, 149(9), 1460-1476, 2009.

1009 Mikutta, R., Kleber, M., Kaiser, K., and Jahn, R.: Review: Organic Matter Removal from  
 1010 Soils using Hydrogen Peroxide, Sodium Hypochlorite, and Disodium Peroxodisulfate. *Soil*  
 1011 *Sci Soc Am J*, 6, 120-135, 2005.

1012 Mitchell, C.P., Stevens, E.A., and Watters, M.P.: Short-rotation forestry— operations,  
 1013 productivity and costs based on experience gained in the UK. *Forest Ecol Manag*, 121(1–2),  
 1014 123–136, 1999.

1015 Moncrieff, J.B., Massheder, J.M., De Bruin, H., Elbers, J., Friborg, T., Heusinkveld, B.,  
 1016 Kabat, P., Scotta, S., Soegaard, H., and Verhoef, A.: A system to measure surface fluxes of  
 1017 momentum, sensible heat, water vapour and carbon dioxide. *J Hydrol*, 188, 589-611, 1997.

1018 Nasso, N., Guidi, W., Ragagnoli, G., Tozzini, C., and Bonari, E.: Biomass  
 1019 production and energy balance of a 12-year-old short-rotation coppice poplar stand under  
 1020 different cutting cycles. *Global Change Biology Bioenergy*, 2(2), 89-97, 2010.

1021 Nasso, N., Bosco, S., Di Bene, C., Coli, A., Mazzoncini, M., and Bonari, E.:  
 1022 Energy efficiency in long-term Mediterranean cropping systems with different management  
 1023 intensities. *Energy*, 36(4), 1924-1930, 2011.

1024 Osborne, B., Saunders, M., Walmsley, D., Jones, M., and Smith, P.: Key questions and  
 1025 uncertainties associated with the assessment of the cropland greenhouse gas balance. *Agr*  
 1026 *Ecosyst Environ*, 139(3), 293-301, 2010.

1027 Pearson, K.: On the criterion that a given system of deviations from the probable in the case  
 1028 of a correlated system of variables is such that it can be reasonably supposed to have arisen  
 1029 from random sampling. The London, Edinburgh, and Dublin Philosophical Magazine and  
 1030 Journal of Science, 50(302), 157-175, 1900.

1031 Papale, D., Reichstein, M., Aubinet, M., Canfora, E., Bernhofer, C., Kutsch, W., Longdoz, B.,  
 1032 Rambal, S., Valentini, R., Vesala, T., and Yakir, D.: Towards a standardized processing of  
 1033 Net Ecosystem Exchange measured with eddy covariance technique: algorithms and  
 1034 uncertainty estimation. Biogeosciences, 3, 571-583, 2006.

1035 Paris, P., Mareschi, L., Sabatti, M., Pisanelli, A., Ecosse, A., Nardin, F., and Scarascia-  
 1036 Mugnozza, G.: Comparing hybrid Populus clones for SRF across northern Italy after two  
 1037 biennial rotations: Survival, growth and yield. Biomass Bioenerg, 35(4), 1524-1532, 2011.

1038 Pihlatie, M., Pumpanen, J., Rinne, J., Ilvesniemi, H., Simojoki, A., Hari, P., and Vesala, T.:  
 1039 Gas concentration driven fluxes of nitrous oxide and carbon dioxide in boreal forest soil.  
 1040 Tellus B, 59(3), 458-469, 2007.

1041 Post, W., and Kwon, K.: Soil carbon sequestration and land-use change: processes and  
 1042 potential. Glob Change Biol, 6, 317-328, 2000.

1043 Price, D., Orchansky, A., Jassal, R., Arevalo, C., and Thomas, B.: Soil respiration in a hybrid  
 1044 poplar plantation located in central Alberta, Alberta Soil Science Workshop 2009, Edmonton,  
 1045 Alberta, CA, 2009.

1046 Reichstein, M., Falge, E., Baldocchi, D., Papale, D., Aubinet, M., Berbigier, P., Bernhofer, C.,  
 1047 Buchmann, N., Gilmanov, T., Granier, A., Grünwald, T., Havránková, K., Ilvesniemi, H.,  
 1048 Janous, D., Knohl, A., Laurila, T., Lohila, A., Loustau, D., Matteucci, G., Meyers, T.,  
 1049 Miglietta, F., Ourcival, J.M., Pumpanen, J., Rambal, S., Rotenberg, E., Sanz, M., Tenhunen,  
 1050 J., Seufert, G., Vaccari, F., Vesala, T., Yakir, D., and Valentini, R.: On the separation of net

ecosystem exchange into assimilation and ecosystem respiration, review and improved algorithm. *Glob Change Biol*, 11, 1424–1439, 2005.

Robertson, G.P., Paul, E.A., and Harwood, R.R.: Greenhouse gases in intensive agriculture: contributions of individual gases to the radiative forcing of the atmosphere. *Science*, 289(5486), 1922-1925, 2000.

Romano, D., Arcarese, C., Bernetti, A., Caputo, A., Córdor, R.D., Contaldi, M., De Lauretis, R., Di Cristofaro, E., Gagna, A., Gonella, B., Lena, F., Liburdi, R., Taurino, E., and Vitullo, M.: Italian Greenhouse Gas Inventory 1990–2012, National Inventory Report 2014, ISPRA – Institute for Environmental Protection and Research, Rome, 2014.

Saidur, R., Abdelaziz, E.A., Demirbas, A., Hossain, M.S., and Mekhilef, S.: A review on biomass as a fuel for boilers. *Renew Sust Energ Rev*, 15(5), 2262-2289, 2011.

Scholz, V., and Ellerbrock, R.: The growth productivity, and environmental impact of the cultivation of energy crops on sandy soil in Germany. *Biomass Bioenerg*, 23(2), 81-92, 2002.

Smith, P., Lanigan, G., Kutsch, W.L., Buchmann, N., Eugster, W., Aubinet, M., Ceschia, E., Béziat, P., Yeluripati, J.B., Osborne, B., Moors, E.J., Brut, A., Wattenbach, M., Saunders, M., and Jones, M.: Measurements necessary for assessing the net ecosystem carbon budget of croplands. *Agr Ecosyst Environ*, 139(3), 302-315, 2010.

Soussana, J.F., Allard, V., Pilegaard, K., Ambus, P., Amman, C., Campbell, C., Ceschia, E., Clifton-Brown, J., Czobel, S., Domingues, R., Flechard, C., Fuhrer, J., Hensen, A., Horvath, L., Jones, M., Kasper, G., Martin, C., Nagy, Z., Neftel, A., Raschi, A., Baronti, S., Rees, R.M., Skiba, U., Stefani, P., Manca, G., Sutton, M., Tuba, Z., and Valentini, R.: Full accounting of the greenhouse gas (CO<sub>2</sub>, N<sub>2</sub>O, CH<sub>4</sub>) budget of nine European grassland sites. *Agr Ecosyst Environ*, 121(1), 121-134, 2007.

1074 Soussana, J.F., Tallec, T., and Blanfort, V.: Mitigating the greenhouse gas balance of  
 1075 ruminant production systems through carbon sequestration in grasslands. *Animal*, 4(03), 334-  
 1076 350, 2010.

1077 Strauss, C.H., and Grado, S.C.: Economics of producing *Populus* biomass for energy and fiber  
 1078 systems, in: *Micropropagation, genetic engineering, molecular biology of Populus*. (eds.  
 1079 Klopfenstein, N.B., Chun, Y.W., Kim, M-S., and Ahuja, M.R.), USDA Forest Service, Fort  
 1080 Collins, CO, Rocky Mountain Forest and Range Experiment Station, pp. 241-248, 1997.

1081 USS Working Group WRB: World Reference Base for Soil Resources 2014. International soil  
 1082 classification system for naming soils and creating legends for soil maps, World Soil  
 1083 Resources Reports, 106, FAO, Rome, Italy, 2014.

1084 Vickers, D., and Mahrt, L.: Quality control and flux sampling problems for tower and aircraft  
 1085 data. *J Atmos Ocean Tech*, 14(3), 512-526, 1997.

1086 Webb, E.K., Pearman, G.I., and Leuning, R.: Correction of flux measurements for density  
 1087 effects due to heat and water vapour transfer. *Q J Roy Meteor Soc*, 106(447), 85-100, 1980.

1088 Weslien, P., Kasimir Klemetsson, Å., Börjesson, G., and Klemetsson, L.: Strong pH  
 1089 influence on N<sub>2</sub>O and CH<sub>4</sub> fluxes from forested organic soils. *Eur J Soil Sci*, 60(3), 311-320,  
 1090 2009.

1091 Wood, S.W., and Cowie, A.: A review of greenhouse gas emission factors for fertiliser  
 1092 production, International Energy Agency Bioenergy Task 38, 1-20, 2004

1093 Zhou, J., Zhang, Z., Sun, G., Fang, X., Zha, T., McNulty, S., Chen, J., Jin, Y., and Noormets,  
 1094 A.: Response of ecosystem carbon fluxes to drought events in a poplar plantation in Northern  
 1095 China. *Forest Ecol Manag*, 300, 33-42, 2013.

1096 Zona, D., Janssens, I.A., Aubinet, M., Vicca, S., Gioli, B., Fichot, R., and Ceulemans, R.:  
 1097 Fluxes of the greenhouse gases (CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O) above a short-rotation poplar plantation  
 1098 after conversion from agricultural land. *Agr Forest Meteorol*, 169, 100-110, 2013.

## 9. Tables

**Table 1 – Instrumental setup of the two towers. SRC = short rotation coppice site; REF = reference site; T = temperature; RH = relative humidity; PAR = photosynthetically active radiation;  $M_{SOIL}$  = soil water content; P = precipitation; EC = eddy covariance; prof = profile. 4-component radiometers were used to measure short- and long-wave radiations, and derive net radiation. SRC site soil profiles were located in irrigated and not-irrigated inter-rows. Precipitation was assumed to be consistent in the two ecosystems.**

|                              | SRC   | REF   |
|------------------------------|---|---|
| <b>Meteo</b>                 |   |   |
| <i>Air T and RH</i>          | MP-100, Rotronic AG, Bassersdorf, CH                    | MP-100, Rotronic AG, Bassersdorf, CH                    |
| <i>PAR</i>                   | Li-190, LI-COR, Lincoln, NE, USA                        | -   |
| <i>Radiations</i>            | CNR-1, Kipp&Zonen, Delft, NL                            | NR01, Hukseflux, Delft, NL                              |
| <i><math>M_{SOIL}</math></i> | CS616, Campbell Scientific, Logan, UT, USA (2<br>prof.) | CS616, Campbell Scientific, Logan, UT, USA (1<br>prof.) |
| <i>Soil T</i>                | 107, Campbell Scientific, Logan, UT, USA (2 prof.)      | 107, Campbell Scientific, Logan, UT, USA (1<br>prof.)   |
| <i>Soil heat flux</i>        | HFT3, REBS Inc., Seattle, WA, USA                       | HFP01, Hukseflux, Delft, NL                             |
| <i>P</i>                     | -   | ARG100, EML, North Shield, UK                           |
| <i>Logger</i>                | CR3000, Campbell Scient., Logan, UT, USA                | CR1000 Campbell Scient. Logan, UT, USA                  |
| <b>EC</b>                    |   |   |
| <i>Anemometer</i>            | CSAT3, Campbell Scientific, Logan, UT, USA              | USA-1, Metek GmbH, Elmshorn, DE                         |
| <i>Gas-Analyser</i>          | LI-7500, LI-COR, Lincoln, NE, USA                       | LI-7500A, LI-COR, Lincoln, NE, USA                      |



1107

**Table 2 – Correction steps applied to the time series using LICOR EddyPro software.**

| <b>Correction</b>                                    | <b>Reference</b>         |
|--|--------------------------|
| Despiking  | Vickers and Mahrt (1997) |
| Density fluctuations                                 | Webb et al. (1980)       |
| Maximisation of covariance for time lag compensation | Aubinet et al. (2000)    |
| Linear detrending for trend removal                  | Gash and Culf (1996)     |
| 2-D coordinate rotation                              | Wilczak et al. (2001)    |
| High-pass filtering effect                           | Moncrieff et al. (1997)  |
| Low-pass filtering effect                            | Ibrom et al. (2007)      |

1108

1109

1110 **Table 3 – Farming activities. Three tractors of different power were normally used to collect chips: two of the type 1,**  
1111 **and one of the type 2. DAP = diammonium phosphate; AN = ammonium nitrate; CAN = calcium ammonium nitrate.**  
1112 **SRC and REF as defined previously. Reported units are intended per hectare and activity.**

| Operation                   | Fuel consumption<br>(unit ha <sup>-1</sup> ) | Input rates<br>(unit ha <sup>-1</sup> ) | Site     |
|-----------------------------|--|---|----------|
| Harvesting – wood chipper   | 30 l diesel                                  | -                                       | SRC      |
| Harvesting – Tractor type 1 | 20 l diesel                                  | -                                       | SRC      |
| Harvesting – Tractor type 2 | 10 l diesel                                  | -                                       | SRC      |
| Shallow tillage             | 8 l diesel                                   | -                                       | SRC, REF |
| Application of insecticide  | 1.125 l diesel                               | 1.25 kg DECIS®                          | SRC      |
| Mechanical weeding          | 4 l diesel                                   | -                                       | SRC      |
| Ploughing                   | 8 l diesel                                   | -                                       | SRC, REF |
| Sowing                      | 2 l diesel                                   | -                                       | REF      |
| Seed covering               | 4 l diesel                                   | -                                       | REF      |
| Application of fertiliser   | 2 l diesel                                   | a. 150 kg DAP                           | a. REF   |
|                             |  | b. 150 kg AN                            | b. REF   |
|                             |  | c. 200 kg CAN                           | c. REF   |
|                             |  | d. 40 kg Urea                           | d. SRC   |
| Reaping                     | 20 l diesel                                  | -                                       | REF      |
| Chemical weeding            | 1.125 l diesel                               | 1 l Buctril®                            | REF      |
| Bale                        | 7.5 l diesel                                 | -                                       | REF      |
| Irrigation                  | a. 471 kWh electricity                       | a. 16 l H <sub>2</sub> O                | a. SRC   |
|                             | b. 149 kWh electricity                       | b. 46 l H <sub>2</sub> O                | b. REF   |

1113

1114



1119 **Table 5 – Soil characteristics of the layer 0-15 cm. SRC and REF as previously defined; SOC = soil organic carbon;**  
 1120  $\rho_{BULK}$  = bulk density. Superscripts a-c indicate statistically significant differences between the means of SOC.

| 1121 | Site  | Variable                            | Value $\pm$ dev. std.           |
|------|-------|-------------------------------------|---------------------------------|
| 1122 | REF   | C (%)                               | 1.46 $\pm$ 0.34                 |
|      |       | $\rho_{BULK}$ (Mg m <sup>-3</sup> ) | 1.00 $\pm$ 0.11                 |
|      |       | SOC (MgC ha <sup>-1</sup> )         | 16.03 $\pm$ 3.76 <sup>(a)</sup> |
|      | SRC   | C (%)                               | 1.05 $\pm$ 0.40                 |
|      |       | $\rho_{BULK}$ (Mg m <sup>-3</sup> ) | 1.12 $\pm$ 0.15                 |
|      |       | SOC (MgC ha <sup>-1</sup> )         | 11.69 $\pm$ 4.42 <sup>(b)</sup> |
|      | O_SRC | C (%)                               | 1.38 $\pm$ 0.27                 |
|      |       | $\rho_{BULK}$ (Mg m <sup>-3</sup> ) | 1.02 $\pm$ 0.11                 |
|      |       | SOC (MgC ha <sup>-1</sup> )         | 14.03 $\pm$ 2.79 <sup>(c)</sup> |

# 10. Figures

Fig. 1 – Scheme of the chronological land cover during the cultivation cycle taken into account for GHG budget calculation in the two ecosystems. The expected harvest of poplar at the beginning of 2014 was postponed of one year: for that reason data from the previous harvest (beginning 2012) were taken into account for GHG budget calculation. Textures indicate different land cover type, symbols mark the most important management practices, straight lines indicate the periods in which sites were irrigated, dashed line period of grazing. SRC = short rotation coppice site; REF = reference site; in the x axis dates are reported as month-year (mm-yy)




Fig. 2 – Boxplot of the 24-month cumulative fluxes of net ecosystem exchange of CO<sub>2</sub> ( $F_{CO_2}$ ) (a), gross primary production (GPP) (b) and ecosystem respiration (Reco) (c) from eddy covariance (EC) data in the REF and SRC sites. Each box represents the range 16th-84th percentile: the central mark is the median, while the whiskers extend to the 5th and 95th percentiles




Fig. 3 – Monthly averages of  $F_{CO_2}$  in the REF and SRC sites (top panel). The bottom panel shows monthly averages of air temperature ( $T_{air}$ ) and soil water content (SWC) at 30 cm depth. In both subplots dotted lines are used for the SRC site and continuous lines for the REF site, while in the bottom panel SWC is in grey and the  $T_{air}$  in black.





Fig. 4 – Fluxes of soil N<sub>2</sub>O (crosses) and CH<sub>4</sub> (circles) in the SRC (a – c) and the REF (b – d) sites. Each marker represents the average of the nine chambers, with bars indicating their standard deviation. First letter of month in the x-axis





**Fig. 5 – N<sub>2</sub>O fluxes from incubation experiment reported in function of the water filled pore space estimated for each single replicate. In (a) data from samples taken in the SRC site are shown, in (b) data from REF site samples**




Fig. 6 – WFPS% in the REF site at three different depths for the 24-month integration periods. Dashed line points to the threshold (70%) unleashing N<sub>2</sub>O from lab incubations. First letter of month in the x-axis




Fig. 7 –GHG emissions of the different farming operations. Harv = harvesting; plow = ploughing; sow = sowing; irr = irrigation; fert = fertilisation; othe= minor contributions. SRC and REF as previously defined




Fig. 8 – Regression line of SOC content in time  $t$  (years). The gap between SOC(0) and SOC content in the REF site represented the loss of SOC for the land use change. Est = estimated values; meas = measured values; SRC and REF as previously defined; O\_SRC is the older short rotation coppice site used to build the regression




Fig. 9 – GHG balances of the SRC and the REF sites: components (left) and net (right).  $F_{CH_4}$  and  $F_{N_2O}$  from soil are negligible and not inserted in the graph.  $F_{MAN}$  = management;  $E_{CH_4}$  = exported biomass reemitted as  $CH_4$  by enteric fermentation;  $E_{CO_2}$  = exported biomass reemitted as  $CO_2$  by sheep respiration;  $F_{SOC}$  = initial SOC change at the installation of cuttings;  $F_{SAV}$  = GHG savings for replacement of fossil fuel use;  $F_{CO_2}$  as previously defined

