

This discussion paper is/has been under review for the journal Biogeosciences (BG).
Please refer to the corresponding final paper in BG if available.

Phytoplankton dynamics in contrasting early stage North Atlantic spring blooms: composition, succession, and potential drivers

C. J. Daniels¹, A. J. Poulton², M. Esposito², M. L. Paulsen³, R. Bellerby^{4,5,6},
M. St. John⁷, and A. P. Martin²

¹Ocean and Earth Sciences, National Oceanography Centre Southampton, University of Southampton, Southampton, UK

²Ocean Biogeochemistry and Ecosystems, National Oceanography Centre, University of Southampton Waterfront Campus, UK

³Department of Biology, Marine Microbiology Department, University of Bergen, Bergen, Norway

⁴Norwegian Institute for Water Research (NIVA), Bergen, Norway

⁵Uni Bjerknnes Centre, Bergen, Norway

⁶Bjerknnes Center for Climate Research, Bergen, Norway

⁷National Institute of Aquatic Resources, Technical University of Denmark, Charlottenlund, Denmark

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Received: 1 December 2014 – Accepted: 5 December 2014 – Published: 6 January 2015

Correspondence to: C. J. Daniels (c.daniels@noc.soton.ac.uk)

Published by Copernicus Publications on behalf of the European Geosciences Union.

BGD

12, 93–133, 2015

**Spring bloom
phytoplankton
composition**

C. J. Daniels et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

The spring bloom is a key annual event in the phenology of pelagic ecosystems, making a major contribution to the oceanic biological carbon pump through the production and export of organic carbon. However, there is little consensus as to the main drivers of spring bloom formation, exacerbated by a lack of in situ observations of the phytoplankton community composition and its evolution during this critical period.

We investigated the dynamics of the phytoplankton community structure at two contrasting sites in the Iceland and Norwegian Basins during the early stage (25 March–25 April) of the 2012 North Atlantic spring bloom. The plankton composition and characteristics of the initial stages of the bloom were markedly different between the two basins. The Iceland Basin (ICB) appeared well mixed to > 400 m, yet surface chlorophyll *a* (0.27–2.2 mg m⁻³) and primary production (0.06–0.66 mmol C m⁻³ d⁻¹) were elevated in the upper 100 m. Although the Norwegian Basin (NWB) had a persistently shallower mixed layer (< 100 m), chlorophyll *a* (0.58–0.93 mg m⁻³) and primary production (0.08–0.15 mmol C m⁻³ d⁻¹) remained lower than in the ICB, with picoplankton (< 2 μm) dominating chlorophyll *a* biomass. The ICB phytoplankton composition appeared primarily driven by the physicochemical environment, with periodic events of increased mixing restricting further increases in biomass. In contrast, the NWB phytoplankton community was potentially limited by physicochemical and/or biological factors such as grazing.

Diatoms dominated the ICB, with the genus *Chaetoceros* (1–166 cells mL⁻¹) being succeeded by *Pseudo-nitzschia* (0.2–210 cells mL⁻¹). However, large diatoms (> 10 μm) were virtually absent (< 0.5 cells mL⁻¹) from the NWB, with only small nano-sized (< 5 μm) diatoms present (101–600 cells mL⁻¹). We suggest micro-zooplankton grazing, potentially coupled with the lack of a seed population of bloom forming diatoms, was restricting diatom growth in the NWB, and that large diatoms may be absent in NWB spring blooms. Despite both phytoplankton communities being in the early stages of bloom formation, different physicochemical and biological factors controlled

BGD

12, 93–133, 2015

Spring bloom phytoplankton composition

C. J. Daniels et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



bloom formation at the two sites. If these differences in phytoplankton composition persist, the subsequent spring blooms are likely to be significantly different in terms of biogeochemistry and trophic interactions throughout the growth season, with important implications for carbon cycling and organic matter export.

1 Introduction

The spring bloom is a key annual event in the phenology of pelagic ecosystems, where a rapid increase in phytoplankton biomass has a significant influence on upper ocean biogeochemistry and food-availability for higher trophic levels (Townsend et al., 1994; Behrenfeld and Boss, 2014). Spring blooms are particularly prevalent in coastal and high latitude waters. The high levels of phytoplankton biomass and primary production that occur during these blooms, and its subsequent export out of the surface ocean, result in a significant contribution to the biological carbon pump (Townsend et al., 1994; Sanders et al., 2014). The North Atlantic spring bloom is one of the largest blooms on Earth, making a major contribution to the annual export of $\sim 1.3 \text{ GtCyr}^{-1}$ from the North Atlantic (Sanders et al., 2014). The timing and magnitude of the spring bloom can have a significant biogeochemical impact (Henson et al., 2009); hence it is important to understand both the controls on, and the variability in, bloom timing and magnitude. Despite its importance, there remains little consensus as to the environmental and ecological conditions required to initiate high latitude spring blooms (Townsend et al., 1994; Behrenfeld, 2010; Taylor and Ferrari, 2011b).

Phytoplankton blooms occur when growth rates exceeds loss rates (i.e. a sustained period of net growth); phytoplankton growth rate constraints include irradiance, nutrient supply, and temperature, while losses can occur through predation, advection, mixing out of the euphotic zone, sinking and viral attack (Miller, 2003). Therefore, the rapid increase in (net) growth rates during the spring bloom must be due to either an alleviation of those factors constraining growth, a reduction in factors determining losses, or (more likely) some combination of both.

Spring bloom phytoplankton composition

C. J. Daniels et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



The critical depth hypothesis (Sverdrup, 1953), the seminal theory of spring bloom initiation, proposes that there exists a critical depth such that when stratification shoals above this depth, phytoplankton growth will exceed mortality and a bloom will occur. However, this hypothesis has been more recently brought into question as bloom formation has been observed to start earlier than expected (Mahadevan et al., 2012), and in the absence of stratification (Townsend et al., 1992; Eilertsen, 1993). Several new theories have now been developed to explain these occurrences (reviewed in Behrenfeld and Boss, 2014; Fischer et al., 2014; Lindemann and St. John, 2014).

Eddies and oceanic fronts have both been identified as sources of stratification prior to the wider onset of seasonal stratification (Taylor and Ferrari, 2011a; Mahadevan et al., 2012). However, they do not explain blooms in the complete absence of stratification, which can instead be explained by the critical turbulence hypothesis (Huisman et al., 1999; Taylor and Ferrari, 2011b; Brody and Lozier, 2014). Both of these theories distinguish between a convectively driven actively mixed layer and a density-defined mixed layer such that if convective mixing reduces sufficiently, blooms can occur in the actively mixing layer although the density-defined mixing layer remains deep. Therefore, blooms are able to form in the apparent absence of stratification, as defined by the presence of a thermocline. An alternative to the hypotheses concerning physical controls on bloom formation is that proposed by Behrenfeld (2010), who suggests that the decoupling of phytoplankton and micro-zooplankton contact rates in deep winter mixed layers results in phytoplankton net growth from winter onwards due to reduced mortality (via grazing). It is also possible that there are multiple biological and physical controls, acting on different spatial and temporal scales, that drive the heterogeneous bloom distributions observed via remote sensing (e.g. Lindemann and St. John, 2014).

Significant interannual and decadal variability in the structure and timing of spring blooms in the North Atlantic has been documented (Henson et al., 2009). Such variability in bloom timing has been attributed to the variation in the winter mixed layer depth (WMLD); a deeper WMLD results in a delayed bloom in the subarctic North Atlantic (Henson et al., 2009). A strong latitudinal trend exists in the North Atlantic where

Spring bloom phytoplankton composition

C. J. Daniels et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the spring bloom propagates north due to seasonal relief from light limitation at high latitudes (Siegel et al., 2002; Henson et al., 2009). Both the role of the WMLD in interannual variability in bloom timing and the northwards progression of bloom start dates highlight how physical processes have a clear and significant impact on bloom formation. The controls on the variability in bloom magnitude are less certain, although it appears to be a combination of WMLD variability influencing the start date as well as biological factors such as phytoplankton composition and grazing (Henson et al., 2009).

Despite considerable discussion on the various factors that may or may not influence bloom initiation, timing, magnitude and phenology, few studies have actually examined the in situ phytoplankton community. Instead, because of the need for temporally resolved data, satellite-derived products and models have been used in much of the previous work on spring blooms. However, such methods cannot address the potential influence of the complex plankton community structure on the development of a spring bloom.

The traditional text book view of a phytoplankton spring bloom is that the pre-bloom pico-phytoplankton (cells $< 2 \mu\text{m}$) dominated community is directly succeeded by a diatom dominated community (Margalef, 1978; Barber and Hiscock, 2006); as conditions become more favourable for growth, a diatom bloom develops, “suppressing” growth of other phytoplankton groups. Through either increased predation, nutrient stress or a changing physical environment (Margalef, 1978), diatoms decline and are then replaced by other phytoplankton such as dinoflagellates and coccolithophores (Lochte et al., 1993; Leblanc et al., 2009). In this way, a series of phytoplankton functional type successions occur as the spring bloom develops. That diatoms often dominate intense spring blooms is well accepted (Lochte et al., 1993; Rees et al., 1999), however the dynamics of the interplay between diatoms and the rest of the community have been questioned (Barber and Hiscock, 2006). The rapid proliferation of diatoms in a spring bloom does not necessarily suppress other phytoplankton (Lochte et al., 1993; Barber and Hiscock, 2006), and the “rising tide” hypothesis states that instead of succession,

**Spring bloom
phytoplankton
composition**

C. J. Daniels et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

the favourable conditions for diatoms also favour other phytoplankton groups and therefore all phytoplankton will respond positively and grow (Barber and Hiscock, 2006). The apparent suppression of the phytoplankton community by diatoms is due to the relatively high intrinsic growth rates of diatoms resulting in concentrations dwarfing the rest of the community. The “rising tide” hypothesis is a contrasting theory to succession, however it may be that the phytoplankton community response will not be universal, with some groups being succeeded due to competition or increased grazing (Brown et al., 2008). Furthermore, succession may appear to occur if phytoplankton loss rates are taxonomically specific, such that while many phytoplankton groups concurrently grow, successive loss of specific groups occurs.

The overall goal of our study was to determine the phytoplankton community structure, and its evolution. During the spring bloom in the North Atlantic, linking the community structure to the physical environment and examining whether succession to a diatom dominated environment would occur early in the growth season (March–April). Sampling for this study was carried out as part of the multidisciplinary EuroBASIN “Deep Convection Cruise”. The timing and location of this cruise (19 March–2 May 2012) was chosen to try and observe the transition from deep winter convection to spring stratification, and examine the physical controls on the dynamics of phytoplankton, carbon export and trophic interactions. A recent study has previously suggested that winter convection in the North Atlantic and Norwegian Sea sustains an overwintering phytoplankton population, thus providing an inoculum for the spring bloom (Backhaus et al., 2003), although this transition has not been explicitly examined before.

2 Methods

2.1 Sampling

The Deep Convection cruise repeatedly sampled two pelagic locations in the North Atlantic (Fig. 1), sited in the Iceland (ICB, 61.50° N, 11.00° W) and Norwegian (NWB,

62.83° N, 2.50° W) Basins, onboard the R/V *Meteor*. Samples were collected from multiple casts of a conductivity–temperature–depth (CTD)–Niskin rosette, equipped with a fluorometer, at each station. Water samples for rates of primary production (PP), community structure and ancillary parameters (chlorophyll *a* [Chl *a*], calcite [PIC], particulate silicate [bSiO_2] and macronutrient concentrations) were collected from predawn (02:30–05:00 GMT) casts from six light depths (55, 20, 14, 7, 5 and 1 % of incidental PAR). The depth of 1 % incident irradiance was assumed to equate to the depth of the euphotic zone (e.g. Poulton et al., 2010). Optical depths were determined from a daytime CTD cast on preceding days at each site. Additional samples for community structure and ancillary parameters were collected from a second CTD cast, while samples for detailed size fractionated Chl *a* were collected from a third cast.

2.2 Primary production

Carbon fixation rates were determined using the ^{13}C stable isotope method (Legendre and Gosseline, 1996). Water samples (1.2 L) collected from the six irradiance depths were inoculated with $45\text{--}46\ \mu\text{mol L}^{-1}$ ^{13}C labelled sodium bicarbonate, representing 1.7–1.8 % of the ambient dissolved inorganic carbon pool. Samples were incubated in an on-deck incubator, chilled with sea surface water, and light depths were replicated using optical filters (Misty-blue and Grey, LEETM). Incubations were terminated after 24 h by filtration onto pre-ashed ($> 400^\circ\text{C}$, > 4 h) Whatman GF/F filters. Acid-labile carbon (PIC) was removed by adding 1–2 drops of 1 % HCl to the filter followed by extensive rinsing with freshly filtered (Fisherbrand MF300, $\sim 0.7\ \mu\text{m}$ pore size) unlabelled seawater. Filters were oven dried (40°C , 8–12 h) and stored in Millipore PetriSlidesTM. A parallel 55 % bottle for size fractionated primary production ($< 10\ \mu\text{m}$) was incubated alongside the other samples, with the incubation terminated by pre-filtration through $10\ \mu\text{m}$ polycarbonate (NucleporeTM) filters and the filtrate was filtered and processed as above.

The isotopic analysis was performed on an Automated Nitrogen and Carbon Analysis prep system with a 20–20 Stable Isotope Analyser (PDZ Europa Scientific Instru-

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ments). The ^{13}C -carbon fixation rate was calculated using the equations described in Legendre and Gosseline (1996). The $> 10\ \mu\text{m}$ PP fraction was calculated as the difference between total PP and $< 10\ \mu\text{m}$ PP.

2.3 Community structure

Water samples for diatom and micro zooplankton counts, collected from the predawn casts, were preserved with acidic Lugol's solution (2% final solution) in 100 mL amber glass bottles. Cells were counted in 50 mL Hydro-Bios chambers using a Brunel SP-95-I inverted microscope (X200; Brunel Microscopes Ltd). Samples for flow cytometry were fixed with glutaraldehyde (0.5% final solution) and stored at -80°C before being analysed using a FACS Calibur (Beckton Dickinson) flow cytometer (Zubkov et al., 2007).

Water samples (0.5–1 L) for coccolithophore cell numbers and species identification were collected onto cellulose nitrate filters ($0.8\ \mu\text{m}$ pore size, Whatman), oven dried and stored in Millipore PetriSlides™. Permanent slides of filter halves were prepared and analysed using polarizing light microscopy following Poulton et al. (2010). Coccolithophores were analysed to species level following Frada et al. (2010). For confirmation of species identification, a subset of filter halves were analysed by scanning electron microscope (SEM) following Daniels et al. (2012). Coccolithophore species were identified according to Young et al. (2003).

2.4 Chlorophyll *a*

Water samples (250 mL) for total Chl *a* analysis were filtered onto Fisherbrand MF300 filters. Parallel samples were filtered onto polycarbonate filters ($10\ \mu\text{m}$) for $> 10\ \mu\text{m}$ Chl *a*. Samples for detailed size fractionated Chl *a*, collected from a single depth in the upper water column (12–35 m), were filtered in parallel onto polycarbonate filters of various pore size (2, 10, $20\ \mu\text{m}$) and MF300 filters (effective pore size $0.7\ \mu\text{m}$). Filters were extracted in 8 mL of 90% acetone (Sigma) for 20–24 h (dark, 4°C). Measurements of

BGD

12, 93–133, 2015

Spring bloom phytoplankton composition

C. J. Daniels et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Chl *a* fluorescence were analysed on a Turner Designs Trilogy Fluorometer, calibrated using a solid standard and a chlorophyll *a* extract.

2.5 Ancillary parameters

Particulate inorganic carbon (PIC) measurements were made on water samples (500 mL) filtered onto polycarbonate filters (0.8 μm pore-size, Whatman), rinsed with trace ammonium solution (pH \sim 10) and oven-dried (6–8 h, 30–40 $^{\circ}\text{C}$). The analysis was carried out following Daniels et al. (2012) except that extractions were carried out in 5.0 mL of 0.4 mol L^{-1} nitric acid, erroneously reported as 0.5 mL in Daniels et al. (2012). Particulate silicate (bSiO₂) samples were collected in the same manner as PIC, extracted in 0.2 mol L^{-1} and neutralised with 0.2 mol L^{-1} hydrochloric acid (Brown et al., 2003). The solutions were analysed using a SEAL QuAAtro autoanalyser. Macronutrients (nitrate, phosphate, silicic acid) concentrations were determined following Sanders et al. (2007) on a Skalar autoanalyser.

Samples for total dissolved inorganic carbon (C_T) were drawn into 500 mL borosilicate bottles. No filtering of samples occurred prior to analysis. Samples were stored in the dark and analysed within 12 h of sampling, thus no poisoning was required. C_T was determined using coulometric titration (Johnson et al., 1987) with a precision of $\leq 2 \mu\text{mol kg}^{-1}$. Measurements were calibrated against certified reference material (CRM, Dickson, 2010). Seawater pH_T was measured using the automated marine pH sensor (AMpS) system as described in Bellerby et al. (2002) modified for discrete mode. This system is an automated spectrophotometric pH sensor that makes dual measurements of thymol blue. The pH_T data used in this study were computed using the total hydrogen ion concentration scale and has a precision of 0.0002 pH_T and an estimated accuracy of better than 0.0025 pH_T units against CRM standards. The measured C_T and pH_T , with associated temperatures and salinity, were input to CO2SYS (Lewis and Wallace, 1998) to calculate saturation state of CaCO_3 using the dissociation constants for carbonic acid of Dickson and Millero (1987), boric acid from Dickson

Spring bloom phytoplankton composition

C. J. Daniels et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(1990b), sulphuric acid following Dickson (1990a) and the CO₂ solubility coefficients from Weiss (1974).

Satellite data on Chl *a*, photosynthetically available radiation (PAR) and sea surface temperature (SST) were obtained from the Aqua Moderate Resolution Image Spectroradiometer (MODIS) as 4 km resolution, 8 day composites. Data were extracted as averaged 3 pixel × 3 pixel grids, centred on the sampling locations. Day length was calculated according to Kirk (1994). The R/V *Meteor* was not fitted with a PAR sensor, thus satellite measurements were the only available source of PAR data.

2.6 Data availability

Data included in the paper are available from the data repository PANGAEA via Daniels and Poulton (2013) for the measurements of primary production, chlorophyll *a*, particulate inorganic carbon and particulate silicate, cell counts of coccolithophores, diatoms and microzooplankton; Esposito and Martin (2013) for measurements of nutrients; Paulsen et al. (2014) for measurements of picoplankton and nanoplankton; and Bellerby (2014) for measurements of the carbonate chemistry.

3 Results

3.1 General oceanography

The two sites were characterised by very different water column profiles throughout the study period. In the NWB, a pycnocline persisted over the upper 400 m with a variable mixed layer (20–100 m, Fig. 2d). In contrast, the ICB appeared well mixed over the upper 400 m when considered over the equivalent density range (Fig. 2a). However, weak unstable stratification was observed in the upper 100 m when examined over a much narrower range in density (Fig. 2a inset).

Sea surface temperature (SST) showed little variation at both sites (Table 1), while the ICB (8.6–8.9 °C) was consistently warmer than the NWB (6.5–7.2 °C). Satellite es-

BGD

12, 93–133, 2015

Spring bloom phytoplankton composition

C. J. Daniels et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



5 timates of SST were colder than in situ measurements and exhibited greater variability (Fig. 3a). However, the general pattern of the ICB being warmer than the NWB was observed from both in situ measurements and satellite derived ones. Sea-surface salinity (SSS), pH_T and Ω_{Ca} were relatively stable throughout the study with total ranges of

10 35.1–35.3, 8.0–8.1 and 3.0–3.2, respectively (Table 1). Initial surface water concentrations of nitrate (NO_3) and phosphate (PO_4) were $\sim 12 \text{ mmol N m}^{-3}$ and $\sim 0.7\text{--}0.8 \text{ mmol P m}^{-3}$ at both sites (Table 1). Silicic acid (dSi) was high throughout the study period (mostly $> 4 \text{ mmol Si m}^{-3}$), with slightly higher concentrations in the NWB ($5.3\text{--}5.7 \text{ mmol Si m}^{-3}$) than the ICB ($< 5 \text{ mmol Si m}^{-3}$). Draw-down of 1 mmol m^{-3} of NO_3 and dSi occurred in the ICB between the 19 and 27 April, but then returned to previous levels by 29 April. Nutrient drawdown did not occur in the NWB during the cruise period.

15 Both sites showed a similar trend of increasing daily PAR during the study (Fig. 3b); a twofold increase in the NWB (from 12.3 to $28.4 \text{ mol quanta m}^{-2} \text{ d}^{-1}$) and a slightly smaller increase in the ICB (from 13.5 to $24.3 \text{ mol quanta m}^{-2} \text{ d}^{-1}$). Daily irradiance continued to increase after the cruise finished, peaking around 40–45 days later at values in excess of $40 \text{ mol quanta m}^{-2} \text{ d}^{-1}$ (Fig. 3b). The general trend of increasing PAR was also reflected in the day length (Fig. 3b). At both sites, the euphotic depth shoaled as the study progressed, from 115 to 50 m in the ICB and from 80 to 56 m in the NWB (Table 2). However, the euphotic depth again deepened by 36 m between the 20 3rd and 4th visits to the ICB.

25 For the duration of the cruise until 27 April, surface and euphotic zone integrated particulate silicate (bSiO_2) increased in the ICB, peaking at $0.66 \text{ mmol Si m}^{-3}$ and $37.1 \text{ mmol Si m}^{-2}$, respectively (Fig. 5a, Table 2), with a significant decline in bSiO_2 after this date. Lower values of bSiO_2 , with little temporal variation, were found in the NWB, although a small increase in surface bSiO_2 was observed between the 14 and 22 April (from 0.05 to $0.08 \text{ mmol Si m}^{-3}$, Fig. 5a). Standing stocks of PIC were less variable than bSiO_2 . Highest surface values were observed during the last visit to the NWB

BGD

12, 93–133, 2015

Spring bloom phytoplankton composition

C. J. Daniels et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



($0.20 \text{ mmol C m}^{-3}$), while integrated calcite peaked at 11 mmol C m^{-2} on the 27 April in the ICB (Table 2).

3.2 Chlorophyll *a*

Profiles of CTD fluorescence in the NWB had a relatively consistent structure with high fluorescence in the stratified upper water column (Fig. 2e and f). Intra-site variation can be seen in the relative fluorescence values in surface waters, but a consistent increase over time was not observed. Fluorescence profiles in the ICB were more variable (Fig. 2b and c), ranging from profiles with high surface fluorescence (10 April) to profiles with elevated fluorescence throughout the upper 300 m.

Acetone extracted measurements of chlorophyll *a* (Chl *a*) ranged from 0.1 to 2.3 mg m^{-3} with highest values generally in surface waters (5–15 m). Surface Chl *a* was variable in the ICB, with the lowest surface values (0.27 – 0.31 mg m^{-3}) measured during the first visit (Table 2). Peak Chl *a* values in the ICB occurred on 10 April (2.2 mg m^{-3}), after which Chl *a* declined reaching a low of 0.62 mg m^{-3} by the end of the study (but remaining above initial Chl *a* values). Initial surface Chl *a* values were higher in the NWB (0.58 mg m^{-3}) than the ICB, and generally increased throughout the cruise. However, the magnitude of this increase was significantly smaller than in the ICB, peaking at only 0.93 mg m^{-3} . Euphotic zone integrated Chl *a* showed a similar pattern to surface Chl *a* across both stations, with highest values on 10 April (ICB, 146.4 mg m^{-2}).

Satellite estimates of Chl *a* also showed an increase in Chl *a* at both sites during the cruise (Fig. 3c and d), although these values ($< 0.4 \text{ mg m}^{-3}$) were much lower than measured in situ Chl *a* (Table 2). The large increase in Chl *a* associated with North Atlantic spring blooms occurred between 20 and 30 days after the cruise (Fig. 3c and d). Both sites were characterised by two peaks in Chl *a* throughout the year, one in late spring (mid-June) and another in late summer (mid-August). The largest satellite-derived Chl *a* values occurred in the ICB in late spring (1.7 mg m^{-3} , Fig. 3c), while in the NWB, peak Chl *a* occurred during the late summer bloom (1.6 mg m^{-3} , Fig. 3d).

Spring bloom phytoplankton composition

C. J. Daniels et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Size fractionated Chl *a* revealed very different communities at the two sites (Table 2 and Fig. 4). Initially in the ICB, approximately a quarter of the Chl *a* biomass was derived from the > 10 μm fraction (24–28 %; Table 2, Fig. 4a). On subsequent visits this increased significantly (to 56–94 %; Table 2, Fig. 4a). A general trend of an increasing contribution from the > 10 μm fraction was also observed in those samples collected for more detailed size fractionation (Fig. 4c). The detailed size fractionation showed that excluding the first ICB visit where samples were not collected, the > 10 μm fraction was completely dominated by the > 20 μm fraction in the ICB (Fig. 4c). Conversely, the > 10 μm fraction formed only a minor component (< 21 %) of the Chl *a* biomass in the NWB, although the > 10 μm contribution did increase throughout the cruise (Table 2, Fig. 4b). Detailed size fractionation in the NWB showed that the biggest increase in contribution came from the 2–10 μm fraction, increasing from 14 to 32 % (Fig. 4d), which was due to an increase in the absolute value of 2–10 μm Chl *a* (from 0.09 to 0.31 mg m^{-3}).

3.3 Primary production

Primary production (PP) in surface waters (5–15 m) ranged from 0.41 to 4.89 $\text{mmol C m}^{-3} \text{d}^{-1}$ in this study (Table 2), with PP generally decreasing with depth. Surface PP correlated well with euphotic zone integrated PP ($r = 0.98$, $p < 0.001$, $n = 7$). The largest change in PP occurred in the ICB, between the 26 March and the 10 April, when peak PP rates were observed in both the surface waters (4.89 $\text{mmol C m}^{-3} \text{d}^{-1}$) and integrated over the euphotic zone (221.9 $\text{mmol C m}^{-2} \text{d}^{-1}$, Table 2). Following this peak, PP in the ICB declined, although it generally remained higher than pre-peak PP rates. The > 10 μm PP fraction contributed between 35–61 % of the total PP in the ICB. In contrast, the range and maximum rate of PP in the NWB was much lower than the ICB (0.67–1.11 $\text{mmol C m}^{-3} \text{d}^{-1}$, Table 2) with the > 10 μm PP making up a much smaller fraction (< 20 %). However, a clear increase in the > 10 μm PP fraction was observed between 14 April (5 %) and 25 April (20 %). The general

BGD

12, 93–133, 2015

Spring bloom phytoplankton composition

C. J. Daniels et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



trend in total and size-fractionated PP at both sites reflected that observed in the Chl *a* measurements.

3.4 Community structure

3.4.1 Community structure – picoplankton and nanoplankton

5 Flow cytometry identified *Synechococcus*, autotrophic picoeukaryotes and autotrophic nanoplankton ($< 10 \mu\text{m}$) in relatively high abundance in all samples (Table 3). In general, *Synechococcus* and picoeukaryotes were more abundant in the NWB than the ICB. In the NWB, a contrasting pattern between *Synechococcus*, nanoplankton and picoeukaryotes was observed; while *Synechococcus* and the nanoplankton increased significantly from 2617 to 5483 and 484 to 1384 cells mL^{-1} respectively, a large decrease in picoeukaryotes was also observed, from 18 016 to 8456 cells mL^{-1} . A less coherent pattern was observed in the ICB, where peak concentrations of both *Synechococcus* (2112 cells mL^{-1}) and picoeukaryotes (6982 cells mL^{-1}) occurred on the 19 April, with a general decline after this date.

15 3.4.2 Community structure – coccolithophores

The coccolithophore species identified by polarised light microscopy were: *Emiliana huxleyi*, *Coccolithus pelagicus*, *Calcidiscus leptoporus*, *Coronosphaera mediterranea* and *Syracosphaera pulchra*. More detailed SEM observations found a number of other species at low cell densities not clearly identified by the light microscope: *Algirosphaera robusta*, *Acanthoica quattrosolina*, *Calciopappus caudatus*, *Gephyrocapsa muelleriae*, *Syracosphaera corolla*, *S. marginaporata*, *S. molischii*, *S. nodosa*, *S. ossa* and unidentified *Syracosphaera* spp. Many of these coccolithophore species have cell diameters between 10 and 20 μm , with the notable exceptions of *E. huxleyi*, *G. muelleriae* and the smaller *Syracosphaera* spp. (Young et al., 2003). Two morphotypes of *E. huxleyi* were observed in all samples (A and B) with morphotype A consistently dominant (71–100 %

BGD

12, 93–133, 2015

Spring bloom phytoplankton composition

C. J. Daniels et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



of total *E. huxleyi* numbers). The coccolithophore composition at both sites were similar, with *E. huxleyi* generally the most abundant species (4.4–28.1 cells mL⁻¹, Table 3) at both sites, while *Coccolithus pelagicus* was present in all samples at relatively low cell densities (0.15–2.79 cells mL⁻¹). The NWB was also characterised by the presence of *A. robusta* (2.7–12.7 cells mL⁻¹), while *S. marginaporata* (0–21.3 cells mL⁻¹) was only present in the ICB.

A general increase in coccolithophore abundance was observed in the ICB, with a large increase between 10 and 18 April (7.7–42.8 cells mL⁻¹). *Emiliana huxleyi* abundance decreased between 27 and 29 April (26.7–13.2 cells mL⁻¹), but *C. pelagicus* remained relatively constant (0.81–0.84 cells mL⁻¹). In the NWB, coccolithophores generally followed the trend of increasing Chl *a* with increases in abundance over time (Table 3). Within the coccolithophore communities, the largest relative increase in species abundance was by *C. pelagicus* with a sevenfold increase (0.38 to 2.66 cells mL⁻¹) between 14 and 22 April in the NWB.

3.4.3 Community structure – diatoms and microzooplankton

The diatom taxa identified by light microscopy were: *Chaetoceros*, *Cylindrotheca*, *Dactyliosolen*, *Guinardia striata*, *Leptocylindrus*, *Navicula*, *Pseudo-nitzschia*, *Rhizosolenia*, *Thalassionema*, and *Thalassiosira*. Whilst samples for diatom counts were collected only once per visit to each station, particulate silicate (bSiO₂) samples were collected from two CTD casts per visit. As the major source of bSiO₂, the significant variability observed in bSiO₂ between the station visits (Fig. 5a) suggested a temporal variability in the diatom cell abundance not captured in the Lugol's counts. Therefore, diatom abundance counts were supplemented using SEM image based diatom counts (Fig. 5b). However, due to the relatively smaller volumes examined by SEM (~ 4.2 mL vs. 50 mL), there is a greater inherent error in the counts and as such Lugol's counts were used wherever possible.

The diatom community was highly variable in the ICB (Fig. 5). Initially present only in very low abundances (1.3 cells mL⁻¹, Table 3), a peak concentration of 249 cells mL⁻¹

BGD

12, 93–133, 2015

Spring bloom phytoplankton composition

C. J. Daniels et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



was reached 15 days later on 10 April. The population then decreased over the rest of the study, down to 88 cells mL⁻¹, but remained above initial levels. A shift in composition was observed after the population peaked, from a *Chaetoceros* dominated community on 7 to 10 April (67–71 %) to one dominated by *Pseudo-nitzschia* (65–73 %, Fig. 5b) on the 27 to 29 April. Diatoms were virtually absent from light microscope measurements of the NWB, reaching a maximum of only 0.5 cells mL⁻¹ (Table 3).

The main microzooplankton groups present were planktonic ciliates and small (~ 5–10 µm) naked dinoflagellates (e.g. *Gyrodinium* and *Gymnodinium*). Microzooplankton concentrations were ~ 4 times higher in the NWB (10.8–17.6 cells mL⁻¹, Table 3) than in the ICB (2.5–4.7 cells mL⁻¹, Table 3). Diatoms initially dominated in the NWB (8.5 cells mL⁻¹), but were succeeded by ciliates (11.9–12.9 cells mL⁻¹). Both dinoflagellates and ciliates were present in similar concentrations in the ICB, except for the final station, when dinoflagellates dominated (4.2 cells mL⁻¹).

4 Discussion

4.1 Time series or mixing?

The dynamic nature of the ocean causes inherent difficulties in interpreting data collected from fixed-point, Eulerian time-series, such as those in this study. The distribution of phytoplankton in the ocean exhibits significant heterogeneity, which can be driven by mesoscale physical processes (Martin, 2005). Therefore, Eulerian time-series are vulnerable to advection such that instead of repeatedly sampling the same phytoplankton community, each sample is potentially from a different population, possibly with a different composition. Before examining the development of the phytoplankton community, it is therefore necessary to consider the physicochemical environment. Eddies and other mesoscale features would potentially cause significant variations in measured SST, SSS, nutrients and carbonate chemistry. With the possible exception of the nutrient concentrations, which will also be affected by the biology present, the

BGD

12, 93–133, 2015

Spring bloom phytoplankton composition

C. J. Daniels et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



measured physicochemical parameters were stable throughout the study period (Table 1). Therefore, although we cannot rule out the influence of mesoscale features and advection during the study, the relative consistency of the sampled physicochemical environment suggests that the community structure is representative of the location, rather than from multiple eddies, and thus we can examine how the community developed during the cruise and compare between two geographically separated sites.

4.2 Drivers of the phytoplankton bloom

Density profiles in the Iceland Basin (ICB) were seemingly indicative of a well-mixed water column (Fig. 2a), yet elevated fluorescence in the upper 100 m of the water column suggests that phytoplankton cells were not being evenly mixed throughout the water column (Fig. 2b). A detailed examination of the upper 100 m found small changes in the density profiles (Fig. 2a inset), corresponding to the elevated fluorescence, however the change in density with respect to depth was smaller ($\Delta\sigma_t < 0.025$ over 1 m) than most metrics used to identify mixed layers (e.g. Kara et al., 2000). Elevated fluorescence with only minimal stratification is consistent with the critical turbulence hypothesis (Huisman et al., 1999); here it is likely that active mixing had ceased, allowing phytoplankton net growth, while the response of the physical environment was slower than the biological response, and stratification was only just beginning to develop.

Although ICB upper water column fluorescence was elevated throughout the study, there was significant variation in the magnitude and structure of the fluorescence profiles (Fig. 3b and c), as well as a peak and decline in surface chlorophyll *a* (Chl *a*) and primary production (PP). The general theory of bloom formation is that once conditions are favourable for bloom formation, the pre-bloom winter ecosystem will transition into a blooming ecosystem, identifiable by increasing Chl *a* biomass and PP. However, we did not observe this smooth transition. Instead we observed periods of stability, characterised by increased stratification, Chl *a* and PP, followed by periods of instability where increased mixing weakened the developing stratification. Increased mixing detains phytoplankton out of the surface waters, reducing both Chl *a* biomass and

BGD

12, 93–133, 2015

Spring bloom phytoplankton composition

C. J. Daniels et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



physical environments, the two sites both represented early stages in the development of spring blooms.

Unlike the ICB, the factors limiting bloom formation in the NWB cannot easily be attributed to the physicochemical environment. Irradiance is a key driver of phytoplankton growth and bloom formation; the main spring bloom did not occur until daily PAR reached its seasonal maximum of $45 \text{ mol photons m}^{-2} \text{ d}^{-1}$ (Fig. 3b–d). The general increase in daily PAR over our study period was coupled with an increase in Chl *a* and PP in the NWB, suggesting that despite a stratified environment, irradiance was an important driving factor. Although the magnitude of the daily flux of PAR at both sites was similar, Chl *a* and PP were higher in the less stable ICB than the NWB, suggesting that irradiance was not the only driver of the NWB phytoplankton community. Irradiance levels can also have a secondary influence on the requirements for phytoplankton growth. While macronutrients were replete at both sites, we did not measure micronutrients such as iron (Fe). The cellular Fe demand increases in low light conditions (Moore et al., 2006), and as such Fe may be limiting at this early stage of bloom formation in the Norwegian Basin. However, without measurements of Fe (or phytoplankton photophysiology), we cannot directly test this hypothesis. Although temperature limits phytoplankton gross growth rates (Eppley, 1972), the relatively small difference in temperature between the NWB and the ICB ($\sim 1.5\text{--}2.5^\circ\text{C}$) is unlikely to have a significant impact on gross growth rates (Eppley, 1972).

Besides physicochemical drivers of bloom formation, the plankton community itself can play a large role in the development and formation of a bloom. Physiological parameters such as net growth rates (μ_{Chl}) and “assimilation efficiency” (i.e. PP normalised to biomass, in this case Chl *a*) can provide an insight into the state of the phytoplankton community. The NWB community had a noticeably lower assimilation efficiency ($13.5\text{--}15.8 \text{ g C [g Chl } a]^{-1} \text{ d}^{-1}$) than that in the ICB ($15.7\text{--}27.0 \text{ g C [g Chl } a]^{-1} \text{ d}^{-1}$), thus the relative increase in biomass in the NWB was slower, as reflected in the growth rates where the maximum estimated (net) growth rate in the NWB ($\mu_{\text{Chl}} = 0.05 \text{ d}^{-1}$) was much lower than in the ICB ($\mu_{\text{Chl}} = 0.22 \text{ d}^{-1}$). Assimilation efficiency varies with both

BGD

12, 93–133, 2015

Spring bloom phytoplankton composition

C. J. Daniels et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



environmental conditions and species composition, and therefore the composition of the phytoplankton community is likely to be another key driver behind the contrasting phytoplankton dynamics observed in the ICB and NWB.

4.3 Overall community composition

5 The contrasting structures of Chl *a* and PP size fractions observed at the two sites (Fig. 4, Table 2), were reflected in the contrasting composition of the phytoplankton communities (Table 3). In the ICB, a change in dominance in both Chl *a* and PP, from < 10 to the > 10 μm fraction, occurred as the diatom abundance increased between 26 March and 7 April. An increase in the abundance of the < 10 μm community was also
10 observed during this period, composed mainly of < 2 μm *Synechococcus* and pico-eukaryotes (Table 3, Fig. 4c). However, with most of the diatom population having cells > 20 μm (Fig. 4c), their relatively large size allowed the diatoms to dominate both the Chl *a* and PP while remaining numerically inferior. The decline in total Chl *a* and PP
15 later in our study was reflected by a decreasing abundance of most of the phytoplankton community (Table 3). However, the relative decrease of pico-phytoplankton (*Synechococcus* and picoeukaryotes) was greater than that of the diatoms, such that the > 10 μm fraction increased its dominance for both Chl *a* (94%) and PP (61%). Therefore, although surface Chl *a* and PP declined after the “mini-bloom event” which peaked around 10 April, the community structure did not return to a pre-bloom composition, but
20 instead remained dominated by diatoms.

Interestingly, the phytoplankton response to the increased diatom abundance was not uniform, with the nanoplankton abundance decreasing and *Synechococcus* increasing only after the peak in diatom abundance. Thus, we observed that the phytoplankton community response during the spring bloom was not universal across functional types as has been previously observed elsewhere (Brown et al., 2008).
25

In contrast to the ICB, a large shift in the NWB community was not observed. Pico-eukaryotes dominated both in terms of abundance (Table 3) and Chl *a*, through the < 2 μm fraction (Fig. 4d). This is consistent with previous observations of early stage

BGD

12, 93–133, 2015

Spring bloom phytoplankton composition

C. J. Daniels et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



spring blooms (Joint et al., 1993). Although the $< 2 \mu\text{m}$ Chl *a* fraction showed little variation throughout the study ($0.45\text{--}0.58 \text{ mg m}^{-3}$), variation in the $< 2 \mu\text{m}$ phytoplankton composition did occur, with an apparent succession from pico-eukaryotes to *Synechococcus* and nanoplankton. This may represent a community shift early in development of the spring bloom or may demonstrate the inherent variability within pre-bloom communities.

The increase in total Chl *a* in the NWB was driven primarily by the 2 to $10 \mu\text{m}$ fraction, which was likely composed of the nanoplankton, which itself had a threefold increase in population size (from 484 to $1384 \text{ cells mL}^{-1}$, Table 3). The phytoplankton responsible for the observed increase in the $> 10 \mu\text{m}$ Chl *a* and PP fraction cannot be confidently determined; large diatoms were absent and thus could not contribute. The microzooplankton population consisted of ciliates and dinoflagellates (*Gyrodinium* and *Gymnodinium*), both of which have been reported to be mixotrophic (Putt, 1990; Stoecker, 1999), and thus could potentially have contributed to the Chl *a* measurements. Furthermore, it is possible that part of the nanoplankton community, as measured by flow cytometry, was $> 10 \mu\text{m}$ and thus the increasing concentration of nanoplankton could also contribute to the increase in the $> 10 \mu\text{m}$ fraction.

4.4 Relative independence of the coccolithophore community

The traditional view on the seasonality of coccolithophores is that they succeed the diatom spring bloom, forming coccolithophore blooms in late summer. However, here we observed a typical North Atlantic community of coccolithophores (Savidge et al., 1995; Dale et al., 1999; Poulton et al., 2010), growing alongside the ICB diatom bloom, rather than just succeeding the diatoms. This is consistent with the “rising tide” hypothesis of Barber and Hiscock (2006), as well as observations from both in situ (Leblanc et al., 2009) and satellite measurements (Hopkins et al., 2014) suggesting that coccolithophores are present in North Atlantic spring blooms. Despite the contrasting environment and overall community structure of the NWB, the coccolithophore dynamics were similar, appearing independent of the overall community dynamics. Species-specific

BGD

12, 93–133, 2015

Spring bloom phytoplankton composition

C. J. Daniels et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



growth rates of coccolithophores (calculated from changes in cell concentration) found that *E. huxleyi* had the same net growth rate at both sites ($\mu = 0.06 \text{ d}^{-1}$), while the net growth rate of *C. pelagicus* was comparable to *E. huxleyi* in the ICB, but was slightly higher in the NWB ($\mu = 0.13 \text{ d}^{-1}$). Culture experiments of *E. huxleyi* and *C. pelagicus* have found comparable gross growth rates at temperatures below 10°C (Daniels et al., 2014), and out in situ observations support this conclusion. That *C. pelagicus* has higher net growth rates could also be indicative of higher grazing on the relatively smaller *E. huxleyi* (Daniels et al., 2014).

4.5 Contrasting patterns of diatoms

The diatom bloom in the ICB, which began between 26 March and 7 April, was initially dominated by *Chaetoceros* (71–67% of total cell numbers, Fig. 5b). As the community developed however, *Pseudo-nitzschia* succeeded as the dominant diatom genera (65–73% of total). Both *Chaetoceros* and *Pseudo-nitzschia* are common spring bloom diatoms (Sieracki et al., 1993; Rees et al., 1999; Brown et al., 2003), with *Chaetoceros* often dominant in the earlier stages of North Atlantic spring blooms (Sieracki et al., 1993; Rees et al., 1999). Resting spores of *Chaetoceros* have also been observed to dominate the export flux out of the Iceland Basin during the North Atlantic spring bloom in May 2008 (Ryner et al., 2013), suggesting dominance of the spring bloom prior to this period, consistent with the early community observed in our study.

Pseudo-nitzschia (previously identified as *Nitzschia* in other studies), tends to dominate later in the spring bloom (Sieracki et al., 1993; Moore et al., 2005), also consistent with this study. This suggests that as a genera, *Chaetoceros* are either able to adapt more quickly than *Pseudo-nitzschia*, or that they have a wider niche of growing conditions through a large diversity of species. However, once established, *Pseudo-nitzschia* are able to outcompete *Chaetoceros*, resulting in a community shift. That the succession of the diatom community observed in the ICB is consistent with that expected in the main diatom spring bloom, suggests that a mini-diatom bloom occurred prior to the formation of the main spring bloom.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Spring bloom phytoplankton composition

C. J. Daniels et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The observed variability in the relationship between diatoms (the main source of bSiO_2) and bSiO_2 was likely due to the species-specific variability in the cellular bSiO_2 content of diatoms (Baines et al., 2010). The abundance of *Pseudo-nitzschia*, rather than *Chaetoceros*, best explained the trend in bSiO_2 ($r = 0.92$, $p < 0.001$, $n = 8$), suggesting that while *Chaetoceros* has previously been observed as the major exporter of bSiO_2 (Rynewson et al., 2013), here *Pseudo-nitzschia* was the major producer.

In contrast to the ICB, diatoms appeared to be virtually absent ($< 0.5 \text{ cells mL}^{-1}$) in the NWB. While the $\text{dSi}:\text{NO}_3$ ratio was below the 1 : 1 requirement for diatoms, consistent with previous studies of North Atlantic blooms (Leblanc et al., 2009), dSi did not become depleted (always above 5 mmol Si m^{-3} , Table 1) and thus was not limiting. Furthermore, significant and increasing concentrations of particulate silicate (bSiO_2) were measured throughout the cruise (Fig. 5a). As the main source of bSiO_2 , diatoms would therefore be expected to be present. Although absent in the Lugol's counts, examination of SEM images found significant numbers ($101\text{--}600 \text{ cells mL}^{-1}$) of small ($< 5 \mu\text{m}$) diatoms (e.g. *Minidiscus* spp.) that were too small to be identified by light microscopy. However, they may still constitute an important component of the nanoplankton, as measured by flow cytometry. As a result of their small cell size, nanno-sized diatoms, such as *Minidiscus*, are easily missed when identifying and enumerating the phytoplankton community, and as such their potential biogeochemical importance may be greatly underestimated (Hinz et al., 2012). Other nanno-sized diatom species have been observed as major components of the phytoplankton community on the Patagonian Shelf (Poulton et al., 2013), in the Scotia Sea (Hinz et al., 2012), the northeast Atlantic (Savidge et al., 1995) and in the Norwegian Sea (Dale et al., 1999).

The *Minidiscus* spp. observed in this study exhibited a significant increase in population size during the study, from initial concentrations of $100\text{--}200 \text{ cells mL}^{-1}$, then up to $600 \text{ cells mL}^{-1}$ by the end of the study, and correlating well with both bSiO_2 ($r = 0.93$, $p < 0.01$, $n = 6$), and $\text{Chl } a$ ($r = 0.93$, $p < 0.01$, $n = 6$). Furthermore, the increasing concentration of *Minidiscus* corresponded to the increase in the 2 to $10 \mu\text{m}$ $\text{Chl } a$ size fraction (Fig. 4d). The maximum net growth rate of *Minidiscus*, estimated from changes

in cell abundances ($\mu = 0.13 \text{ d}^{-1}$), was significantly higher than that calculated for the total community using Chl *a* ($\mu_{\text{Chl}} = 0.05 \text{ d}^{-1}$). While different methods were used to determine these growth rates, it does suggest that conditions were favourable for the small nanno-sized diatoms to grow more rapidly than the bulk community.

The question therefore remains as to why the larger ($> 10 \mu\text{m}$) diatoms were virtually absent in an environment that is physically stable and nutrient replete, while small diatoms were able to thrive? The fate and ecology of overwintering oceanic diatoms is poorly understood. Many diatom species, both neritic and pelagic, are capable of forming resting stages that sink post bloom (Smetacek, 1985; Rynearson et al., 2013), yet diatoms must be present in spring when the diatom bloom begins. Therefore, either a diatom population is sustained in the upper water column over winter (Backhaus et al., 2003), or the spring diatom community is sourced from elsewhere (horizontally or vertically). In relatively shallow coastal environments, benthic resting stages overwinter until spring when they are remixed up into the water column, providing the seed population for the spring bloom (McQuoid and Godhe, 2004). It is unlikely that oceanic diatom blooms are seeded from the sediment, as the depths are far too great for remixing. However, viable diatom cells have been observed suspended at depth ($> 1000 \text{ m}$) in the ocean (Smetacek, 1985), and it is possible that these suspended deep populations are remixed to seed the spring bloom. An alternative hypothesis is based on the observation that diatom blooms generally occur first in coastal waters before progressing to the open ocean (Smetacek, 1985), suggesting that coastal diatom populations are horizontally advected into pelagic waters, thus seeding the spring bloom in the open ocean from shelf waters. The location of the source coastal populations, and their transit time to the open ocean location, would then affect the timing of the diatom blooms.

With such low concentrations of $> 10 \mu\text{m}$ diatoms ($< 0.5 \text{ cells mL}^{-1}$) in the NWB, it is possible that the overwintering diatom population was too small to seed the spring bloom. Furthermore, the potential grazing pressure from the significant microzooplankton population ($10.8\text{--}17.6 \text{ cells mL}^{-1}$) suggests that the observed diatom population was unlikely to develop into a diatom bloom. Instead an alternative seed popu-

Spring bloom phytoplankton composition

C. J. Daniels et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

at both sites, independent of the overall community, with similar concentrations of the main species *Emiliana huxleyi* and *Coccolithus pelagicus*.

In terms of the diatom community, *Chaetoceros* initially dominated the ICB diatom bloom, but was replaced by *Pseudo-nitzschia* as the bloom progressed, suggesting *Chaetoceros* as a key species in diatom bloom formation, while *Pseudo-nitzschia* was the major source of particulate silicate (bSiO₂). The lack of large (> 10 μm) bloom forming diatoms in the NWB, while small (< 5 μm) diatoms were present in high numbers (101–600 cells mL⁻¹), suggests that micro-zooplankton grazing, coupled with a potential lack of a seed population, was restricting diatom growth in the NWB, or that large diatoms are absent in NWB spring blooms.

These results suggest that despite both phytoplankton communities being in the early stage of bloom formation and exhibiting positive net growth rates, different physicochemical and biological factors control bloom formation with the resulting blooms likely to be significantly different in terms of biogeochemistry and trophic interactions throughout the growth season. Clearly, more in situ studies are needed in the transitional period between winter and the peak productivity of the spring bloom to examine compositional differences, growth and mortality factors, and how regional variability impacts on upper ocean biogeochemistry and deep-sea fluxes of organic material. Coupled studies of satellite derived products, including bloom phenology and phytoplankton physiology, and in situ processes are needed to examine the full spectrum of factors forming the spring bloom.

Acknowledgements. The “Deep Convection” cruise was funded by the Deutsche Forschungsgemeinschaft in a grant to M. St. John with financial support for this research from the EU Framework 7 EuroBASIN (EUROpean Basin-scale Analysis, Synthesis & Integration) project. C. J. Daniels had additional financial support from the UK Natural Environmental Research Council, via a Studentship, A. J. Poulton and A. P. Martin were also supported by NERC National Capability funding. We thank the officers and crew of the R/V *Meteor*, Theresa Reichelt for running and processing the CTD data, Mark Stinchcombe for bSiO₂ and nutrient measurements, Darryl Green and Matt Cooper for ICP-OES analyses, Stuart Painter, Keith Davidson and Sharon McNeill for ¹³C and POC analyses and Jason Hopkins for assistance with satellite

BGD

12, 93–133, 2015

Spring bloom phytoplankton composition

C. J. Daniels et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



data. Further thanks go to Stephanie Henson for her helpful comments on an early version of the paper. MODIS Aqua data were obtained from the NASA Ocean Color distributed archive (<http://oceancolor.gsfc.nasa.gov/>).

References

- 5 Backhaus, J. O., Hegseth, E. N., Wehde, H., Irigoien, X., Hatten, K., and Logemann, K.: Convection and primary production in winter, *Mar. Ecol.-Prog. Ser.*, 251, 1–14, doi:10.3354/meps251001, 2003.
- Baines, S. B., Twining, B. S., Brzezinski, M. A., Nelson, D. M., and Fisher, N. S.: Causes and biogeochemical implications of regional differences in silicification of marine diatoms, *Global Biogeochem. Cy.*, 24, GB4031, doi:10.1029/2010GB003856, 2010.
- 10 Barber, R. T. and Hiscock, M. R.: A rising tide lifts all phytoplankton: growth response of other phytoplankton taxa in diatom-dominated blooms, *Global Biogeochem. Cy.*, 20, GB4S03, doi:10.1029/2006gb002726, 2006.
- Behrenfeld, M. J.: Abandoning sverdrup's critical depth hypothesis on phytoplankton blooms, *Ecology*, 91, 977–989, doi:10.1890/09-1207.1, 2010.
- 15 Behrenfeld, M. J. and Boss, E. S.: Resurrecting the ecological underpinnings of ocean plankton blooms, *Annual Review of Marine Science*, 6, 167–194, doi:10.1146/annurev-marine-052913-021325, 2014.
- Bellerby, R. G. J.: Discrete carbonate chemistry measured during the M87/1 cruise in April 2012, PANGAEA, doi:<http://doi.pangaea.de/10.1594/PANGAEA.830294>, 2014.
- 20 Bellerby, R. G. J., Olsen, A., Johannessen, T., and Croot, P.: A high precision continuous spectrophotometric method for seawater pH measurements: the automated marine pH sensor (AMpS), *Talanta*, 56, 61–69, doi:10.1016/S0039-9140(01)00541-0, 2002.
- Brody, S. R. and Lozier, M. S.: Changes in dominant mixing length scales as a driver of subpolar phytoplankton bloom initiation in the North Atlantic, *Geophys. Res. Lett.*, 41, 3197–3203, doi:10.1002/2014GL059707, 2014.
- 25 Brown, L., Sanders, R., Savidge, G., and Lucas, C. H.: The uptake of silica during the spring bloom in the Northeast Atlantic Ocean, *Limnol. Oceanogr.*, 48, 1831–1845, doi:10.4319/lo.2003.48.5.1831, 2003.

Spring bloom phytoplankton composition

C. J. Daniels et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Spring bloom phytoplankton composition

C. J. Daniels et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Brown, S. L., Landry, M. R., Selph, K. E., Jin Yang, E., Rii, Y. M., and Bidigare, R. R.: Diatoms in the desert: plankton community response to a mesoscale eddy in the subtropical North Pacific, *Deep-Sea Res. Pt. II*, 55, 1321–1333, doi:10.1016/j.dsr2.2008.02.012, 2008.

Dale, T., Rey, F., and Heimdal, B. R.: Seasonal development of phytoplankton at a high latitude oceanic site, *Sarsia*, 84, 419–435, doi:10.1080/00364827.1999.10807347, 1999.

Daniels, C. J. and Poulton, A. J.: Chlorophyll *a*, primary production rates, carbon concentrations and cell counts of coccolithophores, diatoms and microzooplankton measured in water samples from the North Atlantic during the M87/1 cruise in April 2012, PANGAEA, doi:http://doi.pangaea.de/10.1594/PANGAEA.823595, 2013.

Daniels, C. J., Tyrrell, T., Poulton, A. J., and Pettit, L.: The influence of lithogenic material on particulate inorganic carbon measurements of coccolithophores in the Bay of Biscay, *Limnol. Oceanogr.*, 57, 145–153, doi:10.4319/lo.2012.57.1.0145, 2012.

Daniels, C. J., Sheward, R. M., and Poulton, A. J.: Biogeochemical implications of comparative growth rates of *Emiliania huxleyi* and *Coccolithus* species, *Biogeosciences*, 11, 6915–6925, doi:10.5194/bg-11-6915-2014, 2014.

Dickson, A. G.: Standard potential of the reaction: $\text{AgCl(s)} + 1/2\text{H}_2\text{(g)} = \text{Ag(s)} + \text{HCl(aq)}$, and the standard acidity constant of the ion HSO_4^- in synthetic seawater from 273.15 to 318.15 K, *J. Chem. Thermodyn.*, 22, 113–127, doi:10.1016/0021-9614(90)90074-Z, 1990a.

Dickson, A. G.: Thermodynamics of the dissociation of boric acid in synthetic sea water from 273.15 to 318.15 K, *Deep-Sea Res.*, 37, 755–766, doi:10.1016/0198-0149(90)90004-F, 1990b.

Dickson, A. G.: Standards for ocean measurements, *Oceanography*, 23, 34–47, doi:10.5670/oceanog.2010.22, 2010.

Dickson, A. G. and Millero, F. J.: A comparison of the equilibrium constants for the dissociation of carbonic acid in seawater media, *Deep-Sea Res.*, 34, 1733–1743, doi:10.1016/0198-0149(87)90021-5, 1987.

Eilertsen, H. C.: Spring blooms and stratification, *Nature*, 363, 24–24, doi:10.1038/363024a0, 1993.

Eppley, R. W.: Temperature and phytoplankton growth in the sea, *Fish. B.-NOAA*, 70, 1063–1085, 1972.

Esposito, M. and Martin, A. P.: Nutrient concentrations of water samples collected by CTD-rosettes during the cruise M87/1 in April 2012, PANGAEA, doi:http://doi.pangaea.de/10.1594/PANGAEA.823595, 2013.

Spring bloom phytoplankton composition

C. J. Daniels et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Fischer, A. D., Moberg, E. A., Alexander, H., Brownlee, E. F., Hunter-Cevera, K. R., Pitz, K. J., Rosengard, S. Z., and Sosik, H. M.: Sixty years of Sverdrup: a retrospective of progress in the study of phytoplankton blooms, *Oceanography*, 27, 222–235, doi:10.5670/oceanog.2014.26, 2014.

5 Frada, M., Young, J. R., Cachao, M., Lino, S., Martins, A., Narciso, A., Probert, I., and De Vargas, C.: A guide to extant coccolithophores (Calcihaptophycidae, Haptophyta) using light microscopy, *J. Nanoplankt. Res.*, 31, 58–112, 2010.

Giering, S. L. C., Sanders, R., Martin, A. P., Lindemann, C., Mayor, D. J., and St. John, M. A.: High export via small particles before the onset of the North Atlantic spring bloom, *Global Biogeochem. Cy.*, in review, 2014.

10 Henson, S. A., Dunne, J. P., and Sarmiento, J. L.: Decadal variability in North Atlantic phytoplankton blooms, *J. Geophys. Res.*, 114, C04013, doi:10.1029/2008JC005139, 2009.

Hinz, D. J., Poulton, A. J., Nielsdóttir, M. C., Steigenberger, S., Korb, R. E., Achterberg, E. P., and Bibby, T. S.: Comparative seasonal biogeography of mineralising nanoplankton in the scotia sea: *Emiliania huxleyi*, *Fragilariopsis* spp., and *Tetraparma pelagica*, *Deep-Sea Res. Pt. II*, 59–60, 57–66, doi:10.1016/j.dsr2.2011.09.002, 2012.

15 Hopkins, J., Henson, S. A., Painter, S. C., Tyrrell, T., and Poulton, A. J.: Global characteristics of *Emiliania huxleyi* blooms: insights into phytoplankton succession, *Global Biogeochem. Cy.*, in review, 2014.

20 Huisman, J., van Oostveen, P., and Weissing, F. J.: Critical depth and critical turbulence: two different mechanisms for the development of phytoplankton blooms, *Limnol. Oceanogr.*, 44, 1781–1787, doi:10.4319/lo.1999.44.7.1781, 1999.

Johnson, K. M., Sieburth, J. M., Williams, P. J., and Brandström, L.: Coulometric total carbon analysis for marine studies: automation and calibration, *Mar. Chem.*, 21, 117–133, doi:10.1016/0304-4203(93)90201-X, 1987.

25 Joint, I., Pomroy, A., Savidge, G., and Boyd, P.: Size-fractionated primary productivity in the northeast Atlantic in May–July 1989, *Deep-Sea Res. Pt. II*, 40, 423–440, doi:10.1016/0967-0645(93)90025-i, 1993.

Kara, A. B., Rochford, P. A., and Hurlburt, H. E.: An optimal definition for ocean mixed layer depth, *J. Geophys. Res.-Oceans*, 105, 16803–16821, doi:10.1029/2000JC900072, 2000.

30 Kirk, J. T. O.: *Light and Photosynthesis in Aquatic Ecosystems*, Cambridge University Press, Cambridge, 1994.

**Spring bloom
phytoplankton
composition**

C. J. Daniels et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Leblanc, K., Hare, C. E., Feng, Y., Berg, G. M., DiTullio, G. R., Neeley, A., Benner, I., Sprengel, C., Beck, A., Sanudo-Wilhelmy, S. A., Passow, U., Klinck, K., Rowe, J. M., Wilhelm, S. W., Brown, C. W., and Hutchins, D. A.: Distribution of calcifying and silicifying phytoplankton in relation to environmental and biogeochemical parameters during the late stages of the 2005 North East Atlantic Spring Bloom, *Biogeosciences*, 6, 2155–2179, doi:10.5194/bg-6-2155-2009, 2009.
- Legendre, L. and Gosseline, M.: Estimation of N and C uptake rates by phytoplankton using ^{15}N or ^{13}C : revisiting the usual computation formulae, *J. Plankton Res.*, 19, 263–271, doi:10.1093/plankt/19.2.263, 1996.
- Lewis, E. and Wallace, D. W. R.: Program Developed for CO_2 System Calculations, ORNL/CDIAC-105, Carbon Dioxide Information Analysis Center, ORNL, US Department of Energy, Oak Ridge, Tennessee, 1998.
- Lindemann, C. and St. John, M. A.: A seasonal diary of phytoplankton in the North Atlantic, *Frontiers in Marine Science*, 1, 37, doi:10.3389/fmars.2014.00037, 2014.
- Lochte, K., Ducklow, H. W., Fasham, M. J. R., and Stienen, C.: Plankton succession and carbon cycling at 47°N 20°W during the JGOFS North Atlantic Bloom Experiment, *Deep-Sea Res. Pt. II*, 40, 91–114, doi:10.1016/0967-0645(93)90008-b, 1993.
- Mahadevan, A., D'Asaro, E., Lee, C., and Perry, M. J.: Eddy-driven stratification initiates North Atlantic spring phytoplankton blooms, *Science*, 337, 54–58, doi:10.1126/science.1218740, 2012.
- Margalef, R.: Life-forms of phytoplankton as survival alternatives in an unstable environment, *Oceanol. Acta*, 1, 493–509, 1978.
- Martin, A.: The kaleidoscope ocean, *Philos. T. R. Soc. A*, 363, 2873–2890, doi:10.1098/rsta.2005.1663, 2005.
- McQuoid, M. R. and Godhe, A.: Recruitment of coastal planktonic diatoms from benthic versus pelagic cells: variations in bloom development and species composition, *Limnol. Oceanogr.*, 49, 1123–1133, doi:10.4319/lo.2004.49.4.1123, 2004.
- Miller, C. B.: *Biological Oceanography*, Wiley-Blackwell, Oxford, 2003.
- Moore, C. M., Lucas, M. I., Sanders, R., and Davidson, R.: Basin-scale variability of phytoplankton bio-optical characteristics in relation to bloom state and community structure in the Northeast Atlantic, *Deep-Sea Res. Pt. I*, 52, 401–419, doi:10.1016/j.dsr.2004.09.003, 2005.
- Moore, C. M., Mills, M. M., Milne, A., Langlois, R., Achterberg, E. P., Lochte, K., Geider, R. J., and La Roche, J.: Iron limits primary productivity during spring bloom development in the cen-

tral North Atlantic, *Glob. Change Biol.*, 12, 626–634, doi:10.1111/j.1365-2486.2006.01122.x, 2006.

Paulsen, M. L., Riisgaard, K., and Nielsen, T. G.: Abundance of pico- and nanophytoplankton during the Meteor cruise M87/1, PANGAEA, doi:http://doi.pangaea.de/10.1594/PANGAEA.839416, 2014

Poulton, A. J., Charalampopoulou, A., Young, J. R., Tarran, G. A., Lucas, M. I., and Quartly, G. D.: Coccolithophore dynamics in non-bloom conditions during late summer in the central Iceland Basin (July–August 2007), *Limnol. Oceanogr.*, 55, 1601–1613, doi:10.4319/lo.2010.55.4.1601, 2010.

Poulton, A. J., Painter, S. C., Young, J. R., Bates, N. R., Bowler, B., Drapeau, D., Lyczskowski, E., and Balch, W. M.: The 2008 *Emiliania huxleyi* bloom along the patagonian shelf: ecology, biogeochemistry, and cellular calcification, *Global Biogeochem. Cy.*, 27, 1023–1033, doi:10.1002/2013GB004641, 2013.

Putt, M.: Abundance, chlorophyll content and photosynthetic rates of ciliates in the Nordic Seas during summer, *Deep-Sea Res.*, 37, 1713–1731, doi:10.1016/0198-0149(90)90073-5, 1990.

Rees, A. P., Joint, I., and Donald, K. M.: Early spring bloom phytoplankton-nutrient dynamics at the Celtic Sea Shelf Edge, *Deep-Sea Res. Pt. I*, 46, 483–510, doi:10.1016/s0967-0637(98)00073-9, 1999.

Rynewson, T., Richardson, K., Lampitt, R., Sieracki, M., Poulton, A., Lyngsgaard, M., and Perry, M.: Major contribution of diatom resting spores to vertical flux in the sub-polar North Atlantic, *Deep-Sea Res. Pt. I*, 82, 60–71, doi:10.1016/j.dsr.2013.07.013, 2013.

Sanders, R., Morris, P. J., Stinchcombe, M., Seeyave, S., Venables, H., and Lucas, M.: New production and the *f* ratio around the Crozet Plateau in austral summer 2004–2005 diagnosed from seasonal changes in inorganic nutrient levels, *Deep-Sea Res. Pt. II*, 54, 2191–2207, doi:10.1016/j.dsr2.2007.06.007, 2007.

Sanders, R., Henson, S. A., Koski, M., De La Rocha, C. L., Painter, S. C., Poulton, A. J., Riley, J., Salihoglu, B., Visser, A., Yool, A., Bellerby, R., and Martin, A.: The biological carbon pump in the North Atlantic, *Prog. Oceanogr.*, 129, 200–218, doi:10.1016/j.pocean.2014.05.005, 2014.

Savidge, G., Boyd, P., Pomroy, A., Harbour, D., and Joint, I.: Phytoplankton production and biomass estimates in the northeast Atlantic Ocean, May–June 1990, *Deep-Sea Res. Pt. I*, 42, 599–617, doi:10.1016/0967-0637(95)00016-Y, 1995.

Spring bloom
phytoplankton
composition

C. J. Daniels et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Spring bloom phytoplankton composition

C. J. Daniels et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Siegel, D., Doney, S., and Yoder, J.: The North Atlantic spring phytoplankton bloom and Sverdrup's critical depth hypothesis, *Science*, 296, 730–733, doi:10.1126/science.1069174, 2002.

Sieracki, M. E., Verity, P. G., and Stoecker, D. K.: Plankton community response to sequential silicate and nitrate depletion during the 1989 North Atlantic spring bloom, *Deep-Sea Res. Pt. II*, 40, 213–225, doi:10.1016/0967-0645(93)90014-E, 1993.

Smetacek, V. S.: Role of sinking in diatom life-history cycles: ecological, evolutionary and geological significance, *Mar. Biol.*, 84, 239–251, doi:10.1007/BF00392493, 1985.

Stoecker, D. K.: Mixotrophy among dinoflagellates, *J. Eukaryot. Microbiol.*, 46, 397–401, doi:10.1111/j.1550-7408.1999.tb04619.x, 1999.

Sverdrup, H.: On conditions for the vernal blooming of phytoplankton, *Journal du Conseil International pour l'Exploration de la Mer*, 18, 287–295, 1953.

Taylor, J. R. and Ferrari, R.: Ocean fronts trigger high latitude phytoplankton blooms, *Geophys. Res. Lett.*, 38, L23601, doi:10.1029/2011GL049312, 2011a.

Taylor, J. R. and Ferrari, R.: Shutdown of turbulent convection as a new criterion for the onset of spring phytoplankton blooms, *Limnol. Oceanogr.*, 56, 2293, doi:10.4319/lo.2011.56.6.2293, 2011b.

Townsend, D. W., Cammen, L. M., Holligan, P. M., Campbell, D. E., and Pettigrew, N. R.: Causes and consequences of variability in the timing of spring phytoplankton blooms, *Deep-Sea Res. Pt. I*, 41, 747–765, doi:10.1016/0967-0637(94)90075-2, 1994.

Townsend, D. W., Keller, M. D., Sieracki, M. E., and Ackleson, S. G.: Spring phytoplankton blooms in the absence of vertical water column stratification, *Nature*, 360, 59–62, doi:10.1038/360059a0, 1992.

Weiss, R. F.: Carbon dioxide in water and seawater: the solubility of a non-ideal gas, *Mar. Chem.*, 2, 203–215, doi:10.1016/0304-4203(74)90015-2, 1974.

Young, J. R., Geisen, M., Cros, L., Kleijne, A., Sprengel, C., Probert, I., and Ostergaard, J.: A guide to extant coccolithophore taxonomy, *Journal of Nannoplankton Research, Special Issue 1*, 1–132, 2003.

Zubkov, M. V., Burkill, P. H., and Topping, J. N.: Flow cytometric enumeration of DNA-stained oceanic planktonic protists, *J. Plankton Res.*, 29, 79–86, doi:10.1093/plankt/fbl059, 2007.

Spring bloom phytoplankton composition

C. J. Daniels et al.

Table 1. Physicochemical features of the Iceland Basin and Norwegian Basin stations: SST, sea surface temperature; SSS, sea surface salinity; C_T , dissolved inorganic carbon; Ω_C , calcite saturation state; NO_3 , nitrate; PO_4 , phosphate; dSi, silicic acid.

Location	Sta.	Date	Day of Year	SST ($^{\circ}C$)	SSS	Carbonate Chemistry			Surface Macro-nutrients ($mmolm^{-3}$)		
						C_T ($\mu molm^{-3}$)	pH_T	Ω_C	NO_3	PO_4	dSi
Iceland Basin	1	25 Mar	85	8.7	35.3	2149	8.0	3.1	12.3	0.79	4.7
	1	26 Mar	86	8.7	35.3	2148	8.0	3.1	12.6	0.81	4.7
	2	7 Apr	98	8.7	35.3	2140	8.0	3.1	12.4	0.81	4.5
	2	10 Apr	101	8.7	35.3	2139	8.1	3.2	11.5	0.75	4.3
	3	18 Apr	109	8.8	35.3	2144	8.1	3.2	11.6	0.79	4.3
	3	19 Apr	110	8.7	35.3	2150	8.1	3.2	11.9	0.76	4.1
	4	27 Apr	118	8.9	35.3	2135	8.1	3.2	10.7	0.70	3.1
	4	29 Apr	120	8.6	35.3	2148	–	–	12.0	0.80	4.2
Norwegian Basin	1	30 Mar	90	7.0	35.2	2142	8.1	3.0	12.1	0.67	5.3
	1	31 Mar	91	7.1	35.2	2161	8.1	3.0	12.5	0.81	5.4
	2	12 Apr	103	7.2	35.2	2153	8.1	3.0	13.4	0.84	5.6
	2	14 Apr	105	6.9	35.2	2152	8.1	3.0	13.5	0.82	5.6
	3	22 Apr	113	6.5	35.1	2150	8.1	3.0	12.2	0.79	5.7
	3	25 Apr	116	6.8	35.2	2143	8.1	3.0	12.5	0.82	5.7

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Spring bloom
phytoplankton
composition

C. J. Daniels et al.

Table 2. Biological features of the Iceland Basin and Norwegian Basin stations: Chl *a*, chlorophyll *a*; PP, primary production; bSiO₂, particulate silicate; PIC, particulate inorganic carbon.

Location	Sta.	Date	Surface Chl <i>a</i> (mg m ⁻³)	Surface PP (mmol C m ⁻³ d ⁻¹)	Surface size fractions		Euphotic zone depth (m)	Euphotic zone integrals			
					> 10 μm Chl <i>a</i> (%)	> 10 μm PP (%)		Chl <i>a</i> (mg m ⁻²)	bSiO ₂ (mmol Si m ⁻²)	PIC (mmol C m ⁻²)	PP (mmol C m ⁻² d ⁻¹)
Iceland Basin	1	25 Mar	0.27		28		115	22.3	8.3	7.7	
	1	26 Mar	0.31	0.41	24	35	115	26.5	2.5	4.5	22.2
	2	7 Apr	1.13		80		72	61.4	8.7	8.7	
	2	10 Apr	2.18	4.89	84	61	72	146.4	19.6	6.9	221.9
	3	18 Apr	1.01		56		50	49.2	13.4	6.5	
	3	19 Apr	1.15	2.11	67	40	50	55.6	15.4	5.8	58.0
	4	27 Apr	1.18		–		86	75.7	37.1	11.0	
	4	29 Apr	0.62	1.19	94	61	86	55.3	27.6	8.1	61.5
Norwegian Basin	1	30 Mar	0.58		6		80	34.6	5.5	7.7	
	1	31 Mar	0.59	0.67	7	5	80	39.2	7.0	7.1	27.3
	2	12 Apr	0.54		9		65	32.3	4.4	5.9	
	2	14 Apr	0.69	0.90	13	5	65	37.2	4.4	6.4	38.2
	3	22 Apr	0.93		10		56	46.7	5.0	9.7	
	3	25 Apr	0.84	1.11	21	20	56	40.5	6.4	10.5	39.8

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Spring bloom phytoplankton composition

C. J. Daniels et al.

Table 3. Phytoplankton abundance at the Iceland Basin and Norwegian Basin stations, measured by flow cytometry (*Synechococcus*, pico-eukaryotes and nanoplankton), inverted microscopy (diatoms and microzooplankton) and polarizing light microscopy (coccolithophores).

Location	Sta.	Date	Phytoplankton abundance (cells mL ⁻¹)								
			<i>Synechococcus</i>	Pico-eukaryotes	Nanoplankton (< 10 μm)	Diatoms (< 10 μm)	Micro- zooplankton	Coccolithophores			
								<i>E. huxleyi</i>	<i>C. pelagicus</i>	<i>A. robusta</i>	Others
Iceland Basin	1	25 Mar	–	–	–	–	–	7.5	0.15		1.2
	1	26 Mar	675	2347	1116	1.3	2.5	4.4	0.22		0.5
	2	7 Apr	400	3375	215	–	–	5.2	0.19		4.1
	2	10 Apr	480	6715	813	249.2	4.0	6.8	0.15		0.7
	3	18 Apr	–	–	–	–	–	16.9	0.22		25.6
	3	19 Apr	2112	6962	712	151.3	2.8	21.9	0.69		22.3
	4	27 Apr	1299	1486	298	–	–	26.7	0.81		7.9
	4	29 Apr	782	1215	313	87.8	4.7	13.2	0.84		7.5
Norwegian Basin	1	30 Mar	–	–	–	–	–	6.1	0.09	4.8	2.9
	1	31 Mar	2617	18 016	484	0.2	10.8	7.2	0.28	3.8	1.0
	2	12 Apr	–	–	–	–	–	11.8	0.41	2.7	0.3
	2	14 Apr	3372	10 433	858	0.1	17.6	16.0	0.38	3.7	5.1
	3	22 Apr	–	–	–	–	–	27.9	2.66	12.7	11.7
3	25 Apr	5483	8456	1384	0.5	14.0	28.1	2.79	7.8	8.6	

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)

[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)





Figure 2. Upper water column profiles for the ICB (a–c) and the NWB (d–f), of (a, d) density, (b, e) CTD fluorescence and (c, f) CTD fluorescence normalised to peak CTD fluorescence for each profile.

Spring bloom
phytoplankton
composition

C. J. Daniels et al.




Figure 3. Seasonal variation in (a) satellite sea surface temperature (SST), (b) satellite daily incidental PAR and day length and (c, d) satellite chlorophyll *a* (Chl *a*) for (c) the Iceland Basin (ICB) and (d) the Norwegian Basin (NWB) for 2012. The grey region indicates the period of the cruise. The vertical dotted lines in plots (c) and (d) indicate bloom initiation, calculated following Henson et al. (2009). The insets in (c) and (d) show the variation in satellite chlorophyll during the period of the cruise.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Spring bloom
phytoplankton
composition

C. J. Daniels et al.




Figure 4. Size fractionated chlorophyll *a* (Chl *a*) for (a, c) the Iceland Basin, and (b, d) the Norwegian Basin. Plots (a) and (b) show the <math><10</math> and >math>>10\mu\text{m}</math> fractions, (c) and (d) show the <math><2</math>, $2-10$, $10-20$, and >math>>20\mu\text{m}</math> fractions.

Spring bloom
phytoplankton
composition

C. J. Daniels et al.




Figure 5. (a) Particulate silicate (bSiO₂) and (b) diatom species abundance in the Iceland Basin. Black symbols indicate where diatoms were counted from Lugol's samples, while open symbols indicate SEM counts.