

$\delta^{11}\text{B}$, $^{87}\text{Sr}/^{86}\text{Sr}$ and
 $\delta^{88/86}\text{Sr}$ of cultured
Arctica islandica

Y.-W. Liu et al.

Environmental controls on the boron and strontium isotopic composition of aragonite shell material of cultured *Arctica islandica*

Y.-W. Liu¹, S. M. Aciego¹, and A. D. Wanamaker Jr.²

¹Department of Earth and Environmental Sciences, University of Michigan, 1100 N. University Avenue, Ann Arbor, MI 48109-1005, USA

²Department of Geological and Atmosphere Sciences, Iowa State University, Room 12, Science I, Ames, IA 50011-3212, USA

Received: 3 January 2015 – Accepted: 9 January 2015 – Published: 10 February 2015

Correspondence to: Y.-W. Liu (liuyiwei@umich.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Ocean acidification, the decrease in ocean pH associated with increasing atmospheric CO₂, is likely to impact marine organisms, particularly those that produce carbonate skeletons or shells. Therefore it is important to investigate how environmental factors (seawater pH, temperature and salinity) influence the chemical compositions in biogenic carbonates. In this study we report the first high-resolution strontium (⁸⁷Sr/⁸⁶Sr and δ^{88/86}Sr) and boron (δ¹¹B) isotopic values in the aragonite shell of cultured *Arctica islandica* (*A. islandica*). The ⁸⁷Sr/⁸⁶Sr ratios from both tank water and shell samples show ratios nearly identical to the open ocean, which suggests that the shell material reflects ambient ocean chemistry without terrestrial influence. The ⁸⁴Sr–⁸⁷Sr double spike resolved shell δ^{88/86}Sr and Sr concentration data show no resolvable change throughout the culture period and reflect no theoretical kinetic mass fractionation throughout the experiment despite a temperature change of more than 15°C. The δ¹¹B records from the experiment show at least a 5‰ increase through the culture season (January 2010–August 2010), with low values from beginning to week 19 and higher values hereafter. The larger range in δ¹¹B in this experiment compared to predictions based on other carbonate organisms (2–3‰) suggests that a species-specific fractionation factor may be required. A relatively strong correlation between the ΔpH (pH_{shell} – pH_{sw}) and seawater pH (pH_{sw}) was observed ($R^2 = 0.34$), which suggests that *A. islandica* partly regulates the pH of the extrapallial fluid. However, this proposed mechanism only explains approximately 34% of the variance in the δ¹¹B data. Instead, a rapid rise in δ¹¹B after week 19 suggests that the boron uptake of the shell changes when a temperature threshold of 13°C is reached.

1 Introduction

Since the onset of the Industrial Revolution (ca. AD 1750) the global ocean has absorbed approximately 30% of the emitted anthropogenic carbon dioxide (CO₂)

BGD

12, 2979–3023, 2015

δ¹¹B, ⁸⁷Sr/⁸⁶Sr and
δ^{88/86}Sr of cultured
Arctica islandica

Y.-W. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



causing ocean acidification (IPCC, 2013). The ecological effects from lowering the pH of the surface ocean on marine organisms, especially those which calcify, will likely be substantial (Doney et al., 2009; Hofmann et al., 2010; Orr et al., 2005), highly variable among taxa (Kroeker et al., 2010; Riebesell et al., 2013), and extend throughout the global ocean (Feely et al., 2004; Orr et al., 2005). Furthermore, anthropogenic CO₂ is not evenly distributed among the ocean basins. In particular, the North Atlantic Ocean stores about 23% of anthropogenic CO₂ despite representing only 15% of the global ocean area (Sabine et al., 2004). Consequently, the North Atlantic Basin, compared to other regions, might be more susceptible to ocean acidification. In situ measurements of surface ocean pH are rare prior to about 1990 (Byrne et al., 2010; Dore et al., 2009), however, instrumental data show that the surface global ocean pH has decreased by approximately 0.05 pH units in the last 20–25 years (IPCC, 2013). Time-series data from the North Pacific and North Atlantic Oceans indicate that the surface ocean pH has been changing between –0.0015 and –0.0024 pH units per year (IPCC, 2013; Vázquez-Rodríguez et al., 2012).

~~Given the recent changes in surface ocean pH and projected persistence of this global problem,~~ proxy records from the geologic record sensitive to oceanic carbon dynamics are highly desired to place modern pH trends into context (e.g., Hönisch et al., 2012.). Biogenic proxy archives calcifying within the surface waters of the global oceans have the unique potential to reveal spatial and temporal patterns and trends in pH using boron isotopes (e.g., Anagnostou et al., 2012; Shinjo et al., 2013). However, in the dynamic coastal regions of the global ocean, local and regional processes have the potential to complicate the boron-pH proxy (described in detail below). Freshwater mixing has the potential to change (1) temperature, salinity, and pH, (2) nutrient availability and productivity leading to changes in pH, and (3) local seawater boron isotopic composition. Additionally, potential species-specific biological effects that occur during calcification need to be evaluated. Here we apply multiple isotope systems to evaluate the boron-pH proxy in the northern North Atlantic coastal and shelf-dwelling marine bivalve *Arctica islandica* (*A. islandica*) exposed to ambient

BGD

12, 2979–3023, 2015

$\delta^{11}\text{B}$, $^{87}\text{Sr}/^{86}\text{Sr}$ and $\delta^{88/86}\text{Sr}$ of cultured *Arctica islandica*

Y.-W. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




conditions. We use radiogenic strontium isotopes to assess terrestrial river water influence on seawater and shell geochemistry for elements with long residence times in coastal water, such as boron. We utilize stable strontium isotopes from shell material to evaluate the potential impacts of growth rates during biomineralization.

1.1 Boron isotopes as pH indicators in biogenic carbonates

Boron has two natural stable isotopes, ^{10}B and ^{11}B , which comprise 19.82 and 80.18% of total boron, respectively (Berglund and Wieser, 2011). The dominant aqueous species of boron in seawater are $\text{B}(\text{OH})_3$ and $\text{B}(\text{OH})_4^-$. The relative proportion of these two species in an aqueous environment is a function of pH with the following relationship:



At low pH, boron exists as $\text{B}(\text{OH})_3$ in solution, conversely at high pH, boron exists as $\text{B}(\text{OH})_4^-$. The governing reaction for isotope exchange between these two species is:



The stable isotope ^{11}B is enriched in $\text{B}(\text{OH})_3$ compared to $\text{B}(\text{OH})_4^-$, and the combination of Reactions (R1) and (R2) can be used to determine the distribution of the two boron species and the isotopic composition of each for a given pH. The isotopic composition of boron is generally reported as:

$$\delta^{11}\text{B} = \left[\frac{(^{11}\text{B}/^{10}\text{B})_{\text{sample}}}{(^{11}\text{B}/^{10}\text{B})_{\text{SRM 951a}}} - 1 \right] \cdot 1000 (\text{‰}) \quad (1)$$

where SRM 951 is the internationally recognized boron isotope standard. Because the residence time of seawater boron is approximately 14 million years (Lemarchand et al., 2000), boron isotopic composition in the open ocean is considered constant over this

BGD

12, 2979–3023, 2015

$\delta^{11}\text{B}$, $^{87}\text{Sr}/^{86}\text{Sr}$ and
 $\delta^{88/86}\text{Sr}$ of cultured
Arctica islandica

Y.-W. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



time period, with an average seawater $\delta^{11}\text{B}$ value of 39.61‰ (Foster et al., 2010). Therefore, $\delta^{11}\text{B}$ has the following relationship: at low pH, the isotopic composition of $\text{B}(\text{OH})_3$ is equal to the isotopic composition of the total dissolved boron (39.61‰). In contrast, at high pH, the isotopic composition of $\text{B}(\text{OH})_4^-$ is equal to the isotopic composition of the total dissolved boron. Therefore the $\delta^{11}\text{B}$ is enriched in $\text{B}(\text{OH})_3$ by about 20‰ with respect to $\text{B}(\text{OH})_4^-$ at any equilibrium pH based on a constant fractionation factor.

During growth, marine carbonates primarily incorporate $\text{B}(\text{OH})_4^-$ into the carbonate structure. Building on these relationships, Hemming and Hanson (1992) demonstrated that seawater pH dictates the amount of $\text{B}(\text{OH})_4^-$ in seawater and thus the isotopic composition of boron in marine carbonates. Changes in the $\delta^{11}\text{B}$ composition of shell carbonates are based on the equation:

$$\text{pH} = \text{pK}_b - \log \left(\frac{\delta^{11}\text{B}_{\text{sw}} - \delta^{11}\text{B}_{\text{carbonate}}}{\alpha \delta^{11}\text{B}_{\text{carbonate}} - \delta^{11}\text{B}_{\text{sw}} + 1000(\alpha - 1)} \right) \quad (2)$$

where pK_b is the pK value for boric acid at a given temperature and salinity, and is 8.5975 at 25 °C and 35 PSU salinity (DOE, 1994), $\delta^{11}\text{B}_{\text{sw}}$ is the isotopic composition of seawater, and α is the equilibrium isotopic fractionation factor between boric acid and borate ion, (e.g., Reaction R2). Of these variables, only the seawater composition can be considered known and constant for all geographic locations and carbonate-precipitating species. Temperature, salinity and the fractionation factor must be estimated. Two empirical and analytical values of α are suggested for seawater: (1) $\alpha = 1.0194$, a theoretical result of Kakihana et al. (1977), which has been applied widely on paleo-reconstructions (Hönisch et al., 2004; Sanyal et al., 1995; Kakihana et al., 1977); and (2) $\alpha = 1.0272$, which was empirically obtained from Klochko et al. (2006) and is considered to better describe the distribution of the two boron species in nature today (Foster, 2008; Pagani et al., 2005; Klochko et al., 2006). However, due to the ability of calcifying organisms to buffer their own

 $\delta^{11}\text{B}$, $^{87}\text{Sr}/^{86}\text{Sr}$ and $\delta^{88/86}\text{Sr}$ of cultured *Arctica islandica*

Y.-W. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



$\delta^{11}\text{B}$, $^{87}\text{Sr}/^{86}\text{Sr}$ and $\delta^{88/86}\text{Sr}$ of cultured *Arctica islandica*

Y.-W. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



local environments, species specific fractionation factors and transfer functions are likely more appropriate than theoretical α values (Hönisch et al., 2004; Trotter et al., 2011; Rae et al., 2011; Anagnostou et al., 2012; Krief et al., 2010; Reynaud et al., 2004, 2008). Thus far, the $\text{pH}-\delta^{11}\text{B}$ relationship has been tested extensively on some biogenic marine carbonates (foraminifera and coral) with broad success (Anagnostou et al., 2012; Hönisch et al., 2004; Rae et al., 2011; Reynaud et al., 2004; Ni et al., 2007; Henehan et al., 2013), but few measurements have been made on aragonitic bivalves.

As indicated earlier, one of the primary assumptions in applying the boron isotope/pH proxy technique is that the boron isotopic composition of the seawater from which the biogenic carbonates form remains constant. Therefore, in order to use such a proxy to understand pH changes through the geological past, a technique is required to evaluate variability in local seawater geochemistry.

1.2 Radiogenic strontium isotopes as a water mass tracer

To evaluate the potential impacts of freshwater on local or regional ocean chemistry, it is necessary to differentiate sources that influence the chemical composition of the target water mass. This is especially important in coastal regions where fresh water input from river, surface runoff and/or groundwater mix with seawater often modifying both physical (e.g. temperature, salinity or turbidity) and chemical (e.g. nutrients, trace metal and/or isotopic compositions) conditions. Even for isotopic systems with long residence times in seawater (including boron), observations indicate that large fluxes of freshwater can cause substantial variations in coastal environments where conservative mixing behavior is generally assumed but not always achieved (D'Olivo et al., 2014; Chung et al., 2009; Huang and You, 2007; Huang et al., 2011; Widerlund and Andersson, 2006).

Radiogenic strontium has successfully been used to trace unique water masses. There are four naturally occurring isotopes of strontium: ^{84}Sr , ^{86}Sr , ^{87}Sr and ^{88}Sr , with the only radiogenic isotope being ^{87}Sr , which decays from ^{87}Rb . Therefore, the $^{87}\text{Sr}/^{86}\text{Sr}$ ratio is widely used to trace provenance in geological studies (Jahn et al.,

$\delta^{11}\text{B}$, $^{87}\text{Sr}/^{86}\text{Sr}$ and $\delta^{88/86}\text{Sr}$ of cultured *Arctica islandica*

Y.-W. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2001; Bataille and Bowen, 2012; Huang and You, 2007; Aarons et al., 2013). Because the residence time of Sr in seawater is more than 4 My, similar to the residence time of boron (Goldberg, 1963; Broecker, 1963), $^{87}\text{Sr}/^{86}\text{Sr}$ is considered spatially homogeneous in seawater at any instant of geological time. However, in coastal areas, radiogenic Sr isotopes vary as inputs from continental sources are released from terrestrial sediments to fresh water and then exported to the open ocean (Huang et al., 2011). The variability in $^{87}\text{Sr}/^{86}\text{Sr}$ ratios in natural archives on seasonal and annual timescales has been used as a proxy for quantitatively evaluating proportions of different water mass sources in coastal regions (Huang et al., 2011). The similarity of residence times of Sr and B in the ocean, and potential large differences between ocean and terrestrial isotopic compositions, suggests that the radiogenic strontium composition of shell material can be used to determine if there is a potential offset between local seawater and open-ocean B isotopic composition.

Prior studies have shown that bivalve shells record both ambient seawater composition and mixing of water sources in the ambient seawater. Widerlund and Anderson (2006) developed a two-endmember mixing model of terrestrial fluvial water and seawater radiogenic Sr in the Baltic Sea and compared the modern bivalve $^{87}\text{Sr}/^{86}\text{Sr}$ inferred salinity across the Baltic Sea from inland to the Atlantic coast to the in situ measurements of water salinity. Their results indicate conservative characteristics of $^{87}\text{Sr}/^{86}\text{Sr}$ vs. salinity in the Baltic Sea. Maurer et al. (2012) also measured $^{87}\text{Sr}/^{86}\text{Sr}$ ratio in a freshwater bivalve species: the Sr isotopic ratios recorded in the shells, in both early and late ontogeny ages, agreed with local water samples, indicating their usefulness in investigating the effects of anthropogenic contamination in rivers.

1.3 Stable strontium isotopes in biogenic carbonates

In addition to $^{87}\text{Sr}/^{86}\text{Sr}$, the stable isotopes of Sr ($\delta^{88/86}\text{Sr}$, the deviation in $^{88}\text{Sr}/^{86}\text{Sr}$ of a sample relative to a standard given in parts per mil) in natural archives are potential indicators of environmental controls on vital effects, although the applicability

is less well established and still controversial. Previously reported $\delta^{88/86}\text{Sr}$ values of coral, foraminifera and coccolithophores from the literature show ^{88}Sr depleted results compared to the $\delta^{88/86}\text{Sr}$ value in seawater, which suggest the lighter ^{86}Sr is preferentially incorporated into biogenic carbonates (Böhm et al., 2012; Fietzke and Eisenhauer, 2006; Krabbenhöft et al., 2010; Rüggeberg et al., 2008; Stevenson et al., 2014). If the fractionation of stable Sr isotopes in precipitated biogenic carbonates is dominated by kinetic isotope effects and not under equilibrium conditions, the $\delta^{88/86}\text{Sr}$ ratio would likely have a strong correlation with precipitation temperature and/or precipitation rate. As an environmental proxy, $^{88}\text{Sr}/^{86}\text{Sr}$ in both deep sea and tropical corals has been reported as a temperature proxy (Fietzke and Eisenhauer, 2006; Rüggeberg et al., 2008). However, more recent, higher precision work has indicated either a null relationship (foraminifera) or negative relationship (coccolithophores) between $\delta^{88/86}\text{Sr}$ and ambient seawater temperature, suggesting growth rate controls the uptake of Sr isotopes into biogenic carbonates (Böhm et al., 2012; Stevenson et al., 2014). Although stable strontium applications are in their infancy, the potential for vital effects (growth rate) fractionating $\delta^{88/86}\text{Sr}$ indicates it could be useful for assessing vital effects during calcification in bivalve archives.

1.4 The biogenic archive *Arctica islandica*

The long-lived bivalve mollusk *A. islandica*, common in the shelf seas of the temperate to sub-polar North Atlantic Ocean, is an excellent high-resolution marine archive with a huge potential for monitoring pH as well as other seawater properties (for a recent review, see Schöne, 2013). This stationary benthic clam lives in water depths ranging from ~ 10 m to as deep as 500 m and thrives in full marine conditions, yet can also tolerate salinities as low as 28 PSU for short time intervals (Merrill and Ropes, 1968; Nicol, 1951). *Arctica islandica* lives within the sediment and extends its relatively short siphons into the main water column, exchanging water to feed and remove waste. Weidman (1995) demonstrated that the geochemical signature in the shell material

BGD

12, 2979–3023, 2015

$\delta^{11}\text{B}$, $^{87}\text{Sr}/^{86}\text{Sr}$ and $\delta^{88/86}\text{Sr}$ of cultured *Arctica islandica*

Y.-W. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



$\delta^{11}\text{B}$, $^{87}\text{Sr}/^{86}\text{Sr}$ and $\delta^{88/86}\text{Sr}$ of cultured *Arctica islandica*

Y.-W. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



reflects that of the ambient water conditions and not pore water. *Arctica islandica* is highly suitable for environmental and ocean studies because: (1) *A. islandica* is extremely long-lived – up to 5 centuries (Schöne et al., 2005; Wanamaker Jr. et al., 2008a), (2) it produces annual growth increments in its shell (Jones, 1980), (3) regional increment series can be crossdated, demonstrating a common response to environmental forcing(s) (Schöne et al., 2003), (4) fossil shells can be crossdated and floating shell chronologies can be constructed after radiocarbon dating (Scourse et al., 2006), (5) live-caught shells can be crossdated with fossil shells to assemble very long, absolutely dated growth records (Butler et al., 2009, 2011; Marchitto et al., 2000), (6) master shell chronologies can be created that are as statistically robust as tree ring chronologies (Butler et al., 2010), (7) it precipitates its aragonitic shell in oxygen isotope equilibrium with ambient seawater (Weidman et al., 1994), and (8) the geochemical signature (e.g., ^{14}C , $\delta^{18}\text{O}$, $\delta^{13}\text{C}$) from shell material has been used to reconstruct ocean circulation, hydrographic changes, seasonal changes in ocean conditions, and ecosystem dynamics (Butler et al., 2009; Schöne et al., 2005, 2011a; Scourse et al., 2012; Wanamaker Jr. et al., 2008a, b, 2009, 2011, 2012; Weidman and Jones, 1993; Weidman et al., 1994; Witbaard and Bergman, 2003). Despite their effectiveness as environmental proxies, applications of non-traditional isotopic systems ($\delta^{11}\text{B}$ and $\delta^{88/86}\text{Sr}$) that have the potential to reveal additional environmental factors have not been explored.

2 Methods

2.1 Sample preparation

Living shells from the Gulf of Maine were collected from Jonesport, Maine, USA on 21 November 2009 with a commercial quahog-fishing vessel, FV *Three of A Kind*. The live-caught animals were then transported to the Darling Marine Center (University of Maine) in Walpole, Maine, USA for the culture experiment. Seawater was pumped

$\delta^{11}\text{B}$, $^{87}\text{Sr}/^{86}\text{Sr}$ and $\delta^{88/86}\text{Sr}$ of cultured *Arctica islandica*

Y.-W. Liu et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

from the Damariscotta River estuary from ~ 10 m and delivered to the flowing seawater laboratories. The shells were reared in flow-through seawater tanks without filtration, in which the temperature, salinity and seawater pH were monitored in situ concurrently and continuously (Fig. 1d). The pH of the culture seawater was also measured a total of seven times with a highly accurate Metrohm handheld pH meter (± 0.003 units). Additional details of the culture conditions are given in Beirne et al. (2012). Tank seawater was sampled biweekly throughout the culture period and filtered through a $0.45\ \mu\text{m}$ filter. Two additional samples, one from the Gulf of Maine surface seawater and one from auxiliary water flow pumped into the culture tanks, were also sampled to evaluate if the culture experiment was representative of the natural marine environment. Boron isotopic data from seawater samples during the experiment were previously measured and published by Liu et al. (2013).

Shell subsamples were collected at Iowa State University via a Dremel hand drill, with 10 intervals throughout the 8 month culturing (Fig. 1c). Based on the calcein staining (see Beirne et al., 2012 for details) and natural marking on the external shell, the timing of the winter (January–March), spring (March–May), and summer (May–August) growing seasons were evident (Fig. 1). These markings were used to establish growth rates during each season as well as to provide temporal controls on the sampled shell material. The instrumental data and shell growth-rates have been published by Beirne et al. (2012) and the average seawater salinity, temperature, and pH for shell record comparison are summarized in Table 1.

The boron and strontium concentrations in *A. islandica* shells are about 10 and 1000 ppm, respectively (Zhang, 2009); 1 mg of shell material, after cleaning (details noted below), is required for B (Liu et al., 2013) and Sr (Stevenson et al., 2014) isotopic measurements. Because shell growth rates differ throughout the year and throughout ontogeny (Beirne et al., 2012), we have one subsample from January to March, but we have four subsamples from March to May and five subsamples from May to August.

The subsample cleaning procedure was modified from Barker et al. (2003). In summary: coral and shell powders were first cleaned with Super-Q (SQ) water

(Millipore, > 18.2 MΩ) in an ultrasonic bath three times and the suspension solution was extracted after centrifuging. Samples were then treated with 10% H₂O₂ at room temperature overnight to remove organic particles embedded in skeletons. The samples were rinsed with SQ water, 0.001 N HNO₃ and SQ water again. After drying at 60 °C, samples were weighed and then dissolved in ~1.7 N HCl with a boron concentration equal to about 750 ppb.

Seawater samples were diluted to [B] = ~750 ppb in 1.7 N HCl for boron measurements. Because the strontium concentration in seawater is about 9 ppm, the residual after sublimation (see below) is not enough for strontium isotopes measurements. Therefore for strontium isotope analysis of seawater, an additional 100 μL of seawater sample was dried and redissolved in concentrated HNO₃ three times and brought into solution in 7 N HNO₃ for column chemistry. Spiked samples were mixed to have a sample to spike ratio of 1 : 1, with at least 600 ng of sample Sr in solution prior to preconditioning steps.

Separation of the boron and strontium elemental fractions was achieved using a combination of micro-sublimation (Liu et al., 2013) and elemental specific ion exchange resin. Briefly, < 50 μL of sample solution was loaded in the cap of a conic-bottom vials in an up-side-down position and put into the high-throughput system. After 12 h of sublimation at 70–74 °C, the purified boron sample solution is condensed and collected on the conic tip. To further improve the reproducibility for carbonate samples, an extra 2 μL of 30% H₂O₂ was added for all the runs after 23 April 2014. The cap of the conic vial was then loosened for two hours to reduce the organic levels and liberate the product oxygen gas. The micro-sublimation method only extracts boron from the sample solution, therefore the residual on the cap of the vials is reserved for Sr separation and analysis.

All measurements were conducted on a Thermo Fisher Triton PLUS multicollector thermal ionization mass spectrometer operating in positive ion mode for strontium isotope analysis and negative ion mode for boron isotope analysis at the

BGD

12, 2979–3023, 2015

$\delta^{11}\text{B}$, $^{87}\text{Sr}/^{86}\text{Sr}$ and $\delta^{88/86}\text{Sr}$ of cultured *Arctica islandica*

Y.-W. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.2 Radiogenic and stable strontium isotope analysis

2.2.1 General

5 The residuals of carbonate samples after sublimation were redissolved in concentrated HNO₃ to [Sr] = ~10 ppm, which is approximately the same Sr concentration in seawater. A small aliquot of sample was spiked with our ⁸⁴Sr-⁸⁷Sr double spike solution. Both unspiked normal sample and spiked mixture sample solutions were then dried down three times, then dissolved in 7 N HNO₃ for Sr column chemistry. In order to
10 separate strontium element from matrix, samples were passed through a 50–100 μm Sr-spec resin (Eichrom), and 0.035 N HNO₃ was used to elute Sr after using 7 N HNO₃ to elute the others. The eluted Sr aliquots were refluxed with 30 % H₂O₂ overnight, dried, and finally dissolved in concentrated HNO₃ for loading (Liu, 2010).

15 For radiogenic isotopic measurements, 100–200 ng of Sr sample was loaded onto outgassed Re filaments with TaF₅ activator solution. Each sample was heated to an intensity of ~8 V ⁸⁸Sr. Four hundred cycles of data were collected for each measurement with a ⁸⁵Rb-centered cup configuration. The long-term reproducibility of ⁸⁷Sr/⁸⁶Sr for SRM987 Sr standard was 0.710268 ± 21 (2σ, n = 140) from the time of Triton installation in January 2012–April 2014. In May 2014, problems associated with
20 the H3 Faraday cup resulted in change to a ⁸⁶Sr-centered cup configuration (June–October 2014). After H3 cup replacement, the ⁸⁵Rb-centered cup configuration was established again. The new SRM987 ⁸⁷Sr/⁸⁶Sr average value is 0.719246 ± 13 (2σ, n = 42) for data collected after June 2014 (for both cup configurations) and sample data were normalized based on this new SRM987 standard ratio. The reported ⁸⁷Sr/⁸⁶Sr
25 data in this study were all normalized to SRM987 = 0.710250 for inter-laboratory comparisons.

$\delta^{11}\text{B}$, $^{87}\text{Sr}/^{86}\text{Sr}$ and $\delta^{88/86}\text{Sr}$ of cultured *Arctica islandica*

Y.-W. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.2.2 Double spike (84–87) Sr

High precision $^{88}\text{Sr}/^{86}\text{Sr}$ isotopic compositions are measured by double spike (DS); measurements by DS removes instrumental mass bias associated with thermal ionization during sample runs. The double spike method was first developed in the 1960s: Dodson (1963) outlined a methodology for determining the unknown mass discrimination factor directly if the sample is mixed with a double spike, consisting of an enriched mixture of two stable isotopes. Later, Krogh (1964) worked out a graphical method for a spike enriched in both ^{84}Sr and ^{86}Sr . Finally, Long (1966) showed that the correction factor for mass discrimination can be calculated by using three elements, and it depends on the fractional abundances in the normal and spike elements. A simpler expression was then published by Boelrijk (1968). These pioneering studies founded the basis of the Sr double spike method, and Sr double spikes have already been successfully used to determine the Sr isotopic composition of the early solar system (Patchett, 1980a, b). Optimal spike compositions are determined using a 3-D data reduction method (Galer, 1999). The choice of isotope ratios used in the equations, the mathematical formulation to solve for the mass discrimination factor ε and the influence of the spike-to-sample ratio in the mixture should be taken into consideration. With the addition of “tuning” with IAPSO seawater standards (Krabbenhöft et al., 2009), this double spike method could produce more precise true isotopic compositions in an unknown sample solution.

A ^{84}Sr - ^{87}Sr double spike solution was prepared at GIGL at the University of Michigan followed the method from Liu (2010). The optimal value of the spike depends on the angle of two planes, defined by normal sample (N) fractionation line and spike point and mixture (M) fractionation line and spike point, respectively (Fig. 2), expressed as θ here. Because all the measured points have their own errors, when the angle between these two planes approximate a right angle, the intersected area reaches a minimum and thus that the N-M-S line will be defined precisely (Fig. 2b). Therefore, by checking different portions of spike mixing with normal (Qs) mathematically, the distribution of

BGD

12, 2979–3023, 2015

$\delta^{11}\text{B}$, $^{87}\text{Sr}/^{86}\text{Sr}$ and
 $\delta^{88/86}\text{Sr}$ of cultured
Arctica islandica

Y.-W. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



θ to spike composition can be derived (Fig. 2c). For our ^{84}Sr – ^{87}Sr double spike with an $^{84}\text{Sr}/^{87}\text{Sr}$ ratio of 0.8679, the optimal Q_s is about 0.5, which makes the two planes perpendicular to each other. Within a range of $Q_s = 0.45$ – 0.55 , the θ is still in a range of $\pm 2^\circ$, which supports a tolerance of spiking samples with a slight deviation from a 1 : 1 sample to spike ratio. The stable Sr data were reported as $\delta^{88/86}\text{Sr}$, which was defined as:

$$\delta^{88/86}\text{Sr} = \left[\frac{(^{88}\text{Sr}/^{86}\text{Sr})_{\text{sample}}}{(^{88}\text{Sr}/^{86}\text{Sr})_{\text{SRM 987}}} - 1 \right] \cdot 1000 (\text{‰}) \quad (3)$$

In this study, a Python script was applied to evaluate true normal values of the shells. We assumed the measured isotopic ratios of normal sample and spiked mixture would follow exponential law, and the equations of each ^{87}Sr -based isotopic ratio for both normal and mixture sample could be stated. After inputting the known isotopic composition of the spike, the true value of each isotope concentration can be solved using a least squares minimization of the residual of the non-linear equations.

For the spiked sample solution, 200–250 ng of Sr sample was loaded onto outgassed Re filaments with TaF activator solution and run the same as the unspiked samples described in the previous section. The deconvolved $\delta^{88/86}\text{Sr}$ values for seawater standard IAPSO 141 and an inter-laboratory coral standard JCp-1 are $0.390 \pm 16\text{‰}$ (2σ , $n = 4$) and $0.144 \pm 26\text{‰}$ (2σ , $n = 3$), respectively. According to Krabbenhöft et al. (2009), using the IAPSO seawater standard to fine-tune the Sr double spike composition provides an optimal $\delta^{88/86}\text{Sr}$ result. Here we used the same technique and obtained a compatible $\delta^{88/86}\text{Sr}$ value for the IAPSO seawater standard to the reported values from Krabbenhöft et al. (2009). However, due to the H3 cup deficiency, the deconvolved $\delta^{88/86}\text{Sr}$ value of inter-laboratory carbonate standard JCp-1 is about 0.05‰ lighter than the reported values between April 2014 and June 2014. This offset was fixed, with the alternate cup configuration and replacement of H3 cup,

 $\delta^{11}\text{B}$, $^{87}\text{Sr}/^{86}\text{Sr}$ and
 $\delta^{88/86}\text{Sr}$ of cultured
Arctica islandica

Y.-W. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



after June 2014 and new values of IAPSO = $0.365 \pm 73\text{‰}$ (2σ , $n = 4$) and JCp-1 = $0.195 \pm 21\text{‰}$ (2σ , $n = 4$) (Fig. 3) were obtained.

2.3 Boron isotope analysis

The procedure used for obtaining B isotopic compositions by total evaporation (TE) is described in Liu et al. (2013). In summary: one μL of boron free synthetic seawater matrix was loaded onto outgassed single Re filament at 0.8 A current, followed by 1 μL of sample solution with 30 s of waiting between the two steps. Samples were then dried down at 2 A current for 10 s and then the filaments were flashed to a dull red color in the center of the filament (about 2.5 A) and ready for analysis. Data collection was initiated when the intensity of mass 42 reached 20 mV, and terminated after the signal dropped lower than the initial 20 mV.

The long-term reproducibility (18 months) of $^{11}\text{B}/^{10}\text{B}$ for boric acid standard SRM 951a is 4.0332 ± 0.0064 (2σ , $n = 97$) before treated with 30% H_2O_2 and is 4.0316 ± 0.0084 (2σ , $n = 19$) after the extra treatment. The precisions of $\delta^{11}\text{B}$ for seawater and biogenic carbonate standards without addition of peroxide are $40.46 \pm 1.29\text{‰}$ (2σ , $n = 54$) and $24.94 \pm 2.35\text{‰}$ (2σ , $n = 39$) for IAEA B-1 and JCp-1, respectively; $41.70 \pm 1.13\text{‰}$ (2σ , $n = 8$) and $24.93 \pm 1.83\text{‰}$ (2σ , $n = 18$) for IAEA B-1 and JCp-1, respectively, with H_2O_2 treatment (Fig. 4).

3 Results

The in situ seawater salinity, temperature and pH results are summarized in Table 1. To compare the instrumental data to the shell records, the instrumental results were averaged with respect to the subsampling intervals. The seawater and shell results are summarized in Tables 2 and 3, respectively.

BGD

12, 2979–3023, 2015

$\delta^{11}\text{B}$, $^{87}\text{Sr}/^{86}\text{Sr}$ and
 $\delta^{88/86}\text{Sr}$ of cultured
Arctica islandica

Y.-W. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3.1 $^{87}\text{Sr}/^{86}\text{Sr}$ and $\delta^{88/86}\text{Sr}$

3.1.1 $^{87}\text{Sr}/^{86}\text{Sr}$

The $^{87}\text{Sr}/^{86}\text{Sr}$ ratios of seawater range from 0.709177 to 0.709192, with an average of 0.709185 ± 8 (2σ , $n = 13$). There is no distinguishable difference between samples from offshore Gulf of Maine seawater, auxiliary flow and tank waters (Fig. 5a). For shell carbonate, the $^{87}\text{Sr}/^{86}\text{Sr}$ ratios range from 0.709163 to 0.709210, with an average of 0.709183 ± 23 (2σ , $n = 27$). Both of the seawater and shell $^{87}\text{Sr}/^{86}\text{Sr}$ are identical to the mean seawater values (Fig. 5a and b). Because all the radiogenic Sr results are identical within error, there is no relationship with either seawater salinity, temperature or pH.

3.1.2 $\delta^{88/86}\text{Sr}$

Two sets of shell samples, A103JV and A105JV, were spiked for stable Sr measurements. However, due to the defect of H3 cup, the A103JV double spike results are underestimated and are not listed. The deconvolved $\delta^{88/86}\text{Sr}$ for A105JV and Sr concentrations values for both of A103JV and A105JV from high-resolution shell records are shown in Fig. 5c and d, respectively. The $\delta^{88/86}\text{Sr}$ values range from 0.215 to 0.296‰ with an average of 0.248 ± 48 ‰ (2σ , $n = 10$) and the concentration ranges from 1200 to 1800 ppm. Considering analytical uncertainty, no distinctive differences or trend for either Sr concentration or stable Sr isotopic compositions in the shells are observed throughout the culture season. Similarly, no correlation can be found between stable Sr or Sr concentration to any measured ambient seawater conditions.

BGD

12, 2979–3023, 2015

$\delta^{11}\text{B}$, $^{87}\text{Sr}/^{86}\text{Sr}$ and $\delta^{88/86}\text{Sr}$ of cultured *Arctica islandica*

Y.-W. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3.2 Boron isotopic composition ($\delta^{11}\text{B}$) in ambient seawater the shell and aragonite shell

Boron isotopic compositions of 11 tank water samples are from 37.51 to 47.83‰, and the average for 36 sample runs is 39.20 ± 1.73 ‰. The $\delta^{11}\text{B}$ values for additional seawater samples from the offshore Gulf of Maine and the auxiliary flow to the culture tanks are 36.93 and 40.38‰, respectively (Fig. 6a) (Liu et al., 2013). Similar to what has been observed from radiogenic Sr data in seawater, the boron isotopic composition of our culture seawater is invariant and identical to the open ocean composition reported by Foster et al. (2010).

High-resolution boron isotopic composition records from 4 juvenile shells (A101JV, A102JV, A103JV and A105JV) show nearly identical patterns and trends throughout the experiment. The shell $\delta^{11}\text{B}$ values range from 11.09 to 18.81‰ before week 19 and from 17.41 to 25.12‰ after week 19 (Fig. 6b). Compared to seawater temperature and average shell growth rates in three growth seasons, we found a distinct rise in $\delta^{11}\text{B}$ for temperatures over $\sim 13^\circ\text{C}$. However, this rapid change in $\delta^{11}\text{B}$ did not correlate to the rapid change in shell growth during the culture period.

In order to evaluate the pH dependency of the $\delta^{11}\text{B}$, Eq. (2) is used. Based on in situ temperature, salinity and pH measurements throughout the culture experiment, we calculated the predicted range in $\delta^{11}\text{B}$ of the *A. islandica* shell based on two boron fractionation factors. The lower estimations are based on $\alpha = 1.0194$, a theoretical result of Kakihana et al. (1977), which has been applied widely on paleo-reconstructions (Hönisch et al., 2004; Sanyal et al., 1995; Kakihana et al., 1977). The higher predicted range was calculated with $\alpha = 1.0272$, which was empirically obtained from Klochko et al. (2006) and is considered better to describe the distribution of the two boron species in natural system (Foster, 2008; Pagani et al., 2005; Klochko et al., 2006) (Fig. 6b). The predictions suggest a slight increase in $\delta^{11}\text{B}$ throughout the culture season, primarily due to more than a 15°C temperature increase. A 0.2 pH unit drop, observed between week 24 and 26, should have decreased the $\delta^{11}\text{B}$ value by about

BGD

12, 2979–3023, 2015

$\delta^{11}\text{B}$, $^{87}\text{Sr}/^{86}\text{Sr}$ and $\delta^{88/86}\text{Sr}$ of cultured *Arctica islandica*

Y.-W. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2‰ even with the large temperature change. ~~All of the shell boron isotopic composition data lie between the two prediction lines, with samples before week 19 lying on the $\alpha = 1.0272$ line.~~ After week 19, the $\delta^{11}\text{B}$ deviate significantly and trend toward higher and higher $\delta^{11}\text{B}$ compositions. Duplications on different individual shells all suggest the same trend.

We also calculated shell pH with $\alpha = 1.0272$ based on shell $\delta^{11}\text{B}$ and the corresponding average seawater temperature and salinity values. The results show ~~fairly strong~~ negative relationship between ΔpH and pH_{sw} ($R^2 = 0.34$; p value ≤ 0.001).

4 Discussion

4.1 Radiogenic Sr isotope incorporation into *A. islandica*

In this study, we measured the radiogenic Sr isotope ratios to estimate the source water contributions to the culture site, which is situated within the Damariscotta River estuary. The $^{87}\text{Sr}/^{86}\text{Sr}$ ratios in cultured seawater showed identical values to the open surface seawater ratio. Bedrock types in the Gulf of Maine coastal region are dominated by late Proterozoic and lower Paleozoic sedimentary rocks (Osberg et al., 1985), which would provide a terrestrial source with high $^{87}\text{Sr}/^{86}\text{Sr}$ values. Considering rock type, age and freshwater flux, a recent model of $^{87}\text{Sr}/^{86}\text{Sr}$ for flux-weighted catchment water suggests the $^{87}\text{Sr}/^{86}\text{Sr}$ value to be in a range of 0.7099 to 0.7145 (Bataille and Bowen, 2012). If river fluxes influence the Sr isotopic composition of coastal seawater, the value should be enriched in ^{87}Sr , driving the $^{87}\text{Sr}/^{86}\text{Sr}$ higher than the current seawater ratio. Therefore the Sr isotopic results suggest a negligible amount of fresh water input into the culture water at the Darling Marine Center.

The mean $^{87}\text{Sr}/^{86}\text{Sr}$ ratio in the shell is consistent with the isotopic composition in the culture seawater, but with a relatively larger variation between individual shell samples. Therefore, incorporation of radiogenic Sr ratios into the shells ~~are in equilibrium with~~ ambient seawater. Although the shell $^{87}\text{Sr}/^{86}\text{Sr}$ values have a larger

BGD

12, 2979–3023, 2015

$\delta^{11}\text{B}$, $^{87}\text{Sr}/^{86}\text{Sr}$ and $\delta^{88/86}\text{Sr}$ of cultured *Arctica islandica*

Y.-W. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



SD compared to the seawater values, they are within the range of the long-term precision of Triton plus at the Department of Earth and Environmental Sciences, University of Michigan (see Sect. 2.2.1). The high content of calcium in carbonate samples, which cannot be fully separated using Sr-specific ion exchange column chemistry, may contribute to the larger variation of shell $^{87}\text{Sr}/^{86}\text{Sr}$ compared to seawater.

4.2 Stable Sr isotope incorporation into *A. islandica*, and Sr concentrations

In this study we observed no statistically significant correlation of $\delta^{88/86}\text{Sr}$ or Sr concentrations with respect to seawater temperature. The stable Sr isotopic composition of some biogenic carbonates has been suggested to reflect ambient seawater temperature due to mass dependent kinetic fractionation, in which the relative mass difference of the isotopes involved accounts for the inverse correlation to the ion mass in a kinetic fractionation process (Fietzke and Eisenhauer, 2006; Rüggeberg et al., 2008). However, more recent work has shown no relationship between seawater temperature and $\delta^{88/86}\text{Sr}$ values from various biogenic archives (Böhm et al., 2012; Stevenson et al., 2014; Raddatz et al., 2013; Vollstaedt et al., 2014). Our results support the argument that a simple temperature dependent kinetic effect is not the primary control of stable Sr in the aragonitic shell of *A. islandica*. The lack of a consistent relationship between stable Sr with temperature or shell growth rate during the experiment indicates that stable Sr is not controlled by temperature or growth rates. Thus, it may be possible stable Sr records derived from *A. islandica* shells may reflect ambient seawater conditions, and could therefore be a potential archive for studying the global Sr cycle in the context of chemical weathering (Krabbenhöft et al., 2010; Raddatz et al., 2013; Vollstaedt et al., 2014). More work is needed to fully evaluate this potential proxy.

Our deconvolved shell Sr concentrations show no relationship to the seawater temperature, which may indicate a physiological control on Sr uptake into the bivalve

BGD

12, 2979–3023, 2015

$\delta^{11}\text{B}$, $^{87}\text{Sr}/^{86}\text{Sr}$ and
 $\delta^{88/86}\text{Sr}$ of cultured
Arctica islandica

Y.-W. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



shell material. In general, co-precipitation of Sr to Ca in aragonite decreases with temperature increases due to a declining distribution coefficient, which has been both measured and derived theoretically (Dietzel et al., 2004). The negative correlation between skeletal Sr/Ca ratios observed in some massive corals with ambient seawater temperature has been widely established and applied to reconstruct paleo seawater temperature (Beck et al., 1992; Corrège, 2006; McCulloch et al., 1994, 1996; de Villiers, 1999; Yu et al., 2005; Shen et al., 1996; Yan et al., 2013; Weber, 1973). However, these relationships may be biased by the influence from symbionts potentially causing a vital effect (Böhm et al., 2012; Cohen et al., 2006, 2002; Stevenson et al., 2014). Schöne et al. (2011b) observed a strong physiological regulation to Sr/Ca and Mg/Ca on ontogenically old adult *A. islandica* records. The metal to calcium ratio (Me/Ca) increased with shell age when the annual increment widths were below 30 to 200 μm . However, Schöne et al. (2013) concluded that the faster-growing juvenile portion of the shells showed a weak relationship between Me/Ca and ambient temperature and results from different specimens were variable. Schöne et al. (2013) proposed that pronounced vital effects may control the trace metal uptake in juvenile shells. Our results are also consistent with the study of Schöne et al. (2011b), where the juvenile portion of the two *A. islandica* shells with annual growth increments larger than 750 μm showed no clear relationship between Me/Ca ratios and growth rates.

4.3 Controls on $\delta^{11}\text{B}$ in *A. islandica* and an evaluation of the proxy archive as a seawater pH indicator

The range in measured shell $\delta^{11}\text{B}$ values lies between the prediction lines (Fig. 6b), which suggest that the shell boron content generally reflects the ambient seawater conditions. However, our data do not consistently follow either prediction line. Previous studies on foraminifera have shown offsets between different genera and the empirical $\delta^{11}\text{B}$ -pH relationship of $\alpha = 1.0194$ (Hönisch and Hemming, 2004). The inconsistency between shell $\delta^{11}\text{B}$ and either prediction is therefore strong evidence that a species-specific fractionation factor is required for bivalves.

$\delta^{11}\text{B}$, $^{87}\text{Sr}/^{86}\text{Sr}$ and $\delta^{88/86}\text{Sr}$ of cultured *Arctica islandica*

Y.-W. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



$\delta^{11}\text{B}$, $^{87}\text{Sr}/^{86}\text{Sr}$ and $\delta^{88/86}\text{Sr}$ of cultured *Arctica islandica*

Y.-W. Liu et al.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


The offsets between our shell data and the predicted trends (Fig. 6b) are likely from vital effects during biomineralization. Rollion-Bard and Erez (2010) and Trotter et al. (2011) evaluated vital effects in corals and foraminifera, and the potential use of the $\delta^{11}\text{B}$ -pH relationship in such biogenic carbonates. They observed a pH offset between calcifying fluid and ambient seawater, and this pH discrepancy (ΔpH) increases with decreasing ambient seawater pH (pH_{sw}). As the environment becomes more acidic, marine calcifiers likely adjust their internal microenvironment during calcification, resulting in larger ΔpH values than expected. Under careful culture conditions, species-specific ΔpH - pH_{sw} relationships can be developed and, after calibration, the corresponding ambient seawater pH can be determined.

It has been argued that bivalves have the ability to regulate their inner shell fluid chemistry, more specifically the extrapallial fluid (EPF), to achieve a carbonate saturation state in order to facilitate biomineralization (Crenshaw, 1980). Stemmer (2013) observed pH gradients between inner shell surface and outer mantle epithelium (OME) of *A. islandica* via in situ pH microscopy method. During a short-term monitoring, Stemmer (2013) also observed that rapid pH elevation happened close to the OME. She concluded this elevation can only be due to active proton uptake by epithelium. This result suggests a pH self-regulation occurs for *A. islandica* shell precipitation.

The calculated ΔpH shows a statistically significant negative relationship to pH_{sw} ($R^2 = 0.34$; p value ≤ 0.001). The negative correlation supports the argument that *A. islandica* regulate their EPF pH for calcification. However, only 34% of the variability can be explained by ΔpH , which indicates that the pH regulation in the EPF is likely not the prevailing factor. Instead, we found a rapid increase of the shell $\delta^{11}\text{B}$ when temperature increased over 13°C (Fig. 7a). This rapid change in boron isotopic composition can be explained with respect to two factors: (1) a growth rate controlled vital effect, or (2) a temperature controlled vital effect.

Herfort et al. (2008) suggested that carbonate species are the limiting factor in coral calcium precipitation rather than calcium: when ambient seawater temperature

increases, $[\text{CO}_2]_{\text{aq}}$ decreases and leads to rising $[\text{CO}_3^{-2}]$ as well as calcification rate. However, in our study, the positive shell $\delta^{11}\text{B}$ trend did not correlate with the rapid change of shell growth after the spring bloom (Fig. 7b), which casts doubt on a kinetic isotopic effect. Therefore, combined correlation with temperature and lack of correlation with growth to the shell $\delta^{11}\text{B}$ cannot be explained by a temperature controlled growth/precipitation effect.

Alternatively, a proton removal mechanism via Ca^{+2} -ATpase from the site of calcification has been proposed. This mechanism raises the pH of the calcification solution (Dissard et al., 2012; Rollion-Bard et al., 2011). In this scenario, the activity of the enzyme is enhanced when a certain temperature has been reached, accelerating the proton removal process and resulting in a higher boron isotopic composition in the calcification solution with respect to the elevated pH. We suggest that there may be a temperature threshold of the boron incorporation into the aragonite *A. islandica*. This proposed threshold may be related to the upper-end thermal tolerance of *A. islandica*. For example, below 13°C , the $\delta^{11}\text{B}$ values closely matched the predicted model of Klochko et al. (2006), supporting the assumption that borate is the dominate species incorporated into the shell and so can reflect the ambient seawater pH. At temperatures above 13°C the utility of the $\delta^{11}\text{B}$ values as a pH indicator is questionable and likely unreliable. We suggest that the thermal tolerance of *A. islandica* was exceeded in the summer growing season in the culture conditions, causing biological stress on the animals.

5 Conclusion

Here we examined boron and strontium isotopic compositions recorded in the shell material of cultured aragonitic *A. islandica* with in situ seawater temperature, salinity and pH measurements. Both seawater and shell $^{87}\text{Sr}/^{86}\text{Sr}$ show identical values to the mean global seawater composition, suggesting there is trivial influence from

BDG

12, 2979–3023, 2015

$\delta^{11}\text{B}$, $^{87}\text{Sr}/^{86}\text{Sr}$ and $\delta^{88/86}\text{Sr}$ of cultured *Arctica islandica*

Y.-W. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



local continental runoff. Shell $\delta^{88/86}\text{Sr}$ and Sr concentration values during the culture season were not influenced by seawater temperature or calcification rates. These results suggest that well-preserved sub-fossil specimens may be use to determine the past isotopic composition ($^{87}\text{Sr}/^{86}\text{Sr}$ and $\delta^{88/86}\text{Sr}$) of seawater.

The boron isotope results from the cultured aragonite *A. islandica* shells generally are within the range of two prediction lines utilizing previously published fractionation factors. Although to first order, these results indicate that the shell $\delta^{11}\text{B}$ values reflect ambient conditions, substantial variability not related to pH changes was noted. The 5–8‰ increase in shell $\delta^{11}\text{B}$ values is larger than theoretical predictions based on in situ seawater temperature, salinity, pH and conventional boron fractionation factors for corals and foraminifera. A species-specific $\delta^{11}\text{B}$ –pH transfer function is recommended for bivalve species because of their inherent ability to self regulate calcifying fluids. The fairly strong relationship ($r^2 = 0.34$) between ΔpH and pH_{sw} indicates that *A. islandica* does regulate the EPF pH during calcification, but self-regulation is not the primary control of the shell $\delta^{11}\text{B}$. The largest increase in shell $\delta^{11}\text{B}$ values was observed after crossing an apparent temperature threshold at 13 °C, suggesting a possible influence from biological processes. To better evaluate the potential of $\delta^{11}\text{B}$ as a seawater pH indicator, a pH controlled culture experiment with limited seawater temperature and salinity variation is needed.

Acknowledgements. This project was funded by grants from the Rackham Graduate School and the Department of Earth and Environmental Sciences at the University of Michigan to Y.-W. Liu and the Packard Foundation to S.M. Aciego. We thank Dr. Tsuyoshi Watanabe, Hokkaido University, Japan for providing an aliquot of the international coral standard (JCP-1).

References

Aarons, S. M., Aciego, S. M., and Gleason, J. D.: Variable HfSrNd radiogenic isotopic compositions in a Saharan dust storm over the Atlantic: implications for dust flux

BGD

12, 2979–3023, 2015

$\delta^{11}\text{B}$, $^{87}\text{Sr}/^{86}\text{Sr}$ and $\delta^{88/86}\text{Sr}$ of cultured *Arctica islandica*

Y.-W. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



$\delta^{11}\text{B}$, $^{87}\text{Sr}/^{86}\text{Sr}$ and $\delta^{88/86}\text{Sr}$ of cultured *Arctica islandica*

Y.-W. Liu et al.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)

[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


to oceans, ice sheets and the terrestrial biosphere, *Chem. Geol.*, 349–350, 18–26, doi:10.1016/j.chemgeo.2013.04.010, 2013.

Anagnostou, E., Huang, K. F., You, C. F., Sikes, E. L., and Sherrel, R. M.: Evaluation of boron isotope ratio as a pH proxy in the deep sea coral *Desmophyllum dianthus*: evidence of physiological pH adjustment, *Earth Planet. Sc. Lett.*, 349, 251–260, 2012.

Barker, S., Greaves, M., and Elderfield, H.: A study of cleaning procedures used for foraminiferal Mg/Ca paleothermometry, *Geochem. Geophys. Geosy.*, 4, 8407, doi:10.1029/2003gc000559, 2003.

Bataille, C. P. and Bowen, G. J.: Mapping $^{87}\text{Sr}/^{86}\text{Sr}$ variations in bedrock and water for large scale provenance studies, *Chem. Geol.*, 304–305, 39–52, doi:10.1016/j.chemgeo.2012.01.028, 2012.

Beck, J. W., Edwards, R. L., Ito, E., Taylor, F. W., Recy, J., Rougerie, F., Joannot, P., and Henin, C.: Sea-surface temperature from coral skeletal strontium/calcium ratios, *Science*, 257, 644–647, doi:10.1126/science.257.5070.644, 1992.

Beirne, E.: Pursuing a proxy for carbon cycling in the temperate North Atlantic: Investigation of the utility of *Arctica islandica* shell carbonate to millennial-scale dissolved inorganic carbon reconstructions, M.S. thesis, Geological and Atmospheric Sciences, Iowa State University, Graduate Theses and Dissertations. Paper 10124, Ames, 193 pp., 2011.

Beirne, E. C., Wanamaker Jr., A. D., and Feindel, S. C.: Experimental validation of environmental controls on the $\delta^{13}\text{C}$ of *Arctica islandica* (ocean quahog) shell carbonate, *Geochim. Cosmochim. Ac.*, 84, 395–409, doi:10.1016/j.gca.2012.01.021, 2012.

Berglund, M. and Wieser, M. E.: Isotopic compositions of the elements 2009 (IUPAC Technical Report), *Pure Appl. Chem.*, 83, 397–410, 2011.

Boelrijk, N. A. I. M.: A general formula for “double” isotope dilution analysis, *Chem. Geol.*, 3, 323–325, 1968.

Böhm, F., Eisenhauer, A., Tang, J., Dietzel, M., Krabbenhöft, A., Kisakürek, B., and Horn, C.: Strontium isotope fractionation of planktic foraminifera and inorganic calcite, *Geochim. Cosmochim. Ac.*, 93, 300–314, doi:10.1016/j.gca.2012.04.038, 2012.

Broecker, W. S.: *Radioisotopes and Large-Scale Oceanic Mixing*, Interscience, New York, 88–108, 1963.

Butler, P. G., Scourse, J. D., Richardson, C. A., Wanamaker Jr., A. D., Bryant, C. L., and Bennell, J. D.: Continuous marine radiocarbon reservoir calibration and the ^{13}C Suess effect

$\delta^{11}\text{B}$, $^{87}\text{Sr}/^{86}\text{Sr}$ and $\delta^{88/86}\text{Sr}$ of cultured *Arctica islandica*

Y.-W. Liu et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

in the Irish Sea: results from the first multi-centennial shell-based marine master chronology, *Earth Planet. Sc. Lett.*, 279, 230–241, 2009.

Butler, P. G., Richardson, C. A., Scourse, J. D., Wanamaker Jr., A. D., Shammon, T. M., and Bennell, J. D.: Marine climate in the Irish Sea: analysis of a 489 year marine master chronology derived from growth increments in the shell of the clam *Arctica islandica*, *Quaternary Sci. Rev.*, 29, 1614–1632, 2010.

Butler, P. G., Wanamaker Jr., A. D., Scourse, J. D., Richardson, C. A., and Reynolds, D. J.: Long-term stability of $\delta^{13}\text{C}$ with respect to biological age in the aragonite shell of mature specimens of the bivalve mollusk *Arctica islandica*, *Palaeogeogr. Palaeoclimatol.*, 302, 21–30, 2011.

Byrne, R. H., Mecking, S., Feely, R. A., and Liu, X.: Direct observations of basin-wide acidification of the North Pacific Ocean, *Geophys. Res. Lett.*, 37, L02601, doi:10.1029/2009GL040999, 2010.

Chung, C.-H., You, C.-F., and Chu, H.-Y.: Weathering sources in the Gaoping (Kaoping) river catchments, southwestern Taiwan: insights from major elements, Sr isotopes, and rare earth elements, *J. Marine Syst.*, 76, 433–443, 2009.

Cohen, A. L., Owens, K. E., Layne, G. D., and Shimizu, N.: The effect of algal symbionts on the accuracy of Sr/Ca paleotemperatures from coral, *Science*, 296, 331–333, doi:10.1126/science.1069330, 2002.

Cohen, A. L., Gaetani, G. A., Lundalv, T., Corliss, B. H., and George, R. Y.: Compositional variability in a cold-water scleractinian, *Lophelia pertusa*: new insights into “vital effects”, *Geochem. Geophys. Geosyst.*, 7, Q12004, doi:10.1029/2006GC001354, 2006.

Correge, T.: Sea surface temperature and salinity reconstruction from coral geochemical tracers, *Palaeogeogr. Palaeoclimatol.*, 232, 408–428, doi:10.1016/j.palaeo.2005.10.014, 2006.

Crenshaw, M. A.: Mechanisms of shell formation and dissolution, in: *Skeletal Growth of Aquatic Organisms*, edited by: Rhoads, D. C., and Lutz, R. A., Plenum Press, New York, 115–132 pp., 1980.

de Villiers, S.: Seawater strontium and Sr/Ca variability in the Atlantic and Pacific oceans, *Earth Planet. Sc. Lett.*, 171, 623–634, doi:10.1016/S0012-821X(99)00174-0, 1999.

D’Olivo, J. P., McCulloch, M. T., Eggins, S. M., and Trotter, J.: Coral records of reef-water pH across the central Great Barrier Reef, Australia: assessing the influence of river runoff on inshore reefs, *Biogeosciences Discuss.*, 11, 11443–11479, doi:10.5194/bgd-11-11443-2014, 2014.

$\delta^{11}\text{B}$, $^{87}\text{Sr}/^{86}\text{Sr}$ and
 $\delta^{88/86}\text{Sr}$ of cultured
Arctica islandica

Y.-W. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Dietzel, M., Gussone, N., and Eisenhauer, A.: Co-precipitation of Sr^{2+} and Ba^{2+} with aragonite by membrane diffusion of CO_2 between 10 and 50°C , *Chem. Geol.*, 203, 139–151, doi:10.1016/j.chemgeo.2003.09.008, 2004.

Dissard, D., Douville, E., Reynaud, S., Juillet-Leclerc, A., Montagna, P., Louvat, P., and McCulloch, M.: Light and temperature effects on $\delta^{11}\text{B}$ and B/Ca ratios of the zooxanthellate coral *Acropora* sp.: results from culturing experiments, *Biogeosciences*, 9, 4589–4605, doi:10.5194/bg-9-4589-2012, 2012.

Dodson, M. H.: A theoretical study of the use of internal standards for precise isotopic analysis by the surface ionization technique: Part I – General first-order algebraic solutions, *J. Phys. E Sci. Instrum.*, 40, 289–295, 1963.

DOE: Handbook of methods for the analysis of the various parameters of the carbon dioxide system in sea water, Version 2, ORNL/CDIAC-74, Carbon Dioxide Information and Analysis Center, Oak Ridge, 1994.

Doney, S. C., Fabry, V. J., Feely, R. A., and Kleypas, J. A.: Ocean Acidification: The Other CO_2 Problem, *Annual Review of Marine Science*, 1, 169–192, doi:10.1146/annurev.marine.010908.163834, 2009.

Dore, J. E., Lukas, R., Sadler, D. W., Church, M. J., and Karl, D. M.: Physical and biogeochemical modulation of ocean acidification in the central North Pacific, *P. Natl. Acad. Sci. USA*, 106, 12235–12240, doi: 10.1073/pnas.0906044106, 2009.

Feely, R. A., Sabine, C. L., Lee, K., Berelson, W., Kleypas, J., Fabry, V. J., and Millero, F. J.: Impact of Anthropogenic CO_2 on the CaCO_3 System in the Oceans, *Science*, 305, 362–366, doi:10.1126/science.1097329, 2004

Fietzke, J. and Eisenhauer, A.: Determination of temperature-dependent stable strontium isotope ($^{88}\text{Sr}/^{86}\text{Sr}$) fractionation via bracketing standard MC-ICP-MS, *Geochem. Geophys. Geosy.*, 7, Q08009, doi:10.1029/2006GC001243, 2006.

Foster, G. L.: Seawater pH, $p\text{CO}_2$ and $[\text{CO}_3^{2-}]$ variations in the Caribbean Sea over the last 130 kyr: a boron isotope and B/Ca study of planktic foraminifera, *Earth Planet. Sc. Lett.*, 271, 254–266, 2008.

Foster, G. L., Pogge von Strandmann, P. A. E., and Rae, J. W. B.: Boron and magnesium isotopic composition of seawater, *Geochem. Geophys. Geosy.*, 11, Q08015, doi:10.1029/2010GC003201, 2010.

- Galer, S. J. G.: Optimal double and triple spiking for high precision lead isotopic measurement, *Chem. Geol.*, 157, 255–274, 1999.
- Goldberg, E. D.: *The Oceans as a Chemical System*, Interscience, New York, 3–25, 1963.
- ~~Hemming, N. G. and Hanson, G. N.: Boron isotopic composition and concentration in modern marine carbonates, *Geochim. Cosmochim. Ac.*, 56, 537–543, 1992.~~
- Henehan, M. J., Rae, J. W. B., Foster, G. L., Erez, J., Prentice, K. C., Kucera, M., Bostock, H. C., Martínez-Botí, M. A., Milton, J. A., Wilson, P. A., Marshall, B. J., and Elliott, T.: Calibration of the boron isotope proxy in the planktonic foraminifera *Globigerinoides ruber* for use in palaeo-CO₂ reconstruction, *Earth Planet. Sc. Lett.*, 364, 111–122, doi:10.1016/j.epsl.2012.12.029, 2013.
- Herfort, L., Thake, B., and Taubner, I.: Bicarbonate stimulation of calcification and photosynthesis in two hermatypic corals, *J. Phycol.*, 44, 91–98, doi:10.1111/j.1529-8817.2007.00445.x, 2008.
- Hofmann, G. E., Barry, J. P., Edmunds, P. J., Gates, R. D., Hutchins, D. A., Klinger, T., and Sewell, M. A.: The effect of ocean acidification on calcifying organisms in marine ecosystems: an organism-to-ecosystem perspective, *Annu. Rev. Ecol. Evol. S.*, 41, 127–147, doi:10.1146/annurev.ecolsys.110308.120227, 2010.
- Hönisch, B. and Hemming, N. G.: Ground-truthing the boron isotope-paleo-pH proxy in planktonic foraminifera shells: partial dissolution and shell size effects, *Paleoceanography*, 19, PA4010, doi:10.1029/2004PA001026, 2004.
- Hönisch, B., Hemming, N. G., Grottoli, A. G., Amat, A., Hanson, G. N., and Bijma, J.: Assessing scleractinian corals as recorders for paleo-pH: empirical calibration and vital effects, *Geochim. Cosmochim. Ac.*, 68, 3675–3685, doi:10.1016/j.gca.2004.03.002, 2004.
- Hönisch, B., Ridgwell, A., Schmidt, D. N., Thomas, E., Gibbs, S. J., Sluijs, A., Zeebe, R., Kump, L., Martindale, R. C., and Greene, S. E.: The geological record of ocean acidification, *Science*, 335, 1058–1063, 2012.
- Huang, K.-F. and You, C.-F.: Tracing freshwater plume migration in the estuary after a typhoon event using Sr isotopic ratios, *Geophys. Res. Lett.*, 34, L02403, doi:10.1029/2006GL028253, 2007.
- Huang, K.-F., You, C.-F., Chung, C.-H., and Lin, I.-T.: Nonhomogeneous seawater Sr isotopic composition in the coastal oceans: A novel tool for tracing water masses and submarine groundwater discharge, *Geochem. Geophys. Geosy.*, 12, Q05002, doi:10.1029/2010GC003372, 2011.

 $\delta^{11}\text{B}$, $^{87}\text{Sr}/^{86}\text{Sr}$ and $\delta^{88/86}\text{Sr}$ of cultured *Arctica islandica*

Y.-W. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



IPCC: Climate Change 2013: The Physical Science Basis. Working Group I Contribution to the IPCC 5th Assessment Report – Changes to the Underlying Scientific/Technical Assessment, Cambridge, UK and New York, NY, USA, 2013.

Jahn, B.-M., Gallet, S., and Han, J.: Geochemistry of the Xining, Xifeng and Jixian sections, Loess Plateau of China: eolian dust provenance and paleosol evolution during the last 140 ka, *Chem. Geol.*, 178, 71–94, 2001.

Jones, D. S.: Annual cycle of shell growth increment formation in two continental shelf bivalves and its paleoecologic significance, *Paleobiology*, 6, 331–340, 1980.

Kakihana, H., Kotaka, M., Satoh, S., Nomura, M., and Okamoto, M.: Fundamental studies on the ion-exchange separation of boron isotopes, *B. Chem. Soc. Jpn.*, 50, 158–163, 1977.

Klochko, K., Kaufman, A. J., Yao, W., Byrne, R. H., and Tossell, J. A.: Experimental measurement of boron isotope fractionation in seawater, *Earth Planet. Sc. Lett.*, 248, 276–285, doi:10.1016/j.epsl.2006.05.034, 2006.

Krabbenhöft, A., Fietzke, J., Eisenhauer, A., Liebetrau, V., Böhm, F., and Vollstaedt, H.: Determination of radiogenic and stable strontium isotope ratios ($^{87}\text{Sr}/^{86}\text{Sr}$; $\delta^{88/86}\text{Sr}$) by thermal ionization mass spectrometry applying an $^{87}\text{Sr}/^{84}\text{Sr}$ double spike, *J. Anal. Atom. Spectrom.*, 24, 1267–1271, 2009.

Krabbenhöft, A., Eisenhauer, A., Böhm, F., Vollstaedt, H., Fietzke, J., Liebetrau, V., Augustin, N., Peucker-Ehrenbrink, B., Müller, M. N., Horn, C., Hansen, B. T., Nolte, N., and Wallmann, K.: Constraining the marine strontium budget with natural strontium isotope fractionations ($^{87}\text{Sr}/^{86}\text{Sr}^*$, $\delta^{88/86}\text{Sr}$) of carbonates, hydrothermal solutions and river waters, *Geochim. Cosmochim. Ac.*, 74, 4097–4109, 2010.

Krief, S., Hendy, E. J., Fine, M., Yam, R., Meiborn, A., Foster, G. L., and Shemesh, A.: Physiological and isotopic responses of scleractinian corals to ocean acidification, *Geochim. Cosmochim. Ac.*, 74, 4988–5001, doi:10.1016/j.gca.2010.05.023, 2010.

Kroeker, K. J., Kordas, R. L., Crim, R. N., and Singh, G. G.: Meta-analysis reveals negative yet variable effects of ocean acidification on marine organisms, *Ecol. Lett.*, 13, 1419–1434, doi:10.1111/j.1461-0248.2010.01518.x, 2010.

Krogh, T. E.: Strontium isotopic variation and whole rock isochron studies in the Grenville Province of Ontario, PhD thesis, Department of Geology and Geophysics, Massachusetts Institute of Technology, Cambridge, 1964.

BGD

12, 2979–3023, 2015

$\delta^{11}\text{B}$, $^{87}\text{Sr}/^{86}\text{Sr}$ and $\delta^{88/86}\text{Sr}$ of cultured *Arctica islandica*

Y.-W. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



$\delta^{11}\text{B}$, $^{87}\text{Sr}/^{86}\text{Sr}$ and $\delta^{88/86}\text{Sr}$ of cultured *Arctica islandica*

Y.-W. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Lemarchand, D., Gaillardet, J., Lewin, E., and Allegre, C. J.: The influence of rivers on marine boron isotopes and implications for reconstructing past ocean pH, *Nature*, 408, 951–954, 2000.
- Liu, Y.-W.: Natural variation of Sr isotopes in coral *Porites* collected from Nanwan Bay, southern tip of Taiwan, M.S. thesis, Department of Geosciences, National Taiwan University, Taipei, 1–78 pp., 2010.
- Liu, Y.-W., Aciego, S. M., Wanamaker Jr., A. D., and Sell, B. K.: A high-throughput system for boron microsublimation and isotope analysis by total evaporation thermal ionization mass spectrometry, *Rapid Commun. Mass Sp.*, 27, 1705–1714, doi:10.1002/rcm.6619, 2013.
- Long, L. E.: Isotope dilution analysis of common and radiogenic strontium using ^{84}Sr -enriched spike, *Earth Planet. Sc. Lett.*, 1, 289–292, 1966.
- Marchitto, T. M., Jones, G. A., Goodfriend, G. A., and Weidman, C. R.: Precise temporal correlation of holocene mollusk shells using sclerochronology, *Quaternary Res.*, 53, 236–246, 2000.
- Maurer, A.-F., Galer, S. J. G., Knipper, C., Beierlein, L., Nunn, E. V., Peters, D., Tütken, T., Alt, K. W., and Schöne, B. R.: Bioavailable $^{87}\text{Sr}/^{86}\text{Sr}$ in different environmental samples – Effects of anthropogenic contamination and implications for isoscapes in past migration studies, *Sci. Total Environ.*, 433, 216–229, doi:10.1016/j.scitotenv.2012.06.046, 2012.
- McCulloch, M. T., Mortimer, G., Esat, T., Xianhua, L., Pillans, B., and Chappell, J.: High resolution windows into early Holocene climate: Sr/Ca coral records from the Huon Peninsula, *Earth Planet. Sc. Lett.*, 138, 169–178, 1996.
- McCulloch, M. T., Gagan, M. K., Mortimer, G. E., Chivas, A. R., and Isdale, P. J.: A high-resolution Sr/Ca and $\delta^{18}\text{O}$ coral record from the Great Barrier Reef, Australia, and the 1982–1983 El Niño, *Geochim. Cosmochim. Ac.*, 58, 2747–2754, 1994.
- Merrill, A. and Ropes, J. W.: The general distribution of the surf clam and ocean quahog *Spisula-solidissima Arctica-islandica*, *J. Shellfish Res.*, 59, 40–45, 1968.
- Ni, Y., Foster, G. L., Bailey, T., Elliott, T., Schmidt, D. N., Pearson, P., Haley, B., and Coath, C.: A core top assessment of proxies for the ocean carbonate system in surface-dwelling foraminifers, *Paleoceanography*, 22, PA3212, doi:10.1029/2006PA001337, 2007.
- Nicol, D.: Recent species of the veneroid pelecypod *Arctica*, *J. Wash. Acad. Sci.*, 41, 102–106, 1951.
- Orr, J. C., Fabry, V. J., Aumont, O., Bopp, L., Doney, S. C., Feely, R. A., Gnanadesikan, A., Gruber, N., Ishida, A., Joos, F., Key, R. M., Lindsay, K., Maier-Reimer, E., Matear, R.,

$\delta^{11}\text{B}$, $^{87}\text{Sr}/^{86}\text{Sr}$ and $\delta^{88/86}\text{Sr}$ of cultured *Arctica islandica*

Y.-W. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Monfray, P., Mouchet, A., Najjar, R. G., Plattner, G.-K., Rodgers, K. B., Sabine, C. L., Sarmiento, J. L., Schlitzer, R., Slater, R. D., Totterdell, I. J., Weirig, M.-F., Yamanaka, Y., and Yool, A.: Anthropogenic ocean acidification over the twenty-first century and its impact on calcifying organisms, *Nature*, 437, 681–686, 2005.

5 Osberg, P. H., Hussey, A. M., and Boone, G. M.: Bedrock geologic map of Maine, Maine Geological Survey, Geologic Map Esries BGMM, 1985.

~~Pagani, M., Lemarchand, D., Spivack, A., and Gaillardet, J.: A critical evaluation of the boron isotope pH proxy: the accuracy of ancient ocean pH estimates, *Geochim. Cosmochim. Ac.*, 69, 953–961, 2005.~~

10 Patchett, P. J.: Sr isotopic fractionation in Allende chondrules: a reflection of solar nebular processes, *Earth Planet. Sc. Lett.*, 50, 181–188, 1980a.

Patchett, P. J.: Sr isotopic fractionation in Ca-Al inclusions from the Allende meteorite, *Nature*, 283, 438–441, 1980b.

15 Raddatz, J., Liebetrau, V., Rüggeberg, A., Hathorne, E., Krabbenhöft, A., Eisenhauer, A., Böhm, F., Vollstaedt, H., Fietzke, J., López Correa, M., Freiwald, A., and Dullo, W. C.: Stable Sr-isotope, Sr/Ca, Mg/Ca, Li/Ca and Mg/Li ratios in the scleractinian cold-water coral *Lophelia pertusa*, *Chem. Geol.*, 352, 143–152, doi:10.1016/j.chemgeo.2013.06.013, 2013.

Rae, J. W. B., Foster, G. L., Schmidt, D. N., and Elliott, T.: Boron isotopes and B/Ca in benthic foraminifera: proxies for the deep ocean carbonate system, *Earth Planet. Sc. Lett.*, 302, 403–413, 2011.

20 Reynaud, S., Hemming, N. G., Juillet-Leclerc, A., and Gattuso, J.-P.: Effect of $p\text{CO}_2$ and temperature on the boron isotopic composition of the zooxanthellate coral *Acropora sp.*, *Coral Reefs*, 23, 539–546, doi:10.1007/s00338-004-0399-5, 2004.

Reynaud, S., Rollion-Bard, C., Martin, S., Rolopho-Metalpa, R., and Gattuso, J.-P.: Effect of elevated $p\text{CO}_2$ on the boron isotopic composition into the Mediterranean scleractinian coral *Cladocora caespitosa*, CIESM Workshop Monographs No. 36, Monaco, ZOOREC:ZOOR14704033379, 2008.

25 Riebesell, U., Gattuso, J.-P., Thingstad, T., and Middelburg, J.: Preface “Arctic ocean acidification: pelagic ecosystem and biogeochemical responses during a mesocosm study”, *Biogeosciences*, 10, 5619–5626, 2013, <http://www.biogeosciences.net/10/5619/2013/>.

Rollion-Bard, C. and Erez, J.: Intra-shell boron isotope ratios in the symbiont-bearing benthic foraminiferan *Amphistegina lobifera*: implications for $\delta^{11}\text{B}$ vital effects and paleo-pH reconstructions, *Geochim. Cosmochim. Ac.*, 74, 1530–1536, 2010.

~~Rollion-Bard, C., Chaussidon, M., and France-Lanord, C.: Biological control of internal pH in scleractinian corals: implications on paleo-pH and paleo-temperature reconstructions, *CR Geosci.*, 343, 397–405, 2011.~~

Rüggeberg, A., Fietzke, J., Liebetrau, V., Eisenhauer, A., Dullo, W.-C., and Freiwald, A.: Stable strontium isotopes ($\delta^{88/86}\text{Sr}$) in cold-water corals – a new proxy for reconstruction of intermediate ocean water temperatures, *Earth Planet. Sc. Lett.*, 269, 570–575, doi:10.1016/j.epsl.2008.03.002, 2008.

Sabine, C. L., Feely, R. A., Gruber, N., Key, R. M., Lee, K., Bullister, J. L., Wanninkhof, R., Wong, C., Wallace, D. W., and Tilbrook, B.: The oceanic sink for anthropogenic CO_2 , *Science*, 305, 367–371, 2004.

Sanyal, A., Hemming, N. G., Hanson, G. N., and Broecker, W. S.: Evidence for a higher pH in the glacial ocean from boron isotopes in foraminifera, *Nature*, 373, 234–236, 1995.

Schöne, B. R., Oschmann, W., Rössler, J., Castro, A. D. F., Houk, S. D., Kröncke, I., Dreyer, W., Janssen, R., Rumohr, H., and Dunca, E.: North Atlantic Oscillation dynamics recorded in shells of a long-lived bivalve mollusk, *Geology*, 31, 1037–1040, 2003.

Schöne, B. R., Fiebig, J., Pfeiffer, M., Gleß, R., Hickson, J., Johnson, A. L. A., Dreyer, W., and Oschmann, W.: Climate records from a bivalved *Methuselah* (*Arctica islandica*, Mollusca; Iceland), *Palaeogeogr. Palaeocl.*, 228, 130–148, 2005.

Schöne, B. R., Wanamaker Jr., A. D., Fiebig, J., Thébault, J., and Kreutz, K.: Annually resolved $\delta^{13}\text{C}$ shell chronologies of long-lived bivalve mollusks (*Arctica islandica*) reveal oceanic carbon dynamics in the temperate North Atlantic during recent centuries, *Palaeogeogr. Palaeocl.*, 302, 31–42, 2011a.

Schöne, B. R., Zhang, Z., Radermacher, P., Thébault, J., Jacob, D. E., Nunn, E. V., and Maurer, A.-F.: Sr/Ca and Mg/Ca ratios of ontogenetically old, long-lived bivalve shells (*Arctica islandica*) and their function as paleotemperature proxies, *Palaeogeogr. Palaeocl.*, 302, 52–64, 2011b.

Schöne, B. R., Radermacher, P., Zhang, Z., and Jacob, D. E.: Crystal fabrics and element impurities (Sr/Ca, Mg/Ca, and Ba/Ca) in shells of *Arctica islandica* – Implications for paleoclimate reconstructions, *Palaeogeogr. Palaeocl.*, 373, 50–59, 2013.

BGD

12, 2979–3023, 2015

$\delta^{11}\text{B}$, $^{87}\text{Sr}/^{86}\text{Sr}$ and $\delta^{88/86}\text{Sr}$ of cultured *Arctica islandica*

Y.-W. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



$\delta^{11}\text{B}$, $^{87}\text{Sr}/^{86}\text{Sr}$ and $\delta^{88/86}\text{Sr}$ of cultured *Arctica islandica*

Y.-W. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Scourse, J., Richardson, C., Forsythe, G., Harris, I., Heinemeier, J., Fraser, N., Briffa, K., and Jones, P.: First cross-matched floating chronology from the marine fossil record: data from growth lines of the long-lived bivalve mollusc *Arctica islandica*, *The Holocene*, 16, 967–974, 2006.

5 Scourse, J. D., Wanamaker Jr., A. D., Weidman, C. R., Heinemeier, J., Reimer, P. J., Butler, P. G., Witbaard, R., and Richardson, C. A.: The marine radiocarbon bomb pulse across the temperate North Atlantic: a compilation of $\delta^{14}\text{C}$ time histories from *Arctica islandica* growth increments, *Radiocarbon*, 54, 165–186, 2012.

10 Shen, C.-C., Lee, T., Chen, C.-Y., Wang, C.-H., Dai, C.-F., and Li, L.-A.: The calibration of $\text{D}[\text{Sr}/\text{Ca}]$ versus sea surface temperature relationship for *Porites* corals, *Geochim. Cosmochim. Ac.*, 60, 3849–3858, 1996.

Shinjo, R., Asami, R., Huang, K.-F., You, C.-F., and Iryu, Y.: Ocean acidification trend in the tropical North Pacific since the mid-20th century reconstructed from a coral archive, *Mar. Geol.*, 342, 58–64, doi:10.1016/j.margeo.2013.06.002, 2013.

15 Stemmer, K.: Shell formation and microstructure of the ocean quahog *Arctica islandica*: Does ocean acidification matter?, Ph.D. thesis, Staats- und Universitaetsbibliothek Bremen, Bremen, 143 pp., 2013.

20 Stevenson, E. I., Hermoso, M., Rickaby, R. E. M., Tyler, J. J., Minoletti, F., Parkinson, I. J., Mokadem, F., and Burton, K. W.: Controls on stable strontium isotope fractionation in coccolithophores with implications for the marine Sr cycle, *Geochim. Cosmochim. Ac.*, 128, 225–235, doi:10.1016/j.gca.2013.11.043, 2014.

25 Trotter, J., Montagna, P., McCulloch, M., Silenzi, S., Reynaud, S., Mortimer, G., Martin, S., Ferrier-Pagès, C., Gattuso, J.-P., and Rodolfo-Metalpa, R.: Quantifying the pH “vital effect” in the temperate zooxanthellate coral *Cladocora caespitosa*: validation of the boron seawater pH proxy, *Earth Planet. Sc. Lett.*, 303, 163–173, doi:10.1016/j.epsl.2011.01.030, 2011.

Vázquez-Rodríguez, M., Pérez, F. F., Velo, A., Ríos, A. F., and Mercier, H.: Observed acidification trends in North Atlantic water masses, *Biogeosciences*, 9, 5217–5230, doi:10.5194/bg-9-5217-2012, 2012.

30 Vollstaedt, H., Eisenhauer, A., Wallmann, K., Böhm, F., Fietzke, J., Liebetrau, V., Krabbenhöft, A., Farkaš, J., Tomašových, A., Raddatz, J., and Veizer, J.: The Phanerozoic $\delta^{88/86}\text{Sr}$ record of seawater: new constraints on past changes in oceanic carbonate fluxes, *Geochim. Cosmochim. Ac.*, 128, 249–265, doi:10.1016/j.gca.2013.10.006, 2014.

$\delta^{11}\text{B}$, $^{87}\text{Sr}/^{86}\text{Sr}$ and
 $\delta^{88/86}\text{Sr}$ of cultured
Arctica islandica

Y.-W. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Wanamaker Jr., A. D., Heinemeier, J., Scourse, J. D., Richardson, C. A., Butler, P. G., Eiríksson, J., and Knudsen, K. L.: Very long-lived mollusks confirm 17th century AD tephra-based radiocarbon reservoir ages for north Icelandic shelf waters, *Radiocarbon*, 50, 399–412, 2008a.
- 5 Wanamaker Jr., A. D., Kreutz, K. J., Schöne, B. R., Pettigrew, N., Borns, H. W., Introne, D. S., Belknap, D., Maasch, K. A., and Feindel, S.: Coupled North Atlantic slope water forcing on Gulf of Maine temperatures over the past millennium, *Clim. Dynam.*, 31, 183–194, 2008b.
- Wanamaker Jr., A. D., Kreutz, K. J., Schöne, B. R., Maasch, K. A., Pershing, A. J., Borns, H. W., Introne, D. S., and Feindel, S.: A late Holocene paleo-productivity record in the western
 10 Gulf of Maine, USA, inferred from growth histories of the long-lived ocean quahog (*Arctica islandica*), *Int. J. Earth Sci.*, 98, 19–29, doi:10.1007/s00531-008-0318-z, 2009.
- Wanamaker Jr., A. D., Kreutz, K. J., Schöne, B. R., and Introne, D. S.: Gulf of Maine shells reveal changes in seawater temperature seasonality during the Medieval Climate Anomaly and the Little Ice Age, *Palaeogeogr. Palaeoclimatol.*, 302, 43–51, 2011.
- 15 Wanamaker Jr., A. D., Butler, P. G., Scourse, J. D., Heinemeier, J., Eiríksson, J., Knudsen, K. L., and Richardson, C. A.: Surface changes in the North Atlantic meridional overturning circulation during the last millennium, *Nature Communications*, 3, 899, doi:10.1038/ncomms1901, 2012.
- Weber, J. N.: Incorporation of strontium into reef coral skeletal carbonate, *Geochim. Cosmochim. Ac.*, 37, 2173–2190, doi:10.1016/0016-7037(73)90015-X, 1973.
- 20 Weidman, C. R.: Development and Application of the Mollusc *Arctica Islandica* as a Paleoceanographic Tool for the North Atlantic Ocean, DTIC Document, 1995.
- Weidman, C. R. and Jones, G. A.: A Shell-Derived Time History of Bomb ^{14}C on Georges Bank and Its Labrador Sea Implications, *J. Geophys. Res.*, 98, 14577–514588, 1993.
- 25 Weidman, C. R., Jones, G. A., and Lohman, L.: The long-lived mollusc *Arctica islandica*: a new paleoceanographic tool for the reconstruction of bottom temperatures for the continental shelves of the northern North Atlantic Ocean, *J. Geophys. Res.*, 99, 18305–318314, 1994.
- Widerlund, A. and Andersson, P. S.: Strontium isotopic composition of modern and Holocene mollusc shells as a palaeosalinity indicator for the Baltic Sea, *Chem. Geol.*, 232, 54–66, 2006.
- 30 Witbaard, R. and Bergman, M. J. N.: The distribution and population structure of the bivalve *Arctica islandica* L. in the North Sea: what possible factors are involved?, *J. Sea Res.*, 50, 11–25, 2003.

Yan, H., Shao, D., Wang, Y., and Sun, L.: Sr/Ca profile of long-lived *Tridacna gigas* bivalves from South China Sea: a new high-resolution SST proxy, *Geochim. Cosmochim. Ac.*, 112, 52–65, doi:10.1016/j.gca.2013.03.007, 2013.

5 Yu, K.-F., Zhao, J.-X., Wei, G.-J., Cheng, X.-R., Chen, T.-G., Felis, T., Wang, P.-X., and Liu, T.-S.: $\delta^{18}\text{O}$, Sr/Ca and Mg/Ca records of *Porites lutea* corals from Leizhou Peninsula, northern South China Sea, and their applicability as paleoclimatic indicators, *Palaeogeogr. Palaeoclimatol.*, 218, 57–73, 2005.

10 Zhang, Z.: Geochemical properties of shells of *Arctica islandica* (Bivalvia) – Implications for Environmental and Climatic Change, PhD thesis, Goethe-Universität in Frankfurt am Main, Frankfurt, 109 pp., 2009.

BGD

12, 2979–3023, 2015

$\delta^{11}\text{B}$, $^{87}\text{Sr}/^{86}\text{Sr}$ and $\delta^{88/86}\text{Sr}$ of cultured *Arctica islandica*

Y.-W. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



$\delta^{11}\text{B}$, $^{87}\text{Sr}/^{86}\text{Sr}$ and $\delta^{88/86}\text{Sr}$ of cultured *Arctica islandica*

Y.-W. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 1. Summary of in situ instrumental data of tank seawater salinity, temperature, and pH during culture season .



Week number (dates)	Salinity (PSU) ¹	Average to subsampling interval ³	Temperature (°C) ¹	Average to subsampling interval ³	pH ¹	Average to subsampling interval ³	Predicted $\delta^{11}\text{B}$ with $\alpha = 1.0194^2$	Predicted $\delta^{11}\text{B}$ with $\alpha = 1.0272^2$
1 (1/17–1/23)	31.036	30.608 ± 0.683	3.029	2.960 ± 0.511	–	8.033 ± 0.019	21.78	14.89
2 (1/24 –1/30)	30.508		3.030		–		21.90	15.06
3 (1/31 –2/6)	30.520		2.375		–		22.03	15.25
4 (2/7 –2/13)	31.066		2.369		8.018		21.82	14.95
5 (2/14 –2/20)	31.186		2.776		8.037		21.92	15.10
6 (2/21 –2/27)	30.755		3.367		8.059		22.05	15.27
7 (2/28–3/6)	29.183		3.774		8.019		22.34	15.68
8 (3/7–3/13)	29.288	29.516 ± 0.322	4.596	4.698 ± 0.145	8.029	8.039 ± 0.014	22.27	15.57
9 (3/14–3/19 (3/20))	29.743		4.800		8.049		22.35	15.69
10 (3/21–3/27)	29.751	29.149 ± 0.851	5.646	5.735 ± 0.127	8.094	8.089 ± 0.006	22.36	15.70
11 (3/28–4/3)	28.547		5.825		8.085		22.54	15.95
12 (4/4–4/10)	28.011	28.745 ± 1.037	7.252	7.205 ± 0.066	8.085	8.082 ± 0.005	22.65	16.09
13 (4/11 (4/14)– 4/17)	29.478		7.158		8.078		22.68	16.14
14 (4/18–4/24)	29.751	29.827 ± 0.107	7.594	8.246 ± 0.922	8.102	8.102 ± 0.000	23.02	16.61
15 (4/25–5/1)	29.902		8.898		8.102		23.04	16.64
16 (5/2–5/8)	30.335	30.702 ± 0.341	9.662	9.784 ± 0.494	8.095	8.128 ± 0.029	23.09	16.72
17 (5/9–5/15)	30.762		9.364		8.150		23.18	16.83
18 (5/16–5/22)	31.008		10.328		8.139		22.88	16.42

$\delta^{11}\text{B}$, $^{87}\text{Sr}/^{86}\text{Sr}$ and $\delta^{88}/^{86}\text{Sr}$ of cultured *Arctica islandica*

Y.-W. Liu et al.

Table 1. Continued.

Week number (dates)	Salinity (PSU) ^a	Average to subsampling interval ^c	Temperature (°C) ^a	Average to subsampling interval ^c	pH ^a	Average to subsampling interval ^c	Predicted $\delta^{11}\text{B}$ with $\alpha = 1.0194^b$	Predicted $\delta^{11}\text{B}$ with $\alpha = 1.0272^b$
19 (5/23–5/29)	31.102	31.099 ± 0.129	12.653	13.170 ± 0.450	8.118	8.105 ± 0.025	22.89	16.44
20 (5/30–6/5)	31.227		13.385		8.120		23.31	17.02
21 (6/6–6/12)	30.969		13.472		8.076		23.59	17.40
22 (6/13–6/19)	31.067	30.829 ± 0.454	13.427	14.494 ± 0.989	8.078	8.121 ± 0.041	22.71	16.18
23 (6/20–6/26)	31.115		14.676		8.125		22.44	15.81
24 (6/27–7/3)	30.306		15.380		8.159		22.58	16.01
25 (7/4–7/10)	30.508	30.521 ± 0.030	16.445	17.076 ± 0.577	8.015	7.985 ± 0.028	22.46	15.84
26 (7/11–7/17)	30.555		17.209		7.960		22.57	15.99
27 (7/18–7/24)	30.500		17.576		7.981		21.78	14.89
28 (7/25–7/31)	30.305	30.466 ± 0.227	17.188	17.107 ± 0.115	7.966	7.976 ± 0.013	21.90	15.06
29 (8/1–8/5)	30.626		17.026		7.985		22.03	15.25

^a Data from Beirne et al. (2012).^b Calculation based on Eq. (2).^c SD of the data.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



$\delta^{11}\text{B}$, $^{87}\text{Sr}/^{86}\text{Sr}$ and $\delta^{88}/^{86}\text{Sr}$ of cultured *Arctica islandica*

Y.-W. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 3. Summary of shell data.



Sample ID	Week number	$^{87}\text{Sr}/^{86}\text{Sr}$ (2 SE)	$\delta^{88}/^{86}\text{Sr}$ (‰)	[Sr] (ppm)	$\delta^{11}\text{B}$ (‰)	Average growth rate (mm/week) ¹	pH _{shell}	ΔpH
A101JV-8	22	–	–	–	20.32	–	8.411	0.29
A101JV-9	24.8	–	–	–	23.34	–	8.588	0.603
A102JV-1	4	0.709181 (7)	0.132	1403	15.68	0.27	8.125	0.092
A102JV-2	8.5	0.709210 (10)	0.151	1282	14.94	0.53	8.003	–0.036
A102JV-3	10.5	0.709181 (10)	0.158	1422	14.28	–	7.878	–0.211
A102JV-4	12.5	0.709157 (15)	0.112	1452	–	–	–	–
A102JV-5	14.5	0.709201 (7)	0.118	1435	13.77	–	7.732	–0.371
A102JV-6	16.4	0.709194 (12)	0.124	1319	13.65	0.65	7.679	–0.449
A102JV-7	19.2	0.709169 (13)	0.154	1474	16.80	–	8.121	0.016
A102JV-8	22	0.709170 (5)	0.099	1558	18.04	–	8.226	0.106
A102JV-9	24.8	0.709178 (6)	0.112	1491	20.23	–	8.374	0.389
A102JV-10	27.6	0.709184 (6)	0.097	–	21.93	–	8.495	0.520
A103JV-1	4	–	–	–	16.57	0.24	8.230	0.197
A103JV-2	8.5	–	–	–	16.57	0.51	8.214	0.175
A103JV-3	10.5	0.709191 (7)	–	–	18.41	–	8.380	0.291
A103JV-4	12.5	0.709187 (5)	–	–	16.04	–	8.125	0.044
A103JV-5	14.5	0.709195 (9)	–	–	16.28	–	8.133	0.031
A103JV-6	16.4	0.709204 (15)	–	–	17.71	0.61	8.255	0.128
A103JV-7	19.2	–	–	–	16.57	–	8.096	–0.008
A103JV-8	22	0.709188 (6)	–	–	17.45	–	8.171	0.051
A103JV-9	24.8	0.709182 (5)	–	–	19.21	–	8.296	0.311
A103JV-10	27.6	0.709182 (5)	–	–	25.12	–	8.702	0.726
A105JV-1	4	0.709178 (6)	0.227	1414	14.53	0.24	7.951	–0.082
A105JV-2	8.5	0.709191 (7)	0.255	1534	13.75	0.60	7.777	–0.262
A105JV-3	10.5	0.709183 (6)	0.251	1482	12.22	–	7.076	–1.013
A105JV-4	12.5	0.709184 (5)	0.263	1453	17.15	–	8.247	0.165
A105JV-5	14.5	0.709175 (7)	0.216	1521	14.59	–	7.898	–0.205
A105JV-6	16.4	0.709179 (6)	0.273	2559	11.09	0.68	–	–
A105JV-7	19.2	0.709183 (5)	0.296	2048	12.52	–	7.211	–0.894
A105JV-8	22	0.709163 (5)	0.230	2204	17.41	–	8.168	0.047
A105JV-9	24.8	0.709186 (6)	0.234	1850	19.43	–	8.314	0.328
A105JV-10	27.6	0.709177 (6)	0.238	1795	22.11	–	8.507	0.532

¹ Data from Beirne (2011).




Figure 1. Photos of (a) adult and (b), (c) juvenile *A. islandica* from the culture experiment. (d) shows the corresponding in situ measurements of tank water salinity, temperature and pH during the 31-week culture experiment. The juvenile shells were sampled in 10 intervals for this study (c). Note that because the growth rates differ during the season, each interval represents different durations (c) and (d).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



$\delta^{11}\text{B}$, $^{87}\text{Sr}/^{86}\text{Sr}$ and $\delta^{88}/^{86}\text{Sr}$ of cultured *Arctica islandica*

Y.-W. Liu et al.




Figure 2. The illustrations of (a) the 84–87 Sr double spike method, (b) how the angle between Mixture-Spike plane and Sample-Spike plane can influence the precision of the deconvolved result, and (c) the optimal sample-spike ratio in our study.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



$\delta^{11}\text{B}$, $^{87}\text{Sr}/^{86}\text{Sr}$ and $\delta^{88/86}\text{Sr}$ of cultured *Arctica islandica*

Y.-W. Liu et al.




Figure 3. Stable Sr results for (a) seawater standard IAPSO and (b) inter-laboratory biogenic carbonate standards JCp-1.

$\delta^{11}\text{B}$, $^{87}\text{Sr}/^{86}\text{Sr}$ and $\delta^{88}/^{86}\text{Sr}$ of cultured *Arctica islandica*

Y.-W. Liu et al.




Figure 4. Long-term precision of (a) boric acid standard SRM 951a, (b) seawater standard IAEA B-1, and (c) inter-laboratory carbonate standard JCP-1.

$\delta^{11}\text{B}$, $^{87}\text{Sr}/^{86}\text{Sr}$ and $\delta^{88/86}\text{Sr}$ of cultured *Arctica islandica*

Y.-W. Liu et al.




Figure 5. GoM $^{87}\text{Sr}/^{86}\text{Sr}$ data for (a) seawater samples and (b) shell samples and the double spike deconvolved (c) $\delta^{88/86}\text{Sr}$ values and (d) Sr concentrations for the juvenile shell.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Figure 6. GoM boron data for (a) seawater samples and (b) shell samples. Two prediction lines based on two boron fractionation factors: $\alpha = 1.0272$ (Klochko et al., 2006) and $\alpha = 1.0194$ (Kakihana et al., 1977).



(a)



(b)

Figure 7. The comparisons between the shell $\delta^{11}\text{B}$ to (a) the corresponding culture water temperature and (b) the growth rates for individual shells.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

