Biogeosciences Discuss., 12, C2750–C2754, 2015 www.biogeosciences-discuss.net/12/C2750/2015/ © Author(s) 2015. This work is distributed under the Creative Commons Attribute 3.0 License.



BGD 12, C2750–C2754, 2015

> Interactive Comment

Interactive comment on "The influence of C₃ and C₄ vegetation on soil organic matter dynamics in contrasting semi-natural tropical ecosystems" by G. Saiz et al.

Anonymous Referee #3

Received and published: 12 June 2015

General comments:

The current paper addresses the topic of land use change (i.e. vegetation thickening) and its effects on soil biogeochemical cycling, an important topic considering current and future changes associated with global climate change. The study also discusses the effects of land use changes associated with woody encroachment on belowground C/OM dynamics, a key component, and yet largely overlooked, of the global C cycle. The authors make use of C isotopes (ïĄd'13C and 14C) to quantify the contribution of C3 and C4 vegetation on SOM dynamics in semi-natural tropical (mixed vegetation type) ecosystems along a precipitation gradient in western Africa. The topics addressed are therefore well within the scope of Biogeosciences and would be of interest





to the readers of this journal and the general soil science community. The manuscript is well written and logical throughout. The English is clear and of an acceptable standard (some minor changes are suggested in the technical corrections below).

Specific comments:

(1) In the methodology section (p.8) the soil sampling strategy is described. Three soil samples were collected for the upper soil surface layer (0-0.05m) however just one sample was collected at 0-0.3m. Why? I assume this was mainly related to ensuring enough sample volume for the lab analyses? Please clarify why there is this discrepancy.

(2) In the results section, the authors mention the discrepancy between -T and -G sampling locations at each site was consistently larger at the savanna sites compared with the forest sites – why? Can you provide an explanation for this observation?

(3) In the discussion section (4.2 Differential patterns in SOM dynamics across contrasting C3/C4 mixed ecosystems) the authors mention the difficulties associated with quantifying belowground litter dynamics. Did you make an attempt to quantify rhizodeposition? Such data would very interesting to see.

(4) Figure 2(b) shows the relationship between weighted d13C and fractional vegetation cover for the o-0.05m surface soil layer. Were similar relationships found for the 0-0.3m soil layer? Why is this data not shown?

(5) Figure 6, site BDA-03 has no data point for the deeper samples (i.e. 0.3-0.5 - 0-0.05m). Is this a mistake or are no data present?

(6) The manuscript has a total of 94 references. This is a very large number of references! Perhaps the authors could be a little more selective in some sections.

Technical corrections:

p. 2 line 42: with this trend also being

12, C2750-C2754, 2015

Interactive Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



p2. line 45: to minimise the confounding

p2. line 47-8: even in deep soil layers, while the most stable SOM fraction associated with silt and clay

- p2. line 49: These results, together with...
- p2. line 54: 'are at variance' what does this mean? Please clarify/change text.
- p3. line 79-80: useful tool for investigating the influence...
- p4. line 81: and for identifying recent...
- p4. line 85: utilize
- p4. line 91: those associated with
- p5. line 120: it remains a challenge to assess
- p5. line 124: ecosystem processes and studying the potential impacts
- p5. line 129: this expansive region
- p6. line 137: can help assess

p6. line 140-3: of this study are: (1) delineate the spatial...(2) investigate any potential...(3) unambiguously evaluate...

- p7. line 149-150: present work have been provided in detail
- p7. line 152: Hence, a short summary is provided here
- p7. line 155: Ghana, Burkina Faso, and Mali
- p7. line 160: the latter also being the case for
- p7. line 162: The transect was established on consistently flat terrain
- p7. line 165: nutrient poor Arenosols on the Southern border

C2752

BGD

12, C2750-C2754, 2015

Interactive Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



p8. line 175: that has been proven to be well suited

p8. line 183: 40mm inner diameter (ïĄę). All samples were placed in labelled zip-lock bags.

p9. line 203-4: any traces of salt, dried at 40oC, and the weight of each fraction was determined...

- p10. line 231: depth intervals to help explain potential variations
- p11. line 255: the southern end of the transect
- p11. line 260: lower C/N values were associated with
- p13. line 292: a Sahelian site showed a gradual decrease
- p13. line 297: increase in SOM ïĄď13C values with soil depth
- p15. Line 326-7: reflects current vegetation patterns well
- p16. line 343: major effect on both the physical protection
- p16. line 359: Indeed, there was considerable
- p17. line 364-5: by some earlier studies, suggests that such variation
- p17. line 372: C/N ratios, a feature considered to be highly relevant

p17. line 383-4: soil fraction that best reflects recent organic inputs to the soil, as it includes contributions

p18. line 410: much lower abundance of C4 vegetation which seems is progressively

- p18. line 413: consequently an overall negative
- p19. line 430: litter carbon chemistry is a key factor
- p19. line 437: extrapolated given that these

BGD

12, C2750-C2754, 2015

Interactive Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- p20. line 448: between the surface (0-0.05m)
- p20. line 452: at the wetter sites along the transect
- p20. line 454: main determinants of vegetation type observed at each site
- p20. line 460: agrees well with the long-term persistence
- p22. line 491: fractions associated with silt and clay
- p22. line 505: combined effect of several factors, including OM decomposition
- p22. line 508: influenced by specific properties
- p23. line 521: These sites are characterized by a low abundance
- p23. line 528: root biomass with depth, this trend is more obvious
- p24. line 554: between the two sites is strongly influenced
- p27. line 582: The reference 14CO2
- p27. line 584: Simple interpolation was used to quantify the ïĄť

Interactive comment on Biogeosciences Discuss., 12, 8085, 2015.

BGD

12, C2750-C2754, 2015

Interactive Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

