

Interactive comment on “Global soil nitrous oxide emissions in a dynamic carbon–nitrogen model” by Y. Y. Huang and S. Gerber

Y. Y. Huang and S. Gerber

sgerber@ufl.edu

Received and published: 3 July 2015

Many thanks for reviewing our manuscript and your suggestions are very help in improving our study. Please find our responses below.

Reviewer: This MS presents a module for simulating N₂O fluxes at the global scale based on equations for denitrification and nitrification and considering N₂O and NO_x as fractions of the nitrogen that is processed. Most model elements were borrowed from other models. I have a number of serious problems with this MS:

Response: Thank you for taking time reviewing our paper.

Reviewer: The model description in Appendix A is not complete as the units are not provided.

C3303

Response: We agree. Appendix A is rewritten and moved to sect. 2.1.2 Soil N₂O emission. Details on units and further explanations were added.

Reviewer: It is not clear how model calculations at a resolution of 3.75 by 2.5 degrees can be meaningful, since all data such as weather, soil and vegetation are kind of aggregates for that resolution, and how can this be compared with point measurements.

Response: We agree with the reviewer that evaluations of global simulations against point measurements incur scale mismatches. Soil N₂O fluxes are highly variable even at stand scales. Currently, a higher resolution that is feasible for global simulation (e.g. 0.5 by 0.5 degrees) is still much larger than the area where field measurements take place, and benchmarking global simulations against field measurements still faces scale mismatches. In this way, any model data comparison is incomplete. In the original manuscript, we focus on capturing the average of annual mean emissions across different observations instead of one-by-one comparison. The overall average emission aggregates measurements and provides some information on model performance.

Reviewer: With this spatial resolution, the time step is 30 minutes, but the authors provide annual and seasonal numbers only. It is probably more interesting to compare the model results with temporal distributions from field measurements. This could be done for a number of test sites in a variety of climate and soil conditions. If not available, perhaps seasonal estimates from experimental sites could be used to validate the model.

Response: We have now added monthly and daily measurements to sect. 2.3 and 3.3.

Reviewer: To assess model quality, it is much more interesting to analyze the functioning of the soil-plant nitrogen cycle. How is denitrification compared to field measurements, and leaching, plant uptake, ammonia volatilization, etc. If the large flows in the system are correct, the authors will also be more confident about the small fluxes like N₂O and NO_x.

C3304

Response: Assessing model quality is a challenge for all models, and several reviewers have pointed out the effect of the larger N cycle on nitrification, denitrification and associated N₂O and NO_x fluxes. We now present an extensive sensitivity analysis that evaluates N₂O fluxes in response to variation the relative strength of plant uptake, the overall supply of N to mineralization (by way of modifying biological N fixation), and by specific parameters that deal with nitrification and denitrification. We discuss the hierarchical system of plants and soils N demand are strong sinks for ammonium and nitrate under limiting condition, resulting in leaching and denitrification very small amounts. Excess nitrogen (if plants and soils have sufficient N) leads to leaching and denitrification. There the partition coefficients and the parameters that determine the fate of the excess N become very important. The sensitivity analysis targets exactly that question. We further point out, that we do not invent a new model but put existing formulation (and parameter choices) in a new model. Currently, we do not have large scale observation data such as denitrification and ammonia volatilization in relative pristine ecosystems to support the benchmarking of the global model. Detailed analyses addressing this are now in sect. 2.2.3, 3.4 and the discussion part of the revised manuscript.

Reviewer: Finally, a true sensitivity analysis will also show what the major variables and parameters are. For example, the N₂O and NO_x fractions will probably pop up as important coefficients.

Response: We add more sensitivity tests with regard to plant N uptake, overall N input and losses, nitrification rates, denitrification rates and the fraction of N₂O lost from net nitrification in sect. 2.2.3 and sect.3.4 of the revised manuscript. These analysis help elucidate the dependency of N₂O emissions on specific parts of the N cycle. The fraction of N₂O lost from net nitrification is the most sensitive parameter. However, this fraction is very uncertain based on limited field or laboratory studies. Goodroad and Keeney (1984) suggested a value of 0.1-0.2% , while Khalil et al. (2004) reported a range of 0.16%-1.48% depending on the O₂ concentration. We applied a value of

C3305

contant 0.4% in the default run which embraces large uncertainties in our modelled results.

Reviewer: I fully agree with one of the other reviewers who states that this work is not ready for publication, and I also agree that perhaps Geoscientific Model Development is a more appropriate journal for submitting a revised MS.

Response: Thanks again for the helpful suggestions. We carefully considered the possibility for Geoscientific Model Development but as laid out in the response to Beni Stocker, the nature of this work lends itself to Biogeosciences. One of the reason is that we do not develop a new model, but implement an existing denitrification and N₂O emission model into an also existing coupled carbon-nitrogen model.

References

Gerber, S., Hedin, L. O., Oppenheimer, M., Pacala, S. W., and Shevliakova, E.: Nitrogen cycling and feedbacks in a global dynamic land model, *Global Biogeochemical Cycles*, 24,doi:10.1029/2008gb003336, 2010.

Goodroad, L. L., and Keeney, D. R.: Nitrous oxide emission from forest, marsh, and prairie ecosystems, *Journal of Environmental Quality*, 13, 448-452, 1984. Khalil, K., Mary, B., and Renault, P.: Nitrous oxide production by nitrification and denitrification in soil aggregates as affected by O₂ concentration, *Soil Biology & Biochemistry*, 36, 687-699, doi:10.1016/j.soilbio.2004.01.004, 2004.

Interactive comment on Biogeosciences Discuss., 12, 3101, 2015.

C3306